

Multimode Brownian oscillators: Exact solutions to heat transport

Xin-Hai Tong

*CAS Key Laboratory of Precision and Intelligent Chemistry, University of Science and Technology of China, Hefei, Anhui 230026, China and
School of Physics and Technology, Wuhan University, Wuhan, Hubei 430072, China*

Hong Gong

CAS Key Laboratory of Precision and Intelligent Chemistry, University of Science and Technology of China, Hefei, Anhui 230026, China

Yao Wang,^{a)} Rui-Xue Xu,^{b)} and YiJing Yan

Hefei National Research Center for Physical Sciences at the Microscale, University of Science and Technology of China, Hefei, Anhui 230026, China

(Dated: 3 July 2023)

In this work, we investigate the multimode Brownian oscillators in nonequilibrium scenarios with multiple reservoirs at different temperatures. For this purpose, an algebraic method is proposed. This approach gives the exact time-local equation of motion for reduced density operator, from which we can easily extract not only the reduced system but also hybrid bath dynamical information. The resulted steady-state heat current is found numerically consistent with another discrete imaginary-frequency method followed by the Meir-Wingreen's formula. It is anticipated that the development in this work would constitute an indispensable component to nonequilibrium statistical mechanics for open quantum systems.

I. INTRODUCTION

Open quantum systems play pivotal roles in diversified fields such as nuclear magnetic resonance,^{1–3} condensed matter and material physics,^{4–6} high energy physics,^{7–9} quantum optics,^{10–12} chemical and biological physics,^{13–15} and nonlinear spectroscopies.^{16–18} In most of these studies, the system and its environment constitute a thermodynamic composite. Thermal effects dictate the system-environment entanglement, which is intimately related to the thermodynamic and transport properties.^{19–23} Practically, these properties are under focus in manipulating mesoscopic nanodevices.^{24–29}

For example, the real-time evolution of heavy quarkonium in the quark-gluon plasma was investigated by means of quantum Brownian motion theory to high-light dynamical mechanism of the relative motion of the quarkonium state.⁹ Another example is that when considering a point charge emitting radiation in an electromagnetic field, the internal degree of freedom of a moving atom can be modeled by a harmonic oscillator coupled to a scalar field. Following the same coupling way as scalar electrodynamics, the pathologies of radiation reactions can be recast as non-Markovian dynamics of a Brownian oscillator (BO) in an environment.²⁴ In terms of quantum decoherence and measurement, BO was also introduced as one of toy models to study universalities in the phenomenology of decoherence.²⁵ These continuous studies on BO indicate the significance of this model can be never underestimated.

As a typical open quantum system, BO is the simplest and exactly solvable model. It serves as an elementary consideration in various studies.^{30–36} Exact real-

time dynamics of BO has been widely investigated.^{37–43} Concerning thermodynamics, properties of the reduced BO system had been shown that the steady state could be expressed as the Gibbs state with a renormalized Hamiltonian.⁴⁴ On the other hand, it should be noticed that besides the reduced system, the hybrid bath properties need also be taken into account for the underlying entangled effects.^{23,45,46}

In this work, we develop an algebraic approach to obtain the equation of motion (EOM) for the multimode BO system, from which we can easily extract both the reduced system and hybrid bath dynamical information. Furthermore, we exploit the well-established EOM to study the heat transport problem. We also propose another discrete imaginary-frequency (DIF) method followed by the Meir-Wingreen's formula. Both the algebraic and DIF methods are consistent with each other in evaluating the steady-state heat current.

The remainder of this paper is organized as follows. In Sec. II we present the multimode BO system model and the exact EOM for dynamics. Especially, we emphasize how to extract the hybrid bath information and to evaluate the heat current. More theoretical details are given in Appendix. Numerical results are demonstrated in Sec. III. We summarize the paper in Sec. IV. Throughout this paper we set $\hbar = 1$ and $\beta_\alpha = 1/(k_B T_\alpha)$ with k_B being the Boltzmann constant and T_α the temperature of the α -reservoir.

II. MULTIMODE BROWNIAN OSCILLATOR

A. Hamiltonian and EOM

Let us start with the total system-bath Hamiltonian, $H_T = H_S + H_{SB} + h_B$. The bath $h_B = \sum_\alpha h_\alpha$ is assumed to be non-interacting with different temperatures β_α . The

^{a)}Electronic mail: wy2010@ustc.edu.cn

^{b)}Electronic mail: rxxu@ustc.edu.cn

BO model adopts

$$H_S = \sum_u \frac{\hat{p}_u^2}{2m_u} + \frac{1}{2} \sum_{uv} k_{uv} \hat{q}_u \hat{q}_v, \quad (1)$$

and

$$H_{SB} = \sum_{\alpha u j} \kappa_{\alpha u j} \hat{q}_u \hat{x}_{\alpha j} \equiv \sum_{\alpha u} \hat{Q}_u \hat{F}_{\alpha u}, \quad (2)$$

with

$$\hat{Q}_u = (m_u k_{uu})^{\frac{1}{4}} \hat{q}_u. \quad (3)$$

This is the dimensionless dissipative system mode that coupled to the random force $\hat{F}_{\alpha u}$ as specified in Eq. (2). Note also $\hat{P}_u = (m_u k_{uu})^{-\frac{1}{4}} \hat{p}_u$ so that $[\hat{Q}_u, \hat{P}_v] = i\delta_{uv}$. Thus Eq. (1) is recast as

$$H_S = \frac{1}{2} \sum_u \Omega_u \hat{P}_u^2 + \frac{1}{2} \sum_{uv} V_{uv} \hat{Q}_u \hat{Q}_v \quad (4)$$

with

$$\Omega_u \equiv \sqrt{\frac{k_{uu}}{m_u}} \quad \text{and} \quad V_{uv} \equiv \frac{k_{uv}}{\sqrt{m_u m_v k_{uu} k_{vv}}}. \quad (5)$$

In order to investigate some system-bath entangled properties such as the heat current, we introduce an auxiliary quantity

$$\tilde{\rho}_{\alpha u}(t) \equiv \text{tr}_B[\hat{F}_{\alpha u} \rho_T(t)]. \quad (6)$$

By applying the Liouville equation to the total density operator, $\dot{\rho}_T(t) = -i[H_T, \rho_T(t)]$, we obtain for the reduced $\rho_S(t) = \text{tr}_B \rho_T(t)$ the EOM as

$$\dot{\rho}_S(t) = -i[H_S, \rho_S(t)] - i \sum_{\alpha u} [\hat{Q}_u, \tilde{\rho}_{\alpha u}(t)]. \quad (7)$$

The above equation exists for any given H_S . It leads to the exact hierarchical equations of motion for general systems.⁴⁷ Focusing on the BO system, Eq. (7) can be evaluated in a closed form with

$$\begin{aligned} \tilde{\rho}_{\alpha u}(t) = \sum_v [\tilde{\Gamma}_{\alpha uv}(t) \hat{Q}_v^{\oplus} + \Gamma_{\alpha uv}(t) \hat{P}_v^{\oplus} \\ + \tilde{\zeta}_{\alpha uv}(t) \hat{Q}_v^{\ominus} - \zeta_{\alpha uv}(t) \hat{P}_v^{\ominus}] \rho_S(t) \end{aligned} \quad (8)$$

where

$$\hat{A}^{\oplus} \hat{O} \equiv \frac{1}{2} \{\hat{A}, \hat{O}\} \quad \text{and} \quad \hat{A}^{\ominus} \hat{O} \equiv -i[\hat{A}, \hat{O}]. \quad (9)$$

Apparently, Eq. (7) is trace preserving for the reduced density operator, since $\text{tr}_S \dot{\rho}_S(t) = 0$. The EOM (7)–(9) is exact and non-Markovian, fully taking account of the environmental temperature effects. As an exact approach, it will guarantee the positivity of ρ_S . Determination of the EOM (7)–(9) is detailed in Appendix.

B. Details of the time-dependent coefficients in the EOM

To specify the involved time-dependent coefficients in Eq. (8), let us first introduce a key quantity

$$\chi(t) \equiv \chi_{QQ}(t) \equiv \{\chi_{uv}^{QQ}(t) \equiv i\langle[\hat{Q}_u(t), \hat{Q}_v(0)]\rangle\} \quad (10)$$

with $\hat{Q}_u(t) \equiv e^{iH_T t} \hat{Q}_u e^{-iH_T t}$ and the average $\langle \dots \rangle$ over the steady state of total composite ρ_T^{st} which commutes with H_T . Similarly, we denote

$$\begin{aligned} \chi_{QP}(t) &\equiv \{\chi_{uv}^{QP}(t) \equiv i\langle[\hat{Q}_u(t), \hat{P}_v(0)]\rangle\} = -\dot{\chi}(t) \Omega^{-1}, \\ \chi_{PQ}(t) &\equiv \{\chi_{uv}^{PQ}(t) \equiv i\langle[\hat{P}_u(t), \hat{Q}_v(0)]\rangle\} = \Omega^{-1} \dot{\chi}(t), \\ \chi_{PP}(t) &\equiv \{\chi_{uv}^{PP}(t) \equiv i\langle[\hat{P}_u(t), \hat{P}_v(0)]\rangle\} = -\Omega^{-1} \ddot{\chi}(t) \Omega^{-1}, \end{aligned}$$

where $\Omega \equiv \{\Omega_{uv} \delta_{uv}\}$. The associated initial values are $\chi(0) = \ddot{\chi}(0) = \mathbf{0}$ and $\dot{\chi}(0) = \Omega$.

The time evolutions of system coordinates and momenta can be resolved as^{32,43,48}

$$\begin{bmatrix} \hat{Q}(t) \\ \hat{P}(t) \end{bmatrix} = \mathbf{T}(t) \begin{bmatrix} \hat{Q}(0) \\ \hat{P}(0) \end{bmatrix} - \sum_{\alpha} \int_0^t d\tau \mathbf{T}(t-\tau) \begin{bmatrix} \mathbf{0} \\ \hat{\mathbf{F}}_{\alpha}^{\text{B}}(\tau) \end{bmatrix}. \quad (11)$$

For compactness, we have introduced vectors, $\hat{\mathbf{P}} \equiv \{\hat{P}_u\}$, $\hat{\mathbf{Q}} \equiv \{\hat{Q}_u\}$, $\hat{\mathbf{F}}_{\alpha}^{\text{B}} \equiv \{\hat{F}_{\alpha u}^{\text{B}}\}$, and the matrix

$$\mathbf{T}(t) = \begin{bmatrix} -\chi_{QP}(t) & \chi_{QQ}(t) \\ -\chi_{PP}(t) & \chi_{PQ}(t) \end{bmatrix} = \begin{bmatrix} \dot{\chi}(t) \Omega^{-1} & \chi(t) \\ \Omega^{-1} \dot{\chi}(t) \Omega^{-1} & \Omega^{-1} \dot{\chi}(t) \end{bmatrix}. \quad (12)$$

By taking the time derivative of Eq. (11) and eliminating the initial values using itself, the time-local EOM can be obtained as^{32,43,48}

$$\begin{bmatrix} \dot{\hat{Q}}(t) \\ \dot{\hat{P}}(t) \end{bmatrix} = \mathbf{\Lambda}(t) \begin{bmatrix} \hat{Q}(t) \\ \hat{P}(t) \end{bmatrix} - \sum_{\alpha} \begin{bmatrix} \mathbf{0} \\ \hat{\mathbf{F}}_{\alpha}^{\text{eff}}(t) \end{bmatrix}. \quad (13)$$

Here, the matrix $\mathbf{\Lambda}(t)$ is resulted as

$$\mathbf{\Lambda}(t) = \dot{\mathbf{T}}(t) \mathbf{T}^{-1}(t) = \begin{bmatrix} \mathbf{0} & \Omega \\ -\mathbf{V} - \tilde{\Gamma}(t) & -\Gamma(t) \end{bmatrix} \quad (14)$$

and (denoting $\tilde{\mathbf{V}}(t) \equiv \tilde{\Gamma}(t) + \mathbf{V}$)

$$\begin{aligned} \hat{\mathbf{F}}_{\alpha}^{\text{eff}}(t) = \hat{\mathbf{F}}_{\alpha}^{\text{B}}(t) + \int_0^t d\tau [\tilde{\mathbf{V}}(t) \chi(t-\tau) \\ + \Gamma(t) \Omega^{-1} \dot{\chi}(t-\tau) + \Omega^{-1} \ddot{\chi}(t-\tau)] \hat{\mathbf{F}}_{\alpha}^{\text{B}}(\tau), \end{aligned} \quad (15)$$

where

$$\Omega \Gamma \Omega^{-1} = \ddot{\chi}(\dot{\chi} \chi^{-1} \dot{\chi} - \ddot{\chi})^{-1} - \ddot{\chi}(\dot{\chi} - \ddot{\chi} \chi^{-1} \dot{\chi})^{-1}, \quad (16a)$$

$$\Omega \tilde{\mathbf{V}} = \ddot{\chi}(\chi \dot{\chi}^{-1} \ddot{\chi} - \dot{\chi})^{-1} - \ddot{\chi}(\chi - \dot{\chi} \dot{\chi}^{-1} \dot{\chi})^{-1}. \quad (16b)$$

They are obtained from Eq. (14) by using the time derivative of Eq. (12) and

$$\mathbf{T}^{-1}(t) = \begin{bmatrix} \Omega(\dot{\chi} - \chi \dot{\chi}^{-1} \ddot{\chi})^{-1} & \Omega(\ddot{\chi} - \dot{\chi} \chi^{-1} \dot{\chi})^{-1} \Omega \\ (\chi - \dot{\chi} \dot{\chi}^{-1} \dot{\chi})^{-1} & (\dot{\chi} - \ddot{\chi} \chi^{-1} \dot{\chi})^{-1} \Omega \end{bmatrix}. \quad (17)$$

Note that all the involved elements in $\chi(t)$ and the related $\mathbf{\Gamma}(t)$ and $\tilde{\mathbf{\Gamma}}(t)$ are real.

For the investigation on the heat transport, we shall identify the contribution of each hybridized α -bath to the key functions $\tilde{\mathbf{\Gamma}}(t)$ and $\mathbf{\Gamma}(t)$ in terms of

$$\tilde{\mathbf{\Gamma}}(t) = \sum_{\alpha} \tilde{\mathbf{\Gamma}}_{\alpha}(t) \quad \text{and} \quad \mathbf{\Gamma}(t) = \sum_{\alpha} \mathbf{\Gamma}_{\alpha}(t). \quad (18)$$

Here, $\tilde{\mathbf{\Gamma}}_{\alpha}(t) \equiv \{\tilde{\Gamma}_{\alpha uv}(t)\}$ and $\mathbf{\Gamma}_{\alpha}(t) \equiv \{\Gamma_{\alpha uv}(t)\}$ [cf. Eq. (8)]. They can be attained by

$$[\tilde{\mathbf{\Gamma}}_{\alpha}(t) \quad \mathbf{\Gamma}_{\alpha}(t)] = - \int_0^t d\tau [\phi_{\alpha}(t-\tau) \quad \mathbf{0}] \mathbf{T}(\tau) \mathbf{T}^{-1}(t). \quad (19)$$

Here $\phi_{\alpha}(t) \equiv \{\phi_{\alpha uv}(t)\}$ with

$$\phi_{\alpha uv}(t) \equiv i \langle [\hat{F}_{\alpha u}^{\text{B}}(t), \hat{F}_{\alpha v}^{\text{B}}(0)] \rangle_{\text{B}} \quad (20)$$

and $\hat{F}_{\alpha u}^{\text{B}}(t) \equiv e^{ih_{\alpha}t} \hat{F}_{\alpha u} e^{-ih_{\alpha}t}$. The average is defined as $\langle \dots \rangle_{\text{B}} \equiv \text{tr}_{\text{B}}[\dots \rho_{\text{B}}^{\text{cano}}]$ with $\rho_{\text{B}}^{\text{cano}} = \otimes_{\alpha} \rho_{\alpha}^{\text{B}}(\beta_{\alpha})$ and $\rho_{\alpha}^{\text{B}}(\beta_{\alpha}) = e^{-\beta_{\alpha} h_{\alpha}} / \text{tr}_{\text{B}}(e^{-\beta_{\alpha} h_{\alpha}})$ over the canonical ensembles of baths. We shall also introduce the hybrid bath correlation function

$$c_{\alpha uv}(t) \equiv \langle \hat{F}_{\alpha u}^{\text{B}}(t) \hat{F}_{\alpha v}^{\text{B}}(0) \rangle_{\text{B}}. \quad (21)$$

With the symmetry property $c_{\alpha uv}^*(t) = c_{\alpha vu}(-t)$, we have $\phi_{\alpha uv}(t) = i[c_{\alpha uv}(t) - c_{\alpha uv}^*(t)] = -2\text{Im}c_{\alpha uv}(t) = -\phi_{\alpha vu}(-t)$. Denote also $r_{\alpha uv}(t) \equiv \text{Re}[c_{\alpha uv}(t)] = r_{\alpha vu}(-t)$ for later use. Note that the cross correlations between different baths do not exist. More details for obtaining Eq. (19) are given in Appendix.

Turn now to $\tilde{\zeta}_{\alpha}(t) \equiv \{\tilde{\zeta}_{\alpha uv}(t)\}$ and $\zeta_{\alpha}(t) \equiv \{\zeta_{\alpha uv}(t)\}$ in Eq. (8). Their evaluations can be summarized as

$$\begin{aligned} & [\zeta_{\alpha}(t) \quad \tilde{\zeta}_{\alpha}(t)] \\ &= [\tilde{\mathbf{V}}(t) \quad \mathbf{\Gamma}(t) \Omega^{-1} \quad \Omega^{-1}] \begin{bmatrix} \mathbf{k}_{\alpha}^{00}(t) & \mathbf{k}_{\alpha}^{01}(t) \Omega^{-1} \\ \mathbf{k}_{\alpha}^{10}(t) & \mathbf{k}_{\alpha}^{11}(t) \Omega^{-1} \\ \mathbf{k}_{\alpha}^{20}(t) & \mathbf{k}_{\alpha}^{21}(t) \Omega^{-1} \end{bmatrix} \end{aligned} \quad (22)$$

where (with T denoting the transpose)

$$\mathbf{k}_{\alpha}^{ij}(t) = \int_0^t d\tau [\mathcal{D}_i \mathbf{u}_{\alpha}(\tau) \mathcal{D}_j \chi^T(\tau) + \mathcal{D}_i \chi(\tau) \mathcal{D}_j \mathbf{u}_{\alpha}^T(\tau)], \quad (23)$$

with

$$\mathbf{u}_{\alpha}(t) \equiv \int_0^t d\tau \chi(\tau) \mathbf{r}_{\alpha}(t-\tau), \quad (24)$$

and the notation \mathcal{D}_m for the derivative $d^m/d\tau^m$. Detailed derivations are given in Appendix. For the condition of a single oscillator coupled to a single bath, Eq. (22) will recover the results in the APPENDIX C of Ref. 43, where the expressions had been reorganized via integral by parts. The asymptotic behaviors had been found there associated with the thermal phase space variances.

C. Entangled system–bath properties and heat transport

Consider the heat transport from the α -reservoir to the local impurity system. The heat current operator reads

$$\begin{aligned} \hat{J}_{\alpha} &\equiv -\dot{h}_{\alpha} = -i[H_{\text{T}}, h_{\alpha}] = -i \sum_u \hat{Q}_u [\hat{F}_{\alpha u}, h_{\alpha}] \\ &= -i \sum_u \hat{Q}_u [\hat{F}_{\alpha u}, H_{\text{T}}] = \sum_u \hat{Q}_u \dot{\hat{F}}_{\alpha u}. \end{aligned} \quad (25)$$

Denote $\langle \hat{O} \rangle_t = \text{Tr}[\hat{O} \rho_{\text{T}}(t)]$. We can evaluate the heat current via

$$\begin{aligned} \langle \hat{J}_{\alpha}(t) \rangle_t &= -i \sum_u \langle \hat{Q}_u [\hat{F}_{\alpha u}, H_{\text{T}}] \rangle_t \\ &= -i \sum_u \text{Tr}\{[H_{\text{T}}, \rho_{\text{T}}(t)] \hat{Q}_u \hat{F}_{\alpha u}\} \\ &\quad - i \sum_u \text{Tr}\{\rho_{\text{T}}(t) [H_{\text{T}}, \hat{Q}_u] \hat{F}_{\alpha u}\} \\ &= \sum_u [\dot{\tilde{Q}}_{\alpha u}(t) - \Omega_u \tilde{P}_{\alpha u}(t)], \end{aligned} \quad (26)$$

where [cf. Eqs. (8) and (A.13)]

$$\begin{aligned} \tilde{Q}_{\alpha u} &\equiv \text{tr}_{\text{S}}[\hat{Q}_u \tilde{\rho}_{\alpha u}(t)] \\ &= -\zeta_{\alpha uu}(t) + \sum_v \tilde{\Gamma}_{\alpha uv}(t) [W_{uv}^{QQ}(t) + Q_u(t) Q_v(t)] \\ &\quad + \sum_v \Gamma_{\alpha uv}(t) [W_{uv}^{QP}(t) + Q_u(t) P_v(t)], \end{aligned} \quad (27a)$$

and

$$\begin{aligned} \tilde{P}_{\alpha u} &\equiv \text{tr}_{\text{S}}[\hat{P}_u \tilde{\rho}_{\alpha u}(t)] \\ &= -\tilde{\zeta}_{\alpha uu}(t) + \sum_v \tilde{\Gamma}_{\alpha uv}(t) [W_{uv}^{PQ}(t) + P_u(t) Q_v(t)] \\ &\quad + \sum_v \Gamma_{\alpha uv}(t) [W_{uv}^{PP}(t) + P_u(t) P_v(t)]. \end{aligned} \quad (27b)$$

Involved in Eqs. (27a) and (27b) are also the first order moments, $\{Q_u(t)\}$ and $\{P_u(t)\}$, and the second order moments, $\{W_{uv}^{QQ}\}$, $\{W_{uv}^{QP}\}$, $\{W_{uv}^{PQ}\}$, and $\{W_{uv}^{PP}\}$. Their definitions are given in Eq. (A.11) and Eq. (A.13), respectively. Thus Eq. (26) can be recast in terms of

$$\langle \hat{J}_{\alpha}(t) \rangle_t = \text{trace}[\hat{\mathbf{J}}_{\alpha;Q}(t) - \Omega \mathbf{J}_{\alpha;P}(t)] \quad (28)$$

with

$$\begin{aligned} \mathbf{J}_{\alpha;Q}(t) &= \tilde{\mathbf{\Gamma}}_{\alpha}(t) [\mathbf{W}^{QQ}(t) + \mathbf{Q}^T(t) \mathbf{Q}(t)] - \zeta_{\alpha}(t) \\ &\quad + \mathbf{\Gamma}_{\alpha}(t) [\mathbf{W}^{PQ}(t) + \mathbf{P}^T(t) \mathbf{Q}(t)], \end{aligned} \quad (29a)$$

$$\begin{aligned} \mathbf{J}_{\alpha;P}(t) &= \tilde{\mathbf{\Gamma}}_{\alpha}(t) [\mathbf{W}^{QP}(t) + \mathbf{Q}^T(t) \mathbf{P}(t)] - \tilde{\zeta}_{\alpha}(t) \\ &\quad + \mathbf{\Gamma}_{\alpha}(t) [\mathbf{W}^{PP}(t) + \mathbf{P}^T(t) \mathbf{P}(t)]. \end{aligned} \quad (29b)$$

The steady-state heat current is obtained by $t \rightarrow \infty$.

An alternative approach to evaluate the steady-state heat current is via

$$J_{\alpha} \equiv \langle \hat{J}_{\alpha}(t) \rangle_t|_{t \rightarrow \infty} = -2 \text{Im} \int_0^{\infty} d\tau \text{tr}[\dot{c}_{\alpha}(\tau) \mathbf{C}(\tau)]. \quad (30)$$

Here, $\mathbf{C}(t) \equiv \{C_{uv}(t)\}$ and

$$C_{uv}(t) \equiv \langle \hat{Q}_u(t) \hat{Q}_v(0) \rangle. \quad (31)$$

This is the time-domain Meir-Wingreen's formula⁴⁹ for general systems. This formula can be derived via such as the nonequilibrium Green's function method⁵⁰ or the dissipaton thermofield approach.²³ In the frequency domain, it can be recast as

$$J_\alpha = \frac{2}{\pi} \sum_{uv} \int_{-\infty}^{\infty} d\omega \frac{\omega \Phi_{\alpha uv}(\omega)}{1 - e^{-\beta \alpha \omega}} \text{Re}[\tilde{C}_{vu}(-\omega)], \quad (32)$$

with the hybrid bath spectral density

$$\Phi_{\alpha uv}(\omega) \equiv \frac{1}{2i} \int_{-\infty}^{\infty} dt e^{i\omega t} \phi_{\alpha uv}(t), \quad (33)$$

and the system correlation resolution

$$\tilde{C}_{uv}(\omega) \equiv \int_0^{\infty} dt e^{i\omega t} C_{uv}(t). \quad (34)$$

The hybrid bath correlation function is related to its spectral density via the fluctuation-dissipation theorem^{32,33}

$$c_{\alpha uv}(t) = \frac{1}{\pi} \int_{-\infty}^{\infty} d\omega \frac{e^{-i\omega t} \Phi_{\alpha uv}(\omega)}{1 - e^{-\beta \alpha \omega}}. \quad (35)$$

To evaluate Eq. (32) for the present multimode BO system, we propose a discrete imaginary-frequency (DIF) method. This method is based on three ingredients as follows. (i) The exponential decomposition of the hybrid bath correlation functions (with $\gamma_{\alpha k}$ assumed real),

$$c_{\alpha uv}(t) = \sum_k g_{\alpha kuv} e^{-\gamma_{\alpha k} t}. \quad (36)$$

This can be readily achieved with some sum-over-pole schemes^{51–56} or the time-domain Prony fitting decomposition scheme.⁵⁷ (ii) The relation of entangled system-bath correlations [cf. Eq. (A.6)]

$$\begin{aligned} \ddot{C}_{uv}(t) = & -\Omega_u \sum_{u'} V_{uu'} C_{u'v}(t) - \Omega_u \sum_{\alpha} X_{\alpha uv}(t) \\ & + \Omega_u \sum_{\alpha u'} \int_0^t d\tau \phi_{\alpha uu'}(t - \tau) C_{u'v}(\tau), \end{aligned} \quad (37)$$

where $X_{\alpha uv}(t) \equiv \langle \hat{F}_{\alpha u}^B(t) \hat{Q}_v(0) \rangle$. In terms of the Laplace transform $\tilde{f}(\omega) \equiv \int_0^{\infty} dt e^{i\omega t} f(t)$, it reads

$$\begin{aligned} \tilde{C}(\omega) = & \left[\Omega \mathbf{V} - \omega^2 \mathbf{I} - \Omega \sum_{\alpha} \tilde{\phi}_{\alpha}(\omega) \right]^{-1} \left[\dot{C}(0) - i\omega C(0) \right. \\ & \left. - \Omega \sum_{\alpha} \tilde{\mathbf{X}}_{\alpha}(\omega) \right]. \end{aligned} \quad (38)$$

(iii) The system-bath entanglement theorem [cf. the Eq.(17) of Ref. 23]

$$\mathbf{X}_{\alpha}(t) = 2 \text{Im} \int_0^{\infty} d\tau \mathbf{c}_{\alpha}^T(t + \tau) \mathbf{C}^T(\tau). \quad (39)$$

Here, (ii) is from the BO algebra while (iii) exists for general systems.

Substituting Eq. (36) into Eq. (39), we obtain the following frequency-domain expression

$$\tilde{\mathbf{X}}_{\alpha}(\omega) = \sum_k \frac{2}{\gamma_{\alpha k} - i\omega} \text{Im} [\tilde{\mathbf{C}}(i\gamma_{\alpha k}) \mathbf{g}_{\alpha k}]^T. \quad (40)$$

We find that once $\{\tilde{\mathbf{C}}(i\gamma_{\alpha k})\}$ is known, $\tilde{\mathbf{X}}_{\alpha}(\omega)$ [Eq. (40)] and hence $\tilde{\mathbf{C}}(\omega)$ [Eq. (38)] will be readily obtained. On the other hand, $\{\tilde{\mathbf{C}}(i\gamma_{\alpha k})\}$ can be solved by Eqs. (38) and (40) in a self-consistent manner, i.e.

$$\begin{aligned} \tilde{\mathbf{C}}(i\gamma_{\alpha' k'}) = & \left[\Omega \mathbf{V} + \gamma_{\alpha' k'}^2 \mathbf{I} - \Omega \sum_{\alpha} \tilde{\phi}_{\alpha}(i\gamma_{\alpha' k'}) \right]^{-1} \left[\dot{\mathbf{C}}(0) \right. \\ & \left. + \gamma_{\alpha' k'} \mathbf{C}(0) - \Omega \sum_{\alpha} \tilde{\mathbf{X}}_{\alpha}(i\gamma_{\alpha' k'}) \right], \end{aligned} \quad (41a)$$

$$\tilde{\mathbf{X}}_{\alpha}(i\gamma_{\alpha' k'}) = \sum_k \frac{2}{\gamma_{\alpha k} + \gamma_{\alpha' k'}} \text{Im} [\tilde{\mathbf{C}}(i\gamma_{\alpha k}) \mathbf{g}_{\alpha k}]^T. \quad (41b)$$

Note that for the multimode BO system in this paper, $\phi_{\alpha uv} = \phi_{\alpha vu}$. The steady-state heat current can then be obtained from Eq. (32) via

$$J_{\alpha} = -\frac{2}{\pi} \sum_{uv} \int_{-\infty}^{\infty} d\omega \frac{\omega \Phi_{\alpha uv}(\omega)}{e^{\beta \alpha \omega} - 1} \text{Re}[\tilde{C}_{vu}(\omega)]. \quad (42)$$

By far, we have proposed two different methods to evaluate the steady-state heat current of the multimode BO system. One is the algebraic method with the aid of the EOM of the density matrix. The other is the DIF method followed by the Meir-Wingreen's formula. These two exact methods are numerically consistent in our calculations. As tested, the simulation results are also found consistent with those via the dissipaton theory.^{23,58} Note that there is $\sum_{\alpha} J_{\alpha} = 0$ in the steady state, which is also confirmed in our calculations.




FIG. 1. Schematic diagram of the model for numerical demonstrations.

III. NUMERICAL DEMONSTRATIONS

For numerical demonstrations, we adopt [cf. Eq. (33)]

$$\Phi_\alpha(\omega) = \eta_\alpha \text{Im} \frac{\Omega_B^2}{\Omega_B^2 - \omega^2 - i\omega\gamma_B}, \quad (43)$$

with $\eta_\alpha \equiv \{\eta_{\alpha uv}; \eta_{\alpha uv} = \eta_{\alpha u} \delta_{uv}\}$ specifying the system-bath coupling strengths. Exemplified are the calculations on transient dynamics and heat currents. Selected is a two-mode BO system embedded in two (left and right with $\alpha=L$ and R respectively) reservoirs at different temperatures. The schematic diagram of the model is shown in Fig. 1. From an application point of view, this setup can be experimentally realized via, for example, molecular junctions.^{59–62} The system-bath coupling strength $\{\eta_{\alpha u}\}$ can be experimentally adjusted via manipulating the distances between molecules. We select the parameters $\Omega_1 = \Omega_2 = V_{11} = V_{22} = \Omega_B = 1600 \text{ cm}^{-1}$ and $\gamma_B = 4\Omega_B$. Other parameters are specified in the caption of each figure.




FIG. 2. Evaluations on the transient two-mode BO dynamics in terms of the energies of the system oscillators. See text for details. Parameters are selected as $V_{12} = V_{21} = 800 \text{ cm}^{-1}$, $\eta_{L1} = \eta_{R2} = 320 \text{ cm}^{-1}$, $\eta_{L2} = \eta_{R1} = 160 \text{ cm}^{-1}$, $T_L = 50 \text{ K}$ and $T_R = 500 \text{ K}$.

Figure 2 depicts the transient dynamics of the system energies, $E_S = \langle H_S \rangle$, $E_{i=1,2} = (\Omega_i/2)\langle \hat{P}_i^2 \rangle + (V_{ii}/2)\langle \hat{Q}_i^2 \rangle$, and $E_\pm = \langle \hat{p}_\pm^2 \rangle/2 + (\omega_\pm^2/2)\langle \hat{q}_\pm^2 \rangle$ of the normal modes. We select $V_{12} = V_{21} = 800 \text{ cm}^{-1}$ resulting in $\omega_+ = 1960 \text{ cm}^{-1}$ and $\omega_- = 1130 \text{ cm}^{-1}$. The ω_+ plays the dominate role in realistic observations. The system is initially at the factorized state, $\rho_s(0) \otimes \rho_B^{\text{cano}}$, with $\rho_s(0) = \otimes_{i=1,2} \rho_i$ where ρ_i is the ground state of the i th isolated oscillator but with a dimensionless position displacement of $d = -2$. To be specific, $\rho_i = |\psi^{(i)}\rangle\langle\psi^{(i)}|$, with

$$\psi^{(i)}(Q_i) = \left(\frac{\sqrt{V_{ii}/\Omega_i}}{\pi} \right)^{1/4} \exp \left[-\frac{\sqrt{V_{ii}/\Omega_i}}{2} (Q_i + 2)^2 \right].$$

Apparently, $E_S = E_+ + E_- \neq E_1 + E_2$.




FIG. 3. The $\tilde{C}(\omega)$ evaluated with $V_{12} = V_{21} = 160 \text{ cm}^{-1}$ (black) and 800 cm^{-1} (red) at $T_L = T_R = 275 \text{ K}$. Other parameters are the same as in Fig. 2.

The two panels of Fig. 3 show the spectra and dispersions, i.e. the real and imaginary parts of $\tilde{C}(\omega)$, respectively, evaluated via the DIF method proposed in Sec. II C. Obviously, the splitting of peaks becomes larger when $V_{12} = V_{21} = 800 \text{ cm}^{-1}$, compared to the case of $V_{12} = V_{21} = 160 \text{ cm}^{-1}$. With the equal temperature of both reservoirs, the detailed-balance relation has been satisfied in our evaluations.

Shown in Fig. 4 is the steady-state heat current with the temperature of the right bath, T_R , varying from 25 K to 475 K while that of the left bath, T_L , fixed at 275 K . Apparently, the current increases with the inter-mode coupling strength, V_{12} . Also the current changes direction at the condition $T_R = T_L$. The differential conductance is found of similar behavior which is thus not shown explicitly. From Fig. 4, one can observe that there is an asymmetry in magnitude of current between two scenarios of $T_R < T_L$ and $T_R > T_L$ with the fixed temperature difference. The mechanism can be interpreted as follows. The heat current increases with not only the temperature difference but also the heat capacity. The phonon frequency involved in our simulation is around 1600 cm^{-1} .




FIG. 4. Heat current versus temperature, with $V_{12} = V_{21} = 160\text{cm}^{-1}$ (black) and 800cm^{-1} (red). The temperature of the right bath, T_R , varies from 25K to 475K while that of the left bath, T_L , is fixed at 275K. Other parameters are the same as in Fig. 2. The crosses marked along the curves are results from the DIF method.

The corresponding Einstein temperature is over 2000K. For the temperature range in Fig. 4, the heat capacity increases with the temperature, leading to the observed asymmetry.

IV. SUMMARY

To conclude, we develop an algebraic method to study multimode Brownian oscillators connected to multiple reservoirs at different temperatures. The algebraic approach directly gives the exact time-local EOM for the reduced density operator. Based on this approach, we can easily extract not only the reduced system but also the hybrid bath dynamical informations. We exploit the EOM to study the heat transport problem, i.e. the heat current from the reservoir to the local impurity system. On the other hand, we also adopt another different approach, the Meir-Wingreen's formula which is for general systems, together with the system-bath entanglement theorem to evaluate the steady-state heat current. For its application to the BO system, we propose a discrete imaginary-frequency (DIF) self-consistent evaluation scheme. Both approaches can serve as initial steps to systematically include anharmonic effects. For more generic (non-integrable) bosonic setups, the self-energy term needs to be modified,⁶³ which can be obtained via self-consistent iteration approximately. Work in this direction is in progress. It is anticipated that the methods presented in this work would constitute basic components for nonequilibrium statistical mechanics of open quantum systems. Moreover, it is still difficult to compute the fluctuation of heat current under the present theoretical framework of Eqs.(7)–(9). However, it may be achieved with the help of dissipaton theory.^{23,58} This constitutes

another direction of further development.

ACKNOWLEDGMENTS

Support from the Ministry of Science and Technology of China (Grant No. 2021YFA1200103) and the National Natural Science Foundation of China (Grant Nos. 22103073 and 22173088) is gratefully acknowledged.

Appendix: Related BO algebra and BO properties

The EOM (7)–(9) can be established by the Yan-Mukamel method.^{17,43,48} Here we present another approach via undetermined coefficients. Both methods are on the basis of the EOM of the first and second order moments, which fully characterize the Gaussian wave packet (GWP). Let us start from the Heisenberg EOM for any operator, $\dot{\hat{O}}(t) = i[H_T, \hat{O}(t)]$. We have

$$\dot{\hat{Q}}_u(t) = \Omega_u \hat{P}_u(t), \quad (\text{A.1a})$$

$$\dot{\hat{P}}_u(t) = -\sum_v V_{uv} \hat{Q}_v(t) - \sum_\alpha \hat{F}_{\alpha u}(t). \quad (\text{A.1b})$$

The solution to $\hat{F}_{\alpha u}(t) \equiv e^{iH_T t} \hat{F}_{\alpha u} e^{-iH_T t}$ can be obtained as [cf. the Eq. (7) of Ref. 63]

$$\hat{F}_{\alpha u}(t) = \hat{F}_{\alpha u}^B(t) - \sum_v \int_0^t d\tau \phi_{\alpha uv}(t - \tau) \hat{Q}_v(\tau). \quad (\text{A.2})$$

From Eq. (A.1) we obtain

$$\dot{\chi}_{uv}^{QQ}(t) = \Omega_u \chi_{uv}^{PQ}(t), \quad (\text{A.3a})$$

$$\dot{\chi}_{uv}^{QP}(t) = \Omega_u \chi_{uv}^{PP}(t), \quad (\text{A.3b})$$

and

$$\dot{\chi}_{uv}^{PQ}(t) = -\sum_{v'} V_{uv'} \chi_{v'v}^{QQ}(t) - i \sum_\alpha \langle [\hat{F}_{\alpha u}(t), \hat{Q}_v(0)] \rangle, \quad (\text{A.4a})$$

$$\dot{\chi}_{uv}^{PP}(t) = -\sum_{v'} V_{uv'} \chi_{v'v}^{QP}(t) - i \sum_\alpha \langle [\hat{F}_{\alpha u}(t), \hat{P}_v(0)] \rangle. \quad (\text{A.4b})$$

Substituting Eq. (A.2) into Eq. (A.4), we get

$$\begin{aligned} \dot{\chi}_{uv}^{PQ}(t) = & -\sum_{v'} V_{uv'} \chi_{v'v}^{QQ}(t) \\ & + \sum_{\alpha v'} \int_0^t d\tau \phi_{\alpha uv'}(t - \tau) \chi_{v'v}^{QQ}(\tau), \end{aligned} \quad (\text{A.5a})$$

$$\begin{aligned} \dot{\chi}_{uv}^{PP}(t) = & -\sum_{v'} V_{uv'} \chi_{v'v}^{QP}(t) \\ & + \sum_{\alpha v'} \int_0^t d\tau \phi_{\alpha uv'}(t - \tau) \chi_{v'v}^{QP}(\tau). \end{aligned} \quad (\text{A.5b})$$

The traditional Langevin equation can be recovered by substituting Eq. (A.2) into Eq. (A.1) as

$$\ddot{\hat{Q}}_u(t) = -\Omega_u \sum_v V_{uv} \hat{Q}_v(t) - \Omega_u \sum_\alpha \hat{F}_{\alpha u}^B(t) + \Omega_u \sum_{\alpha v} \int_0^t d\tau \phi_{\alpha uv}(t-\tau) \hat{Q}_v(\tau). \quad (\text{A.6})$$

The solutions to it, i.e. Eq. (11) with Eq. (12), can be proved by Eqs. (A.3) and (A.5) together with initial values $\chi(0) = \ddot{\chi}(0) = \mathbf{0}$ and $\dot{\chi}(0) = \mathbf{\Omega}$. Furthermore, Eqs. (A.3) and (A.5) can be resolved in terms of the Laplace transform $\tilde{f}(\omega) \equiv \int_0^\infty dt e^{i\omega t} f(t)$ as

$$\tilde{\chi}_{QQ}(\omega) = [\mathbf{\Omega V} - \omega^2 \mathbf{I} - \mathbf{\Omega} \sum_\alpha \tilde{\phi}_\alpha(\omega)]^{-1} \mathbf{\Omega}, \quad (\text{A.7a})$$

$$\tilde{\chi}_{QP}(\omega) = i\omega \tilde{\chi}_{QQ}(\omega) \mathbf{\Omega}^{-1}, \quad (\text{A.7b})$$

$$\tilde{\chi}_{PQ}(\omega) = -i\omega \mathbf{\Omega}^{-1} \tilde{\chi}_{QQ}(\omega), \quad (\text{A.7c})$$

$$\tilde{\chi}_{PP}(\omega) = \mathbf{\Omega}^{-1} + \omega^2 \mathbf{\Omega}^{-1} \tilde{\chi}_{QQ}(\omega) \mathbf{\Omega}^{-1}. \quad (\text{A.7d})$$

Equations (A.3) and (A.5) can be recast into the matrix form as

$$\dot{\mathbf{T}}(t) = \begin{bmatrix} \mathbf{0} & \mathbf{\Omega} \\ -\mathbf{V} & \mathbf{0} \end{bmatrix} \mathbf{T}(t) + \sum_\alpha \int_0^t d\tau \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \phi_\alpha(t-\tau) & \mathbf{0} \end{bmatrix} \mathbf{T}(\tau). \quad (\text{A.8})$$

By the first identity of Eq. (14), we can recast the $\mathbf{\Lambda}(t)$ matrix using Eq. (A.8) as

$$\mathbf{\Lambda}(t) = \begin{bmatrix} \mathbf{0} & \mathbf{\Omega} \\ -\mathbf{V} & \mathbf{0} \end{bmatrix} + \sum_\alpha \int_0^t d\tau \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \phi_\alpha(t-\tau) & \mathbf{0} \end{bmatrix} \mathbf{T}(\tau) \mathbf{T}^{-1}(t). \quad (\text{A.9})$$

Equation (19) is resulted from the comparison of Eq. (A.9) to the second identity of Eq. (14) with Eq. (18).

Consider the initial factorized state before the system and baths interact, i.e.

$$\rho_{\text{T}}(0) = \rho_{\text{S}}(0) \otimes \prod_\alpha \rho_\alpha^{\text{B}}(\beta_\alpha). \quad (\text{A.10})$$

We have denoted the average $\langle (\cdot) \rangle_t$ over the total composite state with the above initial total GWP. That is in the Schrödinger picture, $\langle \hat{O} \rangle_t = \text{Tr}[\hat{O} \rho_{\text{T}}(t)]$, which is equal to $\langle \hat{O} \rangle_t = \text{Tr}[\hat{O}(t) \rho_{\text{T}}(0)]$ in the Heisenberg picture. From Eq. (13), the EOM of the first order moments,

$$Q_u(t) \equiv \langle \hat{Q}_u(t) \rangle_t \text{ and } P_u(t) \equiv \langle \hat{P}_u(t) \rangle_t, \quad (\text{A.11})$$

can be obtained as

$$\begin{bmatrix} \dot{Q}(t) \\ \dot{P}(t) \end{bmatrix} = \mathbf{\Lambda}(t) \begin{bmatrix} Q(t) \\ P(t) \end{bmatrix}. \quad (\text{A.12})$$

Note that $\langle \hat{\mathbf{F}}_\alpha^{\text{eff}}(t) \rangle_t = \langle \hat{\mathbf{F}}_\alpha^{\text{B}}(t) \rangle_t = \mathbf{0}$. Actually for any system operator \hat{A} , there is $\langle \hat{\mathbf{F}}_\alpha^{\text{eff}}(t) \hat{A}(0) \rangle_t = \langle \hat{\mathbf{F}}_\alpha^{\text{B}}(t) \hat{A}(0) \rangle_t = \mathbf{0}$.

The EOM of the second order moments,

$$W_{uv}^{QQ}(t) \equiv \frac{1}{2} \langle \{ \delta \hat{Q}_u(t), \delta \hat{Q}_v(t) \} \rangle_t, \quad (\text{A.13a})$$

$$W_{uv}^{QP}(t) \equiv \frac{1}{2} \langle \{ \delta \hat{Q}_u(t), \delta \hat{P}_v(t) \} \rangle_t = W_{vu}^{PQ}(t), \quad (\text{A.13b})$$

$$W_{uv}^{PP}(t) \equiv \frac{1}{2} \langle \{ \delta \hat{P}_u(t), \delta \hat{P}_v(t) \} \rangle_t, \quad (\text{A.13c})$$

with $\delta \hat{Q}_u(t) \equiv \hat{Q}_u(t) - Q_u(t)$ and $\delta \hat{P}_u(t) \equiv \hat{P}_u(t) - P_u(t)$ can also be derived from Eq. (13). For convenience, we recast Eq. (A.13) in the matrix form as

$$\mathbf{W}(t) \equiv \begin{bmatrix} \mathbf{W}^{QQ}(t) & \mathbf{W}^{QP}(t) \\ \mathbf{W}^{PQ}(t) & \mathbf{W}^{PP}(t) \end{bmatrix} = \frac{1}{2} \left\langle \left\{ \begin{bmatrix} \delta \hat{\mathbf{Q}}(t) \\ \delta \hat{\mathbf{P}}(t) \end{bmatrix}, \begin{bmatrix} \delta \hat{\mathbf{Q}}(t) \\ \delta \hat{\mathbf{P}}(t) \end{bmatrix}^T \right\} \right\rangle. \quad (\text{A.14})$$

We obtain from Eq. (13) that

$$\dot{\mathbf{W}}(t) = \mathbf{\Lambda}(t) \mathbf{W}(t) + \mathbf{W}(t) \mathbf{\Lambda}^T(t) + \sum_\alpha \begin{bmatrix} \mathbf{0} & [\zeta_\alpha(t)]^T \\ \zeta_\alpha(t) & \tilde{\zeta}_\alpha(t) + [\tilde{\zeta}_\alpha(t)]^T \end{bmatrix}, \quad (\text{A.15})$$

where

$$\zeta_\alpha(t) = -\text{Re} \langle \hat{\mathbf{F}}_\alpha^{\text{eff}}(t) \hat{\mathbf{Q}}^T(t) \rangle_t, \quad (\text{A.16a})$$

$$\tilde{\zeta}_\alpha(t) = -\text{Re} \langle \hat{\mathbf{F}}_\alpha^{\text{eff}}(t) \hat{\mathbf{P}}^T(t) \rangle_t. \quad (\text{A.16b})$$

The evaluations of $\zeta_\alpha(t)$ and $\tilde{\zeta}_\alpha(t)$ can be done by substituting Eqs. (11) and (15) into Eq. (A.16). Denoting

$$\begin{aligned} \xi_\alpha(\tau; t) &\equiv \mathbf{r}_\alpha(\tau) + \tilde{\mathbf{V}}(t) \int_0^t d\tau' \chi(\tau') \mathbf{r}_\alpha(\tau - \tau') \\ &\quad + \mathbf{\Gamma}(t) \int_0^t d\tau' \mathbf{\Omega}^{-1} \dot{\chi}(\tau') \mathbf{r}_\alpha(\tau - \tau') \\ &\quad + \mathbf{\Omega}^{-1} \int_0^t d\tau' \ddot{\chi}(\tau') \mathbf{r}_\alpha(\tau - \tau'), \end{aligned} \quad (\text{A.17})$$

we have

$$[\zeta_\alpha(t) \quad \tilde{\zeta}_\alpha(t)] = \int_0^t d\tau \xi_\alpha(\tau; t) [\chi^T(\tau) \quad \dot{\chi}^T(\tau) \mathbf{\Omega}^{-1}]. \quad (\text{A.18})$$

Equation (22) can now be obtained by noticing the following relation of double integrals

$$\begin{aligned} &\int_0^t d\tau \int_0^t d\tau' f(\tau') g(\tau - \tau') h(\tau) \\ &= \int_0^t d\tau \int_0^\tau d\tau' [f(\tau') g(\tau - \tau') h(\tau) \\ &\quad + f(\tau) g(\tau' - \tau) h(\tau')]. \end{aligned} \quad (\text{A.19})$$

Note in Eqs. (22)–(24)

$$\dot{\mathbf{u}}_\alpha(t) = \int_0^t d\tau \dot{\chi}(\tau) \mathbf{r}_\alpha(t - \tau), \quad (\text{A.20})$$

$$\ddot{\mathbf{u}}_\alpha(t) = \mathbf{\Omega} \mathbf{r}_\alpha(t) + \int_0^t d\tau \ddot{\chi}(\tau) \mathbf{r}_\alpha(t - \tau). \quad (\text{A.21})$$

Turn now to the EOM of the reduced density operator $\rho_s(t)$. Concerning the Gaussian property and the Liouville-von Neumann equation for the total density operator, $\dot{\rho}_T(t) = -i[H_T, \rho_T(t)]$, the desired EOM is of the form

$$\dot{\rho}_s(t) = -i[H_s, \rho_s(t)] + \sum_{\alpha uv} \hat{Q}_u^\ominus [a_{\alpha uv}^\oplus(t) \hat{Q}_v^\oplus + b_{\alpha uv}^\oplus(t) \hat{P}_v^\oplus + a_{\alpha uv}^\ominus(t) \hat{Q}_v^\ominus + b_{\alpha uv}^\ominus(t) \hat{P}_v^\ominus] \rho_s(t), \quad (\text{A.22})$$

with the coefficients $\{a_{\alpha uv}^\oplus\}$, $\{b_{\alpha uv}^\oplus\}$, $\{a_{\alpha uv}^\ominus\}$, and $\{b_{\alpha uv}^\ominus\}$ to be determined. To do that, consider an arbitrary system operator \hat{A} . We can obtain the EOM of $\langle \hat{A} \rangle_t = \text{Tr}_s[\hat{A} \rho_s(t)]$ from Eq. (A.22) as

$$\frac{d}{dt} \langle \hat{A} \rangle_t = i \langle [H_s, \hat{A}] \rangle_t - \sum_{\alpha uv} \langle [a_{\alpha uv}^\oplus(t) \hat{Q}_v^\oplus + b_{\alpha uv}^\oplus(t) \hat{P}_v^\oplus - a_{\alpha uv}^\ominus(t) \hat{Q}_v^\ominus - b_{\alpha uv}^\ominus(t) \hat{P}_v^\ominus] \hat{Q}_u^\ominus \hat{A} \rangle_t. \quad (\text{A.23})$$

We can now derive the EOM of the first and second order moments from Eq. (A.23) and compare the results to Eqs. (A.12) and (A.15). After some simple steps, the coefficients $\{a_{\alpha uv}^\oplus\}$, $\{b_{\alpha uv}^\oplus\}$, $\{a_{\alpha uv}^\ominus\}$, and $\{b_{\alpha uv}^\ominus\}$ can be determined as $\{\Gamma_{\alpha uv}(t)\}$, $\{\Gamma_{\alpha uv}(t)\}$, $\{\tilde{\zeta}_{\alpha uv}(t)\}$, and $-\{\zeta_{\alpha uv}(t)\}$, respectively. Thus the final EOM of $\rho_s(t)$ is obtained as Eqs. (7)–(9). The effect of the initial state [Eq. (A.10)] vanishes when $t \rightarrow \infty$. The asymptotic behaviors of the coefficients in Eq. (8) can refer to Ref. 43.

- ¹P. Reineker, *Exciton Dynamics in Molecular Crystals and Aggregates: Stochastic Liouville Equation Approach: Coupled Coherent and Incoherent Motion, Optical Line Shapes, Magnetic Resonance Phenomena*, Springer, Berlin, 1982.
- ²C. P. Slichter, *Principles of Magnetic Resonance*, Springer Verlag, New York, 1990.
- ³L. M. K. Vandersypen and I. L. Chuang, “NMR techniques for quantum control and computation,” *Rev. Mod. Phys.* **76**, 1037 (2005).
- ⁴M. Born and K. Huang, *Dynamical Theory of Crystal Lattices*, Oxford University Press, New York, 1985.
- ⁵C. F. Klingshirm, *Semiconductor Optics*, Springer-Verlag, Heidelberg, 1997.
- ⁶J. Rammer, *Quantum Transport Theory*, Perseus Books, Reading, Mass., 1998.
- ⁷Y. Akamatsu, “Heavy quark master equations in the Lindblad form at high temperatures,” *Phys. Rev. D* **91**, 056002 (2015).
- ⁸J.-P. Blaizot and M. A. Escobedo, “Quantum and classical dynamics of heavy quarks in a quark-gluon plasma,” *J. High Energy Phys.* **2018**, 1 (2018).
- ⁹T. Miura, Y. Akamatsu, M. Asakawa, and A. Rothkopf, “Quantum Brownian motion of a heavy quark pair in the quark-gluon plasma,” *Phys. Rev. D* **101**, 034011 (2020).
- ¹⁰H. Haken, *Laser Theory*, Springer, Berlin, 1970.
- ¹¹W. H. Louisell, *Quantum Statistical Properties of Radiation*, Wiley, New York, 1973.
- ¹²M. O. Scully and M. S. Zubairy, *Quantum Optics*, Cambridge University Press, Cambridge, 1997.
- ¹³A. Nitzan, *Chemical Dynamics in Condensed Phases: Relaxation, Transfer and Reactions in Condensed Molecular Systems*, Oxford University Press, New York, 2006.
- ¹⁴C. Creatore, M. A. Parker, S. Emmott, and A. W. Chin, “Efficient biologically inspired photocell enhanced by delocalized quantum states,” *Phys. Rev. Lett.* **111**, 253601 (2013).
- ¹⁵J. Fang, Z.-H. Chen, Y. Su, Z.-F. Zhu, Y. Wang, R.-X. Xu, and Y. J. Yan, “Coherent excitation energy transfer in model photosynthetic reaction center: Effects of non-Markovian quantum environment,” *J. Chem. Phys.* **157**, 084119 (2022).
- ¹⁶Y. R. Shen, *The Principles of Nonlinear Optics*, Wiley, New York, 1984.
- ¹⁷Y. J. Yan and S. Mukamel, “Electronic dephasing, vibrational relaxation, and solvent friction in molecular nonlinear optical lineshapes,” *J. Chem. Phys.* **89**, 5160 (1988).
- ¹⁸S. Mukamel, *The Principles of Nonlinear Optical Spectroscopy*, Oxford University Press, New York, 1995.
- ¹⁹A. Kato and Y. Tanimura, “Quantum heat current under non-perturbative and non-Markovian conditions: Applications to heat machines,” *J. Chem. Phys.* **145**, 224105 (2016).
- ²⁰L. Song and Q. Shi, “Hierarchical equations of motion method applied to nonequilibrium heat transport in model molecular junctions: Transient heat current and high-order moments of the current operator,” *Phys. Rev. B* **95**, 064308 (2017).
- ²¹H. Gong, Y. Wang, H. D. Zhang, Q. Qiao, R. X. Xu, X. Zheng, and Y. J. Yan, “Equilibrium and transient thermodynamics: A unified dissipaton-space approach,” *J. Chem. Phys.* **153**, 154111 (2020).
- ²²S. Koyanagi and Y. Tanimura, “The laws of thermodynamics for quantum dissipative systems: A quasi-equilibrium Helmholtz energy approach,” *J. Chem. Phys.* **157**, 014104 (2022).
- ²³Y. Wang, Z. H. Chen, R. X. Xu, X. Zheng, and Y. J. Yan, “A statistical quasi-particles thermofield theory with Gaussian environments: System-bath entanglement theorem for nonequilibrium correlation functions,” *J. Chem. Phys.* **157**, 044012 (2022).
- ²⁴J.-T. Hsiang and B.-L. Hu, “Non-Markovian Abraham-Lorentz-Dirac equation: Radiation reaction without pathology,” *Phys. Rev. D* **106**, 125018 (2022).
- ²⁵J. R. Anglin, J. P. Paz, and W. H. Zurek, “Deconstructing decoherence,” *Phys. Rev. A* **55**, 4041 (1997).
- ²⁶C. W. Gardiner and P. Zoller, *Quantum Noise*, Springer, Wellington, New Zealand, 2004, 3rd ed.
- ²⁷A. A. Clerk, M. H. Devoret, S. M. Girvin, F. Marquardt, and R. J. Schoelkopf, “Introduction to quantum noise, measurement, and amplification,” *Rev. Mod. Phys.* **82**, 1155 (2010).
- ²⁸A. Soare, H. Ball, D. Hayes, J. Sastrawan, M. C. Jarratt, J. J. McLoughlin, X. Zhen, T. J. Green, and M. J. Biercuk, “Experimental noise filtering by quantum control,” *Nat. Phys.* **10**, 825 (2014).
- ²⁹R. Harper, S. T. Flammia, and J. J. Wallman, “Efficient learning of quantum noise,” *Nat. Phys.* **16**, 1184 (2020).
- ³⁰A. O. Caldeira and A. J. Leggett, “Path integral approach to quantum Brownian motion,” *Physica A* **121**, 587 (1983).
- ³¹H. Grabert, P. Schramm, and G. L. Ingold, “Quantum Brownian motion: The functional integral approach,” *Phys. Rep.* **168**, 115 (1988).
- ³²Y. J. Yan and R. X. Xu, “Quantum mechanics of dissipative systems,” *Annu. Rev. Phys. Chem.* **56**, 187 (2005).
- ³³U. Weiss, *Quantum Dissipative Systems*, World Scientific, Singapore, 2021, 5th ed.
- ³⁴A. Purkayastha, A. Dhar, and M. Kulkarni, “Out-of-equilibrium open quantum systems: A comparison of approximate quantum master equation approaches with exact results,” *Phys. Rev. A* **93**, 062114 (2016).
- ³⁵P. P. Hofer, M. Perarnau-Llobet, L. D. M. Miranda, G. Haack, R. Silva, J. B. Brask, and N. Brunner, “Markovian master equations for quantum thermal machines: local versus global approach,” *New J. Phys.* **19**, 123037 (2017).
- ³⁶D. Farina and V. Giovannetti, “Open-quantum-system dynamics: Recovering positivity of the Redfield equation via the partial secular approximation,” *Phys. Rev. A* **100**, 012107 (2019).
- ³⁷F. Haake and R. Reibold, “Strong damping and low-temperature anomalies for the harmonic oscillator,” *Phys. Rev. A* **32**, 2462 (1985).
- ³⁸W. G. Unruh and W. H. Zurek, “Reduction of a wave packet in quantum Brownian motion,” *Phys. Rev. D* **40**, 1071 (1989).
- ³⁹B. L. Hu, J. P. Paz, and Y. Zhang, “Quantum Brownian motion in a general environment: Exact master equation with nonlocal dissipation and colored noise,” *Phys. Rev. D* **45**, 2843 (1992).
- ⁴⁰J. J. Halliwell and T. Yu, “Alternative derivation of the Hu-Paz-Zhang master equation of quantum Brownian motion,” *Phys. Rev. D* **53**, 1212 (1996).
- ⁴¹R. Karrlein and H. Grabert, “Exact time evolution and master

- equations for the damped harmonic oscillator,” *Phys. Rev. E* **55**, 153 (1997).
- ⁴²G. W. Ford and R. F. O’Connell, “Exact solution to the Hu–Paz–Zhang master equation,” *Phys. Rev. D* **64**, 105020 (2001).
- ⁴³R. X. Xu, B. L. Tian, J. Xu, and Y. J. Yan, “Exact dynamics of driven Brownian oscillators,” *J. Chem. Phys.* **130**, 074107 (2009).
- ⁴⁴W.-M. Huang and W.-M. Zhang, “Nonperturbative renormalization of quantum thermodynamics from weak to strong couplings,” *Phys. Rev. Research* **4**, 023141 (2022).
- ⁴⁵H. Gong, Y. Wang, H. D. Zhang, R. X. Xu, X. Zheng, and Y. J. Yan, “Thermodynamic free-energy spectrum theory for open quantum systems,” *J. Chem. Phys.* **153**, 214115 (2020).
- ⁴⁶H. Gong, Y. Wang, X. Zheng, R. X. Xu, and Y. J. Yan, “Nonequilibrium work distributions in quantum impurity system–bath mixing processes,” *J. Chem. Phys.* **157**, 054109 (2022).
- ⁴⁷Y. Tanimura, “Numerically “exact” approach to open quantum dynamics: The hierarchical equations of motion (HEOM),” *J. Chem. Phys.* **153**, 020901 (2020).
- ⁴⁸R. X. Xu, Y. Mo, P. Cui, S. H. Lin, and Y. J. Yan, “Non-Markovian quantum dissipation in the presence of external fields,” in *Progress in Theoretical Chemistry and Physics, Vol. 12: Advanced Topics in Theoretical Chemical Physics*, edited by J. Maruani, R. Lefebvre, and E. Brändas, pages 7–40, Kluwer, Dordrecht, 2003.
- ⁴⁹Y. Meir and N. S. Wingreen, “Landauer formula for the current through an interacting electron region,” *Phys. Rev. Lett.* **68**, 2512 (1992).
- ⁵⁰T. Ojanen and A.-P. Jauho, “Mesoscopic Photon Heat Transistor,” *Phys. Rev. Lett.* **100**, 155902 (2008).
- ⁵¹J. Hu, R. X. Xu, and Y. J. Yan, “Padé spectrum decomposition of Fermi function and Bose function,” *J. Chem. Phys.* **133**, 101106 (2010).
- ⁵²J. Hu, M. Luo, F. Jiang, R. X. Xu, and Y. J. Yan, “Padé spectrum decompositions of quantum distribution functions and optimal hierarchical equations of motion construction for quantum open systems,” *J. Chem. Phys.* **134**, 244106 (2011).
- ⁵³J. J. Ding, J. Xu, J. Hu, R. X. Xu, and Y. J. Yan, “Optimized hierarchical equations of motion for Drude dissipation with applications to linear and nonlinear optical responses,” *J. Chem. Phys.* **135**, 164107 (2011).
- ⁵⁴J. J. Ding, R. X. Xu, and Y. J. Yan, “Optimizing hierarchical equations of motion for quantum dissipation and quantifying quantum bath effects on quantum transfer mechanisms,” *J. Chem. Phys.* **136**, 224103 (2012).
- ⁵⁵L. Cui, H. D. Zhang, X. Zheng, R. X. Xu, and Y. J. Yan, “Highly efficient and accurate sum-over-poles expansion of Fermi and Bose functions at near zero temperatures: Fano spectrum decomposition scheme,” *J. Chem. Phys.* **151**, 024110 (2019).
- ⁵⁶H. D. Zhang, L. Cui, H. Gong, R. X. Xu, X. Zheng, and Y. J. Yan, “Hierarchical equations of motion method based on Fano spectrum decomposition for low temperature environments,” *J. Chem. Phys.* **152**, 064107 (2020).
- ⁵⁷Z. H. Chen, Y. Wang, X. Zheng, R. X. Xu, and Y. J. Yan, “Universal time-domain Prony fitting decomposition for optimized hierarchical quantum master equations,” *J. Chem. Phys.* **156**, 221102 (2022).
- ⁵⁸Y. Wang, R. X. Xu, and Y. J. Yan, “Entangled system-and-environment dynamics: Phase-space dissipaton theory,” *J. Chem. Phys.* **152**, 041102 (2020).
- ⁵⁹D. Segal and A. Nitzan, “Spin-Boson Thermal Rectifier,” *Phys. Rev. Lett.* **94**, 034301 (2005).
- ⁶⁰F. Giazotto, T. T. Heikkilä, A. Luukanen, A. M. Savin, and J. P. Pekola, “Opportunities for mesoscopics in thermometry and refrigeration: Physics and applications,” *Rev. Mod. Phys.* **78**, 217 (2006).
- ⁶¹O.-P. Saira, M. Meschke, F. Giazotto, A. M. Savin, M. Möttönen, and J. P. Pekola, “Heat Transistor: Demonstration of Gate-Controlled Electronic Refrigeration,” *Phys. Rev. Lett.* **99**, 027203 (2007).
- ⁶²L. E. Bell, “Cooling, Heating, Generating Power, and Recovering Waste Heat with Thermoelectric Systems,” *Science* **321**, 1457 (2008).
- ⁶³P. L. Du, Y. Wang, R. X. Xu, H. D. Zhang, and Y. J. Yan, “System-bath entanglement theorem with Gaussian environments,” *J. Chem. Phys.* **152**, 034102 (2020).