Optimal consumption-investment with coupled constraints on consumption and investment strategies in a regime switching market with random coefficients

Ying Hu * Xiaomin Shi[†] Zuo Quan Xu[‡]
November 11, 2022

This paper studies finite-time optimal consumption-investment problems with power, logarithmic and exponential utilities, in a regime switching market with random coefficients, subject to coupled constraints on the consumption and investment strategies. We provide explicit optimal consumption-investment strategies and optimal values for the problems in terms of the solutions to some diagonally quadratic backward stochastic differential equation (BSDE) systems and linear BSDE systems with unbound coefficients. Some of these BSDEs are new in the literature and solving them is one of the main theoretical contributions of this paper. We accomplish the latter by applying the truncation, approximation technique to get some a priori uniformly lower and upper bounds for their solutions.

Key words. optimal consumption-investment, regime switching, random coefficients, backward stochastic differential equation

Mathematics Subject Classification (2020) 91B16 93E20 60H30 91G10

1 Introduction

In a consumption-investment model, one aims at maximizing his utility of consumption and wealth by choosing the best consumption and investment strategies in a financial market. Following the pioneering work of Merton [26], a large volume of works has been done on the model. In particular, various constraints such as bankruptcy prohibition, subsistence consumption requirement, wealth-dependent investment and consumption constraints are introduced into the model; see, e.g., Guan, Xu and Yi [9], Sethi [30], Xu and Yi [35], Zariphopoulou [38]. All these papers focus on Markovian markets. The model has also been considered in non-Markovian markets with general random

^{*}Univ Rennes, CNRS, IRMAR-UMR 6625, F-35000 Rennes, France. Partially supported by Lebesgue Center of Mathematics "Investissements d'avenir" program-ANR-11-LABX-0020-01. Email: ying.hu@univ-rennes1.fr

[†]School of Statistics and Mathematics, Shandong University of Finance and Economics, Jinan 250100, China. Partially supported by NSFC (No. 11801315), NSF of Shandong Province (No. ZR2018QA001). Email: shixm@mail.sdu.edu.cn

[‡]Department of Applied Mathematics, The Hong Kong Polytechnic University, Kowloon, Hong Kong. Partially supported by NSFC (No. 11971409), Hong Kong RGC (GRF 15202421 and 15204622), The PolyU-SDU Joint Research Center on Financial Mathematics, The CAS AMSS-PolyU Joint Laboratory of Applied Mathematics, The Research Centre for Quantitative Finance, The Hong Kong Polytechnic University. Email: maxu@polyu.edu.hk

coefficients, by Cox and Huang [5], Karatzas, Lehoczky and Shreve [18], Matoussi and Xing [25], Xing [34]. The method used in [5] and [18] is known as martingale method nowadays. Please refer to the seminar monograph Karatzas and Shreve [19] for a systematic account on this method for more advanced (incomplete/constrained) markets.

On the other hand, Markov chain is usually adopted to reflect, at the macroeconomic level, the market status (such as bull and bear) in the literature. This is the famous regime switching model of market; see Hamilton [10]. Utility maximization and mean-variance models in a regime switching market have been studied by many researchers; see, e.g., [2, 24, 31–33, 36, 37, 39, 40]. In these works, the market parameters are assumed to be deterministic functions of time for each given regime. This allows the authors to apply ordinary or partial differential equation (ODE or PDE, for short) method to solve their problems. In practice, however, even in a bull market, the market parameters, such as the interest rate, stock appreciation rates and volatilities are affected by the uncertainties caused by many factors such as politics, economic, legal, military, corporate governance. Thus, it is too restrictive to set market parameters as deterministic function of time even if the market status is known. It is necessary to allow the market parameters to depend on not only the Markov chain but also other random resources. With this in mind, the authors consider stochastic linear quadratic control and mean-variance problems in [13–15]. Along this framework, this paper generalizes the consumption-investment model of Cheridito and Hu [4] to a regime switching market with random coefficients.

This paper studies optimal consumption-investment problems with power, logarithmic and exponential utilities in a regime switching market with parameters depending on both a Markov chain and Brownian motion. Because of the randomness of the market parameters, the ODE and PDE methods will no longer work for our problem. To deal with the randomness, it is necessary to apply the backward stochastic differential equation (BSDE) theory of Pardoux and Peng [28]. Since its inception in 1990, BSDE theory has become a powerful tool in dealing with modern financial engineering problems such as portfolio selection and asset pricing problems. For instance, Rouge and El Karoui [29] characterize the price equation via exponential utility maximization and quadratic BSDEs. By only requiring the strategies take their values in some closed sets, Hu, Imkeller and Müller [11] give solutions to exponential/power/logarithmic utility maximization of terminal wealth and Cheridito and Hu [4] take into account the intermediate consumption. Becherer [3] and Morlais [27] study exponential utility optimization and indifference valuation with jumps using quadratic BSDEs driven by random measures. Taking ambiguity and time-consistent ambiguityaverse preferences into account, Laeven and Stadye [23] investigate the robust portfolio choice and indifference valuation by quadratic BSDEs with infinite activity jumps. Kramkov and Pulido [22] solve a quadratic BSDE system arising from a price impact model. Because of regime switching, the BSDE systems in our problems are all multidimensional, making them harder to study compared to one dimensional case.

Meanwhile, in the study of consumption-investment problems, one often assumes that the consumption and investment strategies are subject to separate constraints; see, e.g., [4]. In practice, however, the constraints on them may be coupled together. For instance, when there is no individual constraint on one's consumption and investment strategies but he is not allowed to borrow money, then the constraints are coupled together. This renders the study of the coupled constraints on the consumption and investment strategies, which is, to the best of our knowledge, still rare in the literature, partially due to its complexity. In this paper, different from [4], we consider general,

not necessarily product, coupled constraints on consumption and investment strategies. This brings new mathematical challenges to our analysis for the related BSDE systems.

We provide explicit optimal consumption and investment strategies in terms of solutions to some BSDE systems. Because of the emergence of regime switching, the BSDE systems in our model are actually diagonally quadratic coupled through the generator of the Markov chain. Different from [4], the quadratic BSDE systems in our paper could not be directly covered by existing literature. Solving these systems is the key theoretical contribution of this paper and we accomplish this by first doing some approximation and truncation of the generators, then finding uniformly lower and upper bounds, and eventually taking limit to obtain the desired solutions. The uniqueness of their solutions are also proved by pure BSDE techniques, so the method may be applied other problems in BSDE theory.

The rest part of this paper is organized as follows. At the end of this section, we introduce some notations and recall some facts about BMO martingales that will be used frequently in the subsequent analysis. In Section 2, we present the financial market and formulate the consumption-investment problem in a regime switching market with random coefficients. In Sections 3, 4, 5, we solve the problem for the power, logarithmic and exponential utilities, respectively. For each utility, we first specify the consumption-investment constraint and define the admissible strategies. We then accomplish the solvability of some related BSDE systems. With the aid of their solutions, we finally provide the optimal consumption and investment strategies and the optimal value of the problem. We conclude the paper in Section 6.

Notation

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a fixed complete probability space on which are defined a standard n-dimensional Brownian motion $W_t = (W_{1,t}, \dots, W_{n,t})'$ and a continuous-time stationary Markov chain α_t valued in a finite state space $\mathcal{M} = \{1, 2, \dots, \ell\}$ with $\ell \geq 1$. We assume $\{W_t\}_{t \geq 0}$ and $\{\alpha_t\}_{t \geq 0}$ are independent processes. The Markov chain has a generator $Q = (q^{ij})_{\ell \times \ell}$ with $q^{ij} \geq 0$ for $i \neq j$ and $\sum_{j=1}^{\ell} q^{ij} = 0$ for every $i \in \mathcal{M}$. Define the filtrations $\mathcal{F}_t = \sigma\{W_s, \alpha_s : 0 \leq s \leq t\} \bigvee \mathcal{N}$ and $\mathcal{F}_t^W = \sigma\{W_s : 0 \leq s \leq t\} \bigvee \mathcal{N}$, where \mathcal{N} is the totality of all the \mathbb{P} -null sets of \mathcal{F} .

Throughout this paper, we denote by \mathbb{R}^n the set of *n*-dimensional column vectors, by \mathbb{R}^n_+ the set of vectors in \mathbb{R}^n whose components are nonnegative, by $\mathbb{R}^{m \times n}$ the set of $m \times n$ real matrices, and by \mathbb{S}^n the set of symmetric $n \times n$ real matrices. For $x \in \mathbb{R}$, we define $x^+ := \max\{x, 0\}$, and $x^- := \max\{-x, 0\}$. If $M = (m_{ij}) \in \mathbb{R}^{m \times n}$, we denote its transpose by M', and its norm by $|M| = \sqrt{\sum_{ij} m_{ij}^2}$. If $M \in \mathbb{S}^n$ is positive definite (positive semidefinite), we write $M > (\geqslant)$ 0. We write $A > (\geqslant)$ B if $A, B \in \mathbb{S}^n$ and $A - B > (\geqslant)$ 0. We use the following spaces throughout the

paper:

$$L_{\mathcal{F}_{T}}^{2}(\Omega;\mathbb{R}) = \left\{ \xi : \Omega \to \mathbb{R} \mid \xi \text{ is } \mathcal{F}_{T}\text{-measurable, and } \mathbb{E}(|\xi|^{2}) < \infty \right\},$$

$$L_{\mathcal{F}_{T}}^{\infty}(\Omega;\mathbb{R}) = \left\{ \xi : \Omega \to \mathbb{R} \mid \xi \text{ is } \mathcal{F}_{T}\text{-measurable, and essentially bounded} \right\},$$

$$L_{\mathcal{F}}^{2}(0,T;\mathbb{R}) = \left\{ \phi : [0,T] \times \Omega \to \mathbb{R} \mid \phi \text{ is an } \{\mathcal{F}_{t}\}_{t \geqslant 0}\text{-predictable process with } \mathbb{E} \int_{0}^{T} |\phi_{t}|^{2} dt < \infty \right\},$$

$$L_{\mathcal{F}}^{1}(0,T;\mathbb{R}) = \left\{ \phi : [0,T] \times \Omega \to \mathbb{R} \mid \phi \text{ is an } \{\mathcal{F}_{t}\}_{t \geqslant 0}\text{-predictable process with } \mathbb{E} \int_{0}^{T} |\phi_{t}| dt < \infty \right\},$$

$$L_{\mathcal{F}}^{\infty}(0,T;\mathbb{R}) = \left\{ \phi : [0,T] \times \Omega \to \mathbb{R} \mid \phi \text{ is an } \{\mathcal{F}_{t}\}_{t \geqslant 0}\text{-predictable essentially bounded process} \right\}.$$

These definitions are generalized in the obvious way to the cases that \mathcal{F} is replaced by \mathcal{F}^W and \mathbb{R} by \mathbb{R}^n , $\mathbb{R}^{n \times m}$ or \mathbb{S}^m .

In our analysis, some arguments such s, t, ω , as well as some terms including "almost surely" (a.s.) and "almost everywhere" (a.e.) may be suppressed for notation simplicity in some circumstances when no confusion occurs.

BMO martingales

We recall some facts about BMO martingales (see Kazamaki [20]). They will be used in our subsequent analysis.

For a $\Lambda \in L^2_{\mathcal{F}^W}(0,T;\mathbb{R}^n)$, the process $\int_0^{\cdot} \Lambda_s' dW_s$ is called a BMO martingale (on [0,T]) if there exists a constant K > 0 such that

$$\mathbb{E}\left[\int_{\tau}^{T} |\Lambda_{s}|^{2} ds \mid \mathcal{F}_{\tau}^{W}\right] \leqslant K$$

for all $\{\mathcal{F}_t^W\}_{t\geqslant 0}$ -stopping times $\tau\leqslant T$. The set of BMO martingales is defined as

$$L_{\mathcal{F}^W}^{2, \text{BMO}}(0, T; \mathbb{R}^n) = \bigg\{ \Lambda \in L_{\mathcal{F}^W}^2(0, T; \mathbb{R}^n) \ \bigg| \ \int_0^{\cdot} \Lambda_s' dW_s \text{ is a BMO martingale on } [0, T] \bigg\}.$$

The following two important properties of BMO martingales will be used in our below arguments without claim.

• If $\int_0^{\cdot} \Lambda'_s dW_s$ is a BMO martingale, then Doléans-Dade stochastic exponential

$$\mathcal{E}\left(\int_0^{\cdot} \Lambda_s' dW_s\right)$$

is a uniformly integrable martingale.

• If $\int_0^{\cdot} \Lambda'_s dW_s$ and $\int_0^{\cdot} Z'_s dW_s$ are both BMO martingales, then $\widetilde{W} := W - \int_0^{\cdot} Z_s ds$ is a standard Brownian motion, and $\int_0^{\cdot} \Lambda'_s d\widetilde{W}_s$ is a BMO martingale under the probability measure $\widetilde{\mathbb{P}}$ defined by

$$\left. \frac{d\widetilde{\mathbb{P}}}{d\mathbb{P}} \right|_{\mathcal{F}_{W}^{W}} = \mathcal{E}\bigg(\int_{0}^{T} Z_{s}' dW_{s}\bigg).$$

2 Problem formulation

2.1 Financial market

Consider a financial market consisting of a risk-free asset (the money market instrument or bond) whose price is S_0 and m risky securities (the stocks) whose prices are S_1, \ldots, S_m . Assume $m \leq n$, i.e., the number of risky securities is no more than that of the market risk resources (namely the Brownian motion). When m < n the market is incomplete. These asset prices are driven by SDEs:

$$\begin{cases} dS_{0,t} = r_t^{\alpha_t} S_{0,t} dt, \\ S_{0,0} = s_0, \end{cases}$$

and

$$\begin{cases} dS_{k,t} = S_{k,t} \left(\mu_{k,t}^{\alpha_t} dt + \sum_{j=1}^n \sigma_{kj,t}^{\alpha_t} dW_{j,t} \right), \\ S_{k,0} = s_k, \end{cases}$$

where r_t^i is the interest rate process and $\mu_{k,t}^i$ and $\sigma_{k,t}^i := (\sigma_{k1,t}^i, \dots, \sigma_{kn,t}^i)$ are the appreciation rate process and volatility rate process of the kth risky security corresponding to a market regime $\alpha_t = i$, for every $k = 1, \dots, m$ and $i \in \mathcal{M}$. Recall that α_t follows a continuous-time stationary Markov chain valued in a finite state space $\mathcal{M} = \{1, 2, \dots, \ell\}$ with $\ell \geqslant 1$. When $\ell = 1$, there is no regime switching and the market becomes the classical Black-Scholes market. In our below argument we assume $\ell > 1$, although all the results remain true if $\ell = 1$.

Define the appreciate vector

$$\mu_t^i = (\mu_{1,t}^i, \dots, \mu_{m,t}^i)',$$

and volatility matrix

$$\sigma_t^i = \begin{pmatrix} \sigma_{1,t}^i \\ \vdots \\ \sigma_{m,t}^i \end{pmatrix} \equiv (\sigma_{kj,t}^i)_{m \times n}, \text{ for each } i \in \mathcal{M}.$$

2.2 Optimal investment-consumption problem

Consider a small investor, whose actions cannot affect the asset prices. He will decide at every time $t \in [0,T]$ the proportion $\pi_{j,t}$ of his wealth to invest in the jth risky asset, $j=1,\ldots,m$, as well as the proportion c_t of his wealth to consume. The vector process $\pi_t := (\pi_{1,t},\ldots,\pi_{m,t})'$ is called a portfolio of the investor. The pair (π,c) is called a consumption-investment strategy. The investor's self-financing wealth process X corresponding to a consumption-investment strategy (π,c) is a strong solution of the SDE (see, e.g., [19]):

$$\begin{cases}
dX_t = X_t [r_t^{\alpha_t} + \pi_t' b_t^{\alpha_t} - c_t] dt + X_t \pi_t' \sigma_t^{\alpha_t} dW_t, \\
X_0 = x > 0, \ \alpha_0 = i_0 \in \mathcal{M},
\end{cases}$$
(2.1)

where $b_t^{\alpha_t} := \mu_t^{\alpha_t} - r_t^{\alpha_t} \mathbf{1}_m$ and $\mathbf{1}_m$ is the *m*-dimensional vector with all entries being one. By Itô's lemma, one can easily show that the process X is always positive, hence the investor would never be bankrupt.

The investor's problem is to maximize

$$J(x, i_0; \pi, c) := \mathbb{E}\left[\int_0^T e^{-\int_0^t \rho_s^{\alpha_s} ds} U(c_t X_t) dt + e^{-\int_0^T \rho_s^{\alpha_s} ds} U(X_T)\right], \quad \text{s.t. } (\pi, c) \in \mathcal{U},$$
 (2.2)

and determine the value function

$$V(x, i_0) := \sup_{(\pi, c) \in \mathcal{U}} J(x, i_0; \pi, c),$$

where $\rho^i \in L^{\infty}_{\mathcal{F}^W}(0,T;\mathbb{R})$, $i \in \mathcal{M}$, are the discount factor processes, U is the utility function of the investor, and \mathcal{U} is the admissible set of consumption-investment strategies. The utility U and admissible set \mathcal{U} will be defined in the sequel case by case. In particular, the consumption-investment strategies can be subject to fairly general constraints. Let Θ denote the constraint set for them, which is assumed to be a given closed nonempty set in $\mathbb{R}^m \times \mathbb{R}_+$.

Example 2.1 Here are some important and interesting examples for the constraint set Θ in financial practice.

• If the investor are not allowed to borrow money, then the total consumption and investment cannot beyond 100%, that is,

$$\Theta = \left\{ (\pi, c) \in \mathbb{R}^m \times \mathbb{R}_+ \mid \sum_{j=1}^m \pi_j + c \leqslant 1 \right\}.$$

Because the constraints on the consumption and investment strategies are coupled together, this will bring some mathematical challenges to our analysis for the related BSDE systems below.

• If no shorting is allowed in the stocks, then

$$\Theta = \left\{ (\pi, c) \in \mathbb{R}^m \times \mathbb{R}_+ \mid \pi_j \geqslant 0, \ j = 1, \dots, m \right\}.$$

In this case Θ is cone.

• There may be restrictions on the investment strategies on some stocks; for instance

$$\Theta = \left\{ (\pi, c) \in \mathbb{R}^m \times \mathbb{R}_+ \mid \pi_j \in [d_j, e_j], \ j = 1, \dots, m \right\};$$

or constraint on the consumption strategies such as

$$\Theta = \left\{ (\pi, c) \in \mathbb{R}^m \times \mathbb{R}_+ \mid c \in [0, \frac{1}{5}] \right\};$$

or coupled constraints on both the investment and consumption strategies

$$\Theta = \left\{ (\pi, c) \in \mathbb{R}^m \times \mathbb{R}_+ \ \middle| \ c \in [0, \frac{1}{5}], \ \pi_j \in [d_j, e_j], \ j = 1, \dots, m \right\}.$$

We put the following standard assumption for the market parameters.

Assumption 1 For all $i \in \mathcal{M}$,

$$\begin{cases} r^i \in L^{\infty}_{\mathbb{F}^W}(0,T;\mathbb{R}), \ \mu^i \in L^{\infty}_{\mathbb{F}^W}(0,T;\mathbb{R}^m), \\ \sigma^i \in L^{\infty}_{\mathbb{F}^W}(0,T;\mathbb{R}^{m \times n}), \ \rho^i \in L^{\infty}_{\mathbb{F}^W}(0,T;\mathbb{R}). \end{cases}$$

Also, there exists a constant $\delta > 0$ such that $\sigma^i(\sigma^i)' \geqslant \delta I_m$ for all $i \in \mathcal{M}$, where I_m denotes the m-dimensional identity matrix.

Due to different features of the power, logarithmic and exponential utilities, they must be dealt with different methods. We study them in the subsequent Sections 3, 4 and 5, respectively.

3 Power utility

In this section, we assume that the investor's utility function is

$$U(x) = \frac{1}{\gamma}x^{\gamma}, \quad x > 0, \quad \gamma \in (-\infty, 0) \cup (0, 1)$$

and the admissible consumption-investment set is defined as

$$\mathcal{U} = \left\{ (\pi, c) \mid \int_0^T (|\pi_t|^2 + c_t) dt < \infty, a.s., \ (\pi_t(\omega), c_t(\omega)) \in \Theta, \ a.e., \ a.s. \right\},$$

if $\gamma \in (0,1)$; and

$$\mathcal{U} = \left\{ (\pi, c) \mid \int_0^T (|\pi_t|^2 + c_t) dt < \infty, a.s., \ (\pi_t(\omega), c_t(\omega)) \in \Theta, \ a.e., \ a.s, \\ (X_t^{\gamma})_{0 \leqslant t \leqslant T} \text{ belongs to class (D) and } \mathbb{E} \left[\int_0^T (c_t X_t)^{\gamma} dt \right] < \infty \right\},$$

if $\gamma < 0.1$

Assumption 2 If $\gamma \in (0,1)$, then no consumption or investment is always permitted, namely $(0\mathbf{1}_m,0) \in \Theta$. If $\gamma < 0$, then positive consumption is always needed (for otherwise the admissible set \mathcal{U} ie empty), there exists $\varepsilon > 0$ such that $(0\mathbf{1}_m,\varepsilon) \in \Theta$.

To tackle problem (2.2), we introduce the following ℓ -dimensional BSDE system:

$$\begin{cases} dP_t^i = -[f^i(P_t^i, \Lambda_t^i) - (\rho^i - \gamma r^i)P^i + \sum_{j=1}^{\ell} q^{ij}P^j]dt + (\Lambda_t^i)'dW, \\ P_T^i = 1, \\ P^i > 0, \quad i \in \mathcal{M}, \end{cases}$$
(3.1)

where, for any $(P, \Lambda) \in \mathbb{R}_+ \times \mathbb{R}^n$,

$$\underline{f^{i}(P,\Lambda)} = \gamma \underset{(\pi,c) \in \Theta}{\operatorname{ess sup}} \left[-\frac{1-\gamma}{2} |\pi'\sigma^{i}|^{2} P + \pi'(Pb^{i} + \sigma^{i}\Lambda) + \frac{c^{\gamma}}{\gamma} - Pc \right].$$

¹We say $(X_t^{\gamma})_{0 \leqslant t \leqslant T}$ belongs to class (D) if $\{X_{\tau}^{\gamma} : \tau \text{ stopping time valued in } [0,T]\}$ is a family of uniformly integrable random variables.

Remark 3.1 If there is no consumption-investment constraint, namely $\Theta = \mathbb{R}^n \times \mathbb{R}_+$, then

$$f^{i}(P,\Lambda) = \frac{1}{2} \frac{\gamma}{1-\gamma} \frac{1}{P} (Pb + \sigma\Lambda)' (\sigma\sigma')^{-1} (Pb + \sigma\Lambda) + (1-\gamma)P^{-\frac{\gamma}{1-\gamma}}.$$

Definition 3.2 A vector process $(P^i, \Lambda^i)_{i \in \mathcal{M}}$ is called a solution to the ℓ -dimensional BSDE (3.1), if it satisfies (3.1), and $(P^i, \Lambda^i) \in L^{\infty}_{\mathcal{F}W}(0, T; \mathbb{R}) \times L^{2, \text{BMO}}_{\mathcal{F}W}(0, T; \mathbb{R}^n)$ for all $i \in \mathcal{M}$. A solution $(P^i, \Lambda^i)_{i \in \mathcal{M}}$ to (3.1) is called uniformly positive if $P^i_t \geqslant \delta$, for all $t \in [0, T]$, a.s. with some deterministic constant $\delta > 0$.

The following comparison theorem for multidimensional BSDE systems firstly appeared in [16] (one can find a concise version in [12, Lemma 2.2] or [13, Lemma 3.4]). We shall use it frequently in the study of BSDEs.

Lemma 3.3 Suppose $(Y^i, Z^i)_{i \in \mathcal{M}}$ and $(\overline{Y}^i, \overline{Z}^i)_{i \in \mathcal{M}}$ satisfy the following two ℓ -dimensional BSDE systems, respectively:

$$Y_t^i = \xi^i + \int_t^T g^i(s, Y_s^i, Y_s^{-i}, Z_s^i) ds - \int_t^T (Z_s^i)' dW_s, \text{ for all } i \in \mathcal{M};$$

and

$$\overline{Y}_t^i = \overline{\xi}^i + \int_t^T \overline{g}^i(s, \overline{Y}_s^i, \overline{Y}_s^{-i}, \overline{Z}_s^i) ds - \int_t^T (\overline{Z}_s^i)' dW_s, \quad for \ all \ i \in \mathcal{M},$$

where $Y_s^{-i} = (Y_s^1, \dots, Y_s^{i-1}, Y_s^{i+1}, \dots, Y_s^{\ell})$. Also suppose that, for all $i \in \mathcal{M}$,

1.
$$\xi^i$$
, $\overline{\xi}^i \in L^2_{\mathcal{F}^W}(\Omega; \mathbb{R})$, and $\xi^i \leqslant \overline{\xi}^i$;

2. there exists a constant K > 0 such that

$$|g^{i}(s, y, z) - g^{i}(s, \overline{y}, \overline{z})| \leq K(|y - \overline{y}| + |z - \overline{z}|),$$

 $\ \, for\,\,any\,\,z,\overline{z}\in\mathbb{R}^n,\,\,y=(y^i,y^{-i}),\,\overline{y}=(\overline{y}^i,\overline{y}^{-i})\in\mathbb{R}^\ell;$

3. $g^{i}(s, y, z)$ is nondecreasing in every y^{j} , $j \neq i \in \mathcal{M}$; and

4.
$$g^i(s, \overline{Y}_s^i, \overline{Y}_s^{-i}, \overline{Z}_s^i) \leqslant \overline{g}^i(s, \overline{Y}_s^i, \overline{Y}_s^{-i}, \overline{Z}_s^i)$$
.

Then $Y_t^i \leqslant \overline{Y}_t^i$ a.s. for all $t \in [0,T]$ and $i \in \mathcal{M}$.

Remark 3.4 If conditions 2 and 3 hold for \overline{g}^i instead of g^i , and condition 4 is replaced by

$$g^i(s,Y_s^i,Y_s^{-i},Z_s^i)\leqslant \overline{g}^i(s,Y_s^i,Y_s^{-i},Z_s^i)$$

in the above lemma, then the conclusion still holds.

Theorem 3.5 Suppose $\gamma \in (0,1)$ and Assumptions 1 and 2 hold. Then there is a unique uniformly positive solution $(P^i, \Lambda^i)_{i \in \mathcal{M}}$ to the BSDE (3.1).

Proof: Let a > 1 be a large constant such that

$$-\rho^i + \gamma r^i \geqslant -a,\tag{3.2}$$

and

$$\frac{\gamma}{2(1-\gamma)}(b^i)'(\sigma^i(\sigma^i)')^{-1}b^i - \rho^i + \gamma r^i \leqslant a(1-\gamma), \tag{3.3}$$

hold simultaneously for all $i \in \mathcal{M}$. Denote $a_1 := e^{-aT}$ and $a_2 := 2e^{aT}$.

Let $g: \mathbb{R} \to [0,1]$ be a smooth truncation function satisfying g(x) = 0 for $x \in (-\infty, \frac{1}{2}a_1]$, and g(x) = 1 for $x \in [a_1, +\infty)$. For $k \ge 1$, $(t, P, \Lambda) \in [0, T] \times \mathbb{R} \times \mathbb{R}^n$, $i \in \mathcal{M}$, define

$$f^{k,i}(t,P,\Lambda) = \operatorname*{ess\ inf}_{\tilde{P} \in \mathbb{R}, \tilde{\Lambda} \in \mathbb{R}^n} \left[f^i(t,\tilde{P},\tilde{\Lambda}) g(\tilde{P}) + k|P - \tilde{P}| + k|\Lambda - \tilde{\Lambda}| \right].$$

Then it is uniformly Lipschitz continuous in (P, Λ) , and increasingly approaches to $f^i(t, P, \Lambda)g(P)$ as k goes to infinity. According to Assumption 2, we have $f^i \ge 0$, hence $f^{k,i} \ge 0$.

The following BSDE system

$$\begin{cases} dP^{k,i} = -\left[f^{k,i}(P^{k,i}, \Lambda^{k,i}) - (\rho^i - \gamma r^i)P^{k,i} + \sum_{j=1}^{\ell} q^{ij}P^{k,j}\right]dt + (\Lambda^{k,i})^{\top}dW, \\ P_T^{k,i} = 1, & \text{for all } i \in \mathcal{M}, \end{cases}$$

has a Lipschitz generator², so it admits a unique solution, denoted by $(P^{k,i}, \Lambda^{k,i})_{i \in \mathcal{M}}$. It is direct to verify that

$$(\underline{P}_t^i, \underline{\Lambda}_t^i) = (e^{-a(T-t)}, 0), \quad i \in \mathcal{M}$$

is the unique solution to the following linear BSDE system:

$$\begin{cases} d\underline{P}^i = -\left[-a\underline{P}^i + \sum_{j=1}^{\ell} q^{ij}\underline{P}^j\right] dt + (\underline{\Lambda}^i)' dW, \\ \underline{P}_T^i = 1, \quad i \in \mathcal{M}. \end{cases}$$

Notice that $f^{k,i}(P,\Lambda) \ge 0$ and recall (3.2), so

$$f^{k,i}(\underline{P}^i,\underline{\Lambda}^i) - (\rho^i - \gamma r^i)\underline{P}^i + \sum_{j=1}^{\ell} q^{ij}\underline{P}^j \geqslant -(\rho^i - \gamma r^i)\underline{P}^i + \sum_{j=1}^{\ell} q^{ij}\underline{P}^j \geqslant -a\underline{P}^i + \sum_{j=1}^{\ell} q^{ij}\underline{P}^j.$$

By Lemma 3.3, we have

$$a_1 \leqslant e^{-a(T-t)} = \underline{P}_t^i \leqslant P_t^{k,i}, \quad i \in \mathcal{M}.$$

On the other hand,

$$(\bar{P}^i, \bar{\Lambda}^i) = \left(\left(e^{a(T-t)} + \frac{1}{a} (e^{a(T-t)} - 1) \right)^{1-\gamma}, \ 0 \right), \quad i \in \mathcal{M},$$

is a solution to the following BSDE system:

$$\begin{cases} d\bar{P}^i = -\left[a(1-\gamma)\bar{P}^i + (1-\gamma)(\bar{P}^i)^{-\frac{\gamma}{1-\gamma}} + \sum_{j=1}^{\ell} q^{ij}\bar{P}^j\right] dt + (\bar{\Lambda}^i)'dW, \\ \bar{P}_T^i = 1, \quad i \in \mathcal{M}. \end{cases}$$

²As for BSDE $Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s' dW_s$, f is called the generator and ξ is called the terminal value.

Notice, for P > 0,

$$f^{k,i}(t,P,\Lambda) \leq f^{i}(t,P,\Lambda)g(P)$$

$$\leq \gamma \underset{(\pi,c)\in\mathbb{R}^{n}\times\mathbb{R}_{+}}{\operatorname{ess sup}} \left[-\frac{1-\gamma}{2}|\pi'\sigma^{i}|^{2}P + \pi'(Pb^{i} + \sigma^{i}\Lambda) + \frac{c^{\gamma}}{\gamma} - Pc \right]g(P)$$

$$= \frac{1}{2}\frac{\gamma}{1-\gamma}\frac{1}{P}(Pb + \sigma\Lambda)'(\sigma\sigma')^{-1}(Pb + \sigma\Lambda)g(P) + (1-\gamma)P^{-\frac{\gamma}{1-\gamma}}g(P)$$

$$\leq \frac{\gamma}{2(1-\gamma)}(b^{i})'(\sigma^{i}(\sigma^{i})')^{-1}b^{i}P + \frac{\gamma}{1-\gamma}(b^{i})'(\sigma^{i}(\sigma^{i})')^{-1}\sigma^{i}\Lambda$$

$$+ \frac{\gamma}{2(1-\gamma)}\frac{1}{P}(\sigma^{i}\Lambda)'(\sigma^{i}(\sigma^{i})')^{-1}\sigma^{i}\Lambda + (1-\gamma)P^{-\frac{\gamma}{1-\gamma}}. \tag{3.4}$$

Hence, by (3.3),

$$f^{k,i}(\bar{P}^i, \bar{\Lambda}^i) - (\rho^i - \gamma r^i)\bar{P}^i + \sum_{j=1}^{\ell} q^{ij}\bar{P}^j$$

$$\leq \frac{\gamma}{2(1-\gamma)} (b^i)' (\sigma^i(\sigma^i)')^{-1} b^i \bar{P}^i + (1-\gamma)(\bar{P}^i)^{-\frac{\gamma}{1-\gamma}} - (\rho^i - \gamma r^i)\bar{P}^i + \sum_{j=1}^{\ell} q^{ij}\bar{P}^j$$

$$\leq a(1-\gamma)\bar{P}^i + (1-\gamma)(\bar{P}^i)^{-\frac{\gamma}{1-\gamma}} + \sum_{j=1}^{\ell} q^{ij}\bar{P}^j.$$

Then by Lemma 3.3 again, we have

$$P_t^{k,i} \leqslant \bar{P}_t^i \leqslant 2e^{aT} = a_2,$$

and $P^{k,i}$ is increasing in k, for each $i \in \mathcal{M}$. Let $P^i_t = \lim_{k \to \infty} P^{k,i}_t$, $i \in \mathcal{M}$. Since a_1 and a_2 are independent of k, $a_1 \leqslant P^i_t \leqslant a_2$. Recalling $f^{k,i} \ge 0$, (3.4) and the role of the truncation function g, we can regard $(P^{k,i}, \Lambda^{k,i})$ as the solution of a scalar-valued quadratic BSDE for each $i \in \mathcal{M}$. Thus by [21, Proposition 2.4], there exists a process $\Lambda \in L^2_{\mathcal{F}^W}(0,T;\mathbb{R}^{n \times \ell})$ such that (P,Λ) satisfies the BSDE (3.1).

Applying Ítô's formula to $(P^i - a_2)^2$, we obtain, for any stopping times $\tau \leqslant T$,

$$\begin{split} \mathbb{E} \bigg[\int_{\tau}^{T} |\Lambda^{i}|^{2} \; \bigg| \; \mathcal{F}_{\tau}^{W} \bigg] &= (1 - a_{2})^{2} - (P_{\tau}^{i} - a_{2})^{2} \\ &+ 2 \mathbb{E} \bigg[\int_{\tau}^{T} (P^{i} - a_{2}) \Big[f^{i}(P^{i}, \Lambda^{i}) - (\rho^{i} - \gamma r^{i}) P^{i} + \sum_{j=1}^{\ell} q^{ij} P^{j} \Big] ds \; \bigg| \; \mathcal{F}_{\tau}^{W} \bigg] \\ &\leqslant (1 - a_{2})^{2} - (P_{\tau}^{i} - a_{2})^{2} \\ &+ 2 \mathbb{E} \bigg[\int_{\tau}^{T} (P^{i} - a_{2}) \Big[- (\rho^{i} - \gamma r^{i}) P^{i} + \sum_{j=1}^{\ell} q^{ij} P^{j} \Big] ds \; \bigg| \; \mathcal{F}_{\tau}^{W} \bigg] \\ &\leqslant (1 - a_{2})^{2} + 2 \mathbb{E} \bigg[\int_{0}^{T} |P^{i} - a_{2}| \Big| - (\rho^{i} - \gamma r^{i}) P^{i} + \sum_{j=1}^{\ell} q^{ij} P^{j} \Big| ds \; \bigg| \; \mathcal{F}_{\tau}^{W} \bigg], \end{split}$$

where we used the fact that $P^i \leqslant a_2$ and $f^i \geqslant 0$ to get the first inequity. Because $a_1 \leqslant P^i \leqslant a_2$ and Assumption 1, we see the right hand side is upper bounded by a constant. Hence $\Lambda^i \in L_{\mathcal{F}W}^{2, \text{BMO}}(0, T; \mathbb{R}^n)$, for all $i \in \mathcal{M}$. We have now established the existence of the solution.

Next, let us prove the uniqueness. Suppose $(P^i, \Lambda^i)_{i \in \mathcal{M}}$, $(\tilde{P}^i, \tilde{\Lambda}^i)_{i \in \mathcal{M}}$ are two uniformly positive solutions of (3.1). For every $i \in \mathcal{M}$, define processes

$$(Y_t^i, Z_t^i) = \left(\ln P_t^i, \frac{\Lambda_t^i}{P_t^i}\right), \ (\tilde{Y}_t^i, \tilde{Z}_t^i) = \left(\ln \tilde{P}_t^i, \frac{\tilde{\Lambda}_t^i}{\tilde{P}_t^i}\right), \ \text{for } t \in [0, T].$$

$$(3.5)$$

Then (Y^i, Z^i) , $(\tilde{Y}^i, \tilde{Z}^i) \in L^{\infty}_{\mathcal{F}^W}(0, T; \mathbb{R}) \times L^{2, \text{BMO}}_{\mathcal{F}^W}(0, T; \mathbb{R}^n)$, for all $i \in \mathcal{M}$. Furthermore, by Itô's formula, $(Y^i, Z^i)_{i \in \mathcal{M}}$ and $(\tilde{Y}^i, \tilde{Z}^i)_{i \in \mathcal{M}}$ satisfy the following BSDE system:

$$\begin{cases} dY^{i} = -\left[F^{i}(Y^{i}, Z^{i}) + \frac{1}{2}|Z^{i}|^{2} - \rho^{i} + \gamma r^{i} + \sum_{j=1}^{\ell} q^{ij} e^{Y^{j} - Y^{i}}\right] dt + (Z^{i})' dW, \\ Y_{T}^{i} = 0, \quad i \in \mathcal{M}. \end{cases}$$
(3.6)

where for any $(Y, Z) \in \mathbb{R} \times \mathbb{R}^n$

$$F^{i}(Y,Z) = \gamma \operatorname*{ess\,sup}_{(\pi,c)\in\Theta} \Big[-\frac{1-\gamma}{2} |\pi'\sigma^{i}|^{2} + \pi'(b^{i} + \sigma^{i}Z) + \frac{c^{\gamma}}{\gamma} e^{-Y} - c \Big].$$

Because $\ln a_1 \leqslant Y^i \leqslant \ln a_2$ and use Assumption 1, for any $(\pi, c) \in \Theta$,

$$-\frac{1-\gamma}{2}|\pi'\sigma^{i}|^{2} + \pi'(b^{i} + \sigma^{i}Z^{i}) + \frac{c^{\gamma}}{\gamma}e^{-Y^{i}} - c$$

$$\leq -\delta \frac{1-\gamma}{2}|\pi|^{2} + K_{1}|\pi|(1+|Z|) + \sup_{c \in \mathbb{R}_{+}} \left(\frac{c^{\gamma}}{\gamma} \frac{1}{a_{1}} - c\right)$$

$$= -\delta \frac{1-\gamma}{2}|\pi|^{2} + K_{1}|\pi|(1+|Z|) + \frac{1-\gamma}{\gamma}a_{1}^{\frac{1}{1-\gamma}}$$

$$< 0.$$

if $|\pi| > K(1+|Z^i|)$ for sufficient large K. Since $0\mathbf{1}_{m+1} \in \Theta$, $F^i(Y,Z) \geqslant 0$. Therefore

$$F^{i}(Y^{i}, Z^{i}) = \gamma \underset{\substack{(\pi, c) \in \Theta \\ |\pi| \leq K(1+|Z^{i}|)}}{\text{ess sup}} \left[-\frac{1-\gamma}{2} |\pi'\sigma^{i}|^{2} + \pi'(b^{i} + \sigma^{i}Z^{i}) + \frac{c^{\gamma}}{\gamma} e^{-Y^{i}} - c \right]. \tag{3.7}$$

This still holds if we replace $K(1 + |Z^i|)$ by any bigger value. A similar expression holds for $F^i(\tilde{Y}^i, \tilde{Z}^i)$.

Set $\overline{Y}^i = Y^i - \tilde{Y}^i$, $\overline{Z}^i = Z^i - \tilde{Z}^i$, for $i \in \mathcal{M}$. From (3.7), we have

$$\left| F^{i}(Y^{i}, Z^{i}) - F^{i}(Y^{i}, \tilde{Z}^{i}) \right| \leqslant \gamma \underset{|\pi| \leqslant K(1+|Z^{i}|+|\tilde{Z}^{i}|)}{\text{ess sup}} |\pi' \sigma^{i} \overline{Z}^{i}| \leqslant K(1+|Z^{i}|+|\tilde{Z}^{i}|) |\overline{Z}^{i}|.$$

Since $Z^i, \tilde{Z}^i \in L^{2, \text{BMO}}_{\mathcal{F}^W}(0, T; \mathbb{R}^n)$, we can define $\beta^i \in L^{2, \text{BMO}}_{\mathcal{F}^W}(0, T; \mathbb{R}^n)$ in an obvious way such that

$$F^{i}(Y^{i}, Z^{i}) - F^{i}(Y^{i}, \tilde{Z}^{i}) = (\beta^{i})'\overline{Z}^{i},$$

³Hereafter, we shall use K to represent a generic positive constant independent of i, m, n and t, which can be different from line to line.

and

$$|\beta^i| \leqslant K(1 + |Z^i| + |\tilde{Z}^i|).$$

Now applying Itô's formula to $(\overline{Y}^i)^2$, we deduce that

$$(\overline{Y}_{t}^{i})^{2} = \int_{t}^{T} \left\{ 2\overline{Y}^{i} \left[F^{i}(Y^{i}, Z^{i}) - F^{i}(\tilde{Y}^{i}, \tilde{Z}^{i}) + \frac{1}{2} (|Z^{i}|^{2} - |\tilde{Z}^{i}|^{2}) + \sum_{i=1}^{\ell} q^{ij} \left(e^{Y^{j} - Y^{i}} - e^{\tilde{Y}^{j} - \tilde{Y}^{i}} \right) \right] - |\overline{Z}^{i}|^{2} \right\} ds - \int_{t}^{T} 2\overline{Y}^{i} (\overline{Z}^{i})' dW.$$
(3.8)

Notice that the map

$$Y \mapsto F^i(Y, Z), Y \in \mathbb{R}_+$$

is non-increasing for every $Z \in \mathbb{R}^n$ and $i \in \mathcal{M}$. Therefore,

$$\begin{split} \overline{Y}^i \big[F^i(Y^i, Z^i) - F^i(\tilde{Y}^i, \tilde{Z}^i) \big] &= \overline{Y}^i \big[F^i(Y^i, Z^i) - F^i(Y^i, \tilde{Z}^i) + F^i(Y^i, \tilde{Z}^i) - F^i(\tilde{Y}^i, \tilde{Z}^i) \big] \\ &\leqslant \overline{Y}^i \big[F^i(Y^i, Z^i) - F^i(Y^i, \tilde{Z}^i) \big] \\ &= \overline{Y}^i (\beta^i)' \overline{Z}^i. \end{split}$$

For each fixed $i \in \mathcal{M}$, let us introduce the process

$$N_t^i = \mathcal{E}\Big(\int_0^t (\beta_s^i + \frac{1}{2}(Z^i + \tilde{Z}^i))'dW_s\Big).$$

Then N_t^i is a uniformly integrable martingale. Notice Y^i , $i \in \mathcal{M}$ are bounded, from (3.8),

$$(\overline{Y}_t^i)^2 \leqslant \int_t^T \left\{ 2\overline{Y}^i \left[(\beta^i)' \overline{Z}^i + \frac{1}{2} (Z^i + \widetilde{Z}^i)' \overline{Z}^i + K \sum_{j \neq i}^{\ell} \overline{Y}^j \right] \right\} ds - \int_t^T 2\overline{Y}^i (\overline{Z}^i)' dW$$

$$= \int_t^T 2K \overline{Y}^i \sum_{i \neq i}^{\ell} \overline{Y}^j ds - \int_t^T 2\overline{Y}^i (\overline{Z}^i)' d\widetilde{W}^i$$

where

$$\widetilde{W}_t^i := W_t - \int_0^t \left(\beta_s^i + \frac{1}{2} (Z^i + \widetilde{Z}^i) \right) ds,$$

is a Brownian motion under the probability $\widetilde{\mathbb{P}}^i$ defined by

$$\left.\frac{d\widetilde{\mathbb{P}^i}}{d\mathbb{P}}\right|_{\mathcal{F}_T^W} = N_T^i.$$

Taking expectation $\widetilde{\mathbb{E}}^i$ w.r.t. the probability measure $\widetilde{\mathbb{P}}^i$,

$$(\overline{Y}_t^i)^2 \leqslant \widetilde{\mathbb{E}}^i \bigg[\int_t^T 2K \overline{Y}^i \sum_{j \neq i}^\ell \overline{Y}^j ds \ \bigg| \ \mathcal{F}_t^W \bigg] \leqslant K \widetilde{\mathbb{E}}^i \bigg[\int_t^T \sum_{j=1}^\ell (\overline{Y}^j)^2 ds \ \bigg| \ \mathcal{F}_t^W \bigg] \leqslant K \int_t^T \sum_{j=1}^\ell E_s^j ds,$$

by the arithmetic-mean and geometric-mean inequality (AM-GM inequality), where

$$E_t^i = \operatorname{ess sup}_{\omega \in \Omega} (\overline{Y}_t^i)^2.$$

Taking essential supreme on both sides, we deduce

$$E_t^i \leqslant K \int_t^T \sum_{j=1}^{\ell} E_s^j ds.$$

Thus

$$0 \leqslant \sum_{j=1}^{\ell} E_t^j \leqslant K\ell \int_t^T \sum_{j=1}^{\ell} E_s^j ds.$$

We infer from Gronwall's inequality that $\sum_{j=1}^{\ell} E_t^j = 0$, so $\overline{Y}_t^i = 0$ a.s. for $t \in [0, T]$ and all $i \in \mathcal{M}$. This completes the proof of the uniqueness.

Remark 3.6 Please note that the solvability of (3.1) and (3.6) cannot be directly covered by Fan, Hu and Tang [8] since the generators violate the locally Lipschitz condition required in [8].

Theorem 3.7 Suppose $\gamma < 0$ and Assumption 1 and 2 hold. Then there is a unique uniformly positive solution $(P^i, \Lambda^i)_{i \in \mathcal{M}}$ to the BSDE (3.1).

Proof: The proof is similar to the procedure of Theorem 3.5 with different uniformly lower bounds and upper bounds. Hence we will only present how to find these bounds. Other details are left to the interested readers.

Let a' > 0 be a large constant such that

$$\frac{\gamma}{2(1-\gamma)}(b^i)'(\sigma^i(\sigma^i)')^{-1}b^i - \rho^i + \gamma r^i \geqslant -a'(1-\gamma),\tag{3.9}$$

and

$$-\rho^i + \gamma r^i - \gamma \varepsilon \leqslant a', \tag{3.10}$$

hold simultaneously for all $i \in \mathcal{M}$, where $\varepsilon > 0$ is given in Assumption 2. Denote $a'_1 := e^{-a'(1-\gamma)T}$ and $a'_2 := e^{a'T} + \frac{\varepsilon^{\gamma}}{a'}(e^{a'T} - 1)$.

Let $g: \mathbb{R} \to [0,1]$ be a smooth truncation function satisfying g(x) = 0 for $x \in (-\infty, \frac{1}{2}a'_1]$, and g(x) = 1 for $x \in [a'_1, +\infty)$. Notice we have $|xg(x)| \leq |x|$ for all $x \in \mathbb{R}$.

For integer $k \geqslant -\gamma \varepsilon$, $(t, P, \Lambda) \in [0, T] \times \mathbb{R} \times \mathbb{R}^n$, $i \in \mathcal{M}$, define

$$f^{k,i}(t,P,\Lambda) = \underset{\tilde{P} \in \mathbb{R}, \tilde{\Lambda} \in \mathbb{R}^n}{\text{ess sup}} \left[f^i(t,\tilde{P},\tilde{\Lambda})g(\tilde{P}) - k|P - \tilde{P}| - k|\Lambda - \tilde{\Lambda}| \right].$$

And consider the following ℓ -dimensional BSDE:

$$\begin{cases} dP^{k,i} = -\Big[f^{k,i}(P^{k,i}, \Lambda^{k,i}) - (\rho^i - \gamma r^i)P^{k,i}g(P^{k,i}) + \sum_{j=1}^{\ell} q^{ij}P^{k,j}\Big]dt + (\Lambda^{k,i})^{\top}dW, \\ P_T^{k,i} = 1, \quad i \in \mathcal{M}. \end{cases}$$

Since its generator is Lipschitz continuous, it admits a unique solution, denoted by $(P^{k,i}, \Lambda^{k,i})_{i \in \mathcal{M}}$. It is easy to verify that

$$(\underline{P}_t^i, \underline{\Lambda}_t^i) = \left(\left(e^{-a'(T-t)} + \frac{1}{a'} (1 - e^{-a'(T-t)}) \right)^{1-\gamma}, 0 \right), \quad i \in \mathcal{M}$$
(3.11)

is a solution to the following BSDE system:

$$\begin{cases} d\underline{P}^i = -\left[-a'(1-\gamma)\underline{P}^i g(\underline{P}^i) + (1-\gamma)(\underline{P}^i)^{-\frac{\gamma}{1-\gamma}} g(\underline{P}^i) + \sum_{j=1}^{\ell} q^{ij}\underline{P}^j\right] dt + (\underline{\Lambda}^i)' dW, \\ \underline{P}_T^i = 1, \quad i \in \mathcal{M}. \end{cases}$$

Since $\gamma < 0$, we have the following estimates for $f^{k,i}(t, P, \Lambda)$ with P > 0:

$$\begin{split} f^{k,i}(t,P,\Lambda) \geqslant f^i(t,P,\Lambda)g(P) \\ &= \gamma \operatorname*{ess\ sup}_{(\pi,c)\in\Theta} \Big[-\frac{1-\gamma}{2} |\pi'\sigma^i|^2 P + \pi'(Pb^i + \sigma^i\Lambda) + \frac{c^\gamma}{\gamma} - Pc \Big] g(P) \\ &\geqslant \gamma \operatorname*{ess\ sup}_{(\pi,c)\in\mathbb{R}^m\times\mathbb{R}_+} \Big[-\frac{1-\gamma}{2} |\pi'\sigma^i|^2 P + \pi'(Pb^i + \sigma^i\Lambda) + \frac{c^\gamma}{\gamma} - Pc \Big] g(P) \\ &= \frac{\gamma}{2(1-\gamma)} \frac{1}{P} (Pb + \sigma\Lambda)'(\sigma\sigma')^{-1} (Pb + \sigma\Lambda)g(P) + (1-\gamma)P^{-\frac{\gamma}{1-\gamma}} g(P). \end{split}$$

Hence, we deduce

$$\begin{split} f^{k,i}(t,\underline{P}^i,\underline{\Lambda}^i) - (\rho^i - \gamma r^i)\underline{P}^i g(\underline{P}^i) + \sum_{j=1}^\ell q^{ij}\underline{P}^j \\ \geqslant \frac{\gamma}{2(1-\gamma)} (b^i)' (\sigma\sigma')^{-1} b^i \underline{P}^i g(\underline{P}^i) + (1-\gamma)\underline{P}^{-\frac{\gamma}{1-\gamma}} g(\underline{P}^i) - (\rho^i - \gamma r^i)\underline{P}^i g(\underline{P}^i) + \sum_{j=1}^\ell q^{ij}\underline{P}^j \\ \geqslant -a'(1-\gamma)\underline{P}^i g(\underline{P}^i) + (1-\gamma)\underline{P}^{-\frac{\gamma}{1-\gamma}} g(\underline{P}^i) + \sum_{j=1}^\ell q^{ij}\underline{P}^j. \end{split}$$

By Lemma 3.3, we have

$$a_1' = e^{-a'(1-\gamma)T} \leqslant \underline{P}_t^i \leqslant P_t^{k,i}, \quad i \in \mathcal{M}.$$

To give an upper bound, we notice that

$$(\bar{P}^i, \bar{\Lambda}^i) = \left(e^{a'(T-t)} + \frac{\varepsilon^{\gamma}}{a'}(e^{a'(T-t)} - 1), 0\right), \quad i \in \mathcal{M}$$
(3.12)

is the unique solution to the following BSDE system:

$$\begin{cases} d\bar{P}^i = -\left[a'\bar{P}^i + \varepsilon^{\gamma} + \sum_{j=1}^{\ell} q^{ij}\bar{P}^j\right] dt + (\bar{\Lambda}^i)'dW, \\ \bar{P}_T^i = 1, \quad i \in \mathcal{M}. \end{cases}$$

Also, $(0\mathbf{1}_m, \varepsilon) \in \Theta$ under Assumption 2, therefore for $k \ge -\gamma \varepsilon$, we have

$$\begin{split} f^{k,i}(t,P,\Lambda) &\leqslant \sup_{\tilde{P} \in \mathbb{R}, \tilde{\Lambda} \in \mathbb{R}^n} \left[(\varepsilon^{\gamma} - \gamma \tilde{P} \varepsilon) g(\tilde{P}) - k |P - \tilde{P}| - k |\Lambda - \tilde{\Lambda}| \right] \\ &\leqslant \sup_{\tilde{P} \in \mathbb{R}} \left[(\varepsilon^{\gamma} - \gamma \tilde{P} \varepsilon) g(\tilde{P}) + \gamma \varepsilon |P - \tilde{P}| \right] \\ &\leqslant \sup_{\tilde{P} \in \mathbb{R}} \left[(\varepsilon^{\gamma} - \gamma \tilde{P} \varepsilon) g(\tilde{P}) + \gamma \varepsilon (|\tilde{P}| - |P|) \right] \\ &\leqslant \varepsilon^{\gamma} - \gamma \varepsilon |P|, \end{split}$$

where the third and fourth inequalities are due to that $\gamma < 0$ and $|xg(x)| \leq |x|$ for all $x \in \mathbb{R}$. Hence, using (3.10) and $\bar{P}^i > 0$, we get

$$f^{k,i}(t,\bar{P}^i,\bar{\Lambda}^i) - (\rho^i - \gamma r^i)\bar{P}^i g(\bar{P}^i) + \sum_{j=1}^{\ell} q^{ij} \bar{P}^j \leqslant \varepsilon^{\gamma} - \gamma \varepsilon |\bar{P}^i| - (\rho^i - \gamma r^i)\bar{P}^i g(\bar{P}^i) + \sum_{j=1}^{\ell} q^{ij} \bar{P}^j \leqslant a' \bar{P}^i + \varepsilon^{\gamma} + \sum_{j=1}^{\ell} q^{ij} \bar{P}^j.$$

By Lemma 3.3 again, we obtain the upper bound

$$P_t^{k,i} \leqslant \bar{P}_t^i \leqslant e^{a'T} + \frac{\varepsilon^{\gamma}}{a'}(e^{a'T} - 1) = a_2', \ t \in [0, T], \quad i \in \mathcal{M}.$$

Based on the above two theorems, we can provide the complete answer to problem (2.2).

Theorem 3.8 Suppose $\gamma \in (-\infty,0) \cup (0,1)$ and Assumptions 1 and 2 hold. Let $(P^i,\Lambda^i)_{i\in\mathcal{M}}$ be the unique solution to (3.1). Then the value function of the optimization problem (2.2) is given by

$$V(x, i_0) = \frac{1}{\gamma} x^{\gamma} P_0^{i_0},$$

and $(\hat{\pi}, \hat{c}) \in \mathcal{U}$ is an optimal investment-consumption pair if and only if

$$(\hat{\pi}_t, \hat{c}_t) \in \underset{(\pi, c) \in \Theta}{\operatorname{argmax}} \left[-\frac{1-\gamma}{2} |\pi' \sigma_t^{\alpha_t}|^2 P_t^{\alpha_t} + \pi' (P_t^{\alpha_t} b_t^{\alpha_t} + \sigma_t^{\alpha_t} \Lambda_t^{\alpha_t}) + \frac{c^{\gamma}}{\gamma} - P_t^{\alpha_t} c \right].$$

Remark 3.9 Let $G: \Omega \times [0,T] \times \mathbb{R}^m \times \mathbb{R}_+ \to \mathbb{R}$ be a $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^m) \otimes \mathcal{B}(\mathbb{R}_+)$ -measurable map, where \mathcal{P} is the σ -field of predictable sets of $\Omega \times [0,T]$, and $\mathcal{B}(\mathbb{R}^m)$ ($\mathcal{B}(\mathbb{R}_+)$) is the Borelian σ -algebra on \mathbb{R}^m (\mathbb{R}_+). By

$$\underset{(\pi,c)\in\Theta}{\operatorname{argmax}}\left[G(\omega,t,\pi,c)\right],$$

we denote the set of all predictable processes valued in Θ which attain the essential supremum of G w.r.t $(\pi,c) \in \Theta$. From a measurable selection theorem (see e.g. [11, Lemma 11], [1, Corollary 18.14] or [7, Proposition 2.4]), the set is not empty. If Θ is further convex, then $\arg\max_{(\pi,c)\in\Theta} \left[G(t,\omega,\pi,c) \right]$ degenerates to a singleton set.

Remark 3.10 If $\Theta = \mathbb{R}^n \times \mathbb{R}_+$, then

$$\hat{\pi} = \frac{1}{1 - \gamma} (\sigma \sigma')^{-1} (b + \sigma \frac{\Lambda}{P}), \quad \hat{c} = P^{\frac{1}{\gamma - 1}}.$$

Proof: For any $(\pi, c) \in \mathcal{U}$, applying Itô's formula to $\frac{1}{\gamma}e^{-\int_0^t \rho_s^{\alpha_s} ds} X_t^{\gamma} P_t^{\alpha_t}$, (here we use Itô's formula for Markovian chain; please refer to [12, Lemma 4.3]), we have

$$\begin{split} &\frac{1}{\gamma}e^{-\int_0^t\rho_s^{\alpha_s}ds}X_t^{\gamma}P_t^{\alpha_t} + \int_0^t\frac{1}{\gamma}e^{-\int_0^s\rho_u^{\alpha_u}du}X_s^{\gamma}c_s^{\gamma}ds\\ &= \frac{1}{\gamma}x^{\gamma}P_0^{i_0} + \int_0^te^{-\int_0^s\rho_u^{\alpha_u}du}X_s^{\gamma}\Big[\frac{1}{\gamma}c^{\gamma} - cP - \frac{1-\gamma}{2}P|\pi'\sigma|^2 + \pi'(Pb + \sigma\Lambda) - \frac{f}{\gamma}\Big]ds\\ &+ \int_0^t\frac{1}{\gamma}e^{-\int_0^s\rho_u^{\alpha_u}du}X_s^{\gamma}(\Lambda' + \gamma P\pi'\sigma)dW\\ &+ \int_0^t\frac{1}{\gamma}e^{-\int_0^s\rho_u^{\alpha_u}du}X_s^{\gamma}\sum_{j,j'\in\mathcal{M}}(P^j - P^{j'})I_{\{\alpha_{s-}=j'\}}d\tilde{N}_s^{j'j}, \end{split}$$

where $(N^{j'j})_{j'j\in\mathcal{M}}$ are independent Poisson processes each with intensity $q^{j'j}$, and $\tilde{N}_t^{j'j} = N_t^{j'j} - q^{j'j}t$, $t \ge 0$ are the corresponding compensated Poisson martingales under the filtration \mathcal{F} . And we drop the superscript α_s in the above integral.

Noting that for any $(\pi, c) \in \mathcal{U}$, the wealth process X is continuous and strictly positive, hence bounded on [0, T] away from 0, a.s. Therefore, the stochastic integrals in the last equation are local martingales. Hence, there exists an increasing sequence of stopping times $(\tau_n)_{n\in\mathbb{N}}$ satisfying $\lim_{n\to\infty} \tau_n = T$, a.s. such that

$$\begin{split} & \mathbb{E}\Big[\frac{1}{\gamma}e^{-\int_0^{\tau_n}\rho_s^{\alpha_s}ds}X_{\tau_n}^{\gamma}P_{\tau_n}^{\alpha_{\tau_n}} + \int_0^{\tau_n}\frac{1}{\gamma}e^{-\int_0^s\rho_u^{\alpha_u}du}X_s^{\gamma}c_s^{\gamma}ds\Big] \\ & = \frac{1}{\gamma}x^{\gamma}P_0^{i_0} + \mathbb{E}\int_0^{\tau_n}e^{-\int_0^s\rho_u^{\alpha_u}du}X_s^{\gamma}\Big[-\frac{1-\gamma}{2}P|\pi'\sigma|^2 + \pi'(Pb+\sigma\Lambda) + \frac{1}{\gamma}c^{\gamma} - cP - \frac{f}{\gamma}\Big]ds \\ & \leqslant \frac{1}{\gamma}x^{\gamma}P_0^{i_0}, \end{split}$$

thanks to the definition of f.

Case $\gamma \in (0,1)$: In this case, $\frac{1}{\gamma}e^{-\int_0^{\tau_n}\rho_s^{\alpha_s}ds}X_{\tau_n}^{\gamma}P_{\tau_n}^{\alpha_{\tau_n}} + \int_0^{\tau_n}\frac{1}{\gamma}e^{-\int_0^s\rho_u^{\alpha_u}du}X_s^{\gamma}c_s^{\gamma}ds$ is bounded from below by 0. Passing to limit and applying Fatou's lemma in the above inequality yields

$$\mathbb{E}\left[\frac{1}{\gamma}e^{-\int_0^T \rho_s^{\alpha_s} ds} X_T^{\gamma} + \int_0^T \frac{1}{\gamma}e^{-\int_0^s \rho_u^{\alpha_u} du} X_s^{\gamma} c_s^{\gamma} ds\right] \leqslant \frac{1}{\gamma} x^{\gamma} P_0^{i_0},\tag{3.13}$$

by virtue of the terminal condition $P_T^{\alpha_T} = 1$. And the inequality (3.13) becomes an equality for some $(\hat{\pi}, \hat{c})$ if and only if

$$(\hat{\pi}, \hat{c}) \in \underset{(\pi, c) \in \Theta}{\operatorname{argmax}} \left[-\frac{1 - \gamma}{2} |\pi' \sigma^{\alpha_t}|^2 P + \pi' (P^{\alpha_t} b^{\alpha_t} + \sigma^{\alpha_t} \Lambda^{\alpha_t}) + \frac{c^{\gamma}}{\gamma} - P^{\alpha_t} c \right].$$

It remains to show that $(\hat{\pi}, \hat{c}) \in \mathcal{U}$. Recall that the unique solution of (3.1) $(P^i, \Lambda^i) \in L^{\infty}_{\mathcal{F}^W}(0, T; \mathbb{R}) \times L^{2, \text{BMO}}_{\mathcal{F}^W}(0, T; \mathbb{R}^n)$ for all $i \in \mathcal{M}$. From (3.7), $|\hat{\pi}| \leq K(1 + |Z^{\alpha_t}|)$, hence $\hat{\pi} \in L^2_{\mathcal{F}}(0, T; \mathbb{R}^m)$. Because P is uniformly positive and bounded, we have

$$-\frac{1-\gamma}{2}|\pi'\sigma^{i}|^{2}P + \pi'(Pb^{i} + \sigma^{i}\Lambda) + \frac{c^{\gamma}}{\gamma} - Pc$$

$$\leq \operatorname{ess\,sup}_{\pi \in \mathbb{R}^{m}} \left[-\frac{1-\gamma}{2}|\pi'\sigma^{i}|^{2}P + \pi'(Pb^{i} + \sigma^{i}\Lambda) \right] + \frac{c^{\gamma}}{\gamma} - Pc$$

$$= \frac{1}{2} \frac{\gamma}{1-\gamma} \frac{1}{P} (Pb + \sigma\Lambda)'(\sigma\sigma')^{-1} (Pb + \sigma\Lambda) + \frac{c^{\gamma}}{\gamma} - Pc$$

$$\leq K_{1}(1+|\Lambda|^{2}) + \frac{c^{\gamma}}{\gamma} - K_{1}c$$

$$< 0,$$

if $c > K(1+|\Lambda|^2)$ for sufficient large K > 0. Hence $\hat{c} \leq K(1+|\Lambda|^2)$ so that $\int_0^T \hat{c}dt < \infty$, a.s. This implies $(\hat{\pi}, \hat{c}) \in \mathcal{U}$.

Case $\gamma < 0$: For any admissible $(\pi, c) \in \mathcal{U}$, $(X^{\gamma})_{0 \leqslant t \leqslant T}$ belongs to class (D) and $\mathbb{E}\left[\int_0^T (cX)^{\gamma} dt\right] < \infty$. So by dominated convergence theorem, we also have (3.13). By a similar argument for the previous case $\gamma \in (0, 1)$, we can prove $\hat{\pi} \in L^2_{\mathcal{F}}(0, T; \mathbb{R}^m)$ and $\int_0^T \hat{c} dt < \infty$, a.s. It remains to prove $(\hat{X}^{\gamma}_t)_{0 \leqslant t \leqslant T}$ belongs to class (D) and $\mathbb{E}\left[\int_0^T (\hat{c}\hat{X})^{\gamma} dt\right] < \infty$.

Since ρ and P are bounded, and

$$\mathbb{E}\left[e^{-\int_{0}^{\tau_{n}}\rho_{s}^{\alpha_{s}}ds}\hat{X}_{\tau_{n}}^{\gamma}P_{\tau_{n}}^{\alpha_{\tau_{n}}} + \int_{0}^{\tau_{n}}e^{-\int_{0}^{s}\rho_{u}^{\alpha_{u}}du}\hat{X}_{s}^{\gamma}\hat{c}_{s}^{\gamma}ds\right] = x^{\gamma}P_{0}^{i_{0}},$$

Fatou's Lemma yields

$$\mathbb{E}\left[\hat{X}_T^{\gamma} + \int_0^T (\hat{c}\hat{X})^{\gamma} dt\right] < \infty. \tag{3.14}$$

Denote $\theta^{\alpha_t} := (\sigma^{\alpha_t})'(\sigma^{\alpha_t}(\sigma^{\alpha_t})')^{-1}$ for all $i \in \mathcal{M}$. By Girsanov's theorem,

$$W_t^{\mathbb{Q}} := W_t + \int_0^t \theta_s^{\alpha_s} ds$$

is a Brownian motion under the measure $\mathbb Q$ defined by

$$\frac{d\mathbb{Q}}{d\mathbb{P}} = \mathcal{E}\Big(-\int_0^T \theta_s^{\alpha_s} dW_s\Big).$$

Under the strategy $(\hat{\pi}, \hat{c})$, we obtain from (2.1) that

$$\hat{X}_t = xe^{\int_0^t (r-\hat{c})ds} \mathcal{E}\Big(\int_0^t \hat{\pi}' \sigma dW^{\mathbb{Q}}\Big).$$

Denote

$$J_t := e^{-\int_0^t \hat{c}ds} \mathcal{E}\Big(\int_0^t \hat{\pi}' \sigma dW^{\mathbb{Q}}\Big).$$

Thus it is sufficient to prove $(J_t^{\gamma})_{0 \leq t \leq T}$ belongs to class (D), as r is bounded.

It can be seen from (3.14) that $\mathbb{E}[J_T^{\gamma}] < \infty$. One obtains for every stopping time $\tau \leqslant T$,

$$J_{\tau}^{\frac{\gamma}{2}} = \left\{ \mathbb{E}^{\mathbb{Q}} \left[e^{-\int_{0}^{\tau} \hat{c}ds} \mathcal{E} \left(\int_{0}^{T} \hat{\pi}' \sigma dW^{\mathbb{Q}} \right) \mid \mathcal{F}_{\tau} \right] \right\}^{\frac{\gamma}{2}}$$

$$\leq \left\{ \mathbb{E}^{\mathbb{Q}} \left[e^{-\int_{0}^{T} \hat{c}ds} \mathcal{E} \left(\int_{0}^{T} \hat{\pi}' \sigma dW^{\mathbb{Q}} \right) \mid \mathcal{F}_{\tau} \right] \right\}^{\frac{\gamma}{2}} = \left\{ \mathbb{E}^{\mathbb{Q}} \left[J_{T} \mid \mathcal{F}_{\tau} \right] \right\}^{\frac{\gamma}{2}} \leq \mathbb{E}^{\mathbb{Q}} \left[J_{T}^{\frac{\gamma}{2}} \mid \mathcal{F}_{\tau} \right], \qquad (3.15)$$

where we used that $\hat{c} > 0$ and $\gamma < 0$ in the first inequality, and Jensen's inequality in the second one. By the Cauchy-Schwarz inequality and the boundedness of θ ,

$$\mathbb{E}^{\mathbb{Q}}\left[J_{T}^{\frac{\gamma}{2}} \mid \mathcal{F}_{\tau}\right] = \mathbb{E}\left[J_{T}^{\frac{\gamma}{2}}\mathcal{E}\left(-\int_{\tau}^{T}\theta_{s}^{\alpha_{s}}dW_{s}\right) \mid \mathcal{F}_{\tau}\right]$$

$$\leq \left\{\mathbb{E}\left[J_{T}^{\gamma} \mid \mathcal{F}_{\tau}\right]\right\}^{\frac{1}{2}}\left\{\mathbb{E}\left[\mathcal{E}\left(-\int_{\tau}^{T}\theta_{s}^{\alpha_{s}}dW_{s}\right)^{2} \mid \mathcal{F}_{\tau}\right]\right\}^{\frac{1}{2}} \leq K\left\{\mathbb{E}\left[J_{T}^{\gamma} \mid \mathcal{F}_{\tau}\right]\right\}^{\frac{1}{2}}. \quad (3.16)$$

Then (3.15) and (3.16) imply that $(J_t^{\gamma})_{0 \leq t \leq T}$ belongs to class (D).

4 Logarithmic utility

Let us now turn to the logarithmic utility $U(x) = \ln x$, x > 0. The admissible set is now defined as

$$\mathcal{U} = \left\{ (\pi, c) \mid \pi \in L^2_{\mathcal{F}}(0, T; \mathbb{R}^m), \ c \in L^1_{\mathcal{F}}(0, T; \mathbb{R}_+), \ (\pi_t(\omega), c_t(\omega)) \in \Theta, \ a.e., \ a.s, \right.$$

$$\left. (\ln X_t)_{0 \leqslant t \leqslant T} \text{ and } \left(\int_0^t \ln(c_s X_s) ds \right)_{0 \leqslant t \leqslant T} \text{ belong to class (D)} \right\}.$$

Clearly \mathcal{U} would be empty if c could not take positive value. Hence, it is reasonable to put the following assumption.

Assumption 3 There exists $\varepsilon > 0$ such that $(0\mathbf{1}_m, \varepsilon) \in \Theta$.

To solve problem (2.2), we introduce the following BSDE system:

$$\begin{cases} dh_t^i = -(1 - \rho^i h_t^i + \sum_{j=1}^{\ell} q^{ij} h^j) dt + (\eta^i)' dW, \\ h_T^i = 1, \quad i \in \mathcal{M}. \end{cases}$$
(4.1)

This is a linear BSDE system with bounded coefficients, so it has a unique solution $(h^i, \eta^i)_{i \in \mathcal{M}}$ such that $(h^i, \eta^i) \in L^{\infty}_{\mathcal{F}^W}(0, T; \mathbb{R}) \times L^{2, \, \text{BMO}}_{\mathcal{F}^W}(0, T; \mathbb{R}^n)$.

We claim that h^i is uniformly positive. Indeed, under Assumption 1, there exists a constant k > 0 such that $-\rho^i \ge -k$ for all $i \in \mathcal{M}$. Notice that

$$\left(e^{-k(T-t)} + \frac{1 - e^{-k(T-t)}}{k}, 0\right)$$

is a solution to the following BSDE:

$$\begin{cases} dh_t = -(1 - kh_t)dt + \eta_t' dW, \\ h_T = 1, \end{cases}$$

From Lemma 3.3, we have $h_t^i \geqslant e^{-k(T-t)} + \frac{1-e^{-k(T-t)}}{k} \geqslant e^{-kT}$.

For the unique solution $(h^i, \eta^i)_{i \in \mathcal{M}}$ of (4.1), consider the following BSDE system:

$$\begin{cases} dP^{i} = -\left[f^{i} - \rho^{i}P^{i} + r^{i}h^{i} + \sum_{j=1}^{\ell} q^{ij}P^{j}\right]dt + (\Lambda^{i})'dW \\ P_{T}^{i} = 0, \quad i \in \mathcal{M}, \end{cases}$$

$$(4.2)$$

where

$$f^{i} := \underset{(\pi,c) \in \Theta}{\text{ess sup}} \left[-\frac{1}{2} h^{i} |\pi' \sigma^{i}|^{2} + \pi' (h^{i} b^{i} + \sigma^{i} \eta^{i}) + \ln c - h^{i} c \right].$$

As $(0\mathbf{1}_m, \varepsilon) \in \Theta$, and $h^i \in L^{\infty}_{\mathcal{F}^W}(0, T; \mathbb{R})$, we have

$$f^{i} \geqslant \ln \varepsilon - h^{i} \varepsilon. \tag{4.3}$$

On the other hand,

$$f^{i} \leqslant \underset{(\pi,c) \in \mathbb{R}^{m} \times \mathbb{R}_{+}}{\operatorname{ess sup}} \left[-\frac{1}{2} h^{i} |\pi' \sigma^{i}|^{2} + \pi' (h^{i} b^{i} + \sigma^{i} \eta^{i}) + \ln c - h^{i} c \right]$$
$$= \frac{1}{2h^{i}} (h^{i} b^{i} + \sigma^{i} \eta^{i})' (\sigma^{i} (\sigma^{i})')^{-1} (h^{i} b^{i} + \sigma^{i} \eta^{i}) - 1 - \ln h^{i}.$$

Thus

$$|f^i| \leqslant K(1 + |\eta^i|^2).$$

Whence (4.2) is an ℓ -dimensional linear BSDE system with unbounded coefficients. By [15, Theorem 3.6], the system of (4.2) admits a unique solution $(P^i, \Lambda^i)_{i \in \mathcal{M}}$ such that

$$(P^i, \Lambda^i) \in L^{\infty}_{\mathcal{F}^W}(0, T; \mathbb{R}) \times L^{2, \, \text{BMO}}_{\mathcal{F}^W}(0, T; \mathbb{R}^n), \text{ for all } i \in \mathcal{M}.$$

Remark 4.1 Set $(Y^i, Z^i) = (-\frac{P^i}{h^i}, \frac{P^i}{(h^i)^2} \eta^i - \frac{1}{h^i} \Lambda^i)$, then (Y^i, Z^i) satisfies

$$\begin{cases}
dY^{i} = -\left\{ \underset{(\pi,c)\in\Theta}{\text{ess inf}} \left[\frac{1}{2} |\pi'\sigma^{i}|^{2} - \pi'(b^{i} + \frac{1}{h^{i}}\sigma^{i}\eta^{i}) - \frac{\ln c}{h^{i}} + c \right] - \frac{1}{h^{i}} (1 + \sum_{j=1}^{\ell} q^{ij}h^{j})Y^{i} \right. \\
\left. + \frac{1}{h^{i}} (\eta^{i})'Z^{i} - r^{i} + \frac{1}{h^{i}} \sum_{j=1}^{\ell} q^{ij}h^{j}Y^{j} \right\} dt + (Z^{i})'dW, \\
Y_{T}^{i} = 0, \quad i \in \mathcal{M}.
\end{cases} \tag{4.4}$$

If there is no regime switching and the ρ is a constant, then $\eta^i \equiv 0$ and (4.4) degenerates to (4.6) in [4].

The following lemma can be found in Page 26 of [20].

Lemma 4.2 Suppose $\phi \in L_{\mathcal{F}^W}^{2, \text{BMO}}(0, T; \mathbb{R}^n)$. Then

$$\mathbb{E}\left[\left(\int_0^T |\phi_s|^2 ds\right)^j\right] < \infty, \quad \text{for any } j > 0.$$

Theorem 4.3 Suppose Assumptions 1 and 3 hold. Let $(h^i, \eta^i)_{i \in \mathcal{M}}$ and $(Y^i, Z^i)_{i \in \mathcal{M}}$ be the unique solutions to (4.1) and (4.4), respectively. Then the value function of the optimization problem (2.2) is given by

$$V(x, i_0) = h_0^{i_0} \ln x + P_0^{i_0},$$

and $(\hat{\pi}, \hat{c}) \in \mathcal{U}$ is an optimal consumption-investment pair if and only if

$$(\hat{\pi}_t, \hat{c}_t) \in \operatorname*{argmax}_{(\pi, c) \in \Theta} \left[-\frac{1}{2} h_t^{\alpha_t} |\pi' \sigma_t^{\alpha_t}|^2 + \pi' (h_t^{\alpha_t} b_t^{\alpha_t} + \sigma_t^{\alpha_t} \eta_t^{\alpha_t}) + \ln c - h_t^{\alpha_t} c \right].$$

Remark 4.4 If $\Theta = \mathbb{R}^m \times \mathbb{R}_+$, then the optimal investment-consumption pair is uniquely given by

$$\hat{\pi}_t = (\sigma^{\alpha_t}(\sigma^{\alpha_t})')^{-1}(b^{\alpha_t} + \frac{1}{h^{\alpha_t}}\sigma^{\alpha_t}\eta^{\alpha_t}), \quad \hat{c}_t = \frac{1}{h^{\alpha_t}}.$$

Proof: For any $(\pi, c) \in \mathcal{U}$, applying Itô's formula to $e^{-\int_0^t \rho_s^{\alpha_s} ds} (h_t^{\alpha_t} \ln X_t + P_t^{\alpha_t})$, we have

$$\begin{split} &e^{-\int_{0}^{t}\rho_{s}^{\alpha s}ds}(h_{t}^{\alpha_{t}}\ln X_{t}+P_{t}^{\alpha_{t}})+\int_{0}^{t}e^{-\int_{0}^{s}\rho_{u}^{\alpha u}du}\ln(c_{s}X_{s})ds\\ &=h_{0}^{i_{0}}\ln x+P_{0}^{i_{0}}+\int_{0}^{t}e^{-\int_{0}^{s}\rho_{u}^{\alpha u}du}\Big[\ln c-ch-\frac{1}{2}|\pi'\sigma|^{2}+\pi'(h\eta+\sigma\eta)-f^{\alpha_{s}}\Big]ds\\ &+\int_{0}^{t}e^{-\int_{0}^{t}\rho_{s}^{\alpha s}ds}\Big[h\sigma'\pi+\eta\ln X+\Lambda\Big]'dW\\ &+\int_{0}^{t}e^{-\int_{0}^{t}\rho_{s}^{\alpha s}ds}\sum_{j,j'\in\mathcal{M}}[(h^{j}\ln X+P^{j})-(h^{j'}\ln X+P^{j'})]I_{\{\alpha_{s-}=j'\}}d\tilde{N}_{s}^{j'j}. \end{split}$$

Then by the definition of f^i , the process

$$e^{-\int_0^t \rho_s^{\alpha_s} ds} (h_t^{\alpha_t} \ln X_t + P_t^{\alpha_t}) + \int_0^t e^{-\int_0^s \rho_u^{\alpha_u} du} \ln(c_s X_s) ds$$

is a local supermartingale. Hence, there exists a sequence of increasing stopping times $(\tau_n)_{n\in\mathbb{N}}$ satisfying $\lim_{n\to\infty} \tau_n = T$, a.s. such that

$$\mathbb{E}\Big[e^{-\int_0^{\tau_n} \rho_s^{\alpha_s} ds} (h_{\tau_n}^{\alpha_{\tau_n}} \ln X_{\tau_n} + P_{\tau_n}^{\alpha_{\tau_n}}) + \int_0^{\tau_n} e^{-\int_0^s \rho_u^{\alpha_u} du} \ln(c_s X_s) ds\Big] \leqslant h_0^{i_0} \ln x + P_0^{i_0}.$$

Recalling the admissibility of (π, c) , we deduce from the dominated convergence theorem that

$$\mathbb{E}\left[e^{-\int_{0}^{T}\rho_{s}^{\alpha_{s}}ds}(h_{T}^{\alpha_{T}}\ln X_{T}+P_{T}^{\alpha_{T}})+\int_{0}^{T}e^{-\int_{0}^{s}\rho_{u}^{\alpha_{u}}du}\ln(c_{s}X_{s})ds\right]\leqslant h_{0}^{i_{0}}\ln x+P_{0}^{i_{0}},\tag{4.5}$$

with the equality holds for some $(\hat{\pi}, \hat{c})$ if and only if

$$(\hat{\pi}_t, \hat{c}_t) \in \operatorname*{argmax}_{(\pi, c) \in \Theta} \left[-\frac{1}{2} h^i |\pi' \sigma^i|^2 + \pi' (h^i b^i + \sigma^i \eta^i) + \ln c - h^i c \right].$$

It remains to prove $(\hat{\pi}, \hat{c}) \in \mathcal{U}$ for such a pair. ⁴

From (4.3) and $h^i \in L^{\infty}_{\mathcal{F}^W}(0,T;\mathbb{R})$, $f^i \geqslant -K_1$ for some $K_1 > 0$. On the other hand, Assumption 1 and $h^i \geqslant e^{-kT}$ imply

$$-\frac{1}{2}h^{i}|\pi'\sigma^{i}|^{2} + \pi'(h^{i}b^{i} + \sigma^{i}\eta^{i}) + \ln c - h^{i}c$$

$$\leq -\frac{1}{2}h^{i}|\pi'\sigma^{i}|^{2} + \pi'(h^{i}b^{i} + \sigma^{i}\eta^{i}) + \underset{c \in \mathbb{R}_{+}}{\operatorname{ess sup}}(\ln c - h^{i}c)$$

$$= -\frac{1}{2}h^{i}|\pi'\sigma^{i}|^{2} + \pi'(h^{i}b^{i} + \sigma^{i}\eta^{i}) - \ln h^{i} - 1$$

$$\leq -K_{2}|\pi|^{2} + K_{3}|\pi|(1 + |\eta^{i}|) + K_{4},$$

for some positive constants K_2 , K_3 , K_4 . Thus

$$-\frac{1}{2}h^{i}|\pi'\sigma^{i}|^{2} + \pi'(h^{i}b^{i} + \sigma^{i}\eta^{i}) + \ln c - h^{i}c < -K_{1} \leqslant f^{i},$$

if $|\pi| > K(1+|\eta^i|)$ with K being sufficient large, which implies

$$|\hat{\pi}| \leqslant K(1+|\eta^i|). \tag{4.6}$$

Similarly, since h^i is uniformly positive, we have for some $K_5, K_6 > 0$,

$$-\frac{1}{2}h^{i}|\pi'\sigma^{i}|^{2} + \pi'(h^{i}b^{i} + \sigma^{i}\eta^{i}) + \ln c - h^{i}c$$

$$\leq \operatorname{ess sup}_{\pi \in \mathbb{R}^{m}} \left[-\frac{1}{2}h^{i}|\pi'\sigma^{i}|^{2} + \pi'(h^{i}b^{i} + \sigma^{i}\eta^{i}) \right] + \ln c - K_{5}c$$

$$= \frac{1}{2h^{i}}(h^{i}b^{i} + \sigma^{i}\eta^{i})'(\sigma^{i}(\sigma^{i})')^{-1}(h^{i}b^{i} + \sigma^{i}\eta^{i}) + \ln c - K_{5}c$$

$$\leq K_{6}(1 + |\eta^{i}|^{2}) + \ln c - K_{5}c$$

$$< -K_{1}$$

$$\leq f^{i},$$

⁴Compared with [4], a new, non-zero term $\eta \in L_{\mathcal{F}W}^{2, \, \text{BMO}}(0, T; \mathbb{R}^n)$ emerges (In [4], ρ is a constant, hence $\eta = 0$). So we need conditions like class (D) to apply the dominated convergence theorem in (4.5). But this brings difficulty in proving the admissibility of $(\hat{\pi}, \hat{c})$.

if $c \ge K(1+|\eta^i|^2)$ or $\ln c \le -K(1+|\eta^i|^2)$ for sufficient large K. Therefore,

$$e^{-K(1+|\eta^i|^2)} \leqslant \hat{c} \leqslant K(1+|\eta^i|^2),$$
 (4.7)

and consequently,

$$|\ln \hat{c}| \le K(1 + |\eta^i|^2).$$
 (4.8)

From (2.1),

$$\ln \hat{X}_t = \ln(x) + \int_0^t (r + \hat{\pi}'b - \hat{c} - \frac{1}{2}|\hat{\pi}'\sigma|^2)ds + \int_0^t \hat{\pi}'\sigma dW.$$

It follows from the boundedness of coefficients and the AM-GM inequality that

$$\sup_{t\leqslant T} |\ln \hat{X}_t| \leqslant K + K \int_0^T (|\hat{c}| + |\hat{\pi}|^2) ds + K \sup_{t\leqslant T} \bigg| \int_0^t \hat{\pi}' \sigma dW \bigg|.$$

Using the AM-GM inequality, the Burkholder-Davis-Gundy inequality, (4.6), (4.7), and Lemma 4.2, we have

$$\mathbb{E}\Big[\sup_{\tau\leqslant T}(\ln\hat{X}_{\tau})^2\Big] \leqslant K + K\mathbb{E}\Big[\Big(\int_0^T (|\hat{c}| + |\hat{\pi}|^2)ds\Big)^2 + \sup_{t\leqslant T} \bigg|\int_0^t \hat{\pi}'\sigma dW\bigg|^2\Big]$$

$$\leqslant K + K\mathbb{E}\Big[\Big(\int_0^T (1 + |\eta^i|^2)ds\Big)^2 + \int_0^T |\hat{\pi}'\sigma|^2 ds\Big]$$

$$\leqslant K + K\mathbb{E}\Big[\Big(\int_0^T (1 + |\eta^i|^2)ds\Big)^2 + \int_0^T (1 + |\eta^i|^2)ds\Big]$$

$$\leqslant K.$$

This shows that $(\ln X_t)_{0 \le t \le T}$ belongs to class (D). By the same argument and using (4.8), we can prove that $\left(\int_0^t \ln(c_s X_s) ds\right)_{0 \le t \le T}$ belongs to class (D) as well.

5 Exponential utility

In this case, we do not consider proportional consumption-investment strategies. Instead, let $\pi_{j,t}$ be the *amount* invest in the jth risky asset, j = 1, ..., m, and let c_t be the *amount* to consume. Then the investor's self-financing wealth process X corresponding to a consumption-investment strategy (π, c) is the unique strong solution of the SDE:

$$\begin{cases} dX_t = (r_t X_t + \pi_t' b_t^{\alpha_t} - c_t) dt + \pi_t' \sigma_t^{\alpha_t} dW_t, \\ X_0 = x, \ \alpha_0 = i_0. \end{cases}$$
 (5.1)

Let Π be a given closed nonempty set in \mathbb{R}^m with $0 \in \Pi$ to represent the constraint set for portfolios. We do not put constraint on consumptions, so $-c_t$ can be interpreted as income if $c_t < 0$. The admissible consumption-investment set is defined as

$$\mathcal{U} = \left\{ (\pi, c) \mid \int_0^T (|\pi_t|^2 + |c_t|) dt < \infty, \ (\pi_t(\omega), c_t(\omega)) \in \Pi \times \mathbb{R}, \ a.e., \ a.s., \right.$$

$$\text{and} \ (e^{-\beta h_t X_t^{\pi,c}})_{0 \leqslant t \leqslant T} \text{ belongs to class (D)} \right\},$$

where the process h_t will be specified case by case in the following subsections.

The investor's problem is to maximize

$$J(x, i_0; \pi, c) := \mathbb{E} \left[\int_0^T e^{-\int_0^t \rho_s^{\alpha_s} ds} U(c_t) dt + e^{-\int_0^T \rho_s^{\alpha_s} ds} U(X_T) \right], \quad \text{s.t. } (\pi, c) \in \mathcal{U},$$
 (5.2)

and determine the value function

$$V(x, i_0) := \sup_{(\pi, c) \in \mathcal{U}} J(x, i_0; \pi, c),$$

where the utility is exponential and given by

$$U(x) = -\exp(-\beta x), \quad x \in \mathbb{R}, \quad \beta > 0.$$

It turns out that whether the interest rate is deterministic or random is crucial, we have to deal with these two cases separately.

5.1 Case I: Deterministic interest rate

Assumption 4 The process r is a deterministic bounded measurable function of t. For all $i \in \mathcal{M}$,

$$\mu^i \in L^{\infty}_{\mathbb{F}^W}(0, T; \mathbb{R}^m), \quad \sigma^i \in L^{\infty}_{\mathbb{F}^W}(0, T; \mathbb{R}^{m \times n}), \quad \rho^i \in L^{\infty}_{\mathbb{F}^W}(0, T; \mathbb{R}),$$

and $\sigma^i(\sigma^i)' \geqslant \delta I_m$ with some constant $\delta > 0$.

Define

$$h_t = \left(e^{-\int_t^T r_s ds} + \int_t^T e^{-\int_t^s r_u du} ds\right)^{-1} > 0.$$
 (5.3)

Then under Assumption 4, h is the unique bounded uniformly positive solution to the following ODE:

$$\begin{cases} dh_t = -h_t(r_t - h_t)dt, \\ h_T = 1. \end{cases}$$

Consider the following BSDE system:

$$\begin{cases}
dP^{i} = -\left\{f^{i}(P^{i}, \Lambda^{i}) - hP^{i} \ln P^{i} - \rho^{i}P^{i} + h(1 - \ln h)P^{i} + \sum_{j=1}^{\ell} q^{ij}P^{j}\right\} dt + (\Lambda^{i})' dW, \\
P_{T}^{i} = 1, \quad i \in \mathcal{M},
\end{cases}$$
(5.4)

where

$$f^i(P,\Lambda) := \beta h \operatorname*{ess\ inf}_{\pi \in \Pi} \Big[\frac{1}{2} \beta h P^i |\pi' \sigma^i|^2 - \pi' (P^i b^i + \sigma^i \Lambda^i) \Big].$$

Thanks to $0 \in \Pi$, we have $f^i \leq 0$.

Solutions to (5.4) are defined as in Definition 3.2.

Theorem 5.1 Under Assumption 4, there is a unique uniformly positive solution $(P^i, \Lambda^i)_{i \in \mathcal{M}}$ to the BSDE (5.4).

Proof: We will only present how to find the uniformly bounds. Details are left to interested readers.

Let $a_1 > 0$, $a_2 > 0$, $a_3 > 0$ be three constants such that

$$-\rho^{i} + h(1 - \ln h) \leqslant a_{1},$$

$$\frac{1}{2}(b^{i})'(\sigma^{i}(\sigma^{i})')^{-1}b^{i} - \rho^{i} + h(1 - \ln h) \geqslant -a_{2},$$

$$h \geqslant a_{3} > 0, \ \forall i \in \mathcal{M}.$$

$$(5.5)$$

Denote $a := e^{a_1 T}$ and $\epsilon := e^{-\frac{a_2}{a_3}(1 - e^{-a_3 T})}$. Note $\epsilon < 1 < a$.

Let $g: \mathbb{R} \to [0,1]$ be a smooth truncation function satisfying g(x) = 0 for $x \in (-\infty, \epsilon/2] \cup [2a, \infty)$, and g(x) = 1 for $x \in [\epsilon, a]$.

For $k \geqslant 1$, $(t, P, \Lambda) \in [0, T] \times \mathbb{R} \times \mathbb{R}^n$, $i \in \mathcal{M}$, define

$$f^{k,i}(t,P,\Lambda) = \sup_{\tilde{P} \in \mathbb{R}, \tilde{\Lambda} \in \mathbb{R}^n} \left[f^i(t,\tilde{P},\tilde{\Lambda}) g(\tilde{P}) - k|P - \tilde{P}| - k|\Lambda - \tilde{\Lambda}| \right].$$

Let $(P^{k,i}, \Lambda^{k,i})_{i \in \mathcal{M}}$ denote the unique solution to the following ℓ -dimensional BSDE system with a Lipschitz generator

$$\begin{cases} dP^{k,i} = -\Big[f^{k,i}(P^{k,i}, \Lambda^{k,i,+}) - hP^{k,i}(\ln P^{k,i})g(P^{k,i}) - \rho^i P^{k,i}g(P^{k,i}) \\ + h(1 - \ln h)P^{k,i}g(P^{k,i}) + \sum_{j=1}^{\ell} q^{ij}P^{k,j}\Big]dt + (\Lambda^{k,i})^{\top}dW, \\ P_T^{k,i} = 1, \quad i \in \mathcal{M}. \end{cases}$$

It is direct to verify that

$$(\underline{P}_t^i, \underline{\Lambda}_t^i) = \left(e^{-\frac{a_2}{a_3}(1 - e^{-a_3(T - t)})}, 0\right), \quad i \in \mathcal{M}$$

$$(5.6)$$

is the unique solution to the following BSDE system:

$$\begin{cases} d\underline{P}^i = -\left[-a_2\underline{P}^i g(\underline{P}^i) - a_3\underline{P}^i (\ln\underline{P}^i) g(\underline{P}^i) + \sum_{j=1}^{\ell} q^{ij}\underline{P}^j\right] dt + (\underline{\Lambda}^i)' dW, \\ \underline{P}_T^i = 1, \quad i \in \mathcal{M}. \end{cases}$$

Notice that $\underline{P}^i \in (0,1)$, and

$$\begin{split} f^{k,i}(t,P,\Lambda) &\geqslant f^i(t,P,\Lambda)g(P) \\ &= \beta h \operatorname*{ess\ inf}_{\pi \in \Pi} \left[\frac{1}{2} \beta h P^i |\pi' \sigma^i|^2 - \pi' (P^i b^i + \sigma^i \Lambda^i) \right] g(P) \\ &\geqslant \beta h \operatorname*{ess\ inf}_{\pi \in \mathbb{R}^m} \left[\frac{1}{2} \beta h P^i |\pi' \sigma^i|^2 - \pi' (P^i b^i + \sigma^i \Lambda^i) \right] g(P) \\ &= \frac{1}{2} (b^i)' (\sigma^i (\sigma^i)')^{-1} b^i P g(P) + (b^i)' (\sigma^i (\sigma^i)')^{-1} \sigma \Lambda g(P) \\ &+ \frac{1}{2P} \Lambda' (\sigma^i)' (\sigma^i (\sigma^i)')^{-1} \sigma^i \Lambda g(P), \end{split}$$

we deduce from (5.5) that

$$f^{k,i}(\underline{P}^{i},\underline{\Lambda}^{i}) - h\underline{P}^{i}(\ln\underline{P}^{i})g(\underline{P}^{i}) - \rho^{i}\underline{P}^{i}g(\underline{P}^{i}) + h(1 - \ln h)\underline{P}^{i}g(\underline{P}^{i}) + \sum_{j=1}^{\ell} q^{ij}\underline{P}^{i}$$

$$\geqslant \frac{1}{2}(b^{i})'(\sigma^{i}(\sigma^{i})')^{-1}b^{i}\underline{P}^{i}g(\underline{P}^{i}) - h\underline{P}^{i}(\ln\underline{P}^{i})g(\underline{P}^{i}) - \rho^{i}\underline{P}^{i}g(\underline{P}^{i}) + h(1 - \ln h)\underline{P}^{i}g(\underline{P}^{i}) + \sum_{j=1}^{\ell} q^{ij}\underline{P}^{i}$$

$$\geqslant -a_{2}\underline{P}^{i}g(\underline{P}^{i}) - a_{3}\underline{P}^{i}(\ln\underline{P}^{i})g(\underline{P}^{i}) + \sum_{j=1}^{\ell} q^{ij}\underline{P}^{j}.$$

By Lemma 3.3, we have

$$P_t^{k,i} \geqslant \underline{P}_t^i \geqslant e^{-\frac{a_2}{a_3}(1 - e^{-a_3 T})} = \epsilon, \quad i \in \mathcal{M}.$$
 (5.7)

This gives a uniform lower bound.

To give a uniform upper bound, we notice

$$(\bar{P}_t^i, \bar{\Lambda}_t^i) = (e^{a_1(T-t)}, 0), \quad i \in \mathcal{M}, \tag{5.8}$$

is a solution to the following BSDE system:

$$\begin{cases} d\bar{P}^i = -\left[a_1\bar{P}^ig(\bar{P}^i) + \sum_{j=1}^{\ell} q^{ij}\bar{P}^j\right]dt + (\bar{\Lambda}^i)'dW, \\ \bar{P}_T^i = 1, \quad i \in \mathcal{M}. \end{cases}$$

Since $0 \in \Pi$, we have $f^i(t, P, \Lambda) \leq 0$, and hence $f^{k,i}(t, P, \Lambda) \leq 0$. Notice that $\bar{P}^i_t \geqslant 1$, thus we have

$$f^{k,i}(\bar{P}^i, \bar{\Lambda}^i) - h\bar{P}^i(\ln \bar{P}^i)g(\bar{P}^i) - \rho^i\bar{P}^ig(\bar{P}^i) + h(1 - \ln h)\bar{P}^ig(\bar{P}^i) + \sum_{j=1}^{\ell} q^{ij}\bar{P}^j$$

$$\leq a_1\bar{P}^ig(\bar{P}^i) + \sum_{j=1}^{\ell} q^{ij}\bar{P}^j.$$

By Lemma 3.3, we have

$$P_t^{k,i} \leqslant \bar{P}_t^i \leqslant e^{a_1 T} = a, \quad i \in \mathcal{M}. \tag{5.9}$$

This gives a uniform upper bound.

Set $Y^i = -\frac{1}{\beta} \ln P^i$, $Z^i = -\frac{\Lambda^i}{\beta P^i}$, then (Y^i, Z^i) is the unique solution to

$$\begin{cases}
dY^{i} = -\left[F^{i}(Z^{i}) - hY^{i} - \frac{\beta}{2}|Z^{i}|^{2} + \frac{\rho^{i}}{\beta} - \frac{h}{\beta}(1 - \ln h) \\
-\frac{1}{\beta}\sum_{j=1}^{\ell} q^{ij} \left(e^{-\beta(Y^{j} - Y^{i})} - 1\right)\right] dt + (Z^{i})' dW, \\
Y_{T}^{i} = 0, \quad i \in \mathcal{M},
\end{cases} (5.10)$$

where

$$F^{i}(z) := h \operatorname{ess \, sup}_{\pi \in \Pi} \left[-\frac{1}{2} \beta h |\pi' \sigma^{i}|^{2} + \pi' (b^{i} - \beta \sigma^{i} z) \right], \quad z \in \mathbb{R}^{n}.$$

From above, we immediately have

Theorem 5.2 Under Assumption 4, there is a unique solution $(Y^i, Z^i)_{i \in \mathcal{M}}$ to the BSDE (5.10) such that $(Y^i, Z^i) \in L^{\infty}_{\mathcal{F}W}(0, T; \mathbb{R}) \times L^{2, \text{BMO}}_{\mathcal{F}W}(0, T; \mathbb{R}^n)$ for all $i \in \mathcal{M}$.

Now we can present the solution to problem (5.2).

Theorem 5.3 Suppose Assumption 4 holds. Let h be defined in (5.3) and $(Y^i, Z^i)_{i \in \mathcal{M}}$ be the unique solution to (5.10). Then the value function of the optimization problem (5.2) is given by

$$V(x, i_0) = -e^{-\beta(h_0 x + Y_0^{i_0})},$$

and the optimal portfolio is

$$\hat{\pi}_t \in \operatorname*{argmax}_{\pi \in \Pi} \left[-\frac{1}{2} \beta h_t |\pi' \sigma_t^{\alpha_t}|^2 + \pi' (b_t^{\alpha_t} - \beta \sigma_t^{\alpha_t} Z_t^{\alpha_t}) \right],$$

the optimal consumption is

$$\hat{c}_t = h_t X_t + Y_t^{\alpha_t} - \frac{1}{\beta} \ln h_t.$$

Proof: For any $(\pi, c) \in \mathcal{U}$, applying Itô's formula to $-e^{-\int_0^t \rho_s^{\alpha_s} ds} e^{-\beta(h_t X_t + Y_t^{\alpha_t})}$, we have

$$-e^{-\int_{0}^{t} \rho_{s}^{\alpha s} ds} e^{-\beta(h_{t}X_{t}+Y_{t}^{\alpha t})} - \int_{0}^{t} e^{-\int_{0}^{s} \rho_{s}^{\alpha u} du} e^{-\beta c_{s}} ds$$

$$= -e^{-\beta(h_{0}x+Y_{0}^{i_{0}})} + \int_{0}^{t} \beta e^{-\int_{0}^{s} \rho_{u}^{\alpha u} du} e^{-\beta(h_{s}X_{s}+Y_{s}^{\alpha s})} \left[-\frac{1}{\beta} e^{\beta(hX+Y)} e^{-\beta c} - hc - \left(\frac{h}{\beta} (\ln h - 1) - h^{2}X - hY \right) + h \left(-\frac{1}{2} \beta h |\pi'\sigma^{i}|^{2} + \pi'(b^{i} - \beta\sigma^{i}z) \right) - F^{i}(Z) \right] ds$$

$$+ \int_{0}^{t} \beta e^{-\int_{0}^{s} \rho_{u}^{\alpha u} du} e^{-\beta(h_{s}X_{s}+Y_{s}^{\alpha s})} (h\pi'\sigma + Z') dW$$

$$+ \int_{0}^{t} e^{-\int_{0}^{s} \rho_{u}^{\alpha u} du} e^{-\beta h_{s}X_{s}} \sum_{i,j' \in \mathcal{M}} (e^{-\beta Y^{j}} - e^{-\beta Y^{i}}) I_{\{\alpha_{s-}=j'\}} d\tilde{N}_{s}^{j'j},$$

where $(N^{j'j})_{j'j\in\mathcal{M}}$ are independent Poisson processes each with intensity $q^{j'j}$, and $\tilde{N}_t^{j'j} = N_t^{j'j} - q^{j'j}t$, $t \ge 0$ are the corresponding compensated Poisson martingales under the filtration \mathcal{F} . And we drop the superscript α_s in the above integral.

Noting that for any $(\pi, c) \in \mathcal{U}$, the wealth process X is continuous, hence bounded on [0, T] a.s. Therefore, the stochastic integrals in the last equation are local martingales. Hence, there exists a sequence of increasing stopping times $(\tau_n)_{n\in\mathbb{N}}$ satisfying $\lim_{n\to\infty} \tau_n = T$, a.s. such that

$$\begin{split} \mathbb{E}\Big[-e^{-\int_{0}^{\tau_{n}} \rho_{s}^{\alpha s} ds} e^{-\beta(h_{\tau_{n}} X_{\tau_{n}} + Y_{\tau_{n}}^{\alpha_{\tau_{n}}})} - \int_{0}^{\tau_{n}} e^{-\int_{0}^{s} \rho_{s}^{\alpha u} du} e^{-\beta c_{s}} ds \Big] \\ = -e^{-\beta(h_{0} x + Y_{0}^{i_{0}})} + \int_{0}^{\tau_{n}} \beta e^{-\int_{0}^{s} \rho_{u}^{\alpha u} du} e^{-\beta(h_{s} X_{s} + Y_{s}^{\alpha_{s}})} \Big[-\frac{1}{\beta} e^{\beta(hX + Y)} e^{-\beta c} - hc \\ - \Big(\frac{h}{\beta} (\ln h - 1) - h^{2} X - hY \Big) + h \Big(-\frac{1}{2} \beta h |\pi' \sigma^{i}|^{2} + \pi' (b^{i} - \beta \sigma^{i} z) \Big) - F^{i}(Z) \Big] ds. \end{split}$$

Because $h, \beta > 0$,

$$\operatorname{ess\,sup}_{c\in\mathbb{R}}\left[-\frac{1}{\beta}e^{\beta(hX+Y)}e^{-\beta c}-hc\right] = \frac{h}{\beta}(\ln h - 1) - h^2X - hY,$$

we obtain from the definition of F^i that

$$\mathbb{E}\bigg[-e^{-\int_{0}^{\tau_{n}}\rho_{s}^{\alpha_{s}}ds}e^{-\beta(h_{\tau_{n}}X_{\tau_{n}}+Y_{\tau_{n}}^{\alpha_{\tau_{n}}})}\bigg] - \mathbb{E}\bigg[\int_{0}^{\tau_{n}}e^{-\int_{0}^{s}\rho_{s}^{\alpha_{u}}du}e^{-\beta c_{s}}ds\bigg] \leqslant -e^{-\beta(h_{0}x+Y_{0}^{i_{0}})},$$

for any $(c, \pi) \in \mathcal{U}$. As $n \to \infty$, the second expectation is convergent by the monotone convergence theorem; while the first expectation, because $(e^{-\beta h_t X_t^{\pi,c}})_{0 \le t \le T}$ belongs to class (D), Y^i , $i \in \mathcal{M}$, are bounded, is also convergent by the dominated convergence theorem. Hence

$$\mathbb{E}\left[-e^{-\int_0^T \rho_s^{\alpha_s} ds} e^{-\beta X_T} - \int_0^T e^{-\int_0^s \rho_s^{\alpha_u} du} e^{-\beta c_s} ds\right] \leqslant -e^{-\beta(h_0 x + Y_0^{i_0})},\tag{5.11}$$

by virtue of the terminal conditions $h_T = 1$ and $Y_T = 0$. Furthermore, the inequality (5.11) becomes an equality for some $(\hat{\pi}, \hat{c})$ if and only if

$$\hat{c}_t = \underset{c \in \mathbb{R}}{\operatorname{argmax}} \left[-\frac{1}{\beta} e^{\beta(hX + Y^{\alpha})} e^{-\beta c} - hc \right] = -\frac{1}{\beta} \ln h + hX + Y^{\alpha},$$

and

$$\hat{\pi}_t \in \operatorname*{argmax}_{\pi \in \Pi} \left[-\frac{1}{2} \beta h |\pi' \sigma^{\alpha}|^2 + \pi' (b^{\alpha} - \beta \sigma^{\alpha} Z^{\alpha}) \right].$$

The admissibility issue $(\hat{\pi}, \hat{c}) \in \mathcal{U}$ is covered in a more general case; please refer to the proof of Theorem 5.8.

5.2 Case II: Random interest rate

Assumption 5 For all $i \in \mathcal{M}$,

$$\rho^i \in L^{\infty}_{\mathbb{F}^W}(0,T;\mathbb{R}).$$

Furthermore, the processes r, μ , σ are independent of the regime $i \in \mathcal{M}$ and such that

$$r \in L^{\infty}_{\mathcal{F}W}(0,T;\mathbb{R}), \ \mu \in L^{\infty}_{\mathbb{R}W}(0,T;\mathbb{R}^m), \ \sigma \in L^{\infty}_{\mathbb{R}W}(0,T;\mathbb{R}^{m \times n}),$$

and $\sigma\sigma' \geqslant \delta I_m$ with some constant $\delta > 0$.

Assumption 6 m = n and $\Pi = \mathbb{R}^m$, i.e. neither the portfolio nor the consumption has constraints.

Under Assumptions 5 and 6, σ is a square invertible matrix.

Consider the following BSDE system:

$$\begin{cases}
dh_t = -[rh - h^2 - \eta' \sigma^{-1} b - \frac{1}{h} |\eta|^2] dt + \eta' dW, \\
h_T = 1.
\end{cases}$$
(5.12)

Theorem 5.4 Under Assumptions 5 and 6, (5.12) admits a unique solution $(h, \eta) \in L^{\infty}_{\mathcal{F}^W}(0, T; \mathbb{R}) \times L^{2, \text{BMO}}_{\mathcal{F}^W}(0, T; \mathbb{R}^n)$ such that $h \geqslant \delta$ for some positive constant δ .

Proof: Let $(p,q) \in L^{\infty}_{\mathcal{F}W}(0,T;\mathbb{R}) \times L^{2, \text{BMO}}_{\mathcal{F}W}(0,T;\mathbb{R}^n)$ denote unique solution to the following linear BSDE system:

$$\begin{cases}
dp_t = -[1 - rp - q'\sigma^{-1}b]dt + q'dW, \\
p_T = 1.
\end{cases}$$
(5.13)

Indeed, p admits the explicit representation:

$$p_t = \widetilde{\mathbb{E}} \left[e^{-\int_t^T r_s ds} + \int_t^T e^{-\int_t^s r_u du} ds \mid \mathcal{F}_t^W \right],$$

where $\widetilde{\mathbb{E}}$ is the expectation with respect to the probability measure $\widetilde{\mathbb{P}}$ defined by

$$\frac{d\widetilde{\mathbb{P}}}{d\mathbb{P}} = \mathcal{E}\bigg(\int_0^T (\sigma^{-1}b)'dW\bigg).$$

It follows from Assumption 5 that $\delta_1 \leqslant p \leqslant \delta_2$ for some $0 < \delta_1 < \delta_2$. Therefore $(h, \eta) := (\frac{1}{p}, -\frac{q}{p^2})$ is well defined and $(h, \eta) \in L^{\infty}_{\mathcal{F}W}(0, T; \mathbb{R}) \times L^{2, \, \mathrm{BMO}}_{\mathcal{F}W}(0, T; \mathbb{R}^n)$ with $h \geqslant \frac{1}{\delta_2} > 0$. It can be directly verified, using Itô's formula, that (h, η) is a solution of (5.12). As for uniqueness, observe that if (h, η) is a solution of (5.12) with $h \geqslant \delta$ for some $\delta > 0$, then $(p, q^i) = (\frac{1}{h}, -\frac{\eta}{h^2})$ is a solution to (5.13). Uniqueness follows from the fact that (5.13) has a unique solution.

Let (h, η) denote the unique solution to (5.12). Consider the following BSDE system:

$$\begin{cases}
dY^{i} = -\left[-hY^{i} - (\sigma^{-1}b + \frac{\eta}{h})'Z^{i} + \frac{1}{2\beta h^{2}}|h\sigma^{-1}b + \eta|^{2} \\
+ \frac{\rho^{i}}{\beta} - \frac{h}{\beta}(1 - \ln h) - \frac{1}{\beta}\sum_{j=1}^{\ell} q^{ij}e^{-\beta(Y^{j} - Y^{i})}\right]dt + (Z^{i})'dW, \\
Y_{T}^{i} = 0, \quad i \in \mathcal{M}.
\end{cases} (5.14)$$

Remark 5.5 Because $\eta \in L_{\mathcal{F}W}^{2, \text{BMO}}(0, T; \mathbb{R}^n)$ is not bounded, we cannot apply [8, Theorem 2.4] to solve (5.14).

Set $(P^i, \Lambda^i) = (\frac{Y^i}{h}, \frac{1}{h}Z^i - \frac{Y^i}{h^2}\eta)$ for all $i \in \mathcal{M}$. Then $(P^i, \Lambda^i)_{i \in \mathcal{M}}$ fulfills the following BSDE system:

$$\begin{cases}
dP^{i} = -\left[-rP^{i} - (\sigma^{-1}b)'\Lambda^{i} + \frac{1}{2\beta h^{3}}|h\sigma^{-1}b + \eta|^{2} \\
+ \frac{\rho^{i}}{\beta h} - \frac{1}{\beta}(1 - \ln h) - \frac{1}{\beta h}\sum_{j=1}^{\ell}q^{ij}e^{-\beta h(P^{j} - P^{i})}\right]dt + (\Lambda^{i})'dW, \\
P_{T}^{i} = 0, \quad i \in \mathcal{M}.
\end{cases} (5.15)$$

Theorem 5.6 Under Assumptions 5 and 6, (5.15) admits a unique solution $(P^i, \Lambda^i)_{i \in \mathcal{M}}$ such that $(P^i, \Lambda^i) \in L^{\infty}_{\mathcal{F}^W}(0, T; \mathbb{R}) \times L^{2, \text{BMO}}_{\mathcal{F}^W}(0, T; \mathbb{R}^n)$ for all $i \in \mathcal{M}$.

Proof: We take the notations in the proof of Theorem 5.4. Under the probability measure $\widetilde{\mathbb{P}}$,

$$\widetilde{W}_t := W_t + \int_0^t \sigma^{-1} b ds,$$

is an *n*-dimensional Brownian motion. Notice that $\eta \in L_{\mathcal{F}W}^{2, \text{BMO}}(0, T; \mathbb{R}^n)$, hence $\int_0^t \eta' d\widetilde{W}_s$ is a BMO martingale under $\widetilde{\mathbb{P}}$.

Let $k_1 > 0$ be a large constant such that, for all $i \in \mathcal{M}$ and $t \in [0, T]$,

$$-r \leqslant k_1, \quad \frac{\rho^i}{\beta h} - \frac{1}{\beta} (1 - \ln h) \geqslant -k_1, \tag{5.16}$$

and

$$\widetilde{\mathbb{E}}\left[\int_{t}^{T} e^{-\int_{t}^{s} r_{u} du} \left(\frac{1}{2\beta h^{3}} |h\sigma^{-1}b + \eta|^{2} + \frac{\rho^{i}}{\beta h} - \frac{1}{\beta} (1 - \ln h) - \frac{1}{\beta h} q^{ii}\right) ds \mid \mathcal{F}_{t}^{W}\right] \leqslant k_{1}.$$

Denote $k := \max\{e^{k_1 T}, k_1\}.$

Define a truncating function $\psi : \mathbb{R} \to \mathbb{R}$ as

$$\psi(y) := \max\{-k, \min\{y, k\}\}.$$

Consider the following truncated system of (5.15),

$$\begin{cases}
dP^{i} = -\left[-rP^{i} - (\sigma^{-1}b)'\Lambda^{i} + \frac{1}{2\beta h^{3}}|h\sigma^{-1}b + \eta|^{2} \\
+ \frac{\rho^{i}}{\beta h} - \frac{1}{\beta}(1 - \ln h) - \frac{1}{\beta h}\sum_{j=1}^{\ell}q^{ij}e^{-\beta h(\psi(P^{j}) - \psi(P^{i}))}\right]dt + (\Lambda^{i})'dW, \\
P_{T}^{i} = 0, \quad i \in \mathcal{M}.
\end{cases} (5.17)$$

The driver is Lipschitz continuous except for $\eta \in L^{2,\,\mathrm{BMO}}_{\mathcal{F}^W}(0,T;\mathbb{R}^n)$ is unbounded. By [6, Theorem 9.3.5], there is a unique solution $(P^{k,i},\Lambda^{k,i})_{i\in\mathcal{M}}$, such that $(P^{k,i},\Lambda^{k,i})\in L^2_{\mathcal{F}^W}(0,T;\mathbb{R})\times L^2_{\mathcal{F}^W}(0,T;\mathbb{R}^n)$ for all $i\in\mathcal{M}$. If we can show $|P^{k,i}|\leqslant k$ and $\Lambda^{k,i}\in L^{2,\,\mathrm{BMO}}_{\mathcal{F}^W}(0,T;\mathbb{R}^n)$, then $\psi(P^{k,i})=P^{k,i}$ and $(P^{k,i},\Lambda^{k,i})_{i\in\mathcal{M}}$ is a solution to (5.15).

We first show that $P^{k,i} \ge -k$. It is direct to verify that

$$(\underline{P},\underline{\Lambda}) = (1 - e^{k_1(T-t)}, 0),$$

is the unique solution to the following BSDE:

$$\begin{cases} d\underline{P} = -(k_1\underline{P} - k_1)dt + \underline{\Lambda}'dW \\ \underline{P}_T = 0. \end{cases}$$
 (5.18)

Notice that $\underline{P} \leqslant 0$ and

$$k_{1}\underline{P} - k_{1} \leqslant -r\underline{P} - (\sigma^{-1}b)'\underline{\Lambda} + \frac{1}{2\beta h^{3}}|h\sigma^{-1}b + \eta|^{2}$$
$$+ \frac{\rho^{i}}{\beta h} - \frac{1}{\beta}(1 - \ln h) - \frac{1}{\beta h}\sum_{j=1}^{\ell}q^{ij}e^{-\beta h(\psi(\underline{P}) - \psi(\underline{P}))},$$

so by Lemma 3.3

$$-k \leqslant -e^{k_1 T} \leqslant \underline{P}_t \leqslant P_t^{k,i}$$
.

We next show that $P^{k,i} \leq k$. From (5.17), we obtain

$$\begin{split} e^{-\int_0^t r_s ds} P_t^{k,i} &= \int_t^T e^{-\int_0^s r_u du} \Big[- (\sigma^{-1}b)' \Lambda^{k,i} + \frac{1}{2\beta h^3} |h\sigma^{-1}b + \eta|^2 \\ &+ \frac{\rho^i}{\beta h} - \frac{1}{\beta} (1 - \ln h) - \frac{1}{\beta h} \sum_{j=1}^\ell q^{ij} e^{-\beta h(\psi(P^{k,j}) - \psi(P^{k,i}))} \Big] ds - \int_t^T (\Lambda^{k,i})' dW \\ &= \int_t^T e^{-\int_0^s r_u du} \Big[\frac{1}{2\beta h^3} |h\sigma^{-1}b + \eta|^2 + \frac{\rho^i}{\beta h} - \frac{1}{\beta} (1 - \ln h) \\ &- \frac{1}{\beta h} \sum_{j=1}^\ell q^{ij} e^{-\beta h(\psi(P^{k,j}) - \psi(P^{k,i}))} \Big] ds - \int_t^T (\Lambda^{k,i})' d\widetilde{W}. \end{split}$$

Taking expectation $\widetilde{\mathbb{E}}$, and noting $q^{ij} \geqslant 0$ for $i \neq j$, we have

$$\begin{split} P_t^{k,i} &= e^{\int_0^t r_s ds} \widetilde{\mathbb{E}} \Bigg[\int_t^T e^{-\int_0^s r_u du} \Big(\frac{1}{2\beta h^3} |h\sigma^{-1}b + \eta|^2 + \frac{\rho^i}{\beta h} - \frac{1}{\beta} (1 - \ln h) \\ &- \frac{1}{\beta h} q^{ii} - \frac{1}{\beta h} \sum_{j \neq i}^\ell q^{ij} e^{-\beta h (\psi(P^{k,j}) - \psi(P^{k,i}))} \Big) ds \, \bigg| \, \mathcal{F}_t^W \Bigg] \\ &\leqslant \widetilde{\mathbb{E}}^i \Bigg[\int_t^T e^{-\int_t^s r_u du} \Big(\frac{1}{2\beta h^3} |h\sigma^{-1}b + \eta|^2 + \frac{\rho^i}{\beta h} - \frac{1}{\beta} (1 - \ln h) - \frac{1}{\beta h} q^{ii} \Big) ds \, \bigg| \, \mathcal{F}_t^W \Bigg] \\ &\leqslant k_1 \leqslant k. \end{split}$$

Therefore $|P^{k,i}| \leq k$. Whence we can drop the superscript k in $(P^{k,i}, \Lambda^{k,i})_{i \in \mathcal{M}}$. Applying Itô's formula to $(P^i)^2$, we have for any stopping time $\tau \leq T$,

$$\begin{split} &(P_{\tau}^{i})^{2} + \mathbb{E}\bigg[\int_{\tau}^{T}|\Lambda^{i}|^{2}ds \ \bigg| \ \mathcal{F}_{\tau}^{W}\bigg] \\ &= \mathbb{E}\bigg[\int_{\tau}^{T}2P^{i}\Big(-rP^{i} - (\sigma^{-1}b)'\Lambda^{i} + \frac{1}{2\beta h^{3}}|h\sigma^{-1}b + \eta|^{2} \\ &\qquad \qquad + \frac{\rho^{i}}{\beta h} - \frac{1}{\beta}(1 - \ln h) - \frac{1}{\beta h}\sum_{j=1}^{\ell}q^{ij}e^{-\beta h(P^{j} - P^{i})}\Big)ds \ \bigg| \ \mathcal{F}_{\tau}^{W}\bigg] \\ &\leqslant \mathbb{E}\bigg[\int_{\tau}^{T}\Big(K + \frac{1}{2}|\Lambda^{i}|^{2} + K|\eta|^{2}\Big)ds \ \bigg| \ \mathcal{F}_{\tau}^{W}\bigg]. \end{split}$$

From which, we obtain

$$\mathbb{E}\left[\int_{\tau}^{T} |\Lambda^{i}|^{2} ds \mid \mathcal{F}_{\tau}^{W}\right] \leqslant K \mathbb{E}\left[\int_{\tau}^{T} \left(1 + |\eta|^{2}\right) ds \mid \mathcal{F}_{\tau}^{W}\right] \leqslant K,$$

therefore $\Lambda^i \in L_{\mathcal{F}^W}^{2, \text{BMO}}(0, T; \mathbb{R}^n)$. The uniqueness comes from [13, Lemma 3.4]. From Theorem 5.4 and Theorem 5.6, we immediately have

Theorem 5.7 Under Assumptions 5 and 6, (5.14) admits a unique solution $(Y^i, Z^i)_{i \in \mathcal{M}}$ such that $(Y^i, Z^i) \in L^{\infty}_{\mathcal{F}W}(0, T; \mathbb{R}) \times L^{2, \text{BMO}}_{\mathcal{F}W}(0, T; \mathbb{R}^n)$ for all $i \in \mathcal{M}$.

Theorem 5.8 Suppose Assumptions 5 and 6 hold. Let (h, η) and $(Y^i, Z^i)_{i \in \mathcal{M}}$ be the unique solutions to (5.12) and (5.14) respectively. Then the value function of the optimization problem (5.2) is given by

$$V(x, i_0) = -e^{-\beta(h_0 x + Y_0^{i_0})},$$

and the optimal portfolio is

$$\hat{\pi}_t = -\frac{1}{\beta h_t^2} (\sigma_t')^{-1} (\beta h_t \eta_t X_t + \beta h_t Z_t^{\alpha_t} - h_t \sigma_t^{-1} b_t - \eta_t), \tag{5.19}$$

the optimal consumption is

$$\hat{c}_t = h_t X_t + Y_t^{\alpha_t} - \frac{1}{\beta} \ln h_t.$$
 (5.20)

Proof: For any $(\pi, c) \in \mathcal{U}$, applying Itô's formula to $-e^{-\int_0^t \rho_s^{\alpha_s} ds} e^{-\beta(h_t X_t + Y_t^{\alpha_t})}$, we obtain

$$\mathbb{E}\Big[-e^{-\int_0^{\tau_n} \rho_s^{\alpha s} ds} e^{-\beta(h_{\tau_n} X_{\tau_n} + Y_{\tau_n}^{\alpha \tau_n})}\Big] - \mathbb{E}\Big[\int_0^{\tau_n} e^{-\int_0^s \rho_s^{\alpha u} du} e^{-\beta c_s} ds\Big] \leqslant -e^{-\beta(h_0 x + Y_0^{i_0})},$$

for a sequence of increasing stopping times $(\tau_n)_{n\in\mathbb{N}}$ satisfying $\lim_{n\to\infty} \tau_n = T$. Same as the deterministic case, the two expectations are both convergence as $n\to\infty$ so that

$$\mathbb{E}\Big[-e^{-\int_{0}^{T}\rho_{s}^{\alpha_{s}}ds}e^{-\beta X_{T}}\Big] - \mathbb{E}\Big[\int_{0}^{T}e^{-\int_{0}^{s}\rho_{s}^{\alpha_{u}}du}e^{-\beta c_{s}}ds\Big] \leqslant -e^{-\beta(h_{0}x+Y_{0}^{i_{0}})},\tag{5.21}$$

thanks to the terminal conditions $h_T = 1$ and $Y_T = 0$. Moreover, the inequality (5.21) becomes an equality for some $(\hat{\pi}, \hat{c})$ if and only if (5.19) and (5.20) hold.

Denote \hat{X} by the corresponding wealth process under the strategy $(\hat{\pi}, \hat{c})$. Then we have

$$\mathbb{E}\Big[-e^{-\int_0^{\tau_n}\rho_s^{\alpha_s}ds}e^{-\beta(h_{\tau_n}^{\alpha\tau_n}\hat{X}_{\tau_n}+Y_{\tau_n}^{\alpha\tau_n})}-\int_0^{\tau_n}e^{-\int_0^s\rho_s^{\alpha_u}du}e^{-\beta\hat{c}_s}ds\Big]=-e^{-\beta(h_0x+Y_0^{i_0})}.$$

Note that $\rho, Y^i, i \in \mathcal{M}$, are bounded processes,

$$\mathbb{E}\Big[e^{-\beta h_{\tau n}^{\alpha \tau_n} \hat{X}_{\tau_n}} + \int_0^{\tau_n} e^{-\beta \hat{c}_s} ds\Big] \leqslant K \mathbb{E}\Big[e^{-\int_0^{\tau_n} \rho_s^{\alpha_s} ds} e^{-\beta (h_{\tau n}^{\alpha \tau_n} \hat{X}_{\tau_n} + Y_{\tau n}^{\alpha \tau_n})} + \int_0^{\tau_n} e^{-\int_0^s \rho_s^{\alpha_u} du} e^{-\beta \hat{c}_s} ds\Big]$$

$$= K e^{-\beta (h_0 x + Y_0^{i_0})}.$$

Applying Fatou's lemma to the above, we obtain

$$\mathbb{E}\left[e^{-\beta\hat{X}_T} + \int_0^T e^{-\beta\hat{c}_s} ds\right] \leqslant K. \tag{5.22}$$

Set $\theta := \frac{1}{h}\sigma^{-1}(hb + \sigma\eta)$, then $\theta \in L_{\mathcal{F}^W}^{2, \text{BMO}}(0, T; \mathbb{R}^n)$ and $\mathcal{E}(\int_0^t \theta_s dW)$ is a uniformly integrable martingale. By Girsanov's theorem,

$$W_t^{\mathbb{Q}} := W_t + \int_0^t \theta_s ds$$

is an *n*-dimensional Brownian motion under the measure \mathbb{Q} defined by

$$\frac{d\mathbb{Q}}{d\mathbb{P}} = \mathcal{E}\Big(-\int_0^T \theta_s dW_s\Big).$$

We need to show that $(e^{-\beta h_t \hat{X}_t})_{0 \leqslant t \leqslant T}$ belongs to class (D). Compared with Section 4 or [4], a new term $\eta \in L_{\mathcal{F}W}^{2, \text{BMO}}(0, T; \mathbb{R}^n)$ emerges (In Section 4 or [4], r is deterministic, hence $\eta = 0$). Applying Itô's formula to $h_t \hat{X}_t$,

$$h_{t}\hat{X}_{t} = h_{0}x + \int_{0}^{t} \left[-\frac{1}{\beta h^{2}} (\beta hZ - h\sigma^{-1}b - \eta)'\sigma^{-1}(hb + \sigma\eta) - h(Y - \frac{1}{\beta}\ln h) \right] ds$$
$$- \int_{0}^{t} \frac{1}{\beta h} (\beta hZ - h\sigma^{-1}b - \eta)' dW$$
$$= h_{0}x + \int_{0}^{t} \left[-h(Y - \frac{1}{\beta}\ln h) \right] ds - \int_{0}^{t} \frac{1}{\beta h} (\beta hZ - h\sigma^{-1}b - \eta)' dW^{\mathbb{Q}}$$

Since $-\frac{1}{\beta h}(\beta hZ - h\sigma^{-1}b - \eta) \in L_{\mathcal{F}}^{2, \text{BMO}}(0, T; \mathbb{R}^n)$, the process

$$L_t := -\int_0^t \frac{1}{\beta h} (\beta h Z - h \sigma^{-1} b - \eta)' dW^{\mathbb{Q}}$$

is a BMO-martingale under \mathbb{Q} . Because $h, Y^i, i \in \mathcal{M}$, are bounded, there exists a positive constant K such that

$$\frac{1}{K}e^{-\beta L_t} \leqslant e^{-\beta h_t \hat{X}_t} \leqslant Ke^{-\beta L_t} \text{ for all } t \in [0, T].$$

Hence our problem reduces to show $(e^{-\beta L_t})_{0 \le t \le T}$ belongs to class (D). By the reverse Hölder inequality, there exists p > 1 such that for all stopping times $\tau \le T$,

$$\left(\mathbb{E}\left[\mathcal{E}\left(-\int_{0}^{T}\theta dW\right)^{p} \mid \mathcal{F}_{\tau}\right]\right)^{\frac{1}{p}} \leqslant K\mathcal{E}\left(-\int_{0}^{\tau}\theta dW\right). \tag{5.23}$$

Let q be the conjugate exponent of p i.e. $\frac{1}{p} + \frac{1}{q} = 1$. Since the function $x \mapsto e^{-\frac{\beta}{q}x}$ is convex, by Jensen's inequality, we have for any stopping time $\tau \leqslant T$,

$$e^{-\frac{\beta}{q}L_{\tau}} \leqslant \mathbb{E}^{\mathbb{Q}} \Big[e^{-\frac{\beta}{q}L_{T}} \mid \mathcal{F}_{\tau} \Big].$$

It follows

$$\begin{split} e^{-\beta L_{\tau}} &\leqslant \left(\mathbb{E}^{\mathbb{Q}} \Big[e^{-\frac{\beta}{q} L_{T}} \ \Big| \ \mathcal{F}_{\tau} \Big] \right)^{q} \\ &= \left(\mathbb{E} \Big[e^{-\frac{\beta}{q} L_{T}} \mathcal{E} \Big(- \int_{0}^{T} \theta dW \Big) \ \Big| \ \mathcal{F}_{\tau} \Big] \right)^{q} \mathcal{E} \Big(- \int_{0}^{\tau} \theta dW \Big)^{-q} \\ &\leqslant \mathbb{E} \Big[e^{-\beta L_{T}} \ \Big| \ \mathcal{F}_{\tau} \Big] \mathbb{E} \Big[\mathcal{E} \Big(- \int_{0}^{T} \theta dW \Big)^{p} \ \Big| \ \mathcal{F}_{\tau} \Big]^{\frac{q}{p}} \mathcal{E} \Big(- \int_{0}^{\tau} \theta dW \Big)^{-q} \\ &\leqslant K \mathbb{E} [e^{-\beta L_{T}} \ \Big| \ \mathcal{F}_{\tau}], \end{split}$$

where we use Hölder's inequality in the second inequality, (5.23) in the last inequality. This shows that $(e^{-\beta h_t \hat{X}_t})_{0 \leqslant t \leqslant T}$ belongs to class (D) noting (5.22), hence the admissibility of $(\hat{\pi}, \hat{c})$.

6 Concluding remarks

In this paper, we studied optimal consumption-investment problems in a market with regime switching and random coefficients with coupled constraints. Explicit solutions in term of the solutions of BSDE systems are given for the cases of power, logarithmic and exponential utilities. Some of these BSDE systems are new in the literature. Their solvability consists the main mathematical contributions and is interesting in its own from the point of view of BSDE theory. Extensions in other directions can be interesting as well; for instance: (1) The problem in infinite time horizon (lifetime). Actually the discount factor ρ was introduced in problems (2.2) and (5.2) for this possible extension. (2) The problem in an incomplete market, i.e. m < n in subsection 5.2 or when r, μ , or σ depends on the regime i.

References

- [1] Aliprantis C, Border K. Infinite dimensional analysis. New York: Springer, 2006.
- [2] Bauerle N, Rieder U. Portfolio optimization with Markov-modulated stock prices and interest rates. IEEE Trans. Automat. Control, 2004, 49(3): 442-447.
- [3] Becherer D. Bounded solutions to backward SDEs with jumps for utility optimization and indifference hedging. Ann. Appl. Probab., 2006, 16(4): 2027-2054.
- [4] Cheridito P, Hu Y. Optimal consumption and investment in incomplete markets with general constraints. Stoch. Dyn., 2011, 11(02n03): 283-299.
- [5] Cox J C, Huang C. Optimal consumption and portfolio policies when asset prices follow a diffusion process. J. Econom. Theory, 1989, 49(1): 33-83.
- [6] Cvitanic J, and Zhang J. Contract theory in continuous-time models, Springer Science and Business Media, 2012.
- [7] Czichowsky C, Schweizer M. Cone-constrained continuous-time Markowitz problems. Ann. Appl. Probab., 2013, 23(2): 764-810.
- [8] Fan S, Hu Y, Tang S. Multi-dimensional backward stochastic differential equations of diagonally quadratic generators: the general result. arXiv:2007.04481, 2020.
- [9] Guan C, Xu Z Q, Yi F. A consumption-investment model with state-dependent lower bound constraint on consumption. J. Math. Anal. Appl., 2022, 516(1): 126511.
- [10] Hamilton J. A new approach to the economic analysis of nonstationary time series and the business cycle. Econometrica, 1989, 57(2), 357-384.
- [11] Hu Y, Imkeller P, Muller M. Utility maximization in incomplete markets. Ann. Appl. Probab., 2005, 15(3): 1691-1712.
- [12] Hu Y, Liang G, Tang S. Systems of ergodic BSDEs arising in regime switching forward performance processes. SIAM J. Control Optim., 2020, 58(4): 2503-2534.

- [13] Hu Y, Shi X, Xu Z Q. Constrained stochastic LQ control with regime switching and application to portfolio selection. Ann. Appl. Probab., 2022, 32 (1):426-460.
- [14] Hu Y, Shi X, Xu Z Q. Constrained stochastic LQ control on infinite time horizon with regime switching, ESAIM Control Optim. Calc. Var., 28 (2022), 5.
- [15] Hu Y, Shi X, Xu Z Q. Mean-variance asset-liability management with regime switching. arXiv:2201.01433v3.
- [16] Hu Y, Peng S. On the comparison theorem for multidimensional BSDEs, C. R. Acad. Sci. Paris, Ser. I, 343 (2006), 135-140.
- [17] Hu Y, Tang S. Multi-dimensional backward stochastic differential equations of diagonally quadratic generators. Stochastic Process. Appl., 2016, 126(4): 1066-1086.
- [18] Karatzas I, Lehoczky J, Shreve S. Optimal portfolio and consumption decisions for a "small investor" on a finite horizon, SIAM J. Control Optim., 25(1987), 1557-1586.
- [19] Karatzas I, Shreve S. Methods of mathematical finance. Springer Science and Business Media, 1998.
- [20] Kazamaki N. Continuous exponential martingales and BMO. Springer, 2006.
- [21] Kobylanski M. Backward stochastic differential equations and partial differential equations with quadratic growth. Ann. Probab., 28 (2000), 558-602.
- [22] Kramkov D, Pulido S. A system of quadratic BSDEs arising in a price impact model. Ann. Appl. Probab., 2016, 26(2): 794-817.
- [23] Laeven R J A, Stadje M. Robust portfolio choice and indifference valuation. Math. Oper. Res., 2014, 39(4): 1109-1141.
- [24] Liu R H. A finite-horizon optimal investment and consumption problem using regime-switching models. Int. J. Theor. Appl. Finance, 2014, 17(04): 1450027.
- [25] Matoussi, A and Xing, H. Convex duality for Epstein-Zin stochastic differential utility. Math. Finance, 2018, 28(4), 991-1019.
- [26] Merton R C. Lifetime portfolio selection under uncertainty: The continuous-time case. Rev. Econ. Stat., 1969: 247-257.
- [27] Morlais M A. A new existence result for quadratic BSDEs with jumps with application to the utility maximization problem. Stochastic Process. Appl., 2010, 120(10): 1966-1995.
- [28] Pardoux E, Peng S. Adapted solution of a backward stochastic differential equation. Systems Control Lett., 1990, 14(1): 55-61.
- [29] Rouge R, El Karoui N. Pricing via utility maximization and entropy. Math. Finance, 2000, 10(2): 259-276.

- [30] Sethi S P. Optimal consumption and investment with bankruptcy. Springer Science & Business Media, 1996.
- [31] Shin Y H. Optimal consumption and investment problem with Regime-Switching and CARA Utility. J. Chungcheong Math. Soc., 2013, 26(1): 85-85.
- [32] Sotomayor L R. Cadenillas A. Explicit solutions of consumption-investment problems in financial markets with regime switching. Math. Finance, 2009, 19(2): 251-279.
- [33] Wei J, Shen Y, Zhao Q. Portfolio selection with regime-switching and state-dependent preferences. J. Comput. Appl. Math., 2020, 365: 112361.
- [34] Xing, H. Consumption-investment optimization with Epstein-Zin utility in incomplete markets. Finance Stoch., 2017, 21, 227-262.
- [35] Xu Z Q, Yi F. An optimal consumption-investment model with constraint on consumption. Math. Control Relat. Fields, 2016, 6(3): 517-534.
- [36] Yin G, and Zhou X. Y. Markowitz's mean-variance portfolio selection with regime switching: from discrete-time models to their continuous-time limits. IEEE Trans. Automat. Contr., 2004, 49(3), 349-360.
- [37] Zariphopoulou T. Investment-consumption models with transaction fees and Markov-chain parameters. SIAM J. Control Optim., 1992, 30(3): 613-636.
- [38] Zariphopoulou T. Consumption-investment model with constraints. SIAM J. Control Optim., 1994, 32(1): 59-85.
- [39] Zhang Q, Yin G. Nearly-optimal asset allocation in hybrid stock investment models. J. Optim. Theory Appl., 2004, 121(2): 419-444.
- [40] Zhou X. Y., and Yin G. Markowitz's mean-variance portfolio selection with regime switching: A continuous-time model. SIAM J. Control Optim., 2003, 42, 1466-1482.