ALGEBRAIC PROPERTIES OF EDGE IDEALS OF CORONA PRODUCT OF CERTAIN GRAPHS

BAKHTAWAR SHAUKAT¹, MUHAMMAD ISHAQ^{1,*}, AHTSHAM UL HAQ¹, AND ZAHID IQBAL²

¹School of Natural Sciences, National University of Sciences and Technology Islamabad, Sector H-12, Islamabad Pakistan.

²Department of Mathematics and Statistics, Institute of Southern Punjab, Multan, Pakistan. * Corresponding email: ishaq_maths@yahoo.com; Tel: +92-51-90855591

Contributing authors emails: bakhtawar.shaukat@sns.nust.edu.pk, ahtsham2192@gmail.com, 786zahidwarraich@gmail.com

ABSTRACT. We study the algebraic invariants namely depth, Stanley depth, regularity and projective dimension of the residue class rings of the edge ideals associated with the corona product of various classes of graphs with any graph. We also give an expression for the Krull dimension of the residue class rings of the edge ideals associated with the corona product of any two graphs. In the end, we discuss when these graphs will be Cohen-Macaulay.

Key Words: Depth; Stanley depth; projective dimension; regularity; Krull dimension; corona product of two graphs.

2010 Mathematics Subject Classification: Primary: 13C15; Secondary: 13P10, 13F20.

INTRODUCTION

Let K be a field and $S := K[x_1, \ldots, x_r]$ denote the polynomial ring in r variables over K with standard grading. For a finitely generated Z-graded S-module A with a minimal free resolution

$$0 \longrightarrow \bigoplus_{j \in \mathbb{Z}} S(-j)^{\beta_{p,j}(A)} \longrightarrow \bigoplus_{j \in \mathbb{Z}} S(-j)^{\beta_{p-1,j}(A)} \longrightarrow \cdots \longrightarrow \bigoplus_{j \in \mathbb{Z}} S(-j)^{\beta_{0,j}(A)} \longrightarrow A \longrightarrow 0,$$

the regularity and projective dimension of A are defined by $\operatorname{reg}(A) = \max\{j - i : \beta_{i,j}(A) \neq 0\}$ and pdim $(A) = \max\{i : \beta_{i,j}(A) \neq 0\}$, respectively. Regularity plays a significant role as one of the keys indicators of a module's complexity and is an important invariant in commutative algebra. Values and bounds for the regularity and projective dimension of edge ideals have been studied by a number of researchers; see for instance [4, 6, 15, 18, 25].

Let $\mathfrak{m} := (x_1, \ldots, x_r)$ be the unique graded maximal ideal of S. The depth of A is defined to be the common length of all maximal M-sequences contained in \mathfrak{m} , for more details related to depth we refer the readers to [5]. For a finitely generated \mathbb{Z}^n -graded S-module A, if $a_i \in A$ is a homogeneous element and $X_i \subset \{x_1, \ldots, x_r\}$, then $a_i K[X_i]$ denotes the K-subspace of A generated by all homogeneous elements of the form $a_i t$, where t is a monomial in $K[X_i]$. The linear K-subspace $a_i K[X_i] \subset A$ is called a Stanley space of dimension $|X_i|$ if $a_i K[X_i]$ is a free $K[X_i]$ -module, where $|X_i|$ denotes the number of indeterminates in X_i . A Stanley decomposition of A is a presentation of the Kvector space A as a finite direct sum of Stanley spaces $\mathcal{D} : A = \bigoplus_{i=1}^r a_i K[X_i]$. The Stanley depth of decomposition \mathcal{D} and Stanley depth of A are defined as sdepth(\mathcal{D}) = min $\{|X_i| : i = 1, \ldots, r\}$ and sdepth(A) = max{sdepth(\mathcal{D}) : \mathcal{D} is a Stanley decomposition of A}, respectively. Herzog et al. showed that the Stanley depth of A = J/I, where $I \subset J \subset S$ can be computed in a finite number of steps by using some posets related to A in [23]. However, it is a difficult task to compute the Stanley depth even by using their method. For some known results, related to values and bounds of depth and Stanley depth, we refer the readers to [8, 28, 29, 35, 37]. Stanley in [38] conjectured that sdepth(A) \geq depth(A). This conjecture was proved for some special cases; see for instance [14, 16, 32, 36]. Later on, this conjecture was disproved by Duval et al. in [11] by providing a counterexample of S/I, where $I \subset S$ is a monomial ideal.

For a simple graph G = (V(G), E(G)) with vertex set $V(G) = \{x_1, \ldots, x_r\}$ and the edge set E(G), we identify the vertices of a graph and variables in S. The edge ideal of a graph G is defined as $I(G) = (x_i x_j : \{x_i, x_j\} \in E(G)) \subset S$. The module S/I(G) is called Cohen-Macaulay if depth $(S/I(G)) = \dim(S/I(G))$. A graph G is said to be Cohen-Macaulay if S/I(G) is Cohen-Macaulay. Characterization and construction of Cohen-Macaulay graphs is one of the fundamental problems with rich literature; see for instance [1, 2, 13, 30, 31, 39]. A broad categorization of Cohen-Macaulay graphs is difficult. Villarreal characterized all Cohen-Macaulay trees [41]. All Cohen-Macaulay chordal graphs are classified by Herzog et al. in [21] and later on Herzog and Hibi classified all Cohen-Macaulay bipartite graphs [22]. The primary goal of this paper is to compute some algebraic invariants and as a result of our findings, we characterize some Cohen-Macaulay graphs.

In this paper, we consider the edge ideals associated to the graph $X \odot H$ called the corona product of X and H. Here H could be any graph and X is either a path, a cycle, a complete graph, a star graph, or a complete bipartite graph. Particularly, if H is a null graph, then the corona product $X \odot H$ is a |V(H)|-fold bristled graph. The first section of this paper contains some background regarding Graph Theory, relevant useful results from Commutative Algebra and all the concepts that will be used throughout the paper. In the second section, we compute depth, projective dimension and lower bound of the Stanley depth for the residue class rings of the edge ideals of $X \odot H$ in terms of corresponding algebraic invariants for the residue class ring of the edge ideal of H; see for instance Theorem 2.5, Theorem 2.9, Theorem 2.13, Theorem 2.17 and Theorem 2.21. In particular, if H is a null graph, then the exact values for Stanley depth are determined. Moreover, we give an explicit formulas for the regularity of residue class rings of the edge ideals of $X \odot H$, see Theorem 2.28, Theorem 2.30, Theorem 2.32, Theorem 2.34 and Theorem 2.36. Third section is devoted to find an expression for Krull dimension of edge ideal of corona product of any two graphs see Theorem 3.1. As a consequence of this theorem, we characterize Cohen-Macaulay $X \odot H$ graphs; see for instance Theorem 3.4.

The corona product $X \odot H$ gives rise to a variety of nice graph structures. Some authors have already discussed depth and Stanley depth of the cyclic modules associated to some specific classes of corona products of graphs; see for instance [10]. To find an expression for the algebraic invariants of the residue class rings of the edge ideals of $X \odot H$ seems to be a challenging task, if Xis any arbitrary graph. Therefore, we initially focus on fixing the underlying graph X to be some well-known graphs, that allow us to gain insight into the properties of these structures and their associated algebraic invariants. Our work may help to extend all the results obtained in this paper to compute the said invariants when X is arbitrary. To this end, we have succeeded in finding an expression for the value of the Krull dimension of the residue class rings of the edge ideal of the graphs $X \odot H$.

1. Preliminaries

In this section, we recall some notions from Graph Theory, review a few auxiliary findings from Commutative Algebra and introduce some notations that are needed in the next sections. Throughout this paper, all graphs are assumed to be finite, simple and may have isolated vertices. For a graph G with $V(G) = \{x_1, \ldots, x_n\}$, adding a whisker to graph G at x_i means adding a new vertex y to V(G) and edge $\{x_i, y\}$ to E(G). A graph obtained by adding a whisker to each vertex of any given graph is called a whisker graph. A graph N_r is said to be a null graph on rvertices if $V(N_r) = \{x_1, \ldots, x_r\}$ and $E(N_r) = \emptyset$. Moreover, if r = 1 then N_1 is also called a trivial graph. The degree of a vertex $x_i \in V(G)$ is the number of adjacent vertices to x_i in graph G and is represented by $\deg_G(x_i)$. Any vertex of degree 1 is called a leaf or pendant vertex of G. For any integer $t \ge 1$, a t-fold bristled graph of G is formed by attaching t pendant vertices to each vertex of G. An internal vertex is a vertex that is not a leaf. A graph is said to be a connected graph if there is a path between any two vertices. For $r \ge 1$, a path P_r is a graph on r vertices such that $E(P_r) = \{\{x_j, x_{j+1}\} : 1 \leq j \leq r-1\} \text{ (if } r = 1, \text{ then } E(P_1) = \emptyset\}. \text{ For } r \geq 3, \text{ a cycle } C_r \text{ on } r \text{ vertices is a graph such that } E(C_r) = \{\{x_j, x_{j+1}\} : 1 \leq j \leq r-1\} \cup \{x_1, x_r\}. \text{ Let } r \geq 1, \text{ a complete graph } K_r \text{ on } r \text{ vertices is a graph in which each pair of vertices is connected by an edge (if } r = 1, \text{ then } E(K_1) = \emptyset\}. A simple and connected graph <math>T_r$ on r vertices is said to be a *tree* if there exists a unique path between any two vertices of T_r . For $k \geq 1$, a k-star is a tree with k leaves and a single vertex with degree k. We denote a k-star by S_k . A *bipartite graph* is a graph in which the set of vertices is partitioned into two disjoint sets called partite sets such that no two vertices of the graph within the same partite set is connected to each vertex of the other partite graph such that every vertex of one partite graph with partite sets $K_u = \{x_1, \ldots, x_u\}$ and $K_v = \{x_{u+1}, \ldots, x_{u+v}\}$. A vertex x_j is a *neighbor* of a vertex x_i in a graph G if $\{x_i, x_j\} \in E(G)$. The *neighborhood* $N_G(x_i)$ of a vertex x_i is the set of all neighbors of x_i , that is, $N_G(x_i) := \{x_j \in V(G) \mid \{x_i, x_j\} \in E(G)\}$. For graph G, a subset W of V(G) is called an *independent set* if no two vertices in W are adjacent.

Lemma 1.1 ([26, Lemma 1]). $\dim(S/I(G)) = \max\{|W|: W \text{ is an independent set of } G\}.$

A subgraph H of a graph G, written as $H \subseteq G$, is a graph such that $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$. For a subset $U \subseteq V(G)$, an *induced subgraph* of G is a graph G' := (U, E(G')), such that $E(G') = \{\{x_i, x_j\} \in E(G) : \{x_i, x_j\} \subseteq U\}$. A graph is said to be *chordal* if it does not contain any induced cycle of length strictly greater than 3. A matching M in a graph G is a subset of E(G) in which no two edges are adjacent in G. An *induced matching* in G is a matching that forms an induced subgraph of G. An *induced matching number* of G is denoted as indmat(G) and defined as

 $indmat(G) = max\{|M|: M \text{ is an induced matching in } G\}.$

Lemma 1.2 ([19, Corollary 6.9]). If G is a chordal graph, then reg(S/I(G)) = indmat(G).

Next we present a few results, these results will play a key role in the proofs of our main results.

Lemma 1.3 ([3, Theorem 2.6 and Theorem 2.7]). If $r \ge 2$ and $I(S_{r-1}) \subset S = K[V(S_r)]$, then

$$depth(S/I(S_{r-1})) = sdepth(S/I(S_{r-1})) = 1.$$

Lemma 1.4. If $0 \longrightarrow X \longrightarrow Y \longrightarrow Z \longrightarrow 0$ is an exact sequence of \mathbb{Z}^n -graded S-module, then

- (a) $\operatorname{depth}(Y) \ge \min\{\operatorname{depth}(X), \operatorname{depth}(Z)\}, [5, \operatorname{Proposition} 1.2.9].$
- (b) $\operatorname{sdepth}(Y) \ge \min\{\operatorname{sdepth}(X), \operatorname{sdepth}(Z)\}, [36, \operatorname{Lemma 2.2}].$

Lemma 1.5. Let $I \subset S$ be a monomial ideal and f be a monomial in S such that $f \notin I$. Then

- (a) $\operatorname{depth}(S/(I:f)) \ge \operatorname{depth}(S/I)$, [36, Corollary 1.3].
- (b) $\operatorname{sdepth}(S/(I:f)) \ge \operatorname{sdepth}(S/I), [7, \operatorname{Proposition} 2.7].$

Lemma 1.6 ([6, Theorem 4.3]). Let I be a monomial ideal and let f be an arbitrary monomial in S. Then

 $depth(S/I) = depth(S/(I:f)) \text{ if } depth(S/(I,f)) \ge depth(S/(I:f)).$

We have a similar result for Stanley depth in the next lemma.

Lemma 1.7. Let I be a monomial ideal and let f be a monomial in S such that $f \notin I$. Then

 $\operatorname{sdepth}(S/I) = \operatorname{sdepth}(S/(I:f))$ if $\operatorname{sdepth}(S/(I,f)) \ge \operatorname{sdepth}(S/(I:f))$.

Proof. Consider the short exact sequence

$$0 \longrightarrow S/(I:f) \xrightarrow{\cdot f} S/I \longrightarrow S/(I,f) \longrightarrow 0.$$

By Lemma 1.4, $\operatorname{sdepth}(S/I) \ge \min\{\operatorname{sdepth}(S/(I : f)), \operatorname{sdepth}(S/(I, f))\}$. If $\operatorname{sdepth}(S/(I, f)) \ge \operatorname{sdepth}(S/(I : f))$, then $\operatorname{sdepth}(S/I) \ge \operatorname{sdepth}(S/(I : f))$. By Lemma 1.5, we get $\operatorname{sdepth}(S/I) \le \operatorname{sdepth}(S/(I : f))$. Thus the required result follows.

Lemma 1.8. Let $I \subset S$ be a monomial ideal, and $\hat{S} = S \otimes_K K[x_{r+1}]$. Then

- (a) depth (\hat{S}/I) = depth(S/I) + 1 and sdepth (\hat{S}/I) = sdepth(S/I) + 1, [23, Lemma 3.6].
- (b) $\operatorname{reg}(\hat{S}/I) = \operatorname{reg}(S/I)$, [33, Lemma 3.6].

Lemma 1.9 ([42, Proposition 2.2.20]). Let $1 \le n < r$. If $S_1 = K[x_1, \ldots, x_n]$ and $S_2 = K[x_{n+1}, \ldots, x_r]$. Then $S/(I+J) \cong S_1/I \otimes_K S_2/J$.

We use Lemma 1.9 and combine it with [42, Proposition 2.2.21] and [36, Theorem 3.1] for depth and Stanley depth, respectively and get the following useful result.

Lemma 1.10. depth_S $(S_1/I \otimes_K S_2/J) = depth_S(S/(I+J)) = depth_{S_1}(S_1/I) + depth_{S_2}(S_2/J)$ and sdepth_S $(S_1/I \otimes_K S_2/J) \ge sdepth_{S_1}(S_1/I) + sdepth_{S_2}(S_2/J).$

Lemma 1.11 ([5, Theorems 1.3.3]). (Auslander–Buchsbaum formula) If R is a commutative Noetherian local ring and M is a non-zero finitely generated R-module of finite projective dimension, then

$$pdim(M) + depth(M) = depth(R).$$

Lemma 1.12 ([27, Lemma 3.2]). If $I \subset S_1 = K[x_1, \ldots, x_n]$ and $J \subset S_2 = K[x_{n+1}, \ldots, x_r]$ are non-zero homogeneous ideals of S_1 and S_2 and regard I + J as a homogeneous ideal of S. Then

$$\operatorname{reg}(S/I + J) = \operatorname{reg}(S_1/I) + \operatorname{reg}(S_2/J).$$

In the next lemma, proof of parts (a) and (c) follows from Corollary 20.19 and Proposition 20.20 of [12], while part (b) follows from [9, Lemma 2.10].

Lemma 1.13 ([6, Theorem 4.7]). Let I be a monomial ideal and x_i be a variable of S. Then

- (a) $\operatorname{reg}(S/I) = \operatorname{reg}(S/(I:x_i)) + 1$, if $\operatorname{reg}(S/(I:x_i)) > \operatorname{reg}(S/(I,x_i))$,
- (b) $\operatorname{reg}(S/I) \in \{\operatorname{reg}(S/(I, x_i)) + 1, \operatorname{reg}(S/(I, x_i))\}, \text{ if } \operatorname{reg}(S/(I : x_i)) = \operatorname{reg}(S/(I, x_i)),$
- (c) $\operatorname{reg}(S/I) = \operatorname{reg}(S/(I, x_i))$ if $\operatorname{reg}(S/(I : x_i)) < \operatorname{reg}(S/(I, x_i))$.

Definition 1.14 ([17]). The corona product of two graphs X and H denoted by $X \odot H$ is a graph constructed by taking one copy of graph X and |V(X)| copies of H, namely, $H_1, H_2, \ldots, H_{|V(X)|}$ and then connecting the i^{th} vertex of graph X to every vertex in H_i .

If X and H are two graphs, then

- $|V(X \odot H)| = |V(X)|(|V(H)| + 1),$
- $|E(X \odot H)| = |E(X)| + |V(X)||E(H)| + |V(X)||V(H)|.$
- If $X \neq H$ then $X \odot H \neq H \odot X$.

For a monomial ideal I we denote the minimal set of monomial generators of I by $\mathcal{G}(I)$. Let |V(X)| = n and |V(H)| = m. Throughout this paper, when we consider $X \odot H$, the vertices of X are labeled as y_1, y_2, \ldots, y_n . Let x_1, x_2, \ldots, x_m be the vertices of H. The vertices of the i^{th} copy H_i of H in $X \odot H$ are labeled as $x_{i1}, x_{i2}, \ldots, x_{im}$ such that $\{x_{il}, x_{ik}\} \in E(H_i)$ iff $\{x_l, x_k\} \in E(H)$. Thus $V(X \odot H) = \{y_1, y_2, \ldots, y_n\} \bigcup_{i=1}^n \{x_{i1}, x_{i2}, \ldots, x_{im}\}$ and

$$E(X \odot H) = E(X) \bigcup_{i=1}^{n} \{\{y_i, x_{i1}\}, \{y_i, x_{i2}\}, \dots, \{y_i, x_{im}\}\} \bigcup_{i=1}^{n} E(H_i).$$

The minimal set of monomial generators of a monomial ideal $I(X \odot H)$ is as follow:

$$\mathcal{G}(I(X \odot H)) = \mathcal{G}(I(X)) \bigcup_{i=1}^{n} \{y_i x_{i1}, y_i x_{i2}, \dots, y_i x_{im}\} \bigcup_{i=1}^{n} \mathcal{G}(I(H_i)).$$

For example, if $X = P_3$ and $H = T_7$, where T_7 is a tree on 7 vertices, then the graph $X \odot H$ is shown in Figure 1.





Remark 1.15. From definition of $X \odot H$, it is clear that if H is any graph and X has s connected components, say X_1, \ldots, X_s , then $X \odot H$ has again s connected components $X_1 \odot H, \ldots, X_s \odot H$. But if X is a connected graph and H has l connected components then $X \odot H$ is still a connected graph. It is obvious that $X \odot H$ has no isolated vertex even if X and H have isolated vertices. Therefore, we allow isolated vertices in both graphs X and H. The isolated vertices of H play a crucial role in our results. Therefore, we introduce the following terminologies.

For any graph H, we denote the set of isolated vertices of H by i(H) and if $A := V(H) \setminus i(H)$ then we denote the induced subgraph of H on A by H'. It is easy to see that |V(H)| = |i(H)| + |A| and $I(H) = (x_i x_j, x_k : \{x_i, x_j\} \in E(H') \text{ and } x_k \in i(H))$ (for instance; if $E(H') = \emptyset$, then $I(H) = (x_k : x_k \in i(H))$). Also $K[V(H)]/I(H) \cong K[V(H')]/I(H')$. Therefore, we have depth(K[V(H)]/I(H)) =depth(K[V(H')]/I(H') and sdepth(K[V(H)]/I(H)) = sdepth(K[V(H')]/I(H'). Similarly, we get reg(K[V(H)]/I(H)) = reg(K[V(H')]/I(H') and dim(K[V(H)]/I(H)) = dim(K[V(H')]/I(H'). If H is a null graph, then |V(H)| = i(H) and depth(K[V(H)]/I(H)) = sdepth(K[V(H)]/I(H)) =reg(K[V(H)]/I(H)) = dim(K[V(H)]/I(H)) = 0. Moreover, if $i(H) = \emptyset$, then in this case we consider $K[i(H)] \cong K$. For example, let $X = C_3$ and H be a graph which is a union of two graphs, one is a complete graph on 4 vertices and the other is a null graph on 3 vertices, that is, $H = K_4 \cup N_3$, as shown in Figure 2(A). Figure 2(B) is the corona product of C_3 and H.



FIGURE 2

For a monomial ideal I, supp $(I) := \{x_i : x_i | f \text{ for some } f \in \mathcal{G}(I)\}.$

Remark 1.16. We associate a graph G_I to the squarefree monomial ideal I with $V(G_I) = \operatorname{supp}(I)$ and $E(G_I) = \{\{x_i, x_j\} : x_i x_j \in \mathcal{G}(I)\}$. Let $x_t \in S$ be a variable of the polynomial ring S such that $x_t \notin I$. Then $(I : x_t)$ and (I, x_t) are the squarefree monomial ideals of S such that $G_{(I:x_t)}$ and $G_{(I,x_t)}$ are subgraphs of G_I . Consider the graph $C_3 \odot H$ as given in Figure 2(B), if $S = K[V(C_3 \odot H)]$, then $G_{(I(C_3 \odot H):y_3)}$ and $G_{(I(C_3 \odot H),y_3)}$ are subgraphs of $G_{I(C_3 \odot H)}$ see Figure 3. It is evident from the Figure 3(A) and Figure 3(B) that we have the following isomorphisms:

$$S/(I(C_3 \odot H): y_3) \cong \bigotimes_{l=1}^2 K[V(K_4)]/I(K_4) \otimes_K K[x_{15}, x_{16}, x_{17}, x_{25}, x_{26}, x_{27}, y_3]$$

and

 $S/(I(C_3 \odot H), y_3) \cong K[V(P_2 \odot H)]/I(P_2 \odot H) \otimes_K K[V(K_4)]/I(K_4) \otimes_K K[x_{35}, x_{36}, x_{37}].$





2. DEPTH, STANLEY DEPTH AND REGULARITY OF RESIDUE CLASS RINGS OF SOME EDGE IDEALS

Let P_n, C_n, K_n, S_n and $K_{u,v}$ are the graphs as defined earlier and H be any graph. In this section, we focus on the invariants depth, Stanley depth, projective dimension and regularity of $S/I(X \odot H)$, where $X \in \{P_n, C_n, K_n, S_n, K_{u,v}\}$.

Remark 2.1. If *H* is a null graph and $X \in \{P_n, C_n, K_n, S_n, K_{u,v}\}$, then $X \odot H$ is a |V(H)|-fold bristled graph of *X* and we denote it by $Br_{|V(H)|}(X)$.

This section is further divided into two subsections. In the first subsection, we discuss depth, Stanley depth and projective dimension of $S/I(X \odot H)$, while in the second subsection, we discuss the regularity.

2.1. Depth, Stanley depth and projective dimension.

. If N_1 denotes a trivial graph, then we have the following K-algebra isomorphisms:

(2.1)
$$K[V(N_1 \odot H)]/(I(N_1 \odot H) : y_1) \cong K[y_1],$$

(2.2)
$$K[V(N_1 \odot H)]/(I(N_1 \odot H), y_1) \cong K[V(H)]/I(H) \otimes_K K[i(H)].$$

Now, we give an elementary lemma that will be used frequently in this subsection.

Lemma 2.2. Let $S = K[V(N_1 \odot H)]$. Then

$$\operatorname{depth}(S/I(N_1 \odot H)) = \operatorname{sdepth}(S/I(N_1 \odot H)) = 1.$$

Proof. First we will prove the result for depth. If H is a null graph, then we have |V(H)| = |i(H)|and $N_1 \odot H \cong S_{|V(H)|}$. Thus by Lemma 1.3, depth $(S/I(N_1 \odot H)) = 1$. Let H is not a null graph, by using Eq. 2.1, we get $\operatorname{depth}(S/(I(N_1 \odot H) : y_1)) = \operatorname{depth}(K[y_1]) = 1$. By using Lemma 1.8 on Eq. 2.2, we have depth $(S/(I(N_1 \odot H), y_1)) = depth(K[V(H)]/I(H)) + |i(H)| \ge 1$. Thus by Lemma 1.6, depth $(S/I(N_1 \odot H)) = 1$. The proof of Stanley depth is similar as depth by using Lemma 1.7 in place of Lemma 1.6.

Corollary 2.3. If $S = K[V(N_1 \odot H)]$, then $pdim(S/I(N_1 \odot H)) = |V(H)|$.

Proof. By using Lemma 1.11 and Lemma 2.2, the required result follows.

Remark 2.4. For n = 0, we define $I(P_0 \odot H) = (0)$, we have $K[V(P_0 \odot H)]/I(P_0 \odot H) \cong K$. Thus $\operatorname{depth}(K[V(P_0 \odot H)]/I(P_0 \odot H)) = \operatorname{sdepth}(K[V(P_0 \odot H)]/I(P_0 \odot H)) = 0.$ Moreover, if n = 1, then $P_1 \odot H \cong N_1 \odot H$.

Let y_n be a leaf of path P_n , we have the following K-algebra isomorphisms: (2.3)

 $K[V(P_n \odot H)]/(I(P_n \odot H) : y_n) \cong K[V(P_{n-2} \odot H)]/(I(P_{n-2} \odot H)) \otimes_K K[V(H)]/(I(H) \otimes_K K[i(H) \cup \{y_n\}],$ (2.4)

$$K[V(P_n \odot H)]/(I(P_n \odot H), y_n) \cong K[V(P_{n-1} \odot H)]/I(P_{n-1} \odot H) \otimes_K K[V(H)]/I(H) \otimes_K K[i(H)].$$

Theorem 2.5. Let $n \ge 1$ and $S = K[V(P_n \odot H)]$. Then

- (a) depth $(S/I(P_n \odot H)) = \lceil \frac{n}{2} \rceil + \lceil \frac{n-1}{2} \rceil (depth(K[V(H)]/I(H)) + |i(H)|).$ (b) sdepth $(S/I(P_n \odot H)) \ge \lceil \frac{n}{2} \rceil + \lceil \frac{n-1}{2} \rceil (sdepth(K[V(H)]/I(H)) + |i(H)|).$ In particular, if H is a null graph, then

$$\operatorname{sdepth}(S/I(P_n \odot H)) = \left\lceil \frac{n}{2} \right\rceil + \left\lceil \frac{n-1}{2} \right\rceil |V(H)|.$$

Proof. Let $t := \operatorname{depth}(K[V(H)]/I(H))$. First we will prove the result for depth by using induction on n. If n = 1, then $P_1 \odot H \cong N_1 \odot H$ and the result follows by Lemma 2.2. Let $n \ge 2$. By induction on n and using Eq. 2.3, Eq. 2.4, Lemma 1.8, Lemma 1.10, we get

$$\begin{split} \operatorname{depth}(S/(I(P_n \odot H) : y_n)) \\ &= \operatorname{depth}(K[V(P_{n-2} \odot H)]/I(P_{n-2} \odot H)) + \operatorname{depth}(K[V(H)]/I(H)) + \operatorname{depth}(K[i(H) \cup \{y_n\}]) \\ &= \lceil \frac{n-2}{2} \rceil + \lceil \frac{n-3}{2} \rceil (t+|i(H)|) + t + |i(H)| + 1 = \lceil \frac{n}{2} \rceil + \lceil \frac{n-1}{2} \rceil (t+|i(H)|), \end{split}$$

$$\operatorname{depth}(S/(I(P_n \odot H), y_n)))$$

$$= \operatorname{depth}(K[V(P_{n-1} \odot H)]/I(P_{n-1} \odot H)) + \operatorname{depth}(K[V(H)]/I(H)) + \operatorname{depth}(K[i(H)])$$
$$= \left\lceil \frac{n-1}{2} \right\rceil + \left\lceil \frac{n-2}{2} \right\rceil \left(t + |i(H)|\right) + t + |i(H)| = \left\lceil \frac{n-1}{2} \right\rceil + \left\lceil \frac{n}{2} \right\rceil (t + |i(H)|).$$

Since $\left\lceil \frac{n}{2} \right\rceil + \left\lceil \frac{n-1}{2} \right\rceil (t+|i(H)|) \leq \left\lceil \frac{n-1}{2} \right\rceil + \left\lceil \frac{n}{2} \right\rceil (t+|i(H)|)$, thus by using Lemma 1.6, we get $\operatorname{depth}(S/(I(P_n \odot H)) = \left\lceil \frac{n}{2} \right\rceil + \left\lceil \frac{n-1}{2} \right\rceil (t+|i(H)|)$.

To prove the result for Stanley depth, consider if H is not a null graph, then by considering Eq. 2.3 and Eq. 2.4 and applying Lemma 1.8, Lemma 1.10 and Lemma 1.4, we get the required inequality for Stanley depth. But if H is a null graph, we have

$$S/(I(P_n \odot H) : y_n) \cong K[V(P_{n-2} \odot H)]/I(P_{n-2} \odot H)) \otimes_K K[V(H) \cup \{y_n\}]$$

$$S/(I(P_n \odot H), y_n) \cong K[V(P_{n-1} \odot H)]/I(P_{n-1} \odot H) \otimes_K K[V(H)],$$

and the proof is similar to depth by replacing Lemma 1.6 by Lemma 1.7.

Corollary 2.6. If Stanley's inequality holds for K[V(H)]/I(H), then it also holds for $S/I(P_n \odot H)$.

Corollary 2.7. If $n \ge 1$, then

$$pdim(S/I(P_n \odot H)) = n(|V(H)| + 1) - \lceil \frac{n}{2} \rceil - \lceil \frac{n-1}{2} \rceil (depth(K[V(H)]/I(H)) + |i(H)|).$$

Proof. As $|V(P_n \odot H)| = n(|V(H)| + 1)$. By using Lemma 1.11 and Theorem 2.5, one can find the required result.

Corollary 2.8. Let $n, m, s \ge 1$ and $S = K[V(P_n \odot P_m)]$. Then

- (a) depth $(S/I(P_n \odot P_m)) = \lceil \frac{n}{2} \rceil + \lceil \frac{n-1}{2} \rceil \lceil \frac{m}{3} \rceil$. (b) pdim $(S/I(P_n \odot P_m)) = n(m+1) \lceil \frac{n}{2} \rceil \lceil \frac{n-1}{2} \rceil \lceil \frac{m}{3} \rceil$. (c) depth $(K[V(Br_s(P_n))]/I(Br_s(P_n))) = \text{sdepth}(K[V(Br_s(P_n))]/I(Br_s(P_n))) = \lceil \frac{n}{2} \rceil + \lceil \frac{n-1}{2} \rceil s$.

For $y_{n-1} \in V(C_n \odot H)$, we have the following K-algebra isomorphisms:

$$(2.5) \quad K[V(C_n \odot H)]/(I(C_n \odot H) : y_{n-1}) \cong K[V(P_{n-3} \odot H)]/I(P_{n-3} \odot H) \bigotimes_{l=1}^2 K[V(H)]/I(H) \bigotimes_{l=1}^2 K[i(H)] \otimes_K K[y_{n-1}],$$

(2.6)

$$K[V(C_n \odot H)]/(I(C_n \odot H), y_{n-1}) \cong K[V(P_{n-1} \odot H)]/I(P_{n-1} \odot H) \otimes_K K[V(H)]/I(H) \otimes_K K[i(H)].$$

Theorem 2.9. If $n \geq 3$ and $S = K[V(C_n \odot H)]$, then

- (a) depth $(S/I(C_n \odot H)) = \lceil \frac{n-1}{2} \rceil + \lceil \frac{n}{2} \rceil (depth(K[V(H)]/I(H)) + |i(H)|).$ (b) sdepth $(S/I(C_n \odot H)) \ge \lceil \frac{n-1}{2} \rceil + \lceil \frac{n}{2} \rceil (sdepth(K[V(H)]/I(H)) + |i(H)|).$ In particular, if H is a null graph, then

$$\operatorname{sdepth}(S/I(C_n \odot H)) = \left\lceil \frac{n-1}{2} \right\rceil + \left\lceil \frac{n}{2} \right\rceil |V(H)|.$$

Proof. Firstly, we discuss the result for depth. Let $t := \operatorname{depth}(K[V(H)]/I(H))$. By using Theorem 2.5, Lemma 1.8 and Lemma 1.10 on Eq. 2.5 and Eq. 2.6, we get

$$\begin{split} \operatorname{depth}(S/(I(C_n \odot H) : y_{n-1})) &= \operatorname{depth}(K[V(P_{n-3} \odot H)]/I(P_{n-3} \odot H)) + 2\operatorname{depth}(K[V(H)]/I(H)) + 2\operatorname{depth}(K[i(H)]) \\ &+ \operatorname{depth}(K[y_{n-1}]) \\ &= \lceil \frac{n-3}{2} \rceil + \lceil \frac{n-4}{2} \rceil (t + |i(H)|) + 2t + 2|i(H)| + 1 = \lceil \frac{n-1}{2} \rceil + \lceil \frac{n}{2} \rceil (t + |i(H)|), \\ \operatorname{depth}(S/(I(C_n \odot H), y_{n-1})) \\ &= \operatorname{depth}(K[V(P_{n-1} \odot H)]/I(P_{n-1} \odot H)) + \operatorname{depth}(K[V(H)]/I(H)) + \operatorname{depth}(K[i(H)]) \\ &= \lceil \frac{n-1}{2} \rceil + \lceil \frac{n-2}{2} \rceil (t + |i(H)|) + t + |i(H)| \\ &= \lceil \frac{n-1}{2} \rceil + \lceil \frac{n}{2} \rceil (t + |i(H)|). \end{split}$$

Using Lemma 1.6, completes the proof for depth.

Now, we prove the result for Stanley depth. If H is not a null graph, then we get the required lower bound for Stanley depth by applying Lemma 1.8, Lemma 1.10, Theorem 2.5 and Lemma 1.4 on Eq. 2.5 and Eq. 2.6. But if H is a null graph, then by Eq. 2.5 and Eq. 2.6 we have the following K-algebra isomorphisms:

$$S/(I(C_n \odot H) : y_{n-1}) \cong K[V(P_{n-3} \odot H)]/I(P_{n-3} \odot H) \bigotimes_{l=1}^{2} K[V(H)] \otimes_{K} K[y_{n-1}],$$
$$S/(I(C_n \odot H), y_{n-1}) \cong K[V(P_{n-1} \odot H)]/I(P_{n-1} \odot H) \otimes_{K} K[V(H)],$$

and the proof is similar to depth by using Lemma 1.7 instead of Lemma 1.6.

Corollary 2.10. Stanley's inequality holds for $S/I(C_n \odot H)$ if it holds for K[V(H)]/I(H).

Corollary 2.11. Let $n \geq 3$. Then

$$\operatorname{pdim}(S/I(C_n \odot H)) = n(|V(H)| + 1) - \left\lceil \frac{n-1}{2} \right\rceil - \left\lceil \frac{n}{2} \right\rceil \big(\operatorname{depth}(K[V(H)]/I(H)) + |i(H)|\big).$$

Proof. As $|V(C_n \odot H)| = n(|V(H)| + 1)$. By using Lemma 1.11 and Theorem 2.9, one can find the required result.

Corollary 2.12. Let $n, m \ge 3$, $s \ge 1$ and $S = K[V(C_n \odot C_m)]$. Then

- (a) depth $(S/I(C_n \odot C_m)) = \left\lceil \frac{n-1}{2} \right\rceil + \left\lceil \frac{n}{2} \right\rceil \left\lceil \frac{m-1}{3} \right\rceil$. (b) pdim $(S/I(C_n \odot C_m)) = n(m+1) \left\lceil \frac{n-1}{2} \right\rceil \left\lceil \frac{n}{2} \right\rceil \left\lceil \frac{m-1}{3} \right\rceil$. (c) depth $(K[V(Br_s(C_n))]/I(Br_s(C_n))) = \text{sdepth}(K[V(Br_s(C_n))]/I(Br_s(C_n))) = \left\lceil \frac{n-1}{2} \right\rceil$ $\left\lfloor \frac{n}{2} \right\rfloor s.$

For $y_j \in V(K_n \odot H)$, where y_j is an arbitrary vertex of K_n , we get the following isomorphisms:

$$(2.7) K[V(K_n \odot H)]/(I(K_n \odot H) : y_j) \cong \bigotimes_{l=1}^{n-1} K[V(H)]/I(H) \bigotimes_{l=1}^{n-1} K[i(H)] \otimes_K K[y_j],$$

$$(2.8)$$

$$K[V(K_n \odot H)]/(I(K_n \odot H), y_j) \cong K[V(K_{n-1} \odot H)]/I(K_{n-1} \odot H) \otimes_K K[V(H)]/I(H) \otimes_K K[i(H)],$$

Theorem 2.13. Let $n \ge 1$ and $S = K[V(K_n \odot H)]$. Then

- (a) depth $(S/I(K_n \odot H)) = 1 + (n-1)(depth(K[V(H)]/I(H)) + |i(H)|).$
- (b) $\operatorname{sdepth}(S/I(K_n \odot H)) \ge 1 + (n-1)(\operatorname{sdepth}(K[V(H)]/I(H)) + |i(H)|).$ In particular, if H is a null graph, then

$$sdepth(S/I(K_n \odot H)) = 1 + (n-1)|V(H)|.$$

Proof. We first prove the result for depth. Let $t := \operatorname{depth}(K[V(H)]/I(H))$. If $n = 1, 2, K_1 \cong N_1$ and $K_2 \cong P_2$, we get the required result by using Lemma 2.2 and Theorem 2.5. By applying Lemma 1.8, Lemma 1.10 on Eq. 2.7 and Eq. 2.8, we have

$$depth(S/(I(K_n \odot H) : y_j)) = (n-1) depth(K[V(H)]/I(H)) + (n-1) depth(K[i(H)]) + depth(K[y_j]) = (n-1)t + (n-1)|i(H)| + 1 = 1 + (n-1)(t + |i(H)|)$$

and by induction on n,

 $\operatorname{depth}(S/(I(K_n \odot H), y_i)) = \operatorname{depth}(K[V(K_{n-1} \odot H)]/I(K_{n-1} \odot H)) + \operatorname{depth}(K[V(H)]/I(H))$ $+ \operatorname{depth}(K[i(H)])$ = 1 + (n-2)(t + |i(H)|) + t + |i(H)|= 1 + (n-1)(t + |i(H)|).

Thus the required result for the depth follows by Lemma 1.6.

Next, to prove the result for Stanley depth, consider if H is not a null graph, then we get the required inequality for Stanley depth by applying Lemma 1.8, Lemma 1.10, and Lemma 1.4 on Eq. 2.7 and Eq. 2.8. If H is a null graph, then by Eq. 2.7 and Eq. 2.8,

$$S/(I(K_n \odot H) : y_j) \cong \bigotimes_{k=1}^{n-1} K[V(H)] \otimes_K K[y_j],$$
$$S/(I(K_n \odot H), y_j) \cong K[V(K_{n-1} \odot H)]/I(K_{n-1} \odot H) \otimes_K K[V(H)]$$

and we get the desired result for Stanley depth similarly as we obtained for depth just by replacing Lemma 1.6 by Lemma 1.7.

Corollary 2.14. Stanley's inequality holds for $S/I(K_n \odot H)$ if it holds for K[V(H)]/I(H).

Corollary 2.15. Let $n \ge 1$. Then

 $pdim(S/I(K_n \odot H)) = n(|V(H)| + 1) - 1 - (n - 1)(depth(K[V(H)]/I(H)) + |i(H)|).$

Proof. As $|V(K_n \odot H)| = n(|V(H)| + 1)$. By using Lemma 1.11 and Theorem 2.13, one can find the required result.

Corollary 2.16. Let $n, m, s \ge 1$ and $S = K[V(K_n \odot K_m)]$. Then

- (a) depth $(S/I(K_n \odot K_m)) = n$.
- (b) $\operatorname{pdim}(S/I(K_n \odot K_m)) = nm.$
- (c) depth $(K[V(Br_s(K_n))]/I(Br_s(K_n))) = \text{sdepth}(K[V(Br_s(K_n))]/I(Br_s(K_n))) = 1 + (n 1)s.$

Theorem 2.17. Let $S = K[V(S_n \odot H)]$. Then

- (a) depth $(S/I(S_n \odot H)) = n + depth(K[V(H)]/I(H)) + |i(H)|.$
- (b) $\operatorname{sdepth}(S/I(S_n \odot H)) \ge n + \operatorname{sdepth}(K[V(H)]/I(H)) + |i(H)|.$
 - In particular, if H is a null graph, then

$$\operatorname{sdepth}(S/I(S_n \odot H)) = n + |V(H)|.$$

Proof. Let $t := \operatorname{depth}(K[V(H)]/I(H))$. First we will prove the depth result. If $1 \le n \le 3$, then the result follows from Lemma 2.2 and Theorem 2.5. Let $n \ge 4$. For $y_{n+1} \in S_n \odot H$, we have the following isomorphisms:

$$S/(I(S_n \odot H) : y_{n+1}) \cong \bigotimes_{k=1}^{n-1} K[V(N_1 \odot H)]/I(N_1 \odot H) \otimes_K K[i(H) \cup \{y_{n+1}\}] \otimes_K K[V(H)]/I(H).$$

Therefore, by Lemma 1.8, Lemma 1.10 and Lemma 2.2, we have

$$depth(S/(I(S_n \odot H) : y_{n+1})) = (n-1) depth(K[V(N_1 \odot H)]/I(N_1 \odot H)) + depth(K[i(H) \cup \{y_{n+1}\}]) + depth(K[V(H)]/I(H)) = n + t + |i(H)|.$$

It can also be seen that

$$S/(I(S_n \odot H), y_{n+1}) \cong K[V(S_{n-1} \odot H)]/I(S_{n-1} \odot H) \otimes_K K[i(H)] \otimes_K K[V(H)]/I(H).$$

Therefore, by using induction, Lemma 1.8 and Lemma 1.10, we get
$$depth(S/(I(S_n \odot H), y_{n+1})) = depth(K[V(S_{n-1} \odot H)]/I(S_{n-1} \odot H)) + depth(K[i(H)])$$

$$\begin{aligned} \operatorname{H}(O/(I(O_n \cup H), g_{n+1})) &= \operatorname{depth}(K[V(B_{n-1} \cup H)]/I(O_{n-1} \cup H)) + \operatorname{depth}(K[V(H)]/I(H)) \\ &= n - 1 + t + |i(H)| + t + |i(H)| \\ &= 2(t + |i(H)|) + n - 1. \end{aligned}$$

By using Lemma 1.4, we get depth $(S/(I(S_n \odot H)) \ge n + t + |i(H)|)$. For the other inequality, let $u := y_2 y_3 \cdots y_{n+1} \notin I(S_n \odot H)$, we have the following isomorphism:

(2.9)
$$S/(I(S_n \odot H) : u) \cong K[\{y_2, y_3, \dots, y_{n+1}\} \cup i(H)] \otimes_K K[V(H)]/I(H).$$

By applying Lemma 1.8 and Lemma 1.5 on Eq. 2.9, we get

$$depth(S/I(S_n \odot H)) \leq depth(S/(I(S_n \odot H) : u))$$

= depth(K[{y₂, y₃, ..., y_{n+1}} $\cup i(H)]) + depth(K[V(H)]/I(H))$
= n + |i(H)| + t.

Hence depth $(S/I(S_n \odot H)) = n + |i(H)| + t$.

Now, we prove the result for Stanley depth. If H is not a null graph, then the required lower bound of Stanley depth is obtain in a similar way as for depth by using induction, Lemma 1.8, Lemma 1.10, Lemma 2.2 and Lemma 1.4. Let H be a null graph, we have a following isomorphisms:

$$S/(I(S_n \odot H), y_{n+1}) \cong K[V(S_{n-1} \odot H)]/I(S_{n-1} \odot H) \otimes_K K[V(H)],$$

$$S/(I(S_n \odot H) : y_{n+1}) \cong \bigotimes_{l=1}^{n-1} K[V(N_1 \odot H)]/I(N_1 \odot H) \otimes_K K[V(H) \cup \{y_{n+1}\}],$$

$$S/(I(S_n \odot H) : u) \cong K[V(H) \cup \{y_2, y_3, \dots, y_{n+1}\}].$$

The proof for Stanley depth is similar to depth.

Corollary 2.18. If Stanley's inequality holds for K[V(H)]/I(H), then it also holds for $S/I(S_n \odot H)$.

Corollary 2.19. If $n \ge 1$, then

$$p\dim(S/I(S_n \odot H)) = (n+1)(|V(H)|+1) - n - depth(K[V(H)]/I(H)) - |i(H)|.$$

Proof. As $|V(S_n \odot H)| = (n+1)(|V(H)|+1)$. By using Lemma 1.11 and Theorem 2.17, the required result follows.

Corollary 2.20. Let $n, m, s \ge 1$ and $S = K[V(S_n \odot S_m)]$. Then

- (a) depth $(S/I(S_n \odot S_m)) = n + 1.$
- (b) $pdim(S/I(S_n \odot S_m)) = (n+1)(m+1).$
- (c) $\operatorname{depth}(K[V(Br_s(S_n))]/I(Br_s(S_n))) = \operatorname{sdepth}(K[V(Br_s(S_n))]/I(Br_s(S_n))) = n + s.$

For $y_{u+v} \in V(K_{u,v} \odot H)$, we have the following isomorphism:

$$(2.10) \quad K[V(K_{u,v} \odot H)]/(I(K_{u,v} \odot H) : y_{u+v}) \cong$$
$$\overset{u+v-1}{\underset{l=u+1}{\otimes}K} K[V(N_1 \odot H)]/I(N_1 \odot H) \overset{u}{\underset{l=1}{\otimes}K} K[i(H)] \overset{u}{\underset{l=1}{\otimes}K} K[V(H)]/I(H) \otimes_K K[y_{u+v}],$$

(2.11)
$$K[V(K_{u,v} \odot H)]/(I(K_{u,v} \odot H), y_{u+v}) \cong K[V(K_{u,v-1} \odot H)]/I(K_{u,v-1} \odot H) \otimes_K K[i(H)] \otimes_K K[V(H)]/I(H).$$

Theorem 2.21. If $u, v \in \mathbb{Z}^+$ and $S = K[V(K_{u,v} \odot H)]$, then

- (a) depth $(S/I(K_{u,v} \odot H)) = \min\{u, v\} (depth(K[V(H)]/I(H)) + |i(H)|) + \max\{u, v\}.$
- (b) $\operatorname{sdepth}(S/I(K_{u,v} \odot H)) \ge \min\{u, v\} (\operatorname{sdepth}(K[V(H)]/I(H)) + |i(H)|) + \max\{u, v\}.$ In particular, if H is a null graph then

$$sdepth(S/I(K_{u,v} \odot H)) = \min\{u, v\}|V(H)| + \max\{u, v\}.$$

Proof. Let $t := \operatorname{depth}(K[V(H)]/I(H))$, without loss of generality, we assume that $v \ge u$. For u = v = 1, the result follows from Theorem 2.5. If u = 1 and $v \ge 1$, then the result follows from Theorem 2.17. Let $u, v \ge 2$. By using induction on v, Lemma 1.8, Lemma 1.10 and Lemma 2.2 on Eq. 2.10 and Eq. 2.11, we have

$$depth(S/(I(K_{u,v} \odot H) : y_{u+v})) = (v-1) depth(K[V(N_1 \odot H)]/I(N_1 \odot H)) + u \cdot depth(K[i(H)]) + u \cdot depth(K[V(H)]/I(H)) + depth(K[y_{u+v}]) = v - 1 + u(|i(H)| + t) + 1 = u(|i(H)| + t) + v,$$

$$depth(S/(I(K_{u,v} \odot H), y_{u+v})) = depth(K[V(K_{u,v-1} \odot H)]/I(K_{u,v-1} \odot H)) + depth(K[i(H)]) + depth(K[V(H)]/I(H)) = u(t + |i(H)|) + v - 1 + t + |i(H)| = (u + 1)(|i(H)| + t) + v - 1.$$

	_	_	_

Since $(u+1)(|i(H)|+t)+v-1 \ge u(|i(H)|+t)+v$. By Lemma 1.4, we get depth $(S/I(K_{u,v} \odot H)) \ge 1$ u(|i(H)|+t)+v. For the other inequality, let $w := y_{u+1}y_{u+2}\cdots y_{u+v} \notin I(K_{u,v} \odot H)$, we have the following *K*-algebra isomorphism:

(2.12)
$$S/(I(K_{u,v} \odot H):w) \cong K[y_{u+1}, y_{u+2}, \dots, y_{u+v}] \bigotimes_{l=1}^{u} K[i(H)] \bigotimes_{l=1}^{u} K[V(H)]/I(H).$$

By applying Lemma 1.8, Lemma 1.10 and Lemma 1.5 on Eq. 2.12, we have

$$depth(S/I(K_{u,v} \odot H)) \leq depth(S/(I(K_{u,v} \odot H) : w))$$

= depth(K[y_{u+1}, y_{u+2}, ..., y_{u+v}]) + u \cdot depth(K[i(H)])
+ u \cdot depth(K[V(H)]/I(H))
= u(|i(H)| + t) + v.

--->

Hence depth $(S/I(K_{u,v} \odot H)) = u(|i(H)| + t) + v$. Now, we prove the result for Stanley depth. If H is not a null graph, then we get the required inequality for Stanley depth in a similar way to depth by using induction on v, Lemma 1.8, Lemma 1.10, Lemma 2.2 and Lemma 1.4 on Eq. 2.10 and Eq. 2.11. But if H is a null graph, then we have the following isomorphisms:

$$S/(I(K_{u,v} \odot H), y_{u+v}) \cong K[V(K_{u,v-1} \odot H)]/I(K_{u,v-1} \odot H) \otimes_K K[V(H)],$$

$$S/(I(K_{u,v} \odot H) : y_{u+v}) \cong \bigotimes_{k=u+1}^{u+v-1} K[V(N_1 \odot H)]/I(N_1 \odot H) \bigotimes_{k=k}^{u} K[V(H)] \otimes_K K[y_{n+1}],$$

$$S/(I(K_{u,v} \odot H) : w) \cong K[y_{u+1}, y_{u+2}, \dots, y_{u+v}] \bigotimes_{l=1}^{u} K[V(H)],$$

and the proof for Stanley depth is similar to the depth.

Corollary 2.22. If Stanley's inequality holds for K[V(H)]/I(H), then it also holds for $S/I(K_{u,v} \odot$ H).

Corollary 2.23. If $S = K[V(K_{u,v} \odot H)]$, then

$$pdim(S/I(K_{u,v} \odot H)) = (u+v)(|V(H)|+1) - \min\{u, v\} (depth(K[V(H)]/I(H)) + |i(H)|) - \max\{u, v\}$$

Proof. As $|V(K_{u,v} \odot H)| = (u+v)(|V(H)|+1)$. By using Lemma 1.11 and Theorem 2.21, we get the required result.

Lemma 2.24 ([34, Lemma 1.1]). If $S = K[V(K_{u,v})]$, then depth $(S/I(K_{u,v})) = 1$.

Corollary 2.25. Let $u, v, m, n, s \in \mathbb{Z}^+$ and $S = K[V(K_{u,v} \odot K_{m,n})]$. Then

- (a) depth $(S/I(K_{u,v} \odot K_{m,n})) = u + v.$
- (b) $pdim(S/I(K_{u,v} \odot K_{m,n})) = (u+v)(m+n).$
- (c) $\operatorname{depth}(K[V(Br_s(K_{u,v}))]/I(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))]/I(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v})))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))]/I(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v})))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v}))) = \operatorname{sdepth}(K[V(Br_s(K_{u,v})$ $\min\{u, v\}s + \max\{u, v\}.$

2.2. Regularity.

. We introduce a lemma at the beginning of this subsection, which will be used very often. It is obvious that $\operatorname{reg}(S) = 0$.

Lemma 2.26. If $S = K[V(N_1 \odot H)]$, then

$$\operatorname{reg}(S/I(N_1 \odot H)) = \begin{cases} 1, & \text{if } H \text{ is a null graph;} \\ \operatorname{reg}(K[V(H)]/I(H)), & \text{otherwise.} \end{cases}$$

Proof. If H is a null graph, then |V(H)| = |i(H)| and $N_1 \odot H \cong S_{|V(H)|}$. By Lemma 1.2, we have $\operatorname{reg}(S/I(N_1 \odot H)) = 1$. If H is not a null graph, then by Eq. 2.1, we get $\operatorname{reg}(S/(I(N_1 \odot H))) = 1$. $(y_1) = \operatorname{reg}(K[y_1]) = 0$. By using Lemma 1.8 in Eq. 2.2, we have $\operatorname{reg}(S/(I(N_1 \odot H), y_1)) = 0$ $\operatorname{reg}(K[V(H)]/I(H)) \ge 1$. By Lemma 1.13(c), we get $\operatorname{reg}(S/I(N_1 \odot H)) = \operatorname{reg}(K[V(H)]/I(H))$.

Remark 2.27. While proving the result by induction on n, we have a module of the type $K[V(P_0 \odot$ $[H]/I(P_0 \odot H)$, so in that case we define $\operatorname{reg}(K[V(P_0 \odot H)]/I(P_0 \odot H)) = 0$.

Theorem 2.28. Let $n \ge 1$. If $S = K[V(P_n \odot H)]$, then

$$\operatorname{reg}(S/I(P_n \odot H)) = \begin{cases} \left\lceil \frac{n}{2} \right\rceil, & \text{if } H \text{ is a null graph;} \\ n \cdot \operatorname{reg}(K[V(H)]/I(H)), & \text{otherwise.} \end{cases}$$

Proof. We consider two cases.

Case 1: Let H is a null graph. If n = 1, result follows by Lemma 2.26. If n = 2, we have $S/(I(P_2 \odot H): y_1) \cong K[V(H)] \otimes_K K[y_1] \text{ and } S/(I(P_2 \odot H), y_1) \cong K[V(N_1 \odot H)]/I(N_1 \odot H)$ H) $\otimes_K K[V(H)]$. We get reg $(S/(I(P_2 \odot H) : y_1)) = 0$ and by Lemma 1.8, Lemma 2.26, we have $\operatorname{reg}(S/(I(P_2 \odot H), y_1)) = \operatorname{reg}(K[V(N_1 \odot H)]/I(N_1 \odot H)) + 0 = 1$. Thus by using Lemma 1.13(c), we get $\operatorname{reg}(S/I(P_2 \odot H)) = 1$. Let $n \ge 3$. For $y_{n-1} \in V(P_n \odot H)$, we have the following K-algebra isomorphisms:

$$S/(I(P_n \odot H) : y_{n-1}) \cong K[V(P_{n-3} \odot H)]/I(P_{n-3} \odot H) \bigotimes_{k=1}^{2} K[V(H)] \otimes_{K} K[y_{n-1}]$$

$$S/(I(P_n \odot H), y_{n-1}) \cong K[V(P_{n-2} \odot H)]/I(P_{n-2} \odot H) \otimes_K K[V(N_1 \odot H)]/I(N_1 \odot H) \otimes_K K[V(H)].$$

Using induction on n and Lemma 1.8, we have

$$\operatorname{reg}(S/(I(P_n \odot H) : y_{n-1})) = \operatorname{reg}(K[V(P_{n-3} \odot H)]/I(P_{n-3} \odot H)) = \lceil \frac{n-3}{2} \rceil$$

By induction on, Lemma 1.8, Lemma 1.12 and Lemma 2.26, we get

$$\operatorname{reg}(S/(I(P_n \odot H), y_{n-1})) = \operatorname{reg}(K[V(P_{n-2} \odot H)]/I(P_{n-2} \odot H)) + \operatorname{reg}(K[V(N_1 \odot H)]/I(N_1 \odot H)) = \lceil \frac{n}{2} \rceil.$$

Since $\lceil \frac{n-3}{2} \rceil < \lceil \frac{n}{2} \rceil$, thus by using Lemma 1.13(c), we get $\operatorname{reg}(S/I(P_n \odot H)) = \lceil \frac{n}{2} \rceil$.

Case 2: Let H is not a null graph and $\operatorname{reg}(K[V(H)]/I(H)) = r$. If n = 1, we get the required result by using Lemma 2.26. Let $n \ge 2$. Considering Eq. 2.3 and Eq. 2.4 and applying Lemma 1.8, Lemma 1.12 and using induction on n, we have

$$\operatorname{reg}(S/(I(P_n \odot H) : y_n)) = \operatorname{reg}(K[V(P_{n-2} \odot H)]/I(P_{n-2} \odot H)) + \operatorname{reg}(K[V(H)]/I(H))$$

= $(n-2)r + r$
= $(n-1)r$,
$$\operatorname{reg}(S/(I(P_n \odot H), y_n)) = \operatorname{reg}(K[V(P_{n-1} \odot H)]/I(P_{n-1} \odot H)) + \operatorname{reg}(K[V(H)]/I(H))$$

= $(n-1)r + r$
= nr .

Hence by using Lemma 1.13(c), we get $\operatorname{reg}(S/I(P_n \odot H)) = nr$. This completes the proof.

Corollary 2.29. Let $n, m \ge 1$ and $q \ge 3$. Then

(a) $\operatorname{reg}(K[V(P_n \odot P_m)]/I(P_n \odot P_m)) = n \cdot \left\lceil \frac{m-1}{3} \right\rceil$. (b) $\operatorname{reg}(K[V(P_n \odot C_q)]/I(P_n \odot C_q)) = n \cdot \left\lfloor \frac{q+1}{3} \right\rfloor$. (c) $\operatorname{reg}(K[V(P_n \odot S_m)]/I(P_n \odot S_m)) = n$.

Theorem 2.30. Let $n \geq 3$. If $S = K[V(C_n \odot H)]$, then

$$\operatorname{reg}(S/I(C_n \odot H)) = \begin{cases} \left\lceil \frac{n-1}{2} \right\rceil, & \text{if } H \text{ is a null graph};\\ n \cdot \operatorname{reg}(K[V(H)]/I(H)), & \text{otherwise.} \end{cases}$$

Proof. We consider two cases.

Case 1: Let *H* is a null graph. For $y_n \in V(C_n \odot H)$, it is easy to see that

$$S/(I(C_n \odot H) : y_n) \cong K[V(P_{n-3} \odot H)]/I(P_{n-3} \odot H) \bigotimes_{l=1}^2 K[V(H)] \otimes_K K[y_n],$$

$$S/(I(C_n \odot H), y_n) \cong K[V(P_{n-1} \odot H)]/I(P_{n-1} \odot H) \otimes_K K[V(H)].$$

By using Lemma 1.8 and Theorem 2.28, we have

$$\operatorname{reg}(S/(I(C_n \odot H): y_n)) = \operatorname{reg}(K[V(P_{n-3} \odot H)]/I(P_{n-3} \odot H)) = \left\lceil \frac{n-3}{2} \right\rceil,$$

$$\operatorname{reg}(S/(I(C_n \odot H), y_n)) = \operatorname{reg}(K[V(P_{n-1} \odot H)]/I(P_{n-1} \odot H)) = \left\lceil \frac{n-1}{2} \right\rceil.$$

Since $\left\lceil \frac{n-3}{2} \right\rceil < \left\lceil \frac{n-1}{2} \right\rceil$. Hence by using Lemma 1.13(c), we have $\operatorname{reg}(S/I(C_n \odot H)) = \left\lceil \frac{n-1}{2} \right\rceil$. **Case 2:** Let *H* is not a null graph and $\operatorname{reg}(K[V(H)]/I(H)) = r$. By using Theorem 2.28,

Lemma 1.8 and Lemma 1.12 on Eq. 2.5 and Eq. 2.6, we have

$$\begin{split} \operatorname{reg}(S/I((C_n \odot H) : y_{n-1})) &= \operatorname{reg}(K[V(P_{n-3} \odot H)]/I(P_{n-3} \odot H)) + 2 \cdot \operatorname{reg}(K[V(H)]/I(H)) \\ &= (n-3)r + 2r \\ &= (n-1)r, \\ \operatorname{reg}(S/(I(C_n \odot H), y_{n-1})) &= \operatorname{reg}(K[V(P_{n-1} \odot H)]/I(P_{n-1} \odot H)) + \operatorname{reg}(K[V(H)]/I(H)) \\ &= (n-1)r + r \\ &= nr. \end{split}$$

The required result follows by using Lemma 1.13(c), thus we have $\operatorname{reg}(S/I(C_n \odot H)) = nr$.

Corollary 2.31. If $n, q \geq 3$ and $m \geq 1$, then

- (a) $\operatorname{reg}(K[V(C_n \odot C_q)]/I(C_n \odot C_q)) = n \cdot \lfloor \frac{q+1}{3} \rfloor.$ (b) $\operatorname{reg}(K[V(C_n \odot P_m)]/I(C_n \odot P_m)) = n \cdot \lceil \frac{m-1}{3} \rceil.$ (c) $\operatorname{reg}(K[V(C_n \odot K_m)]/I(C_n \odot K_m)) = n.$

Theorem 2.32. Let $n \ge 1$. If $S = K[V(K_n \odot H)]$, then

$$\operatorname{reg}(S/I(K_n \odot H)) = \begin{cases} 1, & \text{if } H \text{ is a null graph}, \\ n \cdot \operatorname{reg}(K[V(H)]/I(H)), & \text{otherwise.} \end{cases}$$

Proof. For n = 1, 2, the result follows from Lemma 2.26 and Theorem 2.28. Let $n \ge 3$. We consider two cases here.

Case 1: Let *H* is a null graph. In this case, consider Eq. 2.7 and Eq. 2.8 and applying induction on n and Lemma 1.8, we have

$$\operatorname{reg}(S/(I(K_n \odot H) : y_j)) = 0,$$

 $\operatorname{reg}(S/(I(K_n \odot H), y_j)) = \operatorname{reg}(K[V(K_{n-1} \odot H)]/(I(K_{n-1} \odot H))) = 1.$ Hence by using Lemma 1.13(c), we have $reg(S/I(K_n \odot H)) = 1$.

Case 2: If H is not a null graph and reg(K[V(H)]/I(H)) = r. By using induction, Lemma 1.8 and Lemma 1.12 on Eq. 2.7 and Eq. 2.8, we have

$$reg(S/(I(K_n \odot H) : y_j)) = (n-1) reg(K[V(H)]/I(H)) = (n-1)r,$$

$$reg(S/(I(K_n \odot H), y_j)) = reg(K[V(K_{n-1} \odot H)]/I(K_{n-1} \odot H)) + reg(K[V(H)]/I(H))$$

$$= (n-1)r + r$$

$$= nr.$$

Hence by using Lemma 1.13(c), we have $\operatorname{reg}(S/I(K_n \odot H)) = nr$.

Corollary 2.33. Let $n, m \ge 1$ and $q \ge 3$. Then

- (a) $\operatorname{reg}(K[V(K_n \odot K_m)]/I(K_n \odot K_m)) = n.$
- (b) $\operatorname{reg}(K[V(K_n \odot P_m)]/I(K_n \odot P_m)) = n \cdot \left\lceil \frac{m-1}{3} \right\rceil.$ (c) $\operatorname{reg}(K[V(K_n \odot C_q)]/I(K_n \odot C_q)) = n \cdot \left\lfloor \frac{q+1}{3} \right\rfloor.$

Theorem 2.34. If $S = K[V(S_n \odot H)]$, then

$$\operatorname{reg}(S/I(S_n \odot H)) = \begin{cases} n, & \text{if } H \text{ is a null graph;} \\ (n+1) \cdot \operatorname{reg}(K[V(H)]/I(H)), & \text{otherwise.} \end{cases}$$

Proof. For n = 1, 2, the result follows from Lemma 2.26 and Theorem 2.28. Let y_1 be a vertex of degree n in S_n , we have the following isomorphisms:

(2.13)
$$S/(I(S_n \odot H): y_1) \cong \bigotimes_{l=1}^n K[V(H)]/I(H) \bigotimes_{l=1}^n K[i(H)] \otimes_K K[y_1],$$

$$(2.14) \qquad S/(I(S_n \odot H), y_1) \cong \bigotimes_{l=1}^n K[V(N_1 \odot H)]/I(N_1 \odot H) \otimes_K K[i(H)] \otimes_K K[V(H)]/I(H).$$

We consider two cases here.

Case 1: If H is a null graph, then by using Lemma 1.8, Lemma 1.12 and Lemma 2.26 on Eq. 2.13 and Eq. 2.14, we have

 $\operatorname{reg}(S/(I(S_n \odot H), y_1)) = n \cdot \operatorname{reg}(K[V(N_1 \odot H)]/I(N_1 \odot H)) = n,$

$$\operatorname{reg}(S/(I(S_n \odot H) : y_1)) = 0$$

Hence by using Lemma 1.13(c), we have $\operatorname{reg}(S/I(S_n \odot H)) = n$. **Case 2:** If H is not a null graph and $\operatorname{reg}(K[V(H)]/I(H)) = r$. Applying Lemma 1.8, Lemma 1.12 and Lemma 2.26 on Eq. 2.13 and Eq. 2.14, we get

$$\operatorname{reg}(S/(I(S_n \odot H) : y_1)) = n \cdot \operatorname{reg}(K[V(H)]/I(H)) = nr,$$

 $\operatorname{reg}(S/(I(S_n \odot H), y_1)) = n \cdot \operatorname{reg}(K[V(N_1 \odot H)]/I(N_1 \odot H)) + \operatorname{reg}(K[V(H)]/I(H)) = nr + r.$ By Lemma 1.13(c), we get $\operatorname{reg}(S/I(S_n \odot H)) = (n+1)r$.

Corollary 2.35. If $n, m \ge 1$ and $q \ge 3$, then

- $\begin{array}{ll} \text{(a)} & \operatorname{reg}(K[V(S_n \odot S_m)]/I(S_n \odot S_m)) = n+1. \\ \text{(b)} & \operatorname{reg}(K[V(S_n \odot P_m)]/I(S_n \odot P_m)) = (n+1) \cdot \left\lfloor \frac{m-1}{3} \right\rceil. \\ \text{(c)} & \operatorname{reg}(K[V(S_n \odot C_q)]/I(S_n \odot C_q)) = (n+1) \cdot \left\lfloor \frac{q+1}{3} \right\rfloor. \end{array}$

Theorem 2.36. Let $u, v \in \mathbb{Z}^+$ and $S = K[V(K_{u,v} \odot H)]$. Then

$$\operatorname{reg}(S/I(K_{u,v} \odot H)) = \begin{cases} \max\{u, v\}, & \text{if } H \text{ is a null graph;} \\ (u+v) \cdot \operatorname{reg}(K[V(H)]/I(H)), & \text{otherwise.} \end{cases}$$

Proof. Without loss of generality, we assume that $u \ge v$. If u = v = 1, then the result follows from Theorem 2.28. For $u \ge 1$ and v = 1, the result follows from Theorem 2.34. Let $u, v \ge 2$. Here we consider two cases.

Case 1: Let H be a null graph. Let $y_{u+v} \in V(K_{u,v} \odot H)$. By using induction on v, Lemma 1.8, Lemma 1.12 and Lemma 2.26 on Eq. 2.10 and Eq. 2.11,

$$\operatorname{reg}(S/(I(K_{u,v} \odot H) : y_{u+v})) = (v-1)\operatorname{reg}(K[V(N_1 \odot H)]/I(N_1 \odot H)) = v-1,$$

 $\operatorname{reg}(S/(I(K_{u,v} \odot H), y_{u+v})) = \operatorname{reg}(K[V(K_{u,v-1} \odot H)]/I(K_{u,v-1} \odot H)) = u.$

As u > v, therefore u > v - 1. Thus using Lemma 1.13(c), we get $\operatorname{reg}(S/I(K_{u,v} \odot H)) = u$.

Case 2: Let *H* is not a null graph and $\operatorname{reg}(K[V(H)]/I(H)) = r$. For $y_1 \in V(K_{u,v} \odot H)$, we have the following *K*-algebra isomorphisms:

$$S/(I(K_{u,v} \odot H) : y_1) \cong$$

$$\overset{u-1}{\bigotimes_K} K[V(N_1 \odot H)]/I(N_1 \odot H) \overset{u+v}{\underset{l=u+1}{\otimes_K}} K[i(H)] \overset{u+v}{\underset{l=u+1}{\otimes_K}} K[V(H)]/I(H) \otimes_K K[y_1],$$

$$S/(I(K_{u,v} \odot H), y_1) \cong K[V(K_{u-1,v} \odot H)]/I(K_{u-1,v} \odot H) \otimes_K K[i(H)] \otimes_K K[V(H)]/I(H).$$

By using induction on u, Lemma 1.8, Lemma 1.12 and Lemma 2.26, we have

$$\operatorname{reg}(S/(I(K_{u,v} \odot H) : y_1)) = (u-1)\operatorname{reg}(K[V(N_1 \odot H)]/I(N_1 \odot H)) + v \cdot \operatorname{reg}(K[V(H)]/I(H)) = (u-1)r + vr$$

$$= (u + v - 1)r,$$

$$\operatorname{reg}(S/(I(K_{u,v} \odot H), y_1)) = \operatorname{reg}(K[V(K_{u-1,v} \odot H)]/I(K_{u-1,v} \odot H)) + \operatorname{reg}(K[V(H)]/I(H))$$

$$= (u + v - 1)r + r$$

$$= (u + v)r.$$

Hence by using Lemma 1.13(c), we have $\operatorname{reg}(S/I(K_{u,v} \odot H)) = (u+v)r$. This completes the proof.

Corollary 2.37. Let $u, v, m, n \in \mathbb{Z}^+$ and $q \geq 3$. Then

(a) $\operatorname{reg}(K[V(K_{u,v} \odot K_{m,n})]/I(K_{u,v} \odot K_{m,n})) = u + v.$

(b)
$$\operatorname{reg}(K[V(K_{u,v} \odot P_m)]/I(K_{u,v} \odot P_m)) = (u+v) \cdot \left|\frac{m-1}{3}\right|.$$

(c) $\operatorname{reg}(K[V(K_{u,v} \odot C_q)]/I(K_{u,v} \odot C_q)) = (u+v) \cdot \lfloor \frac{q+1}{3} \rfloor.$

3. Krull dimension and Cohen-Macaulay graphs

In this section, we find an expression for the Krull dimension of residue class rings of edge ideals associated with the corona product of two graphs if the Krull dimension of one graph is given. Moreover, by using the values of depth given in Section 2, we characterize some Cohen-Macaulay graphs.

Theorem 3.1. Let X and H be any two graphs and $S = K[V(X \odot H)]$. Then

 $\dim(S/I(X \odot H)) = |V(X)| \cdot \big(\dim(K[V(H)]/I(H)) + |i(H)|\big).$

Proof. First we assume that X is connected. By Lemma 1.1, if W is a maximum independent set of graph X ⊙ H, then dim(S/I(X ⊙ H)) = |W|. We need to prove that there exists an independent set W such that $|W| = |V(X)| \cdot (\dim(K[V(H)]/I(H)) + |i(H)|)$. Let B be an induced subgraph of X ⊙ H on vertex set $\bigcup_{j=1}^{|V(X)|} V(H_j)$, clearly B is a disjoint union of |V(X)| graphs where each graph is isomorphic to H. If W_j is a maximum independent set of the j^{th} copy of H in B, then clearly $W = W_1 \cup W_2 \cdots \cup W_{|V(X)|}$ is a maximum independent set of B. It is easy to see that $|W_j| = \dim(K[V(H)]/I(H)) + |i(H)|$. We claim that W is a maximum independent set of X ⊙ H such that |W'| > |W|. This implies that there exists another maximum independent set W' of X ⊙ H such that |W'| > |W|. This implies that there exists a vertex $z \in W'$ such that $z \notin W$. Then we have two cases to discuss that is either $z \in V(X)$ or $z \in \bigcup_{j=1}^{|V(X)|} V(H_j)$. If $z \in \bigcup_{j=1}^{|V(X)|} V(H_j)$ then for some $j, z \notin W_j$ where W_j is a maximum independent set of induced subgraph of X ⊙ H on $V(H_j)$. This implies that z is adjacent to some vertex in W_j . This is a contradiction to the assumption that W' is an independent set. Now assume that $z \in V(X)$. This means that for some j, we have $z = y_j$. By Definition 1.14, y_j is pairwise adjacent to each vertex in $V(H_j)$, therefore $W' \subset$ $V(X ⊙ H) \setminus N_{X ⊖ H}(y_j)$. Since $|N_{X ⊖ H}(y_j)| \ge 1$, then $|V(X ⊙ H) \setminus \{y_j\}| \ge |V(X ⊙ H) \setminus N_{X ⊖ H}(y_j)|$.

This implies that the cardinality of the maximum independent set of induced subgraph of $X \odot H$ on $V(X \odot H) \setminus \{y_i\}$ is strictly greater than or equals to the cardinality of a maximum independent set of induced subgraph of $X \odot H$ on $V(X \odot H) \setminus N_{X \odot H}(y_j)$. Clearly, $W \subset V(X \odot H) \setminus \{y_j\}$ and $W' \subset V(X \odot H) \setminus N_{X \odot H}(y_i)$. We get $|W| \geq |W'|$, which contradicts our assumption. If X is a not a connected graph, that is, X is a union of disjoint connected components say X_1, \ldots, X_l , then by [24, Lemma 1.11(3)], we have $\dim(S/I(X \odot H)) = \sum_{k=1}^{l} \dim(K[V(X_k \odot H)]/I(X_k \odot H))$ and our result follows.

Corollary 3.2. Let $m, n, u, v, r \in \mathbb{Z}^+$ and $l, q \geq 3$. Then

- (a) $\dim(K[V(P_n \odot P_m)]/I(P_n \odot P_m)) = n \cdot \left\lceil \frac{m}{2} \right\rceil.$ (b) $\dim(K[V(C_l \odot C_q)]/I(C_l \odot C_q)) = l \cdot \left\lceil \frac{q-1}{2} \right\rceil.$ (c) $\dim(K[V(S_n \odot S_m)]/I(S_n \odot S_m)) = (n+1) \cdot m.$
- (d) $\dim(K[V(K_n \odot K_m)]/I(K_n \odot K_m)) = n.$
- (e) $\dim(K[V(K_{m,n} \odot K_{u,v}))]/I(K_{m,n} \odot K_{u,v})) = (m+n) \cdot \max\{u, v\}.$
- (f) If G is a disjoint union of C_q and N_r , then $\dim(K[V(P_n \odot G))]/I(P_n \odot G)) = n \cdot (\lceil \frac{q-1}{2} \rceil + r)$.

Lemma 3.3. Let H be a non-trivial connected graph and S = K[V(H)]. Then $\dim(S/I(H)) = 1$ iff H is complete graph.

Proof. Let W be the maximum independent set of H. If $\dim(S/I(H)) = 1$ (i.e |W| = 1) implies that all the vertices of H are pairwise adjacent which proves that H is a complete graph. The converse statement easily followed by Lemma 1.1.

If H is a trivial graph, then $X \odot H$ is a whisker graph of X. Villarreal proved in [41, Proposition 2.2] that whisker graph of any graph is Cohen-Macaulay graph. So in our next theorem, we characterize all Cohen-Macaulay graphs $X \odot H$, if $X \in \{P_n, C_n, K_n, S_{n+1}, K_{u,v}\}$.

Theorem 3.4. Let H be any graph, $X \in \{P_n, C_n, K_n, S_{n+1}, K_{u,v}\}$ and $S = K[V(X \odot H)]$. Then $S/I(X \odot H)$ is Cohen-Macaulay iff H is a complete graph.

Proof. By Lemma 3.3, a non-trivial connected graph H is complete iff $\dim(K[V(H)]/I(H)) =$ depth(K[V(H)]/I(H)) = 1. We discuss all the cases one by one as follows:

1: Let $X = P_n$. By Theorem 2.5 and Theorem 3.1, the module $S/I(P_n \odot H)$ is Cohen-Macaulay iff depth $(S/I(P_n \odot H)) = \dim(S/I(P_n \odot H))$ iff

$$\left\lceil \frac{n}{2} \right\rceil + \left\lceil \frac{n-1}{2} \right\rceil \left(\operatorname{depth}(K[V(H)]/I(H)) + |i(H)| \right) = n \cdot \left(\operatorname{dim}(K[V(H)]/I(H)) + |i(H)| \right)$$

iff |i(H)| = 1 and depth $(K[V(H)]/I(H)) = 0 = \dim(K[V(H)]/I(H))$ or |i(H)| = 0 and $\dim(K[V(H)]/I(H)) = \operatorname{depth}(K[V(H)]/I(H)) = 1$ iff H is a complete graph.

2: If $X = C_n$, then by using Theorem 2.9 and Theorem 3.1, the module $S/I(C_n \odot H)$ is Cohen-Macaulay iff depth $(S/I(C_n \odot H)) = \dim(S/I(C_n \odot H))$ iff

$$\lceil \frac{n-1}{2} \rceil + \lceil \frac{n}{2} \rceil \left(\operatorname{depth}(K[V(H)]/I(H)) + |i(H)| \right) = n \cdot \left(\operatorname{dim}(K[V(H)]/I(H)) + |i(H)| \right)$$
 iff $|i(H)| = 1$ and $\operatorname{depth}(K[V(H)]/I(H)) = \operatorname{dim}(K[V(H)]/I(H)) = 0$ or $|i(H)| = 0$ and

$$\dim(K[V(H)]/I(H)) = \operatorname{depth}(K[V(H)]/I(H)) = 1 \text{ iff } H \text{ is a complete graph.}$$

3: Let $X = K_n$. By Theorem 2.13 and Theorem 3.1, the module $S/I(K_n \odot H)$ is Cohen-Macaulay iff depth $(S/I(K_n \odot H)) = \dim(S/I(K_n \odot H))$ iff

$$1 + (n-1)\left(\operatorname{depth}(K[V(H)]/I(H)) + |i(H)|\right) = n \cdot \left(\operatorname{dim}(K[V(H)]/I(H)) + |i(H)|\right)$$

iff |i(H)| = 1 and depth $(K[V(H)]/I(H)) = \dim(K[V(H)]/I(H)) = 0$ or |i(H)| = 0 and $\dim(K[V(H)]/I(H)) = \operatorname{depth}(K[V(H)]/I(H)) = 1$ iff H is a complete graph.

4: If $X = S_n$, then by Theorem 2.17 and Theorem 3.1, we have $S/I(S_n \odot H)$ is Cohen-Macaulay iff depth $(S/I(S_n \odot H)) = \dim(S/I(S_n \odot H))$ iff

 $\begin{aligned} n + \operatorname{depth}(K[V(H)]/I(H)) + |i(H)| &= (n+1) \cdot \left(\dim(K[V(H)]/I(H)) + |i(H)| \right) \\ \operatorname{iff} |i(H)| &= 1 \text{ and } \operatorname{depth}(K[V(H)]/I(H)) = \dim(K[V(H)]/I(H)) = 0 \text{ or } |i(H)| = 0 \text{ and} \\ \operatorname{dim}(K[V(H)]/I(H)) &= \operatorname{depth}(K[V(H)]/I(H)) = 1 \text{ iff } H \text{ is a complete graph.} \end{aligned}$

- 5: Let $X = K_{u,v}$. Without loss of generality, we assume that $v \ge u$. Using Theorem 2.21 and Theorem 3.1, the module $S/I(K_{u,v} \odot H)$ is Cohen-Macaulay iff depth $(S/I(K_{u,v} \odot H)) = \dim(S/I(K_{u,v} \odot H))$ iff
- $$\begin{split} & u \cdot \big(\operatorname{depth}(K[V(H)]/I(H)) + |i(H)| \big) + v = (u+v) \cdot \big(\operatorname{dim}(K[V(H)]/I(H)) + |i(H)| \big) \\ & \text{iff } |i(H)| = 1 \text{ and } \operatorname{depth}(K[V(H)]/I(H)) = \operatorname{dim}(K[V(H)]/I(H)) = 0 \text{ or } |i(H)| = 0 \text{ and } \\ & \operatorname{dim}(K[V(H)]/I(H)) = 1 = \operatorname{depth}(K[V(H)]/I(H)) \text{ iff } H \text{ is a complete graph.} \end{split}$$

Let $G_1 \sqcup G_2$ denotes a disjoint union of two graphs G_1 and G_2 . By using Theorem [20, Theorem 3.2], we have the following corollary.

Corollary 3.5. Let H be any graph and G be a graph such that $G = G_1 \sqcup G_2 \sqcup \cdots \sqcup G_k$, where $G_i \in \{P_n, C_n, K_n, S_{n+1}, K_{u,v}\}$. Then $K[V(G \odot H)]/I(G \odot H)$ is Cohen-Macaulay iff H is a complete graph.

CONCLUSION

In last decades, edge ideals have garnered significant attention; see for instance [40]. Various findings on these edge ideals have demonstrated how combinatorial and algebraic aspects interact. The primary goal of this paper is to compute algebraic invariants such as depth, Stanley depth, regularity, projective dimension and Krull dimension of edge ideals associated with the corona product of two graphs and as a consequence of our findings, we characterize some Cohen–Macaulay graphs. By restricting one of the class to a well-known graph in corona product of two graphs, we are able to handle the algebraic invariants for these structures, and our results gives strong motivation for further studies into this intriguing area to compute the algebraic invariants when both classes in corona product of two graphs are arbitrary.

References

- [1] Ahmad, S., Anwar, I., Abbas, F. (2019). Cohen Macaulay Hybrid Graphs. arXiv preprint arXiv:1904.03824.
- [2] Ahmad, S., Kanwal, S. (2019). A Construction of Cohen-Macaulay Graphs. Studia Scientiarum Mathematicarum Hungarica, 56(4), 492-499.
- [3] Alipour, A., and Tehranian, A. (2017). Depth and Stanley depth of edge ideals of star graphs. International Journal of Applied Mathematics and Statics, 56(4), 63-69.
- [4] Bouchat, R. R. (2010). Free resolutions of some edge ideals of simple graphs. Journal of Commutative Algebra, 2(1), 1-35.
- [5] Bruns, W., Herzog, H. J. (1998). Cohen-Macaulay rings. Cambridge University Press.
- [6] Caviglia, G., Hà, H. T., Herzog, J., Kummini, M., Terai, N., Trung, N. V. (2019). Depth and regularity modulo a principal ideal. *Journal of Algebraic Combinatorics*, 49(1), 1-20.
- [7] Cimpoeas, M. (2012). Several inequalities regarding Stanley depth. Romanian Journal of Math. and Computer Science, 2(1), 28-40.
- [8] Cimpoeas, M. (2013). Stanley depth of squarefree Veronese ideals. Analele Universitatii" Ovidius" Constanta-Seria Matematica, 21(3), 67-72.
- [9] Dao, H., Huneke, C., Schweig, J. (2013). Bounds on the regularity and projective dimension of ideals associated to graphs. Journal of Algebraic Combinatorics, 38(1), 37-55.
- [10] Din, N. U., Ishaq, M., Sajid, Z. (2021). Values and bounds for depth and Stanley depth of some classes of edge ideals. AIMS Mathematics,6(8), 8544-8566.
- [11] Duval, A. M., Goeckner, B., Klivans, C. J., Martin, J. L. (2016). A non-partitionable Cohen–Macaulay simplicial complex. Advances in Mathematics, 299, 381-395.
- [12] Eisenbud, D. (2013). Commutative algebra: with a view toward algebraic geometry (Vol. 150). Springer Science, Business Media.

- [13] Estrada, M., Villarreal, R. H. (1997). Cohen-Macaulay bipartite graphs. Archiv der Mathematik, 68(2), 124-128.
- [14] Fakhari, S. S. (2017). On the Stanley depth of powers of edge ideals. Journal of Algebra, 489, 463-474.
- [15] Faridi, S., Hersey, B. (2017). Resolutions of monomial ideals of projective dimension 1. Communications in Algebra, 45(12), 5453-5464.
- [16] Fouli, L., Morey, S. (2015). A lower bound for depths of powers of edge ideals. Journal of Algebraic Combinatorics, 42(3), 829-848.
- [17] Frucht, R., Harary, F. (1970). On the corona of two graphs, Aeq. Math, 4, 322–32.
- [18] Galetto, F., Hofscheier, J., Keiper, G., Kohne, C., Van Tuyl, A., Paczka, M. E. U. (2019). Betti numbers of toric ideals of graphs: a case study. Journal of Algebra and its Applications, 18(12), 1950226.
- [19] Hà, H. T., Van Tuyl, A. (2008). Monomial ideals, edge ideals of hypergraphs, and their graded Betti numbers. Journal of Algebraic Combinatorics, 27(2), 215-245.
- [20] Haghighi, H., Yassemi, S., Zaare Nahandi, R. (2015). Cohen-Macaulay bipartite graphs in arbitrary codimension. Proceedings of the American Mathematical Society, 143(5), 1981-1989.
- [21] Herzog, J., Hibi, T., Zheng, X. (2006). Cohen-Macaulay chordal graphs. Journal of Combinatorial Theory, 911-916.
- [22] Herzog, J., Hibi, T. (2005). Distributive lattices, bipartite graphs and Alexander duality. Journal of Algebraic Combinatorics, 22(3), 289-302.
- [23] Herzog, J., Vladoiu, M., Zheng, X. (2009). How to compute the Stanley depth of a monomial ideal. Journal of Algebra, 322(9), 3151-3169.
- [24] Hibi, T., Kanno, H., Matsuda, K. (2019). Induced matching numbers of finite graphs and edge ideals. Journal of Algebra, 532, 311-322.
- [25] Hibi, T., Kanno, H., Kimura, K., Matsuda, K., Van Tuyl, A. (2021). Homological invariants of Cameron–Walker graphs. Transactions of the American Mathematical Society, 374(09), 6559-6582.
- [26] Hirano A., Matsuda K. (2021). Matching numbers and dimension edge ideals. Graphs and Combinatorics, 37(3), 761-774. 2021.
- [27] Hoa, L. T., Tam, N. D. (2010). On some invariants of a mixed product of ideals. Archiv der Mathematik, 94(4), 327-337.
- [28] Ishaq, M., Qureshi, M. I. (2013). Upper and lower bounds for the Stanley depth of certain classes of monomial ideals and their residue class rings. *Communications in Algebra*, 41(3), 1107-1116.
- [29] Iqbal, Z., Ishaq, M., Aamir, M. (2018). Depth and Stanley depth of the edge ideals of square paths and square cycles. Communications in Algebra, 46(3), 1188-1198.
- [30] Kiani, D., Saeedi Madani, S. (2015). Some Cohen–Macaulay and unmixed binomial edge ideals. Communications in Algebra, 43(12), 5434-5453.
- [31] Mahmood, H., Anwar, I., Zafar, M. K. (2014). A construction of Cohen–Macaulay f-graphs. Journal of Algebra and its Applications, 13(06), 1450012.
- [32] Morey, S. (2010). Depths of powers of the edge ideal of a tree. Communications in Algebra, 38(11), 4042-4055.
- [33] Morey, S., Villarreal, R. H. (2012). Edge ideals: algebraic and combinatorial properties. Progress in commutative algebra, 1, 85-126.
- [34] Popescu, A. (2010). Special stanley decompositions. Bulletin mathématique de la Société des Sciences Mathématiques de Roumanie, 363-372.
- [35] Pournaki, M., Seyed Fakhari, S. A., Yassemi, S. (2013). Stanley depth of powers of the edge ideal of a forest. Proceedings of the American Mathematical Society, 141(10), 3327-3336.
- [36] Rauf, A. (2010). Depth and Stanley depth of multigraded modules. Communications in Algebra, 38(2), 773-784.
- [37] Shaukat, B., Haq, A. U., Ishaq, M. (2022). Some algebraic Invariants of the residue class rings of the edge ideals of perfect semiregular trees. Communications in Algebra, 1-20.
- [38] Stanley, R. P., RP, S. (1982). Linear Diophantine equations and local cohomology. Invent. Math., 68(2), (1982) 175-193.
- [39] Vander Meulen, K. N., Van Tuyl, A., Watt, C. (2014). Cohen-Macaulay circulant graphs. Communications in Algebra, 42(5), 1896-1910.
- [40] Van Tuyl, A. A beginner's guide to edge and cover ideals. Monomial ideals, computations and applications, 63–94. Lecture Notes in Math, 2083.
- [41] Villarreal, R. H. (1990). Cohen-macaulay graphs. manuscripta mathematica, 66(1), 277-293.
- [42] Villarreal, R. H.(2001). Monomial Algebras. Monographs and Textbooks in Pure and Applied Mathematics. New York: Marcel Dekker, Inc., Vol. 238.