

The reducibility of optimal 1-planar graphs ^{*}

Licheng Zhang [†], Yuanqiu Huang [‡]

[†] School of Mathematics, Hunan University

Changsha, Hunan, 410082, P.R. China

[‡] School of Mathematics and Statistics, Hunan Normal University

Changsha, Hunan, 410081, P.R. China

Abstract

A graph is reducible if it is the lexicographic product of two smaller non-trivial graphs. It is well-known that a 1-planar graph with $n \geq 3$ vertices has at most $4n - 8$ edges, and a 1-planar graph G with n vertices is optimal if G has exactly $4n - 8$ edges. In this paper, we characterize the reducibility of optimal 1-planar graphs. This work is motivated by a problem posed by Bucko and Czap in 2015, which concerns determining the 1-planarity of the lexicographic product of a graph and two isolated vertices.

Keywords: lexicographic product, 1-planarity, reducibility

MSC: 05C10, 05C62, 05C76

1 Introduction

1.1 The study of 1-planarity of graphs

All graphs considered here are simple and finite. Let $G = (V, E)$ be a graph. Its order and size are $|V|$ and $|E|$, respectively. A drawing of G is a mapping D that assigns to each vertex in V a distinct point in the plane and to each edge uv in E a continuous arc connecting $D(u)$ and $D(v)$. A drawing of a graph is *1-planar* if each edge is crossed at most once. A graph is *1-planar* if it has a 1-planar drawing, and such drawing is called a *1-plane graph*. 1-planar graphs were first studied by Ringel (1965) [14]. Since then, many properties of 1-planar graphs have been widely investigated; see [6, 10, 18] for examples, or [11] for a survey.

^{*}E-mail addresses: lc Zhang@163.com (L. Zhang), hyqq@hunnu.edu.cn (Y. Huang, corresponding author).

Determining the 1-planarity of a graph is a fundamental problem in the topic of 1-planar graphs. Unfortunately, determining the 1-planarity of a graph is NP-complete [12], and it has been pointed out in [13] that 1-planar graphs cannot be characterized by using a finite set of forbidden (topological) minors, unlike the well-known criterion (Wagner's theorem and Kuratowski's theorem) for planar graphs. Based on these two main reasons, the 1-planarity of only a few graphs has been determined, and even for graphs with given a small order, their 1-planarity remains unknown [2]. In addition, few necessary conditions for 1-planarity are known. Currently, researchers are focusing on specific classes of graphs (with well-defined structures), including operations on graphs, to determine their 1-planarity. For examples, Czap and Hudák determined the 1-planarity of complete multipartite graphs [5]. Czap, Hudák and Madaras [7] studied the 1-planarity of the join of two graphs.

Definition 1. A lexicographic product $G \circ H$ of two graphs G and H is a graph such that the vertex set of $G \circ H$ is the Cartesian product $V(G) \times V(H)$ and two vertices (g_1, h_1) and (g_2, h_2) are adjacent in $G \circ H$ if and only if

- g_1 is adjacent to g_2 in G , or
- $g_1 = g_2$ and h_1 is adjacent to h_2 in H .

In 2015, Bucko and Czap [4] discussed the 1-planarity of lexicographic products of graphs. The authors propose a conjecture: A graph $G \circ K_2$ is 1-planar if and only if G is a *cactus*, which is a connected graph in which every edge belongs to at most one cycle. Recently, Matsumoto and Suzuki [13] confirmed the conjecture. Thus, combining the works [4] and [13], the 1-planarity of the lexicographic product of two graphs leaves only the following unsolved problem.

Problem 1 ([4]). Let G be a connected graph with maximum degree at least 3. Characterize the 1-planarity of $G \circ 2K_1$.

Note that the lexicographic product is not commutative [9, Chapter 5, Page 56], i.e., $G \circ H \neq H \circ G$ in general. A graph is *trivial* if it has just one vertex or no vertex. A graph is called *reducible* if it is the lexicographic product of two non-trivial graphs, otherwise is called *irreducible*¹. It is well-known that a 1-planar graph with $n(\geq 3)$ vertices has at most $4n - 8$ edges, and a 1-planar graph with n vertices is *optimal* if it has exactly $4n - 8$ edges. This means that optimal 1-planar graphs are a class of 1-planar graphs with the maximum number of edges. In this paper, we focus on characterizing the reducibility of optimal 1-planar graphs. We shall show that, with the exception of a single graph $K_{2,2,2,2}$, all optimal 1-planar

¹In general, determining whether a graph is reducible is as difficult as determining whether two graphs are isomorphic [8].

graphs are not reducible. As a byproduct, we provide the upper bound on the size of the left factor G for the graph $G \circ 2K_1$ to be 1-planar, which will serve as a necessary condition for Problem 1.

Theorem 1. *Any optimal 1-planar graph G is not reducible unless $G \cong K_{2,2,2,2}$.*




Fig. 1. A 1-planar drawing of $K_{2,2,2,2}$

Theorem 2. *Let G be a graph of order $n \geq 5$. If $G \circ 2K_1$ is 1-planar, then $|E(G)| \leq 2n - 3$.*

In a certain sense, Theorem 1 subtly suggests that optimal 1-planar graphs lack strong symmetry (from the perspective of the lexicographic product), as the structure of the graph $G \circ H$ implicitly encodes numerous copies of the right factor H and complete bipartite subgraphs rooted in the left factor G . On the other hand, the upper bound on the size of a reducible 1-planar graph is also naturally given. Interestingly, by reducing the number of edges by just one, there exist infinitely many reducible 1-planar graphs.

Theorem 3. *Let G be a 1-planar graph of order $n \geq 9$. If G is reducible, then $|E(G)| \leq 4n - 9$. Moreover, the bound for n being divisible by 3 is tight.*

1.2 Terminology and notations

The terms not defined here can be found in [17]. Let K_n denote the complete graph of order n , respectively. The disjoint union of k copies of a graph G is denoted by kG . A k -cycle C_k is a cycle of with k vertices.

Let G be a graph. If S is a set of vertices of G , the vertex-induced subgraph $G[S]$ is the subgraph of G that has S as its set of vertices and contains all the edges of G that have both end-vertices in S . We denote by $G \cong H$ that graphs G and H are isomorphic. We denoted by K_{n_1, n_2, \dots, n_k} a complete multipartite graph with k partition classes of sizes n_1, n_2, \dots, n_k , respectively. We call a cycle C of G *even* if $|V(C)|$ is even.

We call G and H the *left* and *right factors* of the lexicographic product $G \circ H$, respectively. The lexicographic product is also known as graph substitution, since

$G \circ H$ is obtained from G by substituting a copy H_u of H for every vertex u of G and then joining all vertices of H_u with all vertices of H_v if $uv \in E(G)$ ².

All graphs considered here are simple. A drawing is *good* if it satisfies the following three conditions:

- No edge crosses itself.
- No two edges cross more than once.
- No two edges incident with a common vertex cross.

Let G be a 1-planar graph with a 1-planar drawing D . The rotation $\text{Rot}_D(u)$ of a vertex u in D as the cyclic permutation that records the (cyclic) counter-clockwise order in which the edges leave u . The rotation system naturally defines two *successive* edges incident with u if they appear consecutively in $\text{Rot}_D(u)$. An edge e of G is called *crossed* in D if it crosses with any other edge in D , and is *uncrossed* in D otherwise. A cycle C of G is *uncrossed* in D if each edge on C is uncrossed in D , and is *crossed* in D otherwise. The *planar skeleton* $S(D)$ of D is the subgraph of G by removing all crossed edges of D . Let H be a subgraph of G . The subdrawing $D|H$ of H induced by D is called a *restricted drawing* of D . Let \mathcal{P} be a plane. Similar to planar drawings, the 1-planar drawing D also defines the *faces*, which are the connected parts of $\mathcal{P} \setminus D$. Each face f contains on its boundary a number of vertices and crossings of D ; these are called the *corners* of f . A face is *uncrossed* if all incident corners are vertices, and is *crossed* otherwise. A *k-face* is a face whose boundary walk has a length of exactly k . Let L be a closed curve in \mathcal{P} that does not cross itself. Thus, L separates \mathcal{P} into two open regions, the bounded one (i.e., the interior of L) and the unbounded one (i.e., the exterior of L). We denote by L_{int} and L_{out} the interior and exterior of L , respectively.

The structure of the remaining sections is as follows. In the next section, we first introduce auxiliary lemmas, including the 1-planarity of lexicographic products and elementary properties on optimal 1-planar graphs. In Section 3 and 4, we characterize the reducibility of optimal 1-planar graphs, depending on the order of the left and right factors. In the proof, we focus on the drawing structures of two types of small subgraphs: C_4 and $K_{3,3}$, along with their surrounding structures.

2 Preliminaries

First, we introduce two lemmas regarding the properties of the lexicographic product of graphs. Lemma 1(i) can be found in [9, Chapter 5, page 58], and Lemma 1(ii) follows directly from the definition of lexicographic product.

²In some literature, lexicographic product $G \circ H$ is a special case of the blow-up of G , where each vertex of G is blown up by the same subgraph H .

Lemma 1 ([9]). *Let G and H be graphs. Then the following statements hold.*

- (i) $G \circ H$ is connected if and only if G is connected, and
- (ii) $|E(G \circ H)| = |V(G)| \times |E(H)| + |V(H)|^2 \times |E(G)|$.

Since optimal 1-planar graphs are connected (in fact, 4-connected [15]), the following lemma naturally follows from Lemma 1(i).

Lemma 2. *Let G be an optimal 1-planar graph. If $G = G_1 \circ G_2$, then G_1 is connected.*

The following Lemma 3 directly follows from the definition of lexicographic products.

Lemma 3. *Let G and H be graphs. Let $V(G) = \{g_1, g_2, \dots, g_s\}$ and $V(H) = \{h_1, h_2, \dots, h_l\}$. If graph $G' = G \circ H$, then the following statements hold.*

- (i) For $1 \leq i \leq j \leq s$, $G'[\{(g_i, h_1), (g_i, h_2), \dots, (g_i, h_l)\}] \cong G'[\{(g_j, h_1), (g_j, h_2), \dots, (g_j, h_l)\}] \cong H$, and
- (ii) if a vertex u in G' is adjacent to (g_k, h_i) where $1 \leq k \leq s$ and $1 \leq i \leq l$, then u is adjacent to each vertex in $\{(g_k, h_1), (g_k, h_2), \dots, (g_k, h_l)\}$.

We present some lemmas on the 1-planarity of lexicographic products of graphs from [4] and [13].

Lemma 4 ([4]). *Let G be a connected graph with $|V(G)| \geq 3$, and let H be a graph with $|V(H)| \geq 4$. Then $G \circ H$ is not 1-planar.*

Lemma 5 ([4]). *Let H be a graph and let $G = K_2$. Then the following statements hold.*

- (i) If $|V(H)| \geq 5$, then $G \circ H$ is not 1-planar, and
- (ii) if $|V(H)| \leq 4$, then H is a subgraph of C_4 or H is a subgraph of C_3 .

Lemma 6 ([4, 13]). *Let G be a graph. Then $G \circ K_2$ is 1-planar if and only if G is a cactus.*

The following result on 1-planar graphs is fundamental and has been independently obtained by many researchers.

Lemma 7 ([3, 15]). *Let G be a 1-planar graph of order n . Then*

$$|E(G)| \leq \begin{cases} \binom{n}{2} & \text{if } n \leq 6; \\ 4n - 9 & \text{if } n = 7 \text{ or } 9; \\ 4n - 8 & \text{otherwise.} \end{cases}$$

Two drawings of a graph are *isomorphic* if there is a homeomorphism of the sphere that maps one drawing to the other; otherwise, the drawings are *non-isomorphic*.³ In this following, we give some properties of optimal 1-planar graphs. A plane graph is a *quadrangulation* if every face is a 4-face. References [3, 15] gave a relationship between optimal 1-planar graphs and 3-connected quadrangulations.

Lemma 8 ([3],[15]). *Let G be an optimal 1-plane graph. Then D is obtained by inserting a pair of crossed edges to each 4-face of a 3-connected quadrangulation. Moreover, the 1-planar drawings of G are unique (up to isomorphism).*

Lemma 8 provides powerful structural information on optimal 1-planar graphs, which directly lead to the following Lemmas 9, 10, and 11.

Lemma 9. *Let G be an optimal 1-planar graph with a 1-planar drawing D . Then edges incident with a vertex $v \in V(G)$ appear successively as crossed and uncrossed edges in $\text{Rot}_D(v)$.*

Lemma 10. *Let G be an optimal 1-planar graph with a 1-planar drawing D . Let v_0v_2 and v_1v_3 be edges of G , which cross each other in point c in D . Then the following two statements hold.*

- (i) *The induced subgraph $H := G[\{v_0, v_1, v_2, v_3\}] \cong K_4$, and any edge in $E(H) \setminus \{v_0v_2, v_1v_3\}$ is uncrossed in D , and*
- (ii) *cv_iv_{i+1} bounds a crossed face of D with indices taken modulo 4.*

Lemma 11 ([16]). *Let G be an optimal 1-planar graph with a 1-planar drawing D . If C is an uncrossed cycle in D . Then C is even.*

The following lemma reveals that optimal 1-planar graphs do not exhibit hereditary properties with respect to optimality⁴, which will be used in the final step of the proof of Proposition 1.

Lemma 12. *Any properly subgraph of an optimal 1-planar graph G is not optimal 1-planar.*

Proof. Let H be a properly subgraph of G . Suppose H is optimal 1-planar with a 1-planar drawing D' . Clearly, since H is optimal, $G[V(H)] = H$. Since D' is unique, then $D|D'$ is fixed. by Lemma 8, every face of D is a crossed 3-face. Since G is 4-connected [15], any 3-face of D' also bounds a 3-face of D , that is, D has no vertices from $G \setminus H$. This implies that $H \cong G$, a contradiction. \square

³In fact, the sphere and the plane differ topologically. However, a graph is 1-planar if and only if it can be embedded on the sphere such that each edge crosses at most once, which can be achieved through stereographic projection. The only difference is that a drawing D in the sphere ensures that each face of D is a finite face. This definition of isomorphism for graph drawings is given on the sphere rather than the plane to avoid concerns about the external infinite face of a 1-planar drawing.

⁴A hereditary property typically refers to a property of a graph that is also preserved in its (induced) subgraphs.

The following lemma which gives two distinct 1-planar drawings of $K_{3,3}$, particularly the first one, will be used in Section 4.

Lemma 13 ([1]). *There are exactly two non-isomorphic 1-planar drawings of $K_{3,3}$, as shown in Fig. 2.*




Fig. 2. Two non-isomorphic 1-planar drawings of $K_{3,3}$

3 Left or right factor with order two

In this section, we shall prove that an optimal 1-planar graph, when reduced into factors where at least one of the left or right factors has order two, can only be isomorphic to $K_{2,2,2,2}$.

Before that, we can easily verify the following simple lemma.

Lemma 14. *The graph $K_{2,2,2,2}$ is optimal 1-planar and a 1-planar drawing of $K_{2,2,2,2}$ is shown in Fig. 1. Moreover $K_{2,2,2,2} = K_2 \circ C_4$ and $K_{2,2,2,2} = K_4 \circ 2K_1$.*

Proposition 1. *Let G be an optimal 1-planar graph. If $G = G_1 \circ G_2$, then, $G \cong K_{2,2,2,2}$ if either of the following holds:*

- (i) $|V(G_1)| = 2$ and $|V(G_2)| \geq 2$;
- (ii) $|V(G_1)| \geq 3$ and $|V(G_2)| = 2$.

Proof. If (i) holds, we have the following Claim.

Claim 1.1. G_2 is C_4 or a C_3 with a pendant edge.

Proof. As G is connected, by Lemma 2, it follows that G_1 is connected, and thus $G_1 \cong K_2$. By Lemma 5, we have $|V(G_2)| \leq 4$. Since $|V(G)| \geq 8$ (by Lemma 7) and $G_1 \cong K_2$, we have $|V(G_2)| \geq 4$. So $|V(G_2)| = 4$. Thus $|V(G)| = 8$. By the optimality of G , we have $|E(G)| = 24$. Hence, by Lemma 1, we have $|E(G_2)| \geq 4$. So G_2 is C_4 or a C_3 with a pendant edge. \square

Furthermore, by Lemma 5(ii), $G_2 \cong C_4$, and thus $G \cong K_{2,2,2,2}$.

Now, we consider the case where condition (ii) of the proposition is satisfied. The situation becomes a bit more complicated. We shall deduce that G is isomorphic to $K_{2,2,2,2}$ through a series of claims, starting from replacing a single edge from the left factor G_1 with a cycle of length four (note that the right factor shall be consist of two isolated vertices by the following first claim). We will see that the drawing structure of the 4-cycle in D and its surrounding elements play a key role.

Claim 1.2. $|V(G_1)| \geq 4$ and $G_2 \cong 2K_1$.

Proof. By Lemma 7, one has $|V(G)| \geq 8$. Furthermore, since $|V(G_2)| = 2$, it follows that $|V(G_1)| \geq 4$. Suppose $G_2 \cong K_2$. By Lemma 1 we have

$$|E(G)| = |V(G_1)| + 4 \times |E(G_1)|. \quad (3.1)$$

On the other hand, by the optimality of G , we have

$$|E(G)| = 4 \times (2 \times |V(G_1)|) - 8. \quad (3.2)$$

By combining (3.1) with (3.2), it follows that

$$|E(G_1)| = \frac{7}{4}|V(G_1)| - 2. \quad (3.3)$$

By Lemma 6, G_1 is a cactus. Then, by a well-known result on the sizes of cactus graphs ([17, Page 160]), we have

$$|E(G_1)| \leq \left\lfloor \frac{3(|V(G_1)| - 1)}{2} \right\rfloor. \quad (3.4)$$

By (3.3) and (3.4), we have $|V(G_1)| \leq 2$, a contradiction to $|V(G_1)| \geq 4$. Hence, $G_2 \cong 2K_1$. \square

So we assume that $V(G_1) := \{u_1, u_2, \dots, u_n\}$ where $n \geq 4$ and $V(G_2) := \{v_1, v_2\}$. By Lemma 1(i), G_1 is connected. Hence, $E(G_1) \neq \emptyset$, and we may assume that $u_i u_j \in E(G_1)$ for $1 \leq i \neq j \leq n$. Then u_i (resp. u_j) is substituted with two distinct vertices (u_i, v_1) and (u_i, v_2) in G (resp. (u_j, v_1) and (u_j, v_2) in G). For simplicity, let $u := (u_i, v_1)$, $v := (u_i, v_2)$, $x := (u_j, v_1)$ and $y := (u_j, v_2)$. Let $V(G) \setminus \{u, v, x, y\} := \{w_1, w_2, \dots, w_{2n-4}\}$.

Claim 1.3. *If a vertex p of G is adjacent to one vertex of $\{u, v\}$ (resp. of $\{x, y\}$), then p is also adjacent to the other vertex of $\{u, v\}$ (resp. of $\{x, y\}$).*

Proof. It follows from Lemma 3 (ii) directly. \square

Claim 1.4. $uv \notin E(G)$ and $xy \notin E(G)$. Furthermore, $G[\{u, v, x, y\}]$ is a 4-cycle $uxvyu$.




Fig. 3. The uncrossed 4-cycle $uxvyu$ with the crossed edge uw_1 in D

Proof. By Claim 1.2 and Lemma 3 (i), $G[\{u, v\}] \cong G[\{x, y\}] \cong 2K_1$. By the definition of lexicographic products, $\{ux, xv, vy, yu\} \subseteq E(G)$, as desired. \square

Now, we denote the 4-cycle $uxvyu$ by C , and our focus below is directed towards C . First, we give Claims 1.5 and 1.6, which discuss the crossing properties of C : C will not be uncrossed, and furthermore C does not self-cross.

Claim 1.5. C is crossed in D .

Proof. Suppose C is uncrossed. We choose a neighbor of u , w_1 , such that the edge uw_1 is successive to ux in $\text{Rot}_D(u)$ in the clockwise direction. By Lemma 9, uw_1 is crossed in D , and thus $w_1 \neq y$ since C is uncrossed. By Claim 1.4, $w_1 \neq v$. So $w_1 \in C_{int}$, shown in Fig 3 (a).

We may assume that uw_1 is crossed by w_2w_3 . Notice that $w_2 \neq v$ and $w_3 \neq v$, otherwise, $uv \in E(G)$ by Lemma 10 (i), a contradiction to Claim 1.4. Furthermore, we discuss the following three cases, each of which leads to contradictions.

Case 1. $w_i \cap \{x, y\} = \emptyset$ for $i = 2, 3$. Denote by c the crossing created by uw_1 and w_2w_3 . By Lemma 10 (i), uw_2 and w_3v are uncrossed in D . By the choice of uw_1 (where uw_1 and ux are successive edges) the drawing shown in Fig. 3 (b) will not occur. Thus uw_2 must be drawn shown in Fig 3 (c), and now $(uw_2cu)_{out}$ (or $(uw_3cu)_{out}$) is not a crossed 3-face of D , a contradiction to Lemma 10 (ii).

Case 2. $w_i \neq x$ and $w_j = y$ for $2 \leq i \neq j \leq 3$. Similar to Case 1, this also leads to a contradiction. We omit the proof.

Case 3. $w_2 = x$ or $w_3 = x$. We may assume, without loss of generality, that $w_2 = x$. By Claim 1.4, $w_3 \neq y$, and thus $w_3 \in C_{int}$; see Fig. 3 (d). By Lemma 10 (i), uw_3 and xw_1 are uncrossed edges in D . By Claim 1.3, $\{w_1y, w_3v, w_3y\} \subseteq E(G)$. As C is

uncrossed in D , w_1y must cross w_3v , and thus w_3y is uncrossed in D by Lemma 10 (i). Now the 3-cycle uyw_3u is uncrossed in D , a contradiction to Lemma 11.

Thus the claim holds. \square

Claim 1.6. *Any two edges of C do not cross each other in D .*

Proof. Since D is good, it is only possible for two non-successive edges of C to cross each other. Then by Lemma 10 (i), $G[u, v, x, y] \cong K_4$, and so $uv \in E(G)$, a contradiction to Claim 1.4. \square

By Claims 1.5 and 1.6, we assume that edge ux (on C) is crossed by some edge, namely w_1w_2 . Furthermore, we claim that neither w_1 nor w_2 belongs to C .

Claim 1.7. $w_i \notin V(C)$ for $i = 1, 2$.

Proof. As D is good, $w_i \neq u$ and $w_i \neq x$ for $i = 1, 2$. If some end-vertex of w_1w_2 , says w_1 , is v , then $w_2 \neq y$, otherwise C is self-crossed in D , a contradiction to Claim 1.6. Thus, $w_2 \notin V(C)$, but now $uv \in E(G)$ by Lemma 10 (i), since $v(= w_1)w_2$ crosses ux ; this contradicts to Claim 1.4. By symmetry of w_1 and w_2 , we also have $w_2 \neq v$. Similarly, $w_i \neq y$ for $i = 1, 2$. Thus, the claim holds. \square

By Claims 1.6 and 1.7, we may assume that $w_1 \in C_{out}$ and $w_2 \in C_{int}$, respectively. Furthermore, we shall prove that C has exactly two crossed edges. In addition, in C , vy , which is not adjacent to ux , is the unique crossed edge other than ux . Before that, we observe the existence of the following subdrawing in D , which is an expansion of the current drawing of C .

Claim 1.8. D contains a subdrawing F , as shown in Fig. 4 (i).

Proof. By Lemma 10 (i), w_1u , w_1x , w_2u and w_2x are uncrossed edges in D . By Claim 1.3, $\{w_1y, w_1v, w_2v, w_2y\} \subseteq E(G)$. Since D is good, w_2v cannot cross vx or vy , and w_2v also cannot cross uy , otherwise $uv \in E(G)$ by Lemma 10 (i), which contradicts Claim 1.4. So w_2v must be within C_{int} . Similarly, w_2y is also within C_{int} . Now observe that neither w_1y nor w_1v can be crossed by any edge among uy , vx , and vy . Therefore, w_1y and w_1v are within $(w_1uyvxw_1)_{out}$. Thus, D contains the subdrawing F , as shown in Fig. 4 (i). \square




Fig. 4. (i) The subdrawing F ; (ii) the edge w_2y is crossed by uw_4

Claim 1.9. *C has at least two crossed edges in D.*

Proof. Suppose ux is a unique crossed edge. By Claim 1.8, D contains F as a subdrawing. We claim that w_2y is crossed by some edge, say w_3w_4 ; otherwise, the 3-cycle uw_2yu is uncrossed in D , a contradiction to Lemma 11. Furthermore, we assert that $w_3 = u$ or $w_4 = u$. If not, then w_3 or w_4 , say w_3 , is within $(uw_2yu)_{int}$. By Lemma 10 (i), $w_3y \in E(G)$, and thus $w_3x \in E(G)$. But w_3x crosses at least two times; this contradicts that D is 1-planar. Thus, $w_3 = u$ or $w_4 = u$. Without loss of generality, assume that $w_3 = u$. By Claim 1.3, $w_4 \neq v$, and thus $w_4 \in (w_2yv w_2)_{int}$; see Fig. 4. Furthermore, w_4y is uncrossed in D by Lemma 10 (i), and thus $w_4x \in E(G)$ by Claim 1.3. Now w_4x must cross w_2v , and hence w_4v is uncrossed in D by Lemma 10 (i). Note that vy is also uncrossed in D . So yvw_4y is uncrossed in D , a contradiction to Lemma 11. Therefore, C has at least two crossed edges in D . \square

Claim 1.10. *vx and uy are uncrossed in D.*

Proof. Note that D contains the subdrawing F . As vx and uy are symmetric in F , we need only consider vx . We assume that vx is crossed by some edge, namely w_3w_4 ; see Fig. 5. Observe that if $w_i \in V(F)$ for $i = 3, 4$, then they can be w_1 or w_2 only. So it is sufficient to consider three cases.

Case 1. $w_3 \notin V(F)$ and $w_4 \notin V(F)$. Without loss of generality, we may assume that $w_3 \in (xvw_2x)_{int}$ and $w_4 \in (xvw_2x)_{out}$; see Fig 5 (a). Since w_3w_4 crosses vx , by Lemma 10 (i) $w_3v \in E(G)$, and thus $w_3u \in E(G)$ by Claim 1.3. But now u_3u will be crossed at least twice, a contradiction.

Case 2. $w_3 \notin V(F)$ and $w_4 = w_1$. See Fig 5 (b). Similar to Case 1, w_3u will be crossed at least twice, a contradiction.

Case 3. $w_3 = w_2$. In this case, $w_4 \notin V(F)$ since G is simple. Since D is 1-planar, $w_4 \in (xvw_1x)_{int}$, as shown in Fig 5 (c). Since $w_2 (= w_3)w_4$ crosses vx , $w_4v \in E(G)$ by Lemma 10 (i), and thus by Claim 1.3, $w_4u \in E(G)$. But now u_4u is crossed (by other edges) at least twice, a contradiction.

Thus, xv is uncrossed in D . Similarly, uy is also uncrossed in D . So the claim follows. \square

By Claims 1.9 and 1.10, the following claim follows directly.

Claim 1.11. *vy is crossed by some edge, namely w_3w_4 .*

Furthermore, we shall prove that $w_3 \notin V(F)$ and $w_4 \notin V(F)$.

Claim 1.12. *$w_3 \notin V(F)$ and $w_4 \notin V(F)$.*




Fig. 5. Three 1-planar drawings involving the crossed edge xv




Fig. 6. The 1-planar drawing F'

Proof. Since w_3w_4 has been crossed by vy , if $w_3 \in V(F)$ or $w_4 \in V(F)$, then they can be w_1 or w_2 only. Without loss of generality, assume that $w_3 = w_1$, and thus $w_4 \neq w_2$ by simplify of G . Thus, $w_4 \in (w_2vyw_2)_{int}$. By Lemma 10 (i), w_1v is uncrossed in D . Note that w_1x and uy are uncrossed in D . So the 3-cycle vw_1yv is uncrossed in D , which contradicts Lemma 11. If $w_3 = w_2$, then similarly, there exists an uncrossed 3-cycle w_2uyw_2 in D , a contradiction. By symmetry, $w_4 \neq w_i$ for $i = 1, 2$. Therefore, $w_3 \notin V(F)$ and $w_4 \notin V(F)$. \square

By Claim 1.12, we can assume that w_3 and w_4 are within $(w_2vyw_2)_{out}$ and $(w_2yv w_2)_{int}$, respectively. Furthermore, since D is 1-planar and w_3w_4 crosses vy , we have $w_3 \in (w_1yv w_1)_{out}$.

Claim 1.13. D contains a 1-planar drawing F' , shown in Fig. 6.

Proof. Similar to Claim 1.8, we obtain this claim. \square

Since w_1v crosses w_3x in F' , by Lemma 10 (i), w_1w_3 is uncrossed in D . Similarly, w_2w_4 is also uncrossed in D . So we obtain graph F'' , which obtained from F' by adding two uncrossed edges w_1w_3 and w_2w_4 . Now $F'' \cong K_{2,2,2,2}$, and F'' is optimal 1-planar. So by Lemma 12, $G \cong F''$. Combining this with Lemma 14, the proposition follows. \square

4 Left and right factors with order at least three

In this section, we shall prove that an optimal 1-planar graph cannot be reduced into factors where both the left and right factors have order greater than three; see Proposition 2. This proof approach of Proposition 2 is somewhat similar to that of Proposition 1(ii), but due to the order (which is three) of the right factor. Our focus is on considering a specific 1-planar drawing of subgraph $K_{3,3}$. In fact, the proof is simpler than in Proposition 1(ii), as there is more direct local information available to utilize compared to the case of a cycle of length four.

Proposition 2. *Let G be an optimal 1-planar graph. If $|V(G_1)| \geq 3$ and $|V(G_2)| \geq 3$, then $G \neq G_1 \circ G_2$.*

Proof. Suppose $G = G_1 \circ G_2$. If $|V(G_2)| \geq 4$, then by Lemma 4, $G_1 \circ G_2$ is not 1-planar, a contradiction. Hence, we only consider $|V(G_2)| = 3$. Let $V(G_1) = \{u_1, u_2, \dots, u_k\}$ where $k \geq 3$ and $V(G_2) = \{v_1, v_2, v_3\}$.

Claim 2.1. *G contains a copy of $K_{3,3}$.*

Proof. By Lemma 2, G_1 is connected. Hence, there is an edge $u_i u_j$ for $1 \leq i \neq j \leq k$ in G_1 . Since $|V(G_2)| \geq 3$, then G contains a copy of $K_{3,3}$. \square

We denote one copy of $K_{3,3}$ by T , where two parts of T are $X := \{(u_i, v_1), (u_i, v_2), (u_i, v_3)\}$ and $Y := \{(u_j, v_1), (u_j, v_2), (u_j, v_3)\}$. For brevity, we relabel $X := \{u, v, w\}$ and $Y := \{x, y, z\}$. Then two observations below follow from Lemma 3 directly.

Observation 1. *If a vertex p of G is adjacent to one vertex of X (resp. Y), then p is adjacent to all vertices of X (resp. Y).*

Observation 2. $G[\{x, y, z\}] \cong G[\{u, v, w\}]$.

Claim 2.2. $D|T \cong A1$.

Proof. By Lemma 13, $D|T$ is isomorphic to $A1$ or $A2$. If $D|T \cong A2$, by Lemma 10 (i), D has the 3-cycle $xyzx$, and it is uncrossed in D , a contradiction to Lemma 11. So $D|T \cong A1$. \square

Claim 2.3. *D contains $D|H$, shown in Fig 8, as a subdrawing.*

Proof. By Claim 2.2, $T \cong A1$. Since xv crosses uy in $D|H$, we have $\{uv, xy\} \subseteq E(G)$. Then uv and xy are uncrossed in D by Lemma 10 (i). Similarly, ux and vy are uncrossed. Let H be the subgraph of G obtained from T by adding xy and uv . So D contains the subdrawing $D|H$. \square

Since uv is uncrossed in D , by Lemma 10 (i), uz or vz is crossed in D . By symmetry, we may assume that uz is crossed by some edge st . By symmetry of s and t , we discuss five cases, each of which leads to a contradiction regarding 1-planarity or an uncrossed 3-cycle.




Fig. 7. Two possible restricted 1-planar drawings of $D|T$

Case 1. $s \notin V(H)$ and $t \notin V(H)$. We may assume that $s \in (uvzu)_{int}$ and $t \in (uvzu)_{out}$, shown in Fig. 9 (1). As st crosses uz , by Lemma 10 (i) su is uncrossed in D . By Observation 1, $\{sw, sv\} \subseteq G$. Since uv is uncrossed in D and uz has been crossed by st , sw crosses vz , and hence sv is uncrossed in D by Lemma 10. Thus, the 3-cycle $uvsu$ is uncrossed in D , a contradiction to Lemma 11.

Case 2. $s \notin V(H)$ and $t = w$. In this case, $s \in (uvzu)_{int}$; see Fig. 9 (2). Then $sz \in E(G)$ by Lemma 10 (i), since sw is crossed by uz . By Observation 1, $sx \in E(G)$. But sx is crossed twice or will make at least one edge on $D|H$ to be crossed at least twice, a contradiction.

Case 3. $s \notin V(H)$ and $t = x$. In this case, s lies in $(uvzu)_{int}$; see Fig. 9 (3). Since sx crosses uz , it follows that $sz, su \in E(G)$ by Lemma 10 (i). By Observation 1, $\{sy, sw\} \subseteq E(G)$. As uz is crossed and uv is uncrossed in D , sy or sw , say sy , must cross vz , and hence sw will cross twice or cause at least one edge on $D|H$ to be crossed at least twice, a contradiction.

Case 4. $s = v$ and $t \notin V(H)$. In this case, uz is crossed by tv , and $t \in (uxwzu)_{out}$. By Lemma 10 (i), $tz \in E(G)$, $tu \in E(G)$, and vz and tu are uncrossed in D . Furthermore, by Observation 1, $ty \in E(G)$. Then sy crosses wz or wx . If ty crosses wz , then yz is uncrossed in D by Lemma 10 (i), and hence the 3-cycle $vyzv$ is uncrossed in D ; see Fig. 9 (4), a contradiction to Lemma 11. If ty crosses wx , then $tx \in E(D)$ and is uncrossed by Lemma 10 (i), as shown in Fig. 9 (5). But now the 3-cycle $tuxt$ is uncrossed in D , a contradiction to Lemma 11.

Case 5. $s = v$ and $t = w$. In this case, since vw crosses uz , uw is an edge of G and vz is uncrossed in D by Lemma 10 (i); see Fig. 9 (6). Now observe that $G[\{u, v, w\}] \cong C_3$, and thus $G[\{x, y, z\}] \cong C_3$ by Observation 2. Hence, $yz \in E(G)$ and $xy \in E(G)$. Then xz must cross wy , and thus yz is uncrossed in D by Lemma 10. Recall that vy is also uncrossed in D . Thus, the 3-cycle $yvzy$ is uncrossed in D , a contradiction to Lemma 11.

Thus, we conclude that $G \neq G_1 \circ G_2$, and with this, the proposition is established. \square




Fig. 8. The subdrawing $D|H$




Fig. 9. The possible six subdrawings involving a crossed edge uz

5 Proof of Theorems 1, 2 and 3

Proof of Theorem 1. The “only if” part of the theorem follows directly from Lemma 14. By combining Proposition 1 with Proposition 2, the “if” part of the theorem is established. \square

Proof of Theorem 2. Since $n \geq 5$, by Theorem 1, $G \circ 2K_1$ is not optimal 1-planar. Thus, $|E(G \circ 2K_1)| \leq 4 \times 2n - 9$. By Lemma 1, $|E(G \circ 2K_1)| = 4m$. So $4m \leq 8n - 9$, and thus $m \leq \lfloor \frac{8n-9}{4} \rfloor = 2n - 3$, as desired. \square

Proof of Theorem 3. If $n \geq 10$, then $m \leq 4n - 8$ by Lemma 7. Theorem 1 implies that any optimal 1-planar graph is irreducible if it has at least 10 vertices, and thus $m \leq 4n - 9$ for $n \geq 10$ (since G is reducible). For if $n = 9$, then any 1-planar




Fig. 10. A 1-planar drawing of $H = P_{\frac{n}{3}} \circ C_3$

graph with 9 vertices has at most $4 \times 9 - 9$ edges, and thus $m \leq 4 \times 9 - 9$ by Lemma 7. Let $P_{\frac{n}{3}}$ be a path with $\frac{n}{3}$ vertices where $n \geq 9$ and $n \equiv 0 \pmod{3}$. Let $V(P_{\frac{n}{3}}) = \{u_1, u_2, \dots, u_{\frac{n}{3}}\}$. Let $H = P_{\frac{n}{3}} \circ C_3$. Clearly, H is reducible, and $|E(H)| = 4n - 9$. Furthermore, a 1-planar drawing of H is shown in Fig. 10, where C_{u_i} is a 3-cycle in H that corresponds to u_i in $P_{\frac{n}{3}}$ for $1 \leq i \leq \frac{n}{3}$. Thus, $4n - 9$ is a tight bound for every $k \geq 3$ and $n = 3k$. \square

Remark 1. We believe that the bound in Theorem 2 we provided is still far from resolving Problem 1.

6 Acknowledgment

We claim that there is no conflict of interest in our paper. No data was used for the research described in the article. The work was supported by the National Natural Science Foundation of China (Grant No. 12271157, 12371346) and the Postdoctoral Science Foundation of China (Grant No. 2024M760867).

References

- [1] P. Angelini, M. A. Bekos, M. Kaufmann, T. Schneck, Efficient generation of different topological representations of graphs beyond-planarity, *J. Graph Algorithms Appl.* **24** (2020), 573–601
- [2] C. Binucci, W. Didimo, F. Montecchiani, 1-planarity testing and embedding: An experimental study, *Comput. Geom.* **108** (2023) 101900.
- [3] R. Bodendiek, H. Schumacher, K. Wagner, Bemerkungen zu einem Sechsfarbenproblem von G. Ringel, *Abh. Math. Semin. Univ. Hamb.* **53** (1983), 41–52.
- [4] J. Bucko, J. Czap, 1-planar lexicographic products of graphs, *Appl. Math. Sci.* **9** (2015), 5441–5449.
- [5] J. Czap, D. Hudák, 1-planarity of complete multipartite graphs, *Discrete Appl. Math.* **160** (2012), 505–512.
- [6] J. Czap, D. Hudák, On drawings and decompositions of 1-planar graphs, *Electron. J. Comb.* **20** (2013), P54.

- [7] J. Czap, D. Hudák, T. Madaras, Joins of 1-planar graphs, *Acta Math. Sin., Engl. Ser.* **30** (2014), 1867–1876.
- [8] J. Feigenbaum, A. A. Schäffer, Recognizing composite graphs is equivalent to testing graph isomorphism, *SIAM J. Comput.* **15** (1986), 619–627.
- [9] R.H. Hammack, W. Imrich, S. Klavžar, W. Imrich, S. Klavžar, *Handbook of Product Graphs*, Vol. 2, CRC press Boca Raton, 2011.
- [10] Y. Huang, Z. Ouyang, F. Dong, On the size of matchings in 1-planar graph with high minimum degree, *SIAM J. Discrete Math.* **36** (2022), 2570–2584.
- [11] S.G. Kobourov, G. Liotta, F. Montecchiani, An annotated bibliography on 1-planarity, *Comput. Sci. Rev.* **25** (2017), 49–67.
- [12] V.P. Korzhik, B. Mohar, Minimal obstructions for 1-immersions and hardness of 1-planarity testing, *J. Graph Theory* **72** (2013), 30–71.
- [13] N. Matsumoto, Y. Suzuki, Non-1-planarity of lexicographic products of graphs, *Discuss. Math. Graph Theory* **41** (2021), 1103–1114.
- [14] G. Ringel, Ein Sechsfarbenproblem auf der Kugel, *Abh. Math. Semin. Univ. Hambg.* **29** (1965), 107–117.
- [15] Y. Suzuki, Re-embeddings of maximum 1-planar graphs, *SIAM J. Discrete Math.* **24** (2010), 1527–1540.
- [16] Y. Suzuki, K_7 -minors in optimal 1-planar graphs, *Discrete Math.* **340** (2017), 1227–1234.
- [17] D.B. West, *Introduction to Graph Theory* (second edition), Prentice Hall, 2001.
- [18] X. Zhang, The edge chromatic number of outer-1-planar graphs, *Discrete Math.* **339** (2016), 1393–1399.