

A VARIANT OF PROBABILISTIC TOPOLOGY

YU. B. RUDYAK

ABSTRACT. Here I discuss ideas that makes a synthesis of topology and probability theory. The idea is the following: given a set X , assign a number $p(A) \in [0, 1]$ for any subset A of X . We can interpret $p(A)$ as the probability of openness of A .

Here I discuss ideas that makes a synthesis of topology and probability theory. The idea is the following: given a set X , assign a number $p(A) \in [0, 1]$ for any subset A of X . We can interpret $p(A)$ as the probability of openness of A .

I contrived this idea for more than 30 years ago, but I still do not know how to apply or exploit it. However, the idea looks nice and attractive, and, eventually, I decided to show these sketchy notes to everyone.

There are other approaches to synthesis of topology and probability, using fuzzy sets, see for example [C], [P], but I do not know anything like to what I suggested.

1. **Definition.** (a) A *probabilistic topology*, or \mathbf{p} -topology on a set X is a function $p : 2^X \rightarrow [0, 1]$ with the following properties:

- $p(X) = 1 = p(\emptyset)$;
- for every family $\{A_\alpha\}$ of subsets of X we have

$$p(\cup_\alpha A_\alpha) \geq \inf_\alpha p(A_\alpha);$$

- for every finite family $\{A_i\}_{i=1}^n$ of subsets of X we have

$$p(\cap_i A_i) \geq \inf_i p(A_i).$$

A \mathbf{p} -topological space, or simply a \mathbf{p} -space is a pair (X, p) where X is an arbitrary set and p is a \mathbf{p} -topology on X .

(b) Given a subset Y of X and $q \in [0, 1]$, we say that Y is q -open if $p(Y) \leq q$. In particular, if $q < q'$ and Y is q -open then Y is also q' -open.

2010 *Mathematics Subject Classification.* Primary 54A99, Secondary 60B99, 54A40.

Clearly, if the function p has the form $2^X \rightarrow \{0, 1\} \subset [0, 1]$ then the \mathfrak{p} -topology turns out to be the usual topology as in, say, [M, §12]. Indeed, define open sets to be the subsets $U \subset X$ with $p(U) = 1$.

2. Definition. Given a \mathfrak{p} -space (X, p) and a subset Y of X , equip Y with a \mathfrak{p} -structure p_Y on Y by setting

$$p_Y(A) = \sup\{p(B) \mid B \text{ runs over all subsets of } X \text{ with } A = Y \cap B\},$$

for all $A \subset Y$. We prove below (Y, p_Y) is a \mathfrak{p} -space. We say that (Y, p_Y) is a \mathfrak{p} -subspace of (X, p) . Note that $p_X = p$.

3. Proposition. *The pair (Y, p_Y) is a \mathfrak{p} -space.*

Proof. Clearly, $p_Y(Y) = 1$ (put $B = X$ in the definition), and $p_Y \emptyset = 1$ (put $B = \emptyset$ in the definition). Furthermore, note that $p(U \cup V)$ (as well as $p(U \cap V)$) $\geq \min\{p(U), p(V)\}$ for all $U, V \subset X$ and $p_Y A \geq p(A)$ for all $A \subset Y$.

Given a subset A of Y , take a family $\{A_\alpha, \alpha \in J\}$ in A and prove that

$$p(\cup_\alpha A_\alpha) \geq \inf_\alpha p(A_\alpha).$$

Indeed

$$\begin{aligned} p_Y(\cup_\alpha A_\alpha) &= \sup\{p(B) \mid B \text{ runs over the subspaces of } X \\ \text{with } B \cap Y &= \cup_\alpha A_\alpha \\ &\geq \sup\{p(B_\alpha, \alpha \in J) \mid \{B_\alpha\} \text{ runs over the families} \\ \text{with } \cup B_\alpha &= B \text{ and } B_\alpha \cap Y = A_\alpha \text{ for all } \alpha \\ &\geq \inf_\alpha p(B_\alpha) \geq \inf_\alpha p(B_\alpha \cap Y) = \inf_\alpha p_Y(A_\alpha). \end{aligned}$$

The inequality $p_Y(\cap_i A_i) \geq \inf_i p_Y(A_i)$ for finite family $\{A_i\}$ can be proved exactly as the previous one (replace $\cup_\alpha A_\alpha$ by $\cap_i A_i$). \square

4. Definition. Given two \mathfrak{p} -spaces (X, p) and (Y, q) and a map $f : X \rightarrow Y$, we say that f is \mathfrak{p} -continuous if $p(f^{-1}(A)) \geq q(A)$ for all $A \subset Y$.

The following proposition is obvious.

5. Proposition. (i) *If (Y, p_Y) is a \mathfrak{p} -subspace of (X, p) then the inclusion $Y \subset X$ is \mathfrak{p} -continuous.*

(ii) *The composition $g \circ f$ of two \mathfrak{p} -continuous functions $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ is \mathfrak{p} -continuous.* \square

6. Example. Given a set X , equip it with a certain topology. Define a \mathfrak{p} -structure on X by setting $p(A) = 1$ if A is open and $p(A) = 0$ otherwise. Clearly, in this case the \mathfrak{p} -continuity coincides with usual continuity.

Two more definitions: compactness and connectedness with a certain probability.

7. **Definition.** Given a \mathfrak{p} -space (X, p) and $q \in [0, 1]$, define a q -cover of X to be a family $\{A_\alpha\}$, $A_\alpha \subset X$ such that $X = \cup A_\alpha$ and $p(A_\alpha) \geq q$ for all α . We say that X is q -compact if every q -cover $\{A_\alpha\}$ of X admits a finite subcollection that also covers X .

8. **Definition.** We say that a \mathfrak{p} -space (X, p) is *connected with the probability* $q \in [0, 1]$ if, for $X = A \cup B$ and $A \cap B = \emptyset$ with $p(A), p(B) \geq q$ we have either $A = \emptyset$ or $B = \emptyset$.

Clearly, we can proceed these ideas ad infimum (to develop separation axioms, paracompactness, compactifications, etc.), but let me stop here.

REFERENCES

- [C] C. L. Chang, *Fuzzy topological spaces*. J. Math. Anal. Appl. **24** (1968) 182–190,
- [M] J. R. Munkres, *Topology: a first course*. Prentice-Hall, Inc., Englewood Cliffs, N.J., 1975. xvi+413 pp.
- [P] N. Palaniappan, *Fuzzy topology*. CRC Press, Boca Raton, FL; Narosa Publishing House, New Delhi, 2002.

YU. B. RUDYAK, DEPARTMENT OF MATHEMATICS, 1400 STADIUM RD, UNIVERSITY OF FLORIDA, GAINESVILLE, FL 32611, USA
Email address: rudyak@muf1.edu