A representation of $Out(F_n)$ by counting subwords of cyclic words

Noam M.D. Kolodner

December 2, 2022

Abstract

We generalize the combinatorial approaches of Rapaport and Higgins-Lyndon to the Whitehead algorithm. We show that for every automorphism φ of a free group F and every word $u \in F$ there exists a finite multiset of words $S_{u,\varphi}$ satisfying the following property: For every cyclic word w, the number of times u appears as a subword of $\varphi(w)$ depends only on the appearances of words in $S_{u,\varphi}$ as subwords of w. We use this fact to construct a faithful representation of Out (F_n) on an inverse limit of \mathbb{Z} -modules, so that each automorphism is represented by sequence of finite rectangular matrices, which can be seen as successively better approximations of the automorphism.

1 Introduction

In [8] Whitehead introduced an algorithm for determining whether a word in the free group is primitive. Whiteheads proof of the algorithm used 3-manifolds. Rapaport [5] Higgins and Lyndon [2] came up with a combinatorial approach for proving the algorithm this is summarized in Lyndon-Schupp [4]. In this paper we present generalization of this combinatorial approach. We will treat words as folded labeled graphs homeomorphic to a compact line segment and cyclic words as folded labeled graphs homeomorphic to a circle. One can interpret the method presented in Lyndon and Schupp in the following way. Let w be a cyclic word and let τ be a Whitehead generator. One can count how many times a specific letter appears in τ (w) by counting how many times appropriate 2-letter words appear in w. Furthermore, we define M_k with k = 1, 2 to be a \mathbb{Z} -module generated freely by words of length k. For every cyclic word w we can assign an integer non negative vector π_k (w) in M_k which counts how many times each subword of length k appears in w. In this interpretation a whitehead generator τ gives rise to a linear function $\tau_1 : M_2 \to M_1$ such that $\tau_1 \pi_2$ (w) = π_1 (τ (w)).

In Section 1 we show that for every automorphism φ and any word u one can count the number of time u appears in $\varphi(w)$ by counting appropriate subwords in w.

Theorem 1. Let φ an automorphism and u a word we show that there is a unique minimal finite multiset of words $S_{u,\varphi}$ such that for every cyclic word w

$$|Hom(u,\varphi(w))| = \sum_{v \in S_{u,\varphi}} |Hom(v,w)|$$
(1.1)

Because we consider words and cyclic words as labeled graphs, by counting the number of graph morphisms between u and w we count the number of time u appears as a subword in w. Given u and φ we will also show how to algorithmically calculate $S_{u,\varphi}$. In Section 2 we use Equation 1.1 to construct a representation of $Out(F_n)$. Let $\mathbb{Z}[C]$ be a \mathbb{Z} -module generated freely by all cyclic words. We extend π_k linearly to an homomorphism $\pi_k : \mathbb{Z}[C] \to M_k$ while extending the definition of M_k for $2 \leq k$.

Theorem 2. There are homomorphisms $p_k : M_k \to M_{k-1}$ that satisfy

$$p_k \circ \pi_k = \pi_{k-1} \tag{1.2}$$

Moreover we construct p_k and prove properties of this construction in Theorem 42. Let φ be an automorphism let $m_k^{\varphi} = \max_{u \in W_k} \max_{v \in S_{u,\varphi}} l(v)$ with W_k denoting the set of words of length k and l(v) denoting the length of v. We will show how to use Equation 1.1 to define homomorphisms $\varphi_k : M_{m_k} \to M_k$ that satisfy

$$\varphi_k \circ \pi_{m_k} = \pi_k \circ \varphi \tag{1.3}$$

Denote $\operatorname{Im} \pi_k = \tilde{M}_k$ and consider $\varprojlim \tilde{M}_k$ the inverse limit of $p_k : \tilde{M}_k \to \tilde{M}_{k-1}$. Equations 1.2 and 1.3 show that $\varphi \mapsto \varprojlim \varphi_k$ is a representation $Out(F_n) \to \operatorname{Aut}(\varprojlim \tilde{M}_k)$.

Theorem 3. The representation $Out(F_n) \to Aut(\varprojlim \tilde{M}_k)$ is a faithful representation

As a corollary the multisets $S_{u,\varphi}$ characterize φ i.e. let φ, ψ automorphisms in different conjugacy classes then there exists a word u such that $S_{u,\varphi} \neq S_{u,\psi}$. Although we have not yet found application for this representation we believe it holds valuable information on outer automorphisms.

2 Preliminaries

Definition 4 (Graphs). We use graphs in the sense of Serre [6]: A graph Γ is a set $V(\Gamma)$ of vertices and a set $E(\Gamma)$ of edges with a function $\iota: E(\Gamma) \to V(\Gamma)$ called the initial vertex map and an involution $\overline{\Box}: E(\Gamma) \to E(\Gamma)$ with $\overline{e} \neq e$ and $\overline{\overline{e}} = e$. A graph morphism $f: \Gamma \to \Delta$ is a pair of set functions $f^E: E(\Gamma) \to E(\Delta)$ and $f^V: V(\Gamma) \to V(\Delta)$ that commute with the structure functions. A path in Γ is a finite sequence $e_1, \ldots, e_n \in E(\Gamma)$ with $\iota(\overline{e}_k) = \iota(e_{k+1})$ for every $1 \leq k < n$. The path is closed if $\iota(\overline{e}_n) = \iota(e_1)$, and reduced if $e_{k+1} \neq \overline{e}_k$ for all k. All the graphs in the paper are assumed to be connected unless specified otherwise, namely, for every $e, f \in E(\Gamma)$ there is a path e, e_1, \ldots, e_n, f .

Definition 5 (Labeled graphs). Let X be a set and let X^{-1} be the set of its formal inverses. We define R_X to be the graph with $E(R_X) = X \cup X^{-1}$, $V(R_X) = \{*\}, \overline{x} = x^{-1}$ and $\iota(x) = *$. An X-labeled graph is a graph Γ together with a graph morphism $\Lambda \colon \Gamma \to R_X$ called the label function. A morphism of X-labeled graphs Γ and Δ is a graph morphism $f \colon \Gamma \to \Delta$ that commutes with the label functions.

Definition 6 (Folding). A labeled graph Γ is folded if $\Lambda(e) \neq \Lambda(f)$ holds for every two edges $e, f \in E(\Gamma)$ with $\iota(f) = \iota(e)$. Or equivalently if Λ is a graph immersion. If Γ is not folded, there exist $e, f \in E(\Gamma)$ s.t. $\iota(e) = \iota(f)$ and $\Lambda(e) = \Lambda(f)$; Let Γ' be the graph obtained by identifying the vertex $\iota(\overline{e})$ with $\iota(\overline{f})$, the edges e with f and \overline{e} with \overline{f} . We say Γ' is the result of folding e and f. The label function Λ factors through Γ' , yielding a label function Λ' on Γ' . The labeled graphs morphism $g: \Gamma \to \Gamma'$ is called a folding. Let Γ be a finite labeled graph. There is a finite sequence of foldings $g_i: \Gamma_{i-1} \to \Gamma_i$ with $\Gamma_0 = \Gamma$, $0 \leq i \leq n$ such that Γ_n is folded. The folded graph Γ_n is unique and it does not depend on the order in which we choose the foldings. See [7].

Remark 7. For reduced words u, v in a free group, we write $u \cdot v$ to indicate that there is no cancellation in their concatenation, namely the first letter in v is not the inverse of the last letter in u.

Definition 8 (Proper subword). Let F_n be a free and let $u_1, u_2, v, w \in F_n$ s.t. $u_1 \cdot v \cdot u_2 = w$. If the multiplication $u_1 \cdot v$ and $v \cdot u_2$ are preformed in F_n without cancellation u_1, v and u_2 are called proper subwords of w. We call u_1 a prefix and u_2 a suffix.

Definition 9 (Unoriented-Word). A unoriented word is a folded labeled graph that is topologically homeomorphic to a connected compact set in \mathbb{R} . Let $u \in F_n$ be a word we denote the unoriented word by $\overline{u} = \overline{u^{-1}}$. And call u and u^{-1} orientations of \overline{u} . When context allows we drop the bar. We abbreviate u-word.

Definition 10 (Cyclic Word). A cyclic word is a folded labeled graph that is topologically homeomorphic to a circle. We denote a cyclic word by a cyclically reduced representative without a bar. This may differ from convention as often cyclic words are defined to be the conjugation class of a word in F_n . Let $w \in F_n$ under our definition w and w^{-1} denote the same cyclic word.

Definition 11 (Subword). Let w be a u-word. A u-word u is called subword of w if there is a graph morphism $u \to w$. Let w be a cyclic word. An u-word \overline{u} will be called a subword of w if there is a labeled graph morphism from u to w (it need not be injective!). Let $x \in F_{\{x,y\}}$ be a one letter cyclic word and let $\overline{x^n}$ be an u-word under this definition $\overline{x^n}$ is a subword of x (as a cyclic word of course). Again this may differ from convention.

Definition 12 (k-Affix). Let $k \in \mathbb{N}$ and let \overline{w} be an u-word with l(w) > k. A k-affix is an u-word \overline{u} of length k s.t. u is either a prefix or suffix of w or w^{-1} . We notice that if u is prefix of w than u^{-1} is a suffix of w^{-1} thus each u-word \overline{w} has exactly two k-affixes. This can be seen as a graph function between u-words that is a local homeomorphism except for exactly one point.

Definition 13. Let $\varphi: F_Y \to F_X$ be a homomorphism such that $\forall y \in Y, \varphi(y) \neq 1$. Let Δ be a Y-labeled graph we construct an X-labeled graph $\mathcal{F}_{\varphi}(\Delta)$ by replacing every edge $e \in E(\Delta)$ by a reduced path labeled $\varphi(\Lambda(e))$. Let Δ and Ξ be Y-labeled graphs, and $f: \Delta \to \Xi$ a graph morphism. The graph morphism $\mathcal{F}_{\varphi}(f): \mathcal{F}_{\varphi}(\Delta) \to \mathcal{F}_{\varphi}(\Xi)$ is defined by gluing the identity maps $\bigsqcup_{e \in E(\Delta)} \varphi(\Lambda(e)) \to \varphi(\Lambda(f(e)))$. We notice that generally $\mathcal{F}_{\varphi}(\Delta)$ is not a folded graph. We denote by $\tilde{\mathcal{F}}_{\varphi}(\Delta)$ the folded graph obtained by folding $\mathcal{F}_{\varphi}(\Delta)$. It is a functor as well. We notice that the folding morphism denoted by $\rho_{\Delta}: \mathcal{F}_{\varphi}(\Delta) \to \tilde{\mathcal{F}}_{\varphi}(\Delta)$ is a natural transformation. This definition is taken from [3].

The motivation for \mathcal{F} is topological: thinking of Δ, R_Y, R_X as topological spaces and of $\Lambda: \Delta \to R_Y, \varphi: R_Y \to R_X$ as continuous functions, we would like to think of Δ as an X-labeled graph with the label function $\varphi \circ \Lambda$. The problem is that $\varphi \circ \Lambda$ does not send edges to edges, and we mend this by splitting edges in Δ to paths representing their images in R_X .

Example 14. Let $\varphi : F_{\{x,y\}} \to F_{\{x,y\}}$ be the automorphism $x \mapsto xy, y \mapsto y$. In Figure 2.1 we see the graph morphism between the u-word \overline{x} and the cyclic word xy and the morphism after applying \mathcal{F}_{φ} and $\tilde{\mathcal{F}}_{\varphi}$.



Figure 2.1

Definition 15. Let W_k be the set of length k u-words. We define $M_k = \operatorname{span}_{\mathbb{Z}} W_k$ the free module generated by W_k . We define M_k^* to be the dual to M_k . If $u \in W_k$ let $u^* \in M_k^*$ be the functional s.t. $u^*(u) = 1$ and $u^*(v) = 0$ for every $v \in W_k$ such that $v \neq u$. Let C the set of cyclic words of F_n and $\mathbb{Z}[C]$ the free model generated by cyclic words. We define a sequence of homomorphisms $\pi_k : \mathbb{Z}[C] \to M_k$. Let $w \in C$ and $u \in W_k$ we define $u^*(\pi_k(w)) = |\operatorname{Hom}(u,w)|$

Example 16. Take $F_{\{x,y\}}$ and the cyclic word $xyx^{-1}y^{-1}$ then $\pi_2(xyx^{-1}y^{-1}) = \overline{xy} + \overline{yx^{-1}} + \overline{x^{-1}y^{-1}} + \overline{y^{-1}x}$ and $\pi_3(xyx^{-1}y^{-1}) = \overline{xyx^{-1}} + \overline{yx^{-1}y^{-1}} + \overline{x^{-1}y^{-1}x} + \overline{yx^{-1}y^{-1}} + \overline{x^{-1}y^{-1}x^{-1}} + \overline{yx^{-1}y^{-1}} + \overline{yx^{-1}$

3 Counting subwords

We start by demonstrating our interpretation of Lyndon-Schupp for the simplest kind of Whitehead generator.

Example 17. Let $\varphi : F_{\{x,y,z\}} \to F_{\{x,y,z\}}$ be the automorphism $x \mapsto xy, y \mapsto y, z \mapsto z$. We construct $\varphi_1 : M_2 \to M_1$ s.t. $\varphi_1 \pi_2 (w) = \pi_1 (\varphi (w))$

Let w be a cyclic word. The letters \overline{x} and \overline{z} (u-words of length 1) will appear the same amount of times in $\varphi(w)$ as they appear in w. Each time x appears in w it is followed by x, y, y^{-1}, z or z^{-1} therefor by counting the number of times the subwords $\overline{xx}, \overline{xy}, \overline{xy}^{-1}, \overline{xz}, \overline{xz^{-1}}$ appear we count the number of times \overline{x} appears. Now we count the number of appearances of \overline{y} in $\varphi(w)$. Every y in w that is not followed by x^{-1} gives a y in $\varphi(w)$. Every x in w gives a y in $\varphi(w)$ as long as it is not followed by y^{-1} . So if we count the number of the subwords $\overline{yx}, \overline{yy}, \overline{yz}, \overline{yz^{-1}}$ and $\overline{xx}, \overline{xy}, \overline{xz}, \overline{xz^{-1}}$ in w this will gives us the number of appearances of y in $\varphi(w)$. We use the identification of Hom (M_2, M_1) with $M_2^* \otimes M_1$ to define φ_1 as follows

$$\varphi_{1} = \left(\overline{xx^{*}} + \overline{xy^{*}} + \overline{xy^{-1}}^{*} + \overline{xz^{*}} + \overline{xz^{-1}}^{*}\right) \otimes \overline{x}$$

$$+ \left(\overline{yx^{*}} + \overline{yy^{*}} + \overline{yz^{*}} + \overline{yz^{-1}}^{*} + \overline{xx^{*}} + \overline{xy^{*}} + \overline{xz^{*}} + \overline{xz^{-1}}^{*}\right) \otimes \overline{y}$$

$$+ \left(\overline{zx^{*}} + \overline{zx^{-1}}^{*} + \overline{zy^{*}} + \overline{zy^{-1}}^{*} + \overline{zz^{*}}\right) \otimes \overline{z}$$

We want to generalize this. We will generalize in a naive intuitive approach then show why this approach does not work and give the right but somewhat complicated and non-intuitive definitions.

Definition (naive ideal-preimage). Let φ be an automorphism and \overline{u} be an uword. An u-word \overline{v} is a φ -ideal preimage of \overline{u} if for every u-word \overline{w} with a graph morphism $\overline{v} \to \overline{w}$ there exists a graph morphism $\overline{u} \to \overline{\varphi}(w)$. Equivalently, let vbe an orientation of \overline{v} . For every words w_1, w_2 with $w_1 \cdot v \cdot w_2$ the u-word \overline{u} is a subword of $\varphi(\overline{w_1 v w_2})$.

We notice that if $\overline{v_1}$ is a φ -ideal preimage of \overline{u} and $\overline{v_2}$ is a u-word such that there is a graph morphism $\overline{v_1} \to \overline{v_2}$ then $\overline{v_2}$ is a φ -ideal preimage of \overline{u} as well.

Definition (naive Minimal ideal-preimage). A φ -ideal preimage \overline{v} of \overline{u} is said to be minimal if there is no $\overline{v'} \to \overline{v}$ such that $\overline{v'}$ is also a φ -ideal preimage of \overline{u}

In Example 17 the minimal φ -ideal preimages of \overline{x} and \overline{z} are \overline{x} and \overline{z} respectively and the minimal φ -ideal preimages of \overline{y} are $\overline{yx}, \overline{yy}, \overline{yz}, \overline{yz^{-1}}, \overline{xx}, \overline{xy}, \overline{xz}, \overline{xz^{-1}}$. Thus imitating Example 17 we count the number of subwords \overline{u} in a cyclic word $\varphi(w)$ by counting the number of minimal φ -ideal preimages in w. The next two example show why a more sophisticated approach is needed

Example 18. Let $\varphi : F_{\{x,y,z\}} \to F_{\{x,y,z\}}$ be the automorphism $x \mapsto xyy, y \mapsto y, z \mapsto z$. We notice that as before in Example 17 the u-word \overline{xz} is a φ -ideal preimage of \overline{y} but we observe that $\varphi(\overline{xz}) = \overline{xyyz}$ and that no matter what words w_1, w_2 with $w_1 \cdot xz \cdot w_2$ then $\varphi(\overline{w_1xzw_2})$ will include 2-copies of \overline{y} . Thus the approach of counting the number of \overline{y} in $\varphi(w)$ by counting minimal ideal preimages will not work here.

Example 19. Let $\varphi : F_{\{x,y,z\}} \to F_{\{x,y,z\}}$ be the automorphism $x \mapsto xy, y \mapsto y, z \mapsto xyz$. Here we give an example of how minimal ideal preimages can be nested in each other. We will state some facts without proof. One can prove these facts using the tools developed in the continuation of this paper. The uword \overline{xz} is a φ -ideal preimage of \overline{xy} . We notice that $\varphi(\overline{xz}) = \overline{xyxyz}$ and for the \overline{xy} marked with an underline \overline{xyxyz} and for any words w_1, w_2 with $w_1 \cdot xz \cdot w_2$ the u-word $\varphi(\overline{w_1xzw_2})$ will contain a copy of the underlined \overline{xy} . As for the other \overline{xy} underlined in \overline{xyxyz} this is not true if we take $w_1 = z^{-1}$ then there is cancellation

$$\varphi\left(\overline{z^{-1}xz}\right) = \overline{z^{-1}y^{-1}x^{-1}xyxyz} = \overline{z^{-1}xyz}$$

But if we take \overline{yxz} then $\varphi(\overline{yxz}) = \overline{yxyxyz}$. Now for every words w_1, w_2 with $w_1 \cdot yxz \cdot w_2$ the u-word $\varphi(\overline{w_1yxzw_2})$ does contains a copy of the \overline{xy} underlined

in \overline{yxyxyz} . Under the naive definition of a minimal ideal preimage \overline{yxz} is not an ideal preimage because it contains the minimal ideal preimage \overline{xz} . But then we cannot count the number of times \overline{xy} appears by counting minimal ideal preimages. Because if w is a cyclic word with a graph morphism $\overline{yxz} \to w$ then \overline{yxz} will contribute two copies of \overline{xy} to $\varphi(w)$ but we will only count one.

Let \overline{u} be a u-word and φ an automorphism and let \overline{v} be a φ -ideal preimage of \overline{u} . We conclude from Examples 18 and 19 that in order to count appearances of \overline{u} in $\varphi(w)$ using ideal preimages our definition need to take into account the specific embedding of \overline{u} in $\varphi(v)$.

Definition 20 (Ideal-preimage). Let φ be an automorphism of a free group. Let u be an u-word. An ideal preimage of u with respect to φ is a pair of a u-word v and a graph morphism $u \to \varphi(v)$ s.t. for every u-word w with $v \to w$ there is a graph morphism $u \to \varphi(w)$ such that the diagram



commutes. Even though we defined an ideal preimage to be a pair $(v, u \to \varphi(v))$ we will denote it $u \to \varphi(v)$ for brevity. We will call the u-word v the base word of $u \to \varphi(v)$.

This definition compares a specific embedding of u in $\varphi(v)$ with a specific embedding of u in $\varphi(w)$ that arises from the specific embedding of $v \to w$. In general a graph morphism $v \to w$ does not give rise to a graph morphisms between $\varphi(v)$ and $\varphi(w)$ but it does give rise to a graph morphism $\tilde{\mathcal{F}}_{\varphi}(v) \to \tilde{\mathcal{F}}_{\varphi}(w)$ cf. Example 14.

Two distinct ideal preimages can have the same base word v but two distinct graph morphisms as in Examples 18 and 19. An ideal preimage $u \to \varphi(v_1)$ is said to be a sub ideal preimage of $u \to \varphi(v_2)$ if there is a graph morphism $v_1 \to v_2$ such that



If $u \to \varphi(v_1)$ and v_2 is a u-word with $v_1 \to v_2$ then there is a morphism $u \to \varphi(v_2)$ satisfying the diagram above thus $u \to \varphi(v_2)$ inherits a structure of ideal preimage. If an ideal preimage has no non-trivial sub ideal preimages we say that it is a minimal ideal preimage.

Proposition 21. Let $u \to \varphi(v)$ be an ideal preimage then for every cyclic word w with $v \to w$ there is a morphism $u \to \varphi(w)$ such that



Proof. Let w' be a word that is a cyclically reduced representative of w. The word $\varphi(w')$ is not necessarily cyclically reduced. We write $\varphi(w') = t^{-1} \cdot \dot{\varphi}(w') \cdot t$ with $\dot{\varphi}(w')$ cyclically reduced. Let k be large enough such that $v \to w'^k$ then by the definition of ideal preimage there is a morphism $u \to \varphi(w')^k$. If we choose k large enough we have a morphism $u \to \dot{\varphi}(w')^k$ and there is a graph morphism from $\dot{\varphi}(w')^k$ to the cyclic word $\varphi(w)$. It is easy to verify that this graph morphism $u \to \varphi(w)$ satisfies the desired diagram.

Definition 22 (Full set of ideal preimages). Let u be a u-word and φ an automorphism. A set of ideal preimages S is called a full set if for every cyclic word w with a morphism $u \to \varphi(w)$ there is a unique ideal preimage $u \to \varphi(v) \in S$ such that $v \to w$ and the diagram



commutes.

Remark 23. It is important that w in the definition is a cyclic word. The definition would be null if we change cyclic word by u-word. Let $\varphi : F_{\{x,y\}} \to F_{\{x,y\}}$ be the automorphism $x \mapsto xy, y \mapsto y, u$ be the u-word $u = \overline{y}$ and w be the u-word $w = \overline{x}$. Then there is a graph morphism $\overline{y} \to \varphi(\overline{x}) = \overline{xy}$ but \overline{x} is not an ideal preimage of \overline{y} .

Definition 24 (A full paradigm). Let $\overline{u}, \overline{v}$ be u-words, φ be an automorphism, $\overline{u} \to \varphi(\overline{v})$ a graph morphism and let v be an orientation of \overline{v} . A full paradigm of $\overline{u} \to \varphi(\overline{v})$ is a set $\overline{u} \to \varphi(\overline{v_1}), \ldots, \overline{u} \to \varphi(\overline{v_{2n-1}})$ such that

$$\{\overline{v_1},\ldots,\overline{v_{2n-1}}\} = \{\overline{vx}|x \in X \cup X^{-1}, v \cdot x\}$$

or

$$\{\overline{v_1},\ldots,\overline{v_{2n-1}}\} = \{\overline{xv}|x \in X \cup X^{-1}, x \cdot v\}$$

and the diagram



commutes.

Remark 25. Let $u \to \varphi(v)$ and let $u \to \varphi(v_1), \ldots, u \to \varphi(v_l)$ be a full paradigm. It is not hard to see that $u \to \varphi(v)$ is an ideal preimage if and only if $u \to \varphi(v_1), \ldots, u \to \varphi(v_l)$ are all ideal preimages.

Remark 26. Let u be a u-word, φ an automorphism, S a set of ideal preimages and $u \to \varphi(v) \in S$. Let

$$S' = (S - \{u \to \varphi(v)\}) \cup \{u \to \varphi(v_1), \dots, u \to \varphi(v_l)\}$$

such that $u \to \varphi(v_1), \ldots, u \to \varphi(v_l)$ is a full paradigm of $u \to \varphi(v)$ then S' is a full set if and only if S is a full set.

Definition 27. A full set is minimal if it does not contain a full paradigm.

Remark 28. Any full set can be reduced to a minimal full set by repeating Remark 26.

Theorem 29. Every u-word u and automorphism φ has a unique finite minimal full set. Moreover it is the set of all minimal ideal preimages.

Theorem 1 follows directly. The multiset $S_{u,\varphi}$ is constructed by taking the base words of the unique minimal full set. We show first that it is unique then we will show it exists.

Fact 30. Let v be a u-word let φ an automorphism recall that we denote the folding morphism by $\rho_v : \mathcal{F}_{\varphi}(v) \to \tilde{\mathcal{F}}_{\varphi}(v)$. We observe that the graph $\tilde{\mathcal{F}}_{\varphi}(v)$ is a tree and that the subgraph $\varphi(v)$ is a line segment. We observe that an edge e in $\tilde{\mathcal{F}}_{\varphi}(v)$ belongs to the subgraph $\varphi(v)$ if and only if the size of the set of preimages $|\rho_v^{-1}(e)|$ is odd.

Proposition 31. Let $\overline{v}, \overline{u}$ be u-words let v be an orientation of \overline{v} s.t. $v = v_1 \cdot v_2$, let $\overline{u} \to \tilde{\mathcal{F}}_{\varphi}(\overline{v})$ and let e be an edge contained in the image of \overline{u} . We clarify the statement $|\rho_v^{-1}(e)| = |\rho_{v_1}^{-1}(e)| + |\rho_{v_2}^{-1}(e)|$ and show it holds.

Proof. First we claim that $\tilde{\mathcal{F}}_{\varphi}(\overline{v_1})$ and $\tilde{\mathcal{F}}_{\varphi}(\overline{v_2})$ both embed in $\tilde{\mathcal{F}}_{\varphi}(\overline{v})$. Because $\tilde{\mathcal{F}}_{\varphi}(\overline{v_1})$ is a folded graph the function $\tilde{\mathcal{F}}_{\varphi}(\overline{v_1}) \to \tilde{\mathcal{F}}_{\varphi}(\overline{v})$ is locally injective. Following the fact that both $\tilde{\mathcal{F}}_{\varphi}(\overline{v_1})$ and $\tilde{\mathcal{F}}_{\varphi}(\overline{v})$ are trees a locally injective function must be globally injective. If there exists an edge e that satisfies



it is unique because all the function in this square are injective. By e in $|\rho_{v_1}^{-1}(e)|$ we mean this unique edge if it exists if it does not exist we consider $|\rho_{v_1}^{-1}(e)| = 0$. We notice that the graph $\mathcal{F}_{\varphi}(\overline{v_1} \cdot v_2)$ is a line segment obtained by gluing $\mathcal{F}_{\varphi}(\overline{v_1})$ and $\mathcal{F}_{\varphi}(\overline{v_2})$ along a vertex. Therefore there are no edges in $\mathcal{F}_{\varphi}(\overline{v_1}) \cap \mathcal{F}_{\varphi}(\overline{v_2})$. Consequently we get

$$\left|\rho_{v}\right|_{\mathcal{F}_{\varphi}(\overline{v_{1}})}^{-1}\left(e\right)\right| + \left|\rho_{v}\right|_{\mathcal{F}_{\varphi}(\overline{v_{2}})}^{-1}\left(e\right)\right| = \left|\rho_{v}^{-1}\left(e\right)\right|$$

Let $\mathcal{F}_{\varphi}(\overline{v_1}) \mathcal{F}_{\varphi}(\overline{v_2})$ be the graph obtained by folding all vertexes in $\mathcal{F}_{\varphi}(\overline{v_1 \cdot v_2})$ besides the vertex connecting $\mathcal{F}_{\varphi}(\overline{v_1})$ and $\mathcal{F}_{\varphi}(\overline{v_2})$. The function $\rho_{v_1 \cdot v_2}$ factors through $\tilde{\mathcal{F}}_{\varphi}(\overline{v_1}) \tilde{\mathcal{F}}_{\varphi}(\overline{v_2})$ because the order of folding does not effect the final folded graph. The following diagram



commutes. Thus $\left|\rho_{v_1 \cdot v_2}\right|_{\mathcal{F}_{\varphi}(\overline{v_1})}^{-1}(e)\right| = \left|\rho_{v_1}^{-1}(e)\right|.$

Lemma 32. Let $\overline{u} \to \varphi(\overline{v})$ be a minimal ideal-preimage and let w be a cyclic word with $\overline{v} \to w$. Let $\overline{u} \to \varphi(\overline{y})$ be an ideal preimage with $\overline{y} \to w$ such that



commutes. Then $\overline{u} \to \varphi(\overline{v})$ is a sub ideal preimage of $\overline{u} \to \varphi(\overline{y})$.

Proof. Assume by contradiction that it is not. Let v, y be orientations of $\overline{v}, \overline{y}$ and let w' be a cyclically reduced representative of w. There are essentially 4 cases up to symmetry.

- 1. y is a proper subword of v.
- 2. Assume l(v) < l(w'), write $w' = v \cdot w_0$ and consider y to be w_0 or a subword of w_0 .
- 3. y includes a subword both of v and of w_0 . Write $v = v_1 \cdot v_2$ and $w_0 = w_{01} \cdot w_{02}$ and $y = v_2 \cdot w_{01}$.
- 4. Assume $l(v) \ge l(w)$ and y is a cyclic permutation of v or a subword of a cyclic permutation of w.



Case 1: is in contradiction to the fact that v is minimal. Case 2: There is graph morphism from $\overline{v \cdot w_0}$ the cyclic word w. Let e be an edge in the image of $u \to \varphi(\overline{v \cdot w_0}) \to \tilde{\mathcal{F}}_{\varphi}(\overline{v \cdot w_0})$. Then $|\rho_{v \cdot w_0}^{-1}(e)|$ is odd. Observe that $|\rho_v^{-1}(e)|$ is odd because v is an ideal preimage. Thus $|\rho_{w_0}^{-1}(e)|$ is even because

$$\left|\rho_{v \cdot w_{0}}^{-1}\left(e\right)\right| = \left|\rho_{v}^{-1}\left(e\right)\right| + \left|\rho_{w_{0}}^{-1}\left(e\right)\right|$$

Therefore there cannot be an ideal preimage $\overline{u} \to \varphi(\overline{w_0})$. Let $\overline{y} \to \overline{w_0}$. If $\overline{u} \to \varphi(\overline{y})$ is an ideal preimage then so is $\overline{u} \to \varphi(\overline{w_0})$ this a contradiction. Case 3: let e be an edge in the image of \overline{u} then $|\rho_{v_2 \cdot w_{01}}^{-1}(e)|$ is odd by assumption. We observe that

$$\left|\rho_{v \cdot w_{01}}^{-1}(e)\right| = \left|\rho_{v}^{-1}(e)\right| + \left|\rho_{w_{01}}^{-1}(e)\right|$$

because $|\rho_{v \cdot w_{01}}^{-1}(e)|$ and $|\rho_{v}^{-1}(e)|$ are odd then $|\rho_{w_{01}}^{-1}(e)|$ is even. We observe that

$$\left|\rho_{v_{2}\cdot w_{01}}^{-1}\left(e\right)\right| = \left|\rho_{v_{2}}^{-1}\left(e\right)\right| + \left|\rho_{w_{01}}^{-1}\left(e\right)\right|$$

and $|\rho_{v_2 \cdot w_{01}}^{-1}(e)|$ is odd and $|\rho_{w_{01}}^{-1}(e)|$ is even then $|\rho_{v_2}^{-1}(e)|$ is odd. We chose e arbitrarily thus for every edge e in the image of \overline{u} we get $|\rho_{v_2}^{-1}(e)|$ is odd. Therefore there is a graph morphism $\overline{u} \to \varphi(\overline{v_2}) \to \tilde{\mathcal{F}}_{\varphi}(\overline{v_2})$. Because $\overline{u} \to \varphi(\overline{v})$ is minimal $\overline{u} \to \varphi(\overline{v_2})$ is not an ideal preimage thus there are words x_1 and x_2 with $x_1 \cdot v_2 \cdot x_2$ and an edge e in the image of \overline{u} s.t. $|\rho_{x_1 \cdot v_2 \cdot x_2}^{-1}(e)|$ is even. From the fact that

$$\left|\rho_{v \cdot x_{2}}^{-1}\left(e\right)\right| = \left|\rho_{v}^{-1}\left(e\right)\right| + \left|\rho_{x_{2}}^{-1}\left(e\right)\right|$$

we get that $|\rho_{x_2}^{-1}(e)|$ is even thus we get that $|\rho_{x_1 \cdot v_2}^{-1}(e)|$ must be even as well. Finally we have

$$\left|\rho_{x_{1}\cdot v_{2}\cdot w_{01}}^{-1}\left(e\right)\right| = \left|\rho_{x_{1}\cdot v_{2}}^{-1}\left(e\right)\right| + \left|\rho_{w_{01}}^{-1}\left(e\right)\right|$$

but we saw that $|\rho_{x_1 \cdot v_2}^{-1}(e)|$ is even and $|\rho_{w_{01}}^{-1}(e)|$ is even thus we get that $|\rho_{x_1 \cdot v_2 \cdot w_{01}}^{-1}(e)|$ is even and this is a contradiction to $v_2 \cdot w_{01}$ being an ideal preimage. Case 4: is proved in an almost identical way to Case 3.

Lemma 33. Let $\overline{u} \to \varphi(\overline{v})$ be a minimal ideal-preimage and let S be a finite full set that does not include $\overline{u} \to \varphi(\overline{v})$. Then S is not minimal i.e. it contains a full paradigm.

Proof. Let $m = \max \{ l(\overline{v}) | \overline{u} \to \varphi(\overline{v}) \in S \}$ the maximal length of the base words in the full set. Let v be an orientation of \overline{v} an let

$$M = \{ \overline{x_1 v x_2} \in F_n | x_1 \cdot v \cdot x_2, l(x_1 v) = l(v x_2) = m \}$$

a set of u-words. We notice that the u-words in M inherit a structure of φ ideal preimages of \overline{u} from the embedding of \overline{v} . We construct a set of cyclic words \overline{M} as following: If $w \in M$ is cyclically reduced we glue its ends; If $w \in M$ is not cyclically reduced we add a letter to the end before we glue. From the construction of \overline{M} every cyclic $w \in \overline{M}$ word has a graph morphism $\overline{v} \to w$. Because $\overline{u} \to \varphi(\overline{v})$ is an ideal preimage every cyclic word $w \in \overline{M}$ has a morphism $\overline{u} \to \varphi(w)$ satisfying the appropriate diagram. From the second property of a full set every $w \in \overline{M}$ has unique ideal preimage $\overline{u} \to \varphi(\overline{v_w})$ with $\overline{v_w} \to w$ satisfying the correct diagram. We show that the set $X = \{\overline{u} \to \varphi(\overline{v_w}) | w \in \overline{M}\} \subset S$ contains a full paradigm. By Lemma 32 every $\overline{u} \to \varphi(\overline{v_w})$ has $\overline{u} \to \varphi(\overline{v})$ as a sub ideal preimage. Particularly there is a graph morphism $\overline{v} \to \overline{v_w} \to w$. By assumption $\Lambda(\overline{v_w}) \leq m$ and by the construction M and \overline{M} there must be a $\overline{y} \in M$ such that $\overline{v} \to \overline{v_w} \to \overline{y} \to w$. We now translate the problem to the language of Posets. We define a Poset on the set $P = \{\overline{x} | y \in M, \overline{v} \to \overline{x} \to \overline{y}\}$ (the composition $\overline{v} \to \overline{x} \to \overline{y}$ is the original embedding of \overline{v} in \overline{y}) by $x_1 \leq x_2$ if and only if



We notice that $\{\overline{v_w} | \overline{u} \to \varphi(\overline{v_w}) \in X\} \subset P$, \overline{v} is the minimal element of P and M is the set of maximal elements of P. As consequence of the second property of a full set and the discussion above for all $y \in M$ there exists a unique word $x \in X$ such that $x \leq y$. We define posets $T_2 = \{x | v \cdot x\}$ with $x_1 \leq x_2$ if and only if x_1 is a prefix of x_2 and $T_1 = \{x | x \cdot v\}$ with $x_1 \leq x_2$ if and only if x_1 is a suffix of x_2 . We define the order $(x_1, x_2) \leq (y_1, y_2)$ if and only if $x_1 \leq y_1 \wedge x_2 \leq y_2$ on $T_1 \times T_2$ and we notice that P can be seen as a suborder of $T_1 \times T_2$. Let T be the poset defined by a directed 2n - 1 regular tree such that the root of T is the minimal element then $T \cong T_1 \cong T_2$. A poset T that is defined by a directed tree is characterized by the following property: for all $a \in T$ the subposet $\{b \in T | b \leq a\}$ is a total order. We denote the root, the minimal element, of T by r. We define $l: T \to \mathbb{N}$ by setting l(a) to be the distance of a from the root of the tree. We notice that l is equivalent to the length of a word in T_1 and T_2 . We say $a, b \in T$ are neighbors if there exists an element $c \in T$ such that $c < a \land c < b$ and l(a) = l(b) = l(c) + 1. In T_2 or T_1 two words are neighbors if they only differ by the last or first letter. This motivates the next definition. Let $a, b_0 \in A$. A subset of $T \times T$ is said to be a full paradigm if it has the form

$$\{(a, b) \in T \times T | b_0 < b, l(b) = l(b_0) + 1\}$$

or

$$\{(b, a) \in T \times T | b_0 < b, l(b) = l(b_0) + 1\}$$

We complete the translation by defining

$$P = \{(a, b) \in T \times T | l(a) + l(b) \le 2m - l(v)\}$$

and

$$M = \{(a, b) \in T \times T | l(a) = l(b) = m - l(v)\}$$

and $X \subset P$ is a set that satisfies for all $y \in M$ there exists a unique element $x \in X$ such that $x \leq y$. We need to show that if $(r, r) \notin X$ then X contains a full paradigm.

Let

$$X_R = \{ b \in T | \exists a \in T \ (a, b) \in X \}$$

and let

$$X_R^m = \{ b \in X_R | \forall c \in X_R \, l \, (c) \le l \, (b) \}$$

the set of elements of maximal length in X_R . Let

$$X_{LX_R^m} = \{a \in T | \exists b \in X_R^m \ (a,b) \in X\}$$

Let $(a_1, b_1) \in X$ such that $b_1 \in X_R^m$ and $a_1 \in X_{LX_R^m}$ is such that $l(a_1)$ is minimal among $X_{LX_R^m}$. Let b_1, \ldots, b_{2n-1} the neighbors of b_1 . Let

$$(a_1^m, b_2^m) \in \{(a, b) \in P | (a_1, b_2) < (a, b)\} \cap M$$

then there is an element in $(x, y) \in X$ such that $(x, y) \leq (a_1^m, b_2^m)$ i.e. $x \leq a_1^m$ and $y \leq b_2^m$. We recall that $\{x \in T | x \leq a_1^m\}$ and $\{y \in T | y \leq b_2^m\}$ are total orders because T is a tree. The element b_2 is a neighbor of b_1 therefore it is maximal and $y \leq b_2$. By way of contradiction suppose $y < b_2$ then $y < b_1$ because b_1 is a neighbor of b_2 . If $x \leq a_1$ then $(x, y) < (a_1, b_1)$ this is a contradiction to the uniqueness in the definition of X. If $x > a_1$ then $(x, y) < (x, b_1)$ and $(a_1, b_1) < (x, b_1)$ also a contradiction to the uniqueness thus $y = b_2$. We chose a_1 to be minimal among $X_{LX_R^m}$ therefore $x \geq a_1$. If $x = a_1$ and $(a_1, b_1), (a_1, b_2), \ldots, (a_1, b_{2n-1}) \in X$ then X has a full paradigm and we are done. If not, without loss of generality, we can assume $(a_1, b_2) \notin X$ and $a_1 < x$. Let

$$X_{Lb_2} = \{ a \in T | (a, b_2) \in X \}$$

and let $c_1 \in X_{Lb_2}$ an element such that $l(c_1)$ is maximal among X_{Lb_2} then $(c_1, b_2) \in X$ and $a_1 < c_1$. Let c_2 be a neighbor of c_1 and let

$$(c_2^m, b_2^m) \in \{(a, b) \in P | (c_2, b_2) < (a, b)\} \cap M$$

then there is an element $(x, y) \in X$ such that $(x, y) \leq (c_2^m, b_2^m)$ (this is a different (x, y) then before). By way of contradiction assume $y < b_2$ then $y < b_1$ because b_1 is a neighbor of b_2 . Because c_2 is a neighbor of c_1 and $a_1 < c_1$ then $a_1 < c_2$. Because $\{x | x \leq c_2^m\}$ is a total order and $a_1 \in \{x | x \leq c_2^m\}$ then either $x < a_1$ or $x \geq a_1$. If $x < a_1$ then $(x, y) < (a_1, b_1)$ and if $a_1 \leq x$ then $(a_1, b_1) \leq (x, b_1)$ and $(x, y) < (x, b_1)$ both contradict the uniqueness condition. Thus $y \geq b_2$ but b_2 is maximal so $y = b_2$. By way of contradiction assume $x < c_2$. Because c_2 is a neighbor of c_1 then $x < c_1$ and we conclude $(x, b_2) < (c_1, b_2)$ which is a contradiction to the uniqueness condition of X. Thus $x \geq c_2$ but c_2 is maximal because it is a neighbor of c_1 which is maximal so $x = c_2$. We get that $(c_2, b_2) \in X$ but c_2 was an arbitrary neighbor of c_1 therefore for all c_i a neighbor of c_1 we have $(c_i, b_2) \in X$ i.e. X contains a full paradigm.

We now show that every automorphism φ and u-word u has a finite full set of ideal preimages. We call an automorphism φ a good automorphism if every u has a finite full set of φ ideal preimages. We will show that if φ and ψ are good automorphisms then $\varphi \circ \psi$ is also a good automorphism. We conclude by showing that the Nielsen move $x_1 \mapsto x_1 x_2, x_j \mapsto x_j$ for $j \neq 1$ is a good automorphism by showing how to construct a full set of ideal preimages for every u. It is trivial that the other Nielsen moves are good. Thus a generating set of automorphisms are good and all automorphisms generated by it.

Proposition 34. Let $\varphi = \psi_1 \circ \psi_2$ be an automorphism such that ψ_1, ψ_2 are both good automorphisms then φ is a good automorphism.

Proof. Let u be a u-word then there is a ψ_1 -full set $u \to \psi_1(v_1), \ldots, u \to \psi_1(v_l)$ and every v_i has a ψ_2 full set $v_i \to \psi_2(v_i^1), \ldots, v_i \to \psi_2(v_i^{l_i})$. Let w be a cyclic word with $v_i^k \to w$ then we have



From $v_i \to \psi_2(w)$ we get the diagram



From $v_i \to \psi_2\left(v_i^k\right)$ we get



We combine these three diagrams to one diagram after applying the functor $\tilde{\mathcal{F}}_{\psi_1}$ to the first diagram and we get:



We notice that generally $\tilde{\mathcal{F}}_{\psi_1} \circ \tilde{\mathcal{F}}_{\psi_2} \neq \tilde{\mathcal{F}}_{\psi_1 \circ \psi_2}$ (see Proposition 3.9 in [3]) thus the previous diagram is not sufficient to show that $u \to \psi_1 \circ \psi_2 \left(v_i^k \right)$ is an ideal preimage. In order to prove that $u \to \psi_1 \circ \psi_2 \left(v_i^k \right)$ is ideal preimage we need

three more commutative diagrams. For every labeled graphs w_1 and w_2 with $w_1 \to w_2$ there is a commutative diagram

$$\begin{array}{cccc} \tilde{\mathcal{F}}_{\psi_{1}\circ\psi_{2}}\left(w_{1}\right) & \longrightarrow \tilde{\mathcal{F}}_{\psi_{1}\circ\psi_{2}}\left(w_{2}\right) \\ & & \downarrow \\ & & \downarrow \\ \tilde{\mathcal{F}}_{\psi_{1}}\circ\tilde{\mathcal{F}}_{\psi_{2}}\left(w_{1}\right) & \longrightarrow \tilde{\mathcal{F}}_{\psi_{1}}\circ\tilde{\mathcal{F}}_{\psi_{2}}\left(w_{2}\right)
\end{array}$$

For every u-word or cyclic word w_1 we have a commutative diagram

$$\begin{array}{c} \psi_{1} \circ \psi_{2}\left(w_{1}\right) \longrightarrow \tilde{\mathcal{F}}_{\psi_{1} \circ \psi_{2}}\left(w_{1}\right) \\ \downarrow \\ \tilde{\mathcal{F}}_{\psi_{1}}\left(\psi_{2}\left(w_{1}\right)\right) \longrightarrow \tilde{\mathcal{F}}_{\psi_{1}} \circ \tilde{\mathcal{F}}_{\psi_{2}}\left(w_{1}\right) \end{array}$$

We combine all the commuting diagrams together to get Diagram 3.3



Using all the commutative diagrams above one can observe that

$$u \to \psi_1 \circ \psi_2 \left(v_i^k \right) \to \tilde{\mathcal{F}}_{\psi_1 \circ \psi_2} \left(v_i^k \right) \to \tilde{\mathcal{F}}_{\psi_1 \circ \psi_2} \left(w \right) \to \tilde{\mathcal{F}}_{\psi_1} \circ \tilde{\mathcal{F}}_{\psi_2} \left(w \right)$$
$$= u \to \psi_1 \circ \psi_2 \left(w \right) \to \tilde{\mathcal{F}}_{\psi_1 \circ \psi_2} \left(w \right) \to \tilde{\mathcal{F}}_{\psi_1} \circ \tilde{\mathcal{F}}_{\psi_2} \left(w \right)$$

But $\tilde{\mathcal{F}}_{\psi_1 \circ \psi_2}(w) \to \tilde{\mathcal{F}}_{\psi_1} \circ \tilde{\mathcal{F}}_{\psi_2}(w)$ is injective because it is a locally injective function between trees. We conclude that the diagram



is commutative. Thus $u \to \psi_1 \circ \psi_2 \left(v_i^k \right)$ is an ideal preimage. We want to show that $u \to \psi_1 \circ \psi_2 \left(v_1^1 \right), \ldots, u \to \psi_1 \circ \psi_2 \left(v_1^{l_1} \right), u \to \psi_1 \circ \psi_2 \left(v_2^1 \right), \ldots, u \to \psi_1 \circ \psi_2 \left(v_2^{l_2} \right), \ldots, u \to \psi_1 \circ \psi_2 \left(v_2^{l_2$

 $\psi_1 \circ \psi_2 \left(v_l^{l_l} \right)$ is a full set. Let $u \to \psi_1 \circ \psi_2 (w)$ then because $u \to \psi_1 (v_1), \ldots, u \to \psi_1 (v_l)$ is a full set there exists a $v_i \to \psi_2 (w)$ satisfying Diagram 3.2. Because $v_i \to \psi_2 \left(v_i^1 \right), \ldots, v_i \to \psi_2 \left(v_i^{l_i} \right)$ is a full set there exists $v_i^k \to w$ satisfying Diagram 3.1 thus for every $u \to \psi_1 \circ \psi_2 (w)$ there exists ideal preimage $u \to \psi_1 \circ \psi_2 (v_i^k)$ with $v_i^k \to w$ that satisfies Diagram 3.4. Suppose there is another $u \to \psi_1 \circ \psi_2 \left(v_{i'}^k \right)$ with $v_{i'}^{k'} \to w$ satisfying Diagram 3.4. But $v_{i'} \to \psi_2 \left(v_{i'}^{k'} \right)$ is an ideal preimage so $v_{i'}^{k'} \to w$ give us $v_{i'} \to \psi_2 (w)$ satisfying Diagram 3.1 and $v_{i'} \to \psi_2 (w)$ give us $u \to \psi_1 \circ \psi_2 (w)$ satisfying Diagram 3.2. But $u \to \psi_1 (v_1), \ldots, u \to \psi_1 (v_l)$ a full set therefor there is a unique ideal preimage satisfying Diagram 3.1 so $v_{i'}^{k'} = v_i^k$.

Lemma 35. Let φ be the automorphism of F_X sending $x_1 \mapsto x_1 x_2, x_i \mapsto x_i$ for $i \neq 1$ (this is a Nielsen move). We show that every u-word \overline{u} has a finite full set of ideal preimages

We notice that φ satisfies the following special case of Thurston's bounded cancellation lemma (see [1]). Let v_1 and v_2 be words such that $v_1 \cdot v_2$. Then at most one letter can be canceled in the multiplication $\varphi(v_1) \varphi(v_2)$. Thus to construct an ideal-preimages for a u-word \overline{u} we need to change at most one letter in each affix of $\varphi^{-1}(\overline{u})$. Let u be an orientation of \overline{u} a suffix of u and a suffix of u^{-1} are said to be pointing out and a prefix of u and prefix of u^{-1} are said to be pointing in. We classify 8 different kinds of affixes pointing out. We classify the affixes by examining the affix starting from the most outer letter looking in until we find the first letter that is not x_2 or x_2^{-1} . These affixes are the minimal subword of w that allow us to know the types of affixes of $\varphi(w)$. A word u can have multiple different ideal preimages. In Table 1 they are either separated by commas or expressed by the letter y standing in for different possible letters. We denote the ideal preimages of u by v_1, \ldots, v_l .

#		suffix of u	suffix of $\varphi^{-1}(u)$	suffixes of v_i	suffixes of $\varphi(v_i)$	
1	$z \neq x_1, x_2, x_2^{-1}$	z	z	z		
2		x_1	$x_1 x_2^{-1}$	x_1	$x_1 x_2$	
3	$z \neq x_1, n \ge 1; y \neq x_1, x_1^{-1}$	zx_2^n	zx_2^n	$zx_2^nx_1, zx_2^ny$	$zx_2^nx_1x_2, zx_2^ny$	
4	$y \neq x_1, x_2^{-1}, x_1^{-1}$	$x_1 x_2$	x_1	x_1x_1, x_1y	$x_1x_2x_1x_2, x_1x_2y$	
5	$n \ge 2; \ y \ne x_1, x_1^{-1}$	$x_1 x_2^n$	$x_1 x_2^{n-1}$	$x_1 x_2^{n-1} x_1, x_1 x_2^{n-1} y$	$x_1x_2^nx_1x_2, x_1x_2^ny$	
6	$z \neq x_1 n \ge 1$	zx_2^{-n}	zx_2^{-n}	$zx_2^{-n}, zx_2^{-n+1}x_1^{-1}$	$zx_2^{-n}, zx_2^{-n}x_1^{-1}$	
7	$n \ge 1$	$x_1 x_2^{-n}$	$x_1 x_2^{-n-1}$	$x_1 x_2^{-n-1}, x_1 x_2^{-n} x_1^{-1}$	$x_1 x_2^{-n}, x_1 x_2^{-n} x_1^{-1}$	

Table 1: φ -ideal preimages

Example 36. We construct the set of ideal preimages of $\overline{u} = \overline{x_2^{-1}x_1^{-1}x_1^{-1}x_3x_1}$ using Table 1. We start by classifying the two affixes of \overline{u} . The first affix is x_1 . It is of type 2. The second is x_1x_2 . It is type 4. We calculate $\varphi^{-1}(\overline{u}) = \overline{x_1^{-1}x_2x_1^{-1}x_3x_1x_2^{-1}}$ we follow the table and we get the following set of ideal

preimages

$$\frac{\overline{x_1^{-1}x_1^{-1}x_2x_1^{-1}x_3x_1}, \overline{x_2^{-1}x_1^{-1}x_2x_1^{-1}x_3x_1},}{\overline{x_3x_1^{-1}x_2x_1^{-1}x_3x_1}, \overline{x_3^{-1}x_1^{-1}x_2x_1^{-1}x_3x_1}, \dots, \overline{x_n^{-1}x_1^{-1}x_2x_1^{-1}x_3x_1}}$$

as for the morphism $\overline{u} \to \overline{\varphi(v_i)}$ we mark \overline{u} in $\overline{\varphi(v_i)}$ by an underline

$$\frac{\overline{x_2^{-1}x_1^{-1}x_2^{-1}x_1^{-1}x_3x_1x_2}, \overline{x_2^{-1}x_2^{-1}x_1^{-1}x_1^{-1}x_3x_1x_2},}{\overline{x_3x_2^{-1}x_1^{-1}x_1^{-1}x_3x_1x_2}, \overline{x_3^{-1}x_2^{-1}x_1^{-1}x_1^{-1}x_3x_1x_2}, \dots, \overline{x_n^{-1}x_2^{-1}x_1^{-1}x_1^{-1}x_3x_1x_2}}$$

Remark 37. Even though affixes were defined to be a proper subword, in these definitions we allow for an affix to be the whole word. For instance if $u = \overline{x_1 x_1^n}$ we classify the two affixes of u to be the following: $x_1 x_2^n$ this is type 5 and x_1^{-1} pointing out this is type 2. Another limit case is the word is $u = \overline{x_2^n}$. In this case we will not find a first letter that is not x_2 or x_2^{-1} . We will treat one affix as type 3 and the other as type 6.

To proof that $u \to \varphi(v_i)$ constructed using Table 1 is an ideal preimage one needs to verify that for every words w_1, w_2 such that $w_1 \cdot v_i \cdot w_2$ no letter of the subword u in $\varphi(v_i)$ cancels in the multiplication $\varphi(w_1)\varphi(v_i)\varphi(w_2)$. This is straight forward verification because one just has to check finite combinations of the affix types mentioned in Table 1. We have verified this but choose not to write it down because of its length. We would like to show that a set of ideal preimages constructed using Table 1 is a full set. We notice that φ^{-1} is the automorphism sending $x_1 \mapsto x_1 x_2^{-1}, x_i \mapsto x_i$ for $i \neq 1$. In order to compute φ^{-1} ideal preimages we can use Table 1 swapping x_2 with x_2^{-1} . Denote by v_1, \ldots, v_l the set of φ -ideal preimages of u we obtain by using Table 1 and denote by $v_i^1, \ldots, v_i^{l_i}$ the φ^{-1} -ideal preimages of v_i obtained using Table 1. In Table 2 we calculate the suffixes of v_i^k .

#		suffix of u	suffixes of v_i	suffixes of v_i^k			
1-2	$z \neq x_2, x_2^{-1}$	z	z	z			
	$r \neq r_1$ $n > 1$		$zx_2^nx_1$	$zx_2^nx_1$			
3	$z \neq x_1, n \geq 1,$ $u \neq x_1, x^{-1}, x_2$	zx_2^n	$zx_2^nx_2$	$zx_2^n x_2, zx_2^n x_1^{-1}$			
	$g \neq x_1, x_1$, x_2		zx_2^ny	zx_2^ny			
	$n \neq$		$x_1 x_1$	$x_1 x_2 x_1$			
4	$y \neq x_1 x_2 x^{-1} x^{-1}$	$x_1 x_2$	$x_1 x_2$	$x_1x_2x_2, x_1x_2x_1^{-1}$			
	x_1, x_2, x_2, x_1		x_1y	x_1x_2y			
	n > 2		$x_1 x_2^{n-1} x_1$	$x_1 x_2^n x_1$			
5	$u \neq x_1 x^{-1} x_2$	$x_1 x_2^n$	$x_1 x_2^{n-1} x_2$	$x_1 x_2^n x_2, x_1 x_2^n x_1^{-1}$			
	$y \neq x_1, x_1$, x_2		$x_1 x_2^{n-1} y$	$x_1 x_2^n y$			
6	$z \neq x_1, n \ge 1,$	$\sim n^{-n}$	$zx_2^{-n+1}x_1^{-1}$	$zx_2^{-n}x_1^{-1}$			
0	$y \neq x_1^{-1}$	zx_2	zx_2^{-n}	$zx_2^{-n}y$			
7	$n > 1$ $u \neq r^{-1}$	$x - x^{-n}$	$x_1 x_2^{-n} x_1^{-1}$	$x_1 x_2^{-n} x_1^{-1}$			
1	$n \ge 1, y \ne x_1$	x_1x_2	$x_1 x_2^{-n-1}$	$x_1 x_2^{-n} y$			

Table 2: φ^{-1} -ideal preimages

Because $v_i \to \varphi^{-1}(v_i^k)$ we get $u \to \varphi \circ \varphi^{-1}(v_i^k) = v_i^k$. Examining Table 2 we notice that in cases 3-7 for every v_i the suffixes of v_i^k for every k form a full

paradigm of the suffixes of u and in cases 1-2 the suffix of v_i^k is the suffix of u.

Thus for every $u \to \varphi(w)$ there exists a v_i^k s.t. $u \to v_i^k \to \varphi(w)$. Let $u \to \varphi(w)$ then there exists $u \to v_i^k \to \varphi(w)$. From v_i^k being an ideal preimage we get that $v_i \to \varphi^{-1} \circ \varphi(w) = w$. If we plug these maps into Diagram 3.3 we get that $v_i \to w$ is the ideal preimage of $u \to \varphi(w)$. This shows that for every $u \to \varphi(w)$ there is an ideal-preimage in the set of ideal preimages constructed using Table 1.

Now we prove that it is unique. Examining Table 2 we notice that if $v_i \neq v_j$ for every k_1, k_2 the push-out of the square



is a folded tree with at least one vertex of degree 3. Let $u \to \varphi(w)$. Assume by way of contradiction that both $v_i \to w$ and $v_i \to w$ are ideal preimages i.e.



commutes. Because v_{i}^{k} were constructed using Table 1 there are $v_{i}^{k_{1}} \rightarrow \varphi(w)$ and $v_{i}^{k_{2}} \rightarrow \varphi\left(w\right)$ such that



commutes. But combining these diagrams and plugging into Diagram 3.3 we

get the diagram



We conclude that the square



commutes. This implies a locally injective morphism $v_i^{k_1} \coprod_u v_j^{k_2} \to \varphi(w)$. All vertices in $\varphi(w)$ are of degree 2 but there is a vertex of degree 3 in $v_i^{k_1} \coprod_u v_j^{k_2}$ which is a contradiction. This concludes the proof of Lemma 35.

Let u be an u-word an φ automorphism. We conclude this part with a method for calculating the unique minimal full set $S_{u,\varphi}$ (if we take the base words of the ideal preimages we get the multiset mentioned in the introduction). First we decompose φ into Nielsen moves $\varphi = \varphi_1 \circ \varphi_2 \circ \cdots \circ \varphi_l$ by combining moves we can assume that every φ_k is of the form $x_i \mapsto x_i x_j, x_m \mapsto x_m$ with $m \neq i$ or $x_i \mapsto x_j x_i, x_m \mapsto x_m$ with $m \neq i$. Now we use Table 1 to construct a full set S_{u,φ_1} of φ_1 ideal preimages of u. Now for every $v \in S_{u,\varphi_1}$ we use Table to construct a full set S_{v,φ_2} of φ_2 ideal preimages of v. We have shown that the set $\bigcup_{v \in S_{u,\varphi_1}} S_{v,\varphi_2}$ is a full set but generally it is not minimal. We can use Remark 26 to replace $\bigcup_{v \in S_{u,\varphi_1}} S_{v,\varphi_2}$ by the minimal set $S_{u,\varphi_1 \circ \varphi_2}$ and now recursively continue this process to φ_3 and onward.

4 Constructions of Inverse limit and Representation

4.1 Construction of inverse limit

We know wish to create an inverse system of modules by defining $p_k : M_k \to M_{k-1}$ such that $p_k \pi_k = \pi_{k-1}$. We make two auxiliary definitions

Definition 38 (U-word orientation). Let f: reduced-words \rightarrow u-words the function that forgets the orientation. A u-word orientation is a section σ :

u-words \rightarrow reduced-words s.t. $f \circ \sigma = \text{Id}$ choosing for every u-word an orientation. Let σ be an u-word orientation, let \overline{u} be an u-word then $\sigma(\overline{u}) = u$ or $\sigma(\overline{u}) = u^{-1}$. Suppose $\sigma(\overline{u}) = u$ we say u is the orientation of \overline{u} . The word u is also the orientation of $\overline{u^{-1}}$. This is an auxiliary definition in order to make definitions precise. We will mostly be interested in objects that are independent of this choice. We choose an arbitrary u-word orientation we will use until the end of the article.

Definition 40. We define two sequences of homomorphisms $\overleftarrow{p}_k : M_k \to M_{k-1}$ and $\overrightarrow{p}_k : M_k \to M_{k-1}$. We first make auxiliary definition. Let u be a length k-1u-word and w a length k u-word we define the functionals $\overleftarrow{\delta}_u, \overrightarrow{\delta}_u \in M_k^*(F_n)$ by

$$\overleftarrow{\delta}_{u}(w) = \begin{cases} 1 & \text{If } u \text{ is } k-1 \text{ affix of } w \text{ pointing outward} \\ 0 & \text{Otherwise} \end{cases}$$
$$\overrightarrow{\delta}_{u}(w) = \begin{cases} 1 & \text{If } u \text{ is } k-1 \text{ affix of } w \text{ pointing inward} \\ 0 & \text{Otherwise} \end{cases}$$

We note that it is impossible for a length k word w to have two k-1-affixes u pointing outwards. We identify $\operatorname{Hom}(M_k, M_{k-1})$ with $M_k^* \otimes M_{k-1}$ and define $\overleftarrow{p}_k, \overrightarrow{p}_k \in M_k^* \otimes M_{k-1}$ to be

$$\overleftarrow{p}_{k} = \sum_{u \in W_{k-1}} \overleftarrow{\delta}_{u} \otimes u, \overrightarrow{p}_{k} = \sum_{u \in W_{k-1}} \overrightarrow{\delta}_{u} \otimes u$$

Example 41. We define an orientation on W_2 by $\sigma(\overline{xx}) = xx$, $\sigma(\overline{yy}) = yy$, $\sigma(\overline{xy}) = xy$, $\sigma(\overline{yx}) = yx$, $\sigma(\overline{y^{-1}x}) = y^{-1}x$, $\sigma(\overline{xy^{-1}}) = xy^{-1}$, we look at $\overleftarrow{p}_3 : M_3 \to M_2$ in matrix form

												-						
	\overline{xxx}	yyy	\overline{xxy}	xxy^{-1}	\overline{yxx}	$y^{-1}xx$	\overline{yyx}	yyx^{-1}	\overline{xyy}	$x^{-1}yy$	\overline{yxy}	$y^{-1}xy^{-1}$	\overline{xyx}	$x^{-1}yx^{-1}$	yxy^{-1}	$y^{-1}xy$	xyx^{-1}	$x^{-1}yx$
\overline{xx}	1	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
\overline{yy}	0	1	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0
\overline{xy}	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0
\overline{yx}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	0	0
$y^{-1}x$	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	0
xy^{-1}	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0

Let $M_k = \mathrm{Im}\pi_k$.

Theorem 42. We state properties of this construction.

- 1. $\tilde{M}_k = Im\pi_k = Eq(\overleftarrow{p}_k, \overrightarrow{p}_k) (= \ker(\overleftarrow{p}_k \overrightarrow{p}_k))$
- 2. $coker(\overleftarrow{p}_{k} \overrightarrow{p}_{k}) \cong \mathbb{Z}/2\mathbb{Z}$ *i.e* $rank\tilde{M}_{k}(F_{n}) = rank ker(\overleftarrow{p}_{k} \overrightarrow{p}_{k}) = rankM_{k}(F_{n}) rankM_{k-1}(F_{n}) = n(2n-2)(2n-1)^{k-2}$
- 3. We denote $p_k = \overleftarrow{p}_k|_{\tilde{M}_k} = \overrightarrow{p}_k|_{\tilde{M}_k}$ then $p_k \pi_k = \pi_{k-1}$

Thus we get an homomorphism $\pi : \mathbb{Z}[C] \to \varprojlim \tilde{M}_k$ We begin by defining gluing.

Definition 43 (Gluing along a k-affix). Let $\overline{u}, \overline{w}_1, \overline{w}_2$ be u-words s.t. \overline{u} is a k-affix situated in opposite intrinsic orientation in \overline{w}_1 vis-à-vis \overline{w}_2 and let v be a u-word s.t. both its k-affixes are u situate in opposite intrinsic orientation. The gluing of w_1 and w_2 along u is the pushout of the square

$$\begin{array}{ccc} u &
ightarrow & w_1 \\ \downarrow & \\ w_2 \end{array}$$

in the category of labeled graphs, the gluing of v to itself along u is the coequalizer of $u \rightrightarrows v$. Notice that because \overline{u} is of opposite intrinsic orientation in \overline{w}_1 and \overline{w}_2 the result of the pushout is an u-word if \overline{u} were to be situated in the same intrinsic orientation we would have gotten a tripod. In the second case the result is a cyclic word if u had the same intrinsic orientation we would have gotten a circle with a tail.

Example 44. We continue Example 16. Recall $\pi_3 (xyx^{-1}y^{-1}) = \overline{xyx^{-1}} + \overline{yx^{-1}y^{-1}} + \overline{x^{-1}y^{-1}x} + \overline{y^{-1}xy}$. We will show that we can reconstruct the cyclic word $xyx^{-1}y^{-1}$ by gluing the components $\pi_3 (xyx^{-1}y^{-1})$ along 2-affixes. We notice that $\overline{yx^{-1}}$ is situated in opposite intrinsic orientation in $\overline{xyx^{-1}}$ vis-à-vis $\overline{yx^{-1}y}$ thus we can glue $\overline{xyx^{-1}}$ and $\overline{yx^{-1}y^{-1}}$ along $\overline{yx^{-1}}$ the result is $\overline{xyx^{-1}y^{-1}}$ (Figure 4.1). We continue and glue $\overline{xyx^{-1}y^{-1}}$ and $\overline{x^{-1}y^{-1}x}$ along $\overline{x^{-1}y^{-1}}$ and $\overline{xyx^{-1}y^{-1}x}$ by glue $\overline{xyx^{-1}y^{-1}}$ and $\overline{xyx^{-1}y^{-1}x}$ by $\overline{yx^{-1}y^{-1}x}$ with $\overline{y^{-1}xy}$ along $\overline{y^{-1}x}$ to get $\overline{xyx^{-1}y^{-1}xy}$. Finally we glue $\overline{xyx^{-1}y^{-1}xy^{-1}xy}$ to itself along \overline{xy} to get the cyclic word $xyx^{-1}y^{-1}$ (Figure 4.2).



We would like to show that any vector in ker $(\overleftarrow{p}_k - \overrightarrow{p}_k)$ can be lifted to a vector in $\mathbb{Z}[C]$. We do this by generalizing the process showed in Example 44. We define a non-negative vector to be a linear combination of basis elements such that all coefficients are non-negative. We think of a non-negative vector in M_k as a graph in the following way: A non negative vector is written as a sum of k length u-words we think of this addition as disjoint union of the graphs thus

a non-negative vector is a disjoint union of *u*-words. We have a problem that gluing two component along a k-1-affix results in a u-word that is longer than k thus it can not be represented in M_k . Therefore we define $\hat{M}_k = \bigoplus_{j \ge k} M_j$ and the dual $\hat{M}_k^* = \prod_{j \ge k} M_j^*$. We extend the homomorphism $\overleftarrow{p}_k - \overrightarrow{p}_k : M_k \rightarrow$ M_{k-1} to \hat{M}_k by extending the auxiliary functionals in Definition 40. Let u be a u-word of length k-1 and w an u-word of length greater or equal to k. We notice that if the length of w is greater or equal to 2k-1 then it is possible that both k-1 affixes of w are u pointing outward. We extend the functional $\overleftarrow{\delta}_u \in M_k^*$ to $\overleftarrow{\delta}_u \in \hat{M}_k^*$ in the following way

$$\widehat{\delta_{u}}(w) = \begin{cases} 2 & \text{If both } k-1 \text{ affixes of } w \text{ are } u \text{ pointing outward} \\ 1 & \text{If one } k-1 \text{ affix of } w \text{ is } u \text{ pointing outward} \\ 0 & \text{Otherwise} \end{cases}$$

we extend $\overrightarrow{\delta}_u$ analogously. Now we extend $\widehat{p}_k = \sum_{u \in W_{k-1}} \overleftarrow{\delta}_u \otimes u, \widehat{p}_k = \sum_{u \in W_{k-1}} \overrightarrow{\delta}_u \otimes u$ finally $\widehat{p}_k - \overrightarrow{p}_k : \widehat{M}_k \to M_{k-1}$ defined to be $\widehat{p}_k - \overrightarrow{p}_k = \widehat{p}_k - \widehat{p}_k$. Clearly $\widehat{p}_k - \overrightarrow{p}_k |_{M_k} = \widehat{p}_k - \overrightarrow{p}_k$. We define a morphism $\widehat{\pi}_k : \widehat{M}_k \to M_k$ analogous to $\pi_k : \mathbb{Z}[C] \to M_k$. Let $w \in \widehat{M}_k$ be an u-word of length greater or equal to k (these words form a basis for \widehat{M}_k) and let u be an u-word of length k - 1. We define $\widehat{\pi}_k : \widehat{M}_k \to M_k$ by $u^*(\widehat{\pi}_k(w)) = |\text{Hom}(u,w)|$ we notice that $\widehat{\pi}_k|_{M_k} = \text{Id}_{M_k}$. We show lemmas about gluing.

Lemma 45 (Gluing Lemma). Let $\overline{w}, \overline{v}$ be u-words with $l(\overline{w}), l(\overline{v}) \geq k$, let \overline{t} be an k-1-affix situated in opposite intrinsic orientation in \overline{w} vis-à-vis \overline{v} , let \overline{u} be an u-word with $l(\overline{u}) \leq k$ and let \overline{s} be the result of gluing \overline{w} and \overline{v} along \overline{t} . We show that

$$Hom\left(\overline{u},\overline{s}\right) \cong \left(Hom\left(\overline{u},\overline{w}\right) \bigsqcup Hom\left(\overline{u},\overline{v}\right)\right) / Hom\left(\overline{u},\overline{t}\right)$$

thus

$$Hom\left(\overline{u},\overline{s}\right)| = |Hom\left(\overline{u},\overline{w}\right)| + |Hom\left(\overline{u},\overline{v}\right)| - |Hom\left(\overline{u},\overline{t}\right)|$$

Proof. Because of general abstract nonsense considerations we get that there is an injective morphism

$$\left(\operatorname{Hom}\left(\overline{u},\overline{w}\right)\bigsqcup\operatorname{Hom}\left(\overline{u},\overline{v}\right)\right)/\operatorname{Hom}\left(\overline{u},\overline{t}\right) \hookrightarrow \operatorname{Hom}\left(u,s\right)$$

. We show it is surjective. Let w be an orientation of \overline{w} , t and orientation of \overline{t} and s and orientation of \overline{s} such that we can write $w = w_0 \cdot t$, $v = t \cdot v_0$ and $s = w_0 \cdot t \cdot v_0$. Assume by way of contradiction that u is a subword of s that is not a subword of either w or v. This means it contains a non trivial segments both of v_0 and of u_0 . We denote them by v_1 and u_1 and notice $l(u_1), l(v_1) \ge 1$. We can write $u = v_1 \cdot t \cdot u_1$, thus $l(u) = l(v_1 \cdot t \cdot u_1) = l(v_1) + l(t) + l(u_1) = k - 1 + l(v_1) + l(u_1) \ge k + 1$ and this is a contradiction since $l(u) \le k$.

Corollary 46. Special case: If $l(\overline{u}) = k$ we get $Hom(\overline{u}, \overline{t}) = \emptyset$. Thus

$$|Hom(\overline{u},\overline{s})| = |Hom(\overline{u},\overline{w})| + |Hom(\overline{u},\overline{v})|$$

Proposition 47. Let \overline{w} be an u-word and let \overline{t} be an u-word situated in \overline{w} twice as two k - 1-affixes in opposite intrinsic orientation, let v be the cyclic word obtained by gluing w to itself along t and let u be an u-word of length k. Then $Hom(u, w) \cong Hom(u, v)$.

Proof. The morphism from $\overline{w} \to v$ is locally injective therefor the morphism $\operatorname{Hom}(u, w) \hookrightarrow \operatorname{Hom}(u, v)$ is injective. We divide into two cases:

- 1. $\frac{l(\overline{w})}{2} > l(\overline{t})$. Let w, t orientation of $\overline{w}, \overline{t}$ and v' a cyclically reduced representative of v such that we can write $w = t \cdot w_0 \cdot t$ and $v' = t \cdot w_0$. There are l(w) k + 1 subwords of length k in w that is $l(t \cdot w_0 \cdot t) k + 1 = 2l(t) + l(w_0) k + 1 = l(t) + l(w_0)$. There are l(v) subwords of length k in v that is $l(t) + l(w_0)$. Because v and \overline{w} have the same amount of subwords this means every subword of v comes from a subword of \overline{w} thus Hom $(u, w) \cong \text{Hom}(u, v)$.
- 2. $\frac{l(\overline{w})}{2} \leq l(\overline{t})$. Let w, t orientation of $\overline{w}, \overline{t}$ such that we can write w in two ways $w = x \cdot t = t \cdot y$. This can be seen as an equation over a free semigroup. The solutions of an equation of this sort has the form $x = x_0y_0, y = y_0x_0, t = (x_0y_0)^{r-1}x_0, w = (x_0y_0)^{r-1}x_0y_0x_0$ with $r \in \mathbb{N}$, the words x_0 and y_0 reduced words s.t. $x_0 \cdot y_0$ and $y_0 \cdot x_0$ where $x_0 \neq 1$ but y_0 possibly trivial. The cyclic word resulting from gluing w to itself along t is $v = x_0y_0$. There are l(w) k + 1 subwords of length k in w that is $l(t \cdot x) k + 1 = l(t) + l(x) k + 1 = l(x)$. There are l(v) subwords of length k in v that is $l(v) = l(x_0y_0) = l(x)$. As before every subword of v comes from w thus Hom $(u, w) \cong$ Hom (u, v),

Proposition 48. Let v be a cyclic word and let the u-word u be a subword of v and let k = l(u) + 1. Then there exists an u-word w with u situated in it twice as two k - 1-affixes in opposite intrinsic orientation such that v is the result of gluing w to itself along u.

Proof. We divide this into two cases:

- 1. u is a proper subword i.e. l(v) > l(u).
- 2. *u* is not a proper subword i.e. $l(u) \ge l(v)$

If u is a proper subword than one can write $v = u \cdot v_0$ and one can check that $w = u \cdot v_0 \cdot u$. If u is not a proper subword than there exists $n \in \mathbb{N}$ a cyclically reduce representative v and a proper subword v_0 s.t. $u = v^n v_0$ then $w = v^{n+1}v_0$. Let v_1 be the proper subword s.t. $v = v_0v_1$ than $w = vv^{n-1}v_0v_1v_0$

Proposition 49. We prove propositions connecting the gluing lemmas with the homomorphisms $\overleftarrow{p}_k - \overrightarrow{p}_k, \hat{\pi}_k, \overleftarrow{p}_k - \overrightarrow{p}_k$,

1. Let $v_0, v_1 \in \hat{M}_k$ be non negative vectors such that v_1 is obtained by gluing two different components of v_0 along a common k-1-affix. Then $\hat{\pi}_k(v_0) = \hat{\pi}_k(v_1)$ and $\overleftarrow{p}_k - \overrightarrow{p}_k(v_0) = \overleftarrow{p}_k - \overrightarrow{p}_k(v_1)$. 2. Let $v \in \hat{M}_k$ be a non-negative vector and let $v_0 \in M_k$ be the vector satisfying $\hat{\pi}_k(v) = v_0$. There is a sequence of vectors $v_0, \ldots, v_n \in \hat{M}$ with $v_n = v$ such that v_{i+1} is obtained from v_i by gluing two different components of v_i along a common k - 1-affix.

3.
$$(\overleftarrow{p}_k - \overrightarrow{p}_k) \circ \hat{\pi}_k = \overleftarrow{p}_k - \overrightarrow{p}_k$$

Proof. (1) The fact that $\hat{\pi}_k(v_0) = \hat{\pi}_k(v_1)$ follows from the gluing lemma (Lemma 45). The fact that $\overleftarrow{p_k} - \overrightarrow{p_k}(v_0) = \overleftarrow{p_k} - \overrightarrow{p_k}(v_1)$ follows because we glue along k-1 affixes situated in opposite intrinsic orientation. (2) We show this for a single connected component i.e. a word of length greater or equal to k the general case follows. Let v be a u-word of length k then $v_0 = v$ and this is the sequence. Let v be a u-word of length k+n and let u_1 be k-affix and u_2 a k+n-1-affix such that u_1 is not a sub word of u_2 (i.e. these are affixes from opposite sides of v). Then u_1, u_2 have a common k-1-affix in opposite orientation s.t. gluing along it gives v. Thus by the gluing lemma (Lemma 45)

$$\hat{\pi}_k(v) = \hat{\pi}_k(u_1) + \hat{\pi}_k(u_2) = u_1 + \hat{\pi}_k(u_2)$$

Thus we write $v_{n-1} = u_1 + u_2$ and continue by induction to decompose u_2 . (3) Let $v \in \hat{M}$ be a non-negative vector, $v_0 = \hat{\pi}_k(v)$ and $v_0, v_1, \ldots, v_n = v$ a sequence as in (2). Then

$$\left(\overleftarrow{p}_{k}-\overrightarrow{p}_{k}\right)\circ\widehat{\pi}_{k}\left(v\right)=\overleftarrow{p}_{k}-\overrightarrow{p}_{k}\left(v_{0}\right)=\overleftarrow{p}_{k}-\overrightarrow{p}_{k}\left(v_{0}\right)=\overleftarrow{p}_{k}-\overrightarrow{p}_{k}\left(v_{i}\right)$$

for every *i*. Thus $(\overleftarrow{p}_k - \overrightarrow{p}_k) \circ \hat{\pi}_k (v) = \overleftarrow{p}_k - \overrightarrow{p}_k (v)$. Lastly \hat{M}_k is generated by non negative vectors therefor $(\overleftarrow{p}_k - \overrightarrow{p}_k) \circ \hat{\pi}_k = \overleftarrow{p}_k - \overrightarrow{p}_k$.

Proposition 50. We show claims about $\overleftarrow{p}_k - \overrightarrow{p}_k : M_k \to M_{k-1}$

- 1. Let $S = \left\{ v \in M_{k-1} | \sum_{u \in W_{k-1}} |u^*(v)|^2 = 2 \right\}$ then $\langle S \rangle = Im(\overleftarrow{p}_k \overrightarrow{p}_k)$.
- 2. $coker(\overleftarrow{p}_k \overrightarrow{p}_k) \cong \mathbb{Z}/2\mathbb{Z}$ (second property in Theorem 42)
- 3. Every $v \in Im(\overleftarrow{p}_k \overrightarrow{p}_k)$ has a non negative preimage.
- 4. ker $\overleftarrow{p}_k \overrightarrow{p}_k$ is generated by non-negative vectors.

Proof. (1) The set S is the set of vectors that are linear combination of exactly two length k - 1 u-words with coefficients 1 or -1. It is clear that $\langle S \rangle \supset$ Im $(\overleftarrow{p}_k - \overrightarrow{p}_k)$ because the u-words of length k generate M_k and every word has exactly two k - 1-affixes. In the other direction let $v \in S$ then there exists $\overline{u}_1, \overline{u}_2 \in W_{k-1}$ and $\epsilon_1, \epsilon_2 \in \{0, 1\}$ such that $v = (-1)^{\epsilon_1} \overline{u}_1 + (-1)^{\epsilon_2} \overline{u}_2$. Let u_1, u_2 be the orientations of $\overline{u}_1, \overline{u}_2$. Then depending on the u-word orientation there exists $\epsilon_3, \epsilon_4 \in \{-1, 1\}$ and a letter x such that there is not cancellation in $u_1^{\epsilon_3} x u_2^{\epsilon_4}$ and

$$(-1)^{\epsilon_1} \overline{u_1} + (-1)^{\epsilon_2} \overline{u_2} = \overleftarrow{p_k} - \overrightarrow{p_k} \left(\overline{u_1^{\epsilon_3} x u_2^{\epsilon_4}} \right) = (\overleftarrow{p_k} - \overrightarrow{p_k}) \circ \hat{\pi}_k \left(\overline{u_1^{\epsilon_3} x u_2^{\epsilon_4}} \right)$$

second equality is (1) in Proposition 49. (2) We define the functional $\lambda \in M_{k-1}^*$ by $\lambda \left(\sum_{u \in W_{k-1}} a_u u \right) = \sum_{u \in W_{k-1}} a_u$. We notice that $\lambda(v)$ is even for every $v \in S$. Let $u_0 \in W_{k-1}$. Because $\lambda(u_0) = 1$ we conclude $u_0 \notin \langle S \rangle$. We show that $S \cup \{u_0\}$ generates M_{k-1} . If $u \in W_{k-1}$ then $u = u_0 + (u - u_0)$ and $(u - u_0) \in S$. We also notice that $2u_0 = (u_0 - u) + (u_0 + u) \in \langle S \rangle$. Thus $\mathbb{Z}/2\mathbb{Z} \cong M_{k-1}/\operatorname{Im}(\overleftarrow{p}_k - \overrightarrow{p}_k)$. (3) Let $v \in \operatorname{Im}(\overleftarrow{p}_k - \overrightarrow{p}_k)$ then by (1) there exists $u_1, \ldots, u_l \in S$ such that $v = \sum c_i u_i$. We notice that S = -S thus we can choose c_i to be positive. We notice that for every $\epsilon_3, \epsilon_4, u_1, u_2$ the vector $\hat{\pi}_k \left(\overrightarrow{u_1^{\epsilon_3} x u_2^{\epsilon_4}} \right)$ is non-negative thus every element in S has a non negative preimage so there exists non-negative vectors $v_1, \ldots, v_l \in M_k$ such that $(\overleftarrow{p}_k - \overrightarrow{p}_k)(v_i) = u_i$. Thus $(\overleftarrow{p}_k - \overrightarrow{p}_k) (\sum c_i v_i) = v$ but $\sum c_i v_i$ is a non-negative combination of non-negative vectors thus it is non-negative. (4) Let $w \in \ker \overleftarrow{p}_k - \overrightarrow{p}_k$ then we can write w = w' - w'' such that w' and w'' are both non negative. If $\overleftarrow{p}_k - \overrightarrow{p}_k (w') = \overleftarrow{p}_k - \overrightarrow{p}_k (w') = 0$ then w is a combination of non negative vectors. Assume $\overleftarrow{p}_k - \overrightarrow{p}_k (-w') = \overleftarrow{p}_k - \overrightarrow{p}_k (-w'') = \overleftarrow{p}_k - \overrightarrow{p}_k (u)$. We write v = (v' + u) - (u + v'') and notice that v' + u, v'' + u are non-negative and $v' + u, v'' + u \in \ker \overleftarrow{p}_k - \overrightarrow{p}_k (u)$.

Proposition 51. We conclude by proving the two remaining statements from Theorem 42

- 1. $Im\pi_k = \tilde{M}_k (F_n) = \ker \overleftarrow{p}_k \overrightarrow{p}_k$
- 2. We denote $p_k = \overleftarrow{p}_k|_{\tilde{M}_k} = \overrightarrow{p}_k|_{\tilde{M}_k}$ we show that $p_k : \tilde{M}_k \to \tilde{M}_{k-1}$ is an inverse system such that $p_k \circ \pi_k = \pi_{k-1}$

Proof. (1) Let w be a cyclic word. Choose an arbitrary subword u of w of length k-1. By Proposition 48 there is an u-word v s.t. if we glue v to itself along u we get w. Then by Proposition 47 we have $\pi_k(w) = \hat{\pi}_k(v)$. Also clearly $\widehat{p}_k - \overrightarrow{p}_k(v) = \overline{u} - \overline{u} = 0$. Thus

$$\left(\overleftarrow{p}_{k}-\overrightarrow{p}_{k}\right)\circ\pi_{k}\left(w\right)=\left(\overleftarrow{p}_{k}-\overrightarrow{p}_{k}\right)\circ\hat{\pi}_{k}\left(v\right)=\overleftarrow{p}_{k}-\overrightarrow{p}_{k}\left(v\right)=0$$

i.e. $\operatorname{Im} \pi_k \subset \ker \overleftarrow{p}_k - \overrightarrow{p}_k$. On the other hand let $v_0 \in \ker \overleftarrow{p}_k - \overrightarrow{p}_k$ be a nonnegative vector. By (4) in Proposition 50 it is enough to show that $v_0 \in \text{Im}\pi_k$. If all the components of v are u-words that have a k-1-affix situated in them twice in opposite intrinsic orientation than by Proposition 47 we can lift v_0 to $v \in \mathbb{Z}[C]$ s.t. $\pi_k(v) = v_0$ by gluing each component to itself along its k-1-affix. If not let w be a component of v that does not have the above property and let u be a k-1affix of w. Because $\overleftarrow{p}_k - \overrightarrow{p}_k(v) = 0$ there exists another component of v with u situated in it in intrinsic orientation opposite to w we denote it by t. Let v_1 be the vector in M_k obtained by gluing w and t along u. By (1) in Proposition 49 $\hat{\pi}_k(v_1) = v_0$ and $\overleftarrow{p}_k - \overrightarrow{p}_k(v_1) = 0$. Now we can recursively apply the previous step until we get v_n s.t. all its components are u-words that have the same k-1-affix situated in them twice in opposite intrinsic orientation. The vector v_n can be lifted to a vector $v \in \mathbb{Z}[C]$ s.t. $\pi_k(v) = \hat{\pi}_k(v_n) = v_0$ by gluing and Proposition 47. (2) We show that $\hat{\pi}_{k-1}|_{M_k} = \overleftarrow{p}_k + \overrightarrow{p}_k$. Let v be a length k u-word then it has exactly two k-1-subwords moreover these are its k-1-affixes therefore $\hat{\pi}_{k-1}(v) = (\overleftarrow{p}_k + \overrightarrow{p}_k)(v)$. We show that $\hat{\pi}_{k-1} \circ \pi_k = 2\pi_{k-1}$. If v is a cyclic word of length 1 then $\pi_k(v) = \overrightarrow{v^k}, \hat{\pi}_{k-1}\left(\overrightarrow{v^k}\right) = 2\overrightarrow{v^{k-1}} = 2\pi_{k-1}(v)$. Let vbe a cyclic word of length greater then 1. Let u be a subword of length k-1 with

a graph morphism $u \to v$ then there are two different length k subwords w_1, w_2 such that $u \to w_1 \to v, u \to w_2 \to v$. We get w_1 by adding the subsequent letter to u in v on one side of u and get w_2 by adding the subsequent letter to u in v on the other side of u. This shows that for every $u \to v$ there are two different morphisms $u \to_1 \pi_k(v)$ and $u \to_2 \pi_k(v)$. But every u that is a subword of $\pi_k(v)$ is also a subword of v. This shows that $\hat{\pi}_{k-1} \circ \pi_k(v) = 2\pi_{k-1}(v)$. We notice that $\hat{\pi}_{k-1}|_{\tilde{M}_k} = p_{k-1}|_{\tilde{M}_k} + p_{k-1}|_{\tilde{M}_k} = 2p_k$ and because $\tilde{M}_k = \mathrm{Im}\pi_k$ we get

$$2\pi_{k-1} = \hat{\pi}_{k-1} \circ \pi_k = \hat{\pi}_{k-1}|_{\tilde{M}_k} \circ \pi_k = 2p_k \circ \pi_k$$

thus $p_k \circ \pi_k = \pi_{k-1}$.

In conclusion we have a homomorphism $\pi : \mathbb{Z}[C] \to \lim \tilde{M}_k$.

4.2 Properties of π

We now try to understand the kernel of π .

Proposition 52. Let w be a cyclically reduced word. We look at w and w^{l} as cyclic words then $l\pi_{k}(w) = \pi_{k}(w^{l})$ for every k. Thus

$$\left< lw - w^l | w \text{ cyclicaly reduced}, l \in \mathbb{N} \right> < \ker \pi$$

Proof. We notice that the morphism $w^l \to w$ is an *l*-cover space of the cyclic word w we also notice that a u-word is simply connected. Let u be a u-word any morphism $u \to w$ has exactly l different lifts to w^l thus

$$\left|\operatorname{Hom}\left(u, w^{l}\right)\right| = l \left|\operatorname{Hom}\left(u, w\right)\right|$$

. Thus we get $l\pi_k(w) = \pi_k(w^l)$ for every k and l.

We would like to show that $\ker \pi = \langle lw - w^l | w$ cyclically reduced, $l \in \mathbb{N} \rangle$ for this we need a lemma with a somewhat technical proof.

Lemma 53. Let w be a cyclically reduced word. The word w is a nontrivial power if and only if for every cyclic permutation w' there are words t and v such that $w' = t \cdot v \cdot t$ where t and v are proper subwords. $t \neq 1$ but v may be trivial.

Proof. One direction is easy: If $w = u^k$ with $k \ge 2$ it can be decomposed as $uu^{k-2}u$. A cyclic permutation of a power is also a power and thus it satisfies the condition. Suppose w satisfies the condition. We say a decomposition $t \cdot v \cdot t$ of a cyclic permutation of w is reducible if t has the form $t = s \cdot t_0 \cdot s$ where s and t_0 are proper subwords ($s \ne 1$, t_0 may be trivial) and irreducible otherwise. Let $t \cdot v \cdot t$ be an irreducible decomposition of a cyclic permutation of w such that l(t) is maximal among all irreducible decompositions of cyclic permutations of w. We want to show that w has a cyclic permutation that is equal to t^k for some k. If v = 1 then w is a power, therefore assume $v \ne 1$. Let us examine the cyclic permutation $v \cdot t \cdot t$. Because w satisfies the condition in the lemma $v \cdot t \cdot t$ has an irreducible decomposition $s \cdot u \cdot s$. Because t is maximal this means that s is a proper suffix of t or s = t. We want to show that s = t.

Remark. If we show that t has proper suffix a that is also a proper prefix we will reach a contradiction. For $l(a) < \frac{l(t)}{2}$ this is clear because we can write $t = a \cdot t_0 \cdot a$ i.e. $t \cdot v \cdot t$ is reducible. For $l(a) \ge \frac{l(t)}{2}$. We write $t = t_1 \cdot a = a \cdot t_2$ this the same kind of equation over a free semigroup as in Proposition 47. As before we have a family of solutions $t_1 = xy, t_2 = yx, a = (xy)^{r-1}x$ with x, y reduced words such that $x \cdot y$ and $y \cdot x$ and $r \in \mathbb{N}$ with $x \neq 1$ and y possibly trivial. Thus $t = at_2 = (xy)^{r-1}xyx = x(y(xy)^{r-1})x$ which is a contradiction to tvt being an irreducible decomposition.

Assume l(s) < l(t). Thus s is a suffix of t and we write $t = t_1 \cdot s$. We divide into three different cases

- 1. l(s) > l(v): We write $s = s_1 \cdot s_2$ such that $s_1 = v$ and $t = s_2 \cdot t_2$. Thus $s_2 \cdot t_2 = t = t_1 \cdot s = t_1 s_1 \cdot s_2$. We see that s_2 is a both proper prefix and a proper suffix of t this is a contradiction.
- 2. l(s) = l(v): Then s = v and we consider the cyclic permutation $t \cdot t \cdot s$. It has an irreducible decomposition $s_1 \cdot u_1 \cdot s_1$. We notice $l(s_1) \leq l(t)$ because t is maximal. We write $t = s_1 \cdot t_2$ ($s_1 \neq 1$ but t_2 is possibly trivial)
 - (a) If $l(s_1) \le l(s) < l(t)$ then s_1 is a prefix of t but it is also a suffix of s (or equal to s) thus a suffix of t this is a contradiction.
 - (b) If $l(s_1) > l(s)$ then we write $s_1 = s_{11} \cdot s_{12}$, $t = t_3 \cdot s_{11}$, $s_{12} = s$. We get $t_3 \cdot s_{11} = t = s_1 \cdot t_2 = s_{11} \cdot s_{12} \cdot t_2$. Thus s_{11} is both a proper prefix and a proper suffix of t this is a contradiction. (This still holds of $t_2 = 1$)
- 3. l(s) < l(v): We write $v = s \cdot v_1$ and $v \cdot t \cdot t = s \cdot v_1 \cdot t \cdot t$. We denote $s = \hat{s}_1$ and proceed by induction.

We examine the permutation $v_k \cdot t \cdot t \cdot \hat{s}_k$. The subword \hat{s}_k satisfies the following properties: (1) It is a concatenation of suffixes of t. (2) If r is a suffix of \hat{s}_k or \hat{s}_k itself and it satisfies l(r) > l(s) then it is a concatenation of two or more suffixes of t. We notice that $\hat{s}_1 = s$ satisfies properties (1) and (2). If u satisfies properties (1) and (2) then any suffix of u also satisfies these properties. If u_1 and u_2 both satisfy properties (1) and (2) then there concatenation $u_1 \cdot u_2$ satisfies the properties as well. The cyclic permutation $v_k \cdot t \cdot t \cdot \hat{s}_k$ has an irreducible decomposition $s_k \cdot u_k \cdot s_k$. Because t is maximal $l(t \cdot \hat{s}_k) > l(s_k)$ this means s_k is suffix of $t \cdot \hat{s}_k$. We notice that s_k satisfies properties (1) and (2). We divide into three cases:

- 1. $l(s_k) > l(v_k)$: Because t is maximal then $l(s_k) < l(v_k \cdot t)$ thus there is suffix of s_k that is a prefix of t denote it by s_{k1} . The subword s_{k1} satisfies properties (1) and (2). Thus s_{k1} has a prefix that is a suffix of t or it is itself a suffix of t. Either way t has a prefix that is also its suffix and this is a contradiction.
- 2. If $l(s_k) = l(v_k)$ then $v_k \cdot t \cdot t \cdot \hat{s}_k = s_k \cdot t \cdot t \cdot \hat{s}_k$ then there is a cyclic permutation $t \cdot t \cdot \hat{s}_k \cdot s_k$. We notice $\hat{s}_k s_k$ satisfies properties (1) and (2). We write $\hat{s}_k s_k = \hat{s}'_k$ and examine $tt\hat{s}'_k$. The cyclic permutation $tt\hat{s}'_k$ has a irreducible decomposition $s'_k \cdot u'_k \cdot s'_k$ we notice that $l(s'_k) < l(t \cdot \hat{s}'_k)$ thus s'_k is a suffix of $t \cdot \hat{s}'_k$ so it satisfies properties (1) and (2). There are two options

- (a) The subword s'_k is not a concatenation of two or more suffixes of t. Because of property (2), it is a suffix of t satisfying $l(s'_k) \leq l(s)$ and by assumption $l(s'_k) \leq l(s) < l(t)$. Thus s'_k is both a prefix and a suffix of t and this is a contradiction.
- (b) Otherwise observe that $l(s'_k) \leq l(t)$. Thus a prefix of s'_k is also a prefix of t. Because s'_k is a concatenation of suffixes of t it has a prefix that is a suffix of t and this is a contradiction.
- 3. If $l(s_k) < l(v_k)$ then we write $v_k = s_k v_{k+1}$ and $v_k \cdot t \cdot \hat{s}_k = s_k \cdot v_{k+1} \cdot t \cdot t \cdot \hat{s}_k$. We examine the cyclic permutation $v_{k+1} \cdot t \cdot t \cdot \hat{s}_k \cdot s_k$. We notice that $\hat{s}_k \cdot s_k$ satisfies properties (1) and (2) and we write $\hat{s}_{k+1} = \hat{s}_k \cdot s_k$. We notice that $l(v_{k+1}) < l(v_k)$ thus case 3 cannot repeat indefinitely and ultimately we reach a contradiction.

We close the argument with another induction. We had a cyclic permutation $v \cdot t \cdot t$ that has a decomposition $s \cdot u \cdot s$ and we concluded that s = t. Because t is irreducible then $l(v) \geq l(t)$. If v = t we are done if not we write $v = tv_1$ and we observe the cyclic permutation v_1t^1tt . This is the base for induction. Let v_kt^ktt be a cyclic permutation. Then writing $v'_k = v_kt^k$ we notice that v'_ktt satisfies the same conditions as vtt thus there is a decomposition sus such that s = t. Because t irreducible then $l(v_k) \geq l(t)$. If $v_k = t$ we are done if not we write $v_k = tv_{k+1}$ and we examine the cyclic permutation $v_{k+1}t^{k+1}tt$. We notice that $l(v_{k+1}) < l(v_k)$ thus this process must end at some point i.e. for some k we have $v_k = t$ thus w has a cyclic permutation that is a power of t.

Proposition 54. Let w_1, \ldots, w_m be cyclic words and let $\sum_{i=1}^m a_i w_i$ a linear combination that is not in $\langle lw - w^l | w$ cyclically reduced, $l \in \mathbb{N} \rangle$. Then there exists a k s.t. $\pi_k \left(\sum_{i=1}^m a_i w_i \right) \neq 0$. This shows that

$$\ker \pi = \langle lw - w^l | w \text{ cyclicaly reduced}, l \in \mathbb{N} \rangle$$

Proof. Using the relations $lw - w^l$ the linear combination $\sum_{i=1}^m a_i w_i$ always has representative $\sum_{i=1}^{\hat{m}} \hat{a}_i \hat{w}_i$ such that $w_1, \ldots, w_{\hat{m}}$ are pairwise distinct words that are not powers. Therefore we assume this for $\sum_{l=1}^m a_l w_l$. Without loss of generality assume w_m is of maximal length. We notice that the length ksubwords of a length k cyclic word are its cyclically reduced representative. If two length k cyclic words have a length k subword in common they are the same cyclic word, therefore we can assume that $l(w_m) > l(w_i)$ for $1 \le i \le m-1$. We notice that if u is a subword of a cyclic word w with l(u) > l(w) then there is a cyclically reduced representative w' of w such that $u = w'^r w_1$ where w_1 is a proper prefix of w' and $r \in \mathbb{N}$ (w_1 is possibly trivial but not when r = 1). Thus u has the form $t \cdot v \cdot t$. Let $l(w_m) = k$. Since w_m is not a power by Lemma 53 it has a subword of length k that does not have the form $t \cdot v \cdot t$. Assume that $\pi_k (\sum_{i=1}^m a_i w_i) = 0$ then $-a_m \pi_k (w_m) = \sum_{i=1}^{m-1} a_i \pi_k (w_i)$ but this is a contradiction because the sum on the left side has a u-word that does not have the form tvt while all the u-word summands on the right side have the form tvt since $k = l(w_m) > l(w_i)$ for $1 \le i \le m-1$.

This proposition is the main component of Theorem 3. We take a moment to talk about the topology of an inverse limit of free modules. **Definition 55.** Let $p_i : M_i \to M_{i-1}$ be an inverse system of finitely generated free \mathbb{Z} -module (vector spaces). We define a valuation V on $\lim_{i \to \infty} M_i$. Let $v = (v_i)_{i \in \mathbb{N}} \in \lim_{i \to \infty} M_i$ i.e. $v_i = p_{i+1} (v_{i+1})$ than we define $V(v) = \max \{k \in \mathbb{N} | v_k = 0\}$. Let $\hat{p}_i : \lim_{i \to \infty} M_i \to M_i$ the natural projection and note that $\hat{p}_i = p_{i+1} \circ \hat{p}_{i+1}$.

Three equivalent ways to define the topology of $\lim M_i$

- 1. We take the discrete topology on each M_i and define the topology of $\varprojlim M_i \leq \prod_i M_i$ to be the induced topology from the product topology (general topology)
- 2. We take topology induced by taking $\ker \hat{p}_{i+1} \leq \ker \hat{p}_i$ to be local base of open neighborhoods of 0. (topological module)
- 3. Let $v \in \varprojlim M_i$ and $n \in \mathbb{N}$ we define $B_{v,n} = \left\{ w \in \varprojlim M_i | V(w-v) \ge n \right\}$. The sets $B_{v,n}$ are a base for the topology. (metric topology)

Let us return to the morphism $\pi : \mathbb{Z}[C] \to \varprojlim \tilde{M}_i$. For every *i* the morphism π_i is onto \tilde{M}_i thus the image of π is dense in $\varprojlim \tilde{M}_i$. The module $\mathbb{Z}[C] / \langle lw - w^l | w$ cyclically reduced, $l \in \mathbb{N} \rangle$ can be seen as the free module generated by cyclic words that are not non-trivial powers. The usual way to define a metric would be $\|\| : \varprojlim \tilde{M}_i \to \mathbb{R}$ by $\|v\| = 2^{-V(v)}$ In this case $\varprojlim \tilde{M}_i$ is the completion of Im π with respect to the metric.

4.3 Representation

We now show how Theorem 1 gives us a representation. Let φ be an automorphism. We recall the definition of m_k^{φ} from the introduction. Let $u \in W_k$, let $S_{\varphi,u}$ be the multiset of words given by Theorem 1, let w a cyclic word and $k_v = l(v)$ for $v \in S_{\varphi,u}$ then by definition of π_k and Theorem 1 we have

$$u^{*}\left(\pi_{k}\left(\varphi\left(w\right)\right)\right) = \left|\operatorname{Hom}\left(u,\varphi\left(w\right)\right)\right| = \sum_{v\in S_{\varphi,u}}\left|\operatorname{Hom}\left(v,w\right)\right| = \sum_{v\in S_{\varphi,u}}v^{*}\left(\pi_{k_{v}}\left(w\right)\right)$$

Because the equation is true for every w we have an equation of functions $u^* \circ \pi_k \circ \varphi = \sum v^* \circ \pi_{k_v}$ equivalently we can write

$$\varphi^{*} \circ \pi_{k}^{*}\left(u^{*}\right) = \sum_{v \in S_{\varphi,u}} \pi_{k_{v}}^{*}\left(v^{*}\right)$$

Where we consider φ as the linear extension of φ to $\mathbb{Z}[C]$. We denote

$$p_{k,l} = p_{l+1} \circ \cdots \circ p_{k-1} \circ p_k$$

thus $p_{k,l} \circ \pi_k = \pi_l$. We recall

$$\overline{p}_{m_k,k_v} \circ \pi_{m_k} = p_{m_k,k_v} \circ \pi_{m_k} = \pi_{k_v}$$

so $\pi_{k_v}^* = \pi_{m_k}^* \circ \overleftarrow{p}_{m_k,k_v}^*$ thus

$$\varphi^* \circ \pi_k^* \left(u^* \right) = \sum_{v \in S_{\varphi, u}} \pi_{k_v}^* \left(v^* \right) = \pi_{m_k}^* \left(\sum_{v \in S_{\varphi, u}} \overleftarrow{p}_{m_k, k_v} \left(v^* \right) \right)$$

We define $\varphi_k^*: M_k^* \to M_{m_k}^*$ by

$$\varphi_{k}^{*}\left(u^{*}\right)=\sum_{v\in S_{\varphi,u}}\overleftarrow{p}_{m_{k},k_{v}}^{*}\left(v^{*}\right)$$

it satisfies $\varphi^* \circ \pi_k^* = \pi_{m_k}^* \circ \varphi_k^*$. The homomorphism φ_k^* defines a homomorphism $\varphi_k : M_{m_k} \to M_k$ that satisfies

$$\varphi_k \circ \pi_{m_k} = \pi_k \circ \varphi$$

We notice that we use $\overleftarrow{p}_{m_k,k_v}^*$ in the definition and we recall that $\overleftarrow{p}_{m_k,k_v}^*$ is defined using an arbitrary u-word orientation. Thus a morphism φ_k that satisfies $\varphi_k \circ \pi_{m_k} = \pi_k \circ \varphi$ is not unique. A different orientation would result in a different morphism φ_k . But if we consider the restriction $\varphi_k|_{\tilde{M}_{m_k}} : \tilde{M}_{m_k} \to \tilde{M}_k$ then it is unique. Let φ_k and φ'_k both satisfy

$$\varphi_k \circ \pi_{m_k} = \pi_k \circ \varphi = \varphi'_k \circ \pi_{m_k}$$

but π_{m_k} is onto on \tilde{M}_{m_k} so $\varphi'_k|_{\tilde{M}_{m_k}} = \varphi_k|_{\tilde{M}_{m_k}}$. We notice that

$$p_k \circ \varphi_k \circ \pi_{m_k} = p_k \circ \pi_k \circ \varphi$$
$$= \pi_{k-1} \circ \varphi$$
$$= \varphi_{k-1} \circ \pi_{m_{k-1}}$$
$$= \varphi_{k-1} \circ p_{m_k, m_{k-1}} \circ \pi_{m_k}$$

But π_{m_k} is onto on M_{m_k} so

$$p_k \circ \varphi_k = \varphi_{k-1} \circ p_{m_k, m_{k-1}} \tag{4.1}$$

This shows that the sequence $\varphi_k : \tilde{M}_{m_k} \to \tilde{M}_k$ defines a homomorphism $\lim \tilde{M}_k \to \lim \tilde{M}_k$. We notice that

$$\begin{aligned} (\varphi \circ \psi)_k \circ p_{m_{m_k}^{\psi}, m_k^{\varphi \circ \psi}} \circ \pi_{m_{m_k}^{\psi}} &= (\varphi \circ \psi)_k \circ \pi_{m_k^{\varphi \circ \psi}} \\ &= \pi_k \circ (\varphi \circ \psi) \\ &= \varphi_k \circ \pi_{m_k^{\psi}} \circ \psi \\ &= \varphi_k \circ \psi_{m_k^{\varphi}} \circ \pi_{m_{m_k}^{\psi}} \end{aligned}$$

as before we get

$$(\varphi \circ \psi)_k \circ p_{m_k^{\psi}, m_k^{\varphi \circ \psi}} = \varphi_k \circ \psi_{m_k^{\varphi}}$$

$$(4.2)$$

We notice that the sequences $(\varphi \circ \psi)_k \circ p_{m_{\tilde{k}}^{\psi}, m_k^{\varphi \circ \psi}}$ and $(\varphi \circ \psi)_k$ define the same homomorphism in the limit. Thus we get that $\operatorname{Out}(F_n) \to \operatorname{Aut}\left(\varprojlim \tilde{M}_k\right)$ is a homomorphism. Lastly let φ, ψ be automorphism from different conjugacy classes then there exists a generator of the free group x such that $\varphi(x), \psi(x)$ are from different conjugacy classes. because φ, ψ are automorphisms $\psi(x), \varphi(x)$ are not powers thus by Proposition 54 there is a k such that $\pi_k \circ \varphi(x) \neq \pi_k \circ \psi(x)$ without loss of generality assume $m_k^{\varphi} \leq m_k^{\psi}$ thus

$$\varphi_{k} \circ p_{m_{k}^{\psi},m_{k}^{\varphi}}\left(\pi_{m_{k}^{\psi}}\left(x\right)\right) \hspace{2mm} \neq \hspace{2mm} \psi_{k}\left(\pi_{m_{k}^{\psi}}\left(x\right)\right)$$

The sequences $\varphi_k \circ p_{m_k^{\psi}, m_k^{\varphi}}$ and φ_k define the same homomorphism in the limit. We conclude that $\operatorname{Out}(F_n) \to \operatorname{Aut}\left(\varprojlim \tilde{M}_k\right)$ is injective (Theorem 3). We note that M_k and M_{m_k} have a given basis thus $\varphi_k : M_{m_k} \to M_k$ is a matrix and φ is represented by a tower of finite matrices. Equation 4.1 shows that φ_{k-1} can be reconstructed given φ_k thus each successive matrix gives more information about φ . Equation 4.2 shows that computations in $\operatorname{Aut}\left(\varprojlim \tilde{M}_k\right)$ can be done via matrix multiplication.

References

- Daryl Cooper. Automorphisms of free groups have finitely generated fixed point sets. J. Algebra, 111(2):453-456, 1987.
- [2] P. J. Higgins and R. C. Lyndon. Equivalence of elements under automorphisms of a free group. J. London Math. Soc. (2), 8:254–258, 1974.
- [3] Noam M. D. Kolodner. On algebraic extensions and decomposition of homomorphisms of free groups. J. Algebra, 569:595–615, 2021.
- [4] Roger C. Lyndon and Paul E. Schupp. Combinatorial group theory. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1977 edition.
- [5] Elvira Strasser Rapaport. On free groups and their automorphisms. Acta Math., 99:139–163, 1958.
- [6] Jean-Pierre Serre. Arbres, amalgames, SL₂. Société Mathématique de France, Paris, 1977. Avec un sommaire anglais, Rédigé avec la collaboration de Hyman Bass, Astérisque, No. 46.
- [7] John R. Stallings. Topology of finite graphs. Invent. Math., 71(3):551-565, 1983.
- [8] J. H. C. Whitehead. On equivalent sets of elements in a free group. Ann. of Math. (2), 37(4):782–800, 1936.