

Determination of Majorana type-phases from the time evolution of lepton numbers

Nicholas J. Benoit^{1*}, Yuta Kawamura^{2†}, Saki Kawano^{3‡},
Takuya Morozumi^{4,5§}, Yusuke Shimizu^{4,5¶}, Kei Yamamoto^{6||}

¹*Graduate School of Science, Hiroshima University, Higashi-Hiroshima 739-8526, Japan*

²*Yokkaichi city, Mie, Japan*

³*Tokyo prefecture, Japan*

⁴*Physics Program, Graduate School of Advanced Science and Engineering,
Hiroshima University, Higashi-Hiroshima 739-8526, Japan*

⁵*Core of Research for the Energetic Universe, Hiroshima University,
Higashi-Hiroshima 739-8526, Japan*

⁶*Department of Global Environment Studies, Hiroshima Institute of Technology,
Saeki-ku, Hiroshima 731-5193, Japan*

Abstract

We have investigated an approach to determine the Majorana type-phases using the time evolution of lepton family numbers. The Majorana type-phases are related to the orientation of unitarity triangles for the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix, and the Majorana phases α_{21} and α_{31} . After taking the second-order time derivative of the lepton family number expectation values, the dependencies on the summation of Majorana type-phases can be determined. Thus allowing for the extraction of the orientation of the unitarity triangles and the Majorana phases. We study how to extract the Majorana type-phases and the lightest neutrino mass for three massive neutrinos, and when a neutrino is massless, i.e., $m_{1,3} = 0$. Our result can be complimentary to using neutrinoless double-beta decay for determining the orientation of PMNS unitarity triangles and the Majorana phases.

*E-mail address: njbenoit@hiroshima-u.ac.jp

†E-mail address: kawamura1994phy@gmail.com

‡E-mail address: sakikwn@gmail.com

§E-mail address: morozumi@hiroshima-u.ac.jp

¶E-mail address: yu-shimizu@hiroshima-u.ac.jp

||E-mail address: k.yamamoto.e3@cc.it-hiroshima.ac.jp

1 Introduction

Despite phenomenal experimental programs in neutrino physics, open questions remain. We know the three active neutrinos of the Standard Model come in three flavors (families) electron neutrino, muon neutrino, and the tauon neutrino[1]. Those flavors are assigned based on the neutrinos weak charged current lepton partner. We also know that neutrinos can oscillate between flavors over macroscopic distances[2, 3]. To describe the oscillations the three neutrino flavors are treated as linear combinations of massive eigenstates, then to explain experimental data at least two of the eigenstates must have small, but non-zero, masses[4, 5]. This is in contrast to the Standard Model, in which the three neutrinos are massless particles. Because neutrino flavor oscillation experiments can only constrain the mass squared differences of the neutrinos $\Delta m_{ij}^2 = m_i^2 - m_j^2$, details of neutrino masses remain unknown. For example neutrinos are neutrally charged particles with non-zero mass, which means they could be their own antiparticles. Under that setup neutrinos would be called Majorana particles and have a Majorana mass[6, 7, 8]. A Dirac particle type and mass are also possible for neutrinos, this would follow the usual Standard Model lepton content. Neither particle type has not been experimentally established for neutrinos despite years of searching for Majorana neutrinos. Perhaps the most famous experiments are searching for neutrinoless double-beta decay[9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23], but there are also neutrinoless quadruple-beta decay[24], neutrinoless double-electron capture[25, 26, 27, 28, 29, 30], proposed processes mediated by the exchange of a pair of virtual neutrinos akin to a Casimir force[31, 32, 33], coherent scattering of neutrinos on nuclei with bremsstrahlung[34], and proposed searches of quantum statics for final state decay processes involving neutrinos[35, 36, 37, 38].

We explore the situation that neutrinos are Majorana particles. In section 2, we summarize how the three neutrino flavors are constructed as linear combinations of the massive eigenstates via a mixing matrix; and we introduce how the mixing matrix can be formed with Majorana type-phases and unitarity triangles. Then, we connect our previous work on the time evolution of lepton family numbers to the Majorana type-phases in section 3. We explain how our previous work can be used to determine the Majorana type-phases and the lightest neutrino mass in section 4. In the last section, section 5, we consider the lightest neutrino to be massless and illustrate how this enhances the predictive power of our work. Finally, we provide some concluding thoughts.

2 Majorana type-phases and unitarity triangles

Neutrino masses are a pillar for physics beyond the Standard Model that started with the discovery of neutrino flavor oscillations [39, 40]. The results of neutrino oscillation experiments are mathematically described by a mixing of basis states via the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix [41, 42, 43]

$$\nu_{L\alpha} = U_{\alpha i} \nu_{Li} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu1} & U_{\mu2} & U_{\mu3} \\ U_{\tau1} & U_{\tau2} & U_{\tau3} \end{pmatrix} \begin{pmatrix} \nu_{L1} \\ \nu_{L2} \\ \nu_{L3} \end{pmatrix}, \quad (1)$$

where $\alpha = e, \mu, \tau$ and $i = 1, 2, 3$. The PMNS matrix $U_{\alpha i}$ can be formed for neutrinos with a Majorana mass through diagonalization of the Majorana mass matrix,

$$(U)_{\alpha i} (m_\nu)_{\alpha\beta} (U)_{\beta j} = (m_\nu)_i \delta_{ij}, \quad (2)$$

using a Takagi Factorization [44]. Then the leptonic weak charged current interaction is written as,

$$\mathcal{L}_I^{(CC)} = \frac{g}{\sqrt{2}} \bar{l}_{L\alpha} \gamma_\mu U_{\alpha i} \nu_{Li} W^{\mu-} + \text{h.c.} \quad (3)$$

The charged leptons are free to be re-phased, because their mass term is invariant under the phase transformations of $l_\alpha \rightarrow l'_\alpha = \exp(i\phi_\alpha) l_\alpha$. But, the neutrino Majorana mass term is not invariant under the phase transformations. This means the standard parametrization of the PMNS matrix [45] must be extended to include two independent CP violating phases. Thus, the unitary 3×3 PMNS matrix depends on three mixing angles and three CP violating phases.

$$U_{\alpha i} = U^D \text{diag} \left(1, e^{i\frac{\alpha_{21}}{2}}, e^{i\frac{\alpha_{31}}{2}} \right), \quad (4)$$

where the Dirac portion U^D is of the form,

$$U^D = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix}. \quad (5)$$

We use the usual notation of $c_{ij} = \cos \theta_{ij}$ and $s_{ij} = \sin \theta_{ij}$ from PDG[45]. The CP violating phases in the diagonal matrix, α_{21} and α_{31} , are the Majorana phases, whereas the CP violating phase of U^D is the Dirac phase δ . The CP violating phases have physical meaning only if all the mixing angles are not 0 or $\pi/2$. This can be seen by constructing the invariants [46, 47, 48, 49] of the PMNS matrix.

We consider an interesting aspect of Majorana neutrinos and the rephasing invariant bilinears [48] that Branco and Rebelo name ‘‘Majorana type-phases’’[50]. The Majorana type-phases are defined by Branco and Rebelo to be the argument of the bilinears $U_{\alpha k} U_{\alpha j}^*$ with no summation over the repeated index α . Then, Branco and Rebelo choose six Majorana type-phases and prove the six phases can be used to exactly reproduce the 3×3 PMNS matrix Eq.(4). We reproduce their [50] six Majorana type-phases for self-containment of our work,

$$\beta_1 = \arg U_{e1} U_{e2}^*, \quad \beta_2 = \arg U_{\mu 1} U_{\mu 2}^*, \quad \beta_3 = \arg U_{\tau 1} U_{\tau 2}^*, \quad (6)$$

$$\gamma_1 = \arg U_{e1} U_{e3}^*, \quad \gamma_2 = \arg U_{\mu 1} U_{\mu 3}^*, \quad \gamma_3 = \arg U_{\tau 1} U_{\tau 3}^*. \quad (7)$$

To reproduce the 3×3 PMNS matrix of Eq.(4), from the Majorana type-phases Eq.(6) and Eq.(7), we must consider the construction of unitarity triangles. Two types of triangles can be created from the PMNS matrix, Dirac and Majorana triangles. We are interested in Majorana triangles and their dependence on the Majorana type-phases. The three Majorana triangles are derived by multiplying the columns of the PMNS matrix,

$$U_{e1} U_{e2}^* + U_{\mu 1} U_{\mu 2}^* + U_{\tau 1} U_{\tau 2}^* = 0, \quad \text{triangle 1,} \quad (8)$$

$$U_{e1} U_{e3}^* + U_{\mu 1} U_{\mu 3}^* + U_{\tau 1} U_{\tau 3}^* = 0, \quad \text{triangle 2,} \quad (9)$$

$$U_{e2}U_{e3}^* + U_{\mu2}U_{\mu3}^* + U_{\tau2}U_{\tau3}^* = 0, \quad \text{triangle 3.} \quad (10)$$

The equation for triangle 1, Eq.(8) is connected with the β_k Majorana type-phases of Eq.(6). We illustrate that connection between the triangle vectors of Eq.(8) and β_k in with figure 1. From figure 1, we can see all sides of the triangle are constructed from the rephasing invariants $U_{\alpha k}U_{\alpha j}^*$. Thus, the Majorana triangles are not free to rotate in the complex plain and their orientation is physically meaningful [50, 51]. In addition, the internal angles of the triangle ζ_k




Figure 1: The first Majorana triangle depends on the Majorana type-phases β_k . The sides of the triangle are constructed from the first two rows of the PMNS matrix. The orientation is physically meaningful and can only be determined by knowledge of the Majorana type-phases.

are,

$$\zeta_1 = \pi - (\beta_3 - \beta_2), \quad \zeta_2 = (\beta_3 - \beta_1) - \pi, \quad \zeta_3 = \pi - (\beta_2 - \beta_1). \quad (11)$$

Triangle 2 is connected to the γ_k of Eq.(7) and behaves similarly to triangle 1 where the orientation is physically meaningful. The internal angles of triangle 2 are calculated by replacing β_k with γ_k in Eq.(11). Lastly, the connection to triangle 3 comes from subtraction between the β_k 's and γ_k 's and its behavior depends on the other two triangles.

Branco and Rebelo relate the mixing angles of the parametrized PMNS matrix in Eq.(5) to the Majorana type-phases using the law of sines on the Majorana triangles. For example, to find θ_{12} they take,

$$\begin{aligned} \tan^2 \theta_{12} &= \frac{|U_{e2}U_{e3}^*|^2}{|U_{e1}U_{e3}^*|^2} \\ &= \frac{|\sin(\gamma_1 - \gamma_3)||\sin(\gamma_2 - \gamma_1)||\sin(\gamma_3 - \gamma_2 - (\beta_3 - \beta_2))|}{|\sin(\gamma_1 - \gamma_3 - (\beta_1 - \beta_3))||\sin(\gamma_2 - \gamma_1 - (\beta_2 - \beta_1))||\sin(\gamma_3 - \gamma_2)|}. \end{aligned} \quad (12)$$

Then they prove how the Dirac phase δ relates to the Majorana type-phases via the common

area of the triangles,

$$\begin{aligned}
A &= \frac{1}{16} |\sin 2\theta_{12} \sin 2\theta_{13} \sin 2\theta_{23} \cos \theta_{13} \sin \delta| \\
&= \frac{1}{2} |\cos \theta_{12} \cos \theta_{13} \sin \theta_{13}|^2 \frac{|\sin(\gamma_1 - \gamma_2)| |\sin(\gamma_1 - \gamma_3)|}{|\sin(\gamma_2 - \gamma_3)|}.
\end{aligned} \tag{13}$$

We use the relations of Branco and Rebelo like Eq.(12) and Eq.(13) to update their analysis [50] of the Majorana triangles based on the recent neutrino oscillation global fit. Specifically, we use the best fit values of NuFITv5.1 calculated by the Nu-Fit collaboration [52]. Compared to the analysis of ref.[50], we include additional data from neutrino oscillation experiments. For simplicity, we consider the Majorana phases to be absent $\alpha_{21} = \alpha_{31} = 0$, and later we will include them. The vectors for Triangle 1 in fig.2 are,

$$\begin{aligned}
U_{e1}U_{e2}^* &= 0.4496 - 0.0i, \\
U_{\mu1}U_{\mu2}^* &= -0.2295 + 0.05711i, \\
U_{\tau1}U_{\tau2}^* &= -0.2200 - 0.05711i,
\end{aligned} \tag{14}$$

where we only write the first four significant figures with no rounding. We treat the vectors




Figure 2: We build the three Majorana triangle using the best fit values of NuFITv5.1[52] for normal hierarchy. The orientation of the triangles depends on the Majorana type-phases; β_k for triangle 1 that is related to the Majorana phase α_{21} , γ_k for triangle 2, that is related to the Majorana phase α_{31} , and for triangle 3 the subtraction between the Majorana type-phases, $\gamma_k - \beta_k$. For this instance, we set the Majorana phase $\alpha_{21} = \alpha_{31} = 0$, because it is experimentally not determined. NuFITv5.1 assumes the PMNS matrix is unitary, so all triangles are completely closed.

for Triangles 2 and 3 of fig.2 similarly as Triangle 1,

$$\left. \begin{aligned}
U_{e1}U_{e3}^* &= -0.07945 - 0.09469i, \\
U_{\mu1}U_{\mu3}^* &= -0.2354 + 0.04261i, \\
U_{\tau1}U_{\tau3}^* &= 0.3149 + 0.05208i;
\end{aligned} \right\} \text{Triangle 2} \tag{15}$$

$$\left. \begin{aligned} U_{e2}U_{e3}^* &= -0.05251 - 0.06258i, \\ U_{\mu2}U_{\mu3}^* &= 0.4339 + 0.02816i, \\ U_{\tau2}U_{\tau3}^* &= -0.3814 + 0.03442i. \end{aligned} \right\} \text{Triangle 3} \quad (16)$$

In all three triangles for fig.2 the vectors completely close to form a triangle. This is because the data we are using from NuFITv5.1 assume the PMNS matrix Eq.(5) is unitary. If the unitary assumption was relaxed, the triangles could be open. Furthermore, all three triangles are scalene, which is different from the isosceles triangles of Branco and Rebelo because we do not consider perturbations around tri-bimaximal mixing.

Lastly, current and near future neutrino oscillation experiments can not determine the orientation of the Majorana triangles. This is because the oscillation experiments measure the Dirac phase δ , but can not measure the Majorana phases α_{21} and α_{31} that determine the triangle orientation. For example, we can change the values of $\Delta\alpha_{21}$ to rotate triangle 1 in figure 3. Branco and Rebelo discussed this problem [50], stating that measurements




Figure 3: We rotate the first Majorana triangle from the best fit values of NuFITv5.1[52] for normal hierarchy. The orientation of the triangle depends on the Majorana type-phases β_k , thus it is rotated by varying between $0 < \alpha_{21} < \pi$.

of the Dirac phase can only provide insight to the differences of the Majorana type-phases, i.e. $\gamma_1 - \gamma_3$; so the triangle orientation can not be determined. Whereas, the addition of Majorana type-phases $\beta_i + \beta_j$ and $\gamma_i + \gamma_j$ are connected to the Majorana phases α_{21} and α_{31} , respectively. They then go on to discuss the possibility of determining the triangle orientation with neutrinoless double-beta decay and flavor sensitive leptogenesis[53, 54, 55]. In this paper, we will describe an additional method to determine the triangle orientation based on our work of lepton number oscillations[56, 57].

3 Time evolution of lepton number and Majorana type-phases

To determine the triangle orientation, we study the time evolution of lepton number focusing on the dependence of the Majorana type-phases. In the works [56, 57], the lepton family numbers L_α^M for the Majorana neutrinos are defined. Then, the time evolution of their expectation values are obtained. We rewrite the Majorana $L_\alpha^M(t)$

$$\begin{aligned}
& \langle \sigma | L_\alpha^M(t) | \sigma \rangle \\
&= \sum_{i,j}^3 \left[\text{Re}(U_{\alpha i}^* U_{\sigma i} U_{\alpha j} U_{\sigma j}^*) \left(\cos\{E_i(\mathbf{q})t\} \cos\{E_j(\mathbf{q})t\} + \frac{\mathbf{q}^2}{E_i(\mathbf{q})E_j(\mathbf{q})} \sin\{E_i(\mathbf{q})t\} \sin\{E_j(\mathbf{q})t\} \right) \right. \\
&\quad \left. - \text{Im}(U_{\alpha i}^* U_{\sigma i} U_{\alpha j} U_{\sigma j}^*) \left(\frac{|\mathbf{q}|}{E_i(\mathbf{q})} \sin\{E_i(\mathbf{q})t\} \cos\{E_j(\mathbf{q})t\} - \frac{|\mathbf{q}|}{E_j(\mathbf{q})} \cos\{E_i(\mathbf{q})t\} \sin\{E_j(\mathbf{q})t\} \right) \right. \\
&\quad \left. - \text{Re}(U_{\alpha i}^* U_{\sigma i}^* U_{\alpha j} U_{\sigma j}) \frac{m_i}{E_i(\mathbf{q})} \frac{m_j}{E_j(\mathbf{q})} \sin\{E_i(\mathbf{q})t\} \sin\{E_j(\mathbf{q})t\} \right]. \quad (17)
\end{aligned}$$

For our notation the initial state σ denotes the neutrino with a definite flavor i.e., $\sigma = e, \mu, \tau$ and the momentum \mathbf{q} . The energy of the mass states we define as $E_{i,j}(\mathbf{q}) = \sqrt{\mathbf{q}^2 + m_{i,j}^2}$. To extract the Majorana type-phases we are interested in the third term of Eq.(17) that depends on the PMNS combination of $\text{Re}(U_{\alpha i}^* U_{\sigma i}^* U_{\alpha j} U_{\sigma j})$. To isolate the third term, consider that the first two terms can be completely determined from neutrino oscillation and neutrino mass experiments. This means we can subtract those terms from the Majorana expectation value to isolate the third term. We define the quantity $L_\alpha^Q(t)$ which denotes the sum of the first and the second term of Eq.(17),

$$\begin{aligned}
& \langle \sigma | L_\alpha^Q(t) | \sigma \rangle \\
&= \sum_{i,j}^3 \left[\text{Re}(U_{\alpha i}^* U_{\sigma i} U_{\alpha j} U_{\sigma j}^*) \left(\cos\{E_i(\mathbf{q})t\} \cos\{E_j(\mathbf{q})t\} + \frac{\mathbf{q}^2}{E_i(\mathbf{q})E_j(\mathbf{q})} \sin\{E_i(\mathbf{q})t\} \sin\{E_j(\mathbf{q})t\} \right) \right. \\
&\quad \left. - \text{Im}(U_{\alpha i}^* U_{\sigma i} U_{\alpha j} U_{\sigma j}^*) \left(\frac{|\mathbf{q}|}{E_i(\mathbf{q})} \sin\{E_i(\mathbf{q})t\} \cos\{E_j(\mathbf{q})t\} - \frac{|\mathbf{q}|}{E_j(\mathbf{q})} \cos\{E_i(\mathbf{q})t\} \sin\{E_j(\mathbf{q})t\} \right) \right]. \quad (18)
\end{aligned}$$

We represent that subtraction process with the difference between the Majorana expectation value Eq.(17) and the quantity Eq.(18),

$$\langle \sigma | L_\alpha^{M-Q}(t) | \sigma \rangle \equiv \langle \sigma | L_\alpha^M(t) | \sigma \rangle - \langle \sigma | L_\alpha^Q(t) | \sigma \rangle. \quad (19)$$

This results in the isolated third term of,

$$\begin{aligned}
\langle \sigma | L_\alpha^{M-Q}(t) | \sigma \rangle &= - \sum_{i,j}^3 \text{Re}(U_{\alpha i}^* U_{\sigma i}^* U_{\alpha j} U_{\sigma j}) \frac{m_i}{E_i(\mathbf{q})} \frac{m_j}{E_j(\mathbf{q})} \sin\{E_i(\mathbf{q})t\} \sin\{E_j(\mathbf{q})t\}, \\
&= - \sum_{i=1}^3 |U_{\alpha i} U_{\sigma i}|^2 \left(\frac{m_i \sin\{E_i(\mathbf{q})t\}}{E_i(\mathbf{q})} \right)^2 \\
&\quad - 2 \sum_{i<j} \text{Re}(U_{\alpha i}^* U_{\alpha j} U_{\sigma i}^* U_{\sigma j}) \frac{m_i}{E_i(\mathbf{q})} \frac{m_j}{E_j(\mathbf{q})} \sin\{E_i(\mathbf{q})t\} \sin\{E_j(\mathbf{q})t\}.
\end{aligned} \tag{20}$$

Let us investigate the difference by focusing on the dependence of the Majorana type-phases from Eq.(6). We take $\sigma = \alpha = e$ to obtain,

$$\begin{aligned}
\langle e | L_e^{M-Q}(t) | e \rangle &= - \sum_{i=1}^3 |U_{ei}|^4 \left(\frac{m_i \sin\{E_i(\mathbf{q})t\}}{E_i(\mathbf{q})} \right)^2 \\
&\quad - 2 \text{Re}\{(U_{e1}^* U_{e2})^2\} \frac{m_1 \sin\{E_1(\mathbf{q})t\}}{E_1(\mathbf{q})} \frac{m_2 \sin\{E_2(\mathbf{q})t\}}{E_2(\mathbf{q})} \\
&\quad - 2 \text{Re}\{(U_{e2}^* U_{e3})^2\} \frac{m_2 \sin\{E_2(\mathbf{q})t\}}{E_2(\mathbf{q})} \frac{m_3 \sin\{E_3(\mathbf{q})t\}}{E_3(\mathbf{q})} \\
&\quad - 2 \text{Re}\{(U_{e1}^* U_{e3})^2\} \frac{m_1 \sin\{E_1(\mathbf{q})t\}}{E_1(\mathbf{q})} \frac{m_3 \sin\{E_3(\mathbf{q})t\}}{E_3(\mathbf{q})}.
\end{aligned} \tag{21}$$

In the last three terms of Eq.(21), the PMNS combinations can be written with the Majorana type-phases $2\beta_1$ and $2\gamma_1$ based on Eq.(6) and Eq.(7). The explicit dependence of Eq.(21) on the Majorana type-phases is then,

$$\begin{aligned}
\langle e | L_e^{M-Q}(t) | e \rangle &= - \sum_{i=1}^3 |U_{ei}|^4 \left(\frac{m_i \sin\{E_i(\mathbf{q})t\}}{E_i(\mathbf{q})} \right)^2 \\
&\quad - 2 |U_{e1} U_{e2}|^2 \cos(2\beta_1) \frac{m_1 \sin\{E_1(\mathbf{q})t\}}{E_1(\mathbf{q})} \frac{m_2 \sin\{E_2(\mathbf{q})t\}}{E_2(\mathbf{q})} \\
&\quad - 2 |U_{e2} U_{e3}|^2 \cos(2(\gamma_1 - \beta_1)) \frac{m_2 \sin\{E_2(\mathbf{q})t\}}{E_2(\mathbf{q})} \frac{m_3 \sin\{E_3(\mathbf{q})t\}}{E_3(\mathbf{q})} \\
&\quad - 2 |U_{e1} U_{e3}|^2 \cos(2\gamma_1) \frac{m_1 \sin\{E_1(\mathbf{q})t\}}{E_1(\mathbf{q})} \frac{m_3 \sin\{E_3(\mathbf{q})t\}}{E_3(\mathbf{q})}.
\end{aligned} \tag{22}$$

We can obtain similar results as Eq.(22) for the muon and tauon numbers of $\alpha = \mu, \tau$,

$$\begin{aligned}
\langle e | L_\mu^{M-Q}(t) | e \rangle &= - \sum_{i=1}^3 |U_{\mu i} U_{e i}|^2 \left(\frac{m_i \sin\{E_i(\mathbf{q})t\}}{E_i(\mathbf{q})} \right)^2 \\
&\quad - 2 |U_{e1}^* U_{e2} U_{\mu 1}^* U_{\mu 2}| \cos(\beta_1 + \beta_2) \frac{m_1 \sin\{E_1(\mathbf{q})t\}}{E_1(\mathbf{q})} \frac{m_2 \sin\{E_2(\mathbf{q})t\}}{E_2(\mathbf{q})} \\
&\quad - 2 |U_{e2}^* U_{e3} U_{\mu 2}^* U_{\mu 3}| \cos(\gamma_1 - \beta_1 + \gamma_2 - \beta_2) \frac{m_2 \sin\{E_2(\mathbf{q})t\}}{E_2(\mathbf{q})} \frac{m_3 \sin\{E_3(\mathbf{q})t\}}{E_3(\mathbf{q})} \\
&\quad - 2 |U_{e1}^* U_{e3} U_{\mu 1}^* U_{\mu 3}| \cos(\gamma_1 + \gamma_2) \frac{m_1 \sin\{E_1(\mathbf{q})t\}}{E_1(\mathbf{q})} \frac{m_3 \sin\{E_3(\mathbf{q})t\}}{E_3(\mathbf{q})}.
\end{aligned} \tag{23}$$

In contrast to Eq.(22), the muon number $L_\mu^{M-Q}(t)$ of Eq.(23) depends on the Majorana type-phases $\beta_1 + \beta_2$ and $\gamma_1 + \gamma_2$. Finally, the tauon number $L_\tau^{M-Q}(t)$ is written as,

$$\begin{aligned}
\langle e | L_\tau^{M-Q}(t) | e \rangle &= - \sum_{i=1}^3 |U_{\tau i} U_{e i}|^2 \left(\frac{m_i \sin\{E_i(\mathbf{q})t\}}{E_i(\mathbf{q})} \right)^2 \\
&\quad - 2 |U_{e1}^* U_{e2} U_{\tau 1}^* U_{\tau 2}| \cos(\beta_1 + \beta_3) \frac{m_1 \sin\{E_1(\mathbf{q})t\}}{E_1(\mathbf{q})} \frac{m_2 \sin\{E_2(\mathbf{q})t\}}{E_2(\mathbf{q})} \\
&\quad - 2 |U_{e2}^* U_{e3} U_{\tau 2}^* U_{\tau 3}| \cos(\gamma_1 - \beta_1 + \gamma_3 - \beta_3) \frac{m_2 \sin\{E_2(\mathbf{q})t\}}{E_2(\mathbf{q})} \frac{m_3 \sin\{E_3(\mathbf{q})t\}}{E_3(\mathbf{q})} \\
&\quad - 2 |U_{e1}^* U_{e3} U_{\tau 1}^* U_{\tau 3}| \cos(\gamma_1 + \gamma_3) \frac{m_1 \sin\{E_1(\mathbf{q})t\}}{E_1(\mathbf{q})} \frac{m_3 \sin\{E_3(\mathbf{q})t\}}{E_3(\mathbf{q})}.
\end{aligned} \tag{24}$$

The last three terms are written with the Majorana type-phases $\beta_1 + \beta_3$ and $\gamma_1 + \gamma_3$. All the lepton family numbers Eqs.(22-24) depend on the sum of Majorana type-phases $\beta_1 + \beta_j$ and $\gamma_1 + \gamma_j$ where $j = 1, 2, 3$. In contrast to the difference of Majorana type-phases in an unitarity triangle such as $\beta_i - \beta_j$ and $\gamma_i - \gamma_j$, the sum of the Majorana type-phases depend on the Majorana phases α_{21} and α_{31} of Eq.(4).

As we have seen in Eqs.(22-24), one can identify the part which depends on the summation of the Majorana type-phases, i.e., $\beta_i + \beta_j$ and $\gamma_i + \gamma_j$. One may determine the combination of Majorana type-phases by fitting the curve for Eq.(20) as we vary those unknown parameters. This method may require the measurement of the long time behaviour of the time evolution. In the next section, we propose other quantities directly related to them from the derivatives of the expectation values with respect to a slice of time.

4 Determination of Majorana type-phases and the lightest neutrino masses with the lepton numbers

In this section, we derive the formula to determine the lightest neutrino mass and Majorana phases with time dependence of the lepton numbers. We consider two cases,

1. there are three massive neutrinos where two Majorana phases are allowed,
2. there is a massless neutrino and only one Majorana phase is allowed.

We take the second-order time derivative of Eq.(17) at the initial time $t = 0$,

$$\begin{aligned} \frac{d^2}{dt^2} \langle \sigma | L_\alpha^M(t) | \sigma \rangle |_{t=0} &= - \sum_{i,j}^3 \text{Re}(U_{\alpha i}^* U_{\sigma i} U_{\alpha j} U_{\sigma j}^*) (m_i^2 + m_j^2) - 2 \sum_{i,j}^3 \text{Re}(U_{\alpha i}^* U_{\sigma i}^* U_{\alpha j} U_{\sigma j}) m_i m_j \\ &= -2 \sum_i^3 \delta_{\alpha\sigma} |U_{\sigma i}|^2 m_i^2 - 2 \sum_{i,j}^3 \text{Re}(U_{\alpha i}^* U_{\sigma i}^* U_{\alpha j} U_{\sigma j}) m_i m_j. \end{aligned} \quad (25)$$

The first term is independent of the Majorana phases and the second term depends on the Majorana phases though the PMNS combination $\text{Re}(U_{\alpha i}^* U_{\sigma i}^* U_{\alpha j} U_{\sigma j})$. Then, the second-order derivative of the total lepton number $L^M(t) = \sum_\alpha L_\alpha^M(t)$ is given by,

$$\frac{d^2}{dt^2} \langle \sigma | L^M(t) | \sigma \rangle |_{t=0} = -4 \sum_i^3 m_i^2 |U_{\sigma i}|^2. \quad (26)$$

We use the total lepton number of Eq.(26) to rewrite the first term of Eq.(25) resulting in,

$$\frac{d^2}{dt^2} \langle \sigma | L_\alpha^M(t) | \sigma \rangle |_{t=0} = \delta_{\alpha\sigma} \frac{1}{2} \frac{d^2}{dt^2} \langle \sigma | L^M(t) | \sigma \rangle |_{t=0} - 2 \sum_{i,j}^3 \text{Re}(U_{\alpha i}^* U_{\sigma i}^* U_{\alpha j} U_{\sigma j}) m_i m_j. \quad (27)$$

Using Eq.(26), one can derive the lightest neutrino mass for the normal hierarchy and the inverted hierarchy cases in terms of the second-order derivative of the total lepton number.

$$m_1^2 = -\frac{1}{4} \frac{d^2}{dt^2} \langle \sigma | L^M(t) | \sigma \rangle |_{t=0} - \Delta m_{21}^2 |U_{\sigma 2}|^2 - \Delta m_{31}^2 |U_{\sigma 3}|^2 \quad \text{Normal}, \quad (28)$$

$$m_3^2 = -\frac{1}{4} \frac{d^2}{dt^2} \langle \sigma | L^M(t) | \sigma \rangle |_{t=0} - \Delta m_{13}^2 |U_{\sigma 1}|^2 - \Delta m_{23}^2 |U_{\sigma 2}|^2 \quad \text{Inverted}. \quad (29)$$

To investigate the Majorana type-phases, we consider two cases for the lepton families. First we set $\alpha = \sigma$ in Eq.(27) and obtain the following formula,

$$\frac{d^2}{dt^2} \langle \sigma | L_\sigma^M(t) | \sigma \rangle |_{t=0} = \frac{1}{2} \frac{d^2}{dt^2} \langle \sigma | L^M(t) | \sigma \rangle |_{t=0} - 2 \sum_{i,j}^3 (U_{\sigma i}^* U_{\sigma j})^2 m_i m_j. \quad (30)$$

Then we set $\alpha \neq \sigma$ in Eq.(27) to obtain,

$$\frac{d^2}{dt^2} \langle \sigma | L_\alpha^M(t) | \sigma \rangle |_{t=0} = -2 \sum_{i,j}^3 \text{Re}(U_{\alpha i}^* U_{\alpha j} U_{\sigma i}^* U_{\sigma j}) m_i m_j. \quad (31)$$

Next, by specifying the family indices σ and α in Eq.(30) and Eq.(31) we can clarify the dependencies on the Majorana type-phases from Eqs.(6-7). We take $\sigma = e$ in Eq.(30) to show the dependencies on the Majorana type-phases β_1 and γ_1 ,

$$\begin{aligned} \frac{d^2}{dt^2} \langle e | L_e^M(t) | e \rangle |_{t=0} &= \frac{1}{2} \frac{d^2}{dt^2} \langle e | L^M(t) | e \rangle |_{t=0} - 2 \sum_{i=1}^3 m_i^2 |U_{ei}|^4 - 4m_1 m_2 |U_{e1} U_{e2}^*|^2 \cos(2\beta_1) \\ &\quad - 4m_2 m_3 |U_{e2} U_{e3}^*|^2 \cos(2\beta_1 - 2\gamma_1) - 4m_3 m_1 |U_{e1} U_{e3}^*|^2 \cos(2\gamma_1). \end{aligned} \quad (32)$$

In contrast, for Eq.(31) we take $\sigma = e$ and $\alpha = \mu$,

$$\begin{aligned} \frac{d^2}{dt^2} \langle e | L_\mu^M(t) | e \rangle |_{t=0} &= -2 \sum_i^3 |U_{\mu i}|^2 |U_{ei}|^2 m_i^2 - 4 |U_{\mu 1}^* U_{\mu 2} U_{e1}^* U_{e2}| \cos(\beta_1 + \beta_2) m_1 m_2 \\ &\quad - 4 |U_{\mu 2}^* U_{\mu 3} U_{e2}^* U_{e3}| \cos(\beta_1 - \gamma_1 + \beta_2 - \gamma_2) m_2 m_3 \\ &\quad - 4 |U_{\mu 3}^* U_{\mu 1} U_{e3}^* U_{e1}| \cos(\gamma_1 + \gamma_2) m_3 m_1, \end{aligned} \quad (33)$$

where we have dependence on the additional Majorana type-phases β_2 and γ_2 compared to Eq.(32). Lastly we take $\alpha = \tau$,

$$\begin{aligned} \frac{d^2}{dt^2} \langle e | L_\tau^M(t) | e \rangle |_{t=0} &= -2 \sum_i^3 |U_{\tau i}|^2 |U_{ei}|^2 m_i^2 - 4 |U_{\tau 1}^* U_{\tau 2} U_{e1}^* U_{e2}| \cos(\beta_1 + \beta_3) m_1 m_2 \\ &\quad - 4 |U_{\tau 2}^* U_{\tau 3} U_{e2}^* U_{e3}| \cos(\beta_1 - \gamma_1 + \beta_3 - \gamma_3) m_2 m_3 \\ &\quad - 4 |U_{\tau 3}^* U_{\tau 1} U_{e3}^* U_{e1}| \cos(\gamma_1 + \gamma_3) m_3 m_1, \end{aligned} \quad (34)$$

which results in the dependence on the Majorana type-phases β_3 and γ_3 .

As discussed near the end of section 2, the Majorana type-phases β_i and γ_i change for variation of the Majorana phases $\Delta\alpha_{21}$ and $\Delta\alpha_{31}$,

$$\beta_i + \beta_j \rightarrow \beta_i + \beta_j - \Delta\alpha_{21}, \quad (35)$$

$$\gamma_i + \gamma_j \rightarrow \gamma_i + \gamma_j - \Delta\alpha_{31}. \quad (36)$$

The time derivatives of all the three lepton family numbers in Eq.(32-34) are dependent on summations of the Majorana type-phases i.e., $\beta_i + \beta_j$ and $\gamma_i + \gamma_j$. From Eqs.(35-36) we know $\beta_i + \beta_j$ and $\gamma_i + \gamma_j$ are sensitive to the Majorana phases $\Delta\alpha_{21}$ and $\Delta\alpha_{31}$. Thus, the two Majorana phases α_{21} and α_{31} can be determined by knowing the time derivatives of the two or more lepton family numbers. In addition, we can resolve the lightest neutrino mass from the time derivatives of the total lepton number in Eqs.(28-29). This is a method to determine the triangle orientations and the lightest neutrino mass based on lepton number oscillations.

5 Considerations when the lightest neutrino is massless

In this section, we consider the lightest neutrino to be massless. As an example, the (3, 2) Type-I seesaw model can predict such a massless neutrino with three active neutrinos and two heavy right-handed Majorana neutrinos[58]. In this framework, which neutrino is massless depends on the hierarchy,

$$m_1 = 0, \quad 0 < m_2 < m_3, \quad \text{Normal hierarchy}, \quad (37)$$

$$m_3 = 0, \quad 0 < m_1 < m_2, \quad \text{Inverted hierarchy.} \quad (38)$$

Because a massless neutrino is free to be re-phased, a few of the six Majorana type-phases from the three massive Majorana neutrinos in the previous section, 4, are no longer invariants. In the normal hierarchy case, the invariant combinations of the three β_i and three γ_i become,

$$\arg(U_{e2}U_{e3}^*) = \gamma_1 - \beta_1, \quad \arg(U_{\mu2}U_{\mu3}^*) = \gamma_2 - \beta_2, \quad \arg(U_{\tau2}U_{\tau3}^*) = \gamma_3 - \beta_3. \quad (39)$$

This is because, β_i and γ_i are no longer re-phasing invariants by themselves and one can form four invariant quartets,

$$\arg(U_{e1}^*U_{e2}U_{\mu1}U_{\mu2}^*) = \beta_2 - \beta_1 \quad (40)$$

$$\arg(U_{e1}^*U_{e2}U_{\tau1}U_{\tau2}^*) = \beta_3 - \beta_1 \quad (41)$$

$$\arg(U_{e1}^*U_{e3}U_{\mu1}U_{\mu3}^*) = \gamma_2 - \gamma_1 \quad (42)$$

$$\arg(U_{e1}^*U_{e3}U_{\tau1}U_{\tau3}^*) = \gamma_3 - \gamma_1. \quad (43)$$

The Majorana type-phases defined in Eq.(39) and the four arguments defined in Eqs.(40-43) are not independent. In addition to a Majorana type-phase $\gamma_1 - \beta_1$ in Eq.(39), one can choose the four arguments of the quartets as independent re-phasing invariant combinations. This means the other two Majorana type-phases can be written using them,

$$\begin{aligned} \gamma_i - \beta_i &= (\gamma_1 - \beta_1) + (\gamma_i - \beta_i) - (\gamma_1 - \beta_1) \\ &= (\gamma_1 - \beta_1) + (\gamma_i - \gamma_1) - (\beta_i - \beta_1) \quad (i = 2, 3). \end{aligned} \quad (44)$$

A similar situation occurs for the inverted hierarchy massless neutrino of $m_3 = 0$. The re-phasing invariant Majorana type-phases become β_i , for $i = 1, 2, 3$. Then, one can choose a Majorana type-phase and the four arguments in Eqs.(40-43) as independent re-phasing invariants.

5.1 Neutrinoless double beta decay and lepton number

In this subsection, we relate one independent Majorana type-phase to physical observables, such as $|m_{\nu ee}|$ related to the neutrinoless double beta decay rate and time evolution of lepton numbers. For the normal hierarchy case, the $|m_{\nu ee}|$ is given by the following formula,

$$|m_{\nu ee}|_{\text{norm}}^2 = m_2^2|U_{e2}|^4 + m_3^2|U_{e3}|^4 + 2m_2m_3|U_{e2}|^2|U_{e3}|^2 \cos 2(\gamma_1 - \beta_1). \quad (45)$$

Whereas for the inverted hierarchy case it is,

$$|m_{\nu ee}|_{\text{inv}}^2 = m_1^2|U_{e1}|^4 + m_2^2|U_{e2}|^4 + 2m_1m_2|U_{e1}|^2|U_{e2}|^2 \cos 2\beta_1. \quad (46)$$

From the expressions in Eq.(45-46), the determination of $|m_{\nu ee}|_{\text{norm,inv}}$ is sufficient to identify a single Majorana type-phase; provided the other moduli of the PMNS matrix elements in Eq.(5) and two non-zero masses of the neutrinos are extracted from the neutrino oscillation experiments.

Next, we investigate the time evolution of lepton number when the lightest neutrino is massless, as in Eqs.(37-38). From Eq.(28), for the normal hierarchy case, the second-order time

derivative of the total lepton number is related to the combination of the two non-vanishing masses;

$$\frac{1}{4} \frac{d^2}{dt^2} \langle \sigma | L^M(t) | \sigma \rangle |_{t=0} = -m_2^2 |U_{\sigma 2}|^2 - m_3^2 |U_{\sigma 3}|^2. \quad (47)$$

Then we can rewrite the second-order time derivatives of the electron, muon, and tauon lepton numbers of Eqs.(32-34) as,

$$\frac{d^2}{dt^2} \langle e | L_e^M(t) | e \rangle |_{t=0} = \frac{d^2}{dt^2} \langle e | L^M(t) | e \rangle |_{t=0} - 4m_2 m_3 |U_{e2} U_{e3}^*|^2 \cos(2\beta_1 - 2\gamma_1), \quad (48)$$

$$\frac{d^2}{dt^2} \langle e | L_\mu^M(t) | e \rangle |_{t=0} = -2 \sum_{i=2}^3 |U_{\mu i}|^2 |U_{ei}|^2 m_i^2 - 4 |U_{\mu 2}^* U_{\mu 3} U_{e2}^* U_{e3}| \cos(\beta_1 - \gamma_1 + \beta_2 - \gamma_2) m_2 m_3, \quad (49)$$

$$\frac{d^2}{dt^2} \langle e | L_\tau^M(t) | e \rangle |_{t=0} = -2 \sum_{i=2}^3 |U_{\tau i}|^2 |U_{ei}|^2 m_i^2 - 4 |U_{\tau 2}^* U_{\tau 3} U_{e2}^* U_{e3}| \cos(\beta_1 - \gamma_1 + \beta_3 - \gamma_3) m_2 m_3. \quad (50)$$

The second-order time derivative for the normal hierarchy in Eq.(48) can be used to determine one of the Majorana like-phases; $\gamma_1 - \beta_1$. The other two Majorana type-phases, $\gamma_i - \beta_i$ ($i = 1, 2$), can be determined through Eq.(44) by knowing the arguments of the quartet in Eq.(40). This allows us to learn of the orientation of triangle 3 in Fig.2. Alternatively, with the three second-order time derivatives of Eqs.(48)-(50) one can determine three Majorana type-phases $\gamma_i - \beta_i$ ($i = 1 - 3$).

In the inverted hierarchy case m_3 is the lightest neutrino mass and the second-order time derivative of the total lepton number becomes,

$$\frac{1}{4} \frac{d^2}{dt^2} \langle \sigma | L^M(t) | \sigma \rangle |_{t=0} = -m_2^2 |U_{\sigma 2}|^2 - m_1^2 |U_{\sigma 1}|^2. \quad (51)$$

The major difference between the normal hierarchy of Eq.(47) and the inverted hierarchy of Eq.(51) is the last mass multiplied by PMNS matrix term. This modifies the second-order time derivatives to become,

$$\frac{d^2}{dt^2} \langle e | L_e^M(t) | e \rangle |_{t=0} = \frac{d^2}{dt^2} \langle e | L^M(t) | e \rangle |_{t=0} - 4m_1 m_2 |U_{e1} U_{e2}^*|^2 \cos(2\beta_1), \quad (52)$$

$$\frac{d^2}{dt^2} \langle e | L_\mu^M(t) | e \rangle |_{t=0} = -2 \sum_i^2 |U_{\mu i}|^2 |U_{ei}|^2 m_i^2 - 4 |U_{\mu 1}^* U_{\mu 2} U_{e1}^* U_{e2}| \cos(\beta_1 + \beta_2) m_1 m_2, \quad (53)$$

$$\frac{d^2}{dt^2} \langle e | L_\tau^M(t) | e \rangle |_{t=0} = -2 \sum_i^2 |U_{\tau i}|^2 |U_{ei}|^2 m_i^2 - 4 |U_{\tau 1}^* U_{\tau 2} U_{e1}^* U_{e2}| \cos(\beta_1 + \beta_3) m_1 m_2. \quad (54)$$

Similar to the normal hierarchy equations of Eqs.(48-50), the second-order time derivatives of the inverted hierarchy in Eqs.(52-54) allow for a complete determination of the Majorana type-phases β_i and the orientation of triangle 1 in Fig.2.

5.2 Numerical illustration for two neutrino generations

In this subsection, we illustrate the analytical results of the relations among the lightest neutrino mass, a Majorana type-phase, and the second-order time derivative of lepton numbers. We illustrate them using the two generation model. In the model, there is only two Majorana type-phases, β_1 and β_2 defined in Eq.(6). They are not independent because the unitary relation

$$U_{e1}U_{e2}^* = -U_{\mu1}U_{\mu2}^*, \quad (55)$$

holds. It leads to the relation between the two Majorana type-phases,

$$\beta_1 = \arg(U_{e1}U_{e2}^*) = \arg(U_{\mu1}U_{\mu2}^*) - \pi = \beta_2 - \pi. \quad (56)$$

Adopting the following parametrization for a 2 by 2 unitary PMNS matrix,

$$\begin{pmatrix} \cos \theta_{12} & \sin \theta_{12} e^{i\frac{\alpha_{21}}{2}} \\ -\sin \theta_{12} & \cos \theta_{12} e^{i\frac{\alpha_{21}}{2}} \end{pmatrix}, \quad (57)$$

the Majorana type-phase β_1 is related to a Majorana phase as, $\beta_1 = -\frac{\alpha_{21}}{2}$. Contrary to the three generation model, one can not write $\tan^2 \theta_{12}$ in terms of the Majorana type-phase.

In this model, the lightest neutrino mass m_1 and a Majorana type-phase β_1 are unknown parameters to be determined by the time evolution of lepton numbers. We assume that a mass squared difference Δm_{21}^2 and a mixing angle θ_{12} are measured by the oscillation experiment. The time evolution of the electron and muon numbers for two generation model are given by,

$$\begin{aligned} \langle e | L_e(t) | e \rangle &= c_{12}^4 \left(1 - \frac{2m_1^2 \sin^2(E_1 t)}{E_1^2} \right) + s_{12}^4 \left(1 - \frac{2m_2^2 \sin^2(E_2 t)}{E_2^2} \right) \\ &+ s_{12}^2 c_{12}^2 \left\{ \left(1 + \frac{q^2 - m_1 m_2 \cos(2\beta_1)}{E_1 E_2} \right) \cos\{(E_1 - E_2)t\} \right. \\ &\left. + \left(1 - \frac{q^2 - m_1 m_2 \cos(2\beta_1)}{E_1 E_2} \right) \cos\{(E_1 + E_2)t\} \right\}, \end{aligned} \quad (58)$$

$$\begin{aligned} \langle e | L_\mu(t) | e \rangle &= c_{12}^2 s_{12}^2 \left(\left(1 - \frac{2m_1^2 \sin^2(E_1 t)}{E_1^2} \right) + \left(1 - \frac{2m_2^2 \sin^2(E_2 t)}{E_2^2} \right) \right) \\ &- s_{12}^2 c_{12}^2 \left\{ \left(1 + \frac{q^2 - m_1 m_2 \cos(2\beta_1)}{E_1 E_2} \right) \cos\{(E_1 - E_2)t\} \right. \\ &\left. + \left(1 - \frac{q^2 - m_1 m_2 \cos(2\beta_1)}{E_1 E_2} \right) \cos\{(E_1 + E_2)t\} \right\}. \end{aligned} \quad (59)$$

One can also compute the expectation values of the total lepton number $L(t) = L_e(t) + L_\mu(t)$ and the difference $L_{e-\mu}(t) = L_e(t) - L_\mu(t)$. The straightforward calculation leads to the second order time derivatives for the total lepton number at $t = 0$ as follows,

$$\frac{d^2}{dt^2} \langle e | L(t) | e \rangle \Big|_{t=0} = -4(m_1^2 c_{12}^2 + m_2^2 s_{12}^2). \quad (60)$$




Figure 4: The time dependence of the total lepton number for different lightest neutrino masses. The momentum $q = 0.002(\text{eV})$. The dashed line shows the case for $m_1 = 0.01(\text{eV})$ and the solid line shows the case for $m_1 = 0.02(\text{eV})$. We use $\Delta m_{21}^2 = 7.42 \times 10^{-5} \text{eV}^2$ and $\sin(\theta_{12}) = 0.551$.

Figure 4 shows that total lepton number sharply decreases at $t = 0$ for the larger the lightest neutrino mass m_1 . This can be also understood by solving the lightest neutrino mass with the second derivatives of Eq.(60), a mixing angle and a mass squared difference.

$$m_1^2 = -\frac{1}{4} \frac{d^2}{dt^2} \langle e|L(t)|e \rangle \Big|_{t=0} - \Delta m_{21}^2 s_{12}^2. \quad (61)$$

The second order time derivative for the electron minus muon number $L_{e-\mu}$ at $t = 0$ is given as follows,

$$\frac{d^2}{dt^2} \langle e|L_{e-\mu}(t)|e \rangle \Big|_{t=0} = -4|m_{\nu ee}|_{\text{two gene.}}^2, \quad (62)$$

where $|m_{\nu ee}|_{\text{two gene.}}^2$ is given by the same formula in Eq.(46) as the inverted hierarchy case with $m_3 = 0$. By substituting the mixing angle to $|U_{ei}|$ ($i = 1, 2$) in Eq.(57), it is given by,

$$|m_{\nu ee}|_{\text{two gene.}}^2 = m_1^2 c_{12}^4 + m_2^2 s_{12}^4 + 2m_1 m_2 c_{12}^2 s_{12}^2 \cos 2\beta_1. \quad (63)$$

Eq.(62) tells us that $L_{e-\mu}$ sharply decreases for the larger $|m_{\nu ee}|$. Fig.(5) shows the dependence of $L_{e-\mu}$ on Majorana type-phase β_1 . Because $|m_{\nu ee}|$ is the largest for $\beta_1 = 0$ and the smallest for $\beta_1 = \pi$, as shown in Eq.(63), Fig.(5) numerically confirms the dependence on the $|m_{\nu ee}|$ of Eq.(62).

One can also solve for a Majorana type-phase β_1 with the second order derivative of Eqs.(62), a mixing angle and a mass squared difference.

$$\cos(2\beta_1) = \frac{-\frac{1}{4} \frac{d^2}{dt^2} \langle e|L_{e-\mu}(t)|e \rangle \Big|_{t=0} - (m_1^2 c_{12}^4 + m_2^2 s_{12}^4)}{2m_1 m_2 c_{12}^2 s_{12}^2}. \quad (64)$$

Eqs.(61-64) can be used to determine the lightest neutrino mass and the Majorana type-phase once the time derivatives for $\ddot{L}|_{t=0}$ and $\ddot{L}_{e-\mu}|_{t=0}$ are measured.




Figure 5: The time dependence of $L_{e-\mu} = L_e - L_\mu$ for different choice of the Majorana type-phase β_1 . $q = 0.002(\text{eV})$ $m_1 = 0.01(\text{eV})$ and $\sin(\theta_{12}) = 0.551$. The dashed line shows the case for $\beta_1 = 0$. The dot-dashed line shows the case for $2\beta_1 = \frac{\pi}{2}$ and solid line shows the case for $2\beta_1 = \pi$. Δm_{21}^2 is the same as in Fig.4.

6 Concluding Remarks

We have investigated an approach to extract the Majorana type-phases of Branco and Rebelo using the time evolution of lepton numbers. The specific combinations of Majorana type-phases in the same triangle are related to the orientation of unitarity triangles for the PMNS matrix, and the Majorana phases α_{21} and α_{31} . After taking the second-order time derivative of the lepton number expectation values, the dependencies on the summation of Majorana type-phases i.e., $\beta_i + \beta_j$ and $\gamma_i + \gamma_j$, can be determined. Thus allowing the extraction of the orientation of unitarity triangles for the PMNS matrix, and the Majorana phases.

We view our result as complimentary to using neutrinoless double-beta decay for determining the orientation of unitarity triangles for the PMNS matrix, and the Majorana phases. We also show the time derivative of the total lepton number is sensitive to the lightest neutrino mass. The above features are also numerically demonstrated for two generation toy model.

For the future, we are interested in identifying a possible experimental setup used to measure the quantities discussed.

Acknowledgement

This work is supported by Japan Society for the Promotion of Science (JSPS) KAKENHI Grant Number JP17K05418 (T.M) and JP21K13923(K.Y). One author, N.J.B, would like to express thanks to the Japanese government Ministry of Education, Culture, Sports, Science, and Technology (MEXT) for the financial support during the writing of this work.

References

- [1] S. Schael *et al.* [ALEPH, DELPHI, L3, OPAL, SLD, LEP Electroweak Working Group, SLD Electroweak Group and SLD Heavy Flavour Group], Phys. Rept. **427**, 257-454

(2006)

- [2] Y. Fukuda *et al.* [Super-Kamiokande], Phys. Rev. Lett. **81**, 1562-1567 (1998)
- [3] Q. R. Ahmad *et al.* [SNO], Phys. Rev. Lett. **89**, 011301 (2002)
- [4] B. Pontecorvo, Zh. Eksp. Teor. Fiz. **53**, 1717-1725 (1967)
- [5] V. N. Gribov and B. Pontecorvo, Phys. Lett. B **28**, 493 (1969)
- [6] E. Majorana, Nuovo Cim. **14**, 171-184 (1937)
- [7] G. Racah, Nuovo Cim. **14**, 322-328 (1937)
- [8] W. H. Furry, Phys. Rev. **54**, 56-67 (1938)
- [9] W. H. Furry, Phys. Rev. **56**, 1184-1193 (1939)
- [10] J. Schechter and J. W. F. Valle, Phys. Rev. D **25**, 2951 (1982)
doi:10.1103/PhysRevD.25.2951
- [11] J. F. Nieves, Phys. Lett. B **147**, 375-379 (1984)
- [12] E. Takasugi, Phys. Lett. B **149**, 372-376 (1984)
- [13] M. J. Dolinski, A. W. P. Poon and W. Rodejohann, Ann. Rev. Nucl. Part. Sci. **69**, 219-251 (2019)
- [14] S. Umehara, T. Kishimoto, I. Ogawa, R. Hazama, H. Miyawaki, S. Yoshida, K. Matsuoka, K. Kishimoto, A. Katsuki and H. Sakai, *et al.* Phys. Rev. C **78**, 058501 (2008)
- [15] M. Agostini *et al.* [GERDA], Phys. Rev. Lett. **125**, no.25, 252502 (2020)
- [16] S. I. Alvis *et al.* [Majorana], Phys. Rev. C **100**, no.2, 025501 (2019)
- [17] A. S. Barabash *et al.* [NEMO], Phys. Atom. Nucl. **74**, 312-317 (2011)
doi:10.1134/S1063778811020062 [arXiv:1002.2862 [nucl-ex]].
- [18] R. Arnold *et al.* [NEMO-3], Phys. Rev. D **92**, no.7, 072011 (2015)
- [19] R. Arnold *et al.* [NEMO-3], Phys. Rev. D **93**, no.11, 112008 (2016)
- [20] R. Arnold *et al.* [NEMO-3], Phys. Rev. D **94**, no.7, 072003 (2016)
- [21] R. Arnold *et al.* [NEMO-3], Phys. Rev. D **95**, no.1, 012007 (2017)
- [22] S. Abe *et al.* [KamLAND-Zen],
- [23] G. Anton *et al.* [EXO-200], Phys. Rev. Lett. **123**, no.16, 161802 (2019)
- [24] R. Arnold *et al.* [NEMO-3], Phys. Rev. Lett. **119**, no.4, 041801 (2017)

- [25] R. G. Winter, Phys. Rev. **100**, 142-144 (1955) doi:10.1103/PhysRev.100.142
- [26] J. Bernabeu, A. De Rujula and C. Jarlskog, Nucl. Phys. B **223**, 15-28 (1983) doi:10.1016/0550-3213(83)90089-5
- [27] Z. Sujkowski and S. Wycech, Phys. Rev. C **70**, 052501 (2004)
- [28] D. Q. Adams *et al.* [CUORE], Phys. Rev. C **105**, 065504 (2022)
- [29] K. Abe *et al.* [XMASS], PTEP **2018**, no.5, 053D03 (2018)
- [30] K. Blaum, S. Eliseev, F. A. Danevich, V. I. Tretyak, S. Kovalenko, M. I. Krivoruchenko, Y. N. Novikov and J. Suhonen, Rev. Mod. Phys. **92**, 045007 (2020)
- [31] J. A. Grifols, E. Masso and R. Toldra, Phys. Lett. B **389**, 563-565 (1996)
- [32] A. Segarra and J. Bernabéu, Phys. Rev. D **101**, no.9, 093004 (2020)
- [33] A. Costantino and S. Fichet, JHEP **09**, 122 (2020) doi:10.1007/JHEP09(2020)122 [arXiv:2003.11032 [hep-ph]].
- [34] A. Millar, G. Raffelt, L. Stodolsky and E. Vitagliano, Phys. Rev. D **98**, no.12, 123006 (2018)
- [35] J. F. Nieves and P. B. Pal, Phys. Rev. D **32**, 1849-1852 (1985)
- [36] T. Chhabra and P. R. Babu, Phys. Rev. D **46**, 903-909 (1992)
- [37] B. Kayser and R. E. Shrock, Phys. Lett. B **112**, 137-142 (1982) doi:10.1016/0370-2693(82)90314-8
- [38] C. S. Kim, M. V. N. Murthy and D. Sahoo, Phys. Rev. D **105**, no.11, 113006 (2022)
- [39] K. Abe *et al.* (Super-Kamiokande), Phys. Rev. D **83**, 052010 (2011)
- [40] B. Aharmim *et al.* (SNO), Phys. Rev. C **81**, 055504 (2010)
- [41] B. Pontecorvo, Sov. Phys. JETP **7**, 172 (1958) [Zh. Eksp. Teor. Fiz. **34**, 247 (1957)].
- [42] Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28**, 870 (1962).
- [43] C. Giunti and C. W. Kim, “Fundamentals of Neutrino Physics and Astrophysics,”
- [44] T. Hahn, [arXiv:physics/0607103 [physics]].
- [45] P.A. Zyla *et al.* [Particle Data Group], PTEP **2020**, no.8, 083C01 (2020)
- [46] C. Jarlskog, Phys. Rev. Lett. **55**, 1039 (1985)
- [47] G. C. Branco, L. Lavoura and M. N. Rebelo, Phys. Lett. B **180**, 264-268 (1986)
- [48] J. F. Nieves and P. B. Pal, Phys. Rev. D **36**, 315 (1987)

- [49] F. Feruglio, C. Hagedorn and R. Ziegler, *JHEP* **07**, 027 (2013) [arXiv:1211.5560 [hep-ph]].
- [50] G. C. Branco and M. N. Rebelo, *Phys. Rev. D* **79**, 013001 (2009) doi:10.1103/PhysRevD.79.013001 [arXiv:0809.2799 [hep-ph]].
- [51] J. A. Aguilar-Saavedra and G. C. Branco, *Phys. Rev. D* **62**, 096009 (2000) doi:10.1103/PhysRevD.62.096009 [arXiv:hep-ph/0007025 [hep-ph]].
- [52] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz and A. Zhou, *JHEP* **09**, 178 (2020) doi:10.1007/JHEP09(2020)178 [arXiv:2007.14792 [hep-ph]], NuFIT 5.1 (2021) www.nu-fit.org.
- [53] M. Fukugita and T. Yanagida, *Phys. Lett. B* **174**, 45-47 (1986) doi:10.1016/0370-2693(86)91126-3
- [54] T. Endoh, T. Morozumi and Z. h. Xiong, *Prog. Theor. Phys.* **111**, 123-149 (2004) doi:10.1143/PTP.111.123 [arXiv:hep-ph/0308276 [hep-ph]].
- [55] T. Fujihara, S. Kaneko, S. K. Kang, D. Kimura, T. Morozumi and M. Tanimoto, *Phys. Rev. D* **72**, 016006 (2005) doi:10.1103/PhysRevD.72.016006 [arXiv:hep-ph/0505076 [hep-ph]].
- [56] A. S. Adam, N. J. Benoit, Y. Kawamura, Y. Matsuo, T. Morozumi, Y. Shimizu, Y. Tokunaga and N. Toyota, *PTEP* **2021**, 5 (2021) doi:10.1093/ptep/ptab025 [arXiv:2101.07751 [hep-ph]].
- [57] A. S. Adam, N. J. Benoit, Y. Kawamura, Y. Matsuo, T. Morozumi, Y. Shimizu and N. Toyota, [arXiv:2106.02783 [hep-ph]].
- [58] E. Ma, D. P. Roy and U. Sarkar, *Phys. Lett. B* **444**, 391-396 (1998) doi:10.1016/S0370-2693(98)01395-1 [arXiv:hep-ph/9810309 [hep-ph]].