

A CUNTZ-KRIEGER UNIQUENESS THEOREM FOR CUNTZ-PIMSNER ALGEBRAS

MENEVŞE ERYÜZLÜ AND MARK TOMFORDE

ABSTRACT. We introduce a property of C^* -correspondences, which we call Condition (S), to serve as an analogue of Condition (L) of graphs. We use Condition (S) to prove a Cuntz-Krieger Uniqueness Theorem for Cuntz-Pimsner algebras and obtain sufficient conditions for simplicity of Cuntz-Pimsner algebras. We also prove that if \mathcal{Q} is a topological quiver with no sinks and $X(\mathcal{Q})$ is the associated C^* -correspondence, then $X(\mathcal{Q})$ satisfies Condition (S) if and only if \mathcal{Q} satisfies Condition (L). Finally, we consider several examples to compare and contrast Condition (S) with Schweizer’s nonperiodic condition.

1. INTRODUCTION

In 1997 Pimsner introduced a method for constructing a C^* -algebra \mathcal{O}_X from a C^* -correspondence X . These C^* -algebras have been named *Cuntz-Pimsner algebras*, and a great deal of research has focused on understanding how the structure of the Cuntz-Pimsner algebra \mathcal{O}_X is determined by properties of the C^* -correspondence X . As with any class of C^* -algebras, one wishes to understand the ideal structure of Cuntz-Pimsner algebras and, in particular, to characterize when Cuntz-Pimsner algebras are simple. Partial results for simplicity of Cuntz-Pimsner algebras were obtained early in the subject’s history: Let X be a C^* -correspondence over a C^* -algebra A . In [4, Corollary 4.4] Kajiwara, Pinzari, and Watatani provided sufficient conditions for simplicity of X when A is unital and X is full and finite projective, and the left action of A on X is injective and by compact operators. In addition, in [9, Theorem 1] Muhly and Solel established sufficient conditions for simplicity of \mathcal{O}_X under the hypotheses that A is unital and X is projective and essential with injective left action. In 2001 Schweizer introduced a property for C^* -correspondences that he called *nonperiodic* and proved that if A is unital and X is full with injective left action, then \mathcal{O}_X is simple if and only if X is nonperiodic and A has no nontrivial invariant ideals [14, Theorem 3.9]. Subsequently, Muhly and Solel wrote in [10] that Schweizer’s result was “the last word on simplicity” for Cuntz-Pimsner algebras. However, as we shall describe in more detail in this paper, when Schweizer’s hypotheses on X and A are removed, it is still an open problem to characterize simplicity of Cuntz-Pimsner algebras, and even more significant, we shall exhibit examples showing that when Schweizer’s hypotheses are removed, the conditions “ X is nonperiodic and A has no nontrivial invariant ideals” are insufficient to imply that \mathcal{O}_X is simple. Consequently, throughout the course of this paper we hope to convince the reader that there remain many open questions concerning

Date: November 15, 2023.

2020 Mathematics Subject Classification. 46L55, 46L08.

Key words and phrases. C^* -algebras, Cuntz-Pimsner algebras, C^* -correspondences.

This work was supported by a grant from the Simons Foundation (#527708 to Mark Tomforde).

simplicity of Cuntz-Pimsner algebras, and Schweizer's result (while significant and elegant) is far from the last word on the topic.

In the development of Cuntz-Pimsner algebras, graph C^* -algebras have provided useful intuition and guidance for the investigation of more general Cuntz-Pimsner algebras. There are two fundamental structure theorems in the theory of graph C^* -algebras: the Gauge-Invariant Uniqueness Theorem and the Cuntz-Krieger Uniqueness Theorem. Katsura has successfully extended the Gauge-Invariant Uniqueness Theorem to all Cuntz-Pimsner algebras [6, Theorem 6.4] and used this result to classify the gauge-invariant ideals in any Cuntz-Pimsner algebra [7, Theorem 8.6]. By contrast, up to this point there has been no extension of the Cuntz-Krieger Uniqueness Theorem to general Cuntz-Pimsner algebras.

A graph is said to satisfy Condition (L) if every cycle in the graph has an exit. The Cuntz-Krieger Uniqueness Theorem for graph C^* -algebras states that if a graph E satisfies Condition (L) and $\phi : C^*(E) \rightarrow A$ is a $*$ -homomorphism that is nonzero on all vertex projections, then ϕ is injective. The Cuntz-Krieger Uniqueness Theorem is used to obtain a characterization of simplicity for graph C^* -algebras, namely $C^*(E)$ is simple if and only if E satisfies Condition (L) and E contains no nontrivial saturated hereditary subsets of vertices.

It would be advantageous to generalize the Cuntz-Krieger Uniqueness Theorem to all Cuntz-Pimsner algebras. Among other applications, such a theorem would provide sufficient conditions for simplicity of general Cuntz-Pimsner algebras as an almost immediate consequence. The main obstacle to developing a generalized Cuntz-Krieger Uniqueness Theorem has been an inability to formulate an appropriate analogue of Condition (L) for C^* -correspondences. While versions of the Cuntz-Krieger Uniqueness Theorem have been obtained by Katsura for topological graph C^* -algebras [5, Theorem 5.12] and by Muhly and the second author for topological quiver C^* -algebras [12, Theorem 6.16], in these situations the C^* -correspondence is built from an object (e.g., graph, topological graph, quiver) providing notions of edges, paths, and cycles for which Condition (L) can be formulated. Up to this point, it has been unclear how to translate Condition (L) to an appropriate property entirely in terms of the C^* -correspondence.

Comparing Schweizer's simplicity result (i.e., a Cuntz-Pimsner algebra is simple if and only if the C^* -correspondence is nonperiodic and A has no nontrivial invariant ideals) to the simplicity result for graph C^* -algebras (i.e., a graph C^* -algebra is simple if and only if the graph satisfies Condition (L) and the graph has no nontrivial saturated hereditary subsets), and using the fact that invariant ideals in A correspond to saturated hereditary subsets in the graph, it is natural to conjecture that nonperiodic is the appropriate analogue of Condition (L) for C^* -correspondences. However, we shall show in this paper that this is not the case. In particular, in Section 6 we show that even under the hypotheses of Schweizer's simplicity theorem there are graphs that do not satisfy Condition (L) but whose C^* -correspondences are nonperiodic. Therefore nonperiodic is an insufficient condition for formulating a general Cuntz-Krieger Uniqueness Theorem for Cuntz-Pimsner algebras.

In this paper we make some of the first progress in establishing an analogue of Condition (L) for C^* -correspondences that do not necessarily come from underlying objects such as graphs, topological graphs, or quivers. In particular, in Section 3 we

define a property of C^* -correspondences that we call Condition (S). We use Condition (S) to prove a generalization of the Cuntz-Krieger Uniqueness Theorem (see Theorem 3.9 and Corollary 3.10) for Cuntz-Pimsner algebras of C^* -correspondences with injective left action: If X satisfies Condition (S) and $\rho : \mathcal{O}_X \rightarrow B$ is a $*$ -homomorphism with $\rho|_{\pi_A(A)}$ injective, then ρ is injective. In Section 4 we use this theorem to deduce sufficient conditions for simplicity of Cuntz-Pimsner algebras; in particular, we show in Theorem 4.3 that if X satisfies Condition (S) and A has no nontrivial invariant ideals, then \mathcal{O}_X is simple. We mention that Condition (S) implicitly implies that the left action of A on X is injective, but other than this assumption of injective left action our theorems hold without imposing any additional hypotheses on X or A . As a consequence, our simplicity theorem applies to substantially broader classes of C^* -correspondences than Schweizer’s simplicity theorem or other existing simplicity results.

In Section 5 we prove that if \mathcal{Q} is a topological quiver with no sinks and $X(\mathcal{Q})$ is the associated C^* -correspondence, then $X(\mathcal{Q})$ satisfies Condition (S) if and only if \mathcal{Q} satisfies Condition (L). (The corresponding results for topological graphs and graphs follow as special cases.) This gives additional credibility to Condition (S) as an appropriate analogue of Condition (L).

In Section 6 we examine Schweizer’s nonperiodic condition in more detail to understand its relationship with Condition (S) and also Condition (L). We produce examples showing that even for finite graphs with no sinks and no sources, nonperiodic is distinct from Condition (L). We also provide examples showing that if any of the hypotheses of Schweizer’s theorem are removed, the conclusion of the theorem need not hold. This shows that a condition other than “nonperiodic” is needed to characterize simplicity of Cuntz-Pimsner algebras when any of the hypotheses of Schweizer’s theorem are removed. The conclusions we draw from our examples are summarized in Remark 6.21.

We end the paper with two open questions related to Condition (S), outlining what remains to be accomplished to develop precisely the correct analogue of Condition (L) for all C^* -correspondences.

2. PRELIMINARIES

Let A be a C^* -algebra, and let X be a (right) Hilbert A -module. Let $\langle \cdot, \cdot \rangle_A$ denote the A -valued inner product on X . We let $\mathcal{L}(X)$ denote the C^* -algebra of adjointable operators on X , and we let $\mathcal{K}(X)$ denote the closed two-sided ideal of compact operators given by

$$\mathcal{K}(X) := \overline{\text{span}}\{\Theta_{\xi,\eta} : \xi, \eta \in X\}$$

where $\Theta_{\xi,\eta}$ is defined by $\Theta_{\xi,\eta}(\zeta) := \xi\langle\eta, \zeta\rangle_A$.

Definition 2.1. Let A be a C^* -algebra. A C^* -correspondence over A is a (right) Hilbert A -module X together with a $*$ -homomorphism $\phi_X : A \rightarrow \mathcal{L}(X)$. We consider ϕ_X as providing a left action of A on X by $a \cdot x := \phi_X(a)x$.

Remark 2.2. When we say that X is a C^* -correspondence over a C^* -algebra A , note that we implicitly have both an A -valued inner product $\langle \cdot, \cdot \rangle_A$ and a left action $\phi_X : A \rightarrow \mathcal{L}(X)$, even if $\langle \cdot, \cdot \rangle_A$ and ϕ_X are not explicitly specified. When no confusion arises (e.g., when we have a single C^* -algebra A and a single C^* -correspondence X) we shall often drop the subscripts on both the inner product and the left action and simply write $\langle \cdot, \cdot \rangle := \langle \cdot, \cdot \rangle_A$ and $\phi := \phi_X$ for ease of notation.

Definition 2.3. Let A be a C^* -algebra and let X and Y be C^* -correspondences over A . A C^* -correspondence isomorphism from X to Y is a bijective map $\Phi : X \rightarrow Y$ such that

- (i) $\Phi(a \cdot x) = a \cdot \Phi(x)$, and
- (ii) $\langle \Phi(x), \Phi(x') \rangle = \langle x, x' \rangle$,

for all $x, x' \in X$ and $a \in A$. We say X and Y are *isomorphic* (or, for emphasis, *isomorphic as C^* -correspondences*) when such an isomorphism exists. Also note that any C^* -correspondence isomorphism automatically preserves the right action (i.e., $\Phi(xa) = \Phi(x)a$ for $x \in X$ and $a \in A$) due to (ii).

Definition 2.4. If X is a C^* -correspondence over a C^* -algebra A , a *representation* of X into a C^* -algebra B is a pair (ψ, π) consisting of a linear map $\psi : X \rightarrow B$ and a $*$ -homomorphism $\pi : A \rightarrow B$ satisfying

- (i) $\psi(\xi)^* \psi(\eta) = \pi(\langle \xi, \eta \rangle_A)$
- (ii) $\psi(\phi(a)\xi) = \pi(a)\psi(\xi)$
- (iii) $\psi(\xi \cdot a) = \psi(\xi)\pi(a)$

for all $\xi, \eta \in X$ and $a \in A$. We often write $(\psi, \pi) : (X, A) \rightarrow B$ to denote this representation, and we let $C^*(\psi, \pi)$ denote the C^* -subalgebra of B generated by $\psi(X) \cup \pi(A)$. For a representation $(\psi, \pi) : (X, A) \rightarrow B$ the relations (i)–(ii) above imply

$$C^*(\pi, t) = \overline{\text{span}}\{\psi^n(\xi)\psi^m(\eta)^* : \xi \in X^{\otimes n}, \eta \in X^{\otimes m}, n, m \geq 0\}.$$

A representation (ψ, π) is said to be *injective* if π is injective. Note that in this case ψ will be isometric (and hence also injective) since

$$\|\psi(\xi)\|^2 = \|\psi(\xi)^* \psi(\xi)\| = \|\pi(\langle \xi, \xi \rangle_A)\| = \|\langle \xi, \xi \rangle_A\| = \|\xi\|^2.$$

Definition 2.5. For a representation $(\psi, \pi) : (X, A) \rightarrow B$ there exists a $*$ -homomorphism $\pi^{(1)} : \mathcal{K}(X) \rightarrow B$ with the property that

$$\pi^{(1)}(\Theta_{\xi, \eta}) = \psi(\xi)\psi(\eta)^*.$$

See [13, p. 202], [4, Lemma 2.2], and [3, Remark 1.7] for details on the existence of this $*$ -homomorphism. Also note that if (ψ, π) is an injective representation, then $\pi^{(1)}$ is injective as well [3, Proposition 1.6(2)].

Definition 2.6. For an ideal I in a C^* -algebra A we define

$$I^\perp := \{a \in A : ab = 0 \text{ for all } b \in I\}$$

and we refer to I^\perp as the *annihilator of I in A* . If X is a C^* -correspondence over A , we define an ideal $J(X)$ of A by

$$J(X) := \phi^{-1}(\mathcal{K}(X)).$$

We also define an ideal J_X of A by

$$J_X := J(X) \cap (\ker \phi)^\perp.$$

Note that $J_X = J(X)$ when ϕ is injective, and that J_X is the maximal ideal of $J(X)$ on which the restriction of ϕ is an injection into $\mathcal{K}(X)$.

Definition 2.7. If X is a C^* -correspondence over a C^* -algebra A , we say that a representation $(\psi, \pi) : (X, A) \rightarrow B$ is *covariant* if

$$\pi^{(1)}(\phi(a)) = \pi(a) \quad \text{for all } a \in J_X.$$

We say that a covariant representation $(\psi_X, \pi_A) : (X, A) \rightarrow B$ is *universal* if whenever $(\psi, \pi) : (X, A) \rightarrow C$ is a covariant representation there exists a $*$ -homomorphism $\rho_{(\psi, \pi)} : C^*(\psi_X, \pi_A) \rightarrow C$ with the property that $\psi = \rho_{(\psi, \pi)} \circ \psi_X$ and $\pi = \rho_{(\psi, \pi)} \circ \pi_A$.

Definition 2.8. If X is a C^* -correspondence over a C^* -algebra A , the *Cuntz-Pimsner algebra*, which we shall denote \mathcal{O}_X , is the C^* -algebra $C^*(\psi_X, \pi_A)$ where (ψ_X, π_A) is a universal covariant representation of X . The existence of \mathcal{O}_X is established in [2, Proposition 1.3], and the universal property ensures \mathcal{O}_X is unique up to isomorphism.

Remark 2.9. The universal property of a Cuntz-Pimsner algebra ensures the existence of a canonical circle action, which is known as the *gauge action*. Let X be a C^* -correspondence over a C^* -algebra A . If \mathcal{O}_X is the associated Cuntz-Pimsner algebra and $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ is a universal covariant representation, then for any $z \in \mathbb{T}$ we have that $(z\psi_X, \pi_A)$ is also a universal covariant representation. Hence by the universal property, there exists a homomorphism $\gamma_z : \mathcal{O}_X \rightarrow \mathcal{O}_X$ such that $\gamma_z(\pi_A(a)) = \pi_A(a)$ for all $a \in A$ and $\gamma_z(\psi_X(\xi)) = z\psi_X(\xi)$ for all $\xi \in X$. Since $\gamma_{z^{-1}}$ is an inverse for this homomorphism, we see that γ_z is an automorphism. Thus we have an action $\gamma : \mathbb{T} \rightarrow \text{Aut } \mathcal{O}_X$ with the property that $\gamma_z(\pi_A(a)) = \pi_A(a)$ for all $a \in A$ and $\gamma_z(\psi_X(\xi)) = z\psi_X(\xi)$ for all $\xi \in X$. Furthermore, a routine $\epsilon/3$ -argument shows that γ is strongly continuous. We call γ the *gauge action on \mathcal{O}_X* .

Because \mathbb{T} is compact, averaging over γ with respect to normalized Haar measure gives a conditional expectation E of \mathcal{O}_X onto the fixed point algebra \mathcal{O}_X^γ by

$$E(x) := \int_{\mathbb{T}} \gamma_z(x) dz \quad \text{for } x \in \mathcal{O}_X,$$

where the integral denotes vector-valued integration in \mathcal{O}_X and dz denotes the normalized Haar measure on \mathbb{T} . The map E is completely positive, has norm 1, and is faithful in the sense that $E(a^*a) = 0$ implies $a = 0$.

Remark 2.10. Let X be a C^* -correspondence over a C^* -algebra A . Recalling that there is $*$ -homomorphism $\phi_X : A \rightarrow \mathcal{L}(X)$ providing a left action of A on X by $a \cdot x := \phi_X(a)(x)$, we let $X^{\otimes n} := X \otimes_A \dots \otimes_A X$ denote the n -fold A -balanced tensor product of X with itself. We note that $X^{\otimes n}$ is a C^* -correspondence over A with right action $(x_1 \otimes \dots \otimes x_n) \cdot a = x_1 \otimes \dots \otimes x_n \cdot a$, left action $a \cdot (x_1 \otimes \dots \otimes x_n) = a \cdot x_1 \otimes \dots \otimes x_n$, and inner product defined inductively by

$$\langle x_1 \otimes \dots \otimes x_n, y_1 \otimes \dots \otimes y_n \rangle := \langle x_2 \otimes \dots \otimes x_n, \langle x_1, y_1 \rangle \cdot y_2 \otimes \dots \otimes y_n \rangle.$$

If $(\psi, \pi) : (X, A) \rightarrow B$ is a representation of X in a C^* -algebra B , then it is shown in [2, Lemma 3.6] that for each $n \in \mathbb{N}$ there is a linear map $\psi^{\otimes n} : X^{\otimes n} \rightarrow B$ satisfying

$$\psi^{\otimes n}(x_1 \otimes \dots \otimes x_n) = \psi(x_1) \dots \psi(x_n) \quad \text{for } x_1, \dots, x_n \in X.$$

Furthermore, $(\psi^{\otimes n}, \pi) : (X^{\otimes n}, A) \rightarrow B$ is a representation and there exists a $*$ -homomorphism $\pi^{(n)} : \mathcal{K}(X^{\otimes n}) \rightarrow B$ such that

$$\pi^{(n)}(\Theta_{x,y}) = \psi^{\otimes n}(x)\psi^{\otimes n}(y)^* \quad \text{for } x, y \in X^{\otimes n}.$$

Suppose X is a C^* -correspondence over a C^* -algebra A , and suppose that the left action $\phi_X : A \rightarrow \mathcal{L}(X)$ is injective. If $(\psi, \pi) : (X, A) \rightarrow B$ is a covariant

representation, we let 1^k denote the identity operator on $X^{\otimes k}$ and define

$$\mathcal{C}_n := A \otimes 1^n + \mathcal{K}(X) \otimes 1^{n-1} + \mathcal{K}(X^{\otimes 2}) \otimes 1^{n-2} + \dots + \mathcal{K}(X^{\otimes n}) \subseteq \mathcal{L}(X^{\otimes n}).$$

Furthermore, the homomorphism $\kappa_n^{\psi, \pi} : \mathcal{C}_n \rightarrow B$ given by

$$\kappa_n^{\psi, \pi}(k_0 \otimes 1^n + k_1 \otimes 1^{n-1} + \dots + k_n) := \pi(k_0) + \pi^{(1)}(k_1) + \dots + \pi^{(n)}(k_n)$$

is injective whenever π is injective [2, Proposition 4.6]. It is also a fact that if $\mathcal{C} := \varinjlim \mathcal{C}_n$ under the isometric homomorphisms $c \in \mathcal{C}_n \mapsto c \otimes 1 \in \mathcal{C}_{n+1}$, then the homomorphisms $\kappa_n^{\psi, \pi} : \mathcal{C}_n \rightarrow B$ induce a map $\kappa^{\psi, \pi} : \mathcal{C} \rightarrow B$, and [2, Corollary 4.8] shows that $\kappa^{\psi, \pi}$ is injective when π is injective. Furthermore, [2, Corollary 4.9] shows that for a universal covariant representation $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ we have that $\kappa^{\psi_X, \pi_A} : \mathcal{C} \rightarrow \mathcal{O}_X$ is an injective map onto the fixed point algebra \mathcal{O}_X^γ .

3. A CUNTZ-KRIEGER UNIQUENESS THEOREM FOR CUNTZ-PIMSNER ALGEBRAS

Definition 3.1. Let X be a C^* -correspondence over A , and let $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ be a universal covariant representation of (X, A) into \mathcal{O}_X . We define a vector $\zeta \in X^{\otimes m}$ with $m \in \mathbb{N}$ to be *nonreturning* if whenever $n \in \{1, \dots, m-1\}$ and $\xi \in X^{\otimes n}$, then

$$\psi_X^{\otimes m}(\zeta)^* \psi_X^{\otimes n}(\xi) \psi_X^{\otimes m}(\zeta) = 0.$$

Remark 3.2. The properties and nomenclature in our definition of a nonreturning vector are inspired by the Katsura's notion of a nonreturning path in a topological graph (see [5, Definition 5.5]). Katsura shows in [5, Lemma 5.6] and [5, Lemma 5.7] that a nonreturning path in a topological graph can be used to construct a compactly supported continuous function that is a nonreturning vector in the sense of our Definition 3.1. (Muhly and the second author have also shown in [11, §6] that Katsura's results on nonreturning paths can be generalized to topological quivers.)

Lemma 3.3. *Let X be a C^* -correspondence over A and let $\zeta \in X^{\otimes m}$ be a nonreturning vector. If $(\psi, \pi) : (X, A) \rightarrow B$ is any covariant representation, then for any $n \in \{1, \dots, m-1\}$ and $\xi \in X^{\otimes n}$, we have*

$$\psi^{\otimes m}(\zeta)^* \psi^{\otimes n}(\xi) \psi^{\otimes m}(\zeta) = 0.$$

Proof. Let $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ be a universal covariant representation of (X, A) into \mathcal{O}_X . If $\rho := \rho_{(\psi, \pi)} : \mathcal{O}_X \rightarrow B$ denotes the $*$ -homomorphism induced by (ψ, π) , then

$$\begin{aligned} \psi^{\otimes m}(\zeta)^* \psi^{\otimes n}(\xi) \psi^{\otimes m}(\zeta) &= (\rho \circ \psi_X^{\otimes m})(\zeta)^* (\rho \circ \psi_X^{\otimes n})(\xi) (\rho \circ \psi_X^{\otimes m})(\zeta) \\ &= \rho(\psi_X^{\otimes m}(\zeta)^* \psi_X^{\otimes n}(\xi) \psi_X^{\otimes m}(\zeta)) = \rho(0) = 0. \quad \square \end{aligned}$$

Condition (S): We say that a C^* -correspondence X over a C^* -algebra A satisfies Condition (S) if whenever $a \in A$ with $a \geq 0$, then for all $n \in \mathbb{N}$ and for all $\epsilon > 0$ there exists $m > n$ and a nonreturning vector $\zeta \in X^{\otimes m}$ with $\|\zeta\| = 1$ and

$$\|\langle \zeta, a\zeta \rangle\| > \|a\| - \epsilon.$$

Remark 3.4. The (S) in Condition (S) is meant to stand for ‘‘simplicity’’.

Proposition 3.5. *If X is a C^* -correspondence over A satisfying Condition (S), then the left action $\phi_X : A \rightarrow \mathcal{L}(X)$ is injective.*

Proof. For each $n \in \mathbb{N}$ let $\phi_n : A \rightarrow \mathcal{L}(X^{\otimes n})$ denote the $*$ -homomorphism defined by $\phi_n(a)(x_1 \otimes \dots \otimes x_n) := a \cdot x_1 \otimes \dots \otimes x_n$. Also define $h : \mathcal{L}(X^{\otimes n}) \rightarrow \mathcal{L}(X^{\otimes n+1})$ by $h(T) := T \otimes I$. Then h is a $*$ -homomorphism, and hence $\|h(T)\| \leq \|T\|$ for all $T \in \mathcal{L}(X^{\otimes n})$. Furthermore, for any $a \in A$ and $n \in \mathbb{N}$, we have $h(\phi_n(a)) = \phi_{n+1}(a)$. Thus $\|\phi_{n+1}(a)\| \leq \|\phi_n(a)\|$. By induction, we conclude that

$$\|\phi_n(a)\| \leq \|\phi_1(a)\| = \|\phi_X(a)\| \quad \text{for all } a \in A \text{ and } n \in \mathbb{N}.$$

For any $b \in A$, let $a := b^*b \geq 0$. For any $\epsilon > 0$, Condition (S) implies there exists $m \in \mathbb{N}$ and a nonreturning vector $\zeta \in X^{\otimes m}$ with $\|\zeta\| = 1$ and

$$\|\langle \zeta, \phi_m(a)\zeta \rangle\| > \|a\| - \epsilon.$$

Since $\|\zeta\| = 1$, we have

$$\begin{aligned} \|\phi_X(b)\|^2 &= \|\phi_1(b)\|^2 \geq \|\phi_m(b)\|^2 \geq \|\phi_m(b)\zeta\|^2 = \|\langle \phi_m(b)\zeta, \phi_m(b)\zeta \rangle\| \\ &= \|\langle \zeta, \phi_m(b^*b)\zeta \rangle\| = \|\langle \zeta, \phi_m(a)\zeta \rangle\| > \|a\| - \epsilon = \|b^*b\| - \epsilon = \|b\|^2 - \epsilon. \end{aligned}$$

Since $\epsilon > 0$ was arbitrary, we deduce $\|\phi_X(b)\|^2 \geq \|b\|^2$ and $\|\phi_X(b)\| \geq \|b\|$. Moreover, because ϕ is a $*$ -homomorphism, and hence norm decreasing, we conclude $\|\phi_X(b)\| = \|b\|$, so that ϕ_X is isometric and hence also injective. \square

Remark 3.6. Proposition 3.5 shows that any C^* -correspondence satisfying Condition (S) necessarily has injective left action. The reader should keep this in mind, and note that in all forthcoming results the hypothesis that a C^* -correspondence satisfies Condition (S) implicitly requires the C^* -correspondence to have injective left action.

Lemma 3.7. *Let X be a C^* -correspondence over a C^* -algebra A , and suppose that X satisfies Condition (S). Let $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ be a universal covariant representation of X in \mathcal{O}_X . Also let $(\psi, \pi) : (X, A) \rightarrow B$ be a covariant representation of X into a C^* -algebra B with the property that $\pi : A \rightarrow B$ is injective. If*

$$x = \sum_{i=1}^N \psi_X^{\otimes n_i}(\xi_i) \psi_X^{\otimes m_i}(\eta_i)^*$$

is positive in \mathcal{O}_X , then

$$y_0 := \sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^*$$

is positive in B , and if $\rho := \rho_{(\psi, \pi)} : \mathcal{O}_X \rightarrow B$ denotes the $$ -homomorphism induced by (ψ, π) , then for all $\epsilon > 0$ there exists $a \in A$ and $b \in B$ with $\|a\| = \|y_0\|$, $\|b\| \leq 1$, and*

$$\|b^* \rho(x) b - \pi(a)\| < \epsilon.$$

Proof. Let $E : \mathcal{O}_X \rightarrow \mathcal{O}_X^\gamma$ denote the conditional expectation determined by the gauge action γ on \mathcal{O}_X and given by $E(x) := \int_{\mathbb{T}} \gamma_z(x) dz$. Then

$$\begin{aligned} y_0 &= \sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* = \sum_{n_i=m_i} (\rho \circ \psi_X)^{\otimes n_i}(\xi_i) (\rho \circ \psi_X)^{\otimes m_i}(\eta_i)^* \\ &= \rho \left(\sum_{n_i=m_i} \psi_X^{\otimes n_i}(\xi_i) \psi_X^{\otimes m_i}(\eta_i)^* \right) = \rho(E(x)). \end{aligned}$$

Since E is a conditional expectation, E is completely positive (as well as positive). Likewise, the $*$ -homomorphism ρ is positive. Since x is positive in \mathcal{O}_X , it follows that $y_0 = \rho(E(x))$ is positive in B .

Given $\epsilon > 0$, set $n := \max\{n_1, \dots, n_N, m_1, \dots, m_N\}$. By [2, Proposition 4.6] the $*$ -homomorphism $\kappa_n^{\psi, \pi} : \mathcal{C}_n \rightarrow B$ is injective and hence isometric. Therefore, since $y_0 = \sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* \in \text{im } \kappa_n^{\psi, \pi}$, there exists $T \in \mathcal{C}_n \subseteq \mathcal{L}(X^{\otimes n})$ with $\kappa_n^{\psi, \pi}(T) = y_0$ and $\|T\| = \|y_0\|$. Moreover, since y_0 is positive in B , we may choose T positive in \mathcal{C}_n : since $y_0 \in \text{im } \kappa_n^{\psi, \pi} \subseteq B$ is positive in B , and hence positive in $\text{im } \kappa_n^{\psi, \pi}$, we may write $y_0 = v^*v$ for $v \in \text{im } \kappa_n^{\psi, \pi}$, choose $S \in \mathcal{C}_n$ with $\kappa_n^{\psi, \pi}(S) = v$, and let $T := S^*S \geq 0$. Then $\kappa_n^{\psi, \pi}(T) = \kappa_n^{\psi, \pi}(S^*S) = \kappa_n^{\psi, \pi}(S)^* \kappa_n^{\psi, \pi}(S) = v^*v = y_0$.

Since $\|T\| = \|y_0\|$ and T is positive, there exists $\xi \in X^{\otimes n}$ with $\|\xi\| = 1$ such that $g := \langle \xi, T\xi \rangle$ is a positive element of A with $\|g\| > \|y_0\| - \epsilon/2$. When $n_i > m_i$ we have

$$\psi^{\otimes n}(\xi)^* \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* \psi^{\otimes n}(\xi) = \psi^{\otimes n'_i}(\xi'_i)$$

for some $\xi'_i \in X^{\otimes n'_i}$, where $n'_i := n_i - m_i$. Similarly, when $n_i < m_i$ we have

$$\psi^{\otimes n}(\xi)^* \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* \psi^{\otimes n}(\xi) = \psi^{\otimes m'_i}(\eta'_i)^*$$

for some $\eta'_i \in X^{\otimes m'_i}$, where $m'_i := m_i - n_i$. Also, when $n_i = m_i$ we have

$$\begin{aligned} \psi^{\otimes n}(\xi)^* \left(\sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* \right) \psi^{\otimes n}(\xi) &= \psi^{\otimes n}(\xi)^* y_0 \psi^{\otimes n}(\xi) \\ &= \psi^{\otimes n}(\xi)^* \kappa_n^{\psi, \pi}(T) \psi^{\otimes n}(\xi) = \psi^{\otimes n}(\xi)^* \psi^{\otimes n}(T\xi) = \pi(\langle \xi, T\xi \rangle) = \pi(g). \end{aligned}$$

Combining these three equations yields

$$\begin{aligned} (3.1) \quad \psi^{\otimes n}(\xi)^* \rho(x) \psi^{\otimes n}(\xi) &= \psi^{\otimes n}(\xi)^* \left(\sum_{i=1}^N \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* \right) \psi^{\otimes n}(\xi) \\ &= \pi(g) + \sum_{n_i > m_i} \psi^{\otimes n'_i}(\xi'_i) + \sum_{n_i < m_i} \psi^{\otimes m'_i}(\eta'_i)^*. \end{aligned}$$

Since $g \geq 0$, by Condition (S) there exists $m > n$ and a nonreturning path $\zeta \in X^{\otimes m}$ with $\|\zeta\| = 1$ and $\|\langle \zeta, g\zeta \rangle\| > \|g\| - \epsilon/2$.

Let $c := \langle \zeta, g\zeta \rangle \in A$. Then

$$\|c\| = \|\langle \zeta, g\zeta \rangle\| > \|g\| - \epsilon/2 > (\|y_0\| - \epsilon/2) - \epsilon/2 = \|y_0\| - \epsilon.$$

In addition,

$$\begin{aligned} \|c\| &= \|\langle \zeta, g\zeta \rangle\| \leq \|g\| \|\langle \zeta, \zeta \rangle\| = \|g\| = \|\langle \xi, T\xi \rangle\| \\ &\leq \|T\| \|\langle \zeta, \zeta \rangle\| = \|T\| = \|y_0\| < \|y_0\| + \epsilon. \end{aligned}$$

These two inequalities show that

$$\left| \|c\| - \|y_0\| \right| < \epsilon.$$

Let $b := \psi^{\otimes n}(\xi) \psi^{\otimes m}(\zeta)$. Then $\|b\| \leq \|\psi^{\otimes n}(\xi)\| \|\psi^{\otimes m}(\zeta)\| = \|\xi\| \|\zeta\| = 1 \cdot 1 = 1$. In addition,

$$\begin{aligned} b^* \rho(x) b &= \psi^{\otimes m}(\zeta)^* \psi^{\otimes n}(\xi)^* \rho(x) \psi^{\otimes n}(\xi) \psi^{\otimes m}(\zeta) \\ &= \psi^{\otimes m}(\zeta)^* \left(\pi(g) + \sum_{n_i > m_i} \psi^{\otimes n'_i}(\xi'_i) + \sum_{n_i < m_i} \psi^{\otimes m'_i}(\eta'_i)^* \right) \psi^{\otimes m}(\zeta) \text{ (by (3.1))} \end{aligned}$$

$$\begin{aligned}
 &= \psi^{\otimes m}(\zeta)^* \pi(g) \psi^{\otimes m}(\zeta) \quad (\text{since } \zeta \text{ is a nonreturning vector}) \\
 &= \psi^{\otimes m}(\zeta)^* \psi^{\otimes m}(g\zeta) \\
 &= \pi(\langle \zeta, g\zeta \rangle) \\
 &= \pi(c).
 \end{aligned}$$

Let $a := \frac{\|y_0\|}{\|c\|}c$. Then $\|a\| = \|y_0\|$, and

$$\begin{aligned}
 \|b^* \rho(x)b - \pi(a)\| &= \|\pi(c) - \pi(a)\| = \|\pi(c - a)\| \leq \|c - a\| = \left\| c - \frac{\|y_0\|}{\|c\|}c \right\| \\
 &= \left| 1 - \frac{\|y_0\|}{\|c\|} \right| \|c\| = \left| \|c\| - \|y_0\| \right| < \epsilon. \quad \square
 \end{aligned}$$

Lemma 3.8. *Let A_0 be a dense $*$ -subalgebra of a C^* -algebra A . If $x \in A$ and $x \geq 0$, then for every $\epsilon > 0$ there exists $x_0 \in A_0$ with $x_0 \geq 0$ and $\|x - x_0\| < \epsilon$.*

Proof. If $x = 0$, the result is trivial, so assume $x \neq 0$. Since $x \geq 0$, we may write $x = v^*v$ for some nonzero $v \in A$. Using the fact that A_0 is dense in A , choose $v_0 \in A_0$ with

$$\|v - v_0\| \leq \min \left\{ \frac{\epsilon}{3\|v\|}, \sqrt{\frac{\epsilon}{3}} \right\}.$$

Let $x_0 := v_0^*v_0 \in A_0$. Then $x_0 \geq 0$, and

$$\begin{aligned}
 \|x - x_0\| &= \|v^*v - v_0^*v_0\| \leq \|v^*v - v^*v_0\| + \|v^*v_0 - v_0^*v_0\| \\
 &\leq \|v^*\| \|v - v_0\| + \|v^* - v_0^*\| \|v_0\| = \|v\| \|v - v_0\| + \|v - v_0\| \|v_0\| \\
 &\leq \|v\| \frac{\epsilon}{3\|v\|} + \|v - v_0\| \|v_0 - v + v\| \leq \epsilon/3 + \|v - v_0\| (\|v_0 - v\| + \|v\|) \\
 &\leq \epsilon/3 + \|v - v_0\|^2 + \|v\| \|v - v_0\| \leq \epsilon/3 + \left(\sqrt{\frac{\epsilon}{3}} \right)^2 + \|v\| \frac{\epsilon}{3\|v\|} \\
 &= \epsilon/3 + \epsilon/3 + \epsilon/3 = \epsilon. \quad \square
 \end{aligned}$$

We are now in a position to prove our main result.

Theorem 3.9 (Cuntz-Krieger Uniqueness for Cuntz-Pimsner Algebras). *Let X be a C^* -correspondence over a C^* -algebra A , and suppose that X satisfies Condition (S). Also let $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ be a universal covariant representation of X into \mathcal{O}_X . If $\rho : \mathcal{O}_X \rightarrow B$ is a $*$ -homomorphism from \mathcal{O}_X into a C^* -algebra B with the property that the restriction $\rho|_{\pi_A(A)}$ is injective, then ρ is injective.*

Proof. Let γ denote the gauge action on \mathcal{O}_X , and let $E : \mathcal{O}_X \rightarrow \mathcal{O}_X^\gamma$ denote the conditional expectation given by

$$E(x) := \int_{\mathbb{T}} \gamma_z(x) dz.$$

We shall show the following statements hold:

- (a) ρ is injective on the fixed point algebra \mathcal{O}_X^γ , and
- (b) $\|\rho(E(x))\| \leq \|\rho(x)\|$ whenever $x \in \mathcal{O}_X$ and $x \geq 0$.

To see (a), let $\psi := \rho \circ \psi_X$ and $\pi := \rho \circ \pi_A$. Then $\rho = \rho_{(\psi, \pi)}$; that is, ρ equal to the $*$ -homomorphism induced by the representation (ψ, π) . For each $n \in \mathbb{N}$, we have $\kappa_n^{\psi, \pi} = \rho \circ \kappa_n^{\psi_X, \pi_A}$, and hence $\kappa^{\psi, \pi} = \rho \circ \kappa^{\psi_X, \pi_A}$. Since $\kappa^{\psi, \pi}$ is injective by [2, Corollary 4.8], it follows that ρ is injective on $\text{im } \kappa^{\psi_X, \pi_A} = \mathcal{O}_X^\gamma$, so (a) holds.

To see (b), note that since

$$\mathcal{O}_X = \overline{\text{span}}\{\psi_X^{\otimes n}(\xi)\psi_X^{\otimes m}(\eta)^* : \xi \in X^{\otimes n}, \eta \in X^{\otimes m}, \text{ and } n, m \geq 0\},$$

Lemma 3.8 implies that any positive element in \mathcal{O}_X may be approximated by a positive element from $\text{span}\{\psi_X^{\otimes n}(\xi)\psi_X^{\otimes m}(\eta)^* : \xi \in X^{\otimes n}, \eta \in X^{\otimes m}, \text{ and } n, m \geq 0\}$. Hence it suffices to prove (b) when x is positive and has the form

$$x = \sum_{i=1}^N \psi_X^{\otimes n_i}(\xi_i)\psi_X^{\otimes m_i}(\eta_i)^*.$$

Keeping the notation $(\psi, \pi) = (\rho \circ \psi_X, \rho \circ \pi_A)$, if we let

$$y := \rho(x) = \sum_{i=1}^N \psi^{\otimes n_i}(\xi_i)\psi^{\otimes m_i}(\eta_i)^*$$

and

$$y_0 := \sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i)\psi^{\otimes m_i}(\eta_i)^*,$$

then Lemma 3.7 shows that for all $\epsilon > 0$ there exists $a \in A$ and $b \in B$ such that $\|a\| = \|y_0\|$, $\|b\| \leq 1$, and $\|b^*yb - \pi(a)\| < \epsilon$. Thus

$$\begin{aligned} \|\rho(E(x))\| &= \left\| \rho \left(\sum_{n_i=m_i} \psi_X^{\otimes n_i}(\xi_i)\psi_X^{\otimes m_i}(\eta_i)^* \right) \right\| = \left\| \sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i)\psi^{\otimes m_i}(\eta_i)^* \right\| \\ &= \|y_0\| = \|a\| = \|\rho(\pi_A(a))\| = \|\pi(a)\| \leq \|b^*yb\| + \|\pi(a) - b^*yb\| \\ &\leq \|b^*\| \|y\| \|b\| + \epsilon \leq \|y\| + \epsilon = \|\rho(x)\| + \epsilon. \end{aligned}$$

Since this inequality holds for all $\epsilon > 0$, we have that $\|\rho(E(x))\| \leq \|\rho(x)\|$. Hence (b) holds.

Finally, given (a) and (b) we see that whenever $\rho(x) = 0$, we have $\rho(x^*x) = 0$ and (b) implies that $\rho(E(x^*x)) = 0$. Thus $E(x^*x) = 0$ by (a), and since E is a faithful conditional expectation, this implies $x = 0$. Thus ρ is injective. \square

Corollary 3.10. *Let X be a C^* -correspondence over a C^* -algebra A , and suppose that X satisfies Condition (S). If $(\psi, \pi) : (X, A) \rightarrow B$ is a covariant representation with the property that π is injective, then the induced $*$ -homomorphism $\rho_{(\psi, \pi)} : \mathcal{O}_X \rightarrow B$ is injective.*

4. SIMPLICITY OF CUNTZ-PIMSNER ALGEBRAS

All ideals in C^* -algebras that we consider will be closed and two-sided. Let X be a C^* -correspondence over A . For any ideal I of A , we define

$$IX := \overline{\text{span}}\{ax : a \in I \text{ and } x \in X\}$$

and

$$XI := \overline{\text{span}}\{xa : a \in I \text{ and } x \in X\}.$$

We also define

$$X_I := \{x \in X : \langle y, x \rangle \in I \text{ for all } y \in X\}.$$

Lemma 4.1. *Let X be a C^* -correspondence over A . For any ideal I of A we have $XI = X_I$.*

Proof. Since X_I has a right Hilbert I -module structure, by the Cohen-Hewitt factorization theorem we have $X_I = X_I I$, which is a subset of XI .

For the reverse inclusion, let $\xi \in XI := \overline{\text{span}}\{xa : a \in I \text{ and } x \in X\}$. Then $\xi = \lim_{n \rightarrow \infty} \sum_{i=1}^{N_n} x_i^n a_i^n$ with each $x_i^n \in X$ and each $a_i^n \in I$. For any $y \in X$ we have

$$\langle y, \xi \rangle = \langle y, \lim_{n \rightarrow \infty} \sum_{i=1}^{N_n} x_i^n a_i^n \rangle = \lim_{n \rightarrow \infty} \langle y, \sum_{i=1}^{N_n} x_i^n a_i^n \rangle = \lim_{n \rightarrow \infty} \sum_{i=1}^{N_n} \langle y, x_i^n \rangle a_i^n \in I.$$

Thus $\xi \in X_I$, and $XI \subseteq X_I$. \square

Let X be a C^* -correspondence over a C^* -algebra A . If I is an ideal in A , we say I is *hereditary* if $\phi(I)X \subseteq XI$. We also say I is *saturated* if for all $a \in J_X$, $\phi(a)X \subseteq XI$ implies $a \in I$. We call an ideal *invariant* when it is both saturated and hereditary.

The following lemma is well-known among researchers studying Cuntz-Pimsner algebras. It can be extracted as a special case of [7, Proposition 8.2], but doing so requires understanding of definitions and preliminary results in [7]. For the convenience of the reader, and the benefit of the Cuntz-Pimsner literature, we provide a direct and self-contained proof for our special case.

Lemma 4.2. *Let X be a C^* -correspondence over a C^* -algebra A , and suppose that $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ is a universal covariant representation. If \mathcal{I} is an ideal in \mathcal{O}_X , then $\pi_A^{-1}(\mathcal{I})$ is an invariant ideal in A .*

Proof. Let $I := \pi_A^{-1}(\mathcal{I})$. The fact that I is an ideal in A follows from the fact π_A is a $*$ -homomorphism. If $a \in I$ and $x \in X$, then for any $y \in X$ we have

$$\pi_A(\langle y, \phi(a)x \rangle) = \psi_X(y)^* \psi_X(ax) = \psi_X(y)^* \pi_A(a) \psi_X(x) \in \mathcal{I}.$$

Thus $\phi(a)x \in X_I = XI$. Hence $\phi(I)X \subseteq XI$, and I is hereditary.

In addition, let $a \in J_X$ with $\phi(a)X \subseteq XI$. Since

$$\begin{aligned} \pi_A^{(1)}(\mathcal{K}(X)) &= \pi_A^{(1)}(\overline{\text{span}}\{\theta_{x,y} : x, y \in X\}) \\ &= \overline{\text{span}}\{\pi_A^{(1)}(\theta_{x,y}) : x, y \in X\} = \overline{\text{span}}\{\psi_X(x)\psi_X(y)^* : x, y \in X\} \end{aligned}$$

we may choose an approximate unit $\{e_\lambda\}_{\lambda \in \Lambda}$ for $\pi^{(1)}(\mathcal{K}(X))$ with

$$e_\lambda := \sum_{i=1}^{N_\lambda} \psi_X(x_i^\lambda) \psi_X(y_i^\lambda)^* \quad \text{for each } \lambda \in \Lambda.$$

Furthermore, since $a \in J_X$, we conclude $\pi_A(a) = \pi^{(1)}(\phi(a)) \in \pi_A(\mathcal{K}(X))$, and

$$\pi_A(a) = \lim_{\lambda} e_\lambda \pi_A(a) e_\lambda.$$

Because $\phi(a)X \subseteq XI = X_I$, for any $x, y \in X$ we have $\phi(a)x \in X_I$ and $\langle y, \phi(a)x \rangle \in I$. Thus for all $x, y \in X$,

$$\psi_X(y)^* \pi_A(a) \psi_X(x) = \pi_A(\langle y, \phi(a)x \rangle) \in \mathcal{I}$$

and since \mathcal{I} is an ideal, it follows that $e_\lambda \pi_A(a) e_\lambda \in \mathcal{I}$ for all $\lambda \in \Lambda$. Hence $\pi_A(a) = \lim_{\lambda} e_\lambda \pi_A(a) e_\lambda \in \mathcal{I}$, so that $a \in I$ and I is saturated. Therefore I is invariant. \square

Theorem 4.3 (Sufficient Conditions for Simplicity of Cuntz-Pimsner Algebras). *Let X be a C^* -correspondence over a C^* -algebra A . If X satisfies Condition (S) and A has no invariant ideals other than A and $\{0\}$, then \mathcal{O}_X is simple.*

Proof. Let $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ be a universal covariant representation. Suppose \mathcal{I} is an ideal in \mathcal{O}_X with $\mathcal{I} \neq \mathcal{O}_X$. Let $I := \pi_A^{-1}(\mathcal{I})$. By Lemma 4.2, I is an invariant ideal of A . Hence our hypothesis implies $I = A$ or $I = \{0\}$. Since $\mathcal{I} \neq \mathcal{O}_X$, we conclude $I \neq A$ and therefore $I = \{0\}$. Define $Q_{\mathcal{I}} : \mathcal{O}_X \rightarrow \mathcal{O}_X/\mathcal{I}$ to be the quotient map $Q_{\mathcal{I}}(x) = x + \mathcal{I}$. Then for any $a \in A$ we have

$$Q_{\mathcal{I}}(\pi_A(a)) = 0 \iff \pi_A(a) + \mathcal{I} = 0 + \mathcal{I} \iff \pi_A(a) \in \mathcal{I} \iff a \in I \iff a = 0.$$

Hence $Q_{\mathcal{I}}|_{\pi_A(A)}$ is injective, and Theorem 3.9 implies $Q_{\mathcal{I}} : \mathcal{O}_X \rightarrow \mathcal{O}_X/\mathcal{I}$ is injective. But the only way this can occur is if $\mathcal{I} = \{0\}$. Hence \mathcal{O}_X is simple. \square

Remark 4.4. Theorem 4.3 gives sufficient conditions for simplicity of a Cuntz-Pimsner algebra. However, at this time we do not know if Condition (S) is necessary for simplicity. If it is, then Condition (S) would be an appropriate generalization of Condition (L) to C^* -correspondences with injective left action.

Remark 4.5 (Comparison of Theorem 4.3 and Schweizer's Simplicity Theorem). Let X be a C^* -correspondence over a C^* -algebra A . Schweizer's Simplicity Theorem states that under the hypotheses that X is full, A is unital, and the left action $\phi_X : A \rightarrow \mathcal{L}(X)$ is injective, the Cuntz-Pimsner algebra \mathcal{O}_X is simple if and only if X is nonperiodic and A has no hereditary ideal other than $\{0\}$ and A .

Our result in Theorem 4.3 shows that X satisfying Condition (S) and A having no invariant ideals implies the Cuntz-Pimsner algebra \mathcal{O}_X is simple. To compare Theorem 4.3 to Schweizer's Simplicity Theorem, let us first consider the difference between the following two hypotheses:

- (1) The only ideals of A that are invariant (i.e., both hereditary and saturated) are $\{0\}$ and A .
- (2) The only ideals of A that are hereditary are $\{0\}$ and A .

One trivially has (2) \implies (1). Thus Theorem 4.3, which is the result

$$\text{Condition (S) and (1)} \implies \mathcal{O}_X \text{ is simple,}$$

is stronger than (and hence implies) the weaker result

$$\text{Condition (S) and (2)} \implies \mathcal{O}_X \text{ is simple.}$$

Schweizer shows that under his hypotheses, simplicity of \mathcal{O}_X is equivalent to “Non-periodic and (1)”. However, when one assumes Schweizer's hypotheses (i.e., X is full, A is unital, and the left action $\phi_X : A \rightarrow \mathcal{L}(X)$ is injective), then (1) and (2) are equivalent. (To see this in the special case of graph algebras, note that Schweizer's three hypotheses are equivalent to the graph having no sources, finitely many vertices, and no sinks, respectively.)

We have chosen to state Theorem 4.3 using the hypothesis (2) for two reasons: First, this formulation produces a stronger sufficiency result. Second, the corresponding simplicity results for C^* -algebras of graph, topological graphs, and topological quivers is “ $C^*(E)$ is simple $\iff E$ satisfies Condition (L) and E^0 has no nontrivial subsets that are both saturated and hereditary”. Our formulation using (2) not only establishes parallelism between these results, but anticipates future simplicity results due to the fact that if necessary and sufficient conditions for simplicity of a Cuntz-Pimsner algebra are found, they will most likely be of the form “some condition on X ” together with “ A has no saturated and hereditary ideals other than $\{0\}$ and A ”.

It remains to compare Schweizer's nonperiodic condition and our Condition (S). This comparison is more involved, and it is the subject of Section 6 in this paper.

5. TOPOLOGICAL QUIVERS AND CONDITION (L)

Definition 5.1. A topological quiver is a quintuple $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ consisting of a second countable locally compact Hausdorff space E^0 (whose elements are called vertices), a second countable locally compact Hausdorff space E^1 (whose elements are called edges), a continuous open map $r : E^1 \rightarrow E^0$, a continuous map $s : E^1 \rightarrow E^0$, and a family of Radon measures $\lambda = \{\lambda_v\}_{v \in E^0}$ on E^1 satisfying the following two conditions:

- (1) $\text{supp } \lambda_v = r^{-1}(v)$ for all $v \in E^0$, and
- (2) $v \mapsto \int_{E^1} \xi(\alpha) d\lambda_v(\alpha)$ is an element of $C_c(E^0)$ for all $\xi \in C_c(E^1)$.

Definition 5.2 (The quiver C^* -correspondence). If $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ is a quiver, we let $A := C_0(E^0)$ and define an A -valued inner product on $C_c(E^1)$ by

$$\langle \xi, \eta \rangle(v) := \int_{r^{-1}(v)} \overline{\xi(\alpha)} \eta(\alpha) d\lambda_v(\alpha) \quad \text{for } v \in E^0 \text{ and } \xi, \eta \in C_c(E^1).$$

We let $X(\mathcal{Q})$ denote the closure of $C_c(E^1)$ with respect to the norm arising from this inner product. We define a right action of A on $X(\mathcal{Q})$ by

$$(\xi \cdot f)(\alpha) := \xi(\alpha) f(r(\alpha)) \quad \text{for } \alpha \in E^1, \xi \in C_c(E^1), \text{ and } f \in C_0(E^0)$$

and extending this action to all of $X(\mathcal{Q})$. We define a left action $\phi : A \rightarrow \mathcal{L}(X(\mathcal{Q}))$ by

$$(\phi(f)\xi)(\alpha) := f(s(\alpha))\xi(\alpha) \quad \text{for } \alpha \in E^1, \xi \in C_c(E^1), \text{ and } f \in C_0(E^0)$$

and extending to all of $X(\mathcal{Q})$. With this inner product and these actions $X(\mathcal{Q})$ is a C^* -correspondence that we call the *quiver C^* -correspondence* of \mathcal{Q} .

Remark 5.3. Although $X(\mathcal{Q}) := \overline{C_c(E^1)}$, as described in [12, Remark 2.2] the elements of $X(\mathcal{Q})$ may be viewed as the continuous sections of a continuous field of Hilbert spaces. In particular, if $\xi \in X(\mathcal{Q})$, then for any $v \in E^0$ the vector ξ determines a function in $L^2(r^{-1}(v), \lambda_v)$. Thus when working with $X(\mathcal{Q})$, the evaluation of the inner product via the integral $\int_{r^{-1}(v)} \overline{\xi(\alpha)} \eta(\alpha) d\lambda_v(\alpha)$ can be performed for all vectors $\xi, \eta \in X(\mathcal{Q})$ and not only those in $C_c(E^1)$.

Definition 5.4 (The quiver C^* -algebra). If $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ is a quiver, let $X(\mathcal{Q})$ denote the quiver C^* -correspondence of \mathcal{Q} . The *quiver C^* -algebra* $C^*(\mathcal{Q})$ is defined to be the Cuntz-Pimsner algebra $\mathcal{O}_{X(\mathcal{Q})}$.

Both topological graph C^* -algebras and graph C^* -algebras are special cases of topological quiver C^* -algebras. Topological graphs and their C^* -algebras are the special case when r is a local homeomorphism and each λ_v is counting measure on $r^{-1}(v)$. Graphs and their C^* -algebras are the special case when E^0 and E^1 both have the discrete topology and each λ_v is counting measure on $r^{-1}(v)$.

If $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ is a topological quiver, we define $E_{\text{sinks}}^0 := E^0 \setminus \overline{s(E^1)}$. An element of E_{sinks}^0 is called a *sink*, and we say \mathcal{Q} has no sinks if E_{sinks}^0 is empty (or, equivalently, if the image of $s : E^1 \rightarrow E^0$ is dense in E^0). It is shown in [12, Proposition 3.15(1)] that $\ker \phi_{X(\mathcal{Q})} = C_0(E_{\text{sinks}}^0)$, and thus the left action $\phi_{X(\mathcal{Q})}$ is injective if and only if \mathcal{Q} has no sinks.

If $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ is a topological quiver, a *path* in \mathcal{Q} is a finite sequence of edges $\alpha := e_1 \dots e_n$ with $r(e_i) = s(e_{i+1})$ for all $1 \leq i \leq n-1$. We say that the path α has length $|\alpha| := n$, and we let E^n denote the set of paths in \mathcal{Q} of length n .

We may extend the maps r and s to $r^n : E^n \rightarrow E^0$ and $s^n : E^n \rightarrow E^0$ by setting $r^n(\alpha) := r(e_n)$ and $s^n(\alpha) := s(e_1)$. When no confusion arises, we shall omit the superscript and simply write r (respectively, s) for r^n (respectively, s^n).

We endow E^n with the topology it inherits as a subspace of the cartesian product $E^1 \times \dots \times E^1$. Since s is continuous and r is continuous and open, we see that s^n is continuous and r^n is continuous and open. We also define a Radon measure λ_v^n on E^n with support in $(r^n)^{-1}(v)$ inductively as follows: For $\xi : E^n \rightarrow \mathbb{C}$ we let

$$\begin{aligned} & \int_{(r^n)^{-1}(v)} \xi(e_1 \dots e_n) d\lambda_v^n(e_1 \dots e_n) \\ & := \int_{r^{-1}(s(e_2))} \dots \int_{r^{-1}(s(e_n))} \int_{r^{-1}(v)} \xi(e_1 \dots e_n) d\lambda_{s(e_2)}(e_1) \dots d\lambda_{s(e_n)}(e_{n-1}) d\lambda_v(e_n). \end{aligned}$$

Since the support of λ_v is $r^{-1}(v)$ for all $v \in E^0$, it follows that the support of λ_v^n is $(r^n)^{-1}(v)$.

If $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ is a topological quiver, then for any $n \in \mathbb{N}$ we have that $\mathcal{Q}^n := (E^0, E^n, r, s, \lambda^n)$ is a topological quiver as well. Furthermore, if $X(\mathcal{Q})$ is the quiver C^* -correspondence for \mathcal{Q} over $A := C_0(E^0)$, then $X(\mathcal{Q})^{\otimes n}$ is isomorphic to $X(\mathcal{Q}^n)$ (cf. [12, Section 6]). Moreover, we see that for $\xi, \eta \in X(\mathcal{Q})^{\otimes n}$ we have that $\langle \xi, \eta \rangle \in C_0(E^0)$ is the function with

$$\langle \xi, \eta \rangle(v) := \int_{(r^n)^{-1}(v)} \overline{\xi(\alpha)} \eta(\alpha) d\lambda_v^n(\alpha) \quad \text{for } v \in E^0.$$

A path $\alpha = e_1 \dots e_n$ is called a *cycle* if $s(e_1) = r(e_n)$, and we call the vertex $s(e_1) = r(e_n)$ the *base point* of the cycle. An *exit* for a cycle $\alpha = e_1 \dots e_n$ is an edge $f \in E^1$ with the property that $s(f) = s(e_i)$ and $f \neq e_i$ for some $1 \leq i \leq n$. (Note: In earlier literature, cycles we often called “loops”. In modern terminology, the typical convention is to use the term “loop” only for cycles of length one.) A cycle $\alpha = e_1 \dots e_n$ is called *simple* when $s(e_1) \neq r(e_i)$ for $1 \leq i \leq n-1$.

Condition (L): If \mathcal{Q} is a topological quiver, we say that \mathcal{Q} satisfies Condition (L) if the set of base points of cycles with no exits has empty interior in E^0 .

Note that a graph satisfies Condition (L) if and only if every cycle in the graph has an exit.

Definition 5.5. A path $\alpha := e_1 \dots e_n \in E^n$ is *returning* if $e_k = e_n$ for some $k \in \{1, 2, \dots, n-1\}$. Otherwise α is said to be *nonreturning*.

Definition 5.6. A nonempty subset $U \subseteq E^n$ is *nonreturning* if whenever $\alpha := e_1 \dots e_n \in U$ and $\beta := f_1 \dots f_n \in U$ we have that $\alpha_n \neq \beta_k$ for all $k \in \{1, \dots, n-1\}$.

Proposition 5.7. *Let $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ be a topological quiver with no sinks, and let $X(\mathcal{Q})$ be the associated quiver C^* -correspondence over $A := C_0(E^0)$. If \mathcal{Q} satisfies Condition (L), then $X(\mathcal{Q})$ satisfies Condition (S).*

Proof. Let $a \in A := C_0(E^0)$ with $a \geq 0$, let $n \in \mathbb{N}$, and let $\epsilon > 0$. Set

$$V := \{v \in E^0 : a(v) > \|a\| - \epsilon\} \subseteq E^0.$$

Since V is a nonempty open subset of E^0 , [12, Lemma 6.11] implies that there exists $m > n$ and a nonreturning path $\alpha \in (s^m)^{-1}(V)$. By [12, Lemma 6.12] there exists an open set $U \subseteq (s^m)^{-1}(V)$ such that $\alpha \in U$ and U is nonreturning. Using the

fact that U is an open subset of the locally compact space E^m , choose a continuous compactly supported function $\zeta : E^m \rightarrow [0, 1]$ with $\|\zeta\| = 1$ and with the support of ζ contained in U . It follows from [12, Lemma 6.11] that ζ is a nonreturning vector in $X(\mathcal{Q})^{\otimes m}$.

Since $\langle \zeta, \zeta \rangle$ is a positive continuous function in $C_0(E^0)$ with $\|\langle \zeta, \zeta \rangle\| = \|\zeta\|^2 = 1$, by the extreme value theorem there exists $v_0 \in E^0$ with $\langle \zeta, \zeta \rangle(v_0) = 1$. In addition, we have

$$\langle \zeta, a\zeta \rangle(v_0) = \int_{(r^m)^{-1}(v_0)} \overline{\zeta(\alpha)} a(s(\alpha)) \zeta(\alpha) d\lambda_{v_0}^n(\alpha).$$

Since ζ is supported in U , whenever $\zeta(\alpha) \neq 0$ we have $\alpha \in U$ and $s(\alpha) \in V$ so that $a(s(\alpha)) > \|a\| - \epsilon$. Thus

$$\begin{aligned} \langle \zeta, a\zeta \rangle(v_0) &= \int_{(r^m)^{-1}(v_0)} \overline{\zeta(\alpha)} a(s(\alpha)) \zeta(\alpha) d\lambda_{v_0}^n(\alpha) \\ &= \int_{(r^m)^{-1}(v_0)} a(s(\alpha)) |\zeta(\alpha)|^2 d\lambda_{v_0}^n(\alpha) \\ &\geq \int_{(r^m)^{-1}(v_0)} (\|a\| - \epsilon) |\zeta(\alpha)|^2 d\lambda_{v_0}^n(\alpha) \\ &= (\|a\| - \epsilon) \int_{(r^m)^{-1}(v_0)} |\zeta(\alpha)|^2 d\lambda_{v_0}^n(\alpha) \\ &= (\|a\| - \epsilon) \int_{(r^m)^{-1}(v_0)} \overline{\zeta(\alpha)} \zeta(\alpha) d\lambda_{v_0}^n(\alpha) \\ &= (\|a\| - \epsilon) \langle \zeta, \zeta \rangle(v_0) \\ &= (\|a\| - \epsilon) \cdot 1 \\ &= (\|a\| - \epsilon). \end{aligned}$$

Hence $\|\langle \zeta, a\zeta \rangle\| \geq \|a\| - \epsilon$, and $X(\mathcal{Q})$ satisfies Condition (S). \square

Proposition 5.8. *Let $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ be a topological quiver, and let $X(\mathcal{Q})$ be the associated quiver C^* -correspondence over $A := C_0(E^0)$. If $X(\mathcal{Q})$ satisfies Condition (S), then \mathcal{Q} satisfies Condition (L).*

Proof. We shall establish the contrapositive: If \mathcal{Q} does not satisfy Condition (L), then $X(\mathcal{Q})$ does not satisfy Condition (S).

Suppose that \mathcal{Q} does not satisfy Condition (L). Then there exists a nonempty open subset $U \subseteq E^0$ such that every vertex in U is the base point of a cycle with no exits. Since E^0 is locally compact, there exists a nonempty precompact subset $V \subseteq U$ with $\overline{V} \subseteq U$. Because every point in U is the base point of a cycle with no exits, we see that $\overline{V} \subseteq U \subseteq \bigcup_{n=1}^{\infty} r((s^n)^{-1}(U))$. In addition, since s^n is continuous and r is open, for each $n \in \mathbb{N}$ the set $r((s^n)^{-1}(U))$ is open. Therefore $\{r((s^n)^{-1}(U)) : n \in \mathbb{N}\}$ is an open cover of \overline{V} , and by the compactness of \overline{V} there exists $N \in \mathbb{N}$ such that $\overline{V} \subseteq \bigcup_{n=1}^N r((s^n)^{-1}(U))$. Thus $V \subseteq \bigcup_{n=1}^N r((s^n)^{-1}(U))$, and hence every vertex in V is the base point of a cycle with no exits having length at most N .

To prove that $X(\mathcal{Q})$ does not satisfy Condition (S), we shall show that there exists $a \in A := C_0(E^0)$ with $a \geq 0$ and there exists $\epsilon > 0$ such that for all $m > N$ when $\zeta \in X(\mathcal{Q})^{\otimes m}$ with $\|\zeta\| = 1$ and $\|\langle \zeta, a\zeta \rangle\| > \|a\| - \epsilon$, then ζ is not a nonreturning vector.

Using the fact that E^0 is locally compact, we choose a nonzero compactly supported $a \in C_0(E^0)$ with $a \geq 0$ and the support of a contained in V . Also let $\epsilon := \|a\|/2$. Suppose $m > N$ and $\zeta \in X(\mathcal{Q})^{\otimes m}$ with $\|\zeta\| = 1$ and $\|\langle \zeta, a\zeta \rangle\| > \|a\| - \epsilon = \|a\|/2$. We shall show that ζ is not a nonreturning vector.

As described in Remark 5.3, ζ may be viewed as a complex-valued function. Choose $v_0 \in E^0$ with $\langle \zeta, a\zeta \rangle(v_0) > \|a\|/2$. Since

$$(5.1) \quad \langle \zeta, a\zeta \rangle(v_0) = \int_{(r^m)^{-1}(v_0)} \overline{\zeta(\alpha)} a(s(\alpha)) \zeta(\alpha) d\lambda_{v_0}^m(\alpha),$$

and a has support contained in V , there exists a path $\alpha_0 \in E^m$ with $s(\alpha_0) \in V$ and $r(\alpha_0) = v_0$. Because every vertex in V is the base point of a cycle with no exits of length at most N , and since $N < m$, we may write

$$\alpha_0 = \underbrace{\beta\beta \dots \beta}_{k \text{ times}} \gamma$$

where β is the unique simple cycle based at $s(\alpha_0) \in W$ and γ is a subpath of β . However, we see that α_0 is then the unique path of length m with source in V and range equal to v_0 . Hence

$$\int_{(r^m)^{-1}(v_0)} \overline{\zeta(\alpha)} a(s(\alpha)) \zeta(\alpha) d\lambda_{v_0}^m(\alpha) = a(s(\alpha_0)) |\zeta(\alpha_0)|^2 \lambda_{v_0}^m(\{\alpha_0\})$$

and (5.1) implies

$$(5.2) \quad a(s(\alpha_0)) |\zeta(\alpha_0)|^2 \lambda_{v_0}^m(\{\alpha_0\}) = \langle \zeta, a\zeta \rangle(v_0) > \|a\|/2 > 0.$$

It follows that $\lambda_{v_0}^m(\{\alpha_0\}) > 0$. Let $n := |\beta|$, and note that $n \leq N < m$. Since $s(\beta) = s(\alpha_0) \in V$, we see that $\beta \in (s^n)^{-1}(V)$. Because E^n is locally compact, there exists a continuous compactly supported function $\xi_\beta : E^n \rightarrow [0, \infty)$ with $\xi_\beta(\beta) = a(s(\alpha_0))$ and the support of ξ_β contained in the open set $(s^n)^{-1}(V)$.

Let $(\psi_X, \pi_A) : (X(\mathcal{Q}), A) \rightarrow \mathcal{O}_{X(\mathcal{Q})} = C^*(\mathcal{Q})$ denote a universal covariant representation of ${}_A X(\mathcal{Q})_A$ into $C^*(\mathcal{Q})$. It follows from [12, Lemma 6.4] that

$$(5.3) \quad \psi_X^{\otimes m}(\zeta) * \psi_X^{\otimes n}(\xi_\beta) \psi_X^{\otimes m}(\zeta) = \psi_X^{\otimes n}(\eta),$$

where $\eta : E^n \rightarrow \mathbb{C}$ is defined by

$$\eta(\alpha) := \int_{(r^m)^{-1}(s(\alpha))} \overline{\zeta(e_1 \dots e_m)} \xi_\beta(e_1 \dots e_n) \zeta(e_{n+1} \dots e_m) d\lambda_{s(\alpha)}^m(e_1 \dots e_m).$$

Recalling that γ is a subpath of the cycle β , we may write $\beta = \gamma\mu$, where μ is a subpath of β with $s(\mu) = r(\gamma)$ and $r(\mu) = s(\beta) = s(\gamma)$. Note that $\mu\gamma$ is a cycle of length n , which is based at $s(\mu) = r(\gamma)$.

As noted before, $\alpha_0 = \beta\beta \dots \beta\gamma$ is the unique path of length m with source in V and range equal to v_0 . Hence if we take $\mu\gamma \in E^n$, then α_0 is also the unique path of length m with source in V and range equal to $r(\mu\gamma) = r(\gamma) = v_0$. Recalling that ξ_β has support contained in $(s^n)^{-1}(V)$ (and hence ξ_β vanishes on any path of length n with source outside V), we conclude

$$\begin{aligned} \eta(\mu\gamma) &= \overline{\zeta(\alpha_0)} \xi_\beta(\beta) \zeta(\underbrace{\beta\beta \dots \beta}_{k-1 \text{ times}} \gamma)(\mu\gamma) \lambda_{v_0}^m(\{\alpha_0\}) \\ &= \overline{\zeta(\alpha_0)} a(s(\alpha_0)) \zeta(\underbrace{\beta\beta \dots \beta}_{k-1 \text{ times}} (\gamma\mu)\gamma) \lambda_{v_0}^m(\{\alpha_0\}) \end{aligned}$$

$$\begin{aligned}
 &= \overline{\zeta(\alpha_0)} a(s(\alpha_0)) \zeta(\underbrace{(\beta\beta \dots \beta\gamma)}_{k \text{ times}}) \lambda_{v_0}^m(\{\alpha_0\}) \\
 &= \overline{\zeta(\alpha_0)} a(s(\alpha_0)) \zeta(\alpha_0) \lambda_{v_0}^m(\{\alpha_0\}) \\
 &= a(s(\alpha_0)) |\zeta(\alpha_0)|^2 \lambda_{v_0}^m(\{\alpha_0\}) \\
 &> 0
 \end{aligned}$$

where the last line follows from (5.2).

Thus $\eta \neq 0$. Since (ψ_X, π_A) is a universal covariant representation of $X(\mathcal{Q})$, both $\psi_X^{\otimes n}$ and π_A are injective, and hence $\psi_X^{\otimes n}(\eta) \neq 0$. It follows from (5.3) that $\psi_X^{\otimes m}(\zeta)^* \psi_X^{\otimes n}(\xi_\beta) \psi_X^{\otimes m}(\zeta) \neq 0$. Thus ζ is not a nonreturning vector. Hence $X(\mathcal{Q})$ does not satisfy Condition (S). \square

Theorem 5.9. *Let $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ be a topological quiver with no sinks, and let $X(\mathcal{Q})$ be the associated quiver C^* -correspondence over $A := C_0(E^0)$. Then \mathcal{Q} satisfies Condition (L) if and only if $X(\mathcal{Q})$ satisfies Condition (S).*

Proof. The result follows from Proposition 5.7 and Proposition 5.8. \square

6. AN EXAMINATION OF SCHWEIZER'S SIMPLICITY RESULT

Definition 6.1. Let X be a C^* -correspondence over the C^* -algebra A . We say X is *full* if

$$\overline{\text{span}}\{\langle x, y \rangle : x, y \in X\} = A.$$

Remark 6.2. Let A be a C^* -algebra. Then A is naturally a C^* -correspondence over itself with the bimodule structure given by multiplication and inner product is defined by $\langle a, b \rangle = a^*b$. This C^* -correspondence is called the *identity C^* -correspondence on A* , and it is used in the definition of Schweizer's "nonperiodic" condition.

Definition 6.3 (Nonperiodic Condition). Let X be a C^* -correspondence over the C^* -algebra A . We say X is *nonperiodic* if for all $n \in \mathbb{N} \cup \{0\}$, $X^{\otimes n} \cong A$ (as C^* -correspondences) implies $n = 0$. We say that a C^* -correspondence is *periodic* if it is not nonperiodic.

Theorem 6.4. (Schweizer's Simplicity Theorem [14, Theorem 3.9].) *Let X be a C^* -correspondence over a C^* -algebra A , and suppose that X is full, A is unital, and the left action $\phi_X : A \rightarrow \mathcal{L}(X)$ is injective. Then \mathcal{O}_X is simple if and only if X is nonperiodic and A has no hereditary ideals other than $\{0\}$ and A .*

Remark 6.5. Although the hypothesis of the injectivity of the left action is not mentioned in the statement of [14, Theorem 3.9], it is used to apply [14, Theorem 3.8] in its proof. Our statement above of Schweizer's theorem makes this hypothesis explicit.

Definition 6.6. If X is a C^* -correspondence over a C^* -algebra A , we say X is *nondegenerate* when $\overline{\text{span}}\{a \cdot x : a \in A \text{ and } x \in X\} = X$.

Definition 6.7. Let $E = (E^0, E^1, r, s)$ be a directed graph. The graph correspondence is the nondegenerate C^* -correspondence $X(E)$ over $C_0(E^0)$ defined by

$$X(E) = \{\xi : E^1 \rightarrow \mathbb{C} : \text{the map } v \mapsto \sum_{r(e)=v} |\xi(e)|^2 \text{ is in } C_0(E^0)\},$$

with module actions and $C_0(E^0)$ -valued inner product given by


$$(a \cdot \xi)(e) = a(s(e))\xi(e), \quad (\xi \cdot b)(e) = \xi(e)b(r(e)), \quad \text{and} \quad \langle \xi, \eta \rangle(v) = \sum_{r(e)=v} \overline{\xi(e)}\eta(e)$$

for $a, b \in C_0(E^0)$, $\xi, \eta \in X(E)$, $e \in E^1$, and $v \in E^0$.

Definition 6.8. If $E = (E^0, E^1, r, s)$ is a graph, then for any $n \in \mathbb{N}$ we let E^n denote the set of paths in E of length n . We also let $E^{\times n} := (E^0, E^n, r, s)$ denote the graph whose vertices are the vertices of E and whose edges are the paths of length n of E .

Remark 6.9. It follows from [1, Proposition 3.3] that if E is a graph, then $X(E)^{\otimes n} \cong X(E^{\times n})$ as C^* -correspondences for all $n \in \mathbb{N}$.

Example 6.10. Let E be the graph




consisting of a simple cycle with n vertices for some $n \in \mathbb{N}$. Note that $A := C(E^0) \cong \mathbb{C}^n$. Since each vertex of E is the base point of a cycle of length n , we see that $E^{\times n}$ is the graph




so that $X(E^{\times n}) \cong \mathbb{C}^n$ with the left and right multiplication occurring component-wise. Thus $X(E)^{\otimes n} \cong X(E^{\times n}) \cong \mathbb{C}^n \cong A$ as C^* -correspondences over A , and $X(E)$ is periodic.

Example 6.11. Let E be the graph



consisting of the disjoint union of three simple cycles of lengths 2, 3, and 4, respectively. If we take the least common multiple of these lengths, $\text{lcm}(2, 3, 4) = 12$, then we see that each vertex is the base point of a cycle of length 12, and hence $E^{\times 12}$ is the graph



In particular, since there are 9 vertices, we have $X(E)^{\otimes 12} \cong X(E^{\times 12}) \cong \mathbb{C}^9 \cong A$ as C^* -correspondences over A , and thus $X(E)$ is periodic.

Proposition 6.12. *Let E be a graph with a finite number of vertices, no sinks, and no sources. (Note that E is not required to have a finite number of edges.) Then the following are equivalent:*

- (1) $X(E)$ is nonperiodic.
- (2) E is not a disjoint union of simple cycles.

Proof. We shall establish the result by proving the contrapositives: $X(E)$ is periodic if and only if E is a disjoint union of cycles.

If E is disjoint union of cycles, then since E has a finite number of vertices there are a finite number of cycles in this disjoint union. Let n be the least common multiple of the lengths of these cycles. Then $E^{\times n}$ is a graph with a single cycle of length one at each vertex. Hence we have the C^* -correspondence isomorphisms $X(E)^{\otimes n} \cong X(E^{\times n}) \cong A$ (cf. Example 6.11). Thus $X(E)$ is periodic.

Conversely, suppose that $X(E)$ is periodic. Then there exists $n \in \mathbb{N}$ such that $X(E)^{\otimes n} \cong A$ as C^* -correspondences. For convenience of notation, label the vertices as $E^0 = \{1, \dots, k\}$. Then $C(E^0) \cong \mathbb{C}^k$, and we have the C^* -correspondence isomorphisms $X(E^{\times n}) \cong X(E)^{\otimes n} \cong \mathbb{C}^k \cong A$. Recalling that $X(E^{\times n}) := C(E^n)$, we conclude there exists a \mathbb{C}^k -linear isomorphism $h : C(E^n) \rightarrow C(E^0) \cong \mathbb{C}^k$. For each $\alpha \in E^n$, let $\delta_\alpha : E^n \rightarrow \mathbb{C}$ be the function with $\delta_\alpha(\alpha) = 1$ and $\delta_\alpha(\beta) = 0$ when $\beta \neq \alpha$. Also, for each $1 \leq i \leq k$, let e_i denote the standard basis element of \mathbb{C}^k having 1 in entry i and 0 in all other entries. Suppose $h(\delta_\alpha) = (z_1, \dots, z_k)$. Then

$$\begin{aligned} e_{r(\alpha)} &= \langle \delta_\alpha, \delta_\alpha \rangle = \langle h(\delta_\alpha), h(\delta_\alpha) \rangle = (z_1, \dots, z_k)^*(z_1, \dots, z_k) \\ &= (\overline{z_1}, \dots, \overline{z_k})(z_1, \dots, z_k) = (|z_1|^2, \dots, |z_k|^2). \end{aligned}$$

Consequently $z_i = 0$ for $i \neq r(\alpha)$, and $z_{r(\alpha)} \neq 0$. Thus $h(\delta_\alpha) = z_{r(\alpha)} e_{r(\alpha)}$. Hence

$$z_{r(\alpha)} e_{r(\alpha)} = h(\delta_\alpha) = h(e_{s(\alpha)} \cdot \delta_\alpha) = e_{s(\alpha)} h(\delta_\alpha) = e_{s(\alpha)} \cdot (z_1, \dots, z_k) = z_{s(\alpha)} e_{s(\alpha)}.$$

Since $z_{r(\alpha)} \neq 0$, the above equation implies $s(\alpha) = r(\alpha)$. Because α is an arbitrary element of E^n , we conclude every element of E^n is a cycle. Moreover, since $C(E^n) \cong C(E^0) \cong \mathbb{C}^k$, we deduce E^n has k elements. Thus $E^n = \{\alpha_1, \dots, \alpha_k\}$, where α_i is a cycle of length n for each $1 \leq i \leq k$. Therefore E is a disjoint union of simple cycles. \square

Definition 6.13. Let $E = (E^0, E^1, r, s)$ be a graph. We say E is *weakly connected* if for all $v, w \in E^0$ there exists a sequence of vertices $v_1, \dots, v_n \in E^0$ with $v_1 = v$, $v_n = w$, and for all $1 \leq i \leq n$ there is either an edge in E from v_i to v_{i+1} or there is an edge in E from v_{i+1} to v_i . We say E is *strongly connected* if for all $v, w \in E^0$ there is a path in E from v to w .

Corollary 6.14. *Let E be a weakly connected graph with a finite number of vertices, no sinks, and no sources. (Note that E is not required to have a finite number of edges.) Then the following are equivalent:*

- (1) $X(E)$ is nonperiodic.
- (2) E is not a simple cycle.

Corollary 6.15. *Let E be a strongly connected graph with a finite number of vertices. (Note that E is not required to have a finite number of edges.) Then the following are equivalent:*

- (1) $X(E)$ is nonperiodic.
- (2) E is not a simple cycle.


(3) E satisfies Condition (L).

Proof. The equivalence (1) \iff (2) follows from Corollary 6.14.

To see (2) \implies (3), suppose E is not a simple cycle. If α is any cycle in E , then since E is not a simple cycle, there exists a vertex $w \in E^0$ that is not on α . Since E is strongly connected, there exists a path from $s(\alpha)$ to w , and some edge of this path must be an exit for α . Hence E satisfies Condition (L), and (3) holds,


To see (3) \implies (2), simply note that if every cycle in E has an exit, then E cannot be a simple cycle. \square

Example 6.16. If E is the graph




then E is weakly connected and does not satisfy Condition (L). In addition, for any $n \in \mathbb{N}$, we see that there are paths of length n in E that are not cycles. Hence, the correspondence $X(E^{\times n})$ is not isomorphic to the correspondence $C_0(E^0)$, so that the correspondence $X(E)^{\otimes n}$ is not isomorphic to the identity correspondence on A . Consequently, $X(E)$ is nonperiodic.

Example 6.17. If E is the graph



then E is weakly connected and does not satisfy Condition (L). In addition, for any $n \in \mathbb{N}$, we see that there are paths of length n in E that are not cycles. Hence, the correspondence $X(E^{\times n})$ is not isomorphic to the correspondence $C_0(E^0)$, so that the correspondence $X(E)^{\otimes n}$ is not isomorphic to the identity correspondence on A . Consequently, $X(E)$ is nonperiodic. Furthermore, E has no nontrivial saturated hereditary subsets, and hence $A \cong \mathbb{C}^2$ has no invariant ideals other than $\{0\}$ and A . Since E does not satisfy Condition (L), we conclude $\mathcal{O}_{X(E)} \cong C^*(E)$ is not simple. (In fact, $C^*(E) \cong M_2(C(\mathbb{T}))$, so $C^*(E)$ is Morita equivalent to $C(\mathbb{T})$, and $C^*(E)$ has uncountably many ideals.)

Example 6.18. If E is the graph



then E is weakly connected, has no sinks and no sources, and does not satisfy Condition (L). In addition, for any $n \in \mathbb{N}$, we see that there are paths of length n in E that are not cycles. Hence, the correspondence $X(E^{\times n})$ is not isomorphic to the correspondence $C_0(E^0)$, so that the correspondence $X(E)^{\otimes n}$ is not isomorphic to the identity correspondence on A . Consequently, $X(E)$ is nonperiodic. Furthermore, E has no nontrivial saturated hereditary subsets, and hence $A = C_0(E^0)$ has no invariant ideals other than $\{0\}$ and A . Since E does not satisfy Condition (L), $\mathcal{O}_{X(E)} \cong C^*(E)$ is not simple. (In fact, $C^*(E) \cong C(\mathbb{T}) \otimes \mathcal{K}$, so $C^*(E)$ is Morita equivalent to $C(\mathbb{T})$, and $C^*(E)$ has uncountably many ideals.)

Example 6.19. For each $k \in \mathbb{N}$ let C_k be the graph that is a simple cycle of length k . If $E := C_1 \sqcup C_2 \sqcup \dots$, then for any $n \in \mathbb{N}$, we see that E contains a path of length

n that is not a cycle. Hence, the correspondence $X(E^{\times n})$ is not isomorphic to the correspondence $C_0(E^0)$, so that the correspondence $X(E)^{\otimes n}$ is not isomorphic to the identity correspondence on A . Consequently, $X(E)$ is nonperiodic. Thus $X(E)$ is nonperiodic even though E is a disjoint union of simple cycles.

Example 6.20. Let E be a strongly connected graph with an infinite number of vertices. If α is any cycle in E , then there exists a vertex $w \in E^0$ that is not on α . Since E is strongly connected, there exists a path from $s(\alpha)$ to w , and some edge of this path must be an exit for α . Hence E satisfies Condition (L).

In addition, since E is strongly connected, for any $n \in \mathbb{N}$ there exists a path of length n in E that is not a cycle. Hence, the correspondence $X(E^{\times n})$ is not isomorphic to the correspondence $C_0(E^0)$, so that the correspondence $X(E)^{\otimes n}$ is not isomorphic to the identity correspondence on A . Consequently, $X(E)$ is nonperiodic.

Therefore if E is any strongly connected graph with infinitely many vertices, then E satisfies Condition (L) and $X(E)$ is nonperiodic.

Remark 6.21. For a graph E , let $X(E)$ denote the graph C^* -correspondence over $A := C_0(E^0)$. Our results and examples from this section can be summarized by the following points:

- Corollary 6.15 states that if E is strongly connected with finitely many vertices, $X(E)$ being nonperiodic is equivalent to E satisfying Condition (L).
- Example 6.16 shows that there exists a weakly connected finite graph with no sinks and no sources such that $X(E)$ is nonperiodic, but E does not satisfy Condition (L). This shows that, even for finite graphs with no sinks and no sources, $X(E)$ being non-periodic does not imply that E satisfies Condition (L). Moreover, if E is the graph of Example 6.16, then $C^*(E)$ has proper non-zero ideals that do not contain any vertex projects. Thus, even in the special case of graph C^* -algebras (indeed, even when the graph is weakly connected, finite, has no sinks, and has no sources), the condition of nonperiodic is insufficient to obtain a Cuntz-Krieger Uniqueness Theorem.
- Example 6.17 shows that there exists a weakly connected finite graph E with no sinks (but having a source) for which $X(E)$ is nonperiodic and A has no invariant ideals other than $\{0\}$ and A , but E does not satisfy Condition (L). Since E does not satisfy Condition (L), the graph C^* -algebra $C^*(E)$ is not simple. This shows that the conditions “ X is nonperiodic and A has no hereditary ideals other than $\{0\}$ and A ” are insufficient to imply simplicity of \mathcal{O}_X when X is not full. Thus a characterization of simplicity for Cuntz-Pimsner algebras of C^* -correspondences that are not full (even under the hypotheses that A unital and the left action of A on X is injective) will require conditions other than “ X is nonperiodic and A has no hereditary ideals other than $\{0\}$ and A ”.
- Example 6.18 shows that there exists a weakly connected graph E with no sinks and no sources (but having a countably infinite number of vertices) for which $X(E)$ is nonperiodic and A has no invariant ideals other than $\{0\}$ and A , but E does not satisfy Condition (L). Since E does not satisfy Condition (L), the graph C^* -algebra $C^*(E)$ is not simple. This shows that the conditions “ X nonperiodic and A has no hereditary ideals other than $\{0\}$ and A ” are insufficient to imply simplicity of \mathcal{O}_X when A is not unital. Thus a characterization of simplicity for Cuntz-Pimsner algebras of

C^* -correspondences over nonunital C^* -algebras (even under the hypotheses that X is full and the left action of A on X is injective) will require conditions other than “ X nonperiodic and A has no hereditary ideals other than $\{0\}$ and A ”.

- Finally, Example 6.20 shows that for strongly connected graphs with an infinite number of vertices, nonperiodic and Condition (L) are both automatic. Moreover, in this situation the fact that the graph is strongly connected implies the vertex set has no hereditary subsets that are proper and nonempty, and hence the associated C^* -algebra is automatically simple.

These observations show that while the notion of nonperiodic is useful for characterizing simplicity under the hypotheses of Schweizer’s Simplicity Theorem, it is insufficient for characterizing simplicity more generally. Moreover, even when Schweizer’s hypotheses hold, the condition of nonperiodic is insufficient for obtaining a Cuntz-Krieger Uniqueness Theorem. Indeed, we see this even in the special case of graph C^* -algebras, where Condition (L) is a necessary and sufficient condition for a Cuntz-Krieger Uniqueness Theorem to hold.

This gives additional credibility for the usefulness of the notion of Condition (S) that we introduced in this paper. First of all, under the hypothesis that the left action is injective, Condition (S) coincides with Condition (L) for graphs, topological graphs, and topological quivers, suggesting it is a reasonable generalization of Condition (L) for general C^* -correspondences. Second, unlike the nonperiodic condition, we have been able to use Condition (S) to obtain a Cuntz-Krieger Uniqueness Theorem for Cuntz-Pimsner algebras of C^* -correspondence with injective left action. Third, our Cuntz-Krieger Uniqueness theorem allows us to obtain sufficient conditions for simplicity of Cuntz-Pimsner algebras in terms of Condition (S).

7. CONCLUDING REMARKS

We conclude with two questions that are important for extending the work in this paper.

Question 1: Let X be a C^* -correspondence over A with injective left action. If \mathcal{O}_X is simple, is it necessary that X satisfies Condition (S)?

When $X = X(\mathcal{Q})$ is the C^* -correspondence of a topological quiver \mathcal{Q} , we know that $\mathcal{O}_{X(E)} = C^*(\mathcal{Q})$ simple implies \mathcal{Q} satisfies Condition (L), which implies $X(\mathcal{Q})$ satisfies Condition (S). Thus we have an affirmative answer to Question (1) for C^* -correspondences of topological quivers, topological graphs, and graphs. If an affirmative answer to Question (1) is obtained in general, then Condition (S) is precisely the correct generalization of Condition (L) for C^* -correspondences with injective left action.

Question 2: How can Condition (S) be appropriately modified to obtain a Cuntz-Krieger Uniqueness Theorem for general C^* -correspondences whose left action is not necessarily injective?

As noted in Proposition 3.5, a C^* -correspondence satisfying Condition (S) necessarily has injective left action. It would be desirable to remove the hypothesis of injective left action and obtain an appropriate condition for general C^* -correspondences.

While there are techniques for extending results for C^* -correspondences with injective left actions to the general situation (e.g., via the method of “adding tails” described in [11]), translating Condition (S) to the non-injective setting has been complicated and elusive, and we have been unable to apply these techniques in any fruitful way. Currently, we are uncertain how to modify Condition (S) when the left action is not injective.

If one is able to find an appropriate formulation of Condition (S) for general C^* -correspondences and prove a Cuntz-Krieger Uniqueness Theorem under this hypothesis, then the result will also provide sufficient conditions for simplicity. In particular, once a general Cuntz-Krieger Uniqueness Theorem is obtained, the proof of Theorem 4.3 will go through verbatim and the modified Condition (S) together with the hypothesis that A has no invariant ideals other than $\{0\}$ and A will imply simplicity of the Cuntz-Pimsner algebra.

REFERENCES

- [1] T.M. Carlsen, A. Dor-On and S. Eilers, *Shift equivalences through the lens of Cuntz-Krieger algebras*, arXiv.2011.10329v5.
- [2] N. Fowler, P.S. Muhly, and I. Raeburn, *Representations of Cuntz-Pimsner algebras*, Indiana Univ. Math. J. **52** (2003), 569–605.
- [3] N. Fowler and I. Raeburn, *The Toeplitz algebra of a Hilbert bimodule*, Indiana Univ. Math. J. **48** (1999), 155–181.
- [4] T. Kajiwara, C. Pinzari, Y. Watatani, *Ideal structure and simplicity of the C^* -algebras generated by Hilbert bimodules*, J. Funct. Anal. **159** (1998), 295–322.
- [5] T. Katsura, *A class of C^* -algebras generalizing both graph algebras and homeomorphism C^* -algebras. I. Fundamental results*, Trans. Amer. Math. Soc. **356** (2004), 4287–4322.
- [6] T. Katsura, *On C^* -algebras associated with C^* -correspondences*, J. Funct. Anal. **217** (2004), 366–401.
- [7] T. Katsura, *Ideal structure of C^* -algebras associated with C^* -correspondences*, Pacific J. Math. **230** (2007), 107–145.
- [8] E. C. Lance, *Hilbert C^* -modules. A toolkit for operator algebraists*. London Mathematical Society Lecture Note Series, **210**. Cambridge University Press, Cambridge, 1995.
- [9] P. S. Muhly and B. Solel, *On the simplicity of some Cuntz-Pimsner algebras*, Math. Scand. **83** (1998), 53–73.
- [10] P. S. Muhly and B. Solel, *Corrigendum to: “On the simplicity of some Cuntz-Pimsner algebras”*, Math. Scand. **91** (2002), 244–246.
- [11] P. S. Muhly and M. Tomforde, *Adding tails to C^* -correspondences*, Documenta Math. **9** (2004), 79–106.
- [12] P.S. Muhly and M. Tomforde, *Topological quivers*, Internat. J. Math. **16** (2005), no. 7, 693–755.
- [13] M. V. Pimsner, *A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by \mathbb{Z}* , Fields Institute Comm. **12** (1997), 189–212.
- [14] J. Schweizer, *Dilations of C^* -correspondences and the simplicity of Cuntz-Pimsner algebras*, J. Funct. Anal. **180** (2001), 404–425.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO BOULDER, BOULDER, CO 80309-0395, USA

Email address: menevse.eryuzlu@colorado.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO, COLORADO SPRINGS, CO 80918-3733, USA

Email address: mtomford@uccs.edu