Generating functions in Riesz spaces

Youssef Azouzi*and Youssef Nasri[†]

Research Laboratory of Algebra, Topology, Arithmetic, and Order Department of Mathematics

COSAEF Faculty of Mathematical, Physical and Natural Sciences of Tunis Tunis-El Manar University, 2092-El Manar, Tunisia

December 2, 2022

Abstract

We introduce and study the concept of generating function for natural elements in a Dedekind complete Riesz space equipped with a conditional expectation operator. This allows to study discrete processes in free-measure setting. In particular we improve a result obtained by Kuo, Vardy and Watson concerning Poisson approximation.

Key words. Generating function, Dedekind complete Riesz space, Conditional expectation, order convergence, unbounded order convergence.

1 Introduction

We introduce in this paper the notion of generating function for special elements of a Riesz space called natural elements. Recently Ben Amor and Omrani published a paper in which they treated the notion of moment generating function in Riesz spaces; the main purpose of their paper is to get a generalization of Chernoff Inequality in the setting of free-measure theory

^{*}The authors are members of the GOSAEF research group

^{\dagger}This document is the results of the research project funded by the National Science Foundation

of Riesz spaces [6]. Our appraoch here is more general and leads to a more general Chernoff Inequality. Also we are able to give a generalization of a result obtained by Kuo, Vardy, and Watson [15]. In these two papers the authors studied only Bernoulli processes and sums of independent Bernoulli processes which give Binomial processes. The main idea in our work here is to develop a general approach to study discrete processes in Riesz spaces. For this reason Functional Calculus via Daniell integral as they are developed in [11] and [5] are more suitable than Functional Calculus as they are developed in [7]. This allows to work in the universal completion E^u of E, and even in its sup-completion E^s . When dealing with natural elements we can use their distributions or their generating functions to define convergence in distribution. This notion can be defined via order and unbounded order convergence and it will be useful to note that these modes of convergence agree for sequences if the space is universally complete; this result is due to Grobler and can be found in [10]. The latter paper contains several important results about unbounded order convergence.

The paper is organized as follows. Section 2 contains some preliminaries needed in the following sections. We prove some results concerning band projections and functional calculus developed using Daniell integral on Riesz spaces following Grobler [11]. In Section 3 we develop some results on series in Riesz spaces. We state and prove versions of Monotone Convergence Theorem, Dominated Convergence Theorem and others and we will make a frequent use of Fubini Theorem.

In section 4 we introduce the notion of generating function in Riesz spaces equipped with a conditional expectation operator and study its basic properties. We prove, in particular, that a natural element x belongs to $L^{p}(T)$ if and only if its generating function g_{x} is an order deravative of rank k at 1.

The last section contains some applications. In particular we give a new short proof of a result established by Kuo, Vardy and Watson in [15] concernin Poisson approximation and we state and prove a Riesz space version of Chernoff Inequality, generalizing a result obtained recently by Ben Amor and Omrani in [6].

For more information about generating function classical probability theory we refer the reader to Feller [9], [17]. All unexplained terminology and notation concerning Riesz spaces an be found in standard references [1] and [19].

2 Preliminaries

Definition 1 We say that (E, e, T) is a conditional Riesz triple if E is a Dedekind complete Riesz space, e is a weak order unit and T is a conditional expectation operator, that is, a positive order continuous projection on E satisfying Te = e, with range, R(T), a Dedekind complete Riesz subspace. We will require in this paper that T is strictly positive, which means that if f > 0 then Tf > 0 (see [5]).

Independence. The concept of independence with respect to a conditional expectation has been introduced in [14, Definition 4.1]. See also [12, Definition 4.1]. We recall here some facts.

A family $(P_{\alpha})_{\alpha \in A}$ of band projections is said to be *T*-independent if for each finite subset $\{\alpha_1, ..., \alpha_n\}$ of *A* we have $TP_{\alpha_1}P_{\alpha_2}...P_{\alpha_n}T = TP_{\alpha_1}TP_{\alpha_2}...TP_{\alpha_2}T$, or equivalently,

$$TP_{\alpha_1}P_{\alpha_2}....P_{\alpha_n}e = TP_{\alpha_1}TP_{\alpha_2}...TP_{\alpha_2}e.$$

A family of closed Riesz subspaces (E_{α}) is said to be *T*-independent if every family of band projections such that $P_{\alpha}e \in E_{\alpha}$ for every α is *T*independent.

A family of elements (f_i) in E is independent with respect to T and e if the family $\{\langle f_i, R(T) \rangle : i \in I\}$ of Dedekind complete Riesz subspaces is independent, where $\langle f, R(T) \rangle$ denotes the closed Riesz subspace generated by f and R(T). It is worth noting that if $x, y \in L^1(T)$ are independent then $xy \in L^1(T)$ and

$$T(xy) = TxTy. \tag{1}$$

Observe that the second member exists as R(T) is an *f*-subalgebra of E^u (see [5, Theorem 3.1]). The formula is true by definition if x and y are components, and hence it extends by linearity for step elements, so for positive elements by Freudenthal Theorem and order continuity of T. Finally we use the linearity to extend it for all elements in $L^1(T)$. Using the fact that Tcan be extended to the whole space of E^s (see [4, Corollary 6]) it easy to see that the formula (1) holds for all elements in E^u_+ .. Here we use the fact that if x and y are T-independent then $x \wedge ne$ and $y \wedge ne$ are T-independent as well. For more information about T-independence the reader is referred to [12].

Bands and projections. As E is Dedekind complete with weak order unit, every band in E is a principal projection band. If P is the band projection on the band B then $P^d = I - P$ is the band projection on B^d . The projection band generated by x is denoted by B_x and the corresponding band projection is denoted by P_x . We recall that for $x, y \in E_+$ we have $B_{x \vee y} = B_{x+y}$. It is easily seen that $P_x^d P_y^d = P_{x \vee y}^d$. If $x, y \in E$ we put $P_{x=y} = P_{(x-y)^+}^d P_{(y-x)^+}^d = P_{|x-y|}^d$. We shall adopt some notations used by Grobler (see for example [11]):

$$P_{x \ge y} = P_{(y-x)^+}^d, \ P_{x > y} = P_{(x-y)^+}, P_{x \ge y, a \ge b} = P_{(y-x)^+}^d P_{(b-a)^+}^d \text{ etc.}$$

Daniell integral. For an element $x \in E$ we denote by I_x the Daniell integral with respect to the right continuous spectral system $(e_t^x)_{t \in \mathbb{R}} := (P_{x \leq te} e)_{t \in \mathbb{R}}$ of x, initially defined on the Riesz space of $\mathcal{F}(\mathbb{R})$ -step functions on \mathbb{R} denoted by $L = L(\mathcal{F}(\mathbb{R}))$ and extended to $L_2(I_x)$, where $\mathcal{F}(\mathbb{R})$ is the algebra consisting of all finite unions of disjoint left open right closed intervals $(a, b], (a, \infty), (-\infty, b]$ with $a, b \in \mathbb{R}$. We just recall that $I_x(f)$ is denoted by f(x) and that $f(x) = e_b^x - e_a^x$ for $f = \mathbf{1}_{(a,b]}$. We refer the reader to [11] and [5] for more details of the subject. We consider here the following spaces

$$L^{\uparrow} = L^{\uparrow}(I_x) := \left\{ f \in \overline{\mathbb{R}}^{\mathbb{R}} : \exists (f_n) \subset L, \ f_n \uparrow f \right\}, \\ L_1^{\uparrow} = L_1^{\uparrow}(I_x) := \left\{ f \in \mathbb{R}^{\mathbb{R}}_+ : f \in L^{\uparrow}(I_x) \text{ and } I_x(f) \in E \right\}, \\ L_1 = L_1(I_x) := L_1^{\uparrow}(I_x) - L_1^{\uparrow}(I_x), \\ L_2 = L_2(I_x) := L^{\uparrow}(I_x) - L_1(I_x) = L^{\uparrow}(I_x)_+ - L_1(I_x).$$

Lemma 2 Let (E, e, T) be a conditional Riesz triple and $x, y, w \in E$. Then the following hold.

- (i) $P_{|x-y|\vee(x-w)^+} = P_{|x-y|\vee(y-w)^+};$
- (*ii*) $P_{x=y,x=w} = P_{x=y,y=w}$;

(iii)
$$P_{x \ge y}P_{x=w} = P_{w \ge y}P_{x=w}$$
 and $P_{y \ge x}P_{x=w} = P_{y \ge w}P_{x=w}$

Proof. (i) It follows from the inequality

$$(x-w)^+ \le (x-y)^+ + (y-w)^+ \le |x-y| + (y-w)^+.$$

that $B_{|x-y|\vee(x-w)^+} \subset B_{|x-y|+(y-w)^+} = B_{|x-y|\vee(y-w)^+}$. We interchange x and y, obtaining the converse inclusion and so the desired equality follows.

(ii) As $P_{x=y,x=w} = P^d_{|x-w| \vee |x-y|}$ it is sufficient to show the inequality

$$P_{|x-w|\vee|x-y|} \le P_{|x-y|\vee|y-w|},$$

as the converse inequality is similar. But this follows easily from the inequality $|x - w| \le |x - y| + |y - w|$. The proof of (iii) is similar.

We need in our work some results on calculus functional from the paper of Grobler [11] and Azouzi and Trabelsi [5]. The next result will be useful and can not be found in those papers and is more general than [5, Lemma 2.7] and [11, Theorem 3.9].

Theorem 3 Let E be a Dedekind complete Riesz space with weak order unit e.

(i) If
$$f, g \in L_2(I_x)^+$$
 then $(fg)(x) = f(x)g(x)$.

(ii) Assume $g \in L_1(I_x)$, $f \in L_2(I_x)$, $g(t) \in dom(f)$ for every t. Then $(f \circ g)(x) = f(g(x)) \in E^s$.

Proof. (i) Assume first that $f = \mathbf{1}_{(-\infty,a)}$ and $g = \mathbf{1}_{(-\infty,b)}$. Then $fg = \mathbf{1}_{(-\infty,a\wedge b)}$. Moreover $f(x) = p_a$, $g(x) = p_b$ and $(fg)(x) = p_{a\wedge b} = p_a p_b$. The formula is then true for all positive linear combination of indicator functions.

If $f \in L^{\uparrow}$ there exists a sequence of step functions (f_n) such that $f_n \uparrow f$ and $g_n \uparrow g$. Now using the previous step we deduce that

$$(fg)(x) = f(x)g(x).$$

This can be extended for every $f, g \in L_2(I_x)^+$.

(ii) This is true if f is continuous by [11, Theorem 3.9]. To see that this is also true for every $f \in L^{\uparrow}$ it is only sufficient to observe that for every $f \in L^{\uparrow}$ there is a sequence of positive continuous functions f_n such that $f_n \uparrow f$. Now by taking the supremum over n in the equality $(f_n \circ g)(x) = f_n(g(x))$ we obtain $(f \circ g)(x) = f(g(x))$. The result for general $f \in L_2(I_x)$ follows easily.

Lemma 4 Let E be a Dedekind complete Riesz space with weak order unit e and $x, y \in E$. Then the following statements hold.

- (i) $P_{x=y}f(x) = P_{x=y}f(y)$, for all $f \in L_2(I_x)$. In particular $P_{x=y}x = P_{x=y}y$.
- (*ii*) $P_{f(x)^+}g(x) = (g\mathbf{1}_{(0,\infty)} \circ f)(x)$ and $P_{f(x)^-}g(x) = (g\mathbf{1}_{(-\infty,0)} \circ f)(x)$, for all $f, g \in L_1(I_x)$ for all $f, g \in L_1(I_x)$.

Proof. (i) We prove first the equality for $f = \mathbf{1}_{(-\infty,a]}, a \in \mathbb{R}$. But in this case $f(x) = P^d_{(x-ae)^+}$ and the result follows from Lemma 2(i). So it can be extended by linearity and then using the continuity of order projection and monotone convergence of Daniell integral to every function in $L_2(I_x)$.

(ii) We have

$$P_{f(x)^{+}}g(x) = g(x)P_{f(x)^{+}}e = g(x).\mathbf{1}_{(0,\infty)}(f(x)) = \left(g.\mathbf{1}_{(0,\infty)} \circ f\right)(x),$$

The second part is similar. \blacksquare

Remark 5 It should be noted that (ii) is not true for f in $L_2(I_x)$. Take for example, $x = e, g = 1, f = \infty$. Then $f \in L_2(I_x)$ and $e = P_{f(x)+}g(x) \neq (g\mathbf{1}_{(0,\infty)} \circ f)(x) = 0$.

Definition 6 Let (E, e, T) be a conditional Riesz triple and $x \in E$.

1. The function

$$\pi_x: t \in \mathbb{R} \longmapsto TP_{x=te}e = F_x(t) - F_x(t^-).$$

is called the (T, e)-mass function of x, where $F_x : t \mapsto TP_{x \leq te}e$ is the T-distribution function of x and $F_x(t^-) = \lim_{s \uparrow t} F_x(s) = TP_{(x-te)^-}e$.

- 2. An element x of E is said to be natural if $\sum_{n=0}^{\infty} \pi_x(n) = e$.
- 3. Note that if x is a natural element in E, then its T-distribution function F_x is piecewise-constant. Bands $\{B_{x=ne} : n \ge 0\}$ are disjoint and it follows from the equality $e = \sum_{n=1}^{\infty} \pi_x(n)$ that $E = [\{B_{x=ne} : n \ge 0\}]$. We derive that for every $a \in (n, n+1)$, $B_{ne < x \le ae} = 0$, which means that $F_x(a) = F_x(n)$ for every $a \in (n, n+1)$.
- 4. In the definition above an element x is said to be natural if $\sum_{n \in \mathbb{N}} \pi_x(n) = e$. As T is supposed to be strictly positive this condition is equivalent to the fact that $\sum_{n \in \mathbb{N}} P_{x=ne}e = e$. But as it is not common to add this assumption we will keep the first definition.

Let (E, e, T) be a conditional Riesz triple. We say that two natural vectors x and y in E have the same conditional distribution with respect to T and we write $x \stackrel{d}{=}_T y$ if they have the same distribution function. Bernoulli processes and Poisson processes have been introduced in [15]. Each one of them is defined as a sequence of elements in E having the same distribution despite the notion of distribution was not explicitly used. Next we define in a more general manner these notions and then explain the link between our definitions and those introduced in [15].

Definition 7 We say that an element $x \in E$ has Bernoulli distribution with parameter 0 with respect to <math>T, and write $x \sim \mathcal{B}(p)$. whenever $\pi_x(1) = p$ and $\pi_x(0) = e - p$. Binomial distribution with parameters $n \in \mathbb{N}$ and $p \in (0, e)$ and Poisson distribution with parameter $g \in E$, g > 0 are defined respectively by their mass function

$$\pi_x (k) = \binom{n}{k} p^k (e-p)^{n-k}, \qquad k = 0, 1, 2, \dots n,$$

$$\pi_x (k) = \frac{g^k}{k!} e^{-g}, \qquad k = 0, 1, 2, \dots$$

We write $x \sim \mathcal{B}(n, p)$ and $x \sim \mathcal{P}(p)$.

Next we show that our definition of Bernoulli processes in vector lattices coincides with that considered in [15].

Proposition 8 Let (E, e, T) be a conditional Riesz triple and $x \in E$. Then $x \sim_T B(p)$ if and only if x = Qe for some band projection Q with TQe = p.

Proof. If x = Qe, then $\pi_x(1) = TP_{x=e}e = TP_{Qe=e}e = TP_{e-Qe}^d e = TQe$, and $\pi_x(0) = TP_{x=0}e = TP_{Qe=0}e = TP_{Qe}^d e = TQ^d e = e - TQe$, Hence $x \sim_T B(p)$ with p = TQe. Conversely assume that $x \sim_T B(p)$. Then $e = TP_{x=e}e + TP_{x=0}e = T(P_{x=e}e + P_{x=0})e$. We will show now that $P_{x=e}e \wedge P_{x=0}e = 0$. But this follows from Lemma 2.(ii) as $P_{x=0}P_{x=e} = P_{0=e}P_{x=e} = 0$. Hence $e \geq P_{x=e}e + P_{x=0}e$; which gives

$$0 = Te - TP_{x=e}e - TP_{x=0}e = T(e - P_{x=e}e - P_{x=0}e).$$

As T is strictly positive and $e - P_{x=e}e - P_{x=0}e \ge 0$ we get $e = P_{x=e}e + P_{x=0}e$. Thus

$$x = P_{x=e}x + P_{x=0}x = P_{x=e}e,$$

and, as required, the band projection $Q = P_{x=e}$ satisfies $TQe = TP_{x=e}e = p$.

We end this preliminary section with a brief discussion concerning order differentiability in Riesz spaces. We need this essentially to get in terms of differentiability of the generating function in 1 of the existence of a moments of a given order for a natural element in E.

Definition 9 Let J be an interval of \mathbb{R} not reduced to a point, s_0 a fixed point in J and E a Riesz space. A function $f: J \longrightarrow E$ is said to be (order) differentiable at s_0 if the order limit $\lim_{s \longrightarrow s_0} \frac{f(s) - f(s_0)}{s - s_0}$ exists in E. This limit is called the (order) derivative of f at s_0 and is denoted by $f'(s_0)$.

Notions of right and left differentiability can be defined in an obvious way. We can define by induction the nth derivative of f by putting $f^{(n+1)} := (f^{(n)})'$ for n = 0, 1, 2, ...

For general information about Riesz spaces we refer the reader to [19] and [1]. We will make use of the notion of sup-completion, a comprehensive presentation of the subject can be found in [2] and [4].

3 Series in Riesz spaces

We will prove in this section some results concerning series in Riesz spaces. We consider only convergence in order, so $\sum_{n=1}^{\infty} x_n$ means the order limit of the sequence (S_n) , where $S_n = \sum_{k=1}^n x_k$. We say that the series $\sum x_n$ is absolutely convergent if the series $\sum |x_n|$ is order convergent. In σ -order complete Riesz spaces every absolutely convergent series is order convergent. In particular if $|x_n| \leq y_n$ for each n and if the series $\sum y_n$ is order convergent, then $\sum x_n$ is order convergent as well. When (f_n) is a sequence of functions from an interval I to a Riesz space E, we say that $\sum f_n$ is order convergent uniformly on t if the series $\sum f_n(t)$ converges in order for every $t \in I$ and there is a net (y_α) such that $y_\alpha \downarrow 0$ and for every α there exists n_0 such that $\left|\sum_{k=n}^{\infty} f_n(t)\right| \leq y_\alpha$ for all $n \geq n_0$. We are interested in this work only in Dedekind complete Riesz spaces; in such spaces the former property is equivalent to the fact that $\sup_{n \ge k} \sup_{t \in I} \left| \sum_{j=n}^{\infty} f_j(t) \right| \downarrow 0$ as $k \longrightarrow \infty$. The following is a version of Monotone Convergence Theorem for series in Riesz spaces.

Theorem 10 Let E be a Dedekind complete Riesz space and $(x_{\alpha}(n))_{\alpha \in \Gamma}$ be a sequence of increasing net in E^s_+ . Then

$$\lim_{\alpha} \sum_{n=1}^{\infty} x_{\alpha}(n) = \sum_{n=1}^{\infty} \lim_{\alpha} x_{\alpha}(n) = \sum_{n=1}^{\infty} x(n).$$

Proof. Recall that suprema commute in E_{+}^{s} and that the limit of an increasing net is its supremum. As $x_{\alpha}(n)$ is positive for all n, α we have

$$\sum_{n=1}^{\infty} x(n) = \sup_{N} \sum_{n=1}^{N} x(n) = \sup_{N} \sup_{\alpha} \sum_{n=1}^{N} x_{\alpha}(n)$$
$$= \sup_{\alpha} \sup_{N} \sum_{n=1}^{N} x_{\alpha}(n) = \sup_{\alpha} \sum_{n=1}^{\infty} x_{\alpha}(n)$$
$$= \lim_{\alpha} \sum_{n=1}^{\infty} x_{\alpha}(n),$$

and we are done. \blacksquare

Next we state a Dominated Convergence Theorem for series in Riesz spaces.

Theorem 11 Let $(x_{\alpha}(n))_{\alpha\in\Gamma}$ be a sequence of nets in E, where E is a Dedekind complete Riesz space. We assume that the following conditions are satisfied.

(i)
$$x_{\alpha}(n) \xrightarrow{o} x(n)$$
 for each n .
(ii) $|x_{\alpha}(n)| \leq y(n)$ for every α , with $\sum_{n=1}^{\infty} y(n) \in E^{u}$.
Then $\lim_{\alpha} \sum_{n=1}^{\infty} x_{\alpha}(n) = \sum_{n=1}^{\infty} \lim_{\alpha} x_{\alpha}(n)$.

Proof. It is sufficient to show that

$$\limsup_{\alpha} \left| \sum_{n=1}^{\infty} x_{\alpha}(n) - \sum_{n=1}^{\infty} x(n) \right| = 0.$$

Observe first that $|x(n)| \leq y(n)$, so the series $\sum x(n)$ is order convergent. Fix an integer N. Then it is clear that

$$\lim_{\alpha} \sum_{n=1}^{N} |x_{\alpha}(n) - x(n)| = 0.$$

It is enough to prove that

$$A := \limsup_{\alpha} \left| \sum_{n=1}^{\infty} x_{\alpha}(n) - \sum_{n=1}^{\infty} x(n) \right| = 0.$$

For this end observe that

$$A \leq \limsup_{\alpha} \sum_{n=1}^{\infty} |x_{\alpha}(n) - x(n)|$$

$$\leq \limsup_{\alpha} \sum_{n=N+1}^{\infty} |x_{\alpha}(n) - x(n)| + \limsup_{\alpha} \sum_{n=1}^{N} |x_{\alpha}(n) - x(n)|$$

$$\leq 2 \sum_{n=N+1}^{\infty} 2 |y(n)|.$$

As this happens for every N, we get A = 0 as required.

We end this section by a version of Fubini Theorem in Riesz spaces. For a family $(x_i)_{i \in J}$ of positive elements in E we write:

$$\sum_{i \in I} x_i = \sup_{F \text{finite } \subset I} \sum_{k \in F} x_k \in E^s.$$

The following properties are very similar to the real case and their proofs will be omitted.

Theorem 12 Let E be a Dedekind complete Riesz space and $(x_i)_{i \in I}$ a family in E^s_+ . Then the following hold.

1. If
$$I = \mathbb{N}$$
 then $\sum_{i \in I} x_i = \lim_{n \to \infty} \sum_{k=1}^n x_k$

2. If $(J_{\alpha})_{\alpha}$ is a partition of I then $\sum_{i \in I} x_i = \sum_{\alpha} \sum_{i \in J_{\alpha}} x_i$. In particular we have

$$\sum_{(i,j)\in\mathbb{N}^2} x_{i,j} = \sum_{i\in\mathbb{N}} \left(\sum_{j\in\mathbb{N}} x_{i,j} \right) = \sum_{j\in\mathbb{N}} \sum_{i\in\mathbb{N}} x_{i,j}.$$
 (2)

3. In the last statement if the family is in E then if $\sum_{(i,j)\in\mathbb{N}^2} |x_{i,j}| \in E^u$ then the formula (2) holds.

4 Generating function

We assume throughout that (E, e, T) is a conditional Riesz triple.

Definition 13 Let x be a T-natural vector in E.

(i) The T-generating function of x, is the function $g_x : [0, \infty) \longrightarrow E^s$ defined by

$$g_x(s) = \sum_{n \ge 0} s^n T P_{x=ne} e = \sum_{n \ge 0} s^n \pi_x(n).$$

(ii) The generalized T-generating function of x is the function $\tilde{g}_x : E_+ \longrightarrow E^s$ defined by

$$\widetilde{g}_x(u) = \sum_{n \ge 0} TP_{x=ne} u^n = \sum_{n \ge 0} \pi_x(n) u^n.$$

Example 14 We refer to Example 7 for the notations below.

- (i) If $x \sim B(p)$ then $g_x(s) = sp + e p$.
- (ii) If $x \sim B(n, p)$ then $g_x(s) = (sp + e p)^n$.
- (iii) If $x \sim \mathcal{P}(p)$ then $g_x(s) = \exp((s-1)g)$.

The following result collects basic properties of generating functions.

Proposition 15 Let (E, e, T) be a conditional Riesz triple and let x be a natural element in E. Then the following hold.

(i)
$$g_x(0) = \pi_x(0), \ g_x(1) = e \text{ and } g_x(s) = \tilde{g}_x(se);$$

- (ii) $0 \le g_x(s) \le e$, $0 \le s \le 1$. and $0 \le \tilde{g}_x(u) \le e$, $0 \le u \le e$.
- (iii) $g_x(s) = Ts^x$, for all $s \ge 0$.
- (iv) g_x and \tilde{g}_x are non-decreasing. Moreover, if x > 0 then g_x and \tilde{g}_x are strictly increasing.
- (v) g_x and \tilde{g}_x are convex. Moreover, if $TP_{x \leq e}e < e$ then g_x and \tilde{g}_x are strictly convex.

Proof. (i), (ii) and (iv) are obvious.

(iii) As $s^{ne} = s^n e$, it follows from Lemma 4 that $P_{x=ne}s^n e = P_{x=ne}s^x$. Hence

$$g_x(s) = \sum_{n=1}^{\infty} TP_{x=ne} s^n e = \sum_{n=1}^{\infty} TP_{x=ne} s^x$$
$$= T\left[\left(\sum_{n=1}^{\infty} P_{x=ne}\right) s^x\right] = Ts^x.$$

(v) This follows from the fact that the map $s \mapsto s^n$ is convex for $n \ge 0$ and strictly convex for $n \ge 2$. Observe that the condition $TP_{x \le e}e < e$ means that $\pi_x(n) > 0$ for some $n \ge 2$.

Chernoff Inequality is stated in general in terms of exponential generating functions (see for instance [16]). We state a version of Chernoff Inequality in Riesz spaces.

Theorem 16 (Chernoff Bounds) Let (E, e, T) be a Riesz conditional triple, x a natural element in E and $u \in R(T)$. Then the following hold.

$$TP_{x \ge u} e \le s^u g_x(s), \quad \forall s > 1,$$

$$TP_{x \le u} e \le s^{-u} g_x(s), \quad \forall 0 < s < 1.$$

Proof. Let s > 1. According to Lemma 2.(iii) we have

$$P_{x \ge u} P_{x=ne} s^u = P_{ne \ge u} P_{x=ne} s^u e \le P_{ne \ge u} P_{x=ne} s^n e.$$

It follows that

$$P_{x \ge u} s^u e = \sum_{n=1}^{\infty} P_{x \ge u} P_{x=ne} s^u e \le \sum_{n=0}^{\infty} P_{ne \ge u} P_{x=ne} s^n e$$
$$= P_{ne \ge u} \sum_{n=0}^{\infty} P_{x=ne} s^n e = P_{ne \ge u} x^s.$$

Applying T to both sides, we get the first inequality. The second one can be proved in a similar way. \blacksquare

Corollary 17 Let x be a natural vector in E and $\alpha \in \mathbb{R}_+$. Then the following hold.

$$TP_{x \ge \alpha e} e \le \frac{g_x(s)}{s^{\alpha}}, \quad \forall s > 1$$
$$TP_{x \le \alpha e} e \le \frac{g_x(s)}{s^{\alpha}}, \quad \forall 0 < s < 1.$$

We will show now that the main result in [6] can be deduced from the previous corollary.

Theorem 18 [6, Theorem 5.3]Let (E, e, T) be Riesz conditional triple. Let (P_j) be a Benoulli process with $TP_j = f$ for all $j \in \mathbb{N}$ for some $f \in E$. Let $S_n = \sum_{k=1}^n P_k$. Then

$$TP_{(S_n-te)^+}e \le \left(\frac{n \|f\|_u \exp 1}{t}\right)^t \exp\left(-nf\right).$$

holds for all $t > n \|f\|_e$, where

$$\|f\|_e := \inf \left\{ \beta \in \mathbb{R} : |f| \le \beta e \right\}.$$

It should be noted that in [6] the authors assume that e is a strong unit for E. This is the case if we work in the ideal generated by e. **Proof.** As $S_n \sim_T B(n, f)$ we have

$$g_{S_n}(s) = (sf + e - f)^n = (e + (s - 1)f)^n.$$

In the sequel we will use the inequality $\log (e + x) \leq x$ for all $x \in E_+$ and the formula (see [11, Theorem 3.9])

$$\exp\left[n\log x\right] = x^n$$

true for any $x \ge \alpha e$, where $\alpha \in (0, \infty)$. Note also that e is a multiplicative unit in the *f*-algebra E_e . The real $s = \frac{t}{n \|f\|_e}$ satisfies s > 1 and we have then

$$g_{S_n}(s) = (e + (s - 1) f)^n = \exp[n \log(e + (s - 1) f)]$$

$$\leq \exp[n (s - 1) f] = \exp[nsf] \exp[-nf]$$

$$\leq \exp(ns ||f||_e e) \exp(-nf) = \exp(t) \exp(-nf)$$

Now using Corollary 17 to derive that

$$TP_{(S_n-te)^+}e \le TP_{S_n \ge te}e \le \frac{g_{S_n}(s)}{s^t} \le \frac{\exp t. \exp(-nf)}{s^t}$$
$$= \exp t. \left(\frac{n \|f\|_e}{t}\right)^t \exp(-nf),$$

which completes the proof. \blacksquare

Lemma 19 Let (E, e, T) be a conditional Riesz triple and x a natural element in E. Then the following hold.

(i) g_x is continuous on [0, 1] and infinitely differentiable on [0, 1). Moreover we have the formula

$$g_x^{(k)}(s) = \sum_{n=k}^{\infty} n(n-1)...(n-k+1)s^{n-k}\pi_x(n)$$
(3)

for every integer $k \ge 0$ and for all $s \in [0, 1)$.

(ii) For each integer k, we have

$$g_x^{(k)}(s) = T \left[x(x-e)...(x-(k-1)e)s^{x-ke} \right]$$

In particular, $g_x^{(k)}(0) = k!\pi_x(k).$

It follows then from the above lemma that

$$g_x(s) = \sum_{n=0}^{\infty} s^n \frac{g_x^{(k)}(0)}{k!}, \qquad s \in [0,1).$$

Proof. (i) As $|s^k \pi_x(k)| \leq \pi_x(k)$ and $\sum \pi_x(k)$ order converges, it follows from Theorem 11 that

$$\lim_{s \to s_0} \sum_{n=0}^{\infty} s^n \pi_x(k) = \sum_{n=0}^{\infty} \lim_{s \to s_0} s^n \pi_x(k) = \sum_{n=0}^{\infty} s_0^n \pi_x(k) ,$$

which proves that g_x is continuous on [0, 1]. Let us now fix a real $s_0 \in [0, 1)$ and a compact neighborhood K of s_0 in [0, 1). We have for $k \in \mathbb{N}$ and $s \in K$,

$$\left|\frac{s^{k}\pi_{x}(k) - s_{0}^{k}\pi_{x}(k)}{s - s_{0}}\right| \le kM^{k-1}\pi_{x}(k) \le kM^{k-1}e,$$

where $M = \sup_{t \in K} |t| < 1$. Using Theorem 11 again we derive that g_x is differentiable on [0, 1) and that formula 3 holds for k = 1. We can then prove by induction on k that g_x is k-differentiable and the formula 3 is valid.

(ii) Let $0 \le s < 1$. We have

$$g_x^{(k)}(s) = \sum_{n=k}^{\infty} n(n-1)...(n-k+1)s^{n-k}\pi_x(n)$$

= $Th_k(x) = T\left[x(x-e)...(x-(k-1)e)s^{x-ke}\right],$

where $h_k(u) = u(u-1)...(u-k+1)\mathbf{1}_{[k,\infty)}(u)$.

The next lemma generalizes a well-known fact in Probability theory and will be needed in the proof of Proposition 21 below.

Lemma 20 If x is a natural element in E then

$$Tx = \sum_{n=1}^{\infty} TP(x \ge ne)e.$$

Proof. We have

$$Tx = \sum_{n=1}^{\infty} n\pi_x(n) = \sum_{n=1}^{\infty} \sum_{k=1}^{n} \pi_x(n) = \sum_{k=1}^{\infty} \sum_{n=k}^{\infty} \pi_x(n) = \sum_{k=1}^{\infty} TP(x \ge ke)e,$$

which proves the lemma. \blacksquare

Proposition 21 Let (E, e, T) be a conditional Riesz triple and let x be a natural element in E. For each $n \in \mathbb{N}$ the following are equivalent:

- (i) $Tx^n \in E^u$ (i.e. $x \in L^n(T)$);
- (ii) $g_x^{(n)}(1)$ exists in E^u (in the sense that the left limit $\lim_{s\uparrow 1} g_x^{(n)}(s)$ exists in E^u).

In either case, we have

$$g_x^{(n)}(1) = T \left[x(x-e)...(x-(n-1)e) \right]$$

= $\sum_{k=n}^{\infty} k(k-1)...(k-n+1)\pi_x(k).$

The quantities Tx^n , $n \ge 1$, are called T-moments of x, and the quantities $Tx, Tx(x-e), Tx(x-e)(x-2e), \dots$ are called T-factorials moments of x. **Proof.** Observe that

$$\frac{g_x(s) - g_x(1)}{s - 1} = \frac{e - g_x(s)}{1 - s} = \sum_{n=1}^{\infty} \frac{1 - s^n}{1 - s} \pi_x(n)$$
$$= \sum_{n=1}^{\infty} \left(1 + s + \dots + s^{n-1}\right) \pi_x(n).$$

So according to Monotone Convergence Theorem (Theorem 10) we have

$$\lim_{s \uparrow 1} \frac{g_x(s) - g_x(1)}{s - 1} = \sum_{n=1}^{\infty} \lim_{s \uparrow 1} \left(1 + s + \dots + s^{n-1} \right) \pi_x(n)$$
$$= \sum_{n=1}^{\infty} n \pi_x(n) = Tx.$$

This proves the equivalence for n = 1.

Assume now that the result is true for rank n. Let us prove it for rank n+1. Assume first that $g^{(n+1)}(1)$ exists. Then

$$g^{(n+1)}(1) = \lim_{s\uparrow 1} \frac{g_x^{(n)}(s) - g_x^{(n)}(1)}{s-1} = \lim_{s\uparrow 1} \sum_{k=n}^{\infty} k (k-1) \dots (k-n+1) \frac{(1-s^{n-k})}{1-s} \pi_x(k)$$
$$= (*) \sum_{k=n}^{\infty} \lim_{s\uparrow 1} k (k-1) \dots (k-n+1) (k-n) \pi_x(k) = T [x (x-e) \dots (x-ne)],$$

where the equality (*) follows from Theorem 10. Now it follows from Lyapunov inequality that if $x \in L^q(T)$ then $x \in L^p(T)$ for all $p \leq q$. So as $x \in L^n(T)$ we have $Tx^{n+1} \in E^u$ if and only if $T[x(x-e)...(x-ne)] \in E^u$. This proves the result.

Assume now that $Tx^{n+1} \in E^u$ then by the above remark, $Tx^n \in E^u$ and then by the induction hypothesis, $g^{(n)}(1)$ exists. Also $T[x(x-e)...(x-ne)] \in E^u$, and then the use of Theorem 10 gives again that

$$g^{(n+1)}(1) = T[x(x-e)...(x-ne)] \in E^{u}.$$

The proof is now complete.

Corollary 22 Let (E, e, T) be a conditional Riesz triple and x a natural element in E. Then $Tx = g'_x(1)$ and $Tx^2 = g'_x(1) + g''_x(1)$ if $Tx^2 \in E^u$.

5 Generating function and T-independence

Theorem 23 Let $(x_k)_{1 \le k \le n}$ be a sequence of *T*-independent natural elements in *E*, and put $S_n := \sum_{i=1}^n x_i$, then the *T*-generating function of S_n is given by

$$g_{S_n}(s) = Ts^{S_n} = g_{x_1}(s)...g_{x_n}(s).$$

Proof. It is enough to prove the result for the case n = 2. Let x and y be two natural elements in E that are T-independent. We have

$$g_{x+y}(s) = \sum_{n \ge 0} s^n T P_{x+y=ne} e = \sum_{n \ge 0} s^n \sum_{k=0}^n T P_{y=ke} P_{x+y=ne} e$$

= $\sum_{n \ge 0} s^n \sum_{k=0}^n T P_{y=ke} P_{x=(n-k)e} e = \sum_{n \ge 0} s^n \sum_{i+j=n} T P_{y=je} e.T P_{x=ie} e$
= $\left(\sum_{i \ge 0} s^i T P_{x=ie} e\right). \left(\sum_{j \ge 0} s^j T P_{y=je} e\right) = g_x(s).g_y(s).$

This proves that $g_{x+y}(s) = g_x(s) g_y(s)$ as required.

Let $N, x_0, x_1, x_2, ...$ be natural elements in E with $x_0 = 0$. Put $x_N := \sum_{n\geq 1} P_{N=ne}x_n$, $S_n := \sum_{k=0}^n x_k$ and $S_N := \sum_{n\geq 1} P_{N=ne}S_n$. It follows from these definitions that $P_{N=ne}x_N = P_{N=ne}x_n$ and $P_{N=ne}S_N = P_{N=ne}S_n$.

Lemma 24 For any $f \in L_2(x)$ the following statements hold.

(i) We have for every $n, k \in \mathbb{N}$

$$P_{N=ne}f(x_N) = P_{N=ne}f(x_n).$$

In particular we have

$$P_{N=ne}P_{x_N=ke}e = P_{N=ne}P_{x_n=ke}e.$$

(ii) We have for every k, n in \mathbb{N} ,

$$P_{N=ne}f(S_N) = P_{N=ne}f(S_n)$$
 and $P_{N=ne}P_{S_N=ke}e = P_{N=ne}P_{S_n=ke}e$.

Proof. (i) It suffices to show these equalities for $f = \mathbf{1}_{(-\infty,a]}$, $a \in \mathbb{R}$. By definition of x_N it follows that $P_{N=ne}x_N = P_{N=ne}x_n$. Using this and noting that $f(x) = P_{x \leq ae}e$, we get

$$P_{N=ne}f(x_N) = P_{N=ne}P_{(x_N-ae)^+}^d e = P_{N=ne}\left(I - P_{(x_N-ae)^+}\right)e$$
$$= P_{N=ne} - P_{P_{N=ne}(x_N-ae)^+}e.$$

Now as

$$P_{N=ne} (x_N - ae)^+ = (P_{N=ne} x_N - P_{N=ne} ae)^+ = (P_{N=ne} x_n - P_{N=ne} ae)^+ = P_{N=ne} (x_n - ae)^+$$

we deduce that

$$P_{N=ne}f(x_N) = P_{N=ne} - P_{P_{N=ne}(x_n-ae)^+}e = P_{N=ne}f(x_n).$$

as required. To get the second equality we just apply the first one for $f = \mathbf{1}_{\{k\}}$.

(ii) The proof is similar to (i). \blacksquare

Lemma 25 With the above notations we have:

- (i) x_N and S_N are natural elements in E.
- (*ii*) $S_N = \sum_{k \ge 1} P_{N \ge ke} x_k.$

Proof. (i) We will show the requested property only for x_N ; the case of S_N is similar. By Lemma 24, we get

$$\sum_{k\geq 0} P_{x_N=ke} e = \sum_{k\geq 0} \sum_{n\geq 0} P_{N=ne} P_{x_N=ke} e = \sum_{k\geq 0} \sum_{n\geq 0} P_{N=ne} P_{x_n=ke} e$$
$$= \sum_{n\geq 0} P_{N=ne} \sum_{k\geq 0} P_{x_n=ke} e = \sum_{n\geq 0} P_{N=ne} e = e.$$

This show that x_N is natural.

(ii) By definition of S_N we have

$$S_{N} = \sum_{n \ge 0} P_{N=ne} S_{n} = \sum_{n \ge 0} P_{N=ne} \sum_{k=1}^{n} x_{k}$$

= $\sum_{k \ge 1} \left(\sum_{n \ge k} P_{N=ne} \right) x_{k} = \sum_{k \ge 1} P_{N \ge ke} x_{k}.$

This gives the desired formula. \blacksquare

Theorem 26 Assume that $x = x_1, x_2, ...$ are *T*-equidistributed and *T*-conditional independent of *N*. The generating function of x_N is given by:

$$g_{x_N} = T(I - P_{N=0})g_x + TP_{N=0}e.$$

Moreover, if $TP_{N>0}e = e$ then $g_{x_N} = g_x$.

Proof. We have

$$g_{x_N}(s) = \sum_{n \ge 0} s^n T P_{x_N = ne} e = \sum_{n \ge 0} s^n \sum_{k \ge 0} T P_{N = ke} P_{x_k = ne} e$$

= $\sum_{n \ge 0} s^n T P_{N=0} P_{x_0 = ne} e + \sum_{n \ge 0} s^n \sum_{k \ge 1} T P_{N = ke} . T P_{x_k = ne} e$
= $T P_{N=0} e + \sum_{k \ge 1} T P_{N = ke} . \sum_{n \ge 0} s^n T P_{x = ne} e.$

Now if we assume that $TP_{N>0}e = e$, we get $TP_{N=0}e = 0$, and then $g_{x_N}(s) = Tg_x(s) = g_x(s)$.

Theorem 27 Let (E, e, T) be a conditional Riesz triple. Assume that $x_1, x_2, ...$ are *T*-equidistributed and *T*-conditional independent of *N*. Then

(i) the generating function of S_N is given by the following formula:

$$g_{S_N} = \widetilde{g}_N \circ g_x$$

- (ii) If N and x belong to $L^{1}(T)$, then $TS_{N} = (TN) \cdot (Tx)$.
- (iii) If N and x belong to $L^{2}(T)$, then

$$\operatorname{Var}_{T}(S_{N}) = (TN) \cdot \operatorname{Var}_{T}(x) + \operatorname{Var}_{T}(N) \cdot (Tx)^{2},$$

where,

$$\operatorname{Var}_{T}(y) := T (y - Ty)^{2} = Ty^{2} - (Ty)^{2}$$

Proof. (i) Using Lemma 24.(ii), we get

$$g_{S_N}(s) = \sum_{n \ge 0} TP_{N=ne} s^{S_n} = \sum_{n \ge 0} T\left(s^{S_n} \cdot P_{N=ne} e\right) = \sum_{n \ge 0} Ts^{S_n} \cdot TP_{N=ne} e = \sum_{n \ge 0} (Ts^x)^n \cdot TP_{N=ne} e = \widetilde{g}_N\left(g_x(s)\right).$$

(ii) As S_n and N are T-independent we have for each n, $TP_{N=ne}S_n = TS_n \cdot TP_{N=ne}e$. Hence

$$TS_N = \sum_{n \ge 1} TP_{N=ne} S_n = \sum_{n \ge 1} TS_n \cdot TP_{N=ne} e = \sum_{n \ge 1} nTx \cdot TP_{N=ne} e$$
$$= Tx \cdot \sum_{n \ge 1} nTP_{N=ne} e = Tx \cdot TN.$$

(iii) We have on one hand,

$$\left(TS_N\right)^2 = \left(Tx\right)^2 \cdot \left(TN\right)^2$$

and on the other hand we have

$$T\left(S_{N}^{2}\right) = \sum_{n\geq 1} Tx_{n}^{2} \cdot TP_{N\geq ne}e + \sum_{n\neq m\geq 1} \left(Tx_{n}x_{m}\right) \cdot TP_{N\geq ne}P_{N\geq me}e$$
$$= Tx^{2} \cdot \sum_{n\geq 1} TP_{N\geq ne}e + \left(Tx\right)^{2} \cdot \sum_{n\neq m\geq 1} TP_{N\geq ne}P_{N\geq me}e$$
$$= \left(Tx^{2}\right) \cdot \left(TN+Z\right).$$

It is enough then to show that $Z := \sum_{n \neq m \geq 1} TP_{N \geq ne} P_{N \geq me} e = TN^2 - TN$. To this end observe that

$$Z = \sum_{k=n \lor m, n \neq m} TP_{N \ge ke} e = \sum_{m=2}^{\infty} 2(m-1)TP_{N \ge ke} e$$
$$= 2\sum_{k>1} \left(\sum_{m=1}^{k-1} m\right) P_{N=ke} e = \sum_{k=1}^{\infty} \left(k^2 - k\right) P_{N=ke} e = TN^2 - TN.$$

This completes the proof. \blacksquare

6 Application to convergence

It is easily seen that if x and y are natural elements in E having the same generating function then x and y have the same T-distribution, namely $TP_{x=ke}e = TP_{y=ke}e$ for k = 0, 1, ... Several notions have been generalized from the classical theory of probability and stochastic processes to the setting of Riesz spaces. One of the challenges is to give a reasonable definition of convergence in distribution. Many equivalent definitions are known in the literature. In [18], David Williams defines the convergence in distribution via distribution function: A sequence (X_n) of real random variables with distribution functions (F_{X_n}) is said to converge in distribution to Xif $F_{X_n}(t) \longrightarrow F(t)$ for every $t \in \text{Cont}(F)$, where F_X is the distribution function of X and Cont(F) is the set of points on which F is continuous. According to [3, Example 2.4], there is a Riesz space E and an element xin E such that F_x is discontinuous everywhere. So the definition above can not be translated to the theory of Riesz spaces. We suggest, however, the following definition in the case of natural elements.

Definition 28 Let (E, e, T) be a conditional Riesz triple and (x_{α}) a net of neutral elements in E. We say that the net $(x_{\alpha})_{\alpha \in A}$ converges in T-distribution to a natural element x in E if $TP_{x_{\alpha}=ke}e \xrightarrow{o} TP_{x=ke}e$ for all $k \geq 0$. We write $x_{\alpha} \xrightarrow{T-d} x$.

Proposition 29 Let (E, e, T) be a conditional Riesz triple, (x_{α}) a net of natural elements in E and $x \in E$. Then the following statements are equivalent.

- (i) The net (x_{α}) converges to x in T-distribution.
- (ii) The net $(g_{x_{\alpha}})$ converges to g_x pointwise on [0,1].

Proof. Let us denote $TP_{x_{\alpha}=ke}e$ briefly by $p_{k,\alpha}$, and $TP_{x=ke}e$ by p_k .

(i) \Longrightarrow (ii) The conclusion is trivial for s = 1. Let s be fixed in [0, 1). We have for each k, $\lim_{\alpha} s^k p_{k,\alpha} = s^k p_k$ and $|s^k p_{k,\alpha}| \leq s^k e$. Now as the series $\sum_{k\geq 0} s^k e$ is order convergent in E_e we can apply Theorem 11 to get $g_{x_\alpha}(s) \xrightarrow{o} g_x(s)$. (ii) \Longrightarrow (i) The proof will be done by induction on k. Observe first that $p_0^{(\alpha)} = g_{x_\alpha}(0) \xrightarrow{o} g_x(0) = p_0$. Assume by induction that $p_i^{(\alpha)} \xrightarrow{o} p_i$ for all i = 0, 1, ...k and let $s \in (0, 1)$. Define

$$A_n^{\alpha}(s) = \left| p_{k+1}^{(\alpha)} - p_{k+1} + \sum_{i \ge k+2} \left(p_i^{(\alpha)} - p_i \right) s^{i-(k+1)} \right|.$$

Then

$$A_{n}^{\alpha}(s) = \left| \frac{g_{x_{\alpha}}(s) - \sum_{i=0}^{k} p_{i}^{(\alpha)} s^{i}}{s^{k+1}} - \frac{g_{x}(s) - \sum_{i=0}^{k} p_{i} s^{i}}{s^{k+1}} \right|$$
$$\leq \left| \frac{g_{x_{\alpha}}(s) - g_{x}(s)}{s^{k+1}} \right| + \left| \frac{\sum_{i=0}^{k} p_{i}^{(\alpha)} s^{i} - \sum_{i=0}^{k} p_{i} s^{i}}{s^{k+1}} \right| \xrightarrow{o} 0$$

It follows that

$$\limsup_{\alpha} \left| p_{k+1}^{(\alpha)} - p_{k+1} \right| \le \left(\frac{s}{1-s} \right) e + \limsup_{\alpha} A_n^{\alpha}(s),$$

for all $s \in (0, 1)$. Hence $p_{k+1}^{(\alpha)} \xrightarrow{o} p_{k+1}$ as required and the proof is complete.

Theorem 30 Suppose that $N \sim_T \mathcal{P}(g)$, and that $x_1, x_2, ...$ are *T*-independent, identically distributed with Bernoulli distribution $\mathcal{B}(p)$ and independent of *N*. Let $S_n = \sum_{i=1}^n x_i$. Then S_N has the Poisson distribution with parameter pg.

Proof. By Theorem 27,

$$g_{S_N}(s) = \tilde{g}_N(g_{x_1}(s)) = \exp((g_{x_1}(s) - e)g) = \exp((e - p + sp - e)g) = \exp((-p + sp)g) = \exp(-pg + spg) = \exp(-(1 - s)pg).$$

This shows that S_N has the Poisson distribution with parameter pg as desired.

As the function exp is convex on \mathbb{R} it follows from [11, Corollary 4.3] that exp (x) is well defined in E^s for all $x \in E$. Observe that if x is positive then exp $(x) = \sup_n v_n$, where $v_n = \sum_{k=0}^n \frac{x^k}{k!}$. We will show that exp x is in fact in E^u , the universal completion of E. To this end it is sufficient to prove that (v_n) is unbounded order Cauchy (see [10, Theorem 3.10] or [2, Theorem 17]). Now observe that for $p \leq q$, and $u \in E^+$,

$$|v_p - v_q| \wedge u = P_{(x-te)^+} \left[(v_q - v_p) \wedge u \right] + P_{(x-te)^+}^d \left[(v_q - v_p) \wedge u \right]$$

$$\leq P_{(x-te)^+} u + \sum_{k=p+1}^q \left(t^k / k! \right) e.$$

Hence

$$\limsup_{p,q} \left[|v_p - v_q| \wedge u \right] \le P_{(x-te)^+} u.$$

As this holds for every $t \ge 0$ we deduce that (v_n) is unbounded order Cauchy as required. It should be noted that if $x \in E_e$, the ideal generated by e, then one can work in that ideal which is lattice isomorphic to some C(K)-space (see for instance [13, Theorems 6.2 and 6.8]) and then use different tool which may seem easier. In both cases the formula $\exp(x) \exp(-x) = e$ is valid. In the sequel we extend a useful result verified by the exponentiation to the setting of Riesz spaces.

Lemma 31 Let E be a Dedekind complete Riesz space with weak order unit e and (x_n) a sequence in E such that $x_n^n \in E$ for each n. Then the following hold.

(i) If $x_n \xrightarrow{o} 0$ then $x_n^n \xrightarrow{uo} 0$.

(ii) If
$$x_n \xrightarrow{o} x$$
 in E_e then $\left(1 + \frac{x_n}{n}\right)^n \xrightarrow{o} \exp(x)$ in E^u .

Proof. (i) As e is a weak order unit it is sufficient to prove that $x_n^n \wedge e \stackrel{o}{\longrightarrow} 0$ [10, Corollary 3.5]. Let us fix a real t in (0, 1). As (x_n) is order null, $P_{(x_n-te)^+}e$ is order null as well. We derive that

$$x_n^n \wedge e = (x_n \wedge e)^n = \left(P_{(x_n - e)^+} (x_n \wedge e) + P_{(x_n - te)^+}^d (x_n \wedge e) \right)^n$$

= $P_{(x_n - te)^+} (x_n \wedge e)^n + P_{(x_n - te)^+}^d (x_n \wedge e)^n$
 $\leq P_{(x_n - te)^+} e + P_{(x_n - te)^+}^d x_n^n \leq P_{(x_n - te)^+} e + t^n e.$

This shows that $x_n^n \wedge e \xrightarrow{o} 0$ and then (i) is proved.

(ii) Write $\left(1 + \frac{x_n}{n}\right)^n = \sum_{k=0}^{\infty} f_n(k)$ where $f_n(k) = \begin{cases} \binom{n}{k} \frac{x_n^k}{n^k} & \text{if } 0 \le k \le n\\ 0 & \text{otherwise} \end{cases}.$

Let M be an upper bound of $(|x_n|)$. Then it is quite easy to see that $|f_n(k)| \leq \frac{M^k}{k!}$ and $f_n(k) \xrightarrow{o} \frac{x^k}{k!}$ as $n \to \infty$. It follows from Theorem 11 that $\left(1 + \frac{x_n}{n}\right)^n \xrightarrow{o} e^x$ in E^u as required.

Remark 32 In (i) the order convergence can not be expected; it is only true in E^u as order convergence and unbounded order convergence agree in E^u for sequences by a result of Grobler (see [10, Corollary 3.12]). The order convergence of (x_n) can not guarantee the order boundedness of the sequence (x_n^n) . Take for instance $E = \ell^\infty$ and $x_n = (1 + 1/\sqrt{n}) e_n$, where (e_n) is the standard basis. Then $x_n \stackrel{o}{\longrightarrow} 0$, however $x_n^n \ge \sqrt{n}e_n$, hence (x_n^n) is not order bounded in ℓ^∞ .

The following theorem generalizes a well-known result in probability theory (see for instance, [17, Example 9.5.2]). A Riesz space version of this result has been proved by Kuo, Vardy, and Watson [15, Theorem 5.6], which is a particular case of our result.

Theorem 33 Let (E, e, T) be a conditional Riesz triple and (y_n) a sequence of natural elements in E. Assume that y_n has a Binomial T-distribution $\mathcal{B}(n, p_n)$ such that $np_n \xrightarrow{o} g$ for some g with $0 < g \in E_e^+$. Then (y_n) converges in T-distribution to $\mathcal{P}(g)$, that is,

$$TP_{y_n=ke}e \xrightarrow{o} \frac{1}{k!}g^n \exp\left(-g\right)$$

for all integer k.

Proof. We have for all $s \in [0, 1]$,

$$g_{y_n}(s) = (e - (1 - s) p_n)^n = \left(e - \frac{(1 - s) n p_n}{n}\right)^n,$$

and by applying Lemma 31 to $x_n = (1 - s) n p_n$ and x = (1 - s) g we get

$$g_{y_n}(s) \xrightarrow{o} \exp((s-1)g)$$
, for all $0 \le s \le 1$,

and the desired result follows. \blacksquare

References

- [1] C.D. Aliprantis and O. Burkinshaw, *Positive Operators*, Springer, 2006.
- [2] Y. Azouzi, Completeness for vector lattices, J. Math. Anal. Appl. 472 (2019) 216-230.

- [3] Y. Azouzi and K. Ramdane, On the distribution function with respect to conditional expectation on Riesz space, Quaest. Math. 41 (2) (2018) 257–264
- [4] Y. Azouzi, Y. Nasri, The sup-completion of a Dedekind complete vector lattice, J. Math. Anal. Appl. 506 (2022)
- [5] Y. Azouzi, M. Trabelsi, Lp-spaces with respect to conditional expectation on Riesz spaces, J. Math. Anal. Appl., 447(2017), 798–816.
- [6] M. Α. Ben Amor & А. Omrani (2022):Chernoff's inequality in Riesz spaces, Quaestiones Mathematicae, DOI: 10.2989/16073606.2022.2113474
- [7] G. Buskes, B. de Pagter, A. van Rooij, Functional calculus on Riesz spaces, Indag. Math. 25 (1991) 275–295.
- [8] K. Donner, Extension of Positive Operators and Korovkin Theorems, Springer (1982).
- [9] W. Feller, An introduction to probability theory ant its applications Vol I, 3^e edition
- [10] N. Gao, V. Troitsky, F. Xanthos, Uo-convergence and its applications to Cesàro means in Banach lattices, Israel J. Math. 220 (2017) 649–689.
- [11] J. J. Grobler, Jensen's and martingale inequalities in Riesz spaces, Indag. Math. 25 (2014) 275–295.
- [12] J.J. Grobler, The Kolmogorov-Centsov theorem and Brownian motion in vector lattices, J. Math. Anal. Appl. 410 (2014) 891–901.
- [13] G.L.M. Groenewegen, A.C.M. van Rooij, Spaces of Continuous Functions
- [14] W.-C. Kuo, C.A. Labuschagne, B.A. Watson, Ergodic theory and the strong law of large numbers on Riesz spaces, J. Math. Anal. Appl. 325 (2007) 422–437.
- [15] W.C. Kuo, J. Vardy, B. A. Watson, Bernoulli Processes in Riesz spaces Ordered Structures and Applications: Positivity VII, Trends in Mathematics (2016), 263–274.

- [16] Michael Mitzenmacher, Eli Upfal, Probability and Computing, Cambridge University Press. 2005
- [17] S. I. Resnick, A Probability Path. Birkhäuser, Basel, 2014, reprint of the 2005 edition.
- [18] D. Williams, Probability with martingales, Cambridge 1991.
- [19] A. C Zaanen. Introduction to operator theory in Riesz spaces. Springer Science & Business Media, 2012.