

Better Together: pooling information in likelihood-free inference

David T. Frazier^{*1}, Christopher Drovandi², and David J. Nott^{3,4}

¹Department of Econometrics and Business Statistics, Monash University,
Clayton VIC 3800, Australia

²School of Mathematical Sciences, Queensland University of Technology,
Brisbane 4000 Australia

³Department of Statistics and Data Science, National University of
Singapore, Singapore 117546

⁴Institute of Operations Research and Analytics, National University of
Singapore, Singapore 119077

December 7, 2022

Abstract

Likelihood-free inference (LFI) methods, such as Approximate Bayesian computation (ABC), are now routinely applied to conduct inference in complex models. While the application of LFI is now commonplace, the choice of which summary statistics to use in the construction of the posterior remains an open question that is fraught with both practical and theoretical challenges. Instead of choosing a single vector of summaries on which to base inference, we suggest a new pooled posterior and show how to optimally combine inferences from different LFI posteriors. This pooled approach to inference obviates the need to choose a single vector of summaries, or even a single LFI algorithm, and delivers guaranteed inferential accuracy without requiring the computational resources associated with sampling LFI posteriors in high-dimensions. We illustrate this approach through a series of benchmark examples considered in the LFI literature.

*Corresponding author: david.frazier@monash.edu

1 Introduction

The use of complex models has led researchers to employ statistical methods that are capable of conducting reliable inference in situations where the likelihood function is intractable. In Bayesian inference, this has led to the use of so-called likelihood free inference (LFI) methods, such as approximate Bayesian computation (ABC), which replace evaluation of the likelihood with model simulation; for a review of ABC see the recent handbook [Sisson et al. \(2018\)](#).

LFI assumes that the observed data is drawn from a given class of models from which it is feasible to generate synthetic data. Common implementations of LFI construct an approximate posterior for the model unknowns by comparing, in a given distance, summary statistics calculated using the observed data and data simulated from the model. While such a procedure allows us to conduct statistical inference in very complex models, the broad applicability of such methods has been hindered by the persistent issue of which summaries one should choose in any given situation.

In general, different choices of summaries result in different posteriors, and can sometimes produce surprisingly different (posterior) inferences. This problem is made more difficult by the fact that a direct comparison of different collections of summaries is not necessarily helpful: under regularity conditions, the LFI posterior variance is (weakly) decreasing in the number of summaries used in the analysis. That is, asymptotically, one can never decrease the information in the LFI posterior by adding summaries (see [Frazier](#)

et al., 2018 for details).¹

This asymptotic perspective, however, stands in direct opposition to the practice of LFI: as one adds more summaries, the computational burden increases. Even if the additional summaries are highly informative, controlling the additional Monte Carlo error introduced into the posterior approximation via their inclusion requires an often untenable increase in computational cost.

In this paper, we make three contributions to the literature on LFI. Firstly, rather than choosing summary statistics, we propose to conduct LFI by combining several posteriors built using different sample information. While it may be possible to fuse posteriors in many different ways, our suggested approach uses linear opinion pools (Stone, 1961), due in part to their parsimonious nature and good performance in a myriad of tasks. Linear opinion pools are known to be useful tools for combining priors beliefs, and evidences, across different individuals, and we refer to Evans and Guo (2022) for a discussion of the latter in likelihood-based Bayesian inference. This linear pooling approach allows us to efficiently combine the information from many different sets of summary statistics to produce posterior inference without requiring the computational resources necessary to produce LFI posteriors conditioned on the entire set of summary statistics, thereby circumventing the curse of dimensionality associated with LFI inference methods (see, e.g., Blum, 2010 for a discussion of the curse of dimensionality in LFI).

¹Moreover, this asymptotic viewpoint also disregards finite-sample differences in the locations and scales of posteriors that can result from employing different collections of summaries.

Secondly, we show that this pooling approach can be applied to effortlessly combine inferences from summary-based LFI posteriors and those built using general discrepancy measures, such as the Wasserstein distance (Bernton et al., 2019); see Drovandi and Frazier (2022) for a review of such approaches in LFI. As a consequence, the latter allows us to combine, in a cohesive fashion, and for the first time, LFI posteriors based on summaries, and those based on a general discrepancy metric. Further, it is possible to pool inferences from different LFI algorithms, such as ABC and Bayesian synthetic likelihood (BSL, Price et al., 2018).

Lastly, we show that this pooling approach produces theoretically guaranteed inferential gains over using the individual collections of summaries by themselves; more specifically, we give conditions under which pooled posteriors will always produce inferences that are more precise than if one were to use just the individual summary collections for inference. In principle, the theoretical guarantees obtained in the case of summaries also carry over to the more general case of combining LFI posteriors of different types. However, such a result would require asymptotic normality of the posterior mean from the discrepancy-based posterior, which has not been theoretically verified in the existing literature. As such, we leave a formal analysis for future research.

In addition, we show that in cases where one set of summaries is incompatible with the assumed model (see, e.g., Marin et al., 2012, Frazier et al., 2020 or Section 3.2 for discussion), the pooled posterior automatically disregards the incompatible set of summaries. In particular, when there is a set of compatible summaries in the pool but the other set is

incompatible, the pooled posterior will (asymptotically) place weight unity on the posterior produced using the compatible summaries.

The remainder of the paper proceeds as follows. Section two contains the motivation and general setup. Section three provides the intuition for the pooling approach, along with a naïve implementation method, and some illustrative examples. Theoretical aspects of the pooling approach are also discussed. Section 4 extends the pooling approach to the case of general discrepancy based measures, and demonstrates the appreciable inferential gains that can be obtained in this setting. Section five concludes with a discussion on future work.

2 LFI inference and the choice of summaries

2.1 LFI

For $n \geq 1$, let $(\Omega_n, \mathcal{F}_n, \mathbb{P}_n)$ denote the intrinsic probability space, with associated expectation operator \mathbb{E}_n , on which all random variables are defined. For simplicity of notation, we drop quantities dependence on n when no confusion will result. Denote by $\mathcal{P}(\mathcal{X})$ the set of probability measures on a space \mathcal{X} . We observe data $\mathbf{y} = (y_1, \dots, y_n)^\top \in \mathcal{Y}^n$, distributed according to some unknown measure $P_0^{(n)}$.

Our beliefs about $P_0^{(n)}$ are specified as a class of parametric models $\mathcal{M}^{(n)} = \{P_\theta^{(n)} : \theta \in \Theta\} \subseteq \mathcal{P}(\mathcal{Y}^n)$, where $\Theta \subseteq \mathbb{R}^{d_\theta}$. We quantify our prior beliefs about θ via a prior distribution $\Pi \in \mathcal{P}(\Theta)$. Even if $\mathcal{M}^{(n)}$ is very complex, it is still feasible to generate

synthetic observations \mathbf{z} according to $P_\theta^{(n)}$, for any $\theta \in \Theta$. Thus, even if the likelihood associated with $P_\theta^{(n)}$ is infeasible to calculate, useful information about the model can still be obtained by comparing observed data, \mathbf{y} , against simulated data, \mathbf{z} .

When $P_\theta^{(n)}$ is intractable, likelihood-free inference (LFI) methods can be used to conduct inference on θ by assigning posterior mass to values of θ that produce simulated data \mathbf{z} which is “close-enough” to \mathbf{y} . To make the problem practical from a computational perspective, LFI often resorts to matching low-dimensional summary statistics, defined by the map $S : \mathcal{Y}^n \rightarrow \mathcal{S} \subseteq \mathbb{R}^{d_s}$, and where we require that $d_s \geq d_\theta$. In what follows, when no confusion will result, we let S denote the summary statistic mapping or the mapping evaluated at the observed data \mathbf{y} .

Given statistics S , the goal of LFI is to construct an approximation to the partial posterior $\pi(\theta|S)$. The two most well-established statistical approaches for constructing this posterior approximation are approximate Bayesian computation (ABC), see [Sisson et al. \(2018\)](#) for a review, and Bayesian synthetic likelihood (BSL), see [Wood \(2010\)](#), and [Price et al. \(2018\)](#). ABC and BSL differ in terms of how the posterior is approximated. In the case of ABC, the posterior is approximated by nonparametrically estimating the likelihood within the algorithm. In BSL, we approximate the intractable likelihood of the summaries using a normal density with mean $b(\theta)$ and variance $\Sigma(\theta)$. Since $b(\theta)$ and $\Sigma(\theta)$ are generally unknown, these are subsequently estimated via Monte Carlo using data simulated iid from $P_\theta^{(n)}$. In what follows, we let $\tilde{\pi}(\theta|S)$ denote an arbitrary approximation to the “exact” partial posterior $\pi(\theta|S)$.

2.2 Choosing Summaries for LFI

Accurately approximating $\pi(\theta|S)$ becomes more computationally costly as the dimension of S , d_s , increases. Thus, the problem is to find a collection of summaries that are both low-dimensional and highly-informative about θ . Several methods have been proposed to select summary statistics; we refer to [Blum et al. \(2013\)](#), and [Prangle \(2018\)](#) for in-depth reviews on different strategies. Several approaches are based on searching for informative subsets of summaries using information criteria such as AIC/BIC ([Blum et al., 2013](#)), or entropy ([Nunes and Balding, 2010](#)), while other approaches are based on approximate sufficiency arguments ([Joyce and Marjoram, 2008](#)). In general, while such approaches can be useful, they lack a rigorous theoretical basis.

Alternatively, projection approaches seek to project S into a lower dimension space, and such methods have obtained much popularity in ABC applications. Arguably, the most celebrated of the projection approaches to summary statistic selection is the semi-automatic approach of [Fearnhead and Prangle \(2012\)](#). [Fearnhead and Prangle \(2012\)](#) consider the problem of choosing summaries by attempting to give a decision rule $\delta \in \Theta$ that minimises the posterior expected loss

$$R_S(\theta, \delta) = \int (\theta - \delta)^\top (\theta - \delta) \tilde{\pi}(\theta|S) d\theta. \quad (1)$$

[Fearnhead and Prangle \(2012\)](#) argue that taking $S = \mathbb{E}[\theta|S(\mathbf{y})]$ results in minimizing $R_S(\theta, \delta)$, and propose to estimate this summary using (non)linear regression methods. However, the goal of [Fearnhead and Prangle \(2012\)](#) is not to choose between summaries,

but to approximate the most informative collection given a fixed set of summaries S . In this way, there is no sense in which the use of posterior expected loss should deliver a helpful criterion for deciding amongst competing collections of summaries.²

2.3 Combining Information: Pooled Posteriors

While it is possible to choose a single vector of summaries to conduct inference, we instead suggest to combine posterior inferences based on distinct sets of low-dimensional summary statistics. Such an approach obviates the need to conduct LFI using a high-dimensional vector of summaries, and still allows us to incorporate information contained across different sets of summaries. To make the following discussion as easily interpretable as possible, we restrict our attention to the case where $S = (S_1^\top, S_2^\top)^\top$, and note here that the methods we develop can be extended to more general settings at the loss of interpretability.

Rather than choosing a single set, or attempting to conduct inference on $S = (S_1^\top, S_2^\top)^\top$, we suggest to pool the inferences obtained from $\tilde{\pi}(\theta|S_1)$ and $\tilde{\pi}(\theta|S_2)$ using a linear opinion pool (Stone, 1961):

$$\tilde{\pi}_\omega(\theta|S) := (1 - \omega)\tilde{\pi}(\theta|S_1) + \omega\tilde{\pi}(\theta|S_2),$$

where $\omega \in [0, 1]$ controls the amount of mass assigned to each posterior. In particular, for a fixed pooling weight, ω , the above posteriors can be sampled by generating posterior draws from $\tilde{\pi}(\theta|S_1)$ and $\tilde{\pi}(\theta|S_2)$, and mixing the draws with probability ω . For a fixed

²Additional discussion regarding the inadequacy of $R_S(\theta, \delta)$ as a mechanism for choosing S is given in Appendix A.1.

computational budget, obtaining samples from $\tilde{\pi}(\theta|S_1)$ and $\tilde{\pi}(\theta|S_2)$ separately, which can be done in parallel, will likely be simpler than attempting to approximating the posterior $\tilde{\pi}(\theta|S_1, S_2)$.

To the best of our knowledge, the only other approach that considers a pooled posterior approach in the context of LFI is the work of [Chakraborty et al. \(2022\)](#), in which the authors are concerned with the application of LFI methods in the case of modular inference, and construct a linear pool *over a subset of posterior elements*. [Chakraborty et al. \(2022\)](#) propose to select the pooling weight through prior-to-posterior conflict checks (see, e.g., [Nott et al., 2020](#) for a discussion of such methods). In contrast, we consider an approach that can guaranteed accuracy in terms of frequentist asymptotic risk.

3 Optimality of pooled posteriors

In this section, we define an optimal pooling weight, and describe how it can be estimated. While it is in principle feasible to optimally combine many different sets of summaries, to maintain expositional clarity we only consider optimally combining inferences based on two sets of summaries.

To make the results in this section easier to state and follow, we maintain the following simplifying notations. For $x \in \mathbb{R}^d$, $\|x\|$ denotes the Euclidean norm of x . Throughout, C denotes a generic positive constant that can change with each use. For real-valued sequences $\{a_n\}_{n \geq 1}$ and $\{b_n\}_{n \geq 1}$: for x_n a random variable, $x_n = o_p(a_n)$ if $\lim_{n \rightarrow \infty} \text{pr}(|x_n/a_n| \geq C) = 0$

for any $C > 0$, and $x_n = O_p(a_n)$ if for any $C > 0$ there exists a finite $M > 0$ and a finite n' such that, for all $n > n'$, $\text{pr}(|x_n/a_n| \geq M) \leq C$. All limits are taken as $n \rightarrow \infty$, so that, when there is no confusion, \lim_n denotes $\lim_{n \rightarrow \infty}$. The notation \Rightarrow denotes weak convergence. Let $\text{Int}(\Theta)$ denote the interior of the set Θ . For any matrix $M \in \mathbb{R}^{d \times d}$, we define $|M|$ as the determinant of M , and, let $\lambda_{\max}(M)$ and $\lambda_{\min}(M)$ be the maximal and minimal eigenvalues, respectively. For $f : \mathbb{R}^d \rightarrow \mathbb{R}$ a differentiable function of $x \in \mathbb{R}^d$, we take $\nabla_x f(x)$ to be the gradient and $\nabla_{xx}^2 f(x)$ the Hessian. For a distribution F , we let $\mathbb{E}_F[X]$ denote the expectation of X under F . When confusion is unlikely to result, we use $\mathbb{E}[X]$ to denote the expectation under the true distribution $P_0^{(n)}$. The supplementary material contains proofs of all stated results.

3.1 Optimal ω

To define an optimal weight, first let us follow [Fearnhead and Prangle \(2012\)](#) and consider the problem of choosing summaries by attempting to give a decision rule $\delta \in \Theta$ that minimises the posterior expected loss $\int L(\theta, \delta) \tilde{\pi}(\theta|S) d\theta$, where $L : \Theta \times \Theta \mapsto \mathbb{R}_+$ is a user-chosen loss function of interest, which satisfies the following assumption.

Assumption 1. *For any $\theta, \theta' \in \Theta$, $L(\theta, \theta') = \ell(\|\theta - \theta'\|)$, for some known function $\ell(\cdot)$ such that $\ell(0) = 0$. There exists a $\delta > 0$, such that for all $\theta \in \Theta$ with $\|\theta - \theta_0\| \leq \delta$, $\ell(\|\theta_0 - \theta\|)$ is three times continuously differentiable in θ with: (i) $\nabla_{\theta} \ell(\|\theta - \theta_0\|)|_{\theta=\theta_0} = 0$; (ii) For $H(\theta) = \nabla_{\theta\theta}^2 L(\theta_0, \theta)$, $H_0 := H(\theta_0)$ is positive-definite; (iii) for each $i = 1, \dots, d_{\theta}$, and $\|\theta - \theta_0\| \leq \delta$, $|(\theta - \theta_0)^{\top} \{dH(\theta)/d\theta_i\}(\theta - \theta_0)| \leq M \|\theta - \theta_0\|^2$.*

Under quadratic loss, $\ell(\|\theta - \theta_0\|) = \|\theta - \theta_0\|^2$, [Fearnhead and Prangle \(2012\)](#) show that the posterior mean $\bar{\theta} = \int \theta \tilde{\pi}(\theta|S) d\theta$ yields the smallest posterior expected loss, and, under regularity conditions, this result extends to any loss $L(\cdot, \cdot)$ satisfying Assumption 1; we refer to [Rousseau \(1997\)](#) for details. However, as discussed in Section 2.2 and Appendix A.1, posterior expected loss does not necessarily deliver a helpful criterion for deciding amongst competing collections of summaries.

Herein, we maintain the spirit of the minimum loss suggested in [Fearnhead and Prangle \(2012\)](#), but instead define an optimal pooling weight by minimizing the asymptotic expected loss of the posterior mean for the pooled posterior: letting

$$\bar{\theta}(\omega) := \int_{\Theta} \theta \tilde{\pi}_{\omega}(\theta|S) d\theta$$

denote the posterior mean of the pooled posterior, we define the optimal ω as the value that minimizes the trimmed asymptotic loss of the pooled posterior:

$$\mathcal{R}_0(\omega) := \lim_{\nu \rightarrow \infty} \liminf_{n \rightarrow \infty} \mathbb{E} [\min\{nL\{\theta_0, \bar{\theta}(\omega)\}, \nu\}].$$

The asymptotic expected loss $\mathbb{E} [nL\{\theta_0, \bar{\theta}(\omega)\}]$ is trimmed at ν so that $\mathcal{R}_0(\omega)$ is guaranteed to exist. The optimal pooling weight is then defined as

$$\omega^* := \operatorname{argmin}_{\omega \in [0,1]} \mathcal{R}_0(\omega).$$

3.2 Asymptotic framework: Compatible Summaries

Recalling that $P_0^{(n)}$ denotes the true distribution of \mathbf{y} , we let $G_j^{(n)}$ denote the true distribution of $S_j(\mathbf{y})$, the projection of $P_0^{(n)}$ under $S_j : \mathcal{Y}^n \rightarrow \mathcal{S}_j$. Denote the projection of the

assumed model $P_\theta^{(n)}$, under S_j as $F_{j,n}(\cdot|\theta)$. To characterize ω^* , we consider two distinct situations: the first is where both sets of simulated summaries can match the observed summaries, $S(\mathbf{y})$, which has been termed *compatibility* by [Marin et al. \(2012\)](#) (see, also, [Frazier et al., 2020](#)), and the second is the case where only the first set of summaries is compatible. We treat the incompatible case in Section 3.5, and focus here on the compatible case.

To formally define what we mean by compatible, we require some definitions and regularity conditions, which are similar to those encountered elsewhere in the literature on LFI, and which are assumed to hold for $j = 1, 2$.

Assumption 2. *There exists a vector $b_0 := (b_1^\top, b_2^\top)^\top$, such that, $\|S(\mathbf{y}) - b_0\| = o_p(1)$, and a sequence ν_n diverging to $+\infty$ such that $\nu_n\{S(\mathbf{y}) - b_0\} \Rightarrow N(0, V)$, under $P_0^{(n)}$, for some $d_S \times d_S$ matrix $V = [V_1, \Omega_{1,2}; \Omega_{1,2}^\top, V_2]$.*

Assumption 3. *Let $b_j(\theta)$ denote the mean of $S_j(\mathbf{z})$ under $F_{j,n}(\cdot|\theta)$, with $b(\theta) = (b_1(\theta)^\top, b_2(\theta)^\top)^\top$. The following are satisfied for each j : (i) The mapping $\theta \mapsto b_j(\theta)$ is continuous and injective; (ii) For some matrix function $\theta \mapsto V(\theta)$, continuous and positive-definite for all $\theta \in \Theta$, $\nu_n\{S(\mathbf{z}) - b(\theta)\} \Rightarrow N\{0, V(\theta)\}$, under $P_\theta^{(n)}$.*

Definition (Compatibility). The model $P_\theta^{(n)}$ and summaries, S , are *compatible* if there exist $\theta_0 \in \text{Int}(\Theta)$ such that $b(\theta) = b_0 \iff \theta = \theta_0$.

Under the above assumptions, and additional regularity conditions, it is possible to

show that the posteriors $\tilde{\pi}(\theta|S_1)$ and $\tilde{\pi}(\theta|S_2)$ are asymptotically Gaussian.³ Since these additional regularity conditions are not directly relevant to the form of ω^* , and are specific to the precise LFI method employed in the analysis, we eschew these assumptions in favour of the following high-level regularity condition. To state the condition, let $B_j(\theta) = \nabla_{\theta} b_j(\theta)$, $B_j = \lim_n B_j(\theta_n)$, $\Sigma_j = (B_j^{\top} V_j^{-1} B_j)^{-1}$, and let

$$\Omega_{\Sigma} = Q_1 \Omega_{1,2} Q_2^{\top}, \quad Q_j = \Sigma_j B_j^{\top} V_j^{-1}.$$

Likewise, define the local parameter $t_j = \sqrt{n}(\theta - \theta_0) - Q_j \sqrt{n}\{S_j(\mathbf{y}) - b_j(\theta_0)\}$, and let the posterior for the local parameter be $\pi(t_j|S_j) := \pi[\theta_0 + t_j/\sqrt{n} + Q_j \sqrt{n}\{S_j(\mathbf{y}) - b_j(\theta_0)\}|S_j]$.

Assumption 4 (Limiting Posteriors). *For $\tilde{\pi}(t_j|S_j)$ the posterior for t_j , $\int \|t_j\| |\tilde{\pi}(t_j|S_j) - N(t; 0, \Sigma_j^{-1})| dt = o_p(1)$.*

The above assumptions allow us to deduce a simple form for ω^* in the compatible case.

Lemma 1. *Under Assumptions 1-4, $\mathcal{R}_0(\omega)$ is minimised at ω^* , where*

$$\omega^* = \begin{cases} \frac{\text{tr} H_0 (\Sigma_1 - \Omega_{\Sigma})}{\text{tr} H_0 \Sigma_1 + \text{tr} H_0 \Sigma_2 - 2 \text{tr} H_0 \Omega_{\Sigma}} & \text{if } \text{tr} H_0 \Sigma_1 > \text{tr} H_0 \Omega_{\Sigma} \\ 0 & \text{otherwise} \end{cases}. \quad (2)$$

3.3 Alternative pooling weights

A pooled posterior based on ω^* will (asymptotically) have an expected loss that is (weakly) smaller than either individual posterior; i.e., $\mathcal{R}_0(\omega^*) \leq \min\{\mathcal{R}_0(S_1), \mathcal{R}_0(S_2)\}$. In practice,

³In the case of ABC, this result can be achieved under the assumptions of [Frazier et al. \(2018\)](#), and for the case of BSL, see [Frazier et al. \(2022\)](#).

an estimator of ω^* can be obtained using information from both posteriors, and any consistent estimator for the covariance term Ω_Σ .⁴

In particular, for $\bar{\Omega}_\Sigma$ an estimator of the covariance term, and estimators $\bar{\theta}_j = m^{-1} \sum_{i=1}^m \theta_{j,i}$, $\bar{\Sigma}_j = \frac{1}{m} \sum_{i=1}^m (\theta_{j,i} - \bar{\theta}_j)(\theta_{j,i} - \bar{\theta}_j)^\top$, where $\theta_{j,i} \stackrel{iid}{\sim} \tilde{\pi}(\theta|S_j)$, $i = 1, \dots, m$, we can estimate ω^* using $\hat{\omega}_+^* = \min\{1, \hat{\omega}^*\}$, where, for $\bar{H} = H(\bar{\theta}_1)$,

$$\hat{\omega}^* := \begin{cases} \frac{\text{tr}\bar{H}(\bar{\Sigma}_1 - \bar{\Omega}_\Sigma)}{\text{tr}\bar{H}\bar{\Sigma}_1 + \text{tr}\bar{H}\bar{\Sigma}_2 - 2\text{tr}\bar{H}\bar{\Omega}_\Sigma} & \text{tr}\bar{H}\bar{\Sigma}_1 > \text{tr}\bar{H}\bar{\Omega}_\Sigma \\ 0 & \text{otherwise} \end{cases}.$$

In practice, consistent estimation of Ω_Σ is non-trivial, and estimation of Ω_Σ produces additional noise that can degrade the performance of $\hat{\omega}_+^*$; indeed, the weight estimator must be defined as $\hat{\omega}_+^* = \max\{0, \hat{\omega}^*\}$ since using simple point estimators of the variance matrices does not enforce that the numerator is positive.⁵

Given the difficulties associated with estimation of Ω_Σ , and the ensuing possibility of ill-effects, we also propose alternative sets of combination weights that do not require estimation of Ω_Σ . We consider two such alternatives, the first is precisely the weight $\hat{\omega}^*$

⁴We note that bootstrapping procedures can be used to estimate V_1, V_2 and $\Omega_{1,2}$, while numerical methods can be used to estimate the Jacobian terms B_1 and B_2 . In cases where the summaries are not particularly smooth in the unknown parameters, such as the case of quantiles of the data, the methods developed in [Frazier et al. \(2019\)](#) can be used to consistently estimate these components.

⁵The poor empirical performance of this form of plug-in estimator is so ubiquitous in the literature on forecast combination ([Bates and Granger, 1969](#)), see [Wang et al. \(2022\)](#) for a review, it has been termed the combination puzzle.

but where we artificially set $\overline{\Omega}_\Sigma = 0$, to obtain

$$\widehat{\omega} := \frac{\text{tr} \overline{H} \overline{\Sigma}_1}{\text{tr} \overline{H} \overline{\Sigma}_1 + \text{tr} \overline{H} \overline{\Sigma}_2}.$$

The pooling weight $\widehat{\omega}$ disregards the fact that the posteriors $\tilde{\pi}(\theta|S_1)$ and $\tilde{\pi}(\theta|S_2)$ can have distinct locations. To account for this fact, while incorporating the structure of $\widehat{\omega}$, we also suggest the alternative pooling weight

$$\tilde{\omega} := \frac{\text{tr} \overline{H} \overline{\Sigma}_1}{(\bar{\theta}_1 - \bar{\theta}_2)^\top \overline{H} (\bar{\theta}_1 - \bar{\theta}_2) + \text{tr} \overline{H} \overline{\Sigma}_1 + \text{tr} \overline{H} \overline{\Sigma}_2}.$$

The weight $\tilde{\omega}$ is particularly useful when the summary statistics S_1 are thought to provide reliable inferences, and where we are unsure about the compatibility of the summary statistics S_2 . If the posterior locations differ across the component LFI posteriors, a higher weight is assigned to the S_1 component. This means that the two LFI component posteriors are not treated symmetrically under this pooling weight.

Critically, the alternative pooling weights can be estimated using only samples from the constituent posteriors, and no estimation of Ω_Σ is required. Obtaining the pooled posterior, based on $\widehat{\omega}$ or $\tilde{\omega}$ is as simple as sampling $\tilde{\pi}(\theta|S_1)$ and $\tilde{\pi}(\theta|S_2)$. Furthermore, in the case where the summaries are compatible, the two weights $\widehat{\omega}$ and $\tilde{\omega}$ will agree asymptotically: that is, under our assumptions,

$$\widehat{\omega} = \tilde{\omega} + o_p(1) = \omega_0 + o_p(1), \quad \omega_0 := \frac{\text{tr} H_0 \Sigma_1}{\text{tr} H_0 \Sigma_1 + \text{tr} H_0 \Sigma_2}.$$

While the optimal pooling weight depends on the covariance term Ω_Σ , it is not difficult to see that if $\text{tr} H_0 \Omega_\Sigma$ is small, then the simpler pooling weights will be close to the optimal

weight ω^* . More generally, these simpler weights will always perform better than using S_1 or S_2 alone, in terms of risk, in the following empirically relevant scenarios.

Lemma 2. *If $\text{tr}H_0\Omega_\Sigma \leq \frac{1}{2} \min\{\mathcal{R}_0(S_1), \mathcal{R}_0(S_2)\}$, then $\mathcal{R}_0(\hat{\omega}) \leq \min\{\mathcal{R}_0(S_1), \mathcal{R}_0(S_2)\}$.*

Lemma 2 demonstrates that if the trace of the covariance is negative, or small, then the pooled posterior will perform better than using either collection individually. The above condition can be checked in cases where the posterior covariance can be estimated reliably, however, it is not guaranteed to be satisfied in all settings. In Appendix A.2, we give an example where the pooled posterior is outperformed by a particularly informative collection of summaries, which produces a small posterior variance, and has posterior means that are also well-located. As a consequence, the pooled posterior does not produce more accurate inferences than those based solely on the more informative collection. However, in this example we note that the difference between the pooled results and best performing results, as measured by MSE, are minimal, and the pooled posterior still produces accurate inferences.

3.4 Examples

In this section, we compare the three different suggested choices for the pooled posteriors in two commonly encountered toy examples used in the literature on LFI.

3.4.1 Example: g-and-k

The g-and-k model is often used as a test case in the LFI literature (see, e.g., [Fearnhead and Prangle, 2012](#)), since the likelihood of the model is intractable, and the model is often stated in terms of its quantile function:

$$Q\{z(p); \theta\} = a + b \left[1 + c \frac{1 - \exp\{-gz(p)\}}{1 + \exp\{-gz(p)\}} \right] \{1 + z(p)^2\}^k z(p), \quad (3)$$

where $p \in (0, 1)$, $z(p)$ is the quantile function of the standard normal distribution, and the model parameters are $\theta = (a, b, g, k)^\top$, where the parameter c is fixed at 0.8 (see [Rayner and MacGillivray, 2002](#) for discussion).

Several sets of summary statistics have been postulated for this model, including order statistics, and robust summaries based on quantiles of the data ([Drovandi and Pettitt, 2011b](#)), and auxiliary models ([Drovandi and Frazier, 2022](#)). Herein, we compare the pooled posterior approach based on two different sets of summaries: octiles of the data, and the robust summaries proposed in [Drovandi and Pettitt \(2011b\)](#).⁶

We compare the accuracy of four different pooled posteriors against the individual ABC posteriors built from the robust summaries, denoted by S_1 for reference, and the octiles, denoted as S_2 . The first pooled posterior is based on $\hat{\omega}^*$, where we estimate the variance and covariance matrices in Ω_Σ using a standard iid bootstrap based on 10000 replications of

⁶The robust summaries used in [Drovandi and Pettitt \(2011b\)](#) are given by $S_1 = (S_{11}, S_{12}, S_{13}, S_{14})^\top$, where $S_{11} = L_2$, $S_{12} = L_3 - L_1$, $S_{13} = \frac{L_3 + L_1 - 2L_2}{S_2}$, $S_{14} = \frac{E_7 - E_5 + E_3 - E_1}{S_2}$, and where L_i denotes the i -th quartile and E_i the i -th octile.

the observed data;⁷ the second is based on an equally weighted pool, which takes $\omega = 1/2$; the additional two pooled posteriors are constructed using the estimated naive weights $\hat{\omega}$ and $\tilde{\omega}$, respectively. We simulate 100 synthetic samples of size $n = 1000$ from the g-and-k model under true parameter value $\theta_0 = (a_0, b_0, g_0, k_0)^\top = (3, 1, 2, 0.5)^\top$.⁸ Across each method, we report the following averages across the replications for each parameter: the bias of the posterior mean, the posterior standard deviation, the raw MSE of the marginal posterior mean; the overall MSE, i.e., the sum of raw MSE across the different parameters, is also reported in the table caption. We bold the smallest overall MSE across methods for ease of reference.

The results are presented in Table 1. Across all weighting choices, the pooled posterior approach produces inferences that are more accurate than using either individual posterior. Relative to the robust collection of summaries, the pooled posterior based on $\hat{\omega}^*$ obtains a nearly 51% reduction in the MSE across the experiments, while a 31% reduction was achievable relative to the octiles. Moreover, even larger gains are obtained when using pooled posteriors based on other weighting schemes. Indeed, the equally weighted pool produces the smallest average MSE, but delivers point estimators with larger bias and variance than those obtained under the pooling weight $\tilde{\omega}$.

In general terms, comparing the bias and posterior variances, we see that the pooling approach borrows strength from both posteriors, delivering point estimators with low bias,

⁷The gradients are estimated using central finite differences around the mean of the posterior S_1 .

⁸We obtain posterior samples from $\tilde{\pi}(\theta|S_1)$ and $\tilde{\pi}(\theta|S_2)$ using the ABC-SMC algorithm of [Drovandi and Pettitt \(2011a\)](#).

and a posterior with less uncertainty than either individual posterior.

(A)	S_1			S_2			$\hat{\omega}^*$		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
a	0.0008	0.0491	0.0014	0.0002	0.0179	0.0013	0.0001	0.0394	0.0009
b	0.0237	0.0997	0.0082	0.0125	0.0392	0.0070	0.0139	0.0826	0.0048
g	0.0604	0.2867	0.0543	0.0159	0.1041	0.0330	0.0223	0.2266	0.0223
k	-0.0117	0.1113	0.0112	-0.0167	0.0482	0.0120	-0.0142	0.1010	0.0077
(B)	$\omega = 1/2$			$\hat{\omega}$			$\tilde{\omega}$		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
a	0.0005	0.0443	0.0007	0.0009	0.0434	0.0008	-0.0002	0.0434	0.0008
b	0.0181	0.0947	0.0036	0.0201	0.0920	0.0043	0.0175	0.0920	0.0042
g	0.0382	0.2610	0.0193	0.0450	0.2544	0.0274	0.0372	0.2544	0.0197
k	-0.0142	0.1150	0.0046	-0.0129	0.1112	0.0054	-0.0143	0.1112	0.0062

Table 1: Posterior accuracy results in the g-and-k model under the base set of summaries S_1 (robust summaries), the alternative set S_2 (octiles), and the pooled posteriors (ω). Bias is the bias of the posterior mean for θ_0 across the replications. Std is the average posterior standard deviation across the replications. The average MSE over the replications is: $S_1 - 0.0751$; $S_2 - 0.0553$; $\hat{\omega}^* - 0.0369$; $\omega = 1/2 - \mathbf{0.0283}$; $\hat{\omega} - 0.0380$; $\tilde{\omega} - 0.0309$.

3.4.2 Example: Stochastic Volatility Model

An additional example we consider is the simple stochastic volatility model of order one, where observed data is generated according to

$$y_t = \exp(h_t/2)e_t, \quad h_t = \omega + \rho h_{t-1} + \sigma_v \nu_t, \quad t = 1, \dots, n; \quad (4)$$

and where $(e_t, \nu_t)^\top$ are iid standard normal, and the unknown parameters are $\theta = (\omega, \rho, \sigma_v)^\top$. Our prior distribution for θ is uniform over $(-1, 1) \times (0, 1) \times (0, 1)$.

[Martin et al. \(2019\)](#) demonstrate that useful summary statistics for this model can be obtained by first taking squares and logarithms of the process to notice that

$$y_t^* = \log y_t^2 = \log e_t^2 + \omega + \rho h_{t-1}^2 + \sigma_v \nu_t,$$

which resembles a latent autoregressive process of order one. Consequently, we can use summary statistics, in $\log y_t^2$, that would identify the parameters of an observable autoregressive model. In particular, we consider the following auxiliary autoregressive model:

$$y_t^* = \beta^\top X_t + \epsilon_t, \quad X_t = [1, \log y_{t-1}^2, \log y_{t-2}^2]^\top, \quad t = 3, \dots, n.$$

With $\widehat{\beta}$ the estimated regression coefficient based on the observed data, the sample moments are then taken to be the sample moment conditions from the least squares estimator. We denote the estimated sample score equations from this auxiliary regression model as S_1 .⁹

In addition to sample moments from an auxiliary model, unconditional sample moments for data from the stochastic volatility model are known to provide reliable point estimators of the unknown parameters ([Andersen and Sørensen, 1996](#)), and so matching sample moments of the data should also provide reliable summary statistics. We consider the seven sample moments: four moments based on the absolute value of powers of the observed data, i.e., $|y_t^k|$, $k = 1, 2, 3, 4$, and the first three autocovariances, i.e., $y_t y_{t-k}$, $k = 1, 2, 3$. The sample versions of these moments are denoted by S_2 .

⁹That is, the observed summaries are given by $S_1(\mathbf{y}) = \sum_{t=3}^T X_t (y_t^* - \widehat{\beta}^\top X_t)$, while $S_1(\mathbf{z})$ replaces y_t^* and X_t with their simulated counterparts.

We now apply the pooling approach to 100 synthetic datasets of size $n = 1000$ generated from (4) under the true parameter value $\theta_0 = (-0.74, 0.90, 0.36)^\top$. Similar to the previous experiment, we apply four different pooling approaches: the three estimated weights, and the fixed weight $\omega = 1/2$. The results are presented in Table 2. Similar to the g-and-k example, the pooled posteriors are more accurate than either individual posterior.¹⁰ Relative to the sample moment summaries, S_1 , the pooled posterior based on $\hat{\omega}^*$ obtains a nearly 20% reduction in the MSE across the experiments, while a 66% reduction was achievable relative to the auxiliary moment summaries, S_2 . Furthermore, the reduction in MSE are all greater for the naive weights and the fixed weights.

¹⁰Posterior samples are again generated using the sequential Monte Carlo ABC (ABC-SMC) algorithm of [Drovandi and Pettitt \(2011a\)](#).

(A)	S_1			S_2			$\hat{\omega}^*$		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
ω	-0.0338	0.0814	0.0122	-0.0090	0.1444	0.0101	-0.0204	0.1302	0.0082
ρ	-0.0078	0.0184	0.0003	-0.0014	0.0197	0.0002	-0.0038	0.0196	0.0002
σ_v	-0.0127	0.0970	0.0058	-0.0016	0.0658	0.0028	-0.0048	0.0825	0.0027
(B)	$\omega = 1/2$			$\hat{\omega}$			$\tilde{\omega}$		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
ω	-0.0214	0.1335	0.0067	-0.0279	0.1313	0.0075	-0.0208	0.1313	0.0077
ρ	-0.0046	0.0213	0.0001	-0.0056	0.0204	0.0002	-0.0042	0.0204	0.0002
σ_v	-0.0071	0.0926	0.0025	-0.0085	0.0876	0.0029	-0.0050	0.0876	0.0025

Table 2: Posterior accuracy results in the stochastic volatility model under the base set of summaries S_1 (sample moments), the alternative set S_2 (autoregressive summaries), and the pooled posterior (ω). The remaining information is as in Table 1. The average MSE over the replications is: $S_1 - 0.0183$; $S_2 - 0.0131$; $\hat{\omega}^* - 0.0110$; $\omega = 1/2 - \mathbf{0.0094}$; $\hat{\omega} - 0.0105$; $\tilde{\omega} - 0.0103$.

3.5 Incompatible summaries

We now study the case where only one set of summaries is compatible, while the other is incompatible. We assume that, either by prior knowledge or previous studies, there is a subset S_1 of S , with $S_1 \in \mathcal{S}_1 \subseteq \mathbb{R}^{d_1}$, and $d_1 \geq d_\theta$, that is compatible, but the set S_2 is incompatible: there exist $\theta_0 \in \Theta$ such that $b_1(\theta) = b_{0,1} \iff \theta = \theta_0$, and $b_2(\theta_0) \neq b_{0,2}$.

In this case, we can show that (in large samples) the pooled posterior approach based

on $\tilde{\omega}$, places zero weight on the second set of summaries.

Corollary 1. *Assume that Assumptions 1-4 are satisfied for $\tilde{\pi}(\theta|S_1)$. If there exists some $\theta^* \neq \theta^0$ such that $\sqrt{n}(\bar{\theta}_2 - \theta^*) = O_p(1)$, and $\bar{\Sigma}_2 = \Sigma_2 + o_p(1)$, $\|\Sigma_2\| > 0$, then*

$$\omega^* = 0, \text{ and } \tilde{\omega} = o_p(1).$$

Corollary 1 demonstrates that if the summaries S_1 are compatible, but S_2 are incompatible, then the pooling weight converges to zero in probability; i.e., in large samples the pooled posterior places weight unity on the compatible set S_1 . Of course, such a result requires that S_1 is compatible. Luckily, this assumption is not difficult to check empirically: the summaries are compatible with the data if the observed summaries fall within the high-mass region of posterior support under the posterior predictive distribution (of the summaries). The summaries are then compatible if this behaviour persists as n increases. That is, one only needs to obtain $\tilde{\pi}(\theta|S_1)$ and calculate the posterior predictive distribution of the summaries to see if $S_1(\mathbf{y})$ is indeed compatible. Alternatively, one can use the methods suggested by [Marin et al. \(2012\)](#), [Frazier and Drovandi \(2021\)](#) or [Ramírez-Hassan and Frazier \(2022\)](#) to determine whether or not the summaries S_1 are compatible.

3.6 Example: individual-based model of toad movement

Here we consider the individual-based movement model of Fowler’s Toads (*Anaxyrus fowleri*) of [Marchand et al. \(2017\)](#), which has also been used as an illustrative example in other likelihood-free research (e.g. [Drovandi and Frazier, 2022](#)). Here we only provide minimalist

details of the example and refer to [Marchand et al. \(2017\)](#) and [Drovandi and Frazier \(2022\)](#) for more information.

The model has three parameters, $\theta = (\alpha, \xi, p_0)^\top$. The overnight displacement for each toad is drawn from a Levy alpha-stable distribution, parameterised by α and ξ . [Marchand et al. \(2017\)](#) consider three models for how each toad takes refuge during the day. Here we consider their ‘Model 2’ since there is evidence that the model does not provide a good fit to the data. In this model, each toad will take refuge at the closest refuge site it has previously visited with a probability p_0 , otherwise it will take refuge at the new location. The data consist of GPS location data for 66 toads for 63 days. In [Marchand et al. \(2017\)](#) the data is summarised down to four sets comprising the relative moving distances for time lags of 1, 2, 4, 8 days. For each lag, we record the number of returns and the distances for the non-returns. We further summarise the vector of non-return distances by 11 equally spaced quantiles. For each time lag, there are thus 12 summary statistics (including the number of returns).

We might anticipate that the model can capture data related to a lag of 1 day, but might be sceptical that the model provide a good fit for longer time lags. Thus we run two separate ABC analyses, one which just includes lag 1 summaries and another that includes summaries for the remaining lags. In each case we use SMC ABC to sample the approximate posterior. We find that the observed summaries for lag 1 are compatible with the model, while some summaries for the remaining lags lie in the tail of the posterior distribution of the summaries. The estimated univariate posteriors of the parameters are




Figure 1: Estimated univariate posterior distributions for the parameters of the toad example. Shown are the results for lag 1 summaries (solid black) and the results for the remaining lags (dash blue).

shown in Figure 1. There is some indication of a difference in the posteriors between the two ABC analyses. From pooling the two ABC analyses, an estimated $\tilde{\omega} = 0.061$ is obtained, which suggests placing a large weight on the ABC results based on the compatible lag 1 summaries, consistent with the theoretical results above.

4 Pooling different types of posteriors

Whilst the above analysis has so far focused on combing LFI posteriors built using different summary statistics, the pooled posterior approach is also applicable if we wish to combine summary statistic-based posteriors and posteriors built using general discrepancy measures between the observed and simulated data. Recently, several authors have suggested replacing the distance and summary statistics under which LFI is usually implemented with

distances based on empirical measures. For a review of such methods, we refer to [Drovandi and Frazier \(2022\)](#). The benefit of such methods are that they do not require a choice of summary statistics, however, as documented by [Drovandi and Frazier \(2022\)](#), such methods may deliver inferences that are not as precise as those obtained under an informative set of summaries.

Let $\mathcal{D} : \mathcal{Y}^n \times \mathcal{Y}^n \rightarrow \mathbb{R}_+$ denote a discrepancy function used to measure the difference between the observed data \mathbf{y} and data \mathbf{z} simulated under the model $P_\theta^{(n)}$. An ABC-based posterior for θ under $\mathcal{D}(\mathbf{y}, \mathbf{z})$ can then be sampled using a number of different algorithms, such as accept/reject ABC or Markov chain Monte Carlo ABC (ABC-MCMC). In the experiments that follow we use a tuned version of the ABC-MCMC algorithm, see [Sisson and Fan \(2011\)](#) for a review, to obtain samples from the approximate posterior $\tilde{\pi}(\theta|\mathcal{D})$.

Given a posterior based on summaries S , $\tilde{\pi}(\theta|S)$, and a posterior based on \mathcal{D} , $\tilde{\pi}(\theta|\mathcal{D})$, we can pool the posteriors via

$$\tilde{\pi}_\omega(\theta|\mathbf{y}) := \omega\tilde{\pi}(\theta|S) + (1 - \omega)\tilde{\pi}(\theta|\mathcal{D}).$$

Denoting the estimated posterior variance obtained under $\tilde{\pi}(\theta|\mathcal{D})$ by $\bar{\Sigma}_{\mathcal{D}}$, and $\bar{\Sigma}_S$ that obtained under $\tilde{\pi}(\theta|S)$, estimated pooling weights can be constructed in the same manner as the summary-based case. Namely, considering expected risk, so that $L(\theta, \theta_0) = \|\theta - \theta_0\|^2$, and $H_0 = I$, we can pool posteriors using the weights $\hat{\omega}$ and $\tilde{\omega}$ given earlier, which yields

$$\hat{\omega} = 1 - \frac{\bar{\Sigma}_{\mathcal{D}}}{\bar{\Sigma}_{\mathcal{D}} + \bar{\Sigma}_S}, \quad \text{and} \quad \tilde{\omega} = 1 - \frac{\bar{\Sigma}_{\mathcal{D}}}{(\bar{\theta}_S - \bar{\theta}_{\mathcal{D}})^\top (\bar{\theta}_S - \bar{\theta}_{\mathcal{D}}) \bar{\Sigma}_{\mathcal{D}} + \bar{\Sigma}_S},$$

where $\bar{\theta}_S$ denotes the posterior mean under $\tilde{\pi}(\theta|S)$, and $\bar{\theta}_{\mathcal{D}}$ the posterior mean under $\tilde{\pi}(\theta|\mathcal{D})$.

In the case of standard LFI posteriors, it is not at all clear how to combine inferences based on summaries and general discrepancies. If one were to attempt to construct a combined distance over the summaries and discrepancies, the resulting properties of such a combination are unknown, untested, and presents issues both from a computational, and theoretical standpoint.¹¹ In contrast, it is very simply to sample $\tilde{\pi}(\theta|\mathcal{D})$, and $\tilde{\pi}(\theta|S)$ separately, and fuse them together using $\tilde{\pi}_\omega(\theta|\mathbf{y})$.¹²

4.1 Examples: summaries and discrepancies

We now demonstrate the usefulness of this pooling approach by combining posterior information built across combinations of summaries and discrepancies. In these experiments, we set \mathcal{D} to be the Wasserstein metric, which yields the Wasserstein ABC (W-ABC) posterior studied in [Bernton et al. \(2019\)](#); while other choices are entirely feasible, we maintain this choice as it is a popular metric. In addition, we conduct inference using BSL based on a generic auxiliary model; namely, we consider inference based on the summaries from

¹¹From a practical standpoint, it is not clear how to choose the relative weighting of each component in the combined distance; this can have consequences for the computational behavior of the resulting algorithm.

¹²While the theoretical guarantees we have obtained can, in principle, apply to combining posteriors built from summaries and discrepancies, such a results would require that $\pi(\theta|\mathcal{D})$ satisfies a Bernstein-von Mises type results. However, the validity of such an result is unknown for general choices of \mathcal{D} . Consequently, in what follows our theoretical analysis is based on combining different choices of summaries. We leave the extension of these results to the case of summaries and discrepancies for future research.

a three component Gaussian mixture model. [Drovandi and Frazier \(2022\)](#) demonstrate that this choice performs well across several different experiments in terms of an accuracy comparison across many different likelihood-free approaches.

The choice of BSL for these experiments is deliberate and done to emphasize the practical usefulness of the pooling approach: in the case of BSL, it is not clear how to combine general discrepancies and summaries, since the form of the BSL posterior does not allow the incorporation of discrepancy distances.

4.1.1 Example: g-and-k

For this experiment, we use precisely the same simulated data generated under the g-and-k model in Section 3.4.1, and compare the results for BSL based on the auxiliary model summaries, against those obtained from the W-ABC approach, and the resulting pooled posteriors. We present all the same accuracy information as in Section 3.4.1 in Table 3. However, we note that in the experiments of [Drovandi and Frazier \(2022\)](#), BSL coupled with these particular auxiliary model summaries performed very well, and so we would expect, *a priori*, for the pooling weights to be close to unity across the experiments.

Analyzing Table 3, we see that the pooled posteriors have accuracy measures that are very similar to those obtained from the BSL posterior. While not entirely surprising given the results of [Drovandi and Frazier \(2022\)](#), the results demonstrate that posterior pooling is capable of providing large weight in cases where one set of information is clearly dominant.

(A)	S			\mathcal{D}			N/A		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
a	0.0006	0.0373	0.0013	-0.0024	0.0444	0.0013			
b	0.0103	0.0767	0.0050	0.0161	0.0848	0.0050			
g	0.0200	0.1360	0.0193	0.0712	0.2410	0.0400			
k	-0.0071	0.0461	0.0019	-0.0101	0.0587	0.0023			
(B)	$\omega = 1/2$			$\hat{\omega}$			$\tilde{\omega}$		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
a	-0.0009	0.0413	0.0012	-0.0001	0.0396	0.0012	0.0006	0.0374	0.0013
b	0.0132	0.0814	0.0049	0.0121	0.0795	0.0049	0.0103	0.0768	0.0050
g	0.0456	0.2024	0.0259	0.0311	0.1734	0.0196	0.0201	0.1379	0.0193
k	-0.0086	0.0541	0.0020	-0.0083	0.0512	0.0019	-0.0072	0.0464	0.0019

Table 3: Pooled posterior accuracy results in the g-and-k model under summaries (S) and discrepancies (\mathcal{D}). The remaining information is as in Table 1. The average MSE over the replications is: $S - 0.0275$; $\mathcal{D} - 0.0486$; $\omega = 1/2 - .0341$; $\hat{\omega} - 0.0275$; $\tilde{\omega} - \mathbf{0.0275}$.

4.1.2 Example: M/G/1

An additional example we consider is the M/G/1 queueing model, which is a stochastic single-server queue model with Poisson arrivals and a general service time distribution. We follow existing constructions of this model in the LFI literature and maintain that the service times are $\mathcal{U}(\theta_1, \theta_2)$ (see e.g. [An et al., 2020](#)), while we consider that the inter-arrival times are distributed as $\text{Exp}(\theta_3)$. We take the observed data \mathbf{y} to be the inter-departure times of 51 customers, resulting in 50 observations. We generate 100 synthetic datasets from

this model according to the true parameters $(\theta_1, \theta_2, \theta_3)^\top = (1, 5, 0.2)^\top$. Our prior beliefs on $(\theta_1, \theta_2, \theta_3)$ are given by $\mathcal{U}(0, \min(y_1, y_2, \dots, y_n)) \times \mathcal{U}(0, 10 + \min(y_1, y_2, \dots, y_n)) \times \mathcal{U}(0, 0.5)$.

The summaries used in this example (denoted by S) are again those based on an auxiliary Gaussian mixture, and the discrepancy used (denoted \mathcal{D}) is the Wasserstein distance; i.e., the W-ABC posterior. We note that, in the experiments of [Bernton et al. \(2019\)](#), the W-ABC posterior was shown to perform well against various summary-based counterparts, but in the experiments of [Drovandi and Frazier \(2022\)](#) the BSL posterior based on S_1 performed just as well as the W-ABC posterior. Thus, we expect that the pooling weights between the two posteriors to be non-trivial.

The results across the synthetic datasets are presented in [Table 4](#), and demonstrate that there are (again) appreciable gains to be obtained by using pooled posteriors. In this experiment, using the pooled posterior based on $\tilde{\omega}$ produces a 27% reduction in the risk relative to using the BSL posterior alone, and a 25% reduction in risk relative to using the W-ABC posterior.

(A)	S			\mathcal{D}			N/A		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
θ_1	-0.0627	0.1750	0.0189	-0.1040	0.1845	0.0230			
θ_2	0.0801	0.7260	0.5364	0.3051	0.8476	0.5212			
θ_3	0.0598	0.0281	0.0051	0.0624	0.0332	0.0056			

(B)	$\omega = 1/2$			$\hat{\omega}$			$\tilde{\omega}$		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
θ_1	-0.0834	0.1839	0.0198	-0.0787	0.1815	0.0194	-0.0667	0.1766	0.0190
θ_2	0.1926	0.8383	0.4451	0.1684	0.7898	0.4147	0.0924	0.7384	0.5198
θ_3	0.0611	0.0324	0.0052	0.0611	0.0314	0.0052	0.0605	0.0294	0.0051

Table 4: Pooled posterior accuracy results in the M/G/1 model under summaries (S) and discrepancies (\mathcal{D}). The remaining information is as in Table 1. The average MSE over the replications is: $S - 0.5603$; $\mathcal{D} - 0.5498$; $\omega = 1/2 - 0.4701$; $\hat{\omega} - 0.5439$; $\tilde{\omega} - \mathbf{0.4393}$;

5 Discussion

In this work we propose to combine LFI posteriors based on different summary statistics, or based on summary statistics and general discrepancy measures. A linear opinion pool of the component LFI posteriors is used for the combination, and guaranteed improved performance is achieved for the pooled posterior mean in terms of asymptotic frequentist risk. Additionally, if one of the component LFI posteriors is incompatible (with the model), we demonstrate that this component of the pool will receive zero weight asymptotically. Hence, not only can this pooled posterior produce guaranteed gains over using the individual LFI

posteriors, but it can also be used to guard against the impacts of model incompatibility in LFI, see, e.g., [Frazier et al. \(2020\)](#) and [Frazier et al. \(2021\)](#) for details.

Looking to future work, it is of interest to apply similar methods in the context of modular posterior inferences for LFI ([Chakraborty et al., 2022](#)). For discussions of modular Bayesian inference outside the LFI context see [Liu and Berger \(2009\)](#), [Lunn et al. \(2009\)](#), [Plummer \(2015\)](#), [Jacob et al. \(2017\)](#) and [Carmona and Nicholls \(2020\)](#). In [Chakraborty et al. \(2022\)](#), the authors consider a misspecified model and marginal inferences for a subset φ of the parameters θ . They consider a linear opinion pool as a pooled posterior for φ , with component LFI posteriors employing summary statistics S_1 and S_2 , $S_1 \subset S_2$. The summaries S_1 are chosen to deliver reliable but possibly imprecise inferences about φ , whereas S_2 can deliver more precise inferences, which we feel nevertheless should not be trusted if they are in conflict with the inferences derived from S_1 . [Chakraborty et al. \(2022\)](#) use conflict checks to give an appropriate value for the mixing weight in the pooled posterior, which cannot be used to produce guaranteed inferential gains as is in this current work.

The theory developed here needs to be modified in the case where $S_1 \subset S_2$. In particular, assumptions 2 and 3 in Section 3, which assume that $S = (S_1^\top, S_2^\top)^\top$ has a strictly positive definite limiting covariance matrix under both the true data generating process and under the model, do not hold in this situation. However, perhaps a more significant difficulty is that if the dimension of S_2 is much higher than S_1 , it becomes more delicate to take the different levels of Monte Carlo error in the component LFI posteriors into account in the

estimation of an appropriate mixing weight.

References

- An, Z., Nott, D. J., and Drovandi, C. (2020). Robust Bayesian synthetic likelihood via a semi-parametric approach. *Statistics and Computing*, 30(3):543–557. [29](#)
- Andersen, T. G. and Sørensen, B. E. (1996). GMM estimation of a stochastic volatility model: A Monte Carlo study. *Journal of Business & Economic Statistics*, 14(3):328–352. [20](#)
- Bates, J. M. and Granger, C. W. (1969). The combination of forecasts. *Journal of the Operational Research Society*, 20(4):451–468. [14](#)
- Bernton, E., Jacob, P. E., Gerber, M., and Robert, C. P. (2019). Approximate Bayesian computation with the Wasserstein distance. *Journal of the Royal Statistical Society Series B*, 81(2):235–269. [4](#), [27](#), [30](#)
- Blum, M. G. (2010). Approximate Bayesian computation: A nonparametric perspective. *Journal of the American Statistical Association*, 105(491):1178–1187. [3](#), [44](#)
- Blum, M. G., Nunes, M. A., Prangle, D., and Sisson, S. A. (2013). A comparative review of dimension reduction methods in approximate Bayesian computation. *Statistical Science*, 28(2):189–208. [7](#)
- Browning, A. P., McCue, S. W., Binny, R. N., Plank, M. J., Shah, E. T., and Simpson, M. J. (2018). Inferring parameters for a lattice-free model of cell migration and proliferation using experimental data. *Journal of Theoretical Biology*, 437:251–260. [45](#)
- Carmona, C. and Nicholls, G. (2020). Semi-modular inference: enhanced learning in multi-modular models by tempering the influence of components. In Chiappa, S. and Calandra, R., editors, *Proceedings of the Twenty Third International Conference on Artificial Intelligence and Statistics*, volume 108 of *Proceedings of Machine Learning Research*, pages 4226–4235. PMLR. [32](#)
- Chakraborty, A., Nott, D. J., Drovandi, C., Frazier, D. T., and Sisson, S. A. (2022). Modularized Bayesian analyses and cutting feedback in likelihood-free inference. *arXiv preprint arXiv:2203.09782*. [9](#), [32](#)
- Drovandi, C. and Frazier, D. T. (2022). A comparison of likelihood-free methods with and without summary statistics. *Statistics and Computing*, 32:42. [4](#), [17](#), [23](#), [24](#), [26](#), [28](#), [30](#)
- Drovandi, C. C. and Pettitt, A. N. (2011a). Estimation of parameters for macroparasite population evolution using approximate Bayesian computation. *Biometrics*, 67(1):225–233. [18](#), [21](#), [45](#)
- Drovandi, C. C. and Pettitt, A. N. (2011b). Likelihood-free Bayesian estimation of multivariate quantile distributions. *Computational Statistics & Data Analysis*, 55(9):2541–2556. [17](#)
- Evans, M. and Guo, Y. J. (2022). Combining evidence. *arXiv preprint arXiv:2202.02922*. [3](#)
- Fearnhead, P. and Prangle, D. (2012). Constructing summary statistics for approximate Bayesian computation: semi-automatic approximate Bayesian computation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 74(3):419–474. [7](#), [10](#), [11](#), [17](#), [43](#), [44](#)

- Frazier, D. T. and Drovandi, C. (2021). Robust approximate Bayesian inference with synthetic likelihood. *Journal of Computational and Graphical Statistics*, 30(4):958–976. [23](#)
- Frazier, D. T., Drovandi, C., and Nott, D. J. (2021). Synthetic likelihood in misspecified models: Consequences and corrections. *arXiv preprint arXiv:2104.03436*. [32](#)
- Frazier, D. T., Martin, G. M., Robert, C. P., and Rousseau, J. (2018). Asymptotic properties of approximate Bayesian computation. *Biometrika*, 105(3):593–607. [2](#), [13](#), [44](#)
- Frazier, D. T., Nott, D. J., Drovandi, C., and Kohn, R. (2022). Bayesian inference using synthetic likelihood: asymptotics and adjustments. *Journal of the American Statistical Association*, page in press. [13](#), [36](#)
- Frazier, D. T., Oka, T., and Zhu, D. (2019). Indirect inference with a non-smooth criterion function. *Journal of Econometrics*, 212(2):623–645. [14](#)
- Frazier, D. T., Robert, C. P., and Rousseau, J. (2020). Model misspecification in approximate Bayesian computation: consequences and diagnostics. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*. [4](#), [12](#), [32](#)
- Jacob, P. E., Murray, L. M., Holmes, C. C., and Robert, C. P. (2017). Better together? Statistical learning in models made of modules. *arXiv:1708.08719*. [32](#)
- Joyce, P. and Marjoram, P. (2008). Approximately sufficient statistics and Bayesian computation. *Statistical Applications in Genetics and Molecular Biology*, 7(1). [7](#)
- Lehmann, E. L. and Casella, G. (2006). *Theory of point estimation*. Springer Science & Business Media. [38](#)
- Liu, F. Bayarri, M. J. and Berger, J. O. (2009). Modularization in Bayesian analysis, with emphasis on analysis of computer models. *Bayesian Analysis*, 4(1):119–150. [32](#)
- Lunn, D., Best, N., Spiegelhalter, D., Graham, G., and Neuenschwander, B. (2009). Combining MCMC with ‘sequential’ PKPD modelling. *Journal of Pharmacokinetics and Pharmacodynamics*, 36:19–38. [32](#)
- Marchand, P., Boenke, M., and Green, D. M. (2017). A stochastic movement model reproduces patterns of site fidelity and long-distance dispersal in a population of Fowler’s toads (*Anaxyrus fowleri*). *Ecological Modelling*, 360:63–69. [23](#), [24](#)
- Marin, J.-M., Pudlo, P., Robert, C. P., and Ryder, R. J. (2012). Approximate Bayesian computational methods. *Statistics and Computing*, 22(6):1167–1180. [4](#), [12](#), [23](#)
- Martin, G. M., McCabe, B. P., Frazier, D. T., Maneesoonthorn, W., and Robert, C. P. (2019). Auxiliary likelihood-based approximate Bayesian computation in state space models. *Journal of Computational and Graphical Statistics*, 28(3):508–522. [20](#)
- Nott, D. J., Wang, X., Evans, M., and Englert, B.-G. (2020). Checking for prior-data conflict using prior-to-posterior divergences. *Statistical Science*, 35(2):234–253. [9](#)

- Nunes, M. A. and Balding, D. J. (2010). On optimal selection of summary statistics for approximate Bayesian computation. *Statistical Applications in Genetics and Molecular Biology*, 9(1). 7
- Plummer, M. (2015). Cuts in Bayesian graphical models. *Statistics and Computing*, 25:37–43. 32
- Prangle, D. (2018). Summary statistics. pages 125–152. 7
- Price, L. F., Drovandi, C. C., Lee, A., and Nott, D. J. (2018). Bayesian synthetic likelihood. *Journal of Computational and Graphical Statistics*, 27(1):1–11. 4, 6, 46
- Priddle, J. W., Sisson, S. A., Frazier, D. T., and Drovandi, C. (2022). Efficient Bayesian synthetic likelihood with whitening transformations. *Journal of Computational and Graphical Statistics*, 31(1):50–63. 45
- Ramírez-Hassan, A. and Frazier, D. T. (2022). Testing model specification in approximate bayesian computation. *arXiv preprint arXiv:2210.12589*. 23
- Rayner, G. D. and MacGillivray, H. L. (2002). Numerical maximum likelihood estimation for the g-and-k and generalized g-and-h distributions. *Statistics and Computing*, 12(1):57–75. 17
- Rousseau, J. (1997). Asymptotic Bayes risks for a general class of losses. *Statistics & Probability Letters*, 35(2):115–121. 11
- Sisson, S. A. and Fan, Y. (2011). Likelihood-free MCMC. *Handbook of Markov Chain Monte Carlo*, pages 313–335. 26
- Sisson, S. A., Fan, Y., and Beaumont, M. (2018). *Handbook of Approximate Bayesian Computation*. Chapman and Hall/CRC, New York. 2, 6
- Stone, M. (1961). The opinion pool. *The Annals of Mathematical Statistics*, pages 1339–1342. 3, 8
- Wang, X., Hyndman, R. J., Li, F., and Kang, Y. (2022). Forecast combinations: an over 50-year review. *arXiv preprint arXiv:2205.04216*. 14
- Wood, S. N. (2010). Statistical inference for noisy nonlinear ecological dynamic systems. *Nature*, 466(7310):1102–1104. 6

A Proofs of Main Results

In this section, we prove the main results stated in the paper. However, before doing so, we state a few useful lemmas that allow us to simplify the proofs of certain results.

A.1 Lemmas

The following lemma gives the asymptotic distribution of the posterior means $\bar{\theta}_1 := \mathbb{E}_{\tilde{\pi}(\theta|S_1)}[\theta]$ and $\bar{\theta}_2 := \mathbb{E}_{\tilde{\pi}(\theta|S_2)}[\theta]$ in the compatible case. This result is used in the proof of Lemma 1.

Lemma 3. *Under Assumption 2-4, for $\Omega_\Sigma := Q_2\Omega_{2,1}Q_1^\top$ and $V_\Sigma = [\Sigma_1, \Omega_\Sigma^\top; \Omega_\Sigma, \Sigma_2]$, $(\sqrt{n}(\bar{\theta}_1 - \theta_0)^\top, \sqrt{n}(\bar{\theta}_2 - \theta_0)^\top)^\top \Rightarrow (\xi + \tau)$ where $\xi = (\xi_1^\top, \xi_2^\top)^\top \sim N(0, V_\Sigma)$ and $\tau = (0^\top, [Q_2\tau_2]^\top)^\top$.*

Proof of Lemma 3. The result follows from Assumption 2-4 and similar arguments to and Corollary 1 in Frazier et al. (2022). In particular, following the arguments in Corollary 1 of Frazier et al. (2022), for $Z_{n,j} = Q_j\sqrt{n}\{S_j(\mathbf{y}) - b_j(\theta_0)\}$,

$$\bar{\theta}_j = \int \theta \tilde{\pi}(\theta|S_j) d\theta = \int (\theta_0 + t_j/\sqrt{n} + Z_{n,j}/\sqrt{n}) \tilde{\pi}(t|S_j) dt$$

so that

$$\begin{aligned} \sqrt{n}(\bar{\theta}_j - \theta_0) - Z_{n,j} &= \int t_j \tilde{\pi}(t|S_j) dt \\ &= \int t_j \{\tilde{\pi}(t|S_j) - N(t_j; 0, \Sigma_j)\} dt_j + \int t_j N(t_j; 0, \Sigma_j) dt_j \end{aligned}$$

The second term is zero by definition, while the first term can be bounded as

$$\int t_j \{\tilde{\pi}(t|S_j) - N(t_j; 0, \Sigma_j)\} dt_j \leq \int \|t_j\| |\{\tilde{\pi}(t|S_j) - N(t_j; 0, \Sigma_j)\}| dt_j = o_p(1)$$

where the $o_p(1)$ term follows by Assumption 4.

Thus, it follows that

$$\sqrt{n}(\bar{\theta}_1 - \theta_0) - Q_1\sqrt{n}\{S_1(\mathbf{y}) - b_1(\theta_0)\} = o_p(1)$$

$$\sqrt{n}(\bar{\theta}_2 - \theta_0) - Q_2\sqrt{n}\{S_2(\mathbf{y}) - b_2(\theta_0)\} = o_p(1)$$

However, under Assumptions 2 and 3,

$$\begin{aligned} \sqrt{n}\{S_2(\mathbf{y}) - b_2(\theta_0)\} &= \sqrt{n}\{S_2(\mathbf{y}) - b_{2,0}\} + \sqrt{n}\{b_2(\theta_0) - b_{2,0}\} \\ &= \sqrt{n}\{S_2(\mathbf{y}) - b_{2,0}\} + \sqrt{n}\delta_{2,n} \\ &\Rightarrow \mathcal{N}\{\tau_2, V_2\}, \end{aligned}$$

where the second line follows from the convergence in Assumption 2. From the joint convergence of $S = (S_1^\top, S_2^\top)^\top$ in Assumption 2, the stated joint convergence then follows. \square

To deduce the form of ω^* , it is useful to first deduce the asymptotic expected loss associated with S_1 and S_2 : $\mathcal{R}_0(S_j) = \lim_{\nu \rightarrow \infty} \liminf_{n \rightarrow \infty} \mathbb{E} [\min\{nL\{\theta_0, \bar{\theta}_j\}, \nu\}]$.

Lemma 4. *Consider that Assumptions 1-4 are satisfied, and let $H_0 = H(\theta_0)$. If $\|\tau_2\| < \infty$, $\mathcal{R}_0(S_1) = \text{tr}\{H_0\Sigma_1\}$ and $\mathcal{R}_0(S_2) = \text{tr}\{H_0\Sigma_2\} + \tau_2^\top Q_2^\top H_0 Q_2 \tau_2$.*

Proof of Lemma 4. For $\bar{\theta}$ denoting $\bar{\theta}_1$ or $\bar{\theta}_2$, a second-order Taylor expansion of $L(\theta_0, \bar{\theta})$ around θ_0 , with Lagrange remainder term ϑ satisfying $\|\vartheta - \theta_0\| \leq C\|\theta - \theta_0\|$ for some

$C > 0$, yields

$$\begin{aligned} L(\theta_0, \bar{\theta}) &= L(\theta_0, \theta_0) + \partial L(\theta_0, \theta_0) / \partial \theta^\top (\bar{\theta} - \theta_0) + \frac{1}{2} (\bar{\theta} - \theta_0)^\top H(\theta_0) (\bar{\theta} - \theta_0) \\ &\quad + \frac{1}{2} (\bar{\theta} - \theta_0)^\top [H(\vartheta) - H(\theta_0)] (\bar{\theta} - \theta_0) \\ &\leq \frac{1}{2} \|(\bar{\theta} - \theta_0)\|_{H(\theta_0)}^2 + M \|(\bar{\theta} - \theta_0)\|^3, \end{aligned}$$

where the second line follows from Assumption 1 and the definition of the intermediate value. Hence,

$$nL(\theta_0, \bar{\theta}) = \frac{1}{2} \{\sqrt{n}(\bar{\theta} - \theta_0)\}^\top H(\theta_0) \{\sqrt{n}(\bar{\theta} - \theta_0)\} + o(\|\{\sqrt{n}(\bar{\theta} - \theta_0)\}\|^2)$$

Define $Y_{j,n} := \sqrt{n}(\bar{\theta} - \theta_0)$, and note that, by Lemma 3,

$$Y_{j,n} \Rightarrow Y := \begin{cases} N(0, \Sigma_1) & \text{if } j = 1 \\ N(Q_2 \tau_2, \Sigma_2) & \text{if } j = 2 \end{cases}.$$

For $Q_{j,n} := \|Y_{j,n}\|_H^2$, let $Y_{j,n,\zeta} = Y_{j,n} \mathbb{I}[Q_{j,n} \leq \zeta] + \zeta \mathbb{I}[Q_{j,n} > \zeta]$. By Theorem 1.8.8 of Lehmann and Casella (2006),

$$\lim_{n \rightarrow \infty} \mathbb{E} [\|Y_{j,n,\zeta}\|_{H(\theta_0)}^2] = \mathbb{E} [\|Y_j\|_{H_0}^2 \mathbb{I}(\|Y_j\|_{H_0}^2 \leq \zeta)] + \zeta^2 \Pr(\|Y_j\|_{H_0}^2 > \zeta).$$

For $\zeta \rightarrow \infty$, the RHS of the above converges to

$$\mathbb{E}[\|Y_j\|_{H_0}^2] = \begin{cases} \text{tr} H_0 \Sigma_1 & \text{if } j = 1 \\ \tau_2^\top Q_2^\top H_0 Q_2 \tau_2 + \text{tr} H_0 \Sigma_2 & \text{if } j = 2 \end{cases}.$$

□

A.2 Proofs of Main Results

Proof of Lemma 1. Write

$$\sqrt{n}\{\bar{\theta}(\omega) - \theta_0\} = \sqrt{n}\{(1 - \omega)\bar{\theta}_1 + \omega\bar{\theta}_2 - \theta_0\} = (1 - \omega)\sqrt{n}\{\bar{\theta}_1 - \theta_0\} + \omega\sqrt{n}\{\bar{\theta}_2 - \theta_0\}.$$

Recall the definitions of $Q_j = \Sigma_j B_j^\top V_j^{-1}$, and let $Q_1^* = [Q_1 : \mathbf{0}_{d_\theta \times d_{s_2}}]$ and $Q_2^* = [\mathbf{0}_{d_\theta \times d_{s_1}} : Q_2]$.

For $Z_n = \sqrt{n}\{S(\mathbf{y}) - b(\theta_0)\}$, under Assumptions 2, by Lemma 3,

$$\sqrt{n}\{\bar{\theta}(\omega) - \theta_0\} = (1 - \omega)Q_1^*Z_n + \omega Q_2^*Z_n \Rightarrow Y(\omega) := (1 - \omega)Q_1^*M + \omega Q_2^*M$$

where $M \sim N(\xi, V_{1,2})$ with $\xi = (0^\top, \tau_2^\top)^\top$, and $V_{1,2} = \text{Var}[\sqrt{n}\{S(\mathbf{y}) - b(\theta_0)\}]$.

Following similar arguments to the proof of Lemma 4 yields $\mathcal{R}_0(\omega) = \mathbb{E}[\|Y(\omega)\|_{H_0}^2]$, and writing out $\|Y(\omega)\|_{H_0}^2$, we have

$$\|Y(\omega)\|_{H_0}^2 = (1 - \omega)^2\|Q_1^*M\|_{H_0}^2 + \omega^2\|Q_2^*M\|_{H_0}^2 + 2\omega(1 - \omega)(Q_1^*M)^\top H_0 Q_2^*M.$$

The result follows by taking the expectations of each term, and solving for the optimal ω .

For the first term, write $\|Q_1^*M\|_{H_0}^2 = \|Q_1^*(M - \xi) + Q_1^*\xi\|_{H_0}^2$, and note that $Q_1^*\xi = 0$.

Hence,

$$\mathbb{E}[\|Q_1^*(M - \xi)\|_{H_0}^2] = \text{tr}H_0 Q_1^* V_{1,2} (Q_1^*)^\top = \text{tr}H_0 Q_1 V_1 Q_1^\top = \text{tr}H_0 [B_1^\top V_1^{-1} B_1]^{-1},$$

where the last equality follows from the definition of Q_1 . Applying a similar approach to the second term yields

$$\mathbb{E}\|Q_2^*M\|_{H_0}^2 = \text{tr}H_0 Q_2 V_2 Q_2^\top + \tau_2^\top Q_2^\top H_0 Q_2 \tau = \text{tr}H_0 [B_2^\top V_2^{-1} B_2]^{-1} + \tau_2^\top Q_2^\top H_0 Q_2 \tau$$

For the last term, write

$$\begin{aligned}
\mathbb{E} [(Q_1^* M)^\top H_0 \{Q_2^* M\}] &= \mathbb{E} \text{tr} [H_0 \{Q_2^* M\} M^\top Q_1^{*\top}] \\
&= \text{tr} H_0 Q_2^* \mathbb{E} [M M^\top] Q_1^{*\top} \\
&= \text{tr} H_0 Q_2^* \{\xi \xi^\top + V_{1,2}\} Q_1^{*\top} \\
&= \text{tr} H_0 Q_2 \Omega_{2,1} Q_1^\top
\end{aligned}$$

where we have used $Q_1^* \xi = 0$.

We recall the following notations: $\Sigma_1 := [B_1^\top V_1^{-1} B_1]^{-1}$ and $\Sigma_2 = [B_2^\top V_2^{-1} B_2]^{-1}$. Using this, and the above expectations, $\mathcal{R}_0(\omega)$ becomes

$$\begin{aligned}
\mathcal{R}_0(\omega) &= (1 - \omega)^2 \text{tr} H_0 \Sigma_1 + \omega^2 [\text{tr} H_0 \Sigma_2 + \tau_2^\top Q_2^\top H_0 Q_2 \tau_2] + 2\omega(1 - \omega) \text{tr} H_0 \Omega_\Sigma \\
&\equiv (1 - \omega)^2 \mathcal{R}_0(1) + \omega^2 \mathcal{R}_0(2) + 2\omega(1 - \omega) \text{tr} H_0 \Omega_\Sigma
\end{aligned}$$

To maximize $\mathcal{R}_0(\omega)$ over $\omega \in [0, 1]$ we consider the Lagrangian

$$\mathcal{L}(\omega, \lambda) = \mathcal{R}_0(\omega) + \lambda(1 - \omega),$$

where λ is the multiplier associated to the constraint $(1 - \omega) \geq 0$.

First, consider that ω^* is in the interior of the space, i.e., $0 < \omega^* < 1$. Differentiating the above wrt ω and solving for ω as a function of λ yields the solution:

$$\omega^*(\lambda) = \frac{\lambda}{2\mathcal{J}} + \frac{\mathcal{R}_0(1) - \text{tr} H_0 \Omega_\Sigma}{\mathcal{J}} = \frac{\lambda + 2(\mathcal{R}_0(1) - \text{tr} H_0 \Omega_\Sigma)}{2\mathcal{J}}, \quad (5)$$

where

$$\mathcal{J} := \mathcal{R}_0(1) + \mathcal{R}_0(2) - \text{tr} H_0 \Omega_\Sigma = \text{tr} H_0 \{\Sigma_1 + \Sigma_2 - \Omega_\Sigma\} + \tau_2^\top Q_2^\top H_0 Q_2 \tau_2.$$

The solution $\omega^*(\lambda)$ must obey the complementary slackness condition

$$0 = \lambda(1 - \omega^*(\lambda)) \quad (6)$$

which, for $0 < \omega^* < 1$, is satisfied only at $\lambda^* = 0$.

Plugging in $\lambda^* = 0$ into equation (5), we see that this solution is feasible only when

$$\mathcal{R}_0(1) - \text{tr}H_0\Omega_\Sigma = \text{tr}H_0\Sigma_1 - \text{tr}H_0\Omega_\Sigma > 0, \quad (7)$$

else the solution $\omega^* = \omega^*(0) \leq 0$, violates the constraint $\omega^* \geq 0$. Therefore, when (7) is satisfied we have

$$\omega^* = \omega^*(0) = \frac{\mathcal{R}_0(1) - \text{tr}H_0\Omega_\Sigma}{\mathcal{R}_0(S_1) + \mathcal{R}_0(S_2) - 2\text{tr}H_0\Omega_\Sigma} \equiv \frac{\text{tr}H_0(\Sigma_1 - \Omega_\Sigma)}{\text{tr}H_0(\Sigma_1 + \Sigma_2 - 2\Omega_\Sigma) + \tau_2^\top Q_2^\top H_0 Q_2 \tau_2},$$

which yields the first claimed solution.

Consider that the condition in (7) is violated. Then, for $C = [\mathcal{R}_0(1) - \text{tr}H_0\Omega_\Sigma] \leq 0$, and

$$\begin{aligned} \mathcal{R}_0(\omega) &= \mathcal{R}_0(1) - 2\omega[\mathcal{R}_0(1) - \text{tr}H_0\Omega_\Sigma] + \omega^2 [\mathcal{R}_0(S_1) + \mathcal{R}_0(S_2) - 2\text{tr}H_0\Omega_\Sigma] \\ &= \mathcal{R}_0(1) + \omega \underbrace{[-2\omega C]}_{>0} + \omega^2 \underbrace{\mathcal{J}}_{>0}. \end{aligned}$$

From the above, we see that $\mathcal{R}_0(\omega)$ is minimized at $\omega^* = 0$, which yields the minimal asymptotic expected loss, and the second claimed solution. \square

Proof of Lemma 2. It follows directly from Lemma 3, and Lemma 1 that $\mathcal{R}_0(\hat{\omega}) = \mathcal{R}_0(\omega_0) +$

$o_p(1)$. Now, recall that

$$\begin{aligned}\mathcal{R}_0(\omega) &= (1 - \omega)^2 \text{tr}H_0\Sigma_1 + \omega^2 [\text{tr}H_0\Sigma_2] + 2\omega(1 - \omega)\text{tr}H_0\Omega_\Sigma \\ &\equiv (1 - \omega)^2\mathcal{R}_0(1) + \omega^2\mathcal{R}_0(2) + 2\omega(1 - \omega)\text{tr}H_0\Omega_\Sigma.\end{aligned}$$

Under $\omega = \omega_0$, and for $C_0 = \mathcal{R}_0(1) + \mathcal{R}_0(2)$, we can rewrite the above as

$$\mathcal{R}_0(\omega_0) = \frac{\mathcal{R}_0(2)^2}{C_0^2}\mathcal{R}_0(1) + \frac{\mathcal{R}_0(1)^2}{C_0^2}\mathcal{R}_0(2) + 2\frac{\mathcal{R}_0(1)\mathcal{R}_0(2)}{C_0^2}\text{tr}H_0\Omega_\Sigma. \quad (8)$$

Consider that $\min\{\mathcal{R}_0(1), \mathcal{R}_0(2)\} = \mathcal{R}_0(1)$, and using equation (8) to rewrite $\mathcal{R}_0(w_0) - \mathcal{R}_0(1)$ as

$$\mathcal{R}_0(w_0) - \mathcal{R}_0(1) = \frac{\mathcal{R}_0(1)}{C_0^2} [\mathcal{R}_0(2)^2 + \mathcal{R}_0(1)\mathcal{R}_0(2) + 2\mathcal{R}_0(2)\text{tr}H_0\Omega_\Sigma - \{\mathcal{R}_0(1) + \mathcal{R}_0(2)\}^2].$$

Using the definition of C_0^2 and completing the square, we see that

$$\begin{aligned}\mathcal{R}_0(w_0) - \mathcal{R}_0(1) &= \frac{\mathcal{R}_0(1)}{C_0^2} [C_0^2 - \mathcal{R}_0(1)^2 - \mathcal{R}_0(1)\mathcal{R}_0(2) + 2\mathcal{R}_0(2)\text{tr}H_0\Omega_\Sigma - C_0^2] \\ &= -\frac{\mathcal{R}_0(1)}{C_0^2} [\mathcal{R}_0(1)^2 + \mathcal{R}_0(1)\mathcal{R}_0(2) - 2\mathcal{R}_0(2)\text{tr}H_0\Omega_\Sigma]\end{aligned} \quad (9)$$

Since $\text{tr}H_0\Omega_\Sigma < 0$, the RHS of the above is negative and $\mathcal{R}_0(w_0) - \mathcal{R}_0(1) \leq 0$.

In the case where $\min\{\mathcal{R}_0(1), \mathcal{R}_0(2)\} = \mathcal{R}_0(2)$, repeating the above steps yields

$$\mathcal{R}_0(w_0) - \mathcal{R}_0(2) = -\frac{\mathcal{R}_0(2)}{C_0^2} [\mathcal{R}_0(2)^2 + \mathcal{R}_0(1)\mathcal{R}_0(2) - 2\mathcal{R}_0(1)\text{tr}H_0\Omega_\Sigma].$$

Thus, a sufficient condition for $\mathcal{R}_0(w_0) - \mathcal{R}_0(2)$ is that $[\mathcal{R}_0(2) - 2\text{tr}H_0\Omega_\Sigma] \geq 0$. Hence, so long as $\text{tr}H_0\Omega_\Sigma \leq \frac{1}{2} \min\{\mathcal{R}_0(1), \mathcal{R}_0(2)\}$ the result follows. □

A Supplementary Material

A.1 Additional Discussion on Summary Selection

Recall that [Fearnhead and Prangle \(2012\)](#) consider the problem of choosing summaries by attempting to give a decision rule $\delta \in \Theta$ that minimises the posterior expected loss

$$R_S(\theta, \delta) = \int (\theta - \delta)^\top (\theta - \delta) \tilde{\pi}(\theta|S) d\theta. \quad (10)$$

Viewing the above loss as a function of arbitrary summaries S , [Fearnhead and Prangle \(2012\)](#) argue that taking $S = \mathbb{E}[\theta|S(\mathbf{y})]$ results in minimizing $R_S(\theta, \delta)$. To estimate this summary statistic [Fearnhead and Prangle \(2012\)](#) propose the use of (non)linear regression methods on (powers of) the summary statistics. That is, given training data $\{\theta, S(\mathbf{z})\}$, [Fearnhead and Prangle \(2012\)](#) use as a summary statistic the fitted regression function evaluated at $S(\mathbf{y})$.

It is important to realize that the goal of [Fearnhead and Prangle \(2012\)](#), is not to choose between summaries, but to approximate the most informative collection given a fixed set of summaries S . In this way, there is no sense in which the use of posterior expected loss should deliver a helpful criterion for deciding amongst competing collections of summaries S_1 and S_2 . Indeed, the minimum of $R_{S_j}(\theta, \delta)$ is obtained at $\delta = \bar{\theta}_j = \int \theta \tilde{\pi}(\theta|S_j) d\theta$, for each S_j , and, under quadratic loss, $R_{S_j}(\theta, \bar{\theta}_j) \approx \text{tr} [B_j^\top V_j^{-1} B_j]^{-1} / n$ (i.e., the asymptotic variance of the posterior). Consequently, choosing summaries by comparing $R_{S_j}(\theta, \bar{\theta}_j)$ would lead us to choose whichever collection of summaries delivered the smallest posterior variance. This is not a helpful selection criterion since, under weak regularity conditions, the posterior

variance of $\pi(\theta|S)$ is (asymptotically) a decreasing function of the number of summaries in S ; that is, asymptotically, adding more summaries can never decrease the LFI posterior variance (see, e.g., [Frazier et al., 2018](#) for theoretical justification of this claim).

Consequently, according to posterior expected loss, S would asymptotically produce the smallest loss. The latter decision rule is unhelpful in practice as it completely disregards the fact that the computational resources required to approximate the posterior can increase drastically as the dimension of the summaries increase. While those resources are somewhat mitigated if one considers the [Fearnhead and Prangle \(2012\)](#) approach, it remains that the use of $R_{S_j}(\theta, \bar{\theta}_j)$ completely ignores the difference between posterior locations that arises when using different summaries, i.e., in general $\mathbb{E}_{\pi(\theta|S_1)}(\theta) \neq \mathbb{E}_{\pi(\theta|S_2)}(\theta) \neq \mathbb{E}_{\pi(\theta|S)}(\theta)$.

The asymptotic viewpoint also masks the critically important issue of the (Monte Carlo) accuracy of the resulting posterior approximation for a fixed computational budget. That is, while we can never decrease the asymptotic variance of the posterior by adding summaries, given a finite-time computational budget, the variability of the posterior approximation is an increasing function of the dimension of the summaries (see, e.g., [Blum, 2010](#)). Hence, given a finite computational budget, a posterior approximation based on S , denoted by $\tilde{\pi}(\theta|S)$, can easily have larger amounts of variability than a posterior approximation that targets a lower-dimensional set of summaries, e.g., $\tilde{\pi}(\theta|S_1)$. Hence, in practice large collections of summary statistics are not generally helpful for LFI without the application of adjustment procedures.

A.2 Cell Biology Example

Here we consider the lattice-free collective cell spreading model of [Browning et al. \(2018\)](#). The model permits cells to move freely in continuous space. There are three parameters in the model. There are two parameters that impact the spatial distribution of the cells, m and γ_b . The parameter p affects the number of cells. For specific details on the stochastic model, see [Browning et al. \(2018\)](#).

In the experiments of [Browning et al. \(2018\)](#), images of the cell population are taken every 12 hours with the final image taken at 36 hours. [Browning et al. \(2018\)](#) use the number of cells and the pair correlation computed from each of the three images as the summary statistics, resulting in a six dimensional summary statistic, S . The pair correlation is the ratio of the number of pairs of agents separated by some pre-specified distance to an expected number of cells separated by the same distance if the cells were uniformly distributed in space. In an attempt to learn more about m and γ_b , [Priddle et al. \(2022\)](#) consider a higher dimensional set of statistics summarising the spatial information. They consider Ripley’s K and J functions evaluated at various diameters for each time point. Combined with the same total number of cells at each time point, there are 21 summary statistics in total; see [Priddle et al. \(2022\)](#) for more details. We refer to the two sets of summary statistics as “pair” ([Browning et al., 2018](#)) and “spatial” [Priddle et al. \(2022\)](#), respectively.

We consider two likelihood-free algorithms. One of them uses the SMC ABC replenishment algorithm of [Drovandi and Pettitt \(2011a\)](#) where the algorithm is stopped when the

acceptance rate of the MCMC move step falls below 1%. We also consider the MCMC BSL algorithm of [Price et al. \(2018\)](#). For BSL, we use 10000 iterations MCMC with a random walk covariance matrix tuned using some pilot runs based on a simulated dataset. Our results below are based on 50 independent datasets simulated using $m = 1$, $p = 0.04$ and $\gamma_b = 0.5$. The prior distribution is set as $p \sim \mathcal{U}(0, 10)$, $m \sim \mathcal{U}(0, 0.2)$ and $\gamma_b \sim \mathcal{U}(0, 20)$ with no dependence amongst parameters.

Firstly we consider pooling the results from ABC with the pair correlation statistics (ABC pair) and ABC with the spatial statistics (ABC spatial). We might suspect that the spatial statistics will carry more information about m and γ_b than the pair statistics, but they have a higher dimension and we may be concerned that ABC spatial may produce inferences that are detrimental to p . The results are shown in [Table 5](#). It is evident that the pooled results improve on the inferences for m and γ_b compared to ABC pair (due to the good performance of ABC spatial for these two parameters) and improve on the inferences for p compared to ABC spatial (due to the good performance of ABC pair for this parameter). Since the parameter estimates are on different scales, we also consider pooling separately for each individual parameter. We can see that pooling with the first set of weights produces low relative MSEs for all three parameters compared to the other approaches.

Secondly we consider pooling the results from ABC spatial and BSL spatial. BSL avoids the tolerance error associated with ABC, but we might be concerned about its Gaussian likelihood assumption. It turns out that BSL is very effective in this particular problem,

since Table 6 shows that it produces the smallest MSE for all parameters. However, it can be seen that the pooled results produces small relative MSEs compared to ABC spatial. Thus, there is only a small loss of inefficiency compared to BSL spatial, whilst providing some robustness to the Gaussian assumption by pooling with the ABC results.

(A)	S_1			S_2			N/A		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
m	0.0018	0.58	0.15	0.16	0.41	0.12			
p	-1.1e-4	0.0017	2.9e-6	3.2e-4	0.0030	4.1e-6			
γ_b	2.8	4.09	10	0.35	1.45	1.1			
(B)	$\hat{\omega}$			$\tilde{\omega}$			N/A		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
m	0.17	0.47	0.12	0.09	0.51	0.10			
p	2.5e-4	0.0028	3.8e-6	1.2e-4	0.0026	3.4e-6			
γ_b	0.56	2.1	1.3	1.4	3.1	3.1			
(C)	$\hat{\omega}$			$\tilde{\omega}$			N/A		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
m	0.056	0.49	0.083	0.039	0.51	0.10			
p	-1.6e-5	0.0020	3.0e-6	-2.2e-5	0.0020	3.0e-6			
γ_b	0.54	2.1	1.3	1.4	3.1	3.0			

Table 5: Pooled posterior accuracy results in the cell biology model under the base set of summaries S_1 (pair) and the alternative set S_2 (spatial). The remaining information is as in Table 1. The average value of $\hat{\omega}$ and $\tilde{\omega}$ over the 50 datasets is 0.13 and 0.37, respectively, indicating preference for the inference based on the spatial summaries. (C) shows the same results as (B) except that pooling is done for each individual parameter to help remove the effect of scaling between different parameters.

(A)	ABC S_2			BSL S_2			N/A		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
m	0.16	0.41	0.12	0.10	0.21	0.07			
p	3.2e-4	0.0030	4.1e-6	5.2e-6	0.0014	2.8e-6			
γ_b	0.35	1.4	1.1	-1.2e-4	0.59	0.23			
(B)	$\hat{\omega}$			$\tilde{\omega}$			N/A		
	Bias	Std	MSE	Bias	Std	MSE	Bias	Std	MSE
m	0.12	0.27	0.07	0.12	0.29	0.08			
p	4.6e-5	0.0017	2.8e-6	1.0e-4	0.0020	3.1e-6			
γ_b	0.03	0.80	0.25	0.11	0.95	0.42			

Table 6: Pooled posterior accuracy results in the cell biology model under the base inference method (ABC spatial) and the alternative inference method (BSL spatial). The remaining information is as in Table 1. The average value of $\hat{\omega}$ and $\tilde{\omega}$ over the 50 datasets is 0.16 and 0.26, respectively, indicating preference for the BSL spatial results.