Herivelto Borges\* Instituto de Ciências Matemáticas e de Computação Universidade de São Paulo São Carlos - SP hborges@icmc.usp.br Cirilo Gonçalves<sup>†</sup> Departamento de Formação Geral Centro Federal de Educação Tecnológica de Minas Gerais Araxá - MG cirilo@cefetmg.br

#### ABSTRACT

Let  $\mathbb{K}$  be an algebraically closed field of characteristic p > 0. A pressing problem in the theory of algebraic curves is the determination of the *p*-rank of a (nonsingular, projective, irreducible) curve  $\mathcal{X}$  over  $\mathbb{K}$ . This birational invariant affects arithmetic and geometric properties of  $\mathcal{X}$ , and its fundamental role in the study of the automorphism group  $\operatorname{Aut}(\mathcal{X})$  has been noted by many authors in the past few decades. In this paper, we provide an extensive study of the *p*-rank of curves of Fermat type  $y^m = x^n + 1$  over  $\mathbb{K} = \overline{\mathbb{F}}_p$ . We determine a combinatorial formula for this invariant in the general case and show how this leads to explicit formulas of the *p*-rank of several such curves. By way of illustration, we present explicit formulas for more than twenty subfamilies of such curves, where *m* and *n* are generally given in terms of *p*. We also show how the approach can be used to compute the *p*-rank of other types of curves.

Keywords p-rank · Hasse-Witt invariant · automorphism group · supersingular curves

### **1** Introduction

Let  $\mathcal{X}$  be a nonsingular, projective, irreducible curve of genus  $\mathfrak{g} > 0$  over an algebraically closed field  $\mathbb{K}$  of characteristic p > 0, and let J[p] be the kernel of the multiplication-by-p morphism on the jacobian J of  $\mathcal{X}$ . The integer  $\gamma(\mathcal{X}) \in \{0, \dots, \mathfrak{g}\}$  for which  $J[p](\mathbb{K}) \cong (\mathbb{Z}/p\mathbb{Z})^{\gamma(\mathcal{X})}$  is a birational invariant, called the p-rank of  $\mathcal{X}$ . This number naturally connects with other birational invariants, and its study is fundamental to addressing several problems related to the classification of curves over finite fields.

The fact that the *p*-rank can affect the study of the automorphism group  $\operatorname{Aut}(\mathcal{X})$  of a curve  $\mathcal{X}$  of genus  $\mathfrak{g} \geq 2$  can be easily illustrated by Nakajima's results in [27, Theorems 1, 2, and 3]. There is also a well-known connection between zero *p*-rank and large automorphism groups. Indeed, for p > 2 and  $G \leq \operatorname{Aut}(\mathcal{X})$ , with G solvable or  $\mathfrak{g}$  even, Giulietti and Korchmáros proved that there exists a constant  $c_p > 0$  such that if  $|G| > c_p \mathfrak{g}^2$ , then the *p*-rank of  $\mathcal{X}$  is zero [9]. On the other hand, a conjecture by Guralnick and Zieve, reported in [13], [19], and [21], states that if  $\mathcal{X}$  is ordinary, that is,  $\gamma(\mathcal{X}) = \mathfrak{g}$ , then  $|G| \leq c_p \mathfrak{g}^{8/5}$  for some constant  $c_p > 0$ . With the additional hypothesis that G is solvable, Korchmáros and Montanucci proved that  $|G| \leq 34(\mathfrak{g}+1)^{3/2}$  [21].

Information on the *p*-rank  $\gamma(\mathcal{X})$  contributes to the study of other important notions, such as supersingularity, zeta function, and the *a*-number  $a(\mathcal{X}) \in \{0, \dots, \mathfrak{g} - \gamma(\mathcal{X})\}$ , which is another birational invariant [1], [7],[25], [29].

The characterization of the *p*-rank for a few curves can be found in the literature, of which the case of elliptic curves is well known [32]. It is also known that the *p*-rank of a curve  $\mathcal{X}$  is equal to its Hasse-Witt invariant, and then  $\gamma(\mathcal{X})$  can be investigated through the Hasse-Witt matrix of  $\mathcal{X}$ . In 1972, Miller studied a family of hyperelliptic curves for which this matrix is invertible [24]. When  $\mathcal{X}$  is a Fermat or hyperelliptic curve, Yui provided conditions for which  $\gamma(\mathcal{X}) = 0$  or  $\gamma(\mathcal{X}) = \mathfrak{g}$  [38, 39], and Kodama and Washio investigated relationships between the ranks of certain Hasse-Witt matrices [17, 18]. Some connections between Hasse-Witt matrices and Weierstrass semigroups were explored by

<sup>\*</sup>Avenida Trabalhador São-carlense, 400, São Carlos, CEP 13566-590, SP, Brazil

<sup>&</sup>lt;sup>†</sup>Avenida Ministro Olavo Drummond, 25, Araxá, CEP 38180-510, MG, Brazil

Stöhr and Viana [33]. The *p*-rank of the Fermat curves of prime degree  $l \neq p$  was studied by González in [12], where explicit formulas were obtained for some particular cases. The problem of the existence of hyperelliptic curves with prescribed genus and *p*-rank was solved by Zhu for p = 2 [40] and by Glass and Pries for p > 2 [11]. In 2010, Bassa and Beelen obtained the *p*-rank of the Fermat curve  $\mathcal{F}_{q-1} : y^{q-1} = x^{q-1} + 1$ , where *q* is a power of *p* [2, Theorem 19].

Though restricted to particular families of curves, the aforementioned results are of absolute interest. For instance, using these curves along with the Deuring-Shafarevich formula [34, Theorem 4.2], one can determine the *p*-rank of several other curves that arise as their Galois *p*-power covers. This approach was used by Giulietti, Korchmáros, and Timpanella to compute the *p*-rank of the Dickson-Guralnick-Zieve curve

$$\mathcal{C}:\frac{(x^{q^3}-x)(y^q-y)-(x^q-x)(y^{q^3}-y)}{(x^{q^2}-x)(y^q-y)-(x^q-x)(y^{q^2}-y)}=0$$

in terms of  $\gamma(\mathcal{F}_{q-1})$  [10, Remark 4]. They determined the p-rank of  $\mathcal{C}$  when q = p, concluding that  $\mathcal{C}$  is an ordinary curve with  $|\operatorname{Aut}(\mathcal{C})| \approx c\mathfrak{g}^{8/5}$ , that is,  $\mathcal{C}$  attains the conjectural Guralnick-Zieve bound. It is worth noting that the Deuring-Shafarevich formula is not the only method to obtain the *p*-rank of additional curves (see, e.g., Lemma 2.4, Theorem 3.5, and Corollary 3.6).

In [10, Remark 4], the authors mention that the *p*-rank of  $\mathcal{F}_{q-1} : y^{q-1} = x^{q-1} + 1$  is known only when q = p, although the *p*-rank of this specific Fermat curve was previously obtained in [2, Theorem 19]. These facts highlight not only the interest in the *p*-rank of Fermat-type curves but also the need for further research and discussion.

For integers  $n, m \ge 2$  coprime to p, this work determines the p-rank of the curves  $y^m = x^n + 1$  over  $\mathbb{K} = \overline{\mathbb{F}}_p$ . More precisely, it proves the following main results.

**Theorem 1.1.** Let  $n, m \ge 2$  be integers coprime to p, and let  $h, \alpha, \beta \in \mathbb{Z}_{>0}$  be such that  $\alpha n = \beta m = p^h - 1$ . Then the p-rank of the curve  $\mathcal{F}_{m,n} : y^m = x^n + 1$  is given by

$$\gamma(\mathcal{F}_{m,n}) = \#T - (m + n + \gcd(m, n)),$$

where

$$T = \left\{ (i,j) \in \mathbb{N}^2 \mid 0 \le i\alpha \le j\beta \le p^h - 1 \text{ and } \binom{j\beta}{i\alpha} \not\equiv 0 \pmod{p} \right\}$$

Note that the cardinality of T can be easily calculated for small values of n and m, and even when such values are larger, the task can be computationally simple. Beyond this, Theorem 1.1 effectively provides explicit formulas for the p-rank of curves of type  $\mathcal{F}_{m,n}$ . To illustrate its reach, we provide Tables 1—5 below, summarizing a few cases.

Table 1: The	<i>p</i> -rank of $y^{m_1}$	$=x^{n_1}+1.$	Throughout the table, $d$	= gcc	d(m)	, n)
--------------	-----------------------------	---------------	---------------------------	-------	------	------

$m_1$	$n_1$	$\gamma(\mathcal{F}_{m_1,n_1})$	Theorem
$\begin{array}{c} m, \\ m \mid p-1 \end{array}$	$\begin{array}{c}n,\\n\mid p+1\end{array}$	$2\sum_{i=\lceil m/2\rceil}^{m} \left\lfloor \frac{i(p-1)/m+1}{(p+1)/n} \right\rfloor - (m-\lceil m/2\rceil+2)(n-1) + 1 - d$	4.1
$\begin{array}{c} m, \\ m \mid p-1 \end{array}$	$p^n - 1$	$\sum_{i=1}^{m-1} (i(p-1)/m + 1)^n - 2(m-1)$	4.2
$p^2+p+1$	$p^2 + p + 1$	$p(p+1)(p^2+p+2)/8$	4.3
$p^2+p+1$	$p^{3} - 1$	$p(p+1)(p^3+2p^2+3p-14)/8$	4.3
$(p^n-1)/2$	$(p^n-1)/2$	$(p+1)^n(p^n+3)/2^{n+2} - 3(p^n-1)/2$	4.4
$(p^n-1)/2$	$p^n - 1$	$(p+1)^n (p^n+1)/2^{n+1} - 2(p^n-1)$	4.4
$p^m - 1$	$p^n - 1$	$\left(\sum_{i=0}^{p-1} \left((i+1)^{\frac{m}{d}} - i^{\frac{m}{d}}\right)(p-i)^{\frac{n}{d}}\right)^{d} - (p^{m} + p^{n} + p^{d} - 3)$	4.6
$p^m + 1, \\ p \neq 2$	$\begin{array}{c} p^n-1,\\ n/d \text{ even} \end{array}$	$\left(\sum_{i=0}^{p-1} \left((p-i)(i+1)\right)^{\frac{n}{2d}} + \sum_{1 \le j \le i \le p-1} \left((j+1)^{\frac{m}{d}} - 2j^{\frac{m}{d}} + (j-1)^{\frac{m}{d}}\right) \left((p-i)(i-j+1)\right)^{\frac{n}{2d}}\right)^d - (p^m + p^d)$	4.9
$p^m + 1, \\ p \neq 2$	$p^n - 1, n/d \text{ odd}$	$\frac{1}{2^n} \Big( (p+1)^{\frac{n}{d}} + \sum_{i=1}^{(p-1)/2} \left( (2i+1)^{\frac{m}{d}} - (2i-1)^{\frac{m}{d}} \right) (p-2i+1)^{\frac{n}{d}} \Big)^d - (p^m+1)$	4.9

$m_1$	$n_1$	$\gamma(\mathcal{F}_{m_1,n_1})$	
$m_1 \mid 2^m - 1, \\ p = 2$	$n_1 \mid 2^n - 1, \\ \gcd(n, m) = 1$	0	Corollary 4.7
$m_1 \mid p^m + 1$	$n_1 \mid p^n + 1$	0	Theorem 4.8
$m_1 \mid 2^m + 1, \\ p = 2$	$\begin{array}{c} n_1 \mid 2^n-1, \\ n/\gcd(m,n) \text{ odd} \end{array}$	0	Corollary 4.10

Table 2: The *p*-rank of  $y^{m_1} = x^{n_1} + 1$ 

Table 3: The *p*-rank of  $y^2 = x^n + 1$  – Corollary 5.2.

n	$\gamma(\mathcal{F}_{2,n})$
$2(p^r-1)$	$\left((p+1)/2\right)^r-2$
$2(p^{r}+1)$	$\left((p+1)/2\right)^r$
$p^{2r} + p^r + 1$	$\left((p+3)(p+1)/8\right)^r$
$2(p^{2r} + p^r + 1)$	$2\Big((p+3)(p+1)/8\Big)^r$
$p^{3r} - p^{2r} + p^r - 1$	$\left((p^2 + 2p + 3)(p+1)/12\right)^r - 2$

Despite the vanishing of their *p*-ranks, note that most curves in Table 2 are not supersingular, see Remark 2.8. To complement Table 3, let us mention that in general the *p*-rank of  $\mathcal{F}_{2,n}$ :  $y^2 = x^n + 1$  is given by

$$\gamma(\mathcal{F}_{2,n}) = \frac{n}{p^h - 1} \Big( ((p+1)/2)^h - \delta_n \Big), \tag{1.1}$$

for a suitable integer  $\delta_n \leq ((p+1)/2)^h$  dependent on  $\alpha = (p^h - 1)/n \in \mathbb{Z}_{>0}$  (Theorem 5.4). Table 4 presents some values of  $\delta_n$ .

Table 4: $\delta_i$	<sub>n</sub> for $y^2$	$= x^n$	+1 -	Theorem	5.4.
---------------------	------------------------	---------	------	---------	------

		$\delta_n$
hypotheses	n  odd	n even
$\alpha \mid \frac{p-1}{2}$	not applicable	$\alpha + 1$
$\alpha \mid \frac{p+1}{2}$	$\alpha$	2lpha
$\begin{array}{c} \alpha \mid \frac{p+3}{2}, \\ (p-1) \nmid n \end{array}$	$\alpha + (-1)^h$	$2\alpha + (-1)^h$
$\begin{array}{c} \alpha \mid \frac{p+3}{2}, \\ (p-1) \mid n \end{array}$	$\alpha + (-1)^h (1 - \alpha)$	$2\alpha + (-1)^h (1-\alpha)$
$\alpha \mid p-1$	$(\alpha + 2)/2$	

The curve  $\mathcal{F}_{2,n}: y^2 = x^{r\frac{p^h-1}{p-1}} + 1$ , where r is an even divisor of p-1, typifies the first line of Table 4. In this case, we have  $\delta_n = \alpha + 1 = (p-1)/r + 1$ , and then (1.1) gives

$$\gamma(\mathcal{F}_{2,n}) = \frac{r}{p-1} \left( ((p+1)/2)^h - 1 \right) - \frac{r}{p-1}.$$

In particular, the curve  $y^2 = x^{p^h-1} + 1$  has *p*-rank  $((p+1)/2)^r - 2$ , a fact recently proved by Pries and Ulmer in [30, Proposition 10.3]. Note that this also follows from the second line of Table 1.

As a by-product of the previous results, one can obtain the *p*-rank of other types of curves. This can be illustrated by the following curves

$$\mathcal{D}_n: y^2 = x(x^n + 1).$$

If  $n = (p^h - 1)/\alpha \in \mathbb{Z}_{>0}$  is odd, then  $\gamma(\mathcal{D}_n) = \gamma(\mathcal{C}_n)$  (Corollary 3.6). Otherwise, from Kani and Rosen's result in Theorem 2.3, it follows that  $\gamma(\mathcal{D}_n) = 0$  when  $\alpha$  is odd, and

$$\gamma(\mathcal{D}_n) = \gamma(\mathcal{F}_{2,2n}) - \gamma(\mathcal{F}_{2,n}) = \frac{n}{p^h - 1} \Big( ((p+1)/2)^h - \tilde{\delta}_n \Big),$$

with  $\tilde{\delta}_n = 2\delta_{2n} - \delta_n$ , when  $\alpha$  is even (Theorem 6.3). Some values of  $\tilde{\delta}_n$  are displayed in Table 5.

Table 5:	$\tilde{\delta}_n$ for	$y^2 = x$	$x(x^n +$	1) -	Theorem	6.3.
	10		· ·			

hypotheses	$\tilde{\delta}_n$
$\alpha \mid \frac{p-1}{2}$	1
$\alpha \mid \frac{p+1}{2}$	0
$\begin{array}{c} \alpha \mid \frac{p+3}{2}, \text{ and } \\ (p-1) \nmid 2n \text{ or } \\ (p-1) \mid n \end{array}$	$(-1)^{h}$
$\begin{array}{c c} \alpha \mid \frac{p+3}{2}, \\ (p-1) \mid 2n \text{ and } \\ (p-1) \nmid n \end{array}$	$(-1)^h(1-\alpha)$

This paper is organized as follows. In Section 2, we establish notation and collect some preliminary results. In Section 3, we prove the main result. In Sections 4 and 5, we use the results of Section 3 to present closed formulas for the p-rank of several Fermat-type curves. Finally in Section 6, we illustrate how the results of the previous sections can be used to obtain the p-rank of other types of curves.

### Notation

The following notation is used throughout this text.

- $\mathbb{F}_p$  is the prime field of characteristic p > 0
- $\mathbb{K} = \overline{\mathbb{F}}_p$  is the algebraic closure of  $\mathbb{F}_p$
- $\mathcal{F}_{m,n}$  is the curve  $y^m = x^n + 1$  over  $\mathbb{K}$ , where m and n are not divisible by p
- $C_n$  is the curve  $\mathcal{F}_{2,n}$
- $\mathcal{D}_n$  is the curve  $y^2 = x(x^n + 1)$  over  $\mathbb{K}$ , where p > 2 does not divide n.
- $\gamma(\mathcal{C})$  is the *p*-rank of the curve  $\mathcal{C}$
- $\mathbb{N} = \{0, 1, 2, \ldots\}$  is the set of natural numbers
- $\mathbb{Z}_{>0}$  is the set of positive integers
- $\lfloor x \rfloor = \max\{m \in \mathbb{Z} \mid m \le x\}$  and  $\lceil x \rceil = \min\{m \in \mathbb{Z} \mid m \ge x\}$ , where x is a real number
- [r, s], where  $r \leq s$  are integers, is the set  $\{r, \ldots, s\}$
- $[\![r,s]\!]^h$ , with  $h \in \mathbb{Z}_{>0}$ , is the cartesian product  $[\![r,s]\!] \times \cdots \times [\![r,s]\!]$  (h times).

# 2 Preliminaries

Let  $\mathcal{X}$  be a projective, nonsingular, and irreducible algebraic curve of genus  $\mathfrak{g} > 0$  defined over  $\mathbb{K}$ . Let  $\mathbb{K}(\mathcal{X}) = \mathbb{K}(x, y)$  be the function field of  $\mathcal{X}$ , where  $x \in \mathbb{K}(\mathcal{X})$  is a separating variable. Let  $\Delta_{\mathcal{X}} = \{\varphi dx : \varphi \in \mathbb{K}(\mathcal{X})\}$  be the module of differentials of  $\mathbb{K}(\mathcal{X})$ . For any  $\omega \in \Delta_{\mathcal{X}}$ , we say that

•  $\omega$  is holomorphic if  $\operatorname{div}(\omega)$  is effective

- $\omega$  is exact if  $\omega = df$  for some  $f \in \mathbb{K}(\mathcal{X})$
- $\omega$  is logarithmic if  $\omega = df/f$  for some  $f \neq 0$  in  $\mathbb{K}(\mathcal{X})$ .

The Cartier operator  $C : \Delta_{\mathcal{X}} \to \Delta_{\mathcal{X}}$  is defined as follows. For each  $\omega \in \Delta_{\mathcal{X}}$ , let  $f_0, \ldots, f_{p-1} \in \mathbb{K}(\mathcal{X})$  be such that  $\omega = (f_0^p + f_1^p x + \cdots + f_{p-1}^p x^{p-1}) dx$ , and set

$$C(\omega) := f_{p-1}dx.$$

This operator has the following properties:

- (i) The value of  $C(\omega)$  does not depend on the choice of x
- (ii) C is 1/p-linear, that is, C is additive and  $C(f^p\omega) = fC(\omega)$  for all  $f \in \mathbb{K}(\mathcal{X})$
- (iii)  $C(f^{p-1}df) = df$  for all  $f \in \mathbb{K}(\mathcal{X})$
- (iv)  $C(\omega) = 0$  if and only if  $\omega$  is exact
- (v)  $C(\omega) = \omega$  if and only if  $\omega$  is logarithmic
- (vi) If  $\omega$  is holomorphic, then so is  $C(\omega)$ .

For  $n, i \in \mathbb{Z}_{>0}$ , using the base-p expansion of i and the previous properties, one can easily check that

$$C^{n}(x^{i}dx) = \begin{cases} 0 & \text{if } i \not\equiv -1 \pmod{p^{n}} \\ x^{s-1}dx & \text{if } i+1 = sp^{n}, \end{cases}$$
(2.1)

where  $C^n = C \circ \cdots \circ C$  (*n* times). For further details, see [8], [15], and [31, §10 and §11].

Let  $\Delta_{\mathcal{X}}^{(1)}$  be the g-dimensional K-vector space of the holomorphic differentials on  $\mathcal{X}$ , and let  $\Delta_{\mathcal{X}}^{\log}$  be the subspace of  $\Delta_{\mathcal{X}}^{(1)}$  spanned by the logarithmic differentials.

**Definition 2.1.** The dimension  $\gamma(\mathcal{X})$  of  $\Delta_{\mathcal{X}}^{\log}$  is the *p*-rank of  $\mathcal{X}$ .

Let  $\Delta^0_{\mathcal{X}}$  be the subspace of the differentials  $\omega \in \Delta^{(1)}_{\mathcal{X}}$  for which  $C^n(\omega) = 0$  for some integer n > 0. Since the Cartier operator is 1/p-linear, the following canonical decomposition holds [14].

$$\Delta_{\mathcal{X}}^{(1)} = \Delta_{\mathcal{X}}^{\log} \oplus \Delta_{\mathcal{X}}^{0}.$$
(2.2)

**Theorem 2.2.** Let  $\omega \in \Delta_{\mathcal{X}}^{(1)}$  be a nonzero holomorphic differential. If  $h \in \mathbb{Z}_{>0}$  is such that  $C^h(\omega) = \alpha \omega$  for some  $\alpha \in \mathbb{K}$ , then  $\omega \in \Delta_{\mathcal{X}}^{\log}$  if and only if  $\alpha \neq 0$ . In particular, for any basis  $B = \{\omega_1, \ldots, \omega_g\}$  of  $\Delta_{\mathcal{X}}^{(1)}$ , where  $C^h(\omega_i) = \alpha_i \omega_i, \alpha_i \in \mathbb{K}$ , we have

$$\gamma(\mathcal{X}) = \#\{\omega_i \in B \mid i \in [[1, \mathfrak{g}]] \text{ and } \alpha_i \neq 0\}.$$

*Proof.* Suppose  $\alpha \neq 0$ . The decomposition (2.2) gives  $\omega = u + v$ , with  $u \in \Delta^{\log}_{\mathcal{X}}$  and  $v \in \Delta^{0}_{\mathcal{X}}$ . Thus  $C^{ht}(v) = 0$  for some integer t > 0, and  $C(\Delta^{\log}_{\mathcal{X}}) \subseteq \Delta^{\log}_{\mathcal{X}}$  implies  $C^{ht}(u) \in \Delta^{\log}_{\mathcal{X}}$ . Since  $C^{h}(\omega) = \alpha \omega$ , with  $\alpha \neq 0$ , the 1/p-linearity of C yields  $C^{ht}(w) = \beta w = \beta(u + v)$  for some  $\beta \neq 0$ . Hence

$$\beta v = C^{ht}(w) - \beta u = C^{ht}(u+v) - \beta u = C^{ht}(u) - \beta u \in \Delta_{\mathcal{X}}^{\log}$$

gives v = 0, and then  $\omega = u \in \Delta_{\mathcal{X}}^{\log}$ . The converse is clear. For the final assertion, note that the basis B for  $\Delta_{\mathcal{X}}^{(1)} = \Delta_{\mathcal{X}}^{\log} \oplus \Delta_{\mathcal{X}}^{0}$  is such that

$$B_1 = \{\omega_i \in B \mid i \in [1, \mathfrak{g}] \text{ and } \alpha_i \neq 0\} \subseteq \Delta_{\mathcal{X}}^{\log}$$

and  $B \setminus B_1 \subseteq \Delta^0_{\mathcal{X}}$ . Therefore,  $B_1$  is a basis of  $\Delta^{\log}_{\mathcal{X}}$ , and then  $\gamma(\mathcal{X}) = \#B_1$ .

Let us recall that the *p*-rank  $\gamma(J)$  of a jacobian *J*, more generally, of an abelian variety, is invariant under isogeny, and  $\gamma(J_1 \times J_2) = \gamma(J_1) + \gamma(J_2)$  for jacobians  $J_1$ ,  $J_2$  defined over  $\mathbb{K}$  [5, 26]. In particular, the following result is suitable to investigate the *p*-rank of a curve  $\mathcal{X}$ .

**Theorem 2.3** ([16], Theorem B). Let  $G \leq Aut(\mathcal{X})$  be a (finite) subgroup such that  $G = H_1 \cup ... \cup H_t$ , where the subgroups  $H_i \leq G$  satisfy  $H_i \cap H_j = \{1\}$  for  $i \neq j$ . Then we have the isogeny relation

$$J_{\mathcal{X}}^{t-1} \times J_{\mathcal{X}/G}^g \sim J_{\mathcal{X}/H_1}^{h_1} \times \cdots \times J_{\mathcal{X}/H_t}^{h_t},$$

where g = |G|,  $h_i = |H_i|$ , and  $J^n = J \times \cdots \times J$  (n times).

The following relates the *p*-rank of a curve to the *p*-rank of any of its (sub)covers.

**Lemma 2.4** ([2], Lemma 6). Let C and D be curves of genus  $\mathfrak{g}(C)$  and  $\mathfrak{g}(D)$ , respectively, defined over a finite field. *If*  $f : C \longrightarrow D$  *is a nonconstant morphism, then* 

- (i)  $\gamma(\mathcal{D}) \leq \gamma(\mathcal{C}) \leq \gamma(\mathcal{D}) + \mathfrak{g}(\mathcal{C}) \mathfrak{g}(\mathcal{D})$
- (ii) if  $\gamma(\mathcal{C}) = 0$ , then  $\gamma(\mathcal{D}) = 0$
- (iii) if C is ordinary, then so is D.

The following well-known facts will be useful in obtaining formulas for the *p*-rank of curves of type  $y^m = x^n + 1$ . Lemma 2.5 ([6, 22], Lucas' Lemma). Let *p* be a prime, and let  $a_0, b_0, \ldots, a_r, b_r \in [0, p-1]$ . Then

$$\begin{pmatrix} a_r p^r + \dots + a_1 p + a_0 \\ b_r p^r + \dots + b_1 p + b_0 \end{pmatrix} \equiv \prod_{i=0}^r \begin{pmatrix} a_i \\ b_i \end{pmatrix} \pmod{p}$$

**Lemma 2.6** ([4], Lemma 2.6; [23], p. 175). *Let* p *be a prime, and*  $m, n \in \mathbb{Z}_{>0}$ . *If* d = gcd(m, n), *then* 

$$\gcd(p^m+1,p^n-1) = \begin{cases} 1 & \text{if } n/d \text{ is odd and } p = 2\\ 2 & \text{if } n/d \text{ is odd and } p > 2\\ p^d+1 & \text{if } n/d \text{ is even }. \end{cases}$$

A curve  $\mathcal{X}$  defined over a finite field k is called supersingular if it is Hasse-Weil minimal over some extension of k, and it is well known that supersingular curves have zero p-rank [29, 32]. Though not affecting our results, it is worth noting that [28, Theorem 1.2], with some simple adjustments, yields the following (see also [36, Theorem 5]).

**Theorem 2.7.** Let  $n, m \in \mathbb{Z}_{>0}$  be such that  $\mathcal{F}_{m,n} : y^m = x^n + 1$  is an irreducible curve over  $\overline{\mathbb{F}}_p$  of genus  $\mathfrak{g} > 0$ . Then  $\mathcal{F}_{m,n}$  is supersingular if and only if there exists  $h \in \mathbb{Z}_{>0}$  such that n and m divide  $p^h + 1$ .

*Remark* 2.8. It follows from Lemma 2.6 and Theorem 2.7 that the curves  $y^{p^m+1} = x^{p^n+1} + 1$  are supersingular if and only if  $\nu_2(n) = \nu_2(m)$ , where  $\nu_2 : \mathbb{Z} \to \mathbb{N} \cup \{\infty\}$  is the 2-adic valuation. In the same vein, one sees that the only nonrational curve of type  $y^{p^m \pm 1} = x^{p^n-1} + 1$  that is supersingular is  $y^{3^m+1} = x^2 + 1$ .

### **3** The p-rank of Fermat-type curves

Throughout this section  $n, m \ge 2$  are integers coprime to  $p, \mathcal{F}_{m,n} : y^m = x^n + 1$  is a curve over  $\mathbb{K}$ , and  $q = p^h$  is such that n and m are divisors of q - 1. Additionally, let  $\mathbb{K}(\mathcal{F}_{m,n}) = \mathbb{K}(x, y)$  be the function field of  $\mathcal{F}_{m,n}$ . The genus of  $\mathcal{F}_{m,n}$  is given by

$$\mathfrak{g} = \frac{(m-1)(n-1) + 1 - \gcd(m,n)}{2}$$

and the set

$$B = \left\{ \omega_{i,j} = \frac{x^{i-1}}{y^j} dx \mid (i,j) \in \mathbb{N}^2 \text{ with } m \le im < jn \le n(m-1) \right\}$$

is a basis for the vector space of holomorphic differentials of  $\mathcal{F}_{m,n}$ . For further details on the basic properties of the curve  $\mathcal{F}_{m,n}$ , see [20] and [37].

Note that for  $(i, j) \in \mathbb{N}^2$  and  $m \leq im < jn \leq n(m-1)$ , the equation  $y^m = x^n + 1$  yields

$$\frac{x^{i-1}}{y^j} = \frac{x^{i-1}}{y^{jq}} (x^n + 1)^{j(q-1)/m} = \sum_{r=0}^{\frac{j(q-1)}{m}} {\binom{j(q-1)}{m} \choose r} \frac{x^{rn+i-1}}{y^{jq}}.$$

That is, the elements of B can be written as

$$\omega_{i,j} = \sum_{r=0}^{\frac{j(q-1)}{m}} {\binom{j(q-1)}{m} r} \frac{x^{rn+i-1}}{y^{jq}} dx.$$
(3.1)

We now restate and prove our main result for the *p*-rank of Fermat-type curves.

**Theorem 3.1.** Let  $n, m \ge 2$  be integers coprime to p, and let  $h, \alpha, \beta \in \mathbb{Z}_{>0}$  be such that  $\alpha n = \beta m = p^h - 1$ . Then the p-rank of the curve  $\mathcal{F}_{m,n} : y^m = x^n + 1$  is given by

$$\gamma(\mathcal{F}_{m,n}) = \#T - (m+n+\gcd(m,n))$$

where

$$T = \left\{ (i,j) \in \mathbb{N}^2 \mid 0 \le i\alpha \le j\beta \le p^h - 1 \text{ and } \binom{j\beta}{i\alpha} \not\equiv 0 \pmod{p} \right\}$$

*Proof.* Let  $q = p^h$ . The 1/p-linearity of the Cartier operator and (3.1) imply

$$C^{h}(\omega_{i,j}) = \sum_{r=0}^{\frac{j(q-1)}{m}} {\binom{j(q-1)}{m} \choose r} \frac{C^{h}(x^{rn+i-1}dx)}{y^{j}}.$$
(3.2)

It follows from (2.1) that  $C^h(x^{rn+i-1}dx)$  is nonzero if and only if  $nr + i \equiv 0 \pmod{q}$ . Note that  $r \leq j(q-1)/m$  and  $m \leq im < jn \leq n(m-1)$  entail  $1 \leq nr + i \leq nq - 1$ . Thus  $nr + i \equiv 0 \pmod{q}$  if and only if  $nr + i \in \{q, 2q, \dots, (n-1)q\}$ . The latter is equivalent to

$$r \in \left\{\frac{q-i}{n}, \frac{2q-i}{n}, \dots, \frac{(n-1)q-i}{n}\right\} \cap \mathbb{Z}$$

and thus  $C^h(x^{rn+i-1}dx)$  is nonzero if and only if  $r = \frac{i(q-1)}{n}$ . Therefore, equation (3.2) reduces to

$$C^{h}(\omega_{i,j}) = \left(\frac{\frac{j(q-1)}{m}}{\frac{i(q-1)}{n}}\right)\omega_{i,j},$$

and Theorem 2.2 yields

$$\gamma(\mathcal{F}_{m,n}) = \#\left\{ (i,j) \in \mathbb{N}^2 \mid m \le im < jn \le n(m-1) \text{ and } \binom{j\beta}{i\alpha} \ne 0 \pmod{p} \right\}.$$
(3.3)

Note that the pairs  $(i, j) \in T$  not occurring in (3.3) are those for which j = m, i = 0, or im = jn. One can check that the number of such pairs is m + n + gcd(n, m), and this concludes the proof.

*Remark* 3.2. In Theorem 3.1, note that one can always define  $h = \operatorname{lcm}(h_1, h_2)$ , where  $h_1$  and  $h_2$  are the orders of p in  $(\mathbb{Z}/m\mathbb{Z})^{\times}$  and  $(\mathbb{Z}/n\mathbb{Z})^{\times}$ , respectively. Then  $\alpha = (p^h - 1)/n$  and  $\beta = (p^h - 1)/m$ .

The subsequent particular cases of Theorem 3.1 are worth highlighting.

**Corollary 3.3.** Let  $n \in \mathbb{Z}_{>0}$ , with gcd(n, p) = 1, and let  $q = p^h$  be such that  $\alpha = (q-1)/n \in \mathbb{Z}_{>0}$ . Then the *p*-rank of the Fermat curve  $\mathcal{F}_n : y^n = x^n + 1$  is given by

$$\gamma(\mathcal{F}_n) = \#T - 3n,$$

where

$$T = \left\{ (i,j) \in \mathbb{N}^2 \mid 0 \le i \le j \le n \text{ and } \binom{j\alpha}{i\alpha} \not\equiv 0 \pmod{p} \right\}.$$

**Corollary 3.4.** Let  $n \in \mathbb{Z}_{>0}$ , with gcd(n, p) = 1, and let  $q = p^h$  be such that  $\alpha = (q-1)/n \in \mathbb{Z}_{>0}$ . Then the *p*-rank of  $C_n : y^2 = x^n + 1$  is given by

$$\gamma(\mathcal{C}_n) = \begin{cases} \#S - 1 & \text{if } n \text{ is odd} \\ \#S - 2 & \text{if } n \text{ is even }, \end{cases}$$

where

$$S = \left\{ i \in \left[\!\left[0, \lfloor n/2 \rfloor\right]\!\right] \mid \begin{pmatrix} \frac{q-1}{2} \\ i\alpha \end{pmatrix} \not\equiv 0 \pmod{p} \right\}.$$

Proof. From Theorem 3.1, the problem reduces to determining the cardinality of

$$T = \left\{ (i,j) \in \mathbb{N}^2 \mid 0 \le i\alpha \le \frac{j(q-1)}{2} \le q-1 \text{ and } \begin{pmatrix} \frac{j(q-1)}{2} \\ i\alpha \end{pmatrix} \ne 0 \pmod{p} \right\}.$$

Note that  $(i, j) \in T$  implies  $j \in \{0, 1, 2\}$ , and  $\#\{(i, j) \in T : j \in \{0, 2\}\} = n + 2$ . In addition, since  $(q - 1)/(2\alpha) = n/2$ , we have that  $(i, 1) \in T$  is equivalent to  $i \in [0, \lfloor n/2 \rfloor]$  and  $\binom{q-1}{2}{i\alpha} \not\equiv 0 \pmod{p}$ . Therefore, #T = #S + n + 2, and then

$$\gamma(\mathcal{C}_n) = \#T - (n+2+\gcd(2,n)) = \begin{cases} \#S-1 & \text{if } n \text{ is odd} \\ \#S-2 & \text{if } n \text{ is even} \end{cases}.$$

In what follows, we use Theorem 2.3 to relate the *p*-ranks of the curves  $\mathcal{F}_{m,n}$  and  $y^m = x^r(x^{m^{k-1}u} + 1)$ , with  $r \in [0, m-1]$ .

**Theorem 3.5.** Let  $m \neq p$  be a prime, and let  $u \geq 1$  be an integer coprime with p and m. Then for each integer  $k \geq 1$  and  $r \in [0, m-1]$ , the curves

$$\mathcal{F}: y^m = x^{m^k u} + 1 \text{ and } \mathcal{G}_r: y^m = x^r (x^{m^{k-1} u} + 1)$$

are such that

$$\gamma(\mathcal{F}) = \sum_{r=0}^{m-1} \gamma(\mathcal{G}_r)$$

*Proof.* Let  $\zeta$  be a primitive *m*-th root of unity. For  $k \ge 1$ , note that

$$\pi: (x, y) \mapsto (\zeta x, y) \text{ and } \sigma: (x, y) \mapsto (x, \zeta^{m-1}y)$$

are automorphisms of  $\mathcal{F}$  and that  $G = \langle \pi, \sigma \rangle \cong \mathbb{Z}_m \times \mathbb{Z}_m$  is a subgroup of  $\operatorname{Aut}(\mathcal{F})$ , with

$$G = \langle \sigma \rangle \cup \langle \pi \rangle \cup \langle \pi \circ \sigma \rangle \cup \cdots \cup \langle \pi \circ \sigma^{m-1} \rangle.$$

Clearly, each pairwise intersection of these m+1 subgroups is trivial. Moreover, one can readily check that the quotient curves  $\mathcal{F}/G$  and  $\mathcal{F}/\langle\sigma\rangle$  have genus zero. For  $r \in [\![0, m-1]\!]$ , note that  $\varepsilon = x^m$  and  $\eta = x^r y$  are elements in  $\mathbb{K}(x, y)$  that are fixed by  $\pi \circ \sigma^r$ , that is,  $\mathbb{K}(\varepsilon, \eta) \subseteq \mathbb{K}(x, y)^{\langle \pi \circ \sigma^r \rangle}$ . Since  $\mathbb{K}(x, y) = \mathbb{K}(\varepsilon, \eta)(x)$  and as  $T^m - \varepsilon \in \mathbb{K}(\varepsilon, \eta)[T]$  vanishes at x, we have that

$$\mathbb{K}(x,y):\mathbb{K}(\varepsilon,\eta)] = [\mathbb{K}(x,y):\mathbb{K}(x,y)^{\langle \pi \circ \sigma' \rangle}] = m$$

and then  $\mathbb{K}(x,y)^{\langle \pi \circ \sigma^r \rangle} = \mathbb{K}(\varepsilon,\eta)$ . Therefore, the quotient curve  $\mathcal{F}/\langle \pi \circ \sigma^r \rangle$  is isomorphic to the curve given by  $\eta^m = \varepsilon^r (\varepsilon^{m^{k-1}u} + 1)$ . Now Theorem 2.3 yields

$$J^m_{\mathcal{F}} \sim J^m_{\mathcal{F}/\langle \pi \rangle} \times J^m_{\mathcal{F}/\langle \pi \circ \sigma \rangle} \times \cdots \times J^m_{\mathcal{F}/\langle \pi \circ \sigma^{m-1} \rangle},$$
  
 $\chi(\mathcal{G}_r).$ 

and then  $\gamma(\mathcal{F}) = \sum_{r=0}^{m-1} \gamma(\mathcal{G}_r).$ 

**Corollary 3.6.** Let  $u \ge 1$  be an odd integer coprime to p > 2. Then for each integer  $k \ge 0$ , the curves

$$\mathcal{F}_k: y^2 = x^{2^k u} + 1 \text{ and } \mathcal{H}_k: y^2 = x(x^{2^k u} + 1)$$

are such that

(i) 
$$\gamma(\mathcal{F}_0) = \gamma(\mathcal{H}_0)$$
  
(ii)  $\gamma(\mathcal{H}_{k-1}) = \gamma(\mathcal{F}_k) - \gamma(\mathcal{F}_{k-1}), \text{ for } k \ge 1$   
(iii)  $\gamma(\mathcal{F}_k) = \gamma(\mathcal{F}_0) + \sum_{i=0}^{k-1} \gamma(\mathcal{H}_i), \text{ for } k \ge 1$   
(iv)  $\gamma(\mathcal{F}_1) = 2\gamma(\mathcal{F}_0).$ 

*Proof.* Note that the map  $(x, y) \mapsto \left(\frac{1}{x}, \frac{y}{x^{(u+1)/2}}\right)$  from  $\mathcal{H}_0$  to  $\mathcal{F}_0$  gives  $\mathcal{H}_0 \cong \mathcal{F}_0$ , which proves the first assertion. By Theorem 3.5, we have

$$\gamma(\mathcal{F}_k) = \gamma(\mathcal{F}_{k-1}) + \gamma(\mathcal{H}_{k-1}), \tag{3.4}$$

which proves (ii). Assertion (iii) follows from the recursive relation in (3.4), and the last assertion follows from (i) and the case k = 1 in (ii).

## 4 Formulas for the p-rank of Fermat-type curves

We have seen in Theorem 3.1 that the *p*-rank of  $\mathcal{F}_{m,n}$  can be obtained from the cardinality of a set *T*, given in terms of certain binomial coefficients. In this section, we show how one can effectively turn this information into formulas for the *p*-rank of  $\mathcal{F}_{m,n}$ .

**Theorem 4.1.** If m|(p-1) and n|(p+1), then  $\mathcal{F}_{m,n}: y^m = x^n + 1$  has p-rank

$$\gamma(\mathcal{F}_{m,n}) = 2\sum_{j=\lceil m/2\rceil}^{m} \left\lfloor \frac{jm_0 + 1}{n_0} \right\rfloor - (m - \lceil m/2\rceil + 2)(n - 1) + 1 - \gcd(m, n),$$
(4.1)

where  $m_0 = (p-1)/m$  and  $n_0 = (p+1)/n$ . In particular,

$$\gamma(\mathcal{F}_{m,p+1}) = \begin{cases} m_0(m-1)^2/4 + (m-1)/2 & \text{if } m \text{ is odd} \\ m_0(m^2 - 2m)/4 + (m-2)/2 & \text{if } m \text{ is even} \end{cases}$$
(4.2)

and

$$\gamma(\mathcal{F}_{p-1,n}) = \begin{cases} n_0(n^2 - 1)/4 - n + 1 & \text{if } n \text{ is odd} \\ n_0 n^2/4 - n & \text{if } n \text{ is even.} \end{cases}$$
(4.3)

*Proof.* For  $\alpha = n_0(p-1), \beta = m_0(p+1)$ , and

$$\Gamma = \left\{ (i,j) \in \mathbb{N}^2 \mid 0 \le i\alpha \le j\beta \le p^2 - 1 \text{ and } \binom{j\beta}{i\alpha} \not\equiv 0 \pmod{p} \right\},$$

Theorem 3.1 gives

$$\gamma(\mathcal{F}_{m,n}) = \#T - (m+n + \gcd(m,n)). \tag{4.4}$$

Note that the values i = 0 and i = n give rise to a total of m + 2 pairs  $(i, j) \in T$ . For  $1 \le i \le n - 1$ , the base-*p* expansions of  $i\alpha$  and  $j\beta$  are given by

$$i\alpha = (in_0 - 1)p + p - in_0$$
 and  $j\beta = jm_0p + jm_0$ 

and Lucas' Lemma entails  $(i, j) \in T$  if and only if  $p - jm_0 \le in_0 \le jm_0 + 1$ , that is,

$$n - \left\lfloor \frac{jm_0 + 1}{n_0} \right\rfloor \le i \le \left\lfloor \frac{jm_0 + 1}{n_0} \right\rfloor.$$

Hence  $\#T = m + 2 + \sum_{j=\lceil m/2 \rceil}^{m} \left( 2 \left\lfloor \frac{jm_0+1}{n_0} \right\rfloor - n + 1 \right)$ , and then equation (4.4) concludes the proof of (4.1). The particular cases corresponding to the curves  $\mathcal{F}_{m,p+1} : y^m = x^{p+1} + 1$  and  $\mathcal{F}_{p-1,n} : y^{p-1} = x^n + 1$  follow directly from the computation of  $\sum_{j=\lceil m/2 \rceil}^{m} \left\lfloor \frac{jm_0+1}{n_0} \right\rfloor$ . Case (4.2) follows immediately from

$$\sum_{j=\lceil m/2 \rceil}^{m} (jm_0+1) = (m - \lceil m/2 \rceil + 1)(m_0m + m_0\lceil m/2 \rceil + 2)/2$$

Case (4.3) follows from the basic properties of the floor function, including the identity  $\sum_{i=0}^{rs-1} \left\lfloor \frac{i}{s} \right\rfloor = sr(r-1)/2$ , from which one obtains

$$\sum_{j=\lceil (p-1)/2\rceil}^{p-1} \left\lfloor \frac{j+1}{n_0} \right\rfloor = \begin{cases} n_0(3n^2 - 2n - 1)/8 & \text{if } n \text{ is odd} \\ n_0(3n^2 - 2n)/8 & \text{if } n \text{ is even.} \end{cases}$$

**Theorem 4.2.** Let  $m, h \in \mathbb{Z}_{>0}$  be such that m > 1 divides  $p^h - 1$ . The *p*-rank of  $\mathcal{F}_{m,p^h-1} : y^m = x^{p^h-1} + 1$  is

$$\gamma = \sum_{j=1}^{m-1} \left( \prod_{r=1}^{h} (a_{j,r} + 1) \right) - 2(m-1), \tag{4.5}$$

where  $a_{j,1}, \ldots, a_{j,h}$  are the coefficients in the base-*p* expansion of  $j(p^h - 1)/m$ . In particular, if *m* divides p - 1, then

$$\gamma = \sum_{j=1}^{m-1} \left( \frac{j(p-1)}{m} + 1 \right)^h - 2(m-1).$$
(4.6)

*Proof.* From Theorem 3.1, we have  $\beta = (p^h - 1)/m$  and  $\gamma = \#T - (2m + p^h - 1)$ , where

$$T = \left\{ (i,j) \in \mathbb{N}^2 \mid 0 \le i \le j\beta \le p^h - 1 \text{ and } \binom{j\beta}{i} \not\equiv 0 \pmod{p} \right\}.$$

For each  $j \in [0, m]$ , consider the base-*p* expansion of  $j\beta = a_{j,1}p^{h-1} + \cdots + a_{j,h}$ . By Lucas' Lemma, the integers  $i \in [0, j\beta]$  for which  $\binom{j\beta}{i} \not\equiv 0 \pmod{p}$  are given by  $i = b_1p^{h-1} + \cdots + b_{h-1}p + b_h$ , where  $b_r \in [0, a_{j,r}]$  and  $1 \leq r \leq h$ . Clearly, there exist  $\prod_{r=1}^{h} (a_{j,r} + 1)$  such integers. Therefore,

$$\gamma = \sum_{j=0}^{m} \left( \prod_{r=1}^{h} (a_{j,r}+1) \right) - (2m+p^{h}-1) = \sum_{j=1}^{m-1} \left( \prod_{r=1}^{h} (a_{j,r}+1) \right) - 2(m-1).$$

Note that if  $m \mid (p-1)$ , then  $a_{j,r} = j(p-1)/m$  for all  $r \in [[1, h]]$ . Thus (4.6) follows immediately from (4.5).

**Theorem 4.3.** The *p*-rank of the curves

$$y^{p^2+p+1} = x^{p^2+p+1} + 1$$
 and  $y^{p^2+p+1} = x^{p^3-1} + 1$ 

are given by

$$\gamma(\mathcal{F}_{p^2+p+1}) = p(p+1)(p^2+p+2)/8$$
 and  $\gamma(\mathcal{F}_{p^2+p+1,p^3-1}) = p(p+1)(p^3+2p^2+3p-14)/8$ ,

respectively.

*Proof.* It follows from (3.3) that  $\gamma(\mathcal{F}_{p^2+p+1}) = #A$ , where

$$A = \left\{ (i,j) \in \mathbb{N}^2 \mid 1 \le i < j \le p^2 + p \text{ and } \binom{j(p-1)}{i(p-1)} \not\equiv 0 \pmod{p} \right\}.$$

For each  $i, j \in [[1, p^2 + p]]$ , write  $j = a_1p + a_0$  and  $i = b_1p + b_0$ , with  $a_r, b_r \in [[0, p]]$  and  $a_0b_0 \neq 0$ , and consider the following base-p expansions

$$j(p-1) = \begin{cases} (a_1-1)p^2 + (p+a_0-a_1-1)p + (p-a_0) & \text{if } a_0 \le a_1\\ a_1p^2 + (a_0-a_1-1)p + (p-a_0) & \text{if } a_0 > a_1 \end{cases}$$
(4.7)

and

$$i(p-1) = \begin{cases} (b_1-1)p^2 + (p+b_0-b_1-1)p + (p-b_0) & \text{if } b_0 \le b_1 \\ b_1p^2 + (b_0-b_1-1)p + (p-b_0) & \text{if } b_0 > b_1. \end{cases}$$

By Lucas' Lemma, there is no pair  $(i, j) = (b_1p + b_0, a_1p + a_0) \in A$  for which  $a_0 > a_1$  or  $b_0 \le b_1$ . Indeed, direct inspection shows that  $\binom{j(p-1)}{i(p-1)} \not\equiv 0 \pmod{p}$  implies that i = j when either  $a_0 > a_1$  and  $b_0 > b_1$  or  $a_0 \le a_1$  and  $b_0 \le b_1$ , and that  $\binom{j(p-1)}{i(p-1)} \equiv 0 \pmod{p}$  whenever  $a_0 > a_1$  and  $b_0 \le b_1$ . Thus if  $(i, j) = (b_1p + b_0, a_1p + a_0) \in A$ , then  $1 \le a_0 \le a_1 \le p$  and  $0 \le b_1 < b_0 \le p$ , and Lucas' Lemma yields

$$A = \left\{ (b_1 p + b_0, a_1 p + a_0) \middle| \begin{array}{l} a_1, b_0 \in \llbracket 1, p \rrbracket, \\ 1 \le a_0, b_1 + 1 \le \min \{a_1, b_0\}, \\ a_1 + b_0 \le p + a_0 + b_1 \end{array} \right\}.$$

Therefore, by Lemma A.1,  $\gamma(\mathcal{F}_{p^2+p+1}) = \#A = p(p+1)(p^2+p+2)/8$ .

For the second curve, note that Theorem 4.2 gives

$$\gamma(\mathcal{F}_{p^2+p+1,p^3-1}) = \sum_{j=1}^{p^2+p} (a_{j,1}+1)(a_{j,2}+1)(a_{j,3}+1) - 2(p^2+p),$$
(4.8)

where  $a_{j,1}, a_{j,2}$ , and  $a_{j,3}$  are the coefficients in the base-*p* expansion of j(p-1). Since  $j \in [\![1, p^2 + p]\!]$ , such coefficients are given by (4.7), where  $a_0, a_1 \in [\![0, p]\!]$  and  $a_0 \neq 0$ . Therefore,

$$\gamma(\mathcal{F}_{p^2+p+1,p^3-1}) + 2(p^2+p) = \sum_{a_1=0}^p \left(\sum_{a_0=1}^{a_1} a_1(p+a_0-a_1)(p-a_0+1) + \sum_{a_0=a_1+1}^p (a_1+1)(a_0-a_1)(p-a_0+1)\right) = p(p+1)^2(p^2+p+2)/8,$$

and then  $\gamma(\mathcal{F}_{p^2+p+1,p^3-1}) = p(p+1)(p^3+2p^2+3p-14)/8.$ 

**Theorem 4.4.** If  $h \in \mathbb{Z}_{>0}$  and  $p \neq 2$ , then the *p*-rank of the curves

$$y^{\frac{p^{h}-1}{2}} = x^{\frac{p^{h}-1}{2}} + 1$$
 and  $y^{\frac{p^{h}-1}{2}} = x^{p^{h}-1} + 1$ 

are given by

$$\gamma(\mathcal{F}_{(p^h-1)/2}) = (p+1)^h (p^h+3)/2^{h+2} - 3(p^h-1)/2$$

and

$$\gamma(\mathcal{F}_{(p^h-1)/2,p^h-1}) = (p+1)^h (p^h+1)/2^{h+1} - 2(p^h-1)$$

respectively.

*Proof.* For each integer  $h \ge 1$ , consider the set

$$T_h = \left\{ (i,j) \in \mathbb{N}^2 \mid 0 \le i \le j \le p^h - 1 \text{ and } \binom{j}{i} \not\equiv 0 \pmod{p} \right\}.$$
(4.9)

From Theorem 3.1, to obtain  $\gamma(\mathcal{F}_{(p^h-1)/2}) = (p+1)^h (p^h+3)/2^{h+2} - 3(p^h-1)/2$ , it suffices to prove that the number of elements  $(i, j) \in T_h$  for which both i and j are even is  $(p+1)^h (p^h+3)/2^{h+2}$ . To this end, first note that  $\#T_1 = p(p+1)/2$ , and that for  $(i, j) \in T_h$ , the base-p expansions  $j = \sum_{r=0}^{h-1} a_r p^r$  and  $i = \sum_{r=0}^{h-1} b_r p^r$  along with Lucas' Lemma entail

$$#T_h = (#T_1)^h = (p(p+1)/2)^h.$$
(4.10)

Now for  $u, v \in \{0, 1\}$ , let us consider the partition of  $T_h$  given by the four sets

$$T_h^{(u,v)} = \{(i,j) \in T_h \mid (i,j) \equiv (u,v) \pmod{2}\},\$$

where  $(i, j) \equiv (u, v) \pmod{2}$  stands for  $i \equiv u \pmod{2}$  and  $j \equiv v \pmod{2}$ . We need to prove that  $\#T_h^{(0,0)} = (p+1)^h(p^h+3)/2^{h+2}$ , and for this we proceed by induction on h. The case h = 1 is given by the following easily checked fact.

$$#T_1^{(u,v)} = \begin{cases} (p+1)(p+3)/8 & \text{if } (u,v) = (0,0) \\ (p^2-1)/8 & \text{otherwise.} \end{cases}$$
(4.11)

Now suppose  $h \ge 2$ . For  $j = \sum_{r=0}^{h-1} a_r p^r$  and  $i = \sum_{s=0}^{h-1} b_s p^s$ , since  $(i, j) \equiv (\sum_{r=0}^{h-1} a_r, \sum_{r=0}^{h-1} b_r) \pmod{2}$ , Lucas' Lemma implies that  $(i, j) \in T_h^{(0,0)}$  if and only if

$$(a_{h-1}, b_{h-1}) \in T_1^{(u,v)}$$
 and  $\left(\sum_{s=0}^{h-2} b_s p^s, \sum_{r=0}^{h-2} a_r p^r\right) \in T_{h-1}^{(u,v)}$ 

for some  $(u, v) \in [0, 1]^2$ . Since  $T_{h-1}$  is the disjoint union of  $T_{h-1}^{(0,0)}$  and  $\bigsqcup_{(u,v)\neq(0,0)} T_{h-1}^{(u,v)}$ , equation (4.11) provides

$$#T_h^{(0,0)} = ((p+1)(p+3)#T_{h-1}^{(0,0)} + (p^2 - 1)(#T_{h-1} - #T_{h-1}^{(0,0)}))/8$$
  
=  $(p+1)(4#T_{h-1}^{(0,0)} + (p-1)#T_{h-1})/8$   
=  $(p+1)^h(p^h+3)/2^{h+2},$ 

where the latter equality follows from  $\#T_{h-1}^{(0,0)} = (p+1)^{h-1}(p^{h-1}+3)/2^{h+1}$ , given by the induction hypothesis, and from  $\#T_{h-1} = (p(p+1)/2)^{h-1}$  in (4.10).

For the curve  $\mathcal{F}_{p^h-1,(p^h-1)/2}$ , in accordance with Theorem 3.1, it only remains to prove that the total number of  $(i,j) \in T_h$  for which j is even, that is,  $\#T_h^{(0,0)} + \#T_h^{(0,1)}$ , is  $(p+1)^h(p^h+1)/2^{h+1}$ . To this end, note that the maps

$$T_{h}^{(1,0)} \longrightarrow T_{h}^{(0,1)}, \ (i,j) \mapsto (p^{h} - 1 - j, p^{h} - 1 - i) \quad \text{and} \quad T_{h}^{(0,1)} \longrightarrow T_{h}^{(1,1)}, \ (i,j) \mapsto (j - i, j)$$

are bijections, given that  $p^h - 1 = \sum_{r=0}^{p-1} (p-1)p^r$  is even and that  $\binom{j}{i} = \binom{j}{j-i}$ . Therefore, since  $\#T_h = p^h (p+1)^h / 2^h$  and  $\#T_h^{(0,0)} = (p+1)^h (p^h+3) / 2^{h+2}$ , we have

$$#T_h^{(1,0)} = #T_h^{(0,1)} = #T_h^{(1,1)} = (p+1)^h (p^h - 1)/2^{h+2}.$$

Hence

$$\#T_h^{(0,0)} + \#T_h^{(0,1)} = (p+1)^h (p^h+1)/2^{h+1}$$

and the result follows.

From Corollary 3.3, the *p*-rank of the Fermat curve  $\mathcal{F}_{p^h-1}$ :  $y^{p^h-1} = x^{p^h-1} + 1$  is  $\#T_h - 3(p^h - 1)$ , where  $T_h$  is the set given in (4.9). Thus equation (4.10) implies that the *p*-rank of  $\mathcal{F}_{p^h-1}$  is  $(p(p+1)/2)^h - 3(p^h - 1)$ , a result proved by Bassa and Beelen in [2, Theorem 19]. In what follows, we provide a preliminary step to obtain formulas for the *p*-rank of the curves

$$y^{p^m \pm 1} = x^{p^n \pm 1} + 1$$

natural generalizations of the Fermat curve  $y^{p^{h}-1} = x^{p^{h}-1} + 1$ .

**Lemma 4.5.** Let  $u, v \in \mathbb{Z}_{>0}$  and w = gcd(u, v). Suppose that  $I, J \in [[0, p^{vu/w} - 1]]$  have base-*p* expansions given by

$$I = \sum_{b=0}^{\frac{v}{w}-1} \sum_{s=0}^{u-1} b_s p^{ub+s} \text{ and } J = \sum_{a=0}^{\frac{u}{w}-1} \sum_{r=0}^{v-1} a_r p^{va+r}.$$
(4.12)

Then the following are equivalent.

- (i)  $\binom{J}{I} \not\equiv 0 \pmod{p}$
- (ii) For every  $(s,t) \in \llbracket 0, u-1 \rrbracket \times \llbracket 0, w-1 \rrbracket$  such that  $s \equiv t \pmod{w}$ , we have

$$0 \le b_s \le \min \left\{ a_{\theta w+t} \mid \theta \in \left[\!\left[0, u/w - 1\right]\!\right] \right\}.$$

*Proof.* Fix  $s \in [[0, u - 1]]$ . In the base-*p* expansion of *I*, the powers  $p^{ub+s}$ , with  $b \in [[0, v/w - 1]]$ , have the same coefficient  $b_s$ . The corresponding coefficients of such powers in the base-*p* expansion of *J* are the v/w coefficients  $a_r$ , with  $r \in [[0, v - 1]]$ , such that  $r \equiv ub + s \pmod{v}$ . One can check that the set of such values of *r* is given by

$$\{r_1, \dots, r_{v/w}\} := \{\theta w + t \mid \theta \in [\![0, v/w - 1]\!]\},\tag{4.13}$$

where  $t \equiv s \pmod{w}$  and  $t \in [\![0, w-1]\!]$ . Note that this set depends only on the reduction of  $s \in [\![0, u-1]\!]$  modulo w. Therefore, for any fixed  $t \in [\![0, w-1]\!]$ , there are exactly u/w values  $s_1, \ldots, s_{u/w} \in [\![0, u-1]\!]$ , with  $s_i \equiv t \pmod{w}$ , that give rise to the same set (4.13). That is, for all  $b \in [\![0, v/w-1]\!]$ , the set of coefficients of  $p^{ub+s_1}, \ldots, p^{ub+s_{n/d}}$  in the base-p expansion of J is given by  $\{a_{r_1}, \ldots, a_{r_{v/w}}\}$ . Thus by Lucas' Lemma,  $\binom{J}{I} \not\equiv 0 \pmod{p}$  if and only if for every  $s \in [\![0, u-1]\!]$ , we have  $0 \le b_s \le \min \{a_{\theta w+t} \mid \theta \in [\![0, u/w-1]\!]\}$ , where  $t \equiv s \pmod{w}$  and  $t \in [\![0, w-1]\!]$ .

**Theorem 4.6.** Let  $n, m \in \mathbb{Z}_{>0}$ , and d = gcd(n, m). Then the *p*-rank of  $\mathcal{F} : y^{p^m-1} = x^{p^n-1} + 1$  is

$$\gamma = \left(\sum_{i=0}^{p-1} \left( (i+1)^{\frac{m}{d}} - i^{\frac{m}{d}} \right) (p-i)^{\frac{n}{d}} \right)^d - (p^m + p^n + p^d - 3)$$

*Proof.* Let  $q = p^{mn/d}$ ,  $\alpha = (q-1)/(p^n-1)$ , and  $\beta = (q-1)/(p^m-1)$ . By Theorem 3.1, the *p*-rank of  $\mathcal{F}$  is  $\gamma = \#T - (p^m + p^n + p^d - 3)$ , where

$$T = \left\{ (i,j) \in \mathbb{N}^2 \mid 0 \le i\alpha \le j\beta \le q-1 \text{ and } \binom{j\beta}{i\alpha} \not\equiv 0 \pmod{p} \right\}$$

For  $(i, j) \in T$ , consider the base-*p* expansions

$$j = a_{m-1}p^{m-1} + \dots + a_1p + a_0$$
 and  $i = b_{n-1}p^{n-1} + \dots + b_1p + b_0$ ,

and note that

$$j\beta = \sum_{a=0}^{\frac{n}{d}-1} \sum_{r=0}^{m-1} a_r p^{ma+r} \text{ and } i\alpha = \sum_{b=0}^{\frac{m}{d}-1} \sum_{s=0}^{n-1} b_s p^{nb+s}$$

Thus by Lemma 4.5, we have #T = #L, where

$$L = \left\{ (a_0, \dots, a_{m-1}, b_0, \dots, b_{n-1}) \middle| \begin{array}{l} a_r \in [\![0, p-1]\!], \\ 0 \le b_s \le \min \{a_{\theta d+t} \mid \theta \in [\![0, m/d-1]\!]\}, \\ t \in [\![0, d-1]\!], \text{ and } t \equiv s \pmod{d} \end{array} \right\}.$$

Let  $t \in [\![0, d-1]\!]$ . Note that for each  $u = (a_0, \ldots, a_{m-1}, b_0, \ldots, b_{n-1}) \in A$ , the set of the indices  $\{r_1, \ldots, r_{m/d}, s_1, \ldots, s_{n/d}\}$  (cf. proof of Lemma 4.5), whose corresponding entries in u are such that  $0 \leq b_{s_j} \leq \min \{a_{r_i} \mid i \in [\![1, m/d]\!]\}$ , is characterized by the condition  $r_i \equiv s_j \equiv t \pmod{d}$  for all  $(i, j) \in [\![1, m/d]\!] \times [\![1, n/d]\!]$ . Thus rearranging the entries of u, the set L can be seen as the cartesian product of the d sets

$$L_{t} = \left\{ \left( a_{r_{1}}, \dots, a_{r_{m/d}}, b_{s_{1}}, \dots, b_{s_{n/d}} \right) \middle| \begin{array}{l} a_{r_{i}} \in \llbracket 0, p-1 \rrbracket, \\ 0 \le b_{s_{j}} \le \min \left\{ a_{r_{i}} \mid i \in \llbracket 1, m/d \rrbracket \right\}, \\ r_{i} \equiv s_{j} \equiv t \pmod{d} \end{array} \right\}$$

each of which has cardinality

$$#L_t = \sum_{i=0}^{p-1} \left( (i+1)^{m/d} - i^{m/d} \right) (p-i)^{n/d},$$

given by Lemma A.2. Therefore,  $\#T = \#L = (\#L_t)^d$ , and the result follows.

**Corollary 4.7.** Let  $n, m \in \mathbb{Z}_{>0}$  be odd, and let u and v be the orders of 2 in  $(\mathbb{Z}/m\mathbb{Z})^{\times}$  and  $(\mathbb{Z}/n\mathbb{Z})^{\times}$ , respectively. If gcd(u, v) = 1, then the 2-rank of  $\mathcal{F} : y^m = x^n + 1$  is zero.

Proof. Considering the morphism

$$\mathcal{F}_{2^u-1,2^v-1} \longrightarrow \mathcal{F}, \quad (x,y) \mapsto (x^{(2^v-1)/n}, y^{(2^u-1)/m}),$$

it follows from item (ii) of Lemma 2.4 that we only need to prove that  $\gamma(\mathcal{F}_{2^u-1,2^v-1}) = 0$ , which is given by Theorem 4.6, since gcd(u, v) = 1.

Let  $n, m \in \mathbb{Z}_{>0}$ . As mentioned in Remark 2.8, the supersingularity of the curves  $\mathcal{F} : y^{p^m+1} = x^{p^n+1} + 1$  depends on whether or not  $\nu_2(n) = \nu_2(m)$ , where  $\nu_2 : \mathbb{Z} \to \mathbb{N} \cup \{\infty\}$  is the 2-adic valuation. Nevertheless, in the following theorem, we show that, in any case, such curves have zero *p*-rank.

**Theorem 4.8.** Let  $n, m \in \mathbb{Z}_{>0}$  be divisors of  $p^u + 1$  and  $p^v + 1$ , respectively, where  $u, v \in \mathbb{Z}_{\geq 0}$ . Then  $\mathcal{F} : y^m = x^n + 1$  has zero *p*-rank.

*Proof.* Considering the map  $\mathcal{F}_{p^v+1,p^u+1} \longrightarrow \mathcal{F}$ ,  $(x,y) \mapsto (x^{(p^u+1)/n}, y^{(p^v+1)/m})$ , and item (ii) of Lemma 2.4, it suffices to prove that  $\gamma(\mathcal{F}_{p^v+1,p^u+1}) = 0$ . From (3.3), we have  $\gamma(\mathcal{F}_{p^v+1,p^u+1}) = \#A$ , where

$$A = \left\{ (i,j) \in \mathbb{N}^2 \mid p^v + 1 \le i(p^v + 1) < j(p^u + 1) \le (p^u + 1)p^v \text{ and } \binom{j\beta}{i\alpha} \not\equiv 0 \pmod{p} \right\},$$

with  $d = \gcd(v, u), q = p^{2vu/d}, \alpha = (q-1)/(p^u+1)$ , and  $\beta = (q-1)/(p^v+1)$ . Suppose  $A \neq \emptyset$ , and let  $(i, j) \in A$ . Since  $j \in [\![1, p^v]\!]$  and  $i \in [\![1, p^u]\!]$ , we may consider the following base-p expansions

$$j - 1 = a_{v-1}p^{v-1} + \dots + a_1p + a_0$$
 and  $i - 1 = b_{u-1}p^{u-1} + \dots + b_1p + b_0.$  (4.14)

Observe that

$$j\beta = \sum_{a=0}^{\frac{n}{d}-1} \sum_{r=0}^{2v-1} c_r p^{2va+r} \text{ and } i\alpha = \sum_{b=0}^{\frac{n}{d}-1} \sum_{s=0}^{2u-1} e_s p^{2ub+s},$$

where

$$c_r = \begin{cases} p-1-a_r & \text{if } r \in \llbracket 0, v-1 \rrbracket \\ a_{r-v} & \text{if } r \in \llbracket v, 2v-1 \rrbracket \end{cases} \text{ and } e_s = \begin{cases} p-1-b_s & \text{if } s \in \llbracket 0, u-1 \rrbracket \\ b_{s-u} & \text{if } s \in \llbracket u, 2u-1 \rrbracket. \end{cases}$$

Let  $(s, w) \in [\![0, u-1]\!] \times [\![0, d-1]\!]$ , with  $s \equiv w \pmod{d}$ . By Lemma 4.5,  $\binom{j\beta}{i\alpha} \not\equiv 0 \pmod{p}$  implies

$$0 \le e_s, e_{u+s} \le \min\{c_{\theta d+w} \mid \theta \in [[0, 2v/d - 1]]\} = \min\{a_{\theta d+w}, p - 1 - a_{\theta d+w} \mid \theta \in [[0, v/d - 1]]\}.$$

In particular,

$$\max\{b_s, p-1-b_s\} \le \min\{a_{\theta d+w}, p-1-a_{\theta d+w} \mid \theta \in [\![0, v/d-1]\!]\}.$$
(4.15)

It is easy to see that (4.15) implies p > 2 and  $b_s = a_{\theta d+w} = (p-1)/2$ . Hence from (4.14), we have  $i = (p^u + 1)/2$  and  $j = (p^v + 1)/2$ , which contradicts the condition  $i(p^v + 1) < j(p^u + 1)$  in the definition of the set A. Therefore,  $\gamma(\mathcal{F}_{p^v+1,p^u+1}) = \#A = 0$ .

**Theorem 4.9.** Let  $m, n \in \mathbb{Z}_{>0}$ ,  $d = \operatorname{gcd}(n, m)$ , and  $\gamma$  be the *p*-rank of

$$\mathcal{F}: y^{p^m + 1} = x^{p^n - 1} + 1.$$

(i) If n/d is even, then

$$\gamma = \Big(\sum_{i=0}^{p-1} \left( (p-i)(i+1) \right)^{\frac{n}{2d}} + \sum_{1 \le j \le i \le p-1} \left( (j+1)^{\frac{m}{d}} - 2j^{\frac{m}{d}} + (j-1)^{\frac{m}{d}} \right) \left( (p-i)(i-j+1) \right)^{\frac{n}{2d}} \Big)^d - (p^m + p^d).$$

(ii) If n/d is odd, then

$$\gamma = \frac{1}{2^n} \Big( (p+1)^{\frac{n}{d}} + \sum_{i=1}^{\frac{p-1}{2}} \big( (2i+1)^{\frac{m}{d}} - (2i-1)^{\frac{m}{d}} \big) (p-2i+1)^{\frac{n}{d}} \Big)^d - (p^m+1),$$

when  $p \neq 2$ , and  $\gamma = 0$  otherwise.

*Proof.* Let  $D = \gcd(2m, n)$ ,  $q = p^{2mn/D}$ ,  $\alpha = (q-1)/(p^n-1)$ , and  $\beta = (q-1)/(p^m+1)$ . By Theorem 3.1,  $\gamma = \#T - (p^m + p^n + \gcd(p^m + 1, p^n - 1))$ , where

$$T = \left\{ (i,j) \in \mathbb{N}^2 \mid 0 \le i\alpha \le j\beta \le q-1 \text{ and } \binom{j\beta}{i\alpha} \not\equiv 0 \pmod{p} \right\}$$

Since  $T \subseteq [\![0, p^n - 1]\!] \times [\![0, p^m + 1]\!]$  and  $\#\{(i, j) \in T \mid j \in \{0, p^m + 1\}\} = p^n + 1$ , we will focus on the pairs  $(i, j) \in T$  with  $j \in [\![1, p^m]\!]$ . Considering the base-p expansions

$$j-1 = a_{m-1}p^{m-1} + \dots + a_1p + a_0$$
 and  $i = b_{n-1}p^{n-1} + \dots + b_1p + b_0$ ,

observe that

$$j\beta = \sum_{a=0}^{\frac{n}{D}-1} \sum_{r=0}^{2m-1} c_r p^{2ma+r} \text{ and } i\alpha = \sum_{b=0}^{\frac{2m}{D}-1} \sum_{s=0}^{n-1} b_s p^{nb+s},$$
(4.16)

where

$$c_r = \begin{cases} p - 1 - a_r & \text{if } r \in [\![0, m - 1]\!] \\ a_{r-m} & \text{if } r \in [\![m, 2m - 1]\!]. \end{cases}$$
(4.17)

By Lemma 4.5,  $\binom{j\beta}{i\alpha} \not\equiv 0 \pmod{p}$  if and only if for every  $s \in \llbracket 0, n-1 \rrbracket$ , we have

$$0 \le b_s \le \min A_t$$

where  $0 \le t \le D - 1$ ,  $t \equiv s \pmod{D}$ , and

$$A_t = \{ c_{D\theta+t} \mid \theta \in [\![0, 2m/D - 1]\!] \}.$$
(4.18)

Now consider the following.

Case n/d even. We have D = 2d, and then (4.18) reads A<sub>t</sub> = {c<sub>2dθ+t</sub> | θ ∈ [[0, m/d - 1]]}. Clearly for any u ∈ N, there exists a unique t ∈ [[0, d - 1]] such that either u ≡ t (mod 2d) or u ≡ t + d (mod 2d). In both cases, we have u ≡ t (mod d), and the latter case occurs if and only if [u/d] is odd. That is, taking into account the parity of [u/d], we may replace reduction modulo 2d by reduction modulo d. This observation underlies the proof of this case.

Note that for  $t \in [0, d-1]$ , considering the sets

$$A_t \cup A_{d+t} = \{ c_{\theta d+t} \mid \theta \in [\![0, 2m/d - 1]\!] \}$$

and the values  $s \in [0, n-1]$ , with  $s \equiv t \pmod{d}$ , the condition  $\binom{j\beta}{i\alpha} \not\equiv 0 \pmod{p}$  is equivalent to

$$0 \le b_s \le \left\{ \begin{array}{ll} \min A_t & \text{if } \lfloor s/d \rfloor \text{ is even} \\ \min A_{d+t} & \text{if } \lfloor s/d \rfloor \text{ is odd.} \end{array} \right.$$

Therefore, Lemma 4.4 gives  $\#T = \#L + p^n + 1$ , where

$$L = \left\{ \begin{array}{c} (a_0, \dots, a_{m-1}, b_0, \dots, b_{n-1}) \\ 0 \le b_s \le \min A_t \text{ if } \lfloor s/d \rfloor \text{ is even,} \\ 0 \le b_s \le \min A_{t+d} \text{ if } \lfloor s/d \rfloor \text{ is odd,} \\ t \in \llbracket 0, d-1 \rrbracket, \text{ and } s \equiv t \pmod{d} \end{array} \right\}.$$

The two sets  $A_t$  and  $A_{d+t}$  are given in terms of the same elements  $a_{\theta d+t}$ , although in different ways. Indeed for  $c_{d\theta+t} \in A_t \cup A_{t+d}$ , equation (4.17) yields

$$c_{d\theta+t} = \begin{cases} p-1-a_{d\theta+t} & \text{if } \theta \le m/d - 1\\ a_{d(\theta-m/d)+t} & \text{if } \theta \ge m/d. \end{cases}$$
(4.19)

Since m/d is odd, collecting the elements  $c_{d\theta+t}$  in (4.19) according to the parity of  $\theta$  gives

$$A_{t} = \left\{ p - 1 - a_{d\theta_{0}+t}, a_{d\theta_{1}+t} \middle| \begin{array}{l} \theta_{i} \in [\![0, m/d - 1]\!], \\ \theta_{i} \equiv i \pmod{2} \end{array} \right\} = \left\{ p - 1 - e_{\theta d+t} \mid \theta \in [\![0, m/d - 1]\!] \right\}$$

and

where

$$\begin{split} A_{d+t} &= \left\{ p - 1 - a_{d\theta_0 + t}, \, a_{d\theta_1 + t} \middle| \begin{array}{l} \theta_i \in \llbracket 0, m/d - 1 \rrbracket, \\ \theta_i &\equiv i \pmod{2} \end{array} \right\} = \left\{ e_{\theta d + t} \mid \theta \in \llbracket 0, m/d - 1 \rrbracket \right\}, \\ e_{\theta d + t} &\coloneqq \left\{ \begin{array}{l} a_{d\theta + t} & \text{if } \theta \text{ is even} \\ p - 1 - a_{d\theta + t} & \text{if } \theta \text{ is odd} \end{array} \right., \text{ with } \theta \in \llbracket 0, m/d - 1 \rrbracket. \end{split}$$

Now for a fixed  $t \in [[0, d-1]]$  and  $u = (e_0, \ldots, e_{m-1}, b_0, \ldots, b_{n-1}) \in L$ , we consider the set of indices  $\{r_1, \ldots, r_{m/d}, s_1, \ldots, s_{n/d}\}$ , whose corresponding entries in u are such that

$$0 \le b_{s_{\lambda}} \le \begin{cases} \min \left\{ e_{r_{\mu}} : \mu \in \llbracket 1, m/d \rrbracket \right\} & \text{if } \lambda \in \llbracket 1, n/(2d) \rrbracket \\ \min \left\{ p - 1 - e_{r_{\mu}} : \mu \in \llbracket 1, m/d \rrbracket \right\} & \text{if } \lambda \in \llbracket n/(2d) + 1, n/d \rrbracket \end{cases}$$

where  $r_{\mu} \equiv s_{\lambda} \equiv t \pmod{d}$  for  $(\mu, \lambda) \in [\![1, m/d]\!] \times [\![1, m/d]\!]$ , and  $\lfloor s_{\lambda}/d \rfloor$  is odd for  $\lambda \leq n/2d$ , and even otherwise. Thus L can be seen as the cartesian product of the d sets

$$L_{t} = \left\{ (e_{r_{1}}, \dots, e_{r_{m/d}}, b_{s_{1}}, \dots, b_{s_{n/d}}) \middle| \begin{array}{l} e_{r_{\mu}} \in \llbracket 0, p - 1 \rrbracket, \\ 0 \le b_{s_{\lambda}} \le \min\{e_{r_{\mu}} \mid \mu \in \llbracket 1, m/d \rrbracket\}, \ \lambda \le \frac{n}{2d}, \\ 0 \le b_{s_{\lambda}} \le \min\{p - 1 - e_{r_{\mu}} \mid \mu \in \llbracket 1, m/d \rrbracket\}, \ \lambda > \frac{n}{2d}, \\ s_{\lambda} \equiv r_{\mu} \equiv t \pmod{d} \end{array} \right\}$$
(4.20)

• Case n/d odd. Now we have D = d, which simplifies the analysis. Indeed equations (4.17) and (4.18) immediately yield

$$A_t = \{a_{\theta d+t}, \, p-1 - a_{\theta d+t} \mid \theta \in [\![0, m/d - 1]\!]\},\$$

and then  $\#T = \#L + p^n + 1$ , where

$$L = \left\{ (a_0, \dots, a_{m-1}, b_0, \dots, b_{n-1}) \middle| \begin{array}{l} a_r \in [\![0, p-1]\!], \\ 0 \le b_s \le \min \{a_{\theta d+t}, p-1 - a_{\theta d+t} \mid \theta \in [\![0, m/d - 1]\!]\}, \\ t \in [\![0, d-1]\!], \text{ and } s \equiv r \equiv t \pmod{d} \end{array} \right\}.$$

As in the first case, L can be seen as the cartesian product of the d sets

$$L_{t} = \left\{ (a_{r_{1}}, \dots, a_{r_{m/d}}, b_{s_{1}}, \dots, b_{s_{n/d}}) \middle| \begin{array}{l} a_{r_{i}} \in \llbracket 0, p-1 \rrbracket, \\ 0 \le b_{s_{j}} \le \min\{a_{r_{i}}, p-1-a_{r_{i}} \mid i \in \llbracket 1, m/d \rrbracket \} \\ s \equiv r \equiv t \pmod{d} \end{array} \right\}.$$
(4.21)

Independent of t, the cardinalities of the sets  $L_t$  in (4.20) and (4.21) are given by Lemmas A.4 and A.3, respectively. Using Lemma 2.6 and  $\#T = (\#L_t)^d + p^n + 1$  for each of the two cases, the result follows.

**Corollary 4.10.** Let  $m, n \in \mathbb{Z}_{>0}$  be divisors of  $2^u + 1$  and  $2^v - 1$ , respectively. If  $v / \operatorname{gcd}(u, v)$  is odd, then  $\mathcal{F} : y^m = x^n + 1$  has zero 2-rank.

*Proof.* As in the proof of Corollary 4.7, it suffices to show that  $\gamma(\mathcal{F}_{2^u+1,2^v-1}) = 0$ , which this follows from (ii) in Theorem 4.9.

# **5** Formulas for the p-rank of $y^2 = x^n + 1$

Throughout this section, p is an odd prime, n is a positive integer coprime to p, and  $\gamma(\mathcal{C}_n)$  is the p-rank of the curve  $\mathcal{C}_n : y^2 = x^n + 1$  defined over  $\mathbb{F}_q$ , where  $q = p^h$  and h is the order of p in  $(\mathbb{Z}/n\mathbb{Z})^{\times}$ . Recall that Corollary 3.4 gives the p-rank of the curve  $\mathcal{C}_n$  in terms of the cardinality of

$$S = \left\{ i \in [\![0, \lfloor n/2 \rfloor]\!] \mid \binom{q-1}{2}{i\alpha} \neq 0 \pmod{p} \right\},\$$

where  $\alpha = (q-1)/n$ . Since  $\lfloor n/2 \rfloor \alpha$  is the largest multiple of  $\alpha$  in [[0, (q-1)/2]], Lucas' Lemma yields the following version of Corollary 3.4.

*Remark* 5.1. If *M* is the number of values  $u = u_1 p^{h-1} + \cdots + u_{h-1} p + u_h$ , with  $u_i \in [[0, (p-1)/2]]$ , that are divisible by  $\alpha = (q-1)/n$ , then

$$\gamma(\mathcal{C}_n) = \begin{cases} M-1 & \text{if } n \text{ is odd} \\ M-2 & \text{if } n \text{ is even} . \end{cases}$$
(5.1)

The following provides some consequences of the results in the previous section.

**Corollary 5.2.** *If*  $r \in \mathbb{Z}_{>0}$ *, then* 

(i)  $\gamma(\mathcal{C}_{p^r-1}) = ((p+1)/2)^r - 2$ 

(ii) 
$$\gamma(\mathcal{C}_{2(p^r+1)}) = ((p+1)/2)^r$$

- (iii)  $\gamma(\mathcal{C}_{2(p^r-1)}) = ((p+1)/2)^r 2$
- (iv)  $\gamma(\mathcal{C}_{p^{2r}+p^r+1}) = ((p+3)(p+1)/8)^r$
- (v)  $\gamma(\mathcal{C}_{2(p^{2r}+p^r+1)}) = 2((p+3)(p+1)/8)^r$

(vi) 
$$\gamma(\mathcal{C}_{p^{3r}-p^{2r}+p^r-1}) = ((p^2+2p+3)(p+1)/12)^r - 2.$$

*Proof.* Assertion (i) follows directly from (4.6) in Theorem 4.2. Assertion (v) follows from (iv) together with  $\gamma(\mathcal{C}_{2(p^{2r}+p^r+1)}) = 2\gamma(\mathcal{C}_{p^{2r}+p^r+1})$ , given by item (iv) in Corollary 3.6. The proofs of (ii) and (iii) are simpler versions of the proofs of (iv) and (vi), respectively. Thus we present proofs for (iv) and (vi) only. Regarding (iv), note that  $q = p^{3r}$  and  $\alpha = (p^{3r} - 1)/(p^{2r} + p^r + 1) = p^r - 1$ . From Remark 5.1, it suffices to count the values

$$u = \sum_{i=1}^{r} a_i p^{3r-i} + \sum_{i=1}^{r} b_i p^{2r-i} + \sum_{i=1}^{r} c_i p^{r-i},$$
(5.2)

with  $a_i, b_i, c_i \in [[0, (p-1)/2]]$ , that are divisible by  $p^r - 1$ . Since

$$u \equiv \sum_{i=1}^{r} (a_i + b_i + c_i) p^{r-i} \pmod{p^r - 1},$$

it follows that  $p^r - 1$  divides u if and only if it divides  $u_0 := \sum_{i=1}^r (a_i + b_i + c_i)p^{r-i}$ . Note that the condition  $a_i, b_i, c_i \in [0, (p-1)/2]$  implies  $u_0 \in [0, 3(p^r - 1)/2]$ , and then  $p^r - 1$  divides  $u_0$  if and only if u = 0 or

 $\sum_{i=1}^{r} (a_i + b_i + c_i) p^{r-i} = p^r - 1.$  Thus we are left with the problem of counting  $a_i, b_i, c_i \in [[0, (p-1)/2]]$  for which  $\sum_{i=1}^{r} (a_i + b_i + c_i) p^{r-i} = p^r - 1.$  The uniqueness of the base-*p* expansion implies that the former equality is equivalent

 $\sum_{i=1}^{n} (a_i + b_i + c_i)p^{-1} = p^{-1}$ . The uniqueness of the base *p* expansion inpress that the former equality is equivalent to  $a_i + b_i + c_i = p - 1$ , given that  $a_i + b_i + c_i \leq 3(p-1)/2 \leq 2(p-1)$ . Since  $c_i = p - 1 - (a_i + b_i)$ , we only need to count the values  $a_i, b_i \in [[0, (p-1)/2]]$  such that  $a_i + b_i \geq (p-1)/2$ . Therefore, the number of values in (5.2) is given by

$$M = 1 + \left(\sum_{a_i=0}^{(p-1)/2} ((p+1)/2 - a_i)\right)^r = 1 + \left(\frac{(p+1)(p+3)}{8}\right)^r,$$

and the result follows from (5.1) as  $n = p^{2r} + p^r + 1$  is odd. The proof of (vi) is similar to that of (iv). Indeed, for item (vi) we have  $q = p^{4r}$  and  $\alpha = p^r + 1$ , and the same argument leads to count  $a_i, b_i, c_i, d_i \in [[0, (p-1)/2]]$  for which

$$u_0 := \left|\sum_{i=1}^r (-a_i + b_i - c_i + d_i)p^{r-i}\right| \le \sum_{i=1}^r |-a_i + b_i - c_i + d_i|p^{r-i} \le \sum_{i=1}^r (p-1)p^{r-i} = p^r - 1$$

is divisible by  $p^r + 1$ . Clearly, this condition is equivalent to  $u_0 = 0$ , that is,  $a_i + c_i = b_i + d_i$  for all  $i \in [\![1, r]\!]$ . Note that for each  $k \in [\![0, p-1]\!]$ , the number of solutions  $(x, y) \in [\![0, (p-1)/2]\!]^2$  to x + y = k is k + 1 if  $k \le (p-1)/2$ , and p - k otherwise. Therefore, the number of values of u, in the notation of Remark (5.1), is given by

$$M = \left( ((p+1)/2)^2 + 2\sum_{i=0}^{(p-1)/2} i^2 \right)^r = \left( \frac{(p^2 + 2p + 3)(p+1)}{12} \right)^r,$$

and since n is even, the result follows.

In the remainder of this section, we incorporate an additional technique to provide a formula for the p-rank  $\gamma(C_n)$  given in (5.1). This will come from observing that the problem of counting the values

$$u = x_1 + x_2 p + \dots + x_h p^{h-1}$$

with  $x_i \in [[0, (p-1)/2]]$ , that are divisible by  $\alpha = (q-1)/n$  can be framed in terms of the problem of finding the number of solutions  $(x_1, \ldots, x_h) \in [[0, \alpha t + s]]^h$  to

$$_1X_1 + \dots + a_hX_h \equiv 0 \pmod{\alpha},$$

where  $a_i \in (\mathbb{Z}/\alpha\mathbb{Z})^{\times}$ , and  $(p-1)/2 = t\alpha + s$  for unique integers  $t \in \mathbb{N}$  and  $s \in [[0, \alpha - 1]]$ . In this regard, the following presents some facts that will be used in the subsequent proofs.

**Lemma 5.3.** Let  $h, t, \alpha \in \mathbb{Z}_{>0}$  and  $a_1, \ldots, a_h \in (\mathbb{Z}/\alpha\mathbb{Z})^{\times}$ . If  $b, s \in [[0, \alpha - 1]]$ , then the number N of solutions  $(x_1, \ldots, x_h) \in [[0, t\alpha + s]]^h$  to

$$a_1 X_1 + \dots + a_h X_h \equiv b \pmod{\alpha}$$
(5.3)

is given by

$$N = \frac{(t\alpha + s + 1)^h - (s + 1)^h}{\alpha} + \delta,$$
(5.4)

where  $\delta \in [[0, (s+1)^{h-1}]]$  is the number of solutions  $(y_1, \ldots, y_h) \in [[0, s]]^h$  to (5.3). Moreovver,

(i) if s = 0, then  $\delta = 1$  when b = 0, and  $\delta = 0$  otherwise

(ii) if 
$$s = \alpha - 1$$
, then  $\delta = \alpha^{h-1}$ 

(iii) if 
$$s = \alpha - 2$$
, then

$$\delta = \begin{cases} \frac{(\alpha - 1)^h - (-1)^h}{\alpha} + (-1)^h & \text{if } \sum_{i=1}^h a_i \equiv -b \pmod{\alpha} \\ \frac{(\alpha - 1)^h - (-1)^h}{\alpha} & \text{otherwise} \end{cases}$$

(iv) if 
$$s = \alpha/2$$
,  $b = 0$ , and h is odd, then  $\delta = \frac{(\alpha/2 + 1)^h - 1}{\alpha} + \frac{1}{2}$ .

*Proof.* For each  $h \in \mathbb{Z}_{>0}$  and  $b \in [[0, \alpha - 1]]$ , let  $N_{h,b}$  and  $\delta(h, b)$  denote the number of solutions to

$$a_1 X_1 + \dots + a_h X_h \equiv b \pmod{\alpha}$$

in  $[0, t\alpha + s]^h$  and  $[0, s]^h$ , respectively. We proceed by induction on h to prove that

$$N_{h,b} = \frac{(t\alpha + s + 1)^h - (s + 1)^h}{\alpha} + \delta(h, b).$$

Clearly,  $N_{1,b} = t + \delta(1, b)$ , which gives the case h = 1. Let us assume the result for  $h-1 \ge 1$ . For each  $j \in [0, t\alpha + s]$ , let  $N_{h,b}^j$  be the number of solutions  $(x_1, \ldots, x_{h-1}) \in [0, t\alpha + s]^{h-1}$  to

$$a_1X_1 + \dots + a_{h-1}X_{h-1} + a_hj \equiv b \pmod{\alpha}$$

Note that  $N_{h,b}^j = N_{h-1,b_j}$ , where  $b_j \in [0, \alpha - 1]$  and  $b_j \equiv b - a_h j \pmod{\alpha}$ , and

$$N_{h,b} = \sum_{j=0}^{t\alpha+s} N_{h,b}^j = t \sum_{j=0}^{\alpha-1} N_{h-1,j} + \sum_{j=0}^s N_{h-1,b_j} = t(t\alpha+s+1)^{h-1} + \sum_{j=0}^s N_{h-1,b_j}.$$

From the induction hypothesis,

$$\sum_{j=0}^{s} N_{h-1,b_j} = (s+1) \cdot \frac{(t\alpha+s+1)^{h-1} - (s+1)^{h-1}}{\alpha} + \sum_{j=0}^{s} \delta(h-1,b_j)$$
$$= \frac{(s+1)(t\alpha+s+1)^{h-1} - (s+1)^h}{\alpha} + \delta(h,b),$$

and then  $N_{h,b} = \frac{(t\alpha + s + 1)^h - (s + 1)^h}{\alpha} + \delta(h, b)$ , which completes the proof. In addition, by the definition of  $\delta(h, b)$ , one can readily see that  $\delta(h, b) \in [0, (s + 1)^{h-1}]$ . Assertions (i) and (ii) are straightforward, and the proof of (iii) follows by induction, as in the proof of (5.4). To prove assertion (iv), let M be the set of the solutions to

$$a_1X_1 + \dots + a_hX_h \equiv s \pmod{2s}$$

in  $\llbracket 0, s \rrbracket^h$ . Since b = 0 and h is odd, we have that  $(x_1, \ldots, x_h) \in \llbracket 0, s \rrbracket^h$  is a solution to (5.3) if and only if  $(s-x_1, \ldots, s-x_h) \in M$ . This implies  $\delta = \#M$ . Note that  $\delta + \#M$  is the number of solutions  $(x_1, \ldots, x_h) \in \llbracket 0, s \rrbracket^h$  to  $a_1X_1 + \cdots + a_hX_h \equiv 0 \pmod{s}$ ,

and then assertion (i) gives  $2\delta = \delta + \#M = \frac{(s+1)^h - 1}{s} + 1$ , that is,  $\delta = \frac{(\alpha/2+1)^h - 1}{\alpha} + \frac{1}{2}$ .

**Theorem 5.4.** Let  $n \in \mathbb{Z}_{>0}$  and  $\alpha = (p^h - 1)/n$ , where h is the order of p in  $(\mathbb{Z}/n\mathbb{Z})^{\times}$ . If  $s \in [0, \alpha - 1]$  is such that  $s \equiv (p-1)/2 \pmod{\alpha}$ , then

$$\gamma(\mathcal{C}_n) = \frac{n}{p^h - 1} \left( \left(\frac{p+1}{2}\right)^h - \delta_n \right), \text{ where } \delta_n = \begin{cases} (s+1)^h - \alpha(\delta-1) & \text{if } n \text{ is odd} \\ (s+1)^h - \alpha(\delta-2) & \text{if } n \text{ is even} \end{cases}$$

and  $\delta \in [\![1,(s+1)^{h-1}]\!]$  is the number of solutions  $(x_1,\ldots,x_h) \in [\![0,s]\!]^h$  to

$$X_1 + pX_2 + \dots + p^{h-1}X_h \equiv 0 \pmod{\alpha}.$$
(5.5)

Moreover,

(i) if 
$$\alpha \mid (p-1)/2$$
, then n is even and  $\delta_n = \alpha + 1$   
(ii) if  $\alpha \mid (p+1)/2$ , then  $\delta_n = \begin{cases} \alpha & \text{if } n \text{ is odd} \\ 2\alpha & \text{if } n \text{ is even} \end{cases}$   
(iii) if  $\alpha \mid (p+3)/2$  and  $p-1 \nmid n$ , then  $\delta_n = \begin{cases} \alpha + (-1)^h & \text{if } n \text{ is odd} \\ 2\alpha + (-1)^h & \text{if } n \text{ is even} \end{cases}$ 

- (iv) if  $\alpha \mid (p+3)/2$  and  $p-1 \mid n$ , then n is even and  $\delta_n = 2\alpha + (-1)^h (1-\alpha)$
- (v) if n is odd and  $\alpha \mid (p-1)$ , then h is odd and  $\delta_n = (\alpha + 2)/2$ .

*Proof.* It follows from Remark 5.1 that

$$\gamma(\mathcal{C}_n) = \begin{cases} M-1 & \text{if } n \text{ is odd} \\ M-2 & \text{if } n \text{ is even} \end{cases},$$

where M is the number of solutions  $(x_1, \ldots, x_h) \in [\![0, (p-1)/2]\!]^h$  to (5.5). Let  $t \in \mathbb{N}$  and  $s \in [\![0, \alpha - 1]\!]$  be such that  $(p-1)/2 = \alpha t + s$ . As  $p^i \in (\mathbb{Z}/\alpha\mathbb{Z})^{\times}$  for all  $i \in [\![0, h-1]\!]$ , Lemma 5.3 gives

$$M = \frac{(t\alpha + s + 1)^h - (s + 1)^h}{\alpha} + \delta = \frac{1}{\alpha} \left( \left(\frac{p+1}{2}\right)^h - \left((s+1)^h - \alpha\delta\right) \right)$$

Therefore,  $\gamma(\mathcal{C}_n) = \frac{n}{p^h - 1} \left( \left( \frac{p+1}{2} \right)^h - \delta_n \right)$ . The proofs of assertions (i) — (v) are all similar and simple. We provide a proof for assertion (iii) only. First note that the conditions

$$s\equiv (p-1)/2 \pmod{\alpha}, \, s\in [\![0,\alpha-1]\!], \text{ and } (p+3)/2\equiv 0 \pmod{\alpha}$$

give  $s = \alpha - 2$ , which leads to item (iii) of Lemma 5.3. Bearing (5.3) and (5.5) in mind, note that  $a_i := p^{i-1} \in (\mathbb{Z}/\alpha\mathbb{Z})^{\times}$  for all  $i \in [1, h]$ , and

$$\sum_{i=1}^{h} a_i = \sum_{i=0}^{h-1} p^i = \frac{p^h - 1}{p-1} = \alpha \left(\frac{n}{p-1}\right) \not\equiv 0 \pmod{\alpha},$$

as  $p-1 \nmid n$ . Hence Lemma 5.3 gives  $\delta = ((\alpha - 1)^h - (-1)^h)/\alpha$ , and then

$$\delta_n = (\alpha - 1)^h - \alpha(\delta - \gcd(n, 2)) = \begin{cases} \alpha + (-1)^h & \text{if } n \text{ is odd} \\ 2\alpha + (-1)^h & \text{if } n \text{ is even} \end{cases}.$$

## 6 Explicit formulas for other types of curves

In the prior sections, we considered an array of curves  $y^m = x^n + 1$  and showed how Theorem 1.1 leads to explicit formulas of their *p*-ranks. In this final section, we further discuss how the previous results can be used to obtain the *p*-rank of other types of curves.

As mentioned in the Introduction, in [10, Remark 4], the authors computed the *p*-rank of the Dickson-Guralnick-Zieve curve

$$\mathcal{C}: \frac{(x^{q^3} - x)(y^q - y) - (x^q - x)(y^{q^3} - y)}{(x^{q^2} - x)(y^q - y) - (x^q - x)(y^{q^2} - y)} = 0$$

in terms of the *p*-rank of  $\mathcal{F}_{q-1}$ :  $y^{q-1} = x^{q-1} + 1$ . More precisely, they used the Deuring-Shafarevich formula [34, Theorem 4.2] to show that

$$\gamma(\mathcal{C}) = 2q^4 + (\gamma(\mathcal{F}_{q-1}) - 4)q^3 + q + 1$$

and computed  $\gamma(\mathcal{C})$  when q = p is prime. From [2, Theorem 19], or its generalization in Theorem 4.6, we have  $\gamma(\mathcal{F}_{q-1}) = (((p+1)/2)^h - 3) q + 3$ , where  $q = p^h$ , which finally gives the *p*-rank

$$\gamma(\mathcal{C}) = \left( ((p+1)/2)^h - 1 \right) q^4 - q^3 + q + 1$$

of the Dickson-Guralnick-Zieve curve in the general case. Note that Theorem 4.6 and the subsequent results in Section 4 also enable us to compute the *p*-rank of possible generalizations of C, arising as *p*-power Galois extensions of  $y^{p^m \pm 1} = x^{p^n \pm 1} + 1$ .

For another example, let us consider the curve over  $\mathbb{F}_q$ , where  $q = p^h$  is odd, given by

$$\mathcal{X} : (x+y)^{q+1} - 2(x^q y^q + xy) = 0.$$

The curve  $\mathcal{X}$ , recently presented and investigated in [3], has a significant role in the classification of certain curves with a large automorphism group. Once again, using the Deuring-Shafarevich formula, it can be shown that the p-rank

of  $\mathcal{X}$  is given by  $\gamma(\mathcal{X}) = (\gamma(\mathcal{C}_{2(q-1)}) + 1)q - 1$ , where  $\mathcal{C}_{2(q-1)}$  is the curve  $y^2 = x^{2(q-1)} + 1$  [3, Proposition 8.12]. Now, after Corollary 5.2, we immediately have

$$\gamma(\mathcal{X}) = (((p+1)/2)^h - 1)q - 1.$$

In addition to the Deuring-Shafarevich formula, Lemma 2.4 and Theorem 2.3 clearly provide methods to compute the p-rank of other types of curves, based on the p-rank of  $y^m = x^n + 1$ . This was already explored in Corollaries 3.6, 4.7 and 4.10 and Theorems 3.5 and 4.8. But now the results of Sections 5 and 6 allow us to go beyond. For instance, let us consider the curve

$$\mathcal{D}_n: y^2 = x(x^n + 1)$$

defined over  $\mathbb{F}_q$ , where q is a power of an odd prime p, and  $n \in \mathbb{Z}_{>0}$  is not divisible by p. Note that (ii) in Corollary 3.6 gives

$$\gamma(\mathcal{D}_n) = \gamma(\mathcal{C}_{2n}) - \gamma(\mathcal{C}_n). \tag{6.1}$$

Let us recall the well-known fact that if  $n \mid q+1$ , then  $C_n$  is  $\mathbb{F}_{q^2}$ -maximal, and that  $\mathcal{D}_n$  is  $\mathbb{F}_{q^2}$ -maximal if and only if  $q \equiv -1$  or  $n+1 \pmod{2n}$  [35, Theorem 1]. In both cases, the curve, being supersingular, has zero *p*-rank. The following are immediate consequences of the aforementioned facts.

**Corollary 6.1.** If  $h \in \mathbb{Z}_{>0}$ , then  $\gamma(\mathcal{D}_{p^{h}+1}) = ((p+1)/2)^{h}$ .

*Proof.* From (6.1), we have  $\gamma(\mathcal{D}_{p^h+1}) = \gamma(\mathcal{C}_{2(p^h+1)}) - \gamma(\mathcal{C}_{p^h+1})$ . Since  $\mathcal{C}_{p^h+1}$  is supersingular, item (ii) of Corollary 5.2 gives  $\gamma(\mathcal{D}_{p^h+1}) = ((p+1)/2)^h$ .

**Corollary 6.2.** Let  $h, \alpha \in \mathbb{Z}_{>0}$  be such that  $\alpha$  is an odd divisor of  $p^h - 1$ . Then  $\gamma(\mathcal{C}_{(p^h-1)/\alpha}) = \gamma(\mathcal{C}_{2(p^h-1)/\alpha})$ .

*Proof.* This follows directly from (6.1) and the supersingularity of the curve  $\mathcal{D}_{(p^h-1)/\alpha}$ .

Now note that by Corollary 3.6, if n is odd, then  $\gamma(\mathcal{D}_n) = \gamma(\mathcal{C}_n)$ . For n even, the p-rank of  $\mathcal{D}_n$  is provided in the following result, compiled in Table 5.

**Theorem 6.3.** Let  $n \in \mathbb{Z}_{>0}$  be even and  $\alpha = (p^h - 1)/n$ , where h is the order of p in  $(\mathbb{Z}/n\mathbb{Z})^{\times}$ . If  $\alpha$  is odd, then  $\gamma(\mathcal{D}_n) = 0$ . Otherwise,

$$\gamma(\mathcal{D}_n) = \frac{n}{p^h - 1} \left( \left(\frac{p+1}{2}\right)^h - \tilde{\delta}_n \right), \text{ with } \tilde{\delta}_n = 2\delta_{2n} - \delta_n$$

where  $\delta_n$  is given as in Theorem 5.4. Moreover,

- (i) if  $\alpha \mid (p-1)/2$ , then  $\tilde{\delta}_n = 1$
- (ii) if  $\alpha \mid (p+1)/2$ , then  $\tilde{\delta}_n = 0$
- (iii) if  $\alpha \mid (p+3)/2$ , then

$$\tilde{\delta}_n = \begin{cases} (-1)^h (1-\alpha) & \text{ if } (p-1) \mid 2n \text{ and } (p-1) \nmid n \\ (-1)^h & \text{ otherwise.} \end{cases}$$

*Proof.* If  $\alpha$  is odd, then  $p^h = 2n((\alpha - 1)/2) + n + 1$ , that is,  $\mathcal{D}_n$  is  $\mathbb{F}_{p^{2h}}$ -maximal. Therefore,  $\gamma(\mathcal{D}_n) = 0$ . Suppose that  $\alpha$  is even. As  $(p^h - 1)/(2n) = \alpha/2 \in \mathbb{Z}_{>0}$ , equation (6.1) and Theorem 5.4 yield

$$\gamma(\mathcal{D}_n) = \frac{2n}{p^h - 1} \left( \left(\frac{p+1}{2}\right)^h - \delta_{2n} \right) - \frac{n}{p^h - 1} \left( \left(\frac{p+1}{2}\right)^h - \delta_n \right)$$
$$= \frac{n}{p^h - 1} \left( \left(\frac{p+1}{2}\right)^h - \tilde{\delta}_n \right).$$

The proofs of the assertions (i) — (iii) are quite similar. We provide the proof of (iii) only. Since  $(p^h - 1)/(2n) = \alpha/2$ , it follows from items (iii) and (iv) of Theorem 5.4 that

$$\delta_{2n} = \begin{cases} \alpha + (-1)^h & \text{if } (p-1) \nmid 2n \\ \alpha + (-1)^h (2-\alpha)/2 & \text{otherwise} \end{cases} \text{ and } \delta_n = \begin{cases} 2\alpha + (-1)^h (1-\alpha) & \text{if } (p-1) \mid n \\ 2\alpha + (-1)^h & \text{otherwise.} \end{cases}$$

Therefore,

$$\tilde{\delta}_n = 2\delta_{2n} - \delta_n = \begin{cases} (-1)^h (1-\alpha) & \text{if } (p-1) \mid 2n \text{ and } (p-1) \nmid n \\ (-1)^h & \text{otherwise.} \end{cases}$$

# 7 A final remark

In Sections 4 and 5, we used a few techniques to show how Theorem 3.1 yields explicit formulas for the *p*-rank of Fermat-type curves, and in Section 6, we demonstrated how the results can be used to arrive at the *p*-rank of other curves. These sections bring a number of new results to the literature, but we should note that they are merely illustrative in the sense that, using Theorem 3.1, there is still room to attain *p*-rank formulas for several other curves. For example, with some of the techniques of Section 4, it can be proved that if *m* and *n* are divisors of p - 1, then the curve  $y^{m(p+1)} = x^{n(p+1)} + 1$  has *p*-rank

$$\frac{1}{4}\left(mnp^{2}+2\left(2mn-m-n-d\right)p+3mn-2m-2n-2d+4\right)-\frac{1}{12mn}\left(m^{2}+n^{2}+d^{2}\right)\left(p-1\right)^{2},$$

where  $d = \gcd(m, n)$ .

# 8 Acknowledgements

We would like to thank Rachel Pries for her comments on a previous version of this manuscript, which enabled us to enhance our text. The first author was supported by CNPq (Brazil), grant 311572/2019-7.

### A The cardinality of a few sets

This appendix provides the cardinality of a few simple sets that appear throughout the proofs of our results in Section 4.

Lemma A.1. If p is a prime and

$$\mathcal{L} = \left\{ (a, b, c, d) \middle| \begin{array}{l} a, b \in \llbracket 1, p \rrbracket, \\ 1 \le c, d \le \min \{a, b\}, \\ a + b \le c + d + p - 1 \end{array} \right\},$$

then  $\#\mathcal{L} = p(p+1)(p^2 + p + 2)/8.$ 

*Proof.* For p = 2, it is straightforward to check that  $\#\mathcal{L} = 6$ . Thus we suppose  $p \ge 3$ , and proceed by fixing  $(a,b) \in [\![1,p]\!]^2$ , and then counting the pairs (c,d) satisfying the conditions given in set  $\mathcal{L}$ .

Suppose a < b. Setting x := a - c and y := a - d, it suffices to count the pairs  $(x, y) \in [[0, a - 1]]^2$  such that  $x + y \le p - 1 - (b - a)$ . Note that (a, b) is such that the square  $[[0, a - 1]]^2$  is below the line x + y = p - 1 - (b - a) if and only if  $a + b \le p + 1$ . Thus such points give rise to

$$\sum_{b=2}^{p} \sum_{a=1}^{\min\{b-1,p+1-b\}} a^2 = (p^4 + 4p^3 + 2p^2 - 4p - 3)/96$$

elements of  $\mathcal{L}$ . In the cases  $a + b \ge p + 2$ , we need to remove the (a + b - p - 1)(a + b - p)/2 points above the line x + y = (p - 1) - (b - a). That is, the points (a, b) for which  $a + b \ge p + 2$  give rise to

$$\sum_{b=\frac{p+3}{2}}^{p} \sum_{a=p+2-b}^{b-1} (a^2 - (a+b-p-1)(a+b-p)/2) = (5p^4 - 4p^3 - 2p^2 + 4p - 3)/96$$

elements of  $\mathcal{L}$ . Adding the quantities above gives us the total of  $(p^4 - 1)/16$  elements  $(a, b, c, d) \in \mathcal{L}$  for which a < b. Clearly, by symmetry, the same holds when a > b. Hence  $(p^4 - 1)/8$  is the number of elements  $(a, b, c, d) \in \mathcal{L}$  for which  $a \neq b$ .

For the cases a = b, considering the line x + y = p - 1, the same procedure gives a total of  $(2p^3 + 3p^2 + 2p + 1)/8$  elements  $(a, b, c, d) \in \mathcal{L}$  for which a = b. Therefore,

$$#\mathcal{L} = (p^4 - 1)/8 + (2p^3 + 3p^2 + 2p + 1)/8 = p(p+1)(p^2 + p + 2)/8.$$

**Lemma A.2.** If  $b, m, n \in \mathbb{Z}_{>0}$  and

$$\mathcal{L} = \left\{ (a_0, \dots, a_{m-1}, b_0, \dots, b_{n-1}) \mid \begin{array}{l} 0 \le a_i \le b - 1\\ 0 \le b_j \le \min\{a_1, \dots, a_{m-1}\} \end{array} \right\},$$
  
then  $\#\mathcal{L} = \sum_{i=0}^{b-1} \left( (i+1)^m - i^m \right) (b-i)^n.$ 

*Proof.* Note that for each  $t \in \{0, \ldots, b-1\}$ , the number of *m*-tuples  $(a_0, \ldots, a_{m-1})$  for which  $\min\{a_0, \ldots, a_{m-1}\} = t$  is  $(b-t)^m - (b-(t+1))^m$ . Since for each such *m*-tuple, we have  $(t+1)^n$  possible *n*-tuples  $(b_0, \ldots, b_{n-1})$ , it follows that

$$#\mathcal{L} = \sum_{t=0}^{b-1} \left( \left( b - t \right)^m - \left( b - t - 1 \right)^m \right) (t+1)^n.$$

Defining i := b - t - 1, the result follows.

**Lemma A.3.** Let  $b, m, n \in \mathbb{Z}_{>0}$ . If  $b \ge 2$  and

$$\mathcal{L} = \left\{ (a_0, \dots, a_{m-1}, b_0, \dots, b_{n-1}) \mid \begin{array}{l} 0 \le a_r \le b - 1, \\ 0 \le b_s \le \min_r \{a_r, b - 1 - a_r\} \end{array} \right\},\$$

then

$$\#\mathcal{L} = \begin{cases} \frac{1}{2^n} \Big( (b+1)^n + \sum_{i=1}^{(b-1)/2} \big( (2i+1)^m - (2i-1)^m \big) (b-2i+1)^n \Big) & \text{if } b \text{ is odd} \\ \\ 2^{m-n} \sum_{i=1}^{b/2} \big( i^m - (i-1)^m \big) (b-2i+2)^n & \text{if } b \text{ is even.} \end{cases}$$

*Proof.* Direct inspection shows that for each  $t \in \{0, ..., b-1\}$ , the number of *m*-tuples  $(a_0, ..., a_{m-1})$  for which  $\min_{x} \{a_r, b - a_r - 1\} = t$  is

$$\begin{cases} (b-2t)^m - (b-2t-2)^m & \text{if } t < (b-1)/2 \\ 1 & \text{if } t = (b-1)/2 \\ 0 & \text{if } t > (b-1)/2. \end{cases}$$

For each such *m*-tuple, we have  $(t+1)^n$  possible *n*-tuples  $(b_0, \ldots, b_{n-1})$ . Since t = (b-1)/2 occurs if and only if *b* is odd, it follows that

$$\#\mathcal{L} = \begin{cases} \left( (b+1)/2 \right)^n + \sum_{t=0}^{(b-3)/2} \left( (b-2t)^m - (b-2t-2)^m \right) (t+1)^n & \text{if } b \text{ is odd} \\ \\ \sum_{t=0}^{(b-2)/2} \left( (b-2t)^m - (b-2t-2)^m \right) (t+1)^n & \text{if } b \text{ is even.} \end{cases}$$

Defining i := (b - 2t - 1)/2 when b is odd, and i := (b - 2t)/2 when b is even, the result follows.

**Lemma A.4.** If 
$$b, m, n \in \mathbb{Z}_{>0}$$
 and

$$\mathcal{L} = \left\{ (a_0, \dots, a_{m-1}, b_0, \dots, b_{2n-1}) \middle| \begin{array}{l} 0 \le a_r \le b - 1\\ 0 \le b_s \le \min\{a_r : r \in [\![0, m-1]\!]\}, \ s \le n - 1,\\ 0 \le b_s \le \min\{b - a_r - 1 : r \in [\![0, m-1]\!]\}, \ s \ge n \end{array} \right\},$$

then

$$#\mathcal{L} = \sum_{i=0}^{b-1} \left( (b-i)(i+1) \right)^n + \sum_{1 \le j \le i \le b-1} \left( (j+1)^m - 2j^m + (j-1)^m \right) \left( (b-i)(i-j+1) \right)^n.$$

*Proof.* Note that for each  $i \in [0, b-1]$  and  $t \in [0, i]$ , the number of *m*-tuples  $(a_0, \ldots, a_{m-1})$  for which  $\max\{a_0, \ldots, a_{m-1}\} = i$  and  $\min\{a_0, \ldots, a_{m-1}\} = t$  is 1 when t = i, and  $(i - t + 1)^m - 2(i - t)^m + (i - t - 1)^m$  when  $t \neq i$ . Since for each such *m*-tuple, we have  $(b - i)^n (t + 1)^n$  possible 2*n*-tuples  $(b_0, \ldots, b_{2n-1})$ , it follows that

$$#\mathcal{L} = \sum_{i=0}^{b-1} (b-i)^n (i+1)^n + \sum_{0 \le t < i \le b-1} \left( (i-t+1)^m - 2(i-t)^m + (i-t-1)^m \right) (b-i)^n (t+1)^n.$$

Defining j := i - t, the result follows.

### References

- [1] Achter, J.D., Pries, R., 2008. Monodromy of the *p*-rank strata of the moduli space of curves. Int. Math. Res. Not. IMRN 2008, 1–25.
- Bassa, A., Beelen, P., 2010. The Hasse-Witt invariant in some towers of function fields over finite fields. Bull. Braz. Math. Soc. (N.S.) 41, 567–582.
- [3] Borges, H., Korchmáros, G., Speziali, P., 2022. Plane curves with a large linear automorphism group in characteristic *p*. Preprint, arXiv:2202.05765.
- [4] Coulter, R.S., 1998. Explicit evaluations of some Weil sums. Acta Arith. 83, 241-251.
- [5] Edixhoven, B., van der Geer, G., Moonen, B., 2022. Abelian varieties. URL: http://van-der-geer.nl/~gerard/AV.pdf. access date: 2022-03-17 14:26:46.
- [6] Fine, N.J., 1947. Binomial coefficients modulo a prime. Amer. Math. Monthly 54, 589–592.
- [7] Friedlander, H., Garton, D., Malmskog, B., Pries, R., Weir, C., 2013. The *a*-number of jacobian of Suzuki curves. Proc. Amer. Math. Soc. 141, 3019–3028.
- [8] Garcia, A., Tafazolian, S., 2008. Certain maximal curves and Cartier operators. Acta Arith 135, 199–218.
- [9] Giulietti, M., Korchmáros, G., 2019. Algebraic curves with many automorphisms. Adv. Math 349, 162–211.
- [10] Giulietti, M., Korchmáros, G., Timpanella, M., 2019. On the Dickson-Guralnick-Zieve curve. J. Number Theory 196, 114–138.
- [11] Glass, D., Pries, R., 2005. Hyperelliptic curves with prescribed *p*-torsion. Manuscripta Math. 117, 299–317.
- [12] Gonzalez, J., 1997. Hasse-Witt matrices for the Fermat curves of prime degree. Tohoku Math. J. 49, 149–163.
- [13] Gunby, G., Smith, A., Yuan, A., 2015. Irreducible canonical representations in positive characteristic. Res. Number Theory 1, 1–25.
- [14] Hasse, H., Witt, E., 1936. Zyklische unverzweigte erweiterungskörper vom primzahlgrade p über einen algebraischen funktionenkörper der characteristik *p*. Monatsh. Math. Phys. 43, 477–492.
- [15] Hirschfeld, J., Korchmáros, G., Torres, F., 2008. Algebraic Curves Over a Finite Field. Princeton Univ. Press, Princeton and Oxford.
- [16] Kani, E., Rosen, M., 1989. Idempotent relations and factors of jacobians. Math. Ann. 284, 307–327.
- [17] Kodama, T., Washio, T., 1986. Hasse-Witt matrices of hyperelliptic function fields. Sci. Bull. Fac. Educ. Nagasaki Univ. 37, 9–15.
- [18] Kodama, T., Washio, T., 1987. Hasse-Witt matrices of Fermat curves. Manuscripta Math. 60, 185–195.
- [19] Kontogeorgis, A., Rotger, V., 2008. On abelian automorphism groups of Mumford curves. Bull. Lond. Math. Soc. 40, 353–362.
- [20] Kontogeorgis, A.I., 1998. The group of automorphisms of the function field of the curve  $x^n + y^m + 1 = 0$ . J. Number Theory 72, 110–136.
- [21] Korchmáros, G., Montanucci, M., 2019. Ordinary algebraic curves with many automorphisms in positive characteristic. Algebra Number Theory 13, 1–18.
- [22] Lucas, E., 1878. Sur les congruences des nombers eulériens et des coefficients différentiels des fonctions trigonométriques, suivant un module premier. Bull. Soc. Math. France 6, 49–54.
- [23] McEliece, R.J., 1987. Finite Fields for Computer Scientists and Engeneers. Springer New York, NY.
- [24] Miller, H., 1972. Curves with invertible Hasse-Witt-matrix. Math. Ann. 197, 123–127.
- [25] Montanucci, M., Speziali, P., 2018. The *a*-numbers of Fermat and Hurwitz curves. J. Pure Appl. Algebra 222, 477–488.

- [26] Mumford, D., 1970. Abelian Varieties. Oxford Univ. Press.
- [27] Nakajima, S., 1987. *p*-ranks and automorphism groups of algebraic curves. Trans. Amer. Math. Soc. 303, 595–607.
- [28] Oliveira, J.A., 2021. On maximal and minimal hypersurfaces of Fermat type. Preprint, arXiv:2110.07452.
- [29] Oort, F., 1974. Subvarieties of moduli spaces. Invent. Math. 24, 95–119.
- [30] Pries, R., Ulmer, D., 2021. On BT<sub>1</sub> group schemes and Fermat curves. New York J. Math. 27, 705–739.
- [31] Serre, J.P., 1958. Sur la topologie des variéts algébriques en caractéristique *p*, Univ. Nac. Aut. México and Unesco. Symposium Internacional de Topología Algebraica. pp. 24–53.
- [32] Silverman, J.H., 1986. The Arithmetic of Elliptic Curves. Springer New York, NY.
- [33] Stöhr, K.O., Viana, P., 1989. A study of Hasse-Witt matrices by local methods. Math. Z. 200, 397-408.
- [34] Subrao, D., 1975. The p-rank of Artin-Schreier curves. manuscripta math. 16, 169–193.
- [35] Tafazolian, S., 2012. A note on certain maximal hyperelliptic curves. Finite Fields Appl. 18, 1013–1016.
- [36] Tafazolian, S., Torres, F., 2013. On maximal curves of Fermat type. Adv. Geom. 13, 613–617.
- [37] Towse, C., 1996. Weierstrass points on cyclic covers of the projective line. Trans. Amer. Math. Soc. 348, 3355– 3378.
- [38] Yui, N., 1978. On the jacobian varieties of hyperelliptic curves over fields of characteristic p > 2. J. Algebra 52, 378–410.
- [39] Yui, N., 1980. On the jacobian variety of the Fermat curve. J. Algebra 65, 1–35.
- [40] Zhu, H.J., 2006. Hyperelliptic curves over  $\mathbb{F}_2$  of every 2-rank without extra automorphisms. Proc. Amer. Math. Soc. 134, 323–331.