UNIVERSALITY OF THE LOCAL LIMIT OF PREFERENTIAL ATTACHMENT MODELS

BY ALESSANDRO GARAVAGLIA^{2,b}, RAJAT SUBHRA HAZRA^{1,a}, REMCO VAN DER HOFSTAD^{2,c} and Rounak Ray^{2,d}

¹University of Leiden, Niels Bohrweg 1, 2333 CA, Leiden, The Netherlands, ^ar.s.hazra@math.leidenuniv.nl

²Department of Mathematics and Computer Science, Eindhoven University of Technology, 5600 MB Eindhoven, The Netherlands, ^bale.garavaglia@gmail.com; ^cr.w.v.d.hofstad@TUE.nl; ^dr.ray@tue.nl

> We study preferential attachment models where vertices enter the network with i.i.d. random numbers of edges that we call the *out-degree*. We identify local limit of such models, substantially extending the work of Berger et al. [8]. The degree distribution of this limiting random graph, which we call the *random Pólya point tree*, has a surprising size-biasing phenomenon.

> Many of the existing preferential attachment models can be viewed as special cases of our preferential attachment model with i.i.d. out-degrees. Additionally, our models incorporates negative values of the preferential attachment fitness parameter, which allows us to consider preferential attachment models with infinite-variance degrees.

> Our proof of local convergence consists of two main steps: a Pólya urn description of our graphs, and an explicit identification of the neighbourhoods in them. We provide a novel and explicit proof to establish a coupling between the preferential attachment model and the Pólya urn graph. Our result proves a density convergence result, for fixed ages of vertices in the local limit.

1. Introduction.

1.1. *Real-world networks and preferential attachment models*. Empirical studies on reallife networks reveal that most of these networks (a) grow with time; (b) are small worlds, meaning that typical distances in the network are small; and (c) have power-law degree sequences. The Barabási-Albert model of [1, 4] is the most popular random graph model for such real-life networks due to the fact that, through a simple dynamic, its properties resemble real-world networks. This model has been generalized in many different ways, creating a wide variety of *preferential attachment models* (PAMs).

By PAMs we denote a class of random graphs with a common dynamics: At every time step, a new vertex appears in the graph, and it connects to $m \ge 1$ existing vertices, with probability proportional to a function of the degrees of the vertices. In other words, when vertex $n \in \mathbb{N}$ appears, it connects to vertex $i \le n$ with probability

(1.1)
$$\mathbb{P}(n \rightsquigarrow i \mid \mathrm{PA}_{n-1}) \propto f(d_i(n-1)),$$

where PA_n is the preferential attachment graph with *n* vertices, $d_i(n-1)$ denotes the degree of vertex *i* in the graph PA_{n-1} , and *f* is some preferential attachment function. We thus in fact consider a whole PA class of random graphs since every function *f* defines a different model. The original Barabási-Albert model is retrieved with f(k) = k. The literature often considers the so-called *affine* PAM, where $f(k) = k + \delta$, for some constant $\delta > -m$.

The constant δ allows for flexibility in the graph structure. In fact, the power-law exponent of the degree distribution is given by $\tau = 3 + \delta/m$ [13, 21, 39], and in general, for $m \ge 2$,

MSC2020 subject classifications: Primary 05C80; secondary 05C82.

Keywords and phrases: local weak convergence, preferential attachment model, Pólya urn.

the typical distance and the diameter are of order $\log \log n$ when $\tau \in (2,3)$, while they are of order $\log n$ when $\tau > 3$. When $\tau = 3$, distances and diameter are of order $\log n / \log \log n$ instead [12, 15, 20, 25].

In PAMs, the degree of a vertex increases over time and higher degree vertices are prone to attract the edges incident to new vertices, increasing their degrees even further. In literature, this is sometimes referred to as *rich-get-richer* effect. The models where the vertex degrees are determined by a *weight* associated to it are sometimes called *rich-by-birth* models.

In [18], the authors have considered a model incorporating both of these effects. The model is a PAM with random out-degrees, i.e., every vertex joins the existing network with a random number of edges that it connects to the existing vertices preferentially to their degrees at that time, as done in usual preferential attachment models. Here, the authors have shown that this system also shows the power-law degree distribution. Jordan [33] considered a special case of this particular model to analyze their results on degree distribution of the random graph. Cooper and Frieze [17] have shown that a further generalised version of this PAM also has a power-law degree distribution. Since most real-life networks are dynamic in nature and have power-law degrees, the PAM is often used to model them. However, in such networks it is never the case that every vertex joins with exactly the same out-degree, and instead i.i.d. out-degrees are more realistic. Gao and van der Vaart [28] have shown that the maximum likelihood estimator of the fitness parameter δ of PAM with random out-degrees is asymptotically normal and has asymptotically minimal variance. There have been further generalisations by allowing younger vertices to have higher degrees in PAMs through a random *fitness* parameter. More precisely, individual factors can be assigned to vertices, obtaining that the probabilities in (1.1) are proportional to $\eta_i f(d_i(n-1))$, thus obtaining PAMs with multiplicative fitness, or proportional to $d_i(n-1) + \eta_i$, giving rise to additive fitness [9, 10, 14, 19, 24].

Similar to many often other random graph models, such as the configuration model, PAMs are called *locally treelike graphs*, meaning that the neighbourhood of the majority of vertices is structured as a tree (up to a certain distance). This idea can be formalized using the notion of local convergence (i.e., the Benjamini-Schramm limit), introduced in [2, 6]. Local convergence turns out to be an extremely versatile tool to understand the geometry of the graph. The basic idea is to explore the neighbourhood of a uniformly chosen vertex of the graph up to a finite distance, to understand its distributional and geometric properties. We refer to, e.g., [40, Chapter 2] for an overview of the theory and applications of the above concept.

Berger et al. initiated the study of local convergence of PAMs in [8]. They showed that the finite neighbourhood of the graph converges to the corresponding neighbourhood of the Pólya point tree (see the description of the Pólya point tree in Section 1.4). The proof uses a Pólya urn representation introduced in [7] to study the spread of viral infections on networks.

Main results and innovation of this paper. The main aim of this article is to extend the local convergence proof in [8] to a more general class of PAMs (including random out-degrees and related dynamics). We achieve this by explicitly computing the density of neighbourhoods of the PAMs. This helps us to extend the result by [8] to models where one can accommodate negative fitness parameters and random out-degrees. The limiting random tree is an extension of the Pólya point tree described in [8], which we call the *random Pólya point tree*.

The randomness of the out-degrees provides a surprising size-biasing effect in the limiting random tree. We show that there is a universal description of the limit by considering many possible affine variants of the PAMs. Additionally, we study the *vertex-marked local convergence in probability* of the PAMs, which is an extension to the local convergence shown in [8]. Here, the marks denote the *ages* of the vertices in the tree, and we prove convergence of the joint *densities* of these ages.

3

In the next section we provide details of the various preferential attachment models considered in this article. We also provide a formal definition of the vertex-marked local limit and the formulation of our main results.

1.2. The models. Several versions of preferential attachment models are available in the literature. We generalize these definitions to the case of random initial number of edges that connect to the already available graph. We refer to these edge numbers as *out-degrees*, even though we consider our model to be *undirected*. Let $(M, m_i)_{i\geq 3}$ be a sequence of i.i.d. N-valued random variables with finite *p*-th moment for some p > 1, and $\delta > -\inf \text{supp}(M)$ be a fixed real number, where supp(M) denotes the support of the random variable M. In our models, every new vertex v joins the graph with m_v many edges incident to it. In the classical preferential attachment models, instead, every new vertex comes with a *fixed* number of edges incident to it. Thus, we can consider the existing models as a degenerate case of ours. Define $m_1 = m_2 = 1$, $\mathbf{m} = (m_1, m_2, m_3, ...)$ and

$$m_{[l]} = \sum_{i \le l} m_i.$$

To describe the edge-connection probabilities and to simplify our calculations, we frequently work with the conditional law given $m = (m_i)_{i \ge 1}$. The conditional measure is denoted by \mathbb{P}_m , i.e., for any event \mathcal{E} ,

(1.2)
$$\mathbb{P}_m(\mathcal{E}) = \mathbb{P}\left(\mathcal{E} \mid \boldsymbol{m}\right).$$

Conditionally on m, we define some versions of the preferential attachment models, special cases of which are equivalent to the models (a-g) described in [29]. We consider the initial graph to be G_0 with 2 vertices with degree a_1 and a_2 , respectively. Throughout the remainder of this paper, we fix $\delta > -\inf \operatorname{supp}(M)$.

Model (A). This is the generalized version of [39, Model (a)]. Conditionally on m, for $v \in \mathbb{N}$ and $j = 1, \ldots, m_v$, the attachment probabilities are given by

(1.3)
$$\mathbb{P}_m\left(v \stackrel{j}{\rightsquigarrow} u \middle| \operatorname{PA}_{v,j-1}^{(A)}(\boldsymbol{m}, \delta)\right) = \begin{cases} \frac{d_u(v, j-1) + \delta}{c_{v,j}} & \text{for } v > u, \\ \frac{d_u(v, j-1) + 1 + \frac{j}{m_u} \delta}{c_{v,j}} & \text{for } v = u, \end{cases}$$

where $v \xrightarrow{j} u$ denotes that vertex v connects to u with its j-th edge, $\operatorname{PA}_{v,j}^{(A)}(\boldsymbol{m}, \delta)$ denotes the graph with v vertices, with the v-th vertex having j out-edges, and $d_u(v, j)$ denotes the degree of vertex u in $\operatorname{PA}_{v,j}^{(A)}(\boldsymbol{m}, \delta)$. We identify $\operatorname{PA}_{v+1,0}^{(A)}(\boldsymbol{m}, \delta)$ with $\operatorname{PA}_{v,m_v}^{(A)}(\boldsymbol{m}, \delta)$. The normalizing constant $c_{v,j}$ in (1.3) equals

(1.4)
$$c_{v,j} := a_{[2]} + 2\left(m_{[v-1]} + j - 2\right) - 1 + (v-1)\delta + \frac{j}{m_v}\delta,$$

where $a_{[2]} = a_1 + a_2$. We denote the above model by $PA_v^{(A)}(\boldsymbol{m}, \delta)$. If we consider M as a degenerate distribution that is equal to m a.s., then $PA_v^{(A)}(\boldsymbol{m}, \delta)$ is essentially equivalent to $PA_v^{(m,\delta)}(a)$ defined in [40]. $PA_v^{(m,0)}(a)$ was informally introduced by [4] and first studied rigorously in [12]. Later the model for general δ was described in [1].

Model (B). This is the generalized version of [39, Model (b)]. Conditionally on m, the attachment probabilities are given by

(1.5)
$$\mathbb{P}_m\left(v \stackrel{j}{\rightsquigarrow} u \middle| \operatorname{PA}_{v,j-1}^{(\mathrm{B})}(\boldsymbol{m}, \delta)\right) = \begin{cases} \frac{d_u(v,j-1)+\delta}{c_{v,j}} & \text{for } v > u, \\ \frac{d_u(v,j-1)+\frac{(j-1)}{m_u}\delta}{c_{v,j}} & \text{for } v = u, \end{cases}$$

where again $\operatorname{PA}_{v,j}^{(B)}(\boldsymbol{m}, \delta)$ denotes the graph with v vertices, with the v-th vertex having j out-edges, and $d_u(v, j)$ denotes the degree of vertex u in $\operatorname{PA}_{v,j}^{(B)}(\boldsymbol{m}, \delta)$. We identify $\operatorname{PA}_{v+1,0}^{(B)}(\boldsymbol{m}, \delta)$ with $\operatorname{PA}_{v,m_v}^{(B)}(\boldsymbol{m}, \delta)$. The normalizing constant $c_{v,j}$ in (1.5) now equals

$$c_{v,j} = a_{[2]} + 2\left(m_{[v-1]} + j - 3\right) + (v-1)\delta + \frac{(j-1)}{m_v}\delta.$$

We denote the above model by $\mathrm{PA}_{v}^{\scriptscriptstyle(\mathrm{B})}(\boldsymbol{m},\delta)$. For M a degenerate distribution which is equal to m a.s., we obtain $\mathrm{PA}_{v}^{\scriptscriptstyle(m,\delta)}(b)$ described in [40] from $\mathrm{PA}_{v}^{\scriptscriptstyle(\mathrm{B})}(\boldsymbol{m},\delta)$.

REMARK 1.1 (Difference between Models (A) and (B)). From the definition of the models (A) and (B), the models are different in that the first edge from every new vertex can create a self-loop in model (A) but not in model (B). Note that the edge probabilities are different for all j.

Model (D). This model is the generalized version of the sequential model described in [8]. We start with a fixed graph G_0 of size 2 and degrees a_1 and a_2 , respectively. Denote the graph by $PA_v^{(D)}(\boldsymbol{m}, \delta)$ when the graph has v vertices.

For $v \ge 3$, vertex v enters the system with m_v many out-edges whose edge-connection probabilities are given by

(1.6)
$$\mathbb{P}_m\left(v \stackrel{j}{\rightsquigarrow} u \middle| \operatorname{PA}_{v,j-1}^{\scriptscriptstyle(\mathrm{D})}\right) = \frac{d_u(v,j-1) + \delta}{c_{v,j}} \quad \text{for} \quad v > u,$$

where $\operatorname{PA}_{v,j}^{(D)}(\boldsymbol{m}, \delta)$ denotes the graph with v vertices, with the v-th vertex having j out-edges, and $d_u(v, j)$ is the degree of the vertex u in the graph $\operatorname{PA}_{v,j}^{(D)}$, and

$$c_{v,j} = a_{[2]} + 2(m_{[v-1]} - 2) + (j-1) + (v-1)\delta.$$

Again we identify $\operatorname{PA}_{v+1,0}^{(D)}(\boldsymbol{m},\delta)$ with $\operatorname{PA}_{v,m_v}^{(D)}(\boldsymbol{m},\delta)$. For M a degenerate random variable that is equal to m a.s., we obtain $\operatorname{PA}_v^{(m,\delta)}(d)$, as described in [40].

Model (E). This model is the independent model proposed by Berger et al. in [18]. We start with the same initial graph G_0 . We name the graph $\operatorname{PA}_v^{(\mathrm{E})}(\boldsymbol{m}, \delta)$ when there are v vertices in the graph. Conditionally on \boldsymbol{m} , every new vertex v is added to the graph with m_v many outedges. Every out-edge $j \in [m_v]$ from v connects to one of the vertex $u \in [v-1]$ independently with the probabilities

(1.7)
$$\mathbb{P}_{m}\left(v \stackrel{j}{\rightsquigarrow} u \middle| \mathrm{PA}_{v-1}^{(\mathrm{E})}(\boldsymbol{m}, \delta)\right) = \frac{d_{u}(v-1) + \delta}{a_{[2]} + 2(m_{[v-1]} - 2) + (v-1)\delta},$$

where $d_u(v-1)$ is the degree of the vertex u in $PA_{v-1}^{(E)}$. Note that the intermediate degree updates are not there in this graph. Similarly as in model (D), no self-loops are allowed here.

5

Model (F). The independent model does not allow for self-loops, but multiple edges between two vertices may still occur. Such multigraphs are, in many applications, unrealistic, and we next propose a model where the graphs obtained are *simple*, i.e., without self-loops and multiple edges. Model (F) is a variation of the independent model where the new vertices connect to the existing vertices in the graph independently but *without replacement* and the edge-connection probabilities are similar to (1.7). Indeed, for $j \ge 2$, the normalization factor changes a little due to the fact that vertices that have already appeared as neighbours are now forbidden. The degenerate case of our PAM, i.e., with fixed out-degree, is studied in [5] for analyzing the largest connected component in the strong friendship subgraph of evolving online social networks.

REMARK 1.2 (The missing model (c)). There is a description of model (c) in [39, Section 8.2], which reduces to model (a), so we refrain from discussing it further in this article. \blacksquare

1.3. *The space of rooted vertex-marked graphs and marked local convergence*. Local convergence of rooted graphs was first introduced by Benjamini and Schramm in [6] and Aldous and Steele in [2]. We now give a brief introduction.

A graph G = (V(G), E(G)) (possibly infinite) is called *locally finite* if every vertex $v \in V(G)$ has finite degree (not necessarily uniformly bounded). A pair (G, o) is called a *rooted* graph, where G is rooted at $o \in V(G)$. For any two vertices $u, v \in V(G)$, $d_G(u, v)$ is defined as the length of the shortest path from u to v in G, i.e., the minimal number of edges needed to connect u and v.

For a rooted graph (G, o), the *r*-neighbourhood of the root o, denoted by $B_r^{(G)}(o)$, is defined as the graph rooted at o with vertex and edge sets given by

$$V(B_r^{(G)}(o)) = \{ v \in V(G) : d_G(o, v) \le r \}, \text{ and }$$

$$E(B_r^{(G)}(o)) = \{\{u, v\} \in E(G) \colon u, v \in V(B_r^{(G)}(o))\}$$

Let (G_1, o_1) and (G_2, o_2) be two rooted locally finite graphs with $G_1 = (V(G_1), E(G_1))$ and $G_2 = (V(G_2), E(G_2))$. Then we say that (G_1, o_1) is *rooted isomorphic* to (G_2, o_2) , which we denote as $(G_1, o_1) \simeq (G_2, o_2)$, when there exists a graph isomorphism between G_1 and G_2 that maps o_1 to o_2 , i.e., when there exists a bijection $\phi \colon V(G_1) \mapsto V(G_2)$ such that

(1.9)
$$\phi(o_1) = o_2$$
, and $\{u, v\} \in E(G_1) \iff \{\phi(u), \phi(v)\} \in E(G_2)$.

Marks are generally images of an injective function \mathcal{M} acting on the vertices of a graph G, as well as the edges. These marks take values in a complete separable metric space (Ξ, d_{Ξ}) . Rooted graphs with only vertices having marks are called *vertex-marked* rooted graphs, and are denoted by $(G, o, \mathcal{M}(G)) = (V(G), E(G), o, \mathcal{M}(V(G)))$. We say that two graphs $(G_1, o_1, \mathcal{M}_1(G_1))$ and $(G_2, o_2, \mathcal{M}_2(G_2))$ are vertex-marked-isomorphic, which we denote as $(G_1, o_1, \mathcal{M}_1(G_1)) \stackrel{\star}{\simeq} (G_2, o_2, \mathcal{M}_2(G_2))$, when there exists a bijective function $\phi: V(G_1) \mapsto V(G_2)$ such that

 $\triangleright \phi(o_1) = o_2;$

(1.8)

$$\triangleright \quad (v_1, v_2) \in E(G_1) \text{ implies that } (\phi(v_1), \phi(v_2)) \in E(G_2);$$

 $\triangleright \mathcal{M}_1(v) = \mathcal{M}_2(\phi(v)) \text{ for all } v \in V(G_1).$

We let \mathcal{G}_{\star} be the vertex-marked isomorphism invariant class of rooted graphs. Similarly as [40, Definition 2.11] that one can define a metric $d_{\mathcal{G}_{\star}}$ as

(1.10)
$$d_{\mathcal{G}_{\star}}\left((G_1, o_1, \mathcal{M}_1(G_1)), (G_2, o_2, \mathcal{M}_2(G_2))\right) = \frac{1}{1 + R^{\star}},$$

where

$$R^{\star} = \sup\{r \ge 0 : B_r^{(G_1)}(o_1) \simeq B_r^{(G_2)}(o_2), \text{ and there exists a bijective function } \phi \text{ from} \}$$

(1.11)
$$V(G_1)$$
 to $V(G_2)$ such that $d_{\Xi}(\mathcal{M}_1(u), \mathcal{M}_2(\phi(u))) \le \frac{1}{r}, \forall u \in V(B_r^{(G_1)}(o_1))\},$

which makes $(\mathcal{G}_{\star}, d_{\mathcal{G}_{\star}})$ a Polish space. We next define the notion of marked local convergence of vertex-marked random graphs on this space. [40, Theorem 2.14] describes various notions of local convergence. For the next two definitions of local convergence, we consider that $\{G_n\}_{n\geq 1}$ is a sequence of (possibly random) vertex-marked graphs with $G_n = (V(G_n), E(G_n), \mathcal{M}_n(V(G_n)))$ that are finite (almost surely). Conditionally on G_n , let o_n be a randomly chosen vertex from $V(G_n)$. Note that $\{(G_n, o_n)\}_{n\geq 1}$ is a sequence of random variables defined on \mathcal{G}_{\star} . Then vertex-marked local convergence is defined as follows:

DEFINITION 1.3 (Vertex marked local convergence).

(a) (Vertex-marked local weak convergence) The sequence $\{(G_n, o_n)\}_{n \ge 1}$ is said to converge vertex-marked locally weakly to a random element $(G, o, \mathcal{M}(V(G))) \in \mathcal{G}_*$ having probability law μ_* , if, for every r > 0, and every $(H_*, \mathcal{M}_{H_*}(H_*)) \in \mathcal{G}_*$, as $n \to \infty$,

$$\mathbb{P}\left(B_{r}^{(G_{n})}(o_{n}) \simeq H_{\star}, \ d_{\mathcal{G}_{\star}}\left((B_{r}^{(G_{n})}(o_{n}), o_{n}, \mathcal{M}(V(B_{r}^{(G_{n})}(o_{n}))), (H_{\star}, \mathcal{M}_{H_{\star}}(H_{\star}))\right) \leq \frac{1}{r}\right)$$
(1.12)
$$\to \mu_{\star}\left(B_{r}^{(G)}(o) \simeq H_{\star}, \ d_{\mathcal{G}_{\star}}\left((B_{r}^{(G)}(o), o, \mathcal{M}(V(B_{r}^{(G)}(o))), (H_{\star}, \mathcal{M}_{H_{\star}}(H_{\star})))\right) \leq \frac{1}{r}\right)$$

(b) (Vertex-marked local convergence) The sequence $\{G_n\}_{n\geq 1}$ is said to converge vertexmarked locally in probability to a random element $(G, o, \mathcal{M}(V(G))) \in \mathcal{G}_*$ having probability law μ_* , when for every r > 0, and for every $(H_*, \mathcal{M}_{H_*}(H_*)) \in \mathcal{G}_*$, as $n \to \infty$,

$$\frac{1}{n} \sum_{\omega \in [n]} \mathbb{1} \left\{ B_r^{(G_n)}(\omega) \simeq H_\star, d_{\mathcal{G}_\star} \left((B_r^{(G_n)}(\omega), \omega, \mathcal{M}(V(B_r^{(G_n)}(\omega))), (H_\star, \mathcal{M}_{H_\star}(H_\star)) \right) \leq \frac{1}{r} \right\}$$
(1.13)
$$\overset{\mathbb{P}}{\to} \mu_\star \left(B_r^{(G)}(o) \simeq H_\star, \ d_{\mathcal{G}_\star} \left((B_r^{(G)}(o), o, \mathcal{M}(V(B_r^{(G)}(o))), (H_\star, \mathcal{M}_{H_\star}(H_\star)) \right) \leq \frac{1}{r} \right).$$

1.4. *Definition of random Pólya point tree.* In this section, we define the random Pólya point tree (RPPT) that will act as the vertex-marked local limit of our preferential attachment graphs. We start by defining the vertex set of this RPPT:

DEFINITION 1.4 (Ulam-Harris set and its ordering). Let $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. The Ulam-Harris set is

$$\mathcal{U} = \bigcup_{n \in \mathbb{N}_0} \mathbb{N}^n.$$

For $x = x_1 \cdots x_n \in \mathbb{N}^n$ and $k \in \mathbb{N}$ we denote the element $x_1 \cdots x_n k$ by $xk \in \mathbb{N}^{n+1}$. The *root* of the Ulam-Harris set is denoted by $\emptyset \in \mathbb{N}^0$.

For any $x \in U$, we say that x has length n if $x \in \mathbb{N}^n$. The lexicographic *ordering* between the elements of the Ulam-Harris set is as follows:

(a) for any two elements $x, y \in \mathcal{U}, x > y$ when the length of x is more than that of y;

(b) if $x, y \in \mathbb{N}^n$ for some n, then x > y if there exists $i \le n$, such that $x_j = y_j \forall j < i$ and $x_i > y_i$.

We use the elements of the Ulam-Harris set to identify nodes in a rooted tree, since the notation in Definition 1.4 allows us to denote the relationships between children and parents, where for $x \in \mathcal{U}$, we denote the k-th child of x by the element xk.

Random Pólya Point Tree (RPPT). The RPPT (M, δ) is an *infinite multi-type rooted random tree* where M is an \mathbb{N} -valued random variable and $\delta > -\inf \operatorname{supp}(M)$. It is a multi-type branching process, with a mixed continuous and discrete type space. We now describe its properties one by one.

Descriptions of the distributions and parameters used.

- $\triangleright \ \text{Define} \ \chi = \tfrac{\mathbb{E}[M] + \delta}{2\mathbb{E}[M] + \delta} \ \text{and} \ \phi = \tfrac{1 \chi}{\chi}.$
- \triangleright Let $\Gamma_{in}(m)$ denote a Gamma distribution with parameters $m + \delta$ and 1, where $m \in \mathbb{N}$.
- \triangleright Let $\Gamma'_{in}(m)$ denote the size-biased distribution of $\Gamma_{in}(m)$ which is also a Gamma distribution with parameters $m + \delta + 1$ and 1.
- \triangleright For $\delta > -\inf \operatorname{supp}(M)$, let $M^{(\delta)}$ be an \mathbb{N} -valued random variable such that

$$\mathbb{P}\left(M^{(\delta)} = m\right) = \frac{m+\delta}{\mathbb{E}[M]+\delta}\mathbb{P}(M=m),$$

i.e., $M^{(\delta)} + \delta$ is a size-biased version of $M + \delta$.

 \triangleright In particular, $M^{(0)}$ is the size-biased distribution of M.

Feature of the vertices of RPPT. Below, to avoid confusion, we use 'node' for a vertex in the RPPT and 'vertex' for a vertex in the PAM. We now discuss the properties of the nodes in $\text{RPPT}(M, \delta)$. Every node except the root in the RPPT has

- \triangleright a *label* ω in the Ulam-Harris set \mathcal{N} (recall Definition 1.4);
- \triangleright an age $A_{\omega} \in [0,1];$
- \triangleright a positive number Γ_{ω} called its *strength*;
- \triangleright a label in {0,Y} depending on the age of the node and its parent, with Y denoting that the node is younger than its parent, and 0 denoting that the node is older than its parent.

Based on its type being 0 or Y, every node ω has an independent \mathbb{N} -valued random variable $m_{-}(\omega)$ associated to it. If ω has type 0, then

- $\triangleright m_{-}(\omega)$ is distributed as $M^{(\delta)}$;
- \triangleright given $m_{-}(\omega), \Gamma_{\omega}$ is distributed as $\Gamma'_{in}(m_{-}(\omega))$.

On the other hand, if ω has type Y, then

- $\triangleright m_{-}(\omega)$ is distributed as $M^{(0)} 1$;
- \triangleright given $m_{-}(\omega), \Gamma_{\omega}$ is distributed as $\Gamma_{in}(m_{-}(\omega)+1)$.

Construction of the RPPT. We next use the above definitions to construct the RPPT using an exploration process. The root is special in the tree. It has label \emptyset and its age A_{\emptyset} is an independent uniform random variable in [0,1]. Since the root \emptyset has no type, $m_{-}(\emptyset)$ is distributed as M. Then the children of the root in the random Pólya point tree are constructed as follows:

- 1. Sample $U_1, \ldots, U_{m_-(\varnothing)}$ uniform random variables on [0, 1], independent of the rest; 2. To nodes $\varnothing 1, \ldots, \varnothing m_-(\varnothing)$ assign the ages $U_1^{1/\chi} A_{\varnothing}, \ldots, U_{m_-(\varnothing)}^{1/\chi} A_{\varnothing}$ and type 0; 3. Assign ages $A_{\varnothing(m_-(\varnothing)+1)}, \ldots, A_{\varnothing(m_-(\varnothing)+d_{\varnothing}^{(in)})}$ to nodes $\varnothing(m_-(\varnothing)+1), \ldots, \varnothing(m_-(\varnothing)+d_{\varnothing})$

 $d_{\alpha}^{(in)}$). These ages are the occurrence times given by a conditionally independent Poisson point process on $[A_{\emptyset}, 1]$ defined by the intensity

(1.14)
$$\rho_{\varnothing}(x) = (1-\chi)\Gamma_{\varnothing}\frac{x^{-\chi}}{A_{\varnothing}^{1-\chi}},$$

where $d_{\varnothing}^{(in)}$ is the total number of points of this process. Assign type Y to them; 4. Draw an edge between \varnothing and each of $\varnothing 1, \ldots, \varnothing (m_{-}(\varnothing) + d_{\varnothing}^{(in)})$; 5. Label \varnothing as explored and nodes $\varnothing 1, \ldots, \varnothing (m_{-}(\varnothing) + d_{\varnothing}^{(in)})$ as unexplored.

Then, recursively over the elements in the set of unexplored nodes we perform the following breadth-first exploration:

- 1. Let ω denote the smallest currently unexplored node;
- Sample m₋(ω) i.i.d. random variables U_{ω1},..., U_{ωm₋(ω)} independently from all the previous steps and from each other, uniformly on [0, 1]. To nodes ω1,..., ωm₋(ω) assign the ages $U_{\omega 1}^{1/\chi} A_{\omega}, \dots, U_{\omega m_{-}(\omega)}^{1/\chi} A_{\omega}$ and type 0 and set them unexplored; 3. Let $A_{\omega(m_{-}(\omega)+1)}, \dots, A_{\omega(m_{-}(\omega)+d_{\omega}^{(in)})}$ be the random $d_{\omega}^{(in)}$ points given by a conditionally
- independent Poisson process on $[A_{\omega}, 1]$ with intensity

(1.15)
$$\rho_{\omega}(x) = (1-\chi)\Gamma_{\omega}\frac{x^{-\chi}}{A_{\omega}^{1-\chi}}.$$

Assign these ages to $\omega(m_{-}(\omega)+1),\ldots,\omega(m_{-}(\omega)+d_{\omega}^{(in)})$, assign them type γ and set them unexplored;

- 4. Draw an edge between ω and each one of the nodes $\omega 1, \ldots, \omega (m_{-}(\omega) + d_{\omega}^{(in)});$
- 5. Set ω as explored.

We call the resulting tree the random Pólya point tree with parameters M and δ , and denote it by $\operatorname{RPPT}(M, \delta)$. Occasionally we drop M and δ while mentioning $\operatorname{RPPT}(M, \delta)$. When M is degenerate and equal to m a.s., we call this the Pólya point tree with parameters m and δ . This coincides with the definition in [8].

1.5. *Main result and discussions.* We now have all the ingredients to state the main result of this article:

THEOREM 1.5 (Local convergence theorem for PA models). Let M be an \mathbb{N} -valued random variable with finite p-th moment for some p > 1 and $\delta > -\inf \operatorname{supp}(M)$. Then models(A), (B), (D), (E) and (F) of the preferential attachment converge vertex-marked locally in probability to the random Pólya point tree with parameters M and δ .

Observations. We make some remarks about the above result:

1. Our proof uses the finiteness of the *p*-th moment for the proofs of some of the concentration bounds around the mean. It would be interesting to identify the precise necessary condition for the local limit result to hold.

- 2. Berger et al. in [8] assumed the fitness parameter δ to be non-negative, but here we allow for negative δ . Note that this accommodates infinite-variance degree distributions used in [33, 35, 42], and suggested in many applied works, see e.g. [26, 27, 41] and the references therein.
- 3. Berger et al. in [8] have shown that $PA_n^{(m,\delta)}(d)$ converges locally in probability to the Pólya point tree with parameters m and δ . Restricting to degenerate distributions, our result can be viewed as an extension of [8] to all preferential attachment models. Moreover our model considers the case where every vertex comes with an i.i.d. number of out-edges which has only finite p-th moment for some p > 1 and we have considered any general starting graph G_0 of size 2. If we do not assume that the initial graph is of size 2, then it increases the computational complexity, and hence we avoid this complication.
- 4. We provide a proof in detail for Model (A). The proof for the models (B) and (D) is very similar and we only indicate the necessary changes. The fact that all these models have the same local limit is a sign of *universality*.
- 5. We prove a *local density* result, which is stronger than in our main result, for models (A), (B) and (E), and interesting in its own right.

Consequences of our main result. We next discuss some consequences of our main result in Theorem 1.5, focusing on degree distributions. Denote

(1.16)
$$\lambda(x) = \frac{1 - x^{1 - \chi}}{x^{1 - \chi}}.$$

It immediately follows from the definition that,

(1.17)
$$\frac{1}{n} \sum_{u} \frac{\mathbb{I}_{\{d_u(n)=k\}}}{m_u} \xrightarrow{\mathbb{P}} p_k$$

where

(1.18)
$$p_k = \mathbb{P}(M + Y(M) = k) = L^{(\emptyset)}(k)k^{-\tau},$$

for some slowly varying function $L^{(\emptyset)}(\cdot)$, with $\tau = \min\{3 + \delta/\mathbb{E}[M], \tau_M\} > 1, \tau_M(\geq p)$ is the power-law exponent of the out-degree distribution (with $\tau_M = +\infty$ when M is light-tailed), and Y(M) is a mixed-Poisson random variable with mixing distribution $\Gamma(M)\lambda(U_{\emptyset})$, where $\Gamma(M)$ is a Gamma variable with parameters $M + \delta$ and 1 and $U_{\emptyset} \sim Unif(0,1)$ and Unif(0,1) is a uniform random variable on (0,1). This was previously established, under related assumptions, for model (E) in [18]. We next extend this result to the convergence of older and younger neighbours of a random vertex:

COROLLARY 1.6 (Asymptotic degree distribution for older and younger neighbours).

(a) As $n \to \infty$, the degree of a uniform older neighbour of a uniform vertex satisfies

(1.19)
$$\frac{1}{n} \sum_{\substack{u,v,j:\\u \stackrel{i}{\to} v}} \frac{\mathbb{I}\left\{d_v(n)=k\right\}}{m_u} \stackrel{\mathbb{P}}{\to} \tilde{p}_k^{(o)} = \mathbb{P}\left(1 + M^{(\delta)} + Y\left(M^{(\delta)} + 1, A_o\right) = k\right), \quad \text{for } k \ge 1$$

where $Y(M^{(\delta)} + 1, A_{\circ})$ is a mixed Poisson random variable with mixing distribution $\Gamma_{in}(M^{(\delta)} + 1)\lambda(A_{\circ})$ and A_{\circ} is distributed as $U_{\varnothing}U_{1}^{1/\chi}$, where U_{\varnothing}, U_{1} are independent Unif(0, 1) random variables.

Similarly, as $n \to \infty$, the degree of a uniform younger neighbour of a uniform vertex satisfies

(1.20)
$$\frac{1}{\sum_{v} \mathbb{1}_{\{d_{v}(n) > m_{v}\}}} \sum_{\substack{u,v,j:\\ u \stackrel{\downarrow}{\to} v}} \frac{\mathbb{1}_{\{d_{v}(n) > m_{v}\}} \mathbb{1}_{\{d_{u}(n) = k\}}}{d_{v}(n) - m_{v}} \\ \xrightarrow{\mathbb{P}} \tilde{p}_{k}^{(\mathsf{Y})} = \mathbb{P}\Big(M^{(0)} + Y\big(M^{(0)}, A_{\mathsf{Y}}\big) = k\Big), \quad \text{for } k \ge 1$$

where $Y(M^{(0)}, A_{\gamma})$ is a mixed Poisson random variable with mixing distribution $\Gamma_{in}(M^{(0)})\lambda(A_{\gamma})$ and A_{γ} has conditional density

(1.21)
$$f_{U_{\varnothing}}(x) = \frac{\rho_{\varnothing}(x)}{\int_{U_{\varnothing}}^{1} \rho_{\varnothing}(s) \, ds} = \frac{x^{-\chi}}{\int_{U_{\varnothing}}^{1} s^{-\chi} \, ds} \,,$$

where $U_{\varnothing} \sim Unif(0,1)$ and $\rho_{\varnothing}(s)$ is as defined in (1.14).

(b) There exist slowly varying functions $L^{(0)}(k)$ and $L^{(Y)}(k)$, such that

(1.22)
$$\tilde{p}_k^{(0)} \sim L^{(0)}(k) k^{-\tau_{(0)}}, \qquad \tilde{p}_k^{(Y)} \sim \Theta(L^{(Y)}(k)) k^{-\tau_{(Y)}} \text{ as } k \to \infty,$$

where $\tau_{\scriptscriptstyle (0)} = \min\{2 + \delta/\mathbb{E}[M], \tau_{\scriptscriptstyle M} - 1\}$ and $\tau_{\scriptscriptstyle (Y)} = \min\{4 + \delta/\mathbb{E}[M], \tau_{\scriptscriptstyle M} - 1\}.$

The convergence is a direct consequence of Theorem 1.5, so what is left is the identification of the limiting probabilities for the RPPT. Thus, the degree distribution of random neighbours is asymptotically *size-biased* compared to the original degree distribution in the graph, as in (1.17)–(1.18). The proof of the corollary is postponed to Section 7. This result is somewhat surprising in the sense that in (1.22), both the tail exponents $\tau_{(o)}$ and $\tau_{(v)}$ contains the $\tau_M - 1$ but the dependence on the PAM power-law exponent is either one larger or one smaller than for the degree distribution of the root.

Idea of proof of Theorem 1.5. For any vertex-marked finite graph $(t, (a_{\omega})_{\omega \in V(t)})$ and $r \in \mathbb{N}$, define

$$N_{r,n}\left(\mathbf{t}, (a_{\omega})_{\omega \in V(\mathbf{t})}\right) = \sum_{v \in [n]} \mathbb{1}_{\left\{B_r^{(G_n)}(v) \simeq \mathbf{t}, |v_{\omega}/n - a_{\omega}| \le 1/r \; \forall \omega \in V(t)\right\}},$$

where G_n is taken as PA_n and v_{ω} is the vertex in G_n corresponding to $\omega \in V(t)$. Then, to prove Theorem 1.5, by Definition 1.3 it is enough to show that $N_{r,n}(t, (a_{\omega})_{\omega \in V(t)})/n$ converges in probability to $\mu(B_r^{(G)}(\emptyset) \simeq t, |A_{\omega} - a_{\omega}| \le 1/r \ \forall \omega \in V(t))$, where μ is the law of the limiting RPPT graph, and A_{ω} is the mark of the vertex in RPPT corresponding to $\omega \in V(t)$ and t is a tree. We aim to prove this convergence using a second moment method, i.e., we will prove that

$$(1.23) \quad \frac{1}{n} \mathbb{E} \left[N_{r,n} \left(\mathbf{t}, (a_{\omega})_{\omega \in V(\mathbf{t})} \right) \right] \to \mu \left(B_r^{(G)}(\varnothing) \simeq \mathbf{t}, |A_{\omega} - a_{\omega}| \le 1/r \; \forall \omega \in V(t) \right),$$

and
$$\frac{1}{n^2} \mathbb{E} \left[N_{r,n} \left(\mathbf{t}, (a_{\omega})_{\omega \in V(\mathbf{t})} \right)^2 \right] \to \mu \left(B_r^{(G)}(\varnothing) \simeq \mathbf{t}, |A_{\omega} - a_{\omega}| \le 1/r \; \forall \omega \in V(t) \right)^2.$$

For proving the first, we show that the *joint density* of the ages of the vertices in the *r*-neighbourhood of a uniformly chosen vertex in PAM converges to that of RPPT. Calculating this joint density explicitly is somewhat complicated because of the dependence structure in the edge-connection probabilities of PAM models. In [7], the authors provide a Pólya urn representation of model (D), which made the proof in [8] possible since this representation implies that the edges are *conditionally independent*. An essential step in our proof is thus

1

11

to construct similar Pólya urn descriptions for models (A), (B) and (D) and show that the models are equal in distribution to the corresponding Pólya urn representations. Sénizergues [38] proved a similar result to the first part of the proof for preferential attachment trees and extended the result for preferential attachment graph with random out-degree. We tend to use a Pólya urn description with a different set of Beta random variables. We do this by explicitly calculating the graph probabilities. Our result works for i.i.d. random δ also. With this distributional equivalence in hand, we can now compute the above joint density, and show that it converges to that of the RPPT.

For the second moment, we first expand the square of the sum arising in the numerator. From the expansion we observe that along with some vanishing terms, we obtain the joint density of the *r*-neighbourhoods of *two* uniformly chosen vertices. Next we prove that the *r*-neighbourhoods of two uniformly chosen vertices are disjoint with high probability (whp). Again, for the joint density calculation, we use the Pólya urn description of our models. Since the edge-connection events are conditionally independent by the Pólya urn representations, the two neighbourhoods are *independent* when they are disjoint. Therefore the joint density factorizes and we obtain the required result.

Though our main steps for proving the theorem are the same as those in [8], our proof techniques differ significantly, for example, we avoid the induction argument in the neighbourhood size and also the coupling of the preferential attachment model with the Pólya urn graphs. To summarize, we have two crucial steps in proving the main theorem:

- (a) Equivalence: there exists Pólya urn representations for models (A), (B) and (D);
- (b) **Convergence:** the joint density of the ages of the vertices in the *r*-neighbourhood of the Pólya urn graphs converges to that of the RPPT.

For model (E), the edges are connected independently and the degrees of the older vertices are updated only after the new vertex is included in the graph with *all* its out-edges. This edge-connection procedure is different from other models and we do not have a Pólya urn representation for this graph. We first prove the convergence of the models having a Pólya urn representation. Then we couple model (D) and (E), such that the probabilities of observing the r-neighbourhoods of a uniformly chosen vertex in the graphs are asymptotically equal. Since for model (D), we have proved the local convergence, convergence for model (E) follows immediately. Note, however, that this does *not* imply the convergence of the joint density of the ages of the vertices in the r-neighbourhood for model (E). Model (F) is dealt with in a similar way as model (E).

Relation to the literature. Pólya urn representations of the preferential attachment models were previously studied by Berger et al. [7] and Sénizergues [38]. Local convergence of models (D) and (E) were first derived by Berger et al. [8] for fixed out-degrees and non-negative δ . Results on local convergence of related PAMs have been established by Y. Y. Lo [34], who analysed the local limit of the preferential attachment trees but i.i.d. random fitness parameter δ . Rudas, Tóth and Valkó [37] proved local convergence almost surely for general preferential attachment trees, based on continuous time embedding, which gives continuous time branching process by Jagers and Nerman [32] in full generality.

After uploading our paper to arXiv, we were informed about the work of Banerjee, Deka and Olvera-Cravioto [3] which also proves local convergence in probability of preferential attachment models with i.i.d. out-degrees. The paper assumes finite first moment of the random out-degree and does not rely on the Pólya urn representation of the preferential attachment model but rather on an extension of the work of the first and third authors that relates preferential attachment models to continuous-time branching processes [30]. Further, in addition to the local convergence, our paper establishes density convergence of the preferential attachment model, and identified the degree distribution of the nodes of the limiting tree. Although the proof techniques are different, both techniques are interesting in their own right.

Open problems. We believe that our results can be an essential ingredient to the proof of various related properties of PAMs, such as the behavior of percolation on, and graph distances in, them. It would further be interesting to extend the work to random fitness distributions, as well as to the model with conditionally independent edges as studied by Dereich and Mörters [21, 22, 23].

Structure of the rest of the article. In Section 2, we extend the definition of Pólya urn graphs to collapsed Pólya urn graphs (for equivalence with models (A) and model (B)). In Section 3 we prove that the preferential attachment models (A), (B) and (D) are equal in distribution to their respective Pólya urn representations. Section 4 deals with some results that are necessary to show the *r*-neighbourhood density convergence. In Section 5 we show the convergence part of the proof for model (A). Using the similarity of the models in Section 3, the proofs for the models (B) and (D) follow in a same way. In Section 6, we show the coupling between models (D), and (E) and (F). Section 7 provides a proof of Corollary 1.6. The details of the proofs for degenerate out-degrees. In Appendix B, we provide the proof of equivalence of models (B) and (D) and their respective Pólya urn representations. These proofs follow the same line as those in Section 3 for the equivalence of model (A) and its Pólya urn representation. Appendix C deals with the adaptation of the coupling between model (D) and (E) done in Section 6, to model (F). The proof of Corollary 1.6(b) is deferred to Appendix D due to similar proof techniques used in the literature.

2. Pólya Urn Graph Description. The first and foremost difficulty in dealing with preferential attachment models is that the edge-connection events are not independent. We now give a representation of our PAMs where these events are *conditionally independent*, extending the results of Berger et al. in [7] that apply to $PA_n^{(m,\delta)}(d)$. Intuitively, every new edge-connection in the preferential attachment model with intermediate degree updates can be viewed as drawing a ball uniformly from a Pólya urn with balls having multiple colours corresponding to the various vertices in the graph. This is the place where the intermediate degree update after every edge-connection step (such as in models (E) and (F)), then the model may seem simpler, but in fact it is harder to work with since we do not get a Pólya urn description for it. We handle these models in Section 6 using a *coupling approach* instead.

2.1. *Pólya Urn Graph.* The sequential preferential attachment models (A), (B) and (D) in Section 1.2 can be interpreted as an experiment with n urns corresponding to the vertices in the graph, where the number of balls in each urn represents the degree of the corresponding vertex in the graph. First, we introduce a new class of random graphs that we later prove (in Section 3) to have the same distribution as our PAMs:

DEFINITION 2.1 (Pólya Urn Graphs). Define the following:

- $\triangleright (U_k)_{k>1}$ is a sequence of i.i.d. uniform random variables in [0, 1];
- ▷ given $\overline{\boldsymbol{m}} = (1, 1, m_3, m_4, \ldots), \boldsymbol{\psi} = (\psi_k)_{k \in \mathbb{N}}$ is a sequence of conditionally independent Beta random variables with support in [0, 1], such that $\mathbb{P}(\psi_k = 1) = 0$ for all $k \ge 2$ and $\psi_1 = 1$ point-wise;
- \triangleright let G_0 be the initial graph of size 2 and $n \in \mathbb{N}$ be the size of the graph.

Define $\mathcal{S}_0^{(n)} = 0$ and $\mathcal{S}_n^{(n)} = 1$ and for $k \in [n-1]$, define

(2.1)
$$S_k^{(n)} = (1 - \psi)_{(k,n]}, \quad \text{and} \quad \mathcal{I}_k^{(n)} = \left[S_{k-1}^{(n)}, S_k^{(n)}\right),$$

where, for $A \subseteq [n]$,

(2.2)
$$(1-\psi)_A = \prod_{a \in A} (1-\psi_a)$$

Then, $\operatorname{PU}_n^{(\operatorname{SL})}(\boldsymbol{m}, \boldsymbol{\psi})$ and $\operatorname{PU}_n^{(\operatorname{NSL})}(\boldsymbol{m}, \boldsymbol{\psi})$, the *Pólya urn graph* of size *n* with and without self-loops, respectively, are defined as follows:

 \triangleright in $\mathrm{PU}_n^{(\mathrm{NSL})}(\boldsymbol{m}, \boldsymbol{\psi})$, the *j*-th edge from vertex k is attached to vertex $u \in [k]$ precisely when

$$(2.3) U_{m_{[k-1]}+j}\mathcal{S}_{k-1}^{(n)} \in \mathcal{I}_u^{(n)}$$

Observe that self-loops are absent here since $(0, \mathcal{S}_{k-1}^{(n)})$ and $\mathcal{I}_k^{(n)}$ are two disjoint sets; \triangleright for $\mathrm{PU}_n^{(\mathrm{SL})}(\boldsymbol{m}, \boldsymbol{\psi})$, the condition (2.3) is replaced by

$$(2.4) U_{m_{[k-1]}+j}\mathcal{S}_k^{(n)} \in \mathcal{I}_u^{(n)}$$

To specify a Pólya urn graph $PU_n(\boldsymbol{m}, \boldsymbol{\psi})$, we need to specify the out-edge distribution M, and the parameters of the Beta variables $\boldsymbol{\psi}$. Berger et al. in [8] have shown that $PA_n^{(m,\delta)}(d)$ is equal in distribution to $PU_n^{(NSL)}(\boldsymbol{m}^{(1)}, \boldsymbol{\psi})$, where $\boldsymbol{\psi}$ is taken as a sequence of independent Beta random variables with certain parameters and M is degenerate at m, i.e. $\boldsymbol{m}^{(1)} = (1, 1, m, m, ...)$. Since $PA_n^{(D)}(\boldsymbol{m}, \delta)$ is a generalized version of $PA_n^{(m,\delta)}(d)$ with i.i.d. random out-degrees, we show that this model also has a Pólya urn description. We sketch an outline in Section 3 [Theorem 3.10] that conditionally on \boldsymbol{m} , $PA_n^{(D)}(\boldsymbol{m}, \delta)$ and $PU_n^{(NSL)}(\boldsymbol{m}, \boldsymbol{\psi})$ are also equal in distribution. On the other hand, it is evident that the deterministic versions of model (B) and (D) are equivalent for m = 1. From the edge-connection probabilities of model (A) and (B), we can observe that they are different only in whether they give rise to self-loops or not. Van der Hofstad in [40, Chapter 5] shows that model (A) and (B) can be obtained by a collapsing procedure for degenerate M. Therefore for obtaining a Pólya urn graph equivalence for models (A) and (B), a generalisation of this collapsing procedure is helpful, and we continue by describing such collapsing procedures.

2.2. Collapsing Operator. Here we generalize the collapsing procedure discussed in [40]. Let $\mathbf{r} = (r_1, r_2, \ldots) \in \mathbb{N}^{\mathbb{N}}$ and let \mathcal{H} be the set of all finite vertex-labelled graphs. Then the collapsing operator C_r , acting on \mathcal{H} and collapsing groups of vertices of size r_i into vertex v_i , is defined as follows:

▷ let $G \in \mathcal{H}$ be a vertex-labelled graph of size n and $n \in (r_{[k-1]}, r_{[k]}]$ for some $k \in \mathbb{N}$, where

$$r_{[l]} = \sum_{i \le l} r_i ;$$

- ▷ group the vertices $\{r_{[i-1]} + 1, ..., r_{[i]}\}$ and name the groups v_i for all i < k, while the group v_k contains the vertices $\{r_{[k-1]} + 1, ..., n\}$;
- \triangleright collapse the vertices of each of these v_k groups into one vertex, and name the new vertex k;
- ▷ edges originating and ending in the same group form self-loops in the new graph;

edges between two different groups form edges between the respective vertices in the new graph.

Note that if we fix $r_1 = r_2 = 1$ and $r_i = m$ for all $i \ge 3$ and suitable G_0 , then we get back the collapsing procedure used first in [12], and further discussed in [40]. We now discuss the construction of models (A) and (B) through this collapsing operator, conditionally on the i.i.d. out-degrees described by m.

PAM construction by collapsing. We start with a vertex-labelled graph G_0 of size 2 and degrees a_1 and a_2 , respectively. First, we explain the construction of model (A) using collapsing.

Every $v \ge 3$ comes with exactly one edge. Given $m = (1, 1, m_3, ...)$, the incoming vertex $v = m_{[i-1]} + j$ for some $i \in [3, n)$ and $j \le m_i$, connects to one of $u \in [v]$ with probability

(2.5)
$$\mathbb{P}_m\left(v \rightsquigarrow u \mid \mathrm{PA}_{v-1}^{(\mathrm{A})}(\boldsymbol{m}, 1, \delta)\right) = \begin{cases} \frac{d_u(v-1) + \delta(u)}{c_{v,j}} & \text{when } v > u, \\ \frac{1 + \delta(u)}{c_{v,j}} & \text{when } v = u. \end{cases}$$

Here $\delta(u) = \delta/m_k$ when $u \in (m_{[k-1]}, m_{[k]}]$,

$$c_{i,j} = a_{[2]} + 2\left(m_{[i-1]} + j - 2\right) - 1 + (i-1)\delta + \frac{j}{m_i}\delta$$

and $\operatorname{PA}_{v-1}^{(A)}(\boldsymbol{m}, 1, \delta)$ denotes the graph formed after the (v-1)-st vertex is added, with $d_u(v-1)$ denoting the degree of the vertex u in $\operatorname{PA}_{v-1}^{(A)}(\boldsymbol{m}, 1, \delta)$. Continue the process until the $m_{[n]}$ -th vertex is added. We obtain $\operatorname{PA}_n^{(A)}(\boldsymbol{m}, \delta)$ by applying $\mathcal{C}_{\boldsymbol{m}}$ on $\operatorname{PA}_{m_{[n]}}^{(A)}(\boldsymbol{m}, 1, \delta)$. Therefore, for model (A), the conditional edge-connection probabilities are given by (1.3), as required.

To construct $PA_n^{(B)}(\boldsymbol{m}, \delta)$ by a collapsing procedure, we do not allow for self-loops for m = 1 as we did in (2.5). The rest of the process remains the same. Starting from the same initial graph G_0 , conditionally on \boldsymbol{m} , every $v \ge 3$ comes with exactly one edge. The incoming vertex $v = m_{[i-1]} + j$ for some $i \in [3, n]$ and $j \le m_i$, connects to $u \in [v - 1]$ with the same probability as in (2.5). Since here we do not allow for any self-loop, the normalising constant in the denominator now is,

$$c_{i,j} = a_{[2]} + 2\left(m_{[i-1]} + j - 3\right) + (i-1)\delta + \frac{(j-1)}{m_i}\delta$$

and $\operatorname{PA}_{v}^{\scriptscriptstyle(B)}(\boldsymbol{m},1,\delta)$ denotes the graph formed after the v-th vertex is added, with $d_{u}(v)$ denoting the degree of the vertex u in $\operatorname{PA}_{v}^{\scriptscriptstyle(B)}(\boldsymbol{m},1,\delta)$. Continue the process until the $m_{[n]}$ -th vertex is added. We obtain $\operatorname{PA}_{n}^{\scriptscriptstyle(B)}(\boldsymbol{m},\delta)$ by applying $\mathcal{C}_{\boldsymbol{m}}$ on $\operatorname{PA}_{m_{[n]}}^{\scriptscriptstyle(B)}(\boldsymbol{m},1,\delta)$. Therefore, for model (B), the edge-connection probabilities are given by (1.5).

REMARK 2.2 (Initial graph is preserved in collapsing). Observe that we always choose $r_1 = r_2 = 1$ while performing the collapsing operator on the pre-collapsed preferential attachment graphs. This is done intentionally to preserve the structure of the initial graph G_0 . If we start with an initial graph of size $\ell \ge 1$ then we can choose $r_1 = r_2 = \ldots = r_\ell = 1$ to preserve the initial graph structure in both collapsed and pre-collapsed preferential attachment graphs.

We will use this collapsing operator to introduce an extension of the Pólya urn graph, the collapsed Pólya urn graph.

2.3. Collapsed Pólya Urn Graphs. $PA_n^{(B)}(m, 1, \delta)$ is essentially the same as model (D) when every vertex comes with exactly one out-edge, but δ is different for every vertex. Therefore, we expect that the Pólya urn graph extends to this graph. Similarly to the construction of model (B) through a collapsing procedure, we collapse the Pólya urn graph.

Conditionally on m, we construct the collapsed Pólya Urn graph by using our collapsing construction on the Pólya Urn Graph defined in Definition 2.1 as follows:

DEFINITION 2.3 (Collapsed Pólya Urn graph). We first construct $PU_{m_{[n]}}(1, \psi)$ with every new vertex having exactly one out-edge and initial graph G_0 of size 2. Conditionally on $m = (1, 1, m_3, m_4, ...)$, the graph $CPU_n(m, \psi)$ is defined as $C_m(PU_{m_{[n]}}(1, \psi))$. The label SL or NSL for the CPU_n will be determined by the label of the $PU_{m_{[n]}}$. We denote the two CPU's with and without self-loops as $CPU_n^{(SL)}$ and $CPU_n^{(NSL)}$ respectively.

REMARK 2.4 (Self-loops for $CPU_n^{(NSL)}$). $CPU_n^{(NSL)}$ may contain self-loops because of collapsing.

We end this section by deriving the connection probabilities for $CPU_n^{(SL)}$ and $CPU_n^{(NSL)}$. For $k \ge 1$ and $j \in [m_k]$, define

(2.6)
$$S_{k,j}^{(n)} = \prod_{l=m_{[k-1]}+j+1}^{m_{[n]}} (1-\psi_l), \quad \text{and} \quad S_k^{(n)} = S_{k,m_k}^{(n)};$$

and the intervals $\mathcal{I}_k^{(n)} = [\mathcal{S}_{k-1}^{(n)}, \mathcal{S}_k^{(n)})$. Let $\mathbb{P}_m^{(\mathrm{SL})}$ and $\mathbb{P}_m^{(\mathrm{NSL})}$ denote the conditional law given \boldsymbol{m} of $\mathrm{CPU}_n^{(\mathrm{SL})}$ and $\mathrm{CPU}_n^{(\mathrm{SL})}$, respectively. Then from the construction of the $\mathrm{CPU}_n^{(\mathrm{SL})}$ it follows that, for $u \geq 3$,

(2.7)
$$\mathbb{P}_{m}^{(\mathrm{SL})}\left(u \stackrel{j}{\rightsquigarrow} v \middle| (\psi_{k})_{k \in \mathbb{N}}\right) = \begin{cases} \frac{\mathcal{S}_{v}^{(n)} - \mathcal{S}_{v-1}^{(n)}}{\mathcal{S}_{u,j}^{(n)}} & \text{for } u > v, \\ \frac{\mathcal{S}_{u,j}^{(n)} - \mathcal{S}_{u-1}^{(n)}}{\mathcal{S}_{u,j}^{(n)}} & \text{for } u = v. \end{cases}$$

This probability is for $CPU_n^{(SL)}$. For $CPU_n^{(NSL)}$ instead, the expression in (2.7) becomes

(2.8)
$$\mathbb{P}_{m}^{(\text{NSL})}\left(u \stackrel{j}{\leadsto} v \middle| (\psi_{k})_{k \in \mathbb{N}}\right) = \begin{cases} \frac{S_{u}^{(n)} - S_{v-1}^{(n)}}{S_{u,j-1}^{(n)}} & \text{for } u > v, \\ \frac{S_{u,j-1}^{(n)} - S_{u-1}^{(n)}}{S_{u,j-1}^{(n)}} & \text{for } u = v. \end{cases}$$

Now that we have introduced the relevant random graph models, in the next section, we will show that the collapsed versions of the Pólya graph models have the same distribution as our preferential attachment models.

3. Equivalence of Preferential Attachment Models. As explained in Section 1.5, there are two key steps in the proof of the local limit result in Theorem 1.5. In the first step, we show that the PAM with random out-degrees is equal in distribution to collapsed Pólya urn graph (CPU) or Pólya urn graph (PU), with the ψ random variables defined appropriately. We bring the notion of CPU in between since it implies the much-appreciated independence structure of the edge-connection events, which is not valid in PAM. We provide explicit calculations for model (A), and state the modifications required for other models.

3.1. Equivalence of model (A) and CPU^(SL). Recall that model (A) has i.i.d. out-degrees for every vertex. Thus, in order to couple it with our CPU, it must have the same out-degrees. Given $\boldsymbol{m} = (m_1, m_2, m_3, \ldots)$, with $m_1 = m_2 = 1$, we thus aim to couple PA^(A)_n(\boldsymbol{m}, δ) and CPU^(SL).

For $i \in [2]$, a_i denotes the degree of vertex i in the initial graph G_0 , while for i > 2 define $a_i \equiv 1$. Let \mathcal{H}_n be the set of all finite vertex-labelled graphs G of size n and let $\mathcal{H}_m(G) = \left\{ G_e \in \mathcal{H}_{m_{[n]}} : \mathcal{C}_m(G_e) \stackrel{\star}{\simeq} G \right\}$ denote the set of graphs that are mapped to G by the collapsing operator \mathcal{C}_m .

For $k \geq 3$ and $l \in [m_k]$, define

(3.1)
$$\psi_{m_{[k-1]}+l} \sim \text{Beta}\left(1 + \frac{\delta}{m_k}, a_{[2]} + 2\left(m_{[k-1]} + l - 3\right) + (k-1)\delta + \frac{(l-1)}{m_k}\delta\right),$$

where $a_{[2]} = a_1 + a_2$ and, for $k \le 2$,

(3.2)
$$\psi_1 \equiv 1$$
, and $\psi_2 \sim \text{Beta}(a_2 + \delta, a_1 + \delta)$

We abbreviate $\psi = (\psi_i)_{i \ge 1}$ for the collection of Beta variables, where we emphasize that these variables are *conditionally independent* given the random out-degrees m. Our main result concerning the relation between collapsed Pólya graphs and model (A) is as follows:

THEOREM 3.1 (Equivalence of model (A) and $CPU^{(SL)}$). For any graph $G \in \mathcal{H}_n$,

(3.3)
$$\mathbb{P}_m\left(\mathrm{PA}_n^{\scriptscriptstyle(\mathrm{A})}(\boldsymbol{m},\delta) \stackrel{\star}{\simeq} G\right) = \mathbb{P}_m\left(\mathrm{CPU}_n^{\scriptscriptstyle(\mathrm{SL})}(\boldsymbol{m},\psi) \stackrel{\star}{\simeq} G\right).$$

We emphasize that Theorem 3.1 describes a *conditional* result given m, i.e., it is conditionally on m.

We prove Theorem 3.1 by proving it for the pre-collapsed version of both graphs, and equating their conditional distributions. The following proposition helps us in equating the conditional probabilities of the pre-collapsed graphs:

PROPOSITION 3.2 (Equivalence of pre-collapsed model (A) and $PU^{(SL)}$). Conditionally on m, for any graph $H \in \mathcal{H}_{m_{[n]}}$,

(3.4)
$$\mathbb{P}_m\left(\mathrm{PA}_n^{\scriptscriptstyle{(A)}}(\boldsymbol{m},1,\delta) \stackrel{\star}{\simeq} H\right) = \mathbb{P}_m\left(\mathrm{PU}_{m_{[n]}}^{\scriptscriptstyle{(\mathrm{SL})}}(\boldsymbol{1},\boldsymbol{\psi}) \stackrel{\star}{\simeq} H\right).$$

PROOF OF THEOREM 3.1 SUBJECT TO PROPOSITION 3.2. Proposition 3.2 essentially provides us with the pre-collapsing equivalence of the graphs. Since, conditionally on m, $\left\{ PA_{m_{[n]}}^{(A)}(m, 1, \delta) \stackrel{\star}{\simeq} H \right\}$ are disjoint events for $H \in \mathcal{H}_{m}(G)$, the probability on the RHS (right hand side) of (3.3) for model (A) can be written in terms of the pre-collapsed graphs as

(3.5)

$$\mathbb{P}_{m}\left(\mathrm{PA}_{n}^{\scriptscriptstyle(A)}(\boldsymbol{m},\delta)\overset{\star}{\simeq}G\right) = \mathbb{P}_{m}\left(\bigcup_{H\in\mathcal{H}_{\boldsymbol{m}}(G)}\left\{\mathrm{PA}_{m_{[n]}}^{\scriptscriptstyle(A)}(\boldsymbol{m},1,\delta)\overset{\star}{\simeq}H\right\}\right)$$

$$=\sum_{H\in\mathcal{H}_{\boldsymbol{m}}(G)}\mathbb{P}_{m}\left(\mathrm{PA}_{m_{[n]}}^{\scriptscriptstyle(A)}(\boldsymbol{m},1,\delta)\overset{\star}{\simeq}H\right).$$

Similarly the probability on the LHS (left hand side) of (3.3) for $CPU_n^{(SL)}$ can be written in terms of $PU_{m_{[n]}}^{(SL)}$ as

(3.6)

$$\mathbb{P}_{m}\left(\operatorname{CPU}_{n}^{(\mathrm{SL})}(\boldsymbol{m},\boldsymbol{\psi})\overset{\star}{\simeq} G\right) = \mathbb{P}_{m}\left(\bigcup_{H\in\mathcal{H}_{m}(G)}\left\{\operatorname{PU}_{m_{[n]}}^{(\mathrm{SL})}(\boldsymbol{1},\boldsymbol{\psi})\overset{\star}{\simeq} H\right\}\right)$$

$$=\sum_{H\in\mathcal{H}_{m}(G)}\mathbb{P}_{m}\left(\operatorname{PU}_{m_{[n]}}^{(\mathrm{SL})}(\boldsymbol{1},\boldsymbol{\psi})\overset{\star}{\simeq} H\right).$$

Now from Proposition 3.2 it follows that the summands in (3.5) and (3.6) are equal. Hence, conditionally on m, $PA_n^{(A)}$ and $CPU_n^{(SL)}$ are equal in distribution.

We now move towards the proof of Proposition 3.2. Berger et al. [8] have proved a version of Theorem 3.1 for model (D) and degenerate out-degrees using an extension to multiple urns of the Pólya urn characterization in terms of conditionally independent events by de Finetti's Theorem. We could adapt this proof. Instead, we prove Proposition 3.2 by explicitly calculating the graph probabilities of both random graphs and equating them term by term, which we now show. This proof is interesting in its own right.

Let v(u) denote the vertex to which the out-edge from u connects in H. Then from (2.5),

(3.7)

$$\mathbb{P}_{m}\left(\mathrm{PA}_{m_{[n]}}^{(A)}(\boldsymbol{m},1,\delta) \stackrel{\star}{\simeq} H\right) = \prod_{u \in [3,n]} \prod_{j \in [m_{u}]} \frac{d_{v(m_{[u-1]}+j)}(m_{[u-1]}+j-1) + \delta(v(m_{[u-1]}+j))}{a_{[2]} + 2(m_{[u-1]}+j-2) + \left((u-1) + \frac{j}{m_{u}}\right)\delta - 1}.$$

The following lemma simplifies and rearranges the factors in the numerator of (3.7):

LEMMA 3.3 (Rearrangement of the numerator of (3.7)). The numerator of (3.7) can be rearranged as

(3.8)
$$\prod_{u \in [3,n]} \prod_{j \in [m_u]} \left(d_{v(m_{[u-1]}+j)}(m_{[u-1]}+j-1) + \delta(v(m_{[u-1]}+j)) \right)$$
$$= \prod_{k \in [n]} \prod_{l \in [m_u]} \prod_{i=a_{m_{[k-1]}+l}}^{d_{m_{[k-1]}+l}(H)-1} \left(i + \frac{\delta}{m_k}\right),$$

where $d_v(H)$ denotes the degree of the vertex v in the graph H.

PROOF. Observe that the factors in the numerator of RHS of (3.7) depend on the receiver's degree. Since the edges from the new vertices connect to one of the existing vertices (or itself), the product in the numerator of (3.7) can be rewritten as

(3.9)
$$\prod_{u \in [3,n]} \prod_{j \in [m_u]} \left(d_{v(m_{[u-1]}+j)}(m_{[u-1]}+j-1) + \delta(v(m_{[u-1]}+j)) \right) \\ = \prod_{s \in [1,m_{[n]}]} \prod_{\substack{u \in [3,n], j \in [m_u] \\ v(m_{[u-1]}+j)=s}} \left(d_s(m_{[u-1]}+j-1) + \delta(s) \right).$$

For the very first incoming edge to $s \in [m_{[n]}]$, we have the factor of $(a_s + \delta(s))$ and for the remaining ones, we have a factor in the RHS of (3.9) with an increment of 1. On the other hand, for the last incoming edge to s in H, we have the factor $(d_s(H) - 1 + \delta(s))$. For any $s \in [m_{[n]}]$, there exists $k \in [n]$ and $l \in [m_u]$ such that $s = m_{[k-1]} + l$ and $\delta(s) = \delta/m_k$. Therefore the LHS of (3.9) can be further simplified as

(3.10)
$$\prod_{s \in [1, m_{[n]}]} \prod_{\substack{u \in [3, n], j \in [m_u] \\ v(m_{[u-1]}+j)=s}} (d_s(m_{[u-1]}+j-1)+\delta(s))$$
$$= \prod_{k \in [n]} \prod_{l \in [m_u]} \prod_{i=a_{m_{[k-1]}+l}}^{d_{m_{[k-1]}+l}(H)-1} \left(i+\frac{\delta}{m_k}\right).$$

By Lemma 3.3 and a rearrangement of numerator, the graph probability in (3.7) can be written as

$$\mathbb{P}_{m}\left(\mathrm{PA}_{m_{[n]}}^{(A)}(\boldsymbol{m},1,\delta) \stackrel{\star}{\simeq} H\right)$$

$$(3.11) = \prod_{u \in [n]} \prod_{j \in [m_{u]}} \prod_{i=a_{m_{[u-1]}+j}}^{d_{m_{[u-1]}+j}(H)-1} \left(i + \frac{\delta}{m_{u}}\right)$$

$$\times \prod_{u \in [3,n]} \prod_{j \in [m_{u]}} \frac{1}{a_{[2]} + 2(m_{[u-1]} + j - 2) + \left((u-1) + \frac{j}{m_{u}}\right)\delta - 1}.$$

Next, we calculate the graph probabilities of $\mathrm{PU}_{m_{[n]}}^{\scriptscriptstyle{(\mathrm{SL})}}(\mathbf{1}, \boldsymbol{\psi})$ and show that these agree. To calculate $\mathbb{P}_m\left(\mathrm{PU}_{m_{[n]}}^{\scriptscriptstyle{(\mathrm{SL})}}(\mathbf{1}, \boldsymbol{\psi}) \stackrel{\star}{\simeq} H\right)$, we condition on the Beta random variables as well. We denote the conditional measure by $\mathbb{P}_{m,\psi}$, i.e., for every event \mathcal{E} ,

$$\mathbb{P}_{m,\psi}(\mathcal{E}) = \mathbb{P}\left(\mathcal{E} \mid (m_k)_{k \ge 3}, (\psi_k)_{k \ge 1}\right)$$

Under this conditioning, the edges of $\mathrm{PU}_{m_{[n]}}^{(\mathrm{SL})}$ are independent. First we calculate the conditional edge-connection probabilities for $\mathrm{PU}_{m_{[n]}}^{(\mathrm{SL})}(\mathbf{1}, \boldsymbol{\psi})$:

LEMMA 3.4 (Conditional edge-connection probability of $PU^{(SL)}$). Conditionally on m and $(\psi_k)_{k\geq 1}$ defined in (3.1) and (3.2), the probability of connecting the edge from v to u in $PU^{(SL)}_{m[n]}(\mathbf{1}, \psi)$ is given by $\psi_v(1 - \psi)_{(v,u]}$.

PROOF. By the construction of $\mathrm{PU}_{m_{[n]}}^{\scriptscriptstyle\mathrm{(SL)}}(\mathbf{1}, oldsymbol{\psi}),$

 $\mathcal{S}_k^{^{(m_{[n]})}} = (1-\psi)_{(k,m_{[n]}]}, \qquad \text{and} \qquad |I_k^{^{(m_{[n]})}}| = \psi_k (1-\psi)_{(k,m_{[n]}]}.$

Taking the ratio gives

$$\mathbb{P}_{m,\psi}\left(u \rightsquigarrow v\right) = \frac{|I_v^{(m_{[n]})}|}{\mathcal{S}_u^{(m_{[n]})}} = \psi_v \frac{(1-\psi)_{(v,m_{[n]}]}}{(1-\psi)_{(u,m_{[n]}]}} = \psi_v (1-\psi)_{(v,u]}.$$

In Pólya urn graphs, conditionally on the Beta random variables, the edges are added independently, leading to the following lemma:

LEMMA 3.5 (Conditional density of $PU^{(SL)}$). For any graph $H \in \mathcal{H}_{m_{[n]}}$,

(3.13)
$$\mathbb{P}_{m,\psi}\left(\mathrm{PU}_{m_{[n]}}^{(\mathrm{SL})}(\mathbf{1},\psi) \stackrel{\star}{\simeq} H\right) = \prod_{s \in [2,n]} \psi_s^{p_s} (1-\psi_s)^{q_s},$$

where

(3.14)
$$p_s = d_s(H) - a_s, \quad and \quad q_s = \sum_{u \in (m_{[2]}, m_{[n]}]} \mathbb{1}_{\{s \in (v(u), u]\}}.$$

PROOF. For every incoming edge of the vertex u, there is a vertex $w \ge u$ such that v(w) = u. Conditionally on m and ψ , the edge-connection events are independent. We then note that there are p_s many incoming edges for vertex s and the factor $\psi_s^{p_s}$ comes from that. Every u such that $s \in (v(u), u]$ gives rise to one factor $1 - \psi_s$, giving rise to q_s many factors $1 - \psi_s$.

Next we aim to take the expectation w.r.t. ψ , and for this, we compute the expectation of powers of Beta variables:

LEMMA 3.6 (Integer moments of Beta distribution). For all $a, b \in \mathbb{N}$ and $\psi \sim \text{Beta}(\alpha, \beta)$,

(3.15)
$$\mathbb{E}\left[\psi^{a}(1-\psi)^{b}\right] = \frac{(\alpha+a-1)_{a}(\beta+b-1)_{b}}{(\alpha+\beta+a+b-1)_{a+b}},$$

where $(n)_k = \prod_{i=0}^{k-1} (n-i)$ for $k \ge 1$.

PROOF. A direct calculation for Beta random variables shows that

$$\mathbb{E}\left[\psi^{a}(1-\psi)^{b}\right] = \frac{B(\alpha+a,\beta+b)}{B(\alpha,\beta)} = \frac{\Gamma(\alpha+\beta)\Gamma(\alpha+a)\Gamma(\beta+b)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\alpha+\beta+a+b)}$$
$$= \frac{(\alpha+a-1)_{a}(\beta+b-1)_{b}}{(\alpha+\beta+a+b-1)_{a+b}}.$$

As a consequence of Lemma 3.5 and 3.6, we can calculate $\mathbb{P}_m\left(\mathrm{PU}_{m_{[n]}}^{(\mathrm{SL})}(\mathbf{1}, \boldsymbol{\psi}) \stackrel{\star}{\simeq} H\right)$:

LEMMA 3.7. For $H \in \mathcal{H}_{\boldsymbol{m}}(G)$,

(3.16)
$$\mathbb{P}_{m}\left(\mathrm{PU}_{m_{[n]}}^{(\mathrm{SL})}(\mathbf{1}, \psi) \stackrel{\star}{\simeq} H\right) = \prod_{s \in [2, m_{[n]})} \frac{(\alpha_{s} + p_{s} - 1)_{p_{s}}(\beta_{s} + q_{s} - 1)_{q_{s}}}{(\alpha_{s} + \beta_{s} + p_{s} + q_{s} - 1)_{p_{s} + q_{s}}},$$

where α_s and β_s are the first and second parameters of the Beta random variables defined in (3.1) and (3.2), p_s, q_s are defined in Lemma 3.5 and $\inf \operatorname{supp}(M)$ is the minimum of the support of the random variable M.

PROOF. Lemma 3.5 describes the conditional probability distribution of the Pólya Urn graphs given the independent Beta random variables. To obtain the unconditional probability, we take the expectation on the RHS of (3.13) w.r.t. the Beta random variables. Since

 $\psi_{m_{[u-1]}+j}$ are independent Beta random variables with parameters $\alpha_{m_{[u-1]}+j}$ and $\beta_{m_{[u-1]}+j}$ respectively,

$$\mathbb{P}_{m}\left(\mathrm{PU}_{m_{[n]}}^{(\mathrm{SL})}(\mathbf{1},\boldsymbol{\psi})\overset{\star}{\simeq} H\right) = \mathbb{E}_{m}\left[\prod_{s\in[2,m_{[n]}]}\psi_{s}^{p_{s}}(1-\psi_{s})^{q_{s}}\right] = \prod_{s\in[2,m_{[n]}]}\mathbb{E}_{m}\left[\psi_{s}^{p_{s}}(1-\psi_{s})^{q_{s}}\right]$$
$$= \frac{(\alpha_{2}+p_{2}-1)_{p_{2}}(\beta_{2}+q_{2}-1)_{q_{2}}}{(\alpha_{2}+\beta_{2}+p_{2}+q_{2}-1)_{p_{2}+q_{2}}}\prod_{s\in[3,m_{[n]}]}\frac{(\alpha_{s}+p_{s}-1)_{p_{s}}(\beta_{s}+q_{s}-1)_{q_{s}}}{(\alpha_{s}+\beta_{s}+p_{s}+q_{s}-1)_{p_{s}+q_{s}}}.$$

By definition, $p_{m_{[n]}} = q_{m_{[n]}} = 0$, so that the last term in the product is 1 and hence we obtain the result in (3.16).

In the next lemma, we derive an alternative expression for q_s defined in (3.14) which helps in understanding the RHS of (3.16):

LEMMA 3.8. The q_s defined in (3.14) can be represented as

(3.17)
$$q_s = \begin{cases} 0 & \text{if } s = 1, \\ d_1(H) - a_1 & \text{if } s = 2, \\ d_{[s-1]}(H) - a_{[2]} - 2(s-3) & \text{if } s \ge 3, \end{cases}$$

where

$$d_{[s-1]}(H) = \sum_{v \in [s-1]} d_v(H), \qquad \textit{and} \qquad a_{[2]} = a_1 + a_2.$$

PROOF. For s = 1, we observe that q_1 is 0 by definition and, for s = 2,

(3.18)
$$q_2 = \sum_{u \in (2, m_{[n]}]} \mathbb{1}_{[2 \in (v(u), u]]} = \sum_{u \in (2, m_{[n]}]} \mathbb{1}_{[v(u)=1]} = d_1(H) - a_1 .$$

Simplifying q_s for $s \ge 3$ gives

$$q_s = \sum_{u \in (m_{[2]}, m_{[n]}]} \mathbb{1}_{[s \in (v(u), u]]} = \sum_{u \in (2, m_{[n]}]} \mathbb{1}_{[s \in (v(u), m_{[n]}]]} - \sum_{u \in (2, m_{[n]}]} \mathbb{1}_{[s \in (u, m_{[n]}]]}$$

$$\begin{split} &= \sum_{u \in (2,m_{[n]}]} \mathbb{1}_{[v(u) \in [s-1]]} - \sum_{u \in (2,m_{[n]}]} \mathbb{1}_{[u \in [s-1]]} \\ &= \Big(\sum_{u \in [m_{[n]}]} \mathbb{1}_{[v(u) \in [s-1]]} - \sum_{u \in [2]} \mathbb{1}_{[v(u) \in [s-1]]}\Big) - \Big(\sum_{u \in [m_{[n]}]} \mathbb{1}_{[u \in [s-1]]} - \sum_{u \in [2]} \mathbb{1}_{[u \in [s-1]]}\Big) \\ &= \sum_{v \in [s-1]} \left(d_v^{(\text{in})}(H) - d_v^{(\text{in})}(G_0)\right) - \left((s-1) - 2\right), \end{split}$$

where $d_v^{(in)}(G)$ is the in-degree of vertex v in the graph G and G_0 is the initial graph we started with. Let $d_v^{(out)}(G)$ denote the out-degree of vertex v in the graph G, so that

$$d_v^{\text{(in)}}(G) + d_v^{\text{(out)}}(G) = d_v(G)$$

Note that the vertices in the initial graph G_0 do not have out-edges directed to any new incoming vertices. Therefore, $d_v^{(\text{out})}(H) = d_v^{(\text{out})}(G_0)$ for all $v \in [2]$.

On the other hand, the new incoming vertices have exactly one out-edge each. Furthermore they are not part of the initial graph. Hence, by definition $d_v(G_0) = 0$ for all $v > 2 = m_{[2]}$. Therefore, both $d_v^{(\text{out})}(G_0)$ and $d_v^{(\text{in})}(G_0)$ are zero for all $v > 2 = m_{[2]}$. Hence, for $s \ge 3$,

$$q_{s} = \sum_{v \in [s-1]} (d_{v}^{(\text{in})}(H) - d_{v}^{(\text{in})}(G_{0})) - ((s-1)-2)$$

$$= \sum_{v \in [s-1]} (d_{v}(H) - d_{v}(G_{0})) - \sum_{v \in [s-1]} (d_{v}^{(\text{out})}(H) - d_{v}^{(\text{out})}(G_{0})) - (s-3)$$

$$= \sum_{v \in [s-1]} (d_{v}(H) - d_{v}(G_{0})) - \sum_{v \in [2]} (d_{v}^{(\text{out})}(H) - d_{v}^{(\text{out})}(G_{0}))$$

$$(3.20) \qquad - \sum_{v \in (2,s-1]} d_{v}^{(\text{out})}(H) - (s-3)$$

$$= \sum_{v \in [s-1]} (d_{v}(H) - d_{v}(G_{0})) - 2(s-3)$$

$$= d_{[s-1]}(H) - a_{[2]} - 2(s-3).$$

Now, we have all the tools to prove Proposition 3.2:

PROOF OF PROPOSITION 3.2. We start by simplifying the RHS of (3.16) and equating term by term. First, we consider the term corresponding to s = 2 in the RHS of (3.16). Using (3.18) and substituting the values of α_2 and β_2 ,

(3.21)
$$\frac{(\alpha_2 + p_2 - 1)_{p_2}(\beta_2 + q_2 - 1)_{q_2}}{(\alpha_2 + \beta_2 + p_2 + q_2 - 1)_{p_2 + q_2}} = \frac{(\delta + d_2(H) - 1)_{d_2(H) - a_2}(d_1(H) + \delta - 1)_{d_1(G) - a_1}}{(d_{[2]}(G) + 2\delta - 1)_{d_{[2]}(G) - a_{[2]}}}$$

$$=\frac{1}{\left(d_{[2]}(H)+2\delta-1\right)_{d_{[2]}(H)-a_{[2]}}}\prod_{i=a_1}^{a_1(H)-1}(i+\delta)\prod_{i=a_2}^{a_2(H)-1}(i+\delta).$$

The factor $(\alpha_s + p_s - 1)_{p_s}$ in the numerator of (3.16) can be simplified as

(3.22)
$$(\alpha_s + p_s - 1)_{p_s} = \prod_{i=0}^{p_s - 1} = (\alpha_s - 1 + i) = \prod_{i=a_s}^{d_s(H) - 1} (i + \alpha_s - 1).$$

The last equality in (3.22) is obtained by substituting the expression for $p_{m_{[u-1]}+j}$ and a simple change of variables. For any $s \in [3, m_{[n]}]$, we can find a $u \in [3, n]$ and $j \in [m_u]$ such that $s = m_{[u-1]} + j$. Therefore,

(3.23)
$$\prod_{s \in [3,m_{[n]}]} (\alpha_s + p_s - 1)_{p_s} = \prod_{u \in [3,n]} \prod_{j \in [m_u]} \prod_{i=a_{m_{[u-1]}+j}}^{d_{m_{[u-1]}+j}(H)-1} \left(i + \frac{\delta}{m_u}\right).$$

So far, from (3.21) and (3.23), we have already obtained the first factor in (3.11) and an additional term in the denominator. Note that, for $s \ge 3$,

(3.24)
$$\alpha_s + \beta_s = \beta_{s+1} - 1$$
, and $p_s + q_s = q_{s+1} + 1$.

(3.25)
$$\prod_{s \in [3,m_{[n]})} \frac{(\beta_s + q_s - 1)_{q_s}}{(\alpha_s + \beta_s + p_s + q_s - 1)_{p_s + q_s}} = \prod_{s \in [3,m_{[n]})} \frac{(\beta_s + q_s - 1)_{q_s}}{(\beta_{s+1} + q_{s+1} - 1)_{q_{s+1}+1}}$$
$$= \prod_{s \in [3,m_{[n]})} \frac{1}{(\beta_{s+1} - 1)} \prod_{s \in [3,m_{[n]})} \frac{(\beta_s + q_s - 1)_{q_s}}{(\beta_{s+1} + q_{s+1} - 1)_{q_{s+1}}}.$$

Since $(\beta_3 + q_3 - 1)_{q_3} = (d_{[2]}(H) + 2\delta - 1)_{d_{[2]}(H) - a_{[2]}}$ and $q_{m_{[n]}} = 0$, we can simplify (3.25) as

$$\prod_{s \in [3,m_{[n]}]} \frac{(\beta_s + q_s - 1)q_s}{(\alpha_s + \beta_s + p_s + q_s - 1)p_{s+q_s}}$$

$$(3.26) = (d_{[2]}(H) + 2\delta - 1)_{d_{[2]}(H) - a_{[2]}}$$

$$\times \prod_{u \in [3,n]} \prod_{j \in [m_u]} \frac{1}{a_{[2]} + 2(m_{[u-1]} + j - 2) + (u - 1)\delta + \frac{j}{m_u}\delta - 1}.$$

Therefore, substituting these simplified expressions from (3.26) and (3.21) in (3.16), we obtain:

$$\mathbb{P}_{m}\left(\mathrm{PU}_{m_{[n]}}^{(\mathrm{SL})}(\mathbf{1},\boldsymbol{\psi}) \stackrel{\star}{\simeq} H\right)$$

$$= \prod_{u \in [1,n]} \prod_{j \in [m_{u}]} \prod_{i=a_{m_{[u-1]}+j}}^{d_{m_{[u-1]}+j}(H)-1} \left(i + \frac{\delta}{m_{u}}\right)$$

$$(3.27) \qquad \qquad \times \prod_{u \in [3,n]} \prod_{j \in [m_{u}]} \frac{1}{a_{[2]} + 2\left(m_{[u-1]} + j - 2\right) + (u-1)\delta + \frac{j}{m_{u}}\delta - 1},$$

which, as required, matches the expression of $\mathbb{P}_m\left(\mathrm{PA}_{m_{[n]}}^{\scriptscriptstyle(A)}(\boldsymbol{m},1,\delta) \stackrel{\star}{\simeq} H\right)$ in (3.11). \Box

3.2. Equivalence of models (B) and (D) with their respective Pólya urn descriptions. Following similar calculations, we can show that model (B) and (D) are equal in distribution to $\text{CPU}^{(\text{NSL})}$ and $\text{PU}^{(\text{NSL})}$, respectively. $\text{PA}_n^{(\text{B})}(\boldsymbol{m}, \delta)$ is equal in distribution to $\text{CPU}_n^{(\text{NSL})}(\boldsymbol{m}, \boldsymbol{\psi})$ where $\boldsymbol{\psi}$ are the Beta random variables defined in (3.1) and (3.2):

THEOREM 3.9 (Equivalence of model (B) and $CPU^{(NSL)}$). Conditionally on m, for any graph $G \in \mathcal{H}_n$,

(3.28)
$$\mathbb{P}_m\left(\mathrm{PA}_n^{\scriptscriptstyle(\mathrm{B})}(\boldsymbol{m},\boldsymbol{\delta})\overset{\star}{\simeq} G\right) = \mathbb{P}_m\left(\mathrm{CPU}_n^{\scriptscriptstyle(\mathrm{NSL})}(\boldsymbol{m},\boldsymbol{\psi})\overset{\star}{\simeq} G\right).$$

Next, we describe the equivalence of $PA_n^{(D)}(\boldsymbol{m}, \delta)$ and $PU_n^{(SL)}(\boldsymbol{m}, \boldsymbol{\psi})$ where $\boldsymbol{\psi}$ is a sequence of independent Beta random variables defined as

(3.29)
$$\begin{aligned} \psi_v &\sim \mathsf{Beta}\left(m_v + \delta, a_{[2]} + 2\left(m_{[v-1]} - 2\right) + m_v + (v-1)\delta\right) & \text{for } v \ge 3, \\ \psi_2 &\sim \mathsf{Beta}(a_2 + \delta, a_1 + \delta), \\ \text{and } \psi_1 = 1. \end{aligned}$$

THEOREM 3.10 (Equivalence of model (D) and $PU^{(NSL)}$). Conditionally on m, for any graph $G \in \mathcal{H}_n$,

(3.30)
$$\mathbb{P}_m\left(\mathrm{PA}_n^{\scriptscriptstyle(\mathrm{D})}(\boldsymbol{m},\delta) \stackrel{\star}{\simeq} G\right) = \mathbb{P}_m\left(\mathrm{PU}_n^{\scriptscriptstyle(\mathrm{NSL})}(\boldsymbol{m},\boldsymbol{\psi}) \stackrel{\star}{\simeq} G\right),$$

where ψ is the sequence of Beta random variables defined in (3.29).

The proofs of Theorem 3.9 and 3.10 follow in exactly the same way as that of Theorem 3.1, and we have included them in Appendix A.

REMARK 3.11 (Models (E) and (F)). We have now presented Pólya Urn representations for models (A), (B) and (D), but not for models (E) and (F). This is due to the fact that models (E) and (F) do not have such representations (except for m = 1). As a result, we handle these models using a coupling to model (D) in Section 6.

4. Preliminary Results. For analysing the convergence to the local limit of CPU, we need some analytical results of both CPU and RPPT. Some of the proofs here follow those in [8], while some include significant novel ideas. We explain how these results can be reproduced for CPU^(NSL) or PU at the end of this section. This section is organised as follows. In Section 4.1, we provide auxiliary results on expected values of random variables. In Section 4.2, we use these to analyze the asymptotics of the positions $S_k^{(n)}$ and for an effective coupling of Beta and Gamma variables. Finally, in Section 4.3, we use these results to study the asymptotics of the attachment probabilities, and we prove some regularity properties of the RPPT.

4.1. Preliminaries on expectations of random variables. Conditionally on m, let $(\chi_k)_{k\in\mathbb{N}}$ be a sequence of independent Gamma random variables with parameters $m_k + \delta$ and 1. Since M has finite p-th moment, the p-th moment of all $(\chi_k)_{k>1}$ are finite as well:

LEMMA 4.1. The random variables $(\chi_k)_{k\geq 1}$ have uniformly bounded *p*-th moment.

PROOF. Fix $u \in \mathbb{N}$. Let, conditionally on m, X_1, \ldots, X_{m_u} be independent Gamma random variables with parameters $1 + \delta/m_u$ and $2\mathbb{E}[M] + \delta$. Hence

$$\mathbb{E}_m[X_1^p] = \frac{\Gamma\left(1 + \frac{\delta}{m_u} + p\right)}{\Gamma\left(1 + \frac{\delta}{m_u}\right)} \le \frac{\Gamma(1 + |\delta| + p)}{A} < K,$$

where

$$A = \min\left\{\Gamma\left(1 + \left|\frac{\delta}{m}\right|\right), \Gamma\left(1 - \left|\frac{\delta}{m}\right|\right)\right\},\,$$

and $m = \inf \operatorname{supp}(M)$. By the triangle inequality, $\mathbb{E}_m[|X_1 - \mathbb{E}_m[X_1]|^p]$ is bounded and the upper bound is independent of u and m_u . By [31, Corollary 8.2], for X_1, \ldots, X_n i.i.d. mean 0 random variables with finite ℓ -th moment, there exists a constant $B_{\ell} < \infty$ depending only on ℓ such that

(4.1)
$$\mathbb{E}[(X_1 + \dots + X_n)^{\ell}] \le \begin{cases} B_{\ell} n \mathbb{E}[|X_1|^{\ell}] & \text{for } 1 \le \ell \le 2, \\ B_{\ell} n^{\ell/2} \mathbb{E}\left[|X_1|^{\ell/2}\right] & \text{for } \ell > 2. \end{cases}$$

Since $\chi_k = X_1 + \cdots + X_{m_k}$ and X_1, \ldots, X_{m_k} are i.i.d. random variables, there exists B_p finite constant depending only on p such that

(4.2)
$$\mathbb{E}_m[|\chi_k - \mathbb{E}_m[\chi_k]|^p] \le B_p m_k^p \mathbb{E}_m[|X_1 - \mathbb{E}_m[X_1]|^p] \le C_1 m_k^p,$$

for some $C_1 > 0$. Using the triangle inequality for the L_p -norm,

(4.3)
$$\mathbb{E}_m\left[\chi_k^p\right] \le \mathbb{E}_m\left[\chi_k\right]^p + C_1 m_k^p \le C m_k^p,$$

for some C > 0. Since we have assumed the existence of the *p*-th moment of M, (4.3) implies that $\mathbb{E} \left[\chi_k^p \right]$ is uniformly bounded from above.

4.2. *Position concentration and Gamma-Beta couplings.* From (2.7) we obtain the conditional edge-probabilities for $CPU^{(SL)}$ as

(4.4)
$$\mathbb{P}_{m,\psi}\left(u \stackrel{j}{\leadsto} v \text{ in } \operatorname{CPU}_{n}^{(\mathrm{SL})}\right) = \begin{cases} \frac{S_{v}^{(n)}}{S_{u,j}^{(n)}} \left(1 - \prod_{l \in [m_{v}]} \left(1 - \psi_{m_{[v-1]+l}}\right)\right) & \text{ for } u > v, \\ 1 - \prod_{l \in [j]} \left(1 - \psi_{m_{[v-1]+l}}\right) & \text{ for } u = v. \end{cases}$$

Similarly, from (2.8), the conditional edge-probabilities for CPU^(NSL) are given by

(4.5)
$$\mathbb{P}_{m,\psi}\left(u \stackrel{j}{\rightsquigarrow} v \text{ in } \operatorname{CPU}_{n}^{(\mathrm{NSL})}\right) = \begin{cases} \frac{S_{v}^{(n)}}{S_{u,j-1}^{(n)}} \left(1 - \prod_{l \in [m_{v}]} \left(1 - \psi_{m_{[v-1]+l}}\right)\right) & \text{ for } u > v, \\ 1 - \prod_{l \in [j-1]} \left(1 - \psi_{m_{[v-1]+l}}\right) & \text{ for } u = v, \end{cases}$$

and for $PU^{\scriptscriptstyle\rm (NSL)}$ it is

(4.6)
$$\mathbb{P}_{m,\psi}\left(u \stackrel{j}{\rightsquigarrow} v \text{ in } \mathrm{PU}_{n}^{(\mathrm{NSL})}\right) = \psi_{v} \frac{\mathcal{S}_{v}^{(n)}}{\mathcal{S}_{u-1}^{(n)}}$$

We thus need to analyze the asymptotics of the $S_k^{(n)}$ of $CPU_n^{(SL)}(\boldsymbol{m}, \boldsymbol{\psi})$ with $\boldsymbol{\psi}$ defined in (3.1) and (3.2), which we do in the following proposition:

PROPOSITION 4.2 (Position Concentration for PU and CPU). Recall that $\chi = \frac{\mathbb{E}[M] + \delta}{2\mathbb{E}[M] + \delta}$. Then, for every $\varepsilon, \omega > 0$, there exists $K < \infty$ such that for all n > K and with probability at least $1 - \varepsilon$, for both PU and CPU,

(4.7)
$$\max_{k\in[n]} \left| \mathcal{S}_k^{(n)} - \left(\frac{k}{n}\right)^{\chi} \right| \le \omega,$$

and

(4.8)
$$\max_{k \in [n] \setminus [K]} \left(\frac{k}{n}\right)^{-\chi} \left| \mathcal{S}_k^{(n)} - \left(\frac{k}{n}\right)^{\chi} \right| \le \omega.$$

The proof to Proposition 4.2 is an adaptation of the proof of [8, Lemma 3.1] by Berger et al. to our setting, see Appendix B.

In collapsed Pólya urn graphs, the sequence $\{S_{k,j}^{(n)}: j \in [m_k]\}$ defined in (2.6) is an increasing sequence and from Proposition 4.2, with probability at least $1 - \varepsilon$, for all n > k > K,

(4.9)

$$\begin{aligned} \mathcal{S}_{k,j}^{(n)} \leq \mathcal{S}_{k}^{(n)} \leq \left(\frac{k}{n}\right)^{\chi} (1+2\omega), \\ \text{and} \qquad \mathcal{S}_{k,j}^{(n)} \geq \mathcal{S}_{k-1}^{(n)} \geq \left(\frac{k-1}{n}\right)^{\chi} (1-\omega) \geq \left(\frac{k}{n}\right)^{\chi} (1-2\omega).
\end{aligned}$$

Therefore it is evident that k for sufficiently large, the variation in $j \mapsto S_{k,j}^{(n)}$ is minor. Hence the differences between the SL and NSL versions of the collapsed Pólya Urn graphs will also be minor. The following proposition provides us with a nice coupling between the Beta random variables ψ and a sequence of Gamma variables. This coupling would be very much useful in analysing the remaining terms in (4.4), (4.5) and (4.6). Before diving into the proposition and its proof, let us denote $p = 1 + \rho$ for some $\rho > 0$.

PROPOSITION 4.3 (Beta-Gamma Coupling for PU and CPU). Consider the sequence $(\psi_k)_{k\in\mathbb{N}}$ for CPU and define the sequence $(\phi_k)_{k\in\mathbb{N}}$ as

(4.10)
$$\phi_v := \sum_{j \in [m_v]} \psi_{m_{[v-1]}+j}$$

Let, conditionally on m, Γ_k has a Gamma distribution with parameters $m_k + \delta$ and $2\mathbb{E}[M] + \delta$ δ . Then, there exist K_{ε} and K_{η} such that

- (i) with probability at least 1ε , $\chi_k \leq k^{1-\varrho/2}$ for all $k \geq K_{\varepsilon}$;
- (ii) consider the function $h_k^{\phi}(x)$ such that $\mathbb{P}(\phi_k \leq h_k^{\phi}(x)) = \mathbb{P}(\chi_k \leq x)$, where conditionally on \mathbf{m} , χ_k has a Gamma distribution with parameters $m_k + \delta$ and 1. Then, for every $\eta \in (0, \frac{1}{2})$, there exists sufficiently large $K_{\eta} \geq 1$ depending on η , such that, for all $k \geq K_{\eta}$ and $x < k^{1-\varrho/2}$.

(4.11)
$$\frac{1-\eta}{k(2\mathbb{E}[M]+\delta)}x \le h_k^{\phi}(x) \le \frac{1+\eta}{k(2\mathbb{E}[M]+\delta)}x$$

Since $(\psi_k)_{k>1}$ is the PU analogue of the sequence $(\phi_k)_{k>1}$, replacing $(\phi_k)_{k>1}$ by the corresponding $(\psi_k)_{k>1}$ of PU defined in (3.29), a similar Beta-Gamma coupling holds for PU.

Berger et al. [8, Lemma 3.2] provide a related result for degenerate M, but our proof technique is a bit different. For obtaining the upper and lower bounds in (4.11), we use a correlation inequality [40, Lemma 1.24] and Chernoff's inequality.

PROOF. First, we start by proving (i). Using Markov's inequality,

(4.12)
$$\mathbb{P}\left(\chi_k \ge k^{1-\varrho/2}\right) = \mathbb{P}\left(\chi_k^{1+\varrho} \ge k^{(1-\varrho/2)(1+\varrho)}\right) \le \mathbb{E}\left(\chi_k^{1+\varrho}\right) k^{-(1+\varrho(1-\varrho)/2)}$$

By Lemma 4.1, $\mathbb{E}(\chi_k^p)$ is uniformly bounded. Therefore the RHS of (4.12) is summable and we obtain (*i*) using the Borel-Cantelli lemma.

We proceed to prove (ii). Let

$$b_{u,j} = a_{[2]} + 2(m_{[u-1]} + j - 3) + (u - 1)\delta + \frac{(j-1)}{m_u}\delta - 1,$$

and, conditionally on m, let $(\chi'_k)_{k\in\mathbb{N}}$ be a sequence of independent Gamma variables such that

(4.13)
$$\chi'_{m_{[u-1]}+j} \sim \operatorname{Gamma}\left(1 + \delta/m_u, 1\right)$$

The function $x \mapsto h_{u,j}^{\phi}(x)$ is defined such that for all $x \leq b_{u,j}^{1-\varrho/2}$,

$$\mathbb{P}\left(\psi_{m_{[u-1]}+j} \le h_{u,j}^{\phi}(x)\right) = \mathbb{P}\left(\chi'_{m_{[u-1]+j}} \le x\right).$$

To prove part (*ii*), we start with the sequence $(\psi_k)_{k\in\mathbb{N}}$ with the aim to couple it with appropriately scaled $(\chi'_k)_{k\in\mathbb{N}}$. We will prove (4.11) using the following claim:

CLAIM 4.4. There exists $K_n \ge 1$ such that for all $x \le (b_{u,j})^{1-\varrho/2}$ and $u > K_n$,

(4.14)
$$\left(1-\frac{\eta}{2}\right)\frac{x}{b_{u,j}} \le h_{u,j}^{\phi}(x) \le \frac{x}{b_{u,j}}$$

Proof of part (*ii*) subject to Claim 4.4. From Claim 4.4 and part (*i*), there exists a K_{η} such that for any $u > K_{\eta}$, we can couple $(\psi_{m_{[u-1]}+j})_{\substack{u>K_{\eta}\\j\in[m_u]}}$ and $(\chi'_{m_{[u-1]}+j})_{\substack{u>K_{\eta}\\j\in[m_u]}}$ with probability at least $1 - \varepsilon$, such that

(4.15)
$$\left(1 - \frac{\eta}{2}\right)\psi_{m_{[u-1]}+j} \le \frac{\chi'_{m_{[u-1]}+j}}{b_{u,j}} \le \psi_{m_{[u-1]}+j}.$$

Since $b_{u,j}$ is random, we replace it with its expectation and encounter some error. By the law of large numbers,

(4.16)
$$\frac{m_{[u-1]}}{u} \stackrel{a.s.}{\to} \mathbb{E}[M],$$

and hence, depending on η , there exists $K_{\eta} \ge 1$ large enough such that, with probability at least $1 - \eta/2$, the error can be bounded as

(4.17)
$$\left|\frac{b_{u,j} - u(\mathbb{E}[M] + \delta)}{u(\mathbb{E}[M] + \delta)}\right| \le \frac{\eta}{2}, \quad \text{for all } u > K_{\eta}.$$

Therefore, from (4.15) and (4.17), with probability at least $1-2\varepsilon$, the sequences $\left(\psi_{m_{[u-1]}+j}\right)_{\substack{u>K_{\eta}\\ j\in m_{[u]}}}$ and $\left(\gamma'$ can be coupled such that

and
$$\left(\chi'_{m_{[u-1]}+j}\right)_{\substack{u>K_{\eta}\\j\in m_{[u]}}}$$
 can be coupled such that

(4.18)
$$(1-\eta)\,\psi_{m_{[u-1]}+j} \le \frac{\chi_{m_{[u-1]}+j}}{u(2\mathbb{E}[M]+\delta)} \le (1+\eta)\psi_{m_{[u-1]}+j}.$$

Thus, summing over all $j \in [m_u]$, we obtain (4.11) subject to (4.14). We are left to proving Claim 4.4:

Proof of Claim 4.4. To prove the claim it is enough to show that for all $x \leq (b_{u,j})^{-\varrho/2}$ and $u > K_\eta \geq 1/\sqrt{\eta}$,

(4.19)
$$\mathbb{P}_m\left(\psi_{m_{[u-1]}+j} \le (1-\eta)x\right) \le \mathbb{P}_m\left(\chi'_{m_{[u-1]}+j} \le b_{u,j}x\right) \le \mathbb{P}_m\left(\psi_{m_{[u-1]}+j} \le x\right).$$

Upper bound in (4.19). With $\alpha_u = 1 + \delta/m_u$, we bound

$$\mathbb{P}_m\left(\chi'_{m_{[u-1]}+j} \le b_{u,j}x\right) = \frac{\int\limits_{0}^{b_{u,j}x} y^{\alpha_u - 1}e^{-y} \, dy}{\int\limits_{0}^{\infty} y^{\alpha_u - 1}e^{-y} \, dy} \le \frac{\int\limits_{0}^{b_{u,j}x} y^{\alpha_u - 1}e^{-y} \, dy}{\int\limits_{0}^{0} y^{\alpha_u - 1}e^{-y} \, dy}.$$

Then, using a change of variables and adjusting the missing factors from the Beta density, we get

$$(4.20) \qquad \mathbb{P}_m\left(\chi'_{m_{[u-1]}+j} \le b_{u,j}x\right) \le \frac{\int\limits_0^1 \mathbbm{1}_{\{y \le x\}} e^{-b_{u,j}y} \left(1-y\right)^{-b_{u,j}} f_{m_{[u-1]}+j}(x) \, dx}{\int\limits_0^1 e^{-b_{u,j}y} \left(1-y\right)^{-b_{u,j}} f_{m_{[u-1]}+j}(x) \, dx},$$

where f_k is the density of ψ_k .

Note that the numerator is $\mathbb{E}_m \left[\mathbb{1}_{[\psi_{m_{[u-1}+j} \leq x]} e^{-b_{u,j}\psi_{m_{[u-1}+j}} \left(1 - \psi_{m_{[u-1}+j}\right)^{-b_{u,j}}\right]$ and the denominator is $\mathbb{E}_m \left[e^{-b_{u,j}\psi_{m_{[u-1}+j}} \left(1 - \psi_{m_{[u-1}+j}\right)^{-b_{u,j}}\right]$. Further, $z \mapsto \mathbb{1}_{\{z \leq x\}}$ is non-increasing,

while $z \mapsto e^{-b_{u,j}z} (1-z)^{-b_{u,j}}$ is increasing. Recall that, for increasing f and decreasing g on the support of X,

(4.21)
$$\mathbb{E}[f(X)g(X)] \le \mathbb{E}[f(X)]\mathbb{E}[g(X)].$$

Therefore, using this correlation inequality in the RHS of (4.20), we have

(4.22)
$$\mathbb{P}_m\left(\chi'_{m_{[u-1]}+j} \le b_{u,j}x\right) \le \mathbb{P}_m\left(\psi_{m_{[u-1]}+j} \le x\right),$$

which proves the upper bound in (4.19).

The lower bound. The lower bound in (4.19) can be handled in a similar, albeit slightly more involved, way. Indeed,

$$\mathbb{P}_m\left(\psi_{m_{[u-1]}+j} \le x(1-\eta)\right) = \frac{\int\limits_0^{x(1-\eta)b_{u,j}} y^{a_u-1} \left(1-\frac{y}{b_{u,j}}\right)^{b_{u,j}} dy}{\int\limits_0^{b_{u,j}} y^{a_u-1} \left(1-\frac{y}{b_{u,j}}\right)^{b_{u,j}} dy}$$

(4.23)

$$= \frac{\mathbb{E}_{m} \left[\mathbbm{1}_{\left\{\chi'_{m_{[u-1]}+j} \le b_{u,j} x(1-\eta)\right\}} e^{\chi'_{m_{[u-1]}+j}} \left(1 - \frac{\chi'_{m_{[u-1]}+j}}{b_{u,j}}\right)^{b_{u,j}} \middle| \chi'_{m_{[u-1]}+j} \le b_{u,j} \right]} \\ \mathbb{E}_{m} \left[e^{\chi'_{m_{[u-1]}+j}} \left(1 - \frac{\chi'_{m_{[u-1]}+j}}{b_{u,j}}\right)^{b_{u,j}} \middle| \chi'_{m_{[u-1]}+j} \le b_{u,j} \right]}$$

Denote $f(z) = \mathbb{1}_{\{z \le b_{u,j}x(1-\eta)\}}$ and $g(z) = e^z \left(1 - \frac{z}{b_{u,z}}\right)^{b_{u,j}}$. Note that f and g are non-increasing and increasing functions, respectively. Therefore, by the correlation inequality in (4.21) once more,

$$(4.24) \quad \mathbb{P}_m\left(\psi_{m_{[u-1]}+j} \le x(1-\eta)\right) \le \mathbb{P}_m\left(\chi'_{m_{[u-1]}+j} \le b_{u,j}x(1-\eta) \left|\chi'_{m_{[u-1]}+j} \le b_{u,j}\right)\right) \,.$$

Thus, to prove the required inequality, it suffices to show that, for all $u \ge K_{\eta}$ and $j \in m_{[u]}$, and $x \le (b_{u,j})^{-\varrho/2}$

$$(4.25) \qquad \mathbb{P}_m\left(\chi'_{m_{[u-1]}+j} \le b_{u,j}x(1-\eta) \middle| \chi'_{m_{[u-1]}+j} \le b_{u,j}\right) \le \mathbb{P}_m\left(\chi'_{m_{[u-1]}+j} \le b_{u,j}x\right).$$

To simplify notation, instead of showing (4.25), we prove the same for any Gamma random variable Z having parameters α and 1, with $\alpha \in (0, 1 + |\delta|)$. Observe that showing (4.25) is equivalent to showing,

(4.26)
$$E(x) = \mathbb{P}\left(Z \le bx\right) \mathbb{P}\left(Z \le b\right) - \mathbb{P}\left(Z \le bx(1-\eta)\right) \ge 0,$$

with $x \leq b^{-\varrho/2}$ for large enough $b > K_{\eta}$.

By definition, E(0) = 0. Therefore, it remains to show that (4.26) holds for $x \in (0, b^{-\varrho/2})$. We simplify

(4.27)
$$E(x) = \mathbb{P}\left((1-\eta)bx \le Z \le bx\right) - \mathbb{P}(Z > b)\mathbb{P}(Z \le bx).$$

Lower bounding the first expression of E(x). Using the properties of the density of the Gamma distribution, we can lower bound the first term as

$$\mathbb{P}\left((1-\eta)bx \le Z \le bx\right) \ge \eta bx \min\left\{\frac{(bx)^{\alpha-1}}{\Gamma(\alpha)}e^{-bx}, \frac{(bx)^{\alpha-1}}{\Gamma(\alpha)}e^{-bx}(1-\eta)^{\alpha-1}e^{bx\eta}\right\}$$

$$= \eta \frac{(bx)^{\alpha}}{\Gamma(\alpha)}e^{-bx}\min\left\{1, (1-\eta)^{\alpha-1}e^{bx\eta}\right\}.$$

With the fact that $\eta < \frac{1}{2}$, and $e^{\eta bx} > 1$, we can lower bound the first term in (4.27) further by

(4.29)
$$\mathbb{P}\left((1-\eta)bx \le Z \le bx\right) \ge \eta \frac{(bx)^{\alpha}}{\Gamma(\alpha)} e^{-bx} \min\{2^{1-\alpha}, 1\} \ge \eta \frac{(bx)^{\alpha}}{\Gamma(\alpha)} e^{-bx} 2^{1-\lceil\alpha\rceil}.$$

Upper bounding the second term of E(x). By Chernoff's inequality,

(4.30)
$$\mathbb{P}(Z > b) = \mathbb{P}\left(e^{\frac{1}{2}Z} > e^{\frac{b}{2}}\right) \le \mathbb{E}\left[e^{\frac{1}{2}Z}\right]e^{-\frac{b}{2}} = 2^{\alpha}e^{-\frac{b}{2}}.$$

We use the fact that $e^{-y} \leq 1$ for $y \geq 0$, for upper bounding the distribution function of Z as

(4.31)
$$\mathbb{P}\left(Z \le bx\right) = \frac{1}{\Gamma(\alpha)} \int_{0}^{bx} y^{\alpha-1} e^{-y} \, dy \le \frac{1}{\Gamma(\alpha)} \int_{0}^{bx} y^{\alpha-1} \, dy = \frac{(bx)^{\alpha}}{\alpha \Gamma(\alpha)}$$

Therefore, the second term in (4.27) can be upper bounded using (4.31) and (4.30), as

(4.32)
$$\mathbb{P}(Z > b)\mathbb{P}(Z \le bx) \le 2^{\alpha} e^{-\frac{b}{2}} \frac{(bx)^{\alpha}}{\alpha \Gamma(\alpha)}.$$

Hence from (4.29) and (4.32), we obtain (4.33)

$$E(x) \ge \eta \frac{(bx)^{\alpha}}{\Gamma(\alpha)} e^{-bx} 2^{1-\lceil \alpha \rceil} - 2^{\alpha} e^{-\frac{b}{2}} \frac{(bx)^{\alpha}}{\alpha \Gamma(\alpha)} = \frac{(bx)^{\alpha}}{\Gamma(\alpha)} e^{-\frac{b}{2}} \left[e^{\left(\frac{1}{2} - x\right)b + \log \eta} 2^{1-\lceil \alpha \rceil} - \frac{2^{\alpha}}{\alpha} \right].$$

Remember that $\eta > b^{-2}$ and, for sufficiently large $b, x \le b^{-\varrho/2} \le \frac{1}{4}$. Therefore, by (4.33), for $b \ge K_{\eta}$,

(4.34)
$$E(x) \ge \frac{(bx)^{\alpha}}{\Gamma(\alpha)} e^{-\frac{b}{2}} \left[2^{1-\lceil \alpha \rceil} e^{\frac{b}{4} - 2\log b} - \frac{2^{\alpha}}{\alpha} \right].$$

Using $\alpha \in (0, 1 + |\delta|)$ and taking *b* large enough, the RHS of (4.34) is non-negative and this proves (4.26). This completes the proof of coupling for CPU.

The proof for PU is similar. Indeed, $b_{u,j}$ is replaced by b_u given by

(4.35)
$$b_u = a_{[2]} + 2\left(m_{[u-1]} - 2\right) + m_u + (u-1)\delta.$$

Now we show that there exists $K_{\eta} \ge 1$ such that for all $u > K_{\eta}$

(4.36)
$$\left(1-\frac{\eta}{2}\right)\frac{x}{b_u} \le h_u^{\psi}(x) \le \frac{x}{b_u}.$$

To prove this we need to show that for $u > K_{\eta} \ge \frac{1}{\sqrt{\eta}}$, and $x \le b_u^{-\varrho/2}$,

$$\mathbb{P}_m\left(\psi_u \le (1-\eta)x\right) \le \mathbb{P}_m\left(\chi_u \le b_u x\right) \le \mathbb{P}_m\left(\psi_u \le x\right),$$

which we have already proved.

4.3. Asymptotics of attachment probabilities and regularity of the RPPT. The coupling arguments in Proposition 4.3 give us a sequence of conditional Gamma random variables $(\hat{\chi}_v)_{v\geq 2}$ corresponding to $(\phi_v)_{v\geq 3}$ such that $\hat{\chi}_v = (2\mathbb{E}[M] + \delta)v\phi_v$ for v large enough (where we assume that $v > K_\eta$). Similarly, for Pólya urn graphs, we can couple $(\hat{\chi}_v)_{v\geq 2}$ and $(\psi_v)_{v\geq 2}$. These relations are crucial to bound the edge-connection probabilities in (4.4) and (4.5), where we recall that $\chi = \frac{\mathbb{E}[M] + \delta}{2\mathbb{E}[M] + \delta}$:

LEMMA 4.5 (Bound on attachment probabilities). For $n \in \mathbb{N}$, consider $v \in [n]$. Then, for every $\varepsilon > 0$ there exists $\omega > 0$ such that, for both CPU and PU, with probability larger than $1 - \varepsilon$, for every $k \ge v \ge K_{\omega}$ and $j \in [m_k]$,

$$(1-\omega)\frac{\hat{\chi}_{v}}{(2\mathbb{E}[M]+\delta)v}\left(\frac{v}{k}\right)^{\chi} \leq \mathbb{P}_{m,\psi}\left(k \stackrel{j}{\leadsto} v\right) \leq (1+\omega)\frac{\hat{\chi}_{v}}{(2\mathbb{E}[M]+\delta)v}\left(\frac{v}{k}\right)^{\chi},$$

where $\hat{\chi}_v$ is a Gamma random variable with parameters $m_v + \delta$ and 1.

PROOF. Fix $n \in \mathbb{N}$ and $\varepsilon, \omega > 0$. The edge-connection probabilities for $CPU^{(SL)}$, $CPU^{(NSL)}$ and $PU^{(NSL)}$ are calculated in (4.4),(4.5) and (4.6), respectively. By Proposition 4.2, with probability at least $1 - \varepsilon/3$,

(4.37)
$$\max_{j \in [m_k]} \left| \frac{\mathcal{S}_v^{(t)}}{\mathcal{S}_{(k,j)}^{(t)}} - \left(\frac{v}{k}\right)^{\chi} \right| \le \left(\frac{\omega}{2}\right) \left(\frac{v}{k}\right)^{\chi},$$

where the value χ comes from the collapsed Pólya Urn graph model.

We next control the remaining terms in the edge-connection probabilities of (4.4), (4.5) and (4.6). For CPU, we rewrite the remaining expression as

(4.38)
$$1 - \prod_{i=1}^{m_v} \left(1 - \psi_{m_{[v-1]}+i} \right) = \sum_{i=1}^{m_v} \psi_{m_{[v-1]}+i} + E_m(v) = \phi_v + E_m(v),$$

which implicitly defines the error term $E_m(v)$. On the other hand, for PU, the remaining expression turns out to be ψ_v . By the assumptions, we can apply Proposition 4.3 to couple the Gamma random variables to ϕ_v and ψ_v of CPU and PU, respectively. As a consequence, with probability at least $1 - \varepsilon/3$, simultaneously for all v sufficiently large,

$$\begin{pmatrix} 1 - \frac{\omega}{2} \end{pmatrix} \frac{\hat{\chi}_v}{2\mathbb{E}[M] + \delta} \le v\phi_v \le \left(1 + \frac{\omega}{2}\right) \frac{\hat{\chi}_v}{2\mathbb{E}[M] + \delta},$$

and
$$\begin{pmatrix} 1 - \frac{\omega}{2} \end{pmatrix} \frac{\hat{\chi}_v}{2\mathbb{E}[M] + \delta} \le v\psi_v \le \left(1 + \frac{\omega}{2}\right) \frac{\hat{\chi}_v}{2\mathbb{E}[M] + \delta},$$

for CPU and PU, respectively, where $\hat{\chi}_v$ is a sequence of independent Gamma distribution with parameters $m_v + \delta$ and 1.

The term $E_m(v)$ in (4.38) is bounded by

(4.39)
$$\sum_{\substack{i\neq j\\i,j\in[m_v]}}\psi_{m_{[v-1]}+i}\psi_{m_{[v-1]}+j} \le \left(\sum_{i\in[m_v]}\psi_{m_{[v-1]}+i}\right)^2 \le (1+2\omega)\left(\frac{\hat{\chi}_v}{v}\right)^2.$$

By Proposition 4.2, $\hat{\chi}_v \leq v^{1-\varrho/2}$ for all $v > K_{\varepsilon}$, with probability at least $1 - \varepsilon$. Therefore, by (4.39),

(4.40)
$$E_m(v) \le \frac{\hat{\chi}_v}{v(2\mathbb{E}[M] + \delta)} O\left(v^{-\varrho/2}\right)$$

This completes the proof.

REMARK 4.6. Observe that the edge-connection probabilities are essentially the same for the three models: $CPU_n^{(SL)}$, $CPU_n^{(NSL)}$ and $PU_n^{(NSL)}$, except for the self-loop creation probability. The latter is insignificant for large graphs. Using Lemma 4.5, we approximate the attachment probabilities of all these models by the same expression, with an error that can be effectively taken care of.

We close this section by recalling a regularity property of the RPPT that is similar to [8, Lemma 3.3] and which is useful in Section 5. We provide a proof to this in the appendix:

LEMMA 4.7 (Regularity of RPPT). Fix $r \ge 0$ and $\varepsilon > 0$. Then there exist $K, C < \infty$ and $\eta(\varepsilon, r) > 0$ such that, with probability at least $1 - \varepsilon$,

 $\begin{array}{ll} 1. \ A_{\omega} \geq \eta(\varepsilon,r), \mbox{ for all } \omega \in B_r^{\scriptscriptstyle (G)}(\varnothing); \\ 2. \ |B_r^{\scriptscriptstyle (G)}(\varnothing)| \leq C; \\ 3. \ \Gamma_{\omega} \leq K, \mbox{ for all } \omega \in B_r^{\scriptscriptstyle (G)}(\varnothing). \end{array}$

5. Local Convergence. We prove local convergence for the vertex-marked preferential attachment model (A). For any finite vertex-marked tree t, with vertex marks in [0, 1], let V(t) denote the set of vertices of t and $\{a_{\omega} \in [0,1] : \omega \in V(t)\}$ denote the age-set of the vertices. Fix $r \in \mathbb{N}$ and $\varepsilon > 0$. Let $G_n = PA_n^{(A)}(\boldsymbol{m}, \delta)$. If $B_r^{(G_n)}(v) \simeq t$ and v_{ω} is the vertex in G_n corresponding to vertex ω of t, then we define

$$N_{r,n}\left(\mathbf{t}, (a_{\omega})_{\omega \in V(\mathbf{t})}\right) = \sum_{v \in [n]} \mathbb{1}_{\left\{B_{r}^{(G_{n})}(v) \simeq \mathbf{t}, |v_{\omega}/n - a_{\omega}| \le 1/r \; \forall \omega \in V(\mathbf{t})\right\}},$$

where $B_r^{(G_n)}(v)$ is the *r*-neighbourhood of v in G_n . With $B_r^{(G)}(\emptyset)$ denoting the *r*-neighbourhood of the RPPT (M, δ) and A_{ω} the age in RPPT (M, δ) of the node ω of t, we aim to show that

(5.1)
$$\frac{N_{r,n}\left(\mathbf{t}, (a_{\omega})_{\omega \in V(\mathbf{t})}\right)}{n} \xrightarrow{\mathbb{P}} \mu\left(B_r^{(G)}(\varnothing) \simeq \mathbf{t}, |A_{\omega} - a_{\omega}| \le 1/r \; \forall \omega \in V(\mathbf{t})\right).$$

To prove (5.1), we use the second moment method, i.e., we prove $\mathbb{E}\left[N_{r,n}\left(t, (a_{\omega})_{\omega \in V(t)}\right)\right]/n$ converges to the limit and that the variance of $N_{r,n}/n$ vanishes for $n \to \infty$. Throughout this section, we consider $\eta = \eta(\varepsilon, r)$ as introduced in Lemma 4.7.

5.1. First Moment Convergence. Here we prove the first moment convergence. Let $(t, (a_{\omega})_{\omega \in V(t)})$ be a vertex-marked tree with marks $(a_{\omega})_{\omega \in V(t)}$ taking values in $[0, 1]^{|V(t)|}$. We compute

(5.2)
$$\frac{1}{n} \mathbb{E}\left[N_{r,n}\left(\mathbf{t}, (a_{\omega})_{\omega \in V(\mathbf{t})}\right)\right] = \mathbb{P}\left(B_r^{(G_n)}(o) \simeq \mathbf{t}, |v_{\omega}/n - a_{\omega}| \le 1/r \; \forall \omega \in V(t)\right),$$

where $o \in [n]$ is chosen uniformly at random. We aim to show that this converges to the RHS of (5.1).

Instead of proving the first moment convergence, we prove the stronger statement that the *age densities* of the vertices in the *r*-neighbourhood of a uniformly chosen vertex in $PA_n^{(A)}(\boldsymbol{m}, \delta)$ converges pointwise to the age density of the nodes in the *r*-neighbourhood of $RPPT(M, \delta)$.

Define $f_{r,t}((A_{\omega})_{\omega \in V(t)})$ as the density of the ages in the RPPT (M, δ) , when the ordered *r*-neighbourhood $B_r^{(G)}(\emptyset)$ is in the same equivalence class as t. Then,

(5.3)
$$\mu\left(B_r^{(G)}(\varnothing)\simeq \mathbf{t}, A_\omega\in da_\omega, \,\forall\omega\in V(\mathbf{t})\right) = f_{r,\mathbf{t}}\left((a_\omega)_{\omega\in V(\mathbf{t})}\right)\prod_{\omega\in V(\mathbf{t})} da_\omega$$

THEOREM 5.1 (First moment density convergence theorem). Fix $\delta > -\inf \operatorname{supp}(M)$, and consider $G_n = \operatorname{PA}_n^{(A)}(\boldsymbol{m}, \delta)$. Uniformly for all $a_{\omega} > \eta$, distinct v_{ω} , and $\hat{\chi}_{v_{\omega}} \leq (v_{\omega})^{1-\frac{\varrho}{2}}$, for all $\omega \in V(\mathfrak{t})$,

(5.4)

$$\mathbb{P}_{m,\psi}\left(B_{r}^{(G_{n})}(v) \simeq t, v_{\omega} = \lceil na_{\omega} \rceil \forall \omega \in V(t)\right) \\
= (1 + o_{\mathbb{P}}(1)) \frac{1}{n^{|V(t)|}} g_{r,t}\left((a_{\omega})_{\omega \in V(t)}; (m_{v_{\omega}}, \hat{\chi}_{v_{\omega}})_{\omega \in V(t)}\right),$$

for some measurable function $g_{r,t}\left((a_{\omega})_{\omega\in V(t)};(m_{v_{\omega}},\hat{\chi}_{v_{\omega}})_{\omega\in V(t)}\right)$, where $(\hat{\chi}_v)_{v\in\mathbb{N}}$ are the Gamma random variables coupled with the corresponding Beta random variables in Proposition 4.3. Consequently, with $(\hat{\chi}_{v_{\omega}})_{\omega\in V(t)}$ a conditionally independent sequence of Gamma $(m_{v_{\omega}} + \delta, 1)$ random variables,

(5.5)
$$\mathbb{E}\left[g_{r,t}\left(\left(a_{\omega}\right)_{\omega\in V(t)};\left(m_{v_{\omega}},\hat{\chi}_{v_{\omega}}\right)_{\omega\in V(t)}\right)\right]=f_{r,t}\left(\left(a_{\omega}\right)_{\omega\in V(t)}\right).$$

By (5.5), Theorem 5.1 can be seen as a *local density limit theorem* for the ages of the vertices in the r-neighbourhoods. This is significantly stronger than local convergence of preferential attachment models. Further, when $(\lceil na_{\omega} \rceil)_{\omega \in V(t)}$ are distinct (which occurs whp), $(\hat{\chi}_{\lceil na_{\omega} \rceil})_{\omega \in V(t)}$ are *conditionally independent* Gamma variables with parameters $m_{v_{\omega}} + \delta$ and 1.

We prove Theorem 5.1 below in several steps. First we calculate the conditional density $f_{r,t}$. Using the equivalence of CPU and model (A) in Proposition 3.2, we compute the explicit expression in (5.4). Let $\partial V(t)$ denotes the leaf nodes of the tree t and $V^{\circ}(t)$ the set of vertices in the interior of the tree t, i.e., $V^{\circ}(t) = V(t) \setminus \partial V(t)$. Then the age densities in RPPT (M, δ) are identified as follows:

PROPOSITION 5.2 (Law of vertex-marked neighbourhood of RPPT). Let M be the law of the out-degrees, and fix $\delta > -\inf \operatorname{supp}(M)$. Let n_{\circ} and n_{\vee} denote the number of \circ and \vee labelled nodes in t and E(t) the edge-set of the tree t. Then,

$$f_{r,t}\left((a_{\omega})_{\omega\in V(t)}\right) = \chi^{n_{0}}(1-\chi)^{n_{Y}}\mathbb{E}\left[\prod_{\omega\in V^{\circ}(t)}\left(m_{\omega}-\mathbb{1}_{\{\omega\ is\ Y\ \}}\right)!\Gamma^{d_{\omega}^{(\mathrm{in})}(t)}_{\omega}\exp\left(-\Gamma_{\omega}\lambda(a_{\omega})\right)\right]$$

$$\times\prod_{(\omega,\omega l)\in E(\mathrm{t})}\left(a_{\omega}\vee a_{\omega l}\right)^{-(1-\chi)}\left(a_{\omega}\wedge a_{\omega l}\right)^{-\chi}\right],$$

where $d_{\omega}^{(in)}(t) = \#\{\omega l : a_{\omega l} > a_{\omega}\}$ denotes the number of γ labelled children of ω , while $(\Gamma_{\omega}, m_{\omega})$ are distributed as in Section 1.4 and we recall from (1.16) that λ is a real-valued function on (0,1) defined as

(5.7)
$$\lambda(x) = \frac{1 - x^{1-\chi}}{x^{1-\chi}}.$$

To prove Proposition 5.2, we need to identify the densities of $(A_{\omega})_{\omega \in V(t)}$. We start by analysing the density of the age of 0 labelled nodes:

LEMMA 5.3 (Conditional age density of 0 labelled children). Conditionally on a_{ω} , the age of its parent $\omega \in V^{\circ}(t)$, the density of an 0 labelled children on $[0, a_{\omega}]$ is given by

(5.8)
$$f_{o}(x) = \chi x^{-(1-\chi)} a_{\omega}^{-\chi}.$$

PROOF. From the construction of RPPT, for any 0 labelled node ωl , its age is given by $U^{1/\chi}A_{\omega}$, where U is uniform in [0,1] independently of A_{ω} . Therefore, conditionally on $A_{\omega} = a_{\omega}$,

$$\mathbb{P}\left(A_{\omega l} \le x \mid A_{\omega} = a_{\omega}\right) = \mathbb{P}\left(A_{\omega}U^{1/\chi} \le x \mid A_{\omega} = a_{\omega}\right) = \mathbb{P}\left(U^{1/\chi} \le xa_{\omega}^{-1}\right) = a_{\omega}^{-\chi}x^{\chi}.$$

Now, differentiating with respect to x, we obtain the conditional density of $A_{\omega l}$ in (5.8).

The ages of the $\mathbf Y$ labelled children of ω follow an inhomogeneous Poisson process with intensity

(5.9)
$$\rho_{\omega}(x) = (1-\chi)\Gamma_{\omega}\frac{x^{-\chi}}{A_{\omega}^{1-\chi}}.$$

This leads to the following density result on the ages of all children of ω :

LEMMA 5.4 (Conditional age density of children). For any $\omega \in V^{\circ}(t)$, conditionally on $(m_{\omega}, A_{\omega}, \Gamma_{\omega})$, the density of the ages of the children of ω is given by

$$f_{\omega}\left((a_{\omega l})_{l \in d_{\omega}(\mathsf{t})} \middle| m_{\omega}, a_{\omega}, \Gamma_{\omega}\right) = m_{-}(\omega)! \prod_{l=1}^{m_{-}(\omega)} \left[\chi(a_{\omega l})^{\chi-1}(a_{\omega})^{-\chi}\right] \exp\left(-\Gamma_{\omega}\lambda(a_{\omega})\right)$$

(5.10)

$$\times \prod_{k=1}^{d_{\omega}^{(\mathrm{in})}(\mathrm{t})} \left[(1-\chi) \left(a_{\omega(m_{-}(\omega)+k)} \right)^{-\chi} (a_{\omega})^{\chi-1} \Gamma_{\omega} \right],$$

where $m_{-}(\omega) = m_{\omega} - \mathbb{1}_{\{\omega \text{ is } \mathsf{r}\}}$ and $d_{\omega}(\mathsf{t}) = m_{-}(\omega) + d_{\omega}^{(\mathrm{in})}(\mathsf{t})$ denotes the number of children of ω .

PROOF. From the construction of the RPPT, we see that γ labelled nodes have $m_{\omega} - 1$ many 0 labelled children, while the other nodes have m_{ω} many 0 labelled children. Since the uniform random variables are chosen independently for obtaining the age of the 0 labelled children, using Lemma 5.3,

(5.11)
$$f\left((a_{\omega l})_{l\in[m_{-}(\omega)]}\middle| m_{\omega}, a_{\omega}, \Gamma_{\omega}\right) = m_{-}(\omega)! \prod_{l=1}^{m_{-}(\omega)} \left[\chi(a_{\omega l})^{\chi-1}(a_{\omega})^{-\chi}\right].$$

Indeed, since there is no particular order for connecting to the older nodes, ω can connect to its older children with its edges in $m_{-}(\omega)!$ different ways.

The label Y children have ages coming from a Poisson process with (random) intensity $x \mapsto \rho_{\omega}(x)$ on $[a_{\omega}, 1]$ defined in (5.9). Therefore for $k \ge 2$, conditionally on $a_{\omega(m_{-}(\omega)+k-1)}, \omega(m_{-}(\omega)+k)$ has an age following a non-homogeneous exponential distribution with intensity $x \mapsto \rho_{\omega}(x)$.

Additionally, there is one more factor arising in this part of the density, which is the nofurther γ labelled child after $\omega d_{\omega}(t)$. Conditionally on $a_{\omega d_{\omega}(t)}$, this no-further child factor is given by

$$\exp\left(-\Gamma_{\omega}\frac{1-a_{\omega d_{\omega}(\mathbf{t})}^{1-\chi}}{a_{\omega}^{1-\chi}}\right).$$

This no-further child part is also independent of the ages of the γ labelled nodes except for $a_{\omega d_{\omega}(t)}$, from the property of the Poisson process. Therefore,

(5.12)

$$f\left(\left(a_{\omega(m_{-}(\omega)+l)}\right)_{l\in[d_{\omega}^{(\mathrm{in})}(\mathrm{t})]}\middle|m_{\omega},a_{\omega},\Gamma_{\omega}\right)$$

$$=\exp\left(-\Gamma_{\omega}\lambda(a_{\omega})\right)\prod_{k=m_{-}(\omega)+1}^{d_{\omega}(\mathrm{t})}\left[(1-\chi)\left(a_{\omega k}\right)^{-\chi}(a_{\omega})^{\chi-1}\Gamma_{\omega}\right].$$

Since the γ labelled nodes connect one by one sequentially in a particular order, ω connects to its γ labelled children in only 1 way. Now, from the construction, the edges to the 0 labelled

children and the edges to the Y labelled children are created independently (conditionally on $(m_{\omega}, A_{\omega}, \Gamma_{\omega})$). Therefore from (5.11) and (5.12) and this independence,

$$f_{\omega}\left((a_{\omega l})_{l\in[d_{\omega}(\mathbf{t})]} \middle| m_{\omega}, a_{\omega}, \Gamma_{\omega}\right)$$
(5.13)
$$= f\left((a_{\omega l})_{l\in[m_{-}(\omega)]} \middle| m_{\omega}, a_{\omega}, \Gamma_{\omega}\right) f\left(\left(a_{\omega(m_{-}(\omega)+l)}\right)_{l\in[d_{\omega}^{(\mathrm{in})}(\mathbf{t})]} \middle| m_{\omega}, a_{\omega}, \Gamma_{\omega}\right),$$
which leads to the required expression in (5.10).

which leads to the required expression in (5.10).

Now we have the required tools to prove Proposition 5.2, for which we use induction on r, the depth of the tree:

Proof of Proposition 5.2. For proving this proposition, we define some notation *that is* used in this proof only. Define t_r to be the r-neighbourhood of the root in t. Therefore,

 $V^{\circ}(\mathbf{t}_{r+1}) = V^{\circ}(\mathbf{t}_r) \cup \partial V(\mathbf{t}_r).$

First, we prove the proposition for r = 1. Here $V^{\circ}(t_1) = \{\emptyset\}$. By Lemma 5.4,

$$f_{\varnothing}\left((a_{\varnothing l})_{l\in[d_{\varnothing}(\mathsf{t})]}\middle| m_{\varnothing}, a_{\varnothing}, \Gamma_{\varnothing}\right) = (m_{\varnothing})! \prod_{l=1}^{m_{\varnothing}} \left[\chi(a_{\varnothing l})^{\chi-1}(a_{\varnothing})^{-\chi}\right]$$

(5.14)

$$\times \prod_{k=1}^{d_{\varnothing}^{(\mathrm{in})}(\mathrm{t})} \left[(1-\chi) \left(a_{\varnothing(m_{\varnothing}+k)} \right)^{-\chi} (a_{\varnothing})^{\chi-1} \Gamma_{\varnothing} \right] \exp\left(-\Gamma_{\varnothing} \lambda(a_{\varnothing})\right).$$

(:--)

Now, observe that for every edge to an o labelled child, we obtain a factor χ and for every edge to Y labelled children, we obtain a factor $(1 - \chi)$ and a factor Γ_{\emptyset} in (5.14). Further note that O labelled children of \emptyset have smaller age than a_{\emptyset} and Y labelled children have higher age than a_{\varnothing} . t_1 has m_{\varnothing} many 0 labelled nodes and $d_{\varnothing}^{(in)}(t)$ many Y labelled nodes. Therefore (5.14) can be rewritten as

$$f_{\varnothing}\left((a_{\varnothing l})_{l\in[d_{\varnothing}(\mathsf{t})]}\middle| m_{\varnothing}, a_{\varnothing}, \Gamma_{\varnothing}\right) = \chi^{n_{1}, \mathsf{o}}(1-\chi)^{n_{1}, \mathsf{v}}(m_{\varnothing})! \Gamma_{\varnothing}^{d_{\varnothing}^{(m)}(\mathsf{t})} \exp\left(-\Gamma_{\varnothing}\lambda(a_{\varnothing})\right)$$

$$(5.15) \qquad \qquad \times \prod_{(\varnothing, \varnothing l)\in E(\mathsf{t}_{1})} \left[(a_{\varnothing} \lor a_{\varnothing l})^{-\chi} (a_{\varnothing} \land a_{\varnothing l})^{-(1-\chi)}\right],$$

where $n_{1,o}$ and $n_{1,v}$ denote the number of o and y labelled nodes in t_1 . Since A_{\varnothing} has a uniform distribution on [0,1], and is independent of m_{\varnothing} and Γ_{\varnothing} ,

$$f_{1,t}\left((a_{\omega})_{\omega\in V(t)}\right) = \mathbb{E}\left[f_{1,t}\left(a_{\varnothing}, (a_{\varnothing l})_{l\in[d_{\varnothing}(t)]} \middle| m_{\varnothing}, \Gamma_{\varnothing}\right)\right] \\ = \mathbb{E}\left[f_{\varnothing}\left((a_{\varnothing l})_{l\in[d_{\varnothing}(t)]} \middle| m_{\varnothing}, a_{\varnothing}, \Gamma_{\varnothing}\right)\right] \\ (5.16) \qquad = \chi^{n_{1,0}}(1-\chi)^{n_{1,Y}}\mathbb{E}\left[\Gamma_{\varnothing}^{d_{\varnothing}^{(\mathrm{in})}(t)}\exp\left(-\Gamma_{\varnothing}\lambda(a_{\varnothing})\right) \right] \\ \times \prod_{(\varnothing, \varnothing l)\in E(t_{1})}\left[(a_{\varnothing} \lor a_{\varnothing l})^{-\chi}(a_{\varnothing} \land a_{\varnothing l})^{-(1-\chi)}\right]\right].$$

The first equality comes from the fact that $V^{\circ}(t_1) = \emptyset$ and hence the proposition is proved for r = 1. We proceed toward the induction step. Let (5.2) be true for $r = k \in \mathbb{N}$. We wish to show that the result holds true for r = k + 1.

We have the distribution for the ages of the nodes in $V(t_k)$. What remains is to compute the density of the boundary conditionally on the ages of $V(t_k)$. Now, the nodes in $\partial V(t_{k+1})$ are the children of the nodes in $\partial V(t_k)$ and their age-distribution is independent of the age of nodes in $V^{\circ}(t_k)$. By Lemma 5.4,

$$f\left((a_{\omega})_{\omega\in\partial V(\mathfrak{t}_{k+1})}\middle|(m_{u},a_{u},\Gamma_{u})_{u\in V(\mathfrak{t}_{k})}\right)$$

$$(5.17) \qquad =f\left((a_{\omega l})_{\omega l\in\partial V(\mathfrak{t}_{k+1})}\middle|(m_{\omega},a_{\omega},\Gamma_{\omega})_{\omega\in\partial V(\mathfrak{t}_{k})}\right)$$

$$=\chi^{n_{k+1,0}}(1-\chi)^{n_{k+1,\gamma}}\prod_{\omega\in\partial V(\mathfrak{t}_{k})}\left[(m_{-}(\omega))!\Gamma_{\omega}^{d_{\omega}^{(\mathrm{in})}(\mathfrak{t}_{k+1})}\exp\left(-\Gamma_{\omega}\lambda(a_{\omega})\right)\right]$$

$$\times\prod_{(\omega,\omega l)\in E(\mathfrak{t}_{k+1})}(a_{\omega}\vee a_{\omega l})^{-\chi}(a_{\omega}\wedge a_{\omega l})^{-(1-\chi)}\right],$$

where $n_{k+1,o}$ and $n_{k+1,v}$ denote the number of o labelled and v labelled nodes in $\partial V(t_{k+1})$. On the other hand, t_k is a rooted tree of depth k. Therefore, using induction on r, the depth of the tree from the root

$$f\left((a_{\omega})_{\omega\in V(\mathbf{t}_{k})}\right) = \chi^{n_{[k],\mathbf{0}}}(1-\chi)^{n_{[k],\mathbf{Y}}}\mathbb{E}\left[\prod_{\omega\in V^{\circ}(\mathbf{t}_{k})}(m_{-}\omega)!\Gamma^{d_{\omega}^{(\mathrm{in})}(t)}_{\omega}\exp\left(-\Gamma_{\omega}\lambda(a_{\omega})\right) \times \prod_{(\omega,\omega l)\in E(\mathbf{t}_{k})}(a_{\omega}\vee a_{\omega l})^{-(1-\chi)}(a_{\omega}\wedge a_{\omega l})^{-\chi}\right],$$
(5.18)

where $n_{[k],o}$ and $n_{[k],v}$ denote the number of O labelled and Y labelled nodes in $V(t_k)$. Since t is tree, $d_{\omega}^{(in)}(t_{k+1}) = d_{\omega}^{(in)}(t)$. Moreover $n_{[k],o} + n_{k+1,o} = n_{[k+1],o}$ and $n_{[k],v} + n_{k+1,v} = n_{[k+1],v}$ and $(m_{\omega}, \Gamma_{\omega})_{\omega \in V(t)}$ are independent random variables. Therefore from (5.17) and (5.18), we obtain the required result for r = k + 1 as

г

$$f_{k+1,t}\left((a_{\omega})_{\omega\in V(t)}\right) = \chi^{n_{[k+1],0}}(1-\chi)^{n_{[k+1],Y}} \mathbb{E}\left[\prod_{\omega\in V^{\circ}(t)} (m_{-}\omega)!\Gamma_{\omega}^{d_{\omega}^{(in)}(t)}\exp\left(-\Gamma_{\omega}\lambda(a_{\omega})\right)\right]$$

$$(5.19) \qquad \qquad \times \prod_{(\omega,\omega l)\in E(t_{k+1})} (a_{\omega}\vee a_{\omega l})^{-(1-\chi)}(a_{\omega}\wedge a_{\omega l})^{-\chi}\right].$$

The general claim follows by induction in r.

By Proposition 5.2, we have the exact expression for the density of the ages of the nodes in the RPPT. To prove Theorem 5.1, we compute the expression for $g_{r,t}$. Instead of finding $g_{r,t}$ directly, we use Theorem 3.1. To make our computation simpler, we first introduce edgemarks in the tree and then lift the edge-marks carefully.

Let $\overline{\mathbf{t}} = (\mathbf{t}, (a_{\omega})_{\omega \in V(\mathbf{t})}, (e_{\omega,\omega j})_{(\omega,\omega j) \in E(\mathbf{t})})$ be the edge-marked version of t, where $E(\mathbf{t})$ is the edge-set of t. We write $\overline{B}_r^{(G_n)}(v) \doteq \overline{\mathbf{t}}$, to denote that the vertex and edge-marks of the *r*-neighbourhood of the vertex *v* in $\operatorname{CPU}_n(\boldsymbol{m}, \boldsymbol{\psi})$ are given by those in t.

PROPOSITION 5.5 (Density of vertex and edge-marked CPU). Let o_n be a uniformly chosen vertex from $\operatorname{CPU}_n^{(SL)}(m, \psi)$. Then,

$$\mathbb{P}_{m,\psi}\left(\overline{B}_r^{(G_n)}(o_n) \doteq \overline{\mathbf{t}}, \ v_\omega = \lceil na_\omega \rceil, \ \forall \omega \in V(\mathbf{t})\right)$$

$$(5.20) = (1 + o_{\mathbb{P}}(1))n^{-|V(t)|} \prod_{\omega \in V^{\circ}(t)} \left[\left(\frac{\hat{\chi}_{v_{\omega}}}{2\mathbb{E}[M] + \delta} \right)^{d_{\omega}^{(in)}(t)} \exp\left(-\hat{\chi}_{v_{\omega}}\lambda(a_{\omega})\right) \right] \\ \times \prod_{(\omega,\omega l)\in E(t)} (a_{\omega} \vee a_{\omega l})^{-\chi} (a_{\omega} \wedge a_{\omega l})^{-(1-\chi)} \prod_{\substack{\omega \in V(t)\\\omega \text{ is o}}} \frac{\hat{\chi}_{v_{\omega}}}{2\mathbb{E}[M] + \delta} .$$

Before we prove Proposition 5.5 we begin with a lemma that gives an estimate of the edgeconnection probabilities and this estimate will be useful in the proof of the above proposition. The estimate is given in terms of the coupled Gamma random variables $(\hat{\chi}_v)_{v\geq 2}$ from Proposition 4.3.

LEMMA 5.6. Conditionally on
$$(\boldsymbol{m}, \boldsymbol{\psi})$$
, for all $\omega \in V^{\circ}(\mathbf{t})$,
(5.21)
$$\sum_{u,j: u \ge v_{\omega}} p^{(j)}(u, v_{\omega}) = (1 + o_{\mathbb{P}}(1))\hat{\chi}_{v_{\omega}}\lambda(a_{\omega}).$$

PROOF. For every u, there are m_u many out-edges from u and using the expression for the edge-connection probabilities in Lemma 4.5,

(5.22)

$$\sum_{u,j:u\geq v_{\omega}} p^{(j)}(u,v_{\omega}) = (1+o_{\mathbb{P}}(1)) \sum_{u\geq v_{\omega}} m_u \frac{\hat{\chi}_{v_{\omega}}}{(2\mathbb{E}[M]+\delta)v_{\omega}} \left(\frac{v_{\omega}}{u}\right)^{\chi}$$

$$= (1+o_{\mathbb{P}}(1)) \frac{\hat{\chi}_{v_{\omega}}}{(2\mathbb{E}[M]+\delta)v_{\omega}^{1-\chi}} \sum_{u\geq v_{\omega}} m_u u^{-\chi}$$

$$= (1+o_{\mathbb{P}}(1)) \frac{\hat{\chi}_{v_{\omega}}}{(2\mathbb{E}[M]+\delta) \left(\frac{v_{\omega}}{n}\right)^{1-\chi}} \left[\frac{1}{n} \sum_{u\geq v_{\omega}} m_u \left(\frac{u}{n}\right)^{-\chi}\right].$$

Define $T_n := \frac{1}{n} \sum_{u \ge v_{\omega}} m_u \left(\frac{u}{n}\right)^{-\chi}$. With $v_{\omega} = \lceil na_{\omega} \rceil$, we aim to show that

(5.23)
$$T_n \xrightarrow{\mathbb{P}} \mathbb{E}[M] \int_{a_\omega}^1 t^{-\chi} dt$$

using a *weighted* strong law of large numbers. Let $(X_i)_{i\geq 1}$ be a sequence of i.i.d. random variables with finite mean. Then from [16, Theorem 5], a sufficient condition for $\sum_{i=1}^{n} a_{i,n}X_i \quad \left(\text{with } \sum_{i=1}^{n} a_{i,n} = 1 \right)$ to converge to $\mathbb{E}[X]$ is that $\max_{i\in[n]} a_{i,n} = O(1/n)$. In our case, consider $a_{u,n} = \frac{\frac{1}{n} \left(\frac{u}{n}\right)^{-\chi}}{\frac{1}{n} \sum_{u \geq v_{ui}} \left(\frac{u}{n}\right)^{-\chi}}$, so that

(5.24)
$$\max_{u \in [v_{\omega}, n]} \frac{\frac{1}{n} \left(\frac{u}{n}\right)^{-\chi}}{\frac{1}{n} \sum_{u \ge v_{\omega}} \left(\frac{u}{n}\right)^{-\chi}} \le \frac{a_{\omega}^{-\chi}}{n(1 - a_{\omega}^{1 - \chi})} = O\left(\frac{1}{n}\right)$$

Therefore, the weighted strong law implies that (5.23) holds. Note that the denominator here is the Riemann sum approximation of $\int_{a_{\omega}}^{1} t^{-\chi} dt$, so that

(5.25)
$$T_n = (1 + o_{\mathbb{P}}(1))\mathbb{E}[M] \int_{a_{\omega}}^{1} t^{-\chi} dt = (1 + o_{\mathbb{P}}(1))\frac{\mathbb{E}[M]}{1 - \chi} \left[1 - a_{\omega}^{1 - \chi}\right].$$

Hence, from (5.22) and (5.25),

(5.26)
$$\sum_{u,j:u\geq v_{\omega}} p^{(j)}(u,v_{\omega}) = (1+o_{\mathbb{P}}(1)) \frac{\hat{\chi}_{v_{\omega}}}{(2\mathbb{E}[M]+\delta)a_{\omega}^{1-\chi}} \frac{\mathbb{E}[M]}{1-\chi} \left[1-a_{\omega}^{1-\chi}\right]$$
$$= (1+o_{\mathbb{P}}(1))\hat{\chi}_{v_{\omega}}\lambda(a_{\omega}),$$

as required.

Proof of Proposition 5.5. The proof of Proposition 5.5 is divided into several steps. Recall that Lemma 4.5 yields the edge-connection probabilities for CPU.

Computing the conditional law of $\overline{B}_r^{(G_n)}(o_n)$. The construction of the CPU implies the conditional independence of the edge-connection events. Conditionally on (m, ψ) , let $p^{(j)}(v, u)$ denote the probability of connecting the *j*-th edge from v to u. Then,

(5.27)
$$\mathbb{P}_{m,\psi}\left(\overline{B}_{r}^{(G_{n})}(o_{n}) \doteq \overline{\mathbf{t}}, v_{\omega} = \lceil na_{\omega} \rceil, \forall \omega \in V(\mathbf{t})\right)$$
$$= \frac{1}{n} \prod_{u \in V(\mathbf{t})} \prod_{\substack{\omega \in V(\mathbf{t}) \\ \omega > u \\ \omega \neq u}} p^{(j)}(v_{\omega}, v_{u}) \prod_{\omega \in V^{\circ}(\overline{\mathbf{t}})} \prod_{\substack{u,j: \\ (u,j,\omega) \notin E(\overline{\mathbf{t}})}} [1 - p^{(j)}(v_{u}, v_{\omega})].$$

The factor of 1/n arises due to the uniform choice of the root and the first product comprises all edge-connection probabilities to make sure that the edges in \overline{t} are there in $\overline{B}_r^{(G_n)}$ keeping the edge-marks the same. The last product ensures that there is no further edge in the (r - 1)-neighbourhood of the randomly chosen vertex o_n , so that the vertex and edge-marks of $\overline{B}_r^{(G_n)}(o_n)$ are same as those in \overline{t} .

The no-further edge probability. We continue by analyzing the second product on the RHS of (5.27) which, for simplicity, we call the *no-further edge probability*. Observe that in this part of the expression, we have not included the edges that are connected in t, i.e., we exclude those factors $[1 - p^{(j)}(v_u, v_\omega)]$ for which $\omega \in V^{\circ}(t)$ and $u \stackrel{j}{\rightarrow} \omega$ in t. Recall from Lemma 4.7 that the minimum age of the vertices in t is η whp. Therefore by Lemma 4.5, for all $u > \eta n$, $p^{(j)}(u, v)$ is $o_{\mathbb{P}}(1)$. Since t is finite, we exclude a finite number of $[1 - p^{(j)}(v_u, v_\omega)]$ factors and hence we can approximate the *no-further edge probability* as

(5.28)
$$\prod_{\omega \in V^{\circ}(\bar{\mathfrak{t}})} \prod_{\substack{u,j:\\(u,j,\omega) \notin E(\bar{\mathfrak{t}})}} [1 - p^{(j)}(v_u, v_{\omega})] = (1 + o_{\mathbb{P}}(1)) \prod_{\omega \in V^{\circ}(\bar{\mathfrak{t}})} \prod_{\substack{u,j:\\u \ge v_{\omega}}} [1 - p^{(j)}(u, v_{\omega})].$$

We can approximate (5.29)

$$\prod_{\substack{u,j:\\u\geq v_{\omega}}} \left[1-p^{(j)}(u,v_{\omega})\right] = \exp\left(\Theta(1)\sum_{u,j:\ u\geq v_{\omega}} p^{(j)}(u,v_{\omega})^2\right) \exp\left(-\sum_{u,j:u\geq v_{\omega}} p^{(j)}(u,v_{\omega})\right).$$

We next investigate the first term in the RHS of (5.29), while the second, and the main, term is estimated using Lemma 5.6. We prove that the first exponential term in (5.29) is $(1 + o_{\mathbb{P}}(1))$.

Using Lemma 4.5, similarly as in (5.22),

(5.30)

$$\sum_{u,j:u\geq v_{\omega}} p^{(j)}(u,v_{\omega})^{2} = (1+o_{\mathbb{P}}(1)) \sum_{u\geq v_{\omega}} m_{u} \frac{\hat{\chi}_{v_{\omega}}^{2}}{(2\mathbb{E}[M]+\delta)^{2}v_{\omega}^{2}} \left(\frac{v_{\omega}}{u}\right)^{2\chi}$$

$$\leq (1+o_{\mathbb{P}}(1)) \frac{\hat{\chi}_{v_{\omega}}^{2}}{(2\mathbb{E}[M]+\delta)^{2}(v_{\omega})^{2-\chi}} \left[\sum_{u\geq v_{\omega}} m_{u} \left(\frac{1}{u}\right)^{\chi}\right]$$

$$= (1+o_{\mathbb{P}}(1)) \frac{(\hat{\chi}_{v_{\omega}}^{2}/n)}{(2\mathbb{E}[M]+\delta)^{2}\left(\frac{v_{\omega}}{n}\right)^{2-\chi}} \left[\frac{1}{n} \sum_{u\geq v_{\omega}} m_{u} \left(\frac{u}{n}\right)^{-\chi}\right].$$

By (5.25) and recalling that $v_{\omega} = \lceil na_{\omega} \rceil$,

$$\frac{1}{n} \sum_{u \ge v_{\omega}} m_u \left(\frac{u}{n}\right)^{-\chi} \stackrel{a.s.}{\to} \frac{\mathbb{E}[M]}{1-\chi} \left[1 - a_{\omega}^{1-\chi}\right] < C,$$

for some constant C > 0. It is enough to show that $\frac{\hat{\chi}_{v_{\omega}}^2}{n} = o_{\mathbb{P}}(1)$. For this, we fix $\varepsilon, \zeta > 0$, and not that, for sufficiently large n,

(5.31)
$$\mathbb{P}\left(\frac{\hat{\chi}_{v_{\omega}}^{2}}{n} \geq \zeta\right) = \mathbb{P}\left(\hat{\chi}_{v_{\omega}}^{p} \geq (n\zeta)^{p/2}\right) \leq \mathbb{E}\left[\hat{\chi}_{v_{\omega}}^{p}\right](n\zeta)^{-p/2} \leq \varepsilon.$$

Therefore, $\sum_{u,j:u\geq v_{\omega}} p^{(j)}(u,v_{\omega})^2 = o_{\mathbb{P}}(1)$ and, by (5.29) and Lemma 5.6,

(5.32)
$$\prod_{\substack{u,j:\\u\geq v_{\omega}}} [1-p^{(j)}(u,v_{\omega})] = (1+o_{\mathbb{P}}(1))\exp\left(-\hat{\chi}_{v_{\omega}}\lambda(a_{\omega})\right).$$

Conclusion of the proof. Substituting the *no-further edge probability* obtained in (5.32) and the conditional probability estimates obtained in Lemma 4.5 and 5.6, we obtain

$$\mathbb{P}_{m,\psi}\left(\overline{B}_{r}^{(G_{n})} \doteq \overline{\mathbf{t}}, v_{\omega} = \lceil na_{\omega} \rceil, \forall \omega \in V(\mathbf{t})\right)$$

$$(5.33) = (1 + o_{\mathbb{P}}(1))\frac{1}{n}\prod_{\omega \in V(\mathbf{t})}\prod_{\substack{u \in V(\mathbf{t}) \\ u > \omega \\ u \stackrel{\lambda}{\neq} \omega}} \frac{\hat{\chi}_{v_{\omega}}}{(2\mathbb{E}[M] + \delta)v_{\omega}} \left(\frac{v_{\omega}}{v_{u}}\right)^{\chi}\prod_{\omega \in V^{\circ}(\mathbf{t})} \exp\left(-\hat{\chi}_{v_{\omega}}\lambda(a_{\omega})\right)$$

$$= (1 + o_{\mathbb{P}}(1))\frac{1}{n}\prod_{\omega \in V(\mathbf{t})}\left(\frac{\hat{\chi}_{v_{\omega}}}{(2\mathbb{E}[M] + \delta)}\right)^{d_{v_{\omega}}^{(\mathrm{in})}(G_{n})}\prod_{\omega \in V^{\circ}(\mathbf{t})} \exp\left(-\hat{\chi}_{v_{\omega}}\lambda(a_{\omega})\right)$$

$$\times \prod_{(\omega,\omega l) \in E(\mathbf{t})}(v_{\omega} \lor v_{\omega l})^{-(1-\chi)}(v_{\omega} \land v_{\omega l})^{-\chi},$$

where $d_{v_{\omega}}^{(in)}(G_n) = d(v_{\omega}) - m_{v_{\omega}}$ is the number of vertices in CPU connected to v_{ω} . Observe that $d_{v_{\omega}}^{(in)}(G_n) = d_{\omega}^{(in)}(t)$ when ω has label Y, and $d_{v_{\omega}}^{(in)}(G_n) = d_{\omega}^{(in)}(t) + 1$ when ω has label O. Further, $d_{v_{\omega}}^{(in)}(G_n) = 1$ for $\omega \in \partial V(t)$ and label O. Therefore, (5.33) can be re-written as

$$\mathbb{P}_{m,\psi}\left(\overline{B}_{r}^{(G_{n})}\doteq\bar{\mathbf{t}}, v_{\omega}=\lceil na_{\omega}\rceil, \forall \omega\in V(\mathbf{t})\right)$$
$$=(1+o_{\mathbb{P}}(1))n^{-(1+|E(\mathbf{t})|)}\prod_{\omega\in V^{\circ}(\mathbf{t})}\left(\frac{\hat{\chi}v_{\omega}}{(2\mathbb{E}[M]+\delta)}\right)^{d_{\omega}^{(\mathrm{in})}(\mathbf{t})}\prod_{\omega\in V^{\circ}(\mathbf{t})}\exp\left(-\hat{\chi}v_{\omega}\lambda(a_{\omega})\right)$$

(5.34)

$$\times \prod_{\substack{\omega \in V(\mathfrak{t})\\\omega \text{ is o}}} \frac{\hat{\chi}_{v_{\omega}}}{(2\mathbb{E}[M] + \delta)} \prod_{(\omega, \omega l) \in E(\mathfrak{t})} (a_{\omega} \vee a_{\omega l})^{-(1-\chi)} (a_{\omega} \wedge a_{\omega l})^{-\chi}.$$

Since t is a tree, |V(t)| = 1 + |E(t)| and hence (5.34) leads to (5.20).

REMARK 5.7 (Density of vertex-marked CPU). Every vertex $\omega \in V^{\circ}(t)$ has $m_{v_{\omega}}$ many out-edges that can be marked in $m_{v_{\omega}}$! different ways. If we sum (5.20) out over the edgemarks, then all such edge-marked graphs produce the same vertex-marked graph. Observe that we have not considered any of the out-edges from the \circ labelled vertices in $\partial V(t)$. Exactly one out-edge from every γ labelled vertex in $\partial V(t)$ is considered and its edge-mark can be labelled in $m_{v_{\omega}}$ many possible ways. Therefore by summing out the edge-marks,

$$\mathbb{P}_{m,\psi}\left(B_{r}^{(G_{n})}(o)\simeq \mathbf{t}, v_{\omega}=\lceil na_{\omega}\rceil, \forall \omega \in V(\mathbf{t})\right)$$
$$= (1+o_{\mathbb{P}}(1))n^{-|V(\mathbf{t})|}\prod_{\omega\in V^{\circ}(\mathbf{t})}\left[(m_{v_{\omega}})!\left(\frac{\hat{\chi}_{v_{\omega}}}{2\mathbb{E}[M]+\delta}\right)^{d_{\omega}^{(\mathrm{in})}(\mathbf{t})}\exp\left(-\hat{\chi}_{v_{\omega}}\lambda(a_{\omega})\right)\right]$$

(5.35)

$$\times \prod_{(\omega,\omega l)\in E(\mathbf{t})} (a_{\omega} \vee a_{\omega l})^{-\chi} (a_{\omega} \wedge a_{\omega l})^{-(1-\chi)} \prod_{\substack{\omega \in V(\mathbf{t})\\\omega \text{ is o}}} \frac{\hat{\chi}_{v_{\omega}}}{2\mathbb{E}[M] + \delta} \prod_{\substack{\omega \in \partial V(\mathbf{t})\\\omega \text{ is o}}} m_{v_{\omega}}.$$

Conditionally on (m, ψ) , we have now obtained the density of the *r*-neighbourhood of a randomly chosen vertex of $PA_n^{(A)}(m, \delta)$. This will be useful in the proof of Theorem 5.1 below.

From Remark 5.7, we have explicitly obtained the age density of the CPU graphs. Now we aim to show that this age density of CPU converges to that of the RPPT.

Proof of Theorem 5.1. We have obtained the explicit form of $f_{r,t}((a_{\omega})_{\omega \in V(t)}; (\hat{\chi}_{v_{\omega}})_{\omega \in V(t)})$ in Proposition 5.2. Remark 5.7 gives us an expression for the LHS of (5.4). Thus, we are left to show that

$$\mathbb{E}\left[\prod_{\omega\in V^{\circ}(\mathbf{t})}\left[(m_{v_{\omega}})!\left(\frac{\hat{\chi}_{v_{\omega}}}{2\mathbb{E}[M]+\delta}\right)^{d_{\omega}^{(\mathrm{in})}(\mathbf{t})}\exp\left(-\hat{\chi}_{v_{\omega}}\lambda(a_{\omega})\right)\right]\prod_{\substack{\omega\in V(\mathbf{t})\\\omega\text{ is o}}}\frac{\hat{\chi}_{v_{\omega}}}{2\mathbb{E}[M]+\delta}\prod_{\substack{\omega\in\partial V(\mathbf{t})\\\omega\text{ is y}}}m_{v_{\omega}}\right]$$

(5.36)

$$=\chi^{n_{\mathsf{o}}}(1-\chi)^{n_{\mathsf{v}}}\mathbb{E}\left[\prod_{\omega\in V^{\circ}(t)}\left(m'_{\omega}-\mathbb{1}_{\{\omega\text{ is }\mathsf{v}\}}\right)!\left(\hat{\chi}'_{\omega}\right)^{d_{\omega}^{(\mathrm{in})}(t)}\exp\left(-\hat{\chi}'_{\omega}\lambda(a_{\omega})\right)\right],$$

where $\hat{\chi}'_{\omega} \sim \text{Gamma}\left(m'_{\omega} + \delta + \mathbb{1}_{\{\omega \text{ is } o\}}, 1\right)$ and $\hat{\chi}_{v_{\omega}} \sim \text{Gamma}\left(m_{v_{\omega}} + \delta, 1\right)$. The distribution of m'_{ω} is same as $M^{(\delta)}$ when ω has label γ , $M^{(0)}$ when ω has label \circ and, M when ω is the root. $m_{v_{\omega}}$ are i.i.d. copies of M. To prove (5.36), we start by simplifying the LHS.

First we rewrite

(5.37)
$$\prod_{\substack{\omega \in V(\mathbf{t})\\\omega \text{ is o}}} \frac{\hat{\chi}_{v_{\omega}}}{2\mathbb{E}[M] + \delta} = \prod_{\substack{\omega \in V(\mathbf{t})\\\omega \text{ is o}}} \left(\frac{\hat{\chi}_{v_{\omega}}}{m_{v_{\omega}} + \delta}\right) \left(\frac{m_{v_{\omega}} + \delta}{\mathbb{E}[M] + \delta}\right) \left(\frac{\mathbb{E}[M] + \delta}{2\mathbb{E}[M] + \delta}\right).$$

38

Remember that $\chi = \frac{\mathbb{E}[M] + \delta}{2\mathbb{E}[M] + \delta}$. Therefore from (5.37), we see that we obtain a factor χ for each of the 0 labelled vertices in t. The first two terms in RHS of (5.37) give rise to the size-biasing of the Gamma random variables and the out-edge distribution of the 0 labelled vertices, as we observed in the definition of RPPT. Now, for $\omega \in \partial V(t)$, there is no other term in the LHS of (5.36) containing $\hat{\chi}_{v_{\omega}}$ and $m_{v_{\omega}}$ and these are independent of the rest. Therefore taking expectation with respect to these $\hat{\chi}_{v_{\omega}}$ and $m_{v_{\omega}}$, the first two terms in (5.37) turn out to be 1. Next, we rewrite

$$\prod_{\substack{\omega \in V^{\circ}(\mathsf{t})\\\omega \text{ is y}}} m_{v_{\omega}}! \prod_{\substack{\omega \in \partial V(\mathsf{t})\\\omega \text{ is y}}} m_{v_{\omega}} = \prod_{\substack{\omega \in V^{\circ}(\mathsf{t})\\\omega \text{ is y}}} (m_{v_{\omega}} - 1)! \prod_{\substack{\omega \text{ is y}}} m_{v_{\omega}}$$

$$(5.38) \qquad \qquad = \prod_{\substack{\omega \in V^{\circ}(\mathsf{t})\\\omega \text{ is y}}} (m_{v_{\omega}} - 1)! \prod_{\substack{\omega \text{ is y}}} \left(\frac{m_{v_{\omega}}}{\mathbb{E}[M]}\right) \left(\frac{\mathbb{E}[M]}{2\mathbb{E}[M] + \delta}\right) (2\mathbb{E}[M] + \delta)^{n_{\mathsf{Y}}}.$$

Again by definition, $1 - \chi = \frac{\mathbb{E}[M]}{2\mathbb{E}[M] + \delta}$. Therefore, similarly as in (5.37), we see that the sizebiased out-degree distributions of the γ labelled vertices arise from the second term in (5.38) and for every γ labelled vertex we obtain a factor $1 - \chi$ as we observe in the RPPT. There is no size-biasing in $\chi_{v_{\varnothing}}$ and $m_{v_{\varnothing}}$. The vertices in $V^{\circ}(t)$ can be partitioned in 3 sets: the root, o labelled vertices and γ labelled vertices. Therefore, using the simplification of expressions in (5.37) and (5.38),

LHS of (5.36) =
$$\chi^{n_0}(1-\chi)^{n_Y}(2\mathbb{E}[M]+\delta)^{n_Y-\sum\limits_{\omega\in V^0(t)}d^{(in)}(t)}$$

(5.39) $\times \mathbb{E}\left[\prod_{\omega\in V^o(t)}(m'_{\omega}-\mathbb{1}_{\{\omega \text{ is } Y\}})!\left(\hat{\chi}'_{\omega}\right)^{d^{(in)}(t)}\exp\left(-\hat{\chi}'_{\omega}\lambda(a_{\omega})\right)\right],$

where $\hat{\chi}'_{\omega}$ is defined as before. Note that $d^{(in)}_{\omega}(t)$ denotes the number of γ labelled children of ω and therefore summing $d^{(in)}_{\omega}(t)$ over all $\omega \in V^{\circ}(t)$ gives the total number of γ labelled vertices in t and hence the terms in the exponent of $2\mathbb{E}[M] + \delta$ cancel. Therefore, (5.36) follows immediately from (5.39), proving the required result.

For obtaining the first moment convergence in (5.2) using Theorem 5.1, we have to sum over the range $(\lceil n(a_{\omega} - 1/r) \rceil, \lceil n(a_{\omega} + 1/r) \rceil)^{|V(t)|}$. Summing the equality over this range, we obtain

$$\frac{1}{n}\mathbb{E}\left[N_{r,n}\left(\mathbf{t},(a_{\omega})_{\omega\in V(\mathbf{t})}\right)\right] \to \mu\left(B_{r}^{(G)}(\varnothing)\simeq\mathbf{t}, |A_{\omega}-a_{\omega}|\leq\frac{1}{r}, \forall \omega\in V(\mathbf{t})\right).$$

5.2. Second Moment Convergence. Here we show that

$$\frac{1}{n^2} \mathbb{E}\left[N_{r,n}\left(\mathbf{t}, (a_{\omega})_{\omega \in V(\mathbf{t})}\right)^2\right] \to \mu\left(B_r^{(G)}(\emptyset) \simeq \mathbf{t}, \ |A_{\omega} - a_{\omega}| \le \frac{1}{r}, \forall \omega \in V(\mathbf{t})\right)^2,$$

or, alternatively, the variance of $N_{r,n}/n$ vanishes. Expanding the double sum yields

$$N_{r,n}\left(\mathbf{t}, (a_{\omega})_{\omega \in V(\mathbf{t})}\right)^{2} = \sum_{v \in [n]} \mathbb{1}_{\left\{B_{r}^{(G_{n})}(v) \simeq \mathbf{t}, |v_{\omega}/n - a_{\omega}| \leq 1/r \, \forall \omega \in V(t)\right\}}$$

$$(5.40) \qquad \qquad + \sum_{\substack{u \neq v \\ u, v \in [n]}} \mathbb{1}_{\left\{B_{r}^{(G_{n})}(u) \simeq \mathbf{t}, B_{r}^{(G_{n})}(v) \simeq \mathbf{t}, \left|\frac{u_{\omega}}{n} - a_{\omega}\right| \leq \frac{1}{r}, \left|\frac{v_{\omega}}{n} - a_{\omega}\right| \leq \frac{1}{r}, \forall \omega \in V(t)\right\}}$$

where v_{ω} and u_{ω} are the vertices in $B_r^{(G_n)}(v)$ and $B_r^{(G_n)}(u)$ corresponding to the vertex $\omega \in V(t)$. Note that the first term in the RHS of (5.40) equals $N_{r,n}(t, (a_{\omega})_{\omega \in V(t)})$. By the results proved earlier in this section, it follows that this term, upon dividing by n^2 , vanishes. Therefore, we are left to analyse the second term on the RHS of (5.40). First, we show that *r*-neighbourhoods of two uniformly chosen vertices are disjoint whp. Note that

$$(5.41) \ \mathbb{P}\left(B_{r}^{(G_{n})}\left(o_{n}^{(1)}\right) \cap B_{r}^{(G_{n})}\left(o_{n}^{(2)}\right) = \varnothing\right) = \mathbb{P}\left(o_{n}^{(2)} \notin B_{2r}^{(G_{n})}\left(o_{n}^{(1)}\right)\right) = 1 - \frac{1}{n}\mathbb{E}\left[\left|B_{2r}^{(G_{n})}\left(o_{n}^{(1)}\right)\right|\right].$$

Previously we have proved that $B_{2r}^{(G_n)}(o_n^{(1)})$ converges in distribution to $B_r^{(G)}(\emptyset)$, where (G, \emptyset) is the random Pólya point tree with parameters M and δ , so that $\{|B_{2r}^{(G_n)}(o_n^{(1)})|\}_{n\in\mathbb{N}}$ is a tight sequence of random variables. Therefore the r-neighbourhood of two uniformly chosen vertices are whp disjoint. Now as we did for the first moment, we use a density argument for the second term:

THEOREM 5.8. Fix $\delta > -\inf \operatorname{supp}(M)$, and let $G_n = \operatorname{PA}_n^{(A)}(\boldsymbol{m}, \delta)$. Let $(\mathfrak{t}_1, (a_\omega)_{\omega \in V(\mathfrak{t}_1)})$ and $(\mathfrak{t}_2, (a_\omega)_{\omega \in V(\mathfrak{t}_2)})$ be two vertex-marked trees of depth r with disjoint vertex marks. If v_ω denotes the vertex in G_n corresponding to $\omega \in V(\mathfrak{t}_1) \cup V(\mathfrak{t}_2)$, then, uniformly for all distinct $v_\omega \geq \eta n$ and $\hat{\chi}_{v_\omega} \leq (v_\omega)^{1-\varrho/2}$,

$$\mathbb{P}_{m,\psi}\left(B_{r}^{(G_{n})}\left(o_{n}^{(1)}\right)\simeq \mathbf{t}_{1},\ B_{r}^{(G_{n})}\left(o_{n}^{(2)}\right)\simeq \mathbf{t}_{2},\ v_{\omega}=\lceil na_{\omega}\rceil,\forall\omega\in V(\mathbf{t}_{1})\cup V(\mathbf{t}_{2})\right)$$

$$=(1+o_{\mathbb{P}}(1))\frac{1}{n^{|V(\mathbf{t}_{1})\cup V(\mathbf{t}_{2})|}}\ g_{r,\mathbf{t}_{1}}\left((a_{\omega})_{\omega\in V(\mathbf{t}_{1})};(m_{v_{\omega}},\hat{\chi}_{v_{\omega}})_{\omega\in V(\mathbf{t}_{1})}\right)$$

$$\times\ g_{r,\mathbf{t}_{2}}\left((a_{\omega})_{\omega\in V(\mathbf{t}_{2})};(m_{v_{\omega}},\hat{\chi}_{v_{\omega}})_{\omega\in V(\mathbf{t}_{2})}\right)$$

where $g_{r,t}(\cdot)$ is as in Theorem 5.1 and $o_n^{(1)}, o_n^{(2)} \in [n]$ are chosen independently and uniformly at random.

PROOF. Since most of the steps in this proof are similar to the proof of Theorem 5.1, we will be more concise. We restrict our analysis to the case where $B_r^{(G_n)}(o_n^{(1)})$ and $B_r^{(G_n)}(o_n^{(2)})$ are disjoint graphs. Using the conditional independence of the edge-connection events for CPU, we can write the LHS of (5.42) as

$$\mathbb{P}_{m,\psi}\left(B_{r}^{(G_{n})}\left(o_{n}^{(1)}\right)\simeq \mathbf{t}_{1},\ B_{r}^{(G_{n})}\left(o_{n}^{(2)}\right)\simeq \mathbf{t}_{2},\ v_{\omega}=\lceil na_{\omega}\rceil,\forall\omega\in V(\mathbf{t}_{1})\cup V(\mathbf{t}_{2})\right)$$

$$(5.43) = \mathbb{P}_{m,\psi}\left(B_{r}^{(G_{n})}\left(o_{n}^{(1)}\right)\simeq \mathbf{t}_{1},\ v_{\omega}=\lceil na_{\omega}\rceil,\forall\omega\in V(\mathbf{t}_{1})\right)$$

$$\times \mathbb{P}_{m,\psi}\left(B_{r}^{(G_{n})}\left(o_{n}^{(2)}\right)\simeq \mathbf{t}_{2},\ v_{\omega}=\lceil na_{\omega}\rceil,\forall\omega\in V(\mathbf{t}_{2})\right).$$

Theorem 5.1 then immediately implies Theorem 5.8.

Considering $t_1 \simeq t_2$ but with different marks, by Theorem 5.8, it follows that

$$\frac{1}{n^2} \mathbb{E}\left[N_{r,n}\left(\mathbf{t}, (a_{\omega})_{\omega \in V(\mathbf{t})}\right)\right] = (1 + o(1))\mu \left(B_r^{(G)}(\varnothing) \simeq \mathbf{t}, |A_{\omega} - a_{\omega}| \le \frac{1}{r}, \forall \omega \in V(\mathbf{t})\right)^2.$$

Therefore the variance of $N_{r,n}(t, (a_{\omega})_{\omega \in V(t)})/n$ vanishes and the local convergence of model (A) to the random Pólya point tree follows immediately.

REMARK 5.9 (Convergence of models (B) and (D)). We have now proved the local convergence of the model (A) which is equal in distribution to $CPU^{(SL)}$. The proof of local convergence of models (B) and (D) follows along the same lines. Theorem 3.9 and 3.10 show

that models (B) and (D) have the same law as $CPU^{(NSL)}$ and $PU^{(NSL)}$, respectively. Recall that by Remark 4.6, the edge-connection probabilities for all the models $CPU^{(SL)}$, $CPU^{(NSL)}$ and $PU^{(NSL)}$ behave similarly. Therefore, upon substitution of these in (5.27), the local convergence of models (B) and (D) follows from the same calculation.

6. Coupling between models (E), (F) and (D). In this section, we prove that models (D), (E) and (F) have the same local limit. We do this by showing that for any fixed $r \ge 1$, we can couple models (D), (E) and (F) in such a way that whp, the *r*-neighbourhood of a uniformly chosen vertex in all these models are identical.

We follow the same coupling used in [8] with proper modifications as required for our models. In the coupling argument we work on the high-probability event $\mathcal{E} = \mathcal{E}_m \cap \mathcal{E}_{\hat{\chi},\psi}$, where \mathcal{E}_m and $\mathcal{E}_{\hat{\chi},\psi}$ are $\{m\}$ and $\{m,\psi\}$ measurable events respectively. In \mathcal{E}_m , following regularity properties hold true:

m_v ≤ v^{1-ℓ/2} for all v > K for some K ∈ N;
 for η' = η(ε/8, 2r), there exists a c₀ such that

$$\frac{1}{n}\sum_{u\geq\eta'n}m_u\left(\frac{u}{n}\right)^{-\chi}\leq c_0;$$

whereas in $\mathcal{E}_{\hat{x},\psi}$, the bounds on the attachment probabilities in Lemma 4.5 hold true. In Section 4, we have proved that $\mathcal{E}_{\hat{x},\psi}$ is a high probability event and from (5.25), substituting a_{ω} by η , we can prove that (2) is also a high probability event. (1) follows whp from the Borel-Cantelli Lemma proving \mathcal{E}_m also a high probability event. Therefore \mathcal{E} is indeed a high probability event and $\mathbb{P}(\mathcal{E}) > 1 - \varepsilon/4$.

The proof for models (E) and (F) are highly similar. Thus, in this section we provide the coupling argument for model (E) in detail, and state the adaptations for model (F) in Appendix C.

For all $j \in [m_n]$, the conditional edge-connection probability for model (E) is given by

(6.1)
$$\mathbb{P}_{m}\left(n \stackrel{j}{\leadsto} v \left| \mathrm{PA}_{n-1}^{(\mathrm{E})}(\boldsymbol{m}, \delta) \right.\right) = \frac{d_{v}(n-1) + \delta}{a_{[2]} + 2\left(m_{[n-1]} - 2\right) + (n-1)\delta}$$

We aim to couple $PA_n^{(E)}(\boldsymbol{m}, \delta)$ with $PA_n^{(D)}(\boldsymbol{m}, \delta)$. Let $V = \{1, 2, ...\}$ be the vertices of the preferential attachment model. For $1 \neq n \in V$ and $i \in [m_n]$, e_n^i and f_n^i denote the vertices in [n-1] to which vertex n connects with its *i*-th edge in models (D) and (E), respectively. Denote

(6.2)
$$\mathbf{e}_n = \left\{ e_n^i \right\}_{i \in [m_n]} \quad \text{and} \quad \mathbf{f}_n = \left\{ f_n^i \right\}_{i \in [m_n]}$$

Since we start with the same initial graph G_0 , $\mathbf{e}_2 = \mathbf{f}_2$. Conditionally on \boldsymbol{m} and $\{\mathbf{e}_l\}_{l < n}$, let \mathbf{e}_n have distribution $D_1^{(n)}$ and, conditionally on $\{\mathbf{f}_l\}_{l < n}$, \mathbf{f}_n have distribution $D_2^{(n)}$. Let $D^{(n)}$ be a coupling of $D_1^{(n)}$ and $D_2^{(n)}$ that minimizes the total variation distance. Then, we choose \mathbf{e}_n and \mathbf{f}_n according to $D^{(n)}$. This provides us with a coupling between models (D) and (E).

PROPOSITION 6.1 (Coupling between models (D) and (E)). Let $(G_n)_{n\geq 2}$ and $(G'_n)_{n\geq 2}$ be the sequences of preferential attachment graphs of models (D) and (E), respectively. Fix $\varepsilon > 0$ and $r \in \mathbb{N}$. Then, there exist a coupling $(\hat{G}_n, \hat{G}'_n)_{n\geq 1}$ and $n_0 \in \mathbb{N}$ (possibly depending on ε and r) such that for all $n > n_0$, with probability at least $1 - \varepsilon$, the r-neighbourhoods of a randomly chosen vertex $o_n \in [n]$ are the same in both G_n and G'_n . To simplify notation, below we write $(G_n)_{n\geq 2}$ and $(G'_n)_{n\geq 2}$ for the coupled graph processes. Fix $\varepsilon > 0$ and $r \in \mathbb{N}$ and let $B_r(v)$ and $B'_r(v)$ denote the *r*-neighbourhood of v in G_n and G'_n , respectively. We denote the set of bad vertices v for which $B_r(v) \neq B'_r(v)$ by

$$\mathcal{B}_{n,r} = \left\{ v \in [n] \colon B_r(v) \neq B'_r(v) \right\}$$

Since o_n is chosen uniformly from the set of vertices, the probability that o_n is a bad vertex equals $\mathbb{E}(|\mathcal{B}_{n,r}|/n)$. Thus, to prove Proposition 6.1, it is enough to show that $\mathbb{E}(|\mathcal{B}_{n,r}|) \leq \varepsilon n$.

We claim that for $B_r(v) \neq B'_r(v)$ to hold, there must be a vertex $u \in B_r(v)$ such that either $e_{n'}^i = u \neq f_{n'}^i$ or $e_{n'}^i \neq u = f_{n'}^i$ for some n' > u and $i \in [m_{n'}]$. To investigate these bad vertices, we thus concentrate on the bad events

(6.3)
$$A_{n'}^{(u)} = \left\{ \exists i \in [m_{n'}] : e_{n'}^i = u \neq f_{n'}^i, \text{ or } e_{n'}^i \neq u = f_{n'}^i \right\}$$

Let us consider $A^{(u)} = \bigcup_{n \ge n' > u} A^{(u)}_{n'}$, i.e., $A^{(u)}$ is the event that at least one of the edges

received by u is different in G_n and G'_n . The 1-neighbourhoods of vertex v are different in both G_n and G'_n precisely when $A^{(u)}$ happens for at least one of the $u \in B_1(v)$. Now inductively on r, it is easy to show that $B_r(v) = B'_r(v)$ unless there exists a $u \in B_r(v)$ for which $A^{(u)}$ holds true.

The following lemma gives us a tool to bound the probabilities of the bad events:

LEMMA 6.2 (Conditional control on bad events). With the coupling of models (E) and (D) defined earlier in this section and fixing $v \ge \eta' n$, we define $A_n = A_n^{(v)}$ to keep the notations simple. Then,

(6.4)
$$\mathbb{P}\left(A_n \cap \mathcal{E} \left| \bigcap_{h=v+1}^{n-1} A_h^c \cap \mathcal{E} \right. \right) = o\left(n^{-1}\right).$$

Lemma 6.2 implies that, conditionally on the event that there have not been any bad events till the (n-1)-st vertex joins the graph, the probability of the bad event taking place when the *n*-th vertex is joining the graph is o(1/n).

We first provide the proof of Proposition 6.1 subject to Lemma 6.2, and then we prove Lemma 6.2. First we construct a set $W_{n,r}$ of well-behaved vertices with the following properties:

 \triangleright for all $v \in W_{n,r}$, $B_{2r}(v)$ has no more than N vertices;

 \triangleright the oldest vertex in $B_{2r}(v)$ is not older than $\eta' n$.

By the proof of local convergence of model (D) and the regularity property of RPPT (Lemma 4.7),

(6.5)
$$\mathbb{E}|W_{n,r}| \ge \left(1 - \frac{\varepsilon}{2}\right)n.$$

Proof of Proposition 6.1. To prove the proposition, it is enough to show that

(6.6)
$$\mathbb{E}\left|W_{n,r}\cap\mathcal{B}_{n,r}\right|\leq\varepsilon n/2\,,$$

i.e., there are few well-behaved vertices that are bad. To prove (6.6), we note that if $v \in W_{n,r} \cap \mathcal{B}_{n,r}$, there must be a $u \in B_r(v)$ such that $A^{(u)}$ is true. Therefore,

(6.7)
$$|W_{n,r} \cap \mathcal{B}_{n,r}| = \sum_{v \in W_{n,r}} \mathbb{1}_{\{v \in \mathcal{B}_{n,r}\}} \leq \sum_{v \in W_{n,r}} \sum_{u \in B_{r}(v)} \mathbb{1}_{A^{(u)}}$$
$$= \sum_{u \in (\eta'n,n]} \mathbb{1}_{A^{(u)}} \sum_{v \in B_{r}(u)} \mathbb{1}_{\{v \in W_{n,r}\}}.$$

Note that the second sum in the RHS of (6.7) produces 0 if there is no well-behaved vertex in $B_r(u)$. Otherwise, the sum can be bounded by $|B_r(u)|$. Again $v \in B_r(u)$ implies that $u \in B_r(v)$ and $B_r(u)$ is a subset of $B_{2r}(v)$. If there is a well-behaved vertex in $B_r(u)$, we can bound $|B_r(u)| \le N$ from the properties of the well-behaved vertices. Therefore the second sum on the RHS of (6.7) can be uniformly bounded by N for all vertices $u > \eta' n$. Hence the cardinality of the well-behaved bad vertices can be further upper-bounded as

(6.8)
$$|W_{n,r} \cap \mathcal{B}_{n,r}| \le N \sum_{u \in (\eta'n,n]} \mathbb{1}_{A^{(u)}},$$

so that

(6.9)
$$\mathbb{E} |W_{n,r} \cap \mathcal{B}_{n,r}| \le N \sum_{u \in (\eta'n,n]} \mathbb{P}(A^{(u)}).$$

By the control on the bad events obtained in Lemma 6.2,

(6.10)
$$\mathbb{P}(A^{(u)}) = \mathbb{P}\left(\bigcup_{l=u+1}^{n} A_{l}^{(u)} \cap \mathcal{E}\right) + \mathbb{P}(\mathcal{E}^{c})$$
$$= \sum_{l=u+1}^{n} \mathbb{P}\left(A_{l}^{(u)} \cap \left(\bigcap_{h=u+1}^{l} \left(A_{h}^{(u)}\right)^{c}\right) \cap \mathcal{E}\right) + \mathbb{P}(\mathcal{E}^{c}) = o(1) + \varepsilon/4.$$

Substituting the bound on $\mathbb{P}(A^{(u)})$ in (6.9),

(6.11)
$$\mathbb{E}\left(|W_{n,r} \cap \mathcal{B}_{n,r}|\right) = o(n) + n\varepsilon/4,$$

which completes the proof of Proposition 6.1 subject to Lemma 6.2.

It remains to prove Lemma 6.2 about bad events, for which we need the following lemma:

LEMMA 6.3 (Bounds on second moments of degrees in model (D)). There exists $c_1 > 0$ such that for all $v \ge \eta' n$,

(6.12)
$$\mathbb{E}_m\left(d_v(n)^2 \mathbb{1}_{\mathcal{E}}\right) \le c_1 m_v^2 \,.$$

PROOF. By definition,

(6.13)
$$d_v(n) = m_v + \sum_{l=v+1}^n X_l, \quad \text{where} \quad X_l = \sum_{j=1}^{m_l} \mathbb{1}_{\{l \stackrel{j}{\to} v\}}$$

By Theorem 3.10, model (D) is equal in distribution to the Pólya urn graph. Therefore,

Now conditionally on $\{m, \psi\}$, $(X_l)_{l \in (v,n]}$ are independent $\operatorname{Binom}(m_l, p_l)$ random variables, where $p_l = \mathbb{P}_{m,\psi}(l \rightsquigarrow v)$. Using this independence in the second term of the RHS

of (6.14),

(6.15)
$$\mathbb{E}_{m,\psi} \Big[\sum_{l=\nu+1}^{n} X_l \Big]^2 = \sum_{l=\nu+1}^{n} \operatorname{Var}_{m,\psi}(X_l) + \Big(\sum_{l=\nu+1}^{n} \mathbb{E}_{m,\psi}[X_l] \Big)^2 \\ = \sum_{l=\nu+1}^{n} m_l p_l (1-p_l) + \Big(\sum_{l=\nu+1}^{n} m_l p_l \Big)^2.$$

Following the same line of proof as in Lemma 5.6 and using properties of $\mathcal{E}_{\hat{x},\psi}$, we can show that conditionally on m, ψ ,

(6.16)
$$\left(\sum_{l=\nu+1}^{n} m_l p_l (1-p_l)\right) \mathbb{1}_{\mathcal{E}_{\hat{\chi},\psi}} \leq \left(\sum_{l=\nu+1}^{n} m_l p_l\right) \mathbb{1}_{\mathcal{E}_{\hat{\chi},\psi}} \leq 2\hat{\chi}_{\nu} \left(\frac{1}{n} \sum_{u \geq \eta' n} m_u \left(\frac{u}{n}\right)^{-\chi}\right).$$

Therefore, using the property (2) of \mathcal{E}_m and substituting the bound of (6.16) in (6.15),

(6.17)
$$\mathbb{E}_m\left[\left(\sum_{l=v+1}^n m_l p_l\right)^2 \mathbb{1}_{\mathcal{E}_{\hat{\chi},\psi}}\right] \mathbb{1}_{\mathcal{E}_m} \le 4c_0^2 \mathbb{E}_m[\hat{\chi}_v^2] \le c_2 m_v^2$$

for some $c_2 > 0$. Choosing $c_1 = 2(1 + c_0 + c_2^2)$, for $v \ge \eta' n$, (6.12) holds true.

Proof of Lemma 6.2.. To prove Lemma 6.2 we bound the LHS of (6.4) by the total variation distance between $D_1^{(n)}$ and $D_2^{(n)}$.

In $PA_n^{(E)}(\boldsymbol{m}, \delta)$, conditionally on \boldsymbol{m} and $d_v(n-1) = d$, the probability of having k connections from n to v and $m_n - k$ connections to other vertices in [n-1] is given by

(6.18)
$$Q_k = \binom{m_n}{k} (p')^k (1-p')^{m_n-k},$$

where $p' = (d + \delta)/(a_{[2]} + 2(m_{[n-1]} - 2) + (n-1)\delta)$ is as defined in (6.1). On the other hand, in $PA_n^{(D)}(\boldsymbol{m}, \delta)$, conditionally on \boldsymbol{m} and $d_v(n-1) = d$, the probability of the same event is

(6.19)
$$T_k = \binom{m_n}{k} \prod_{l=0}^{k-1} p_l(k) \prod_{l=k}^{m_n-1} (1 - p_l(k)),$$

where

$$p_l(k) = \begin{cases} & \frac{d + l + \delta}{a_{[2]} + 2(m_{[n-1]} - 2) + l + (n-1)\delta}, & \text{for } l < k, \\ & \frac{d + k + \delta}{a_{[2]} + 2(m_{[n-1]} - 2) + l + (n-1)\delta}, & \text{for } l \ge k. \end{cases}$$

The exchangeability of model (D) implies that the probability of connecting any k edges from n to v is the same, which explains (6.19). Then, again conditionally on m, and $d_v(n-1) = d$, the total variation distance between $D_1^{(n)}$ and $D_2^{(n)}$ is given by (6.20)

$$d_{\rm TV}\left(D_1^{(n)}, D_2^{(n)}\right) = \frac{1}{2} \sum_{k=0}^{m_n} |Q_k - T_k| \le \frac{1}{2} \left[|Q_0 - T_0| + |Q_1 - T_1| + \sum_{k=2}^{m_n} Q_k + \sum_{k=2}^{m_n} T_k \right].$$

For the first two terms we use the telescoping bound in [8, equation (27)] that reads

(6.21)
$$|Q_k - T_k| \le \binom{m_n}{k} \left[\tilde{p}^{k-1} \sum_{l=0}^{k-1} \left| p' - p_l(k) \right| + (p')^k \sum_{l=k}^{m_n} \left| p' - p_l(k) \right| \right],$$

44

where $\tilde{p} = \max\{p', p_0(k), \dots, p_{m_n}(k)\}$. For k = 0 and 1, the above bound gives

(6.22)
$$|Q_0 - T_0| + |Q_1 - T_1| \le c_0 m_n^2 (d + \delta + 1)^2 n^{-2}.$$

For the last term in the RHS of (6.20), we use that T_k equals the probability that exactly k out-edges from vertex n connect to vertex v in model (D). Conditionally on m, let $\mathbb{P}_m^{(D)}$ denote the law for model (D). Next we condition on the event that the first 2 out-edges from n connect to v, which is denoted by $n \xrightarrow{1,2} v$.

(6.23)

$$\sum_{k=2}^{m_n} T_k = \sum_{k=2}^{m_n} \binom{m_n}{k} \mathbb{P}_m^{(D)} (n \rightsquigarrow v \text{ with its first } k \text{ edges})$$

$$\leq \frac{m_n^2}{2} \mathbb{P}_m^{(D)} \left(n \stackrel{1,2}{\rightsquigarrow} v \right) \sum_{k=2}^{m_n} \binom{m_n - 2}{k - 2} \mathbb{P}_m^{(D)} \left(n \rightsquigarrow v \text{ with first } k \text{ edges} \left| n \stackrel{1,2}{\rightsquigarrow} v \right)$$

$$\leq \frac{m_n^2 (d + \delta + 1)^2}{2n^2}.$$

The last inequality is obtained by substituting the first and second edge-connection probabilities from n to v and the conditional probabilities sum to 1. Similarly for model (E), if $\mathbb{P}_m^{(E)}$ denotes the conditional law for model (E), then the probability of connecting more than 1 edge from n to v can be bounded as

(6.24)

$$\sum_{k=2}^{m_n} Q_k = \sum_{k=2}^{m_n} \binom{m_n}{k} \mathbb{P}_m^{(\mathrm{E})} (n \rightsquigarrow v \text{ with its first } k \text{ edges})$$

$$\leq \frac{m_n^2}{2} \mathbb{P}_m^{(\mathrm{E})} \left(n \stackrel{1,2}{\rightsquigarrow} v \right) \sum_{k=2}^{m_n} \binom{m_n - 2}{k - 2} \mathbb{P}_m^{(\mathrm{E})} \left(n \rightsquigarrow v \text{ with first } k \text{ edges } \left| n \stackrel{1,2}{\rightsquigarrow} v \right)$$

$$\leq \frac{m_n^2 (d+1)^2}{2n^2}.$$

On the other hand, the total variation distance is bounded by 1. Therefore, collecting the bounds for all the contributions, conditionally on m, we obtain

(6.25)
$$d_{\scriptscriptstyle \mathrm{TV}}(D_1^{\scriptscriptstyle (n)}, D_2^{\scriptscriptstyle (n)}) \le 1 \wedge \frac{cm_n^2(d_v(n-1) + \delta + \delta)^2}{n^2} ,$$

for some constant c > 0. Therefore,

$$\mathbb{P}\left(A_n \cap \mathcal{E}\left|\bigcap_{h=v+1}^{n-1} A_h^c \cap \mathcal{E}\right.\right) \le \mathbb{E}\left[d_{\scriptscriptstyle \mathrm{TV}}\left(D_1^{\scriptscriptstyle (n)}, D_2^{\scriptscriptstyle (n)}\right) \mathbbm{1}_{\mathcal{E}}\right] \le \mathbb{E}\left[1 \wedge \frac{cm_n^2 \mathbb{E}_m[d_v(n-1)^2 \mathbbm{1}_{\mathcal{E}}]}{n^2}\right] \,.$$

By Lemma 6.3, we can bound the conditional probability on the LHS of (6.26) by

(6.27)
$$\mathbb{E}\left[1 \wedge \frac{cc_1^2 m_n^2 m_v^2}{n^2}\right] = \mathbb{P}\left(m_n m_v > \frac{n}{c_1 \sqrt{c}}\right) + \frac{cc_1^2}{n^2} \mathbb{E}\left[m_n^2 m_v^2 \mathbb{1}_{\{m_v m_n \le \frac{n}{c_1 \sqrt{c}}\}}\right],$$

where c_0 is as defined in Lemma 6.3. Since $(m_v)_{v\geq 3}$ have finite (1+p)-th moment, by Markov's inequality,

$$\mathbb{P}(m_v > x) \le c_2 x^{-(1+p)}.$$

Using the independence of m_n and m_v and the above inequality, we can bound the first probability in (6.27) by Markov's inequality as

(6.28)
$$\mathbb{P}\left(m_n m_v > \frac{n}{c_1 \sqrt{c}}\right) \le c_3 \mathbb{E}\left[(m_n m_v)^{1+p}\right] n^{-(1+p)} = o(n^{-1}) .$$

The second term in (6.27) can be written explicitly as (6.29)

$$\mathbb{E}\left[m_n^2 m_v^2 \mathbb{1}_{\{m_v m_n \le x\}}\right] \le \sum_{k=1}^x k \mathbb{P}(m_n m_v > k) \le c_3 \mathbb{E}\left[(m_n m_v)^{1+p}\right] \sum_{k=1}^x k^{-p} \le c_3 x^{1-p} .$$

Substituting the bounds obtained from (6.28) and (6.29) in (6.26) yields,

$$\mathbb{P}\left(A_n \cap \mathcal{E}\left|\bigcap_{h=v+1}^{n-1} A_h^c \cap \mathcal{E}\right.\right) \le c_4 n^{-(1+p)} = o\left(n^{-1}\right).$$

This completes the proof of Lemma 6.2.

The only difference between model (E) and (F) is that model (F) does not allow for multiple connections between two vertices, while model (E) does. Thus, model (F) and (D) could be coupled in a similar way with some minor adaptations:

PROPOSITION 6.4 (Coupling between models (D) and (F)). *The statement in Proposition* 6.1 also holds for models (D) and (F).

We defer the discussion of the necessary adaptations to Appendix C.

7. Proof of Corollary 1.6. In this section, we prove Corollary 1.6 as a consequence of Theorem 1.5.

Proof of Corollary 1.6. As a consequence of local convergence, if G_n converges locally in probability to (G, o) having law μ , then for any bounded continuous function $h : \mathcal{G}_{\star} \mapsto \mathbb{R}$,

(7.1)
$$\frac{1}{n} \sum_{u \in [n]} h(G_n, u) \xrightarrow{\mathbb{P}} \mathbb{E}_{\mu}[h(G, o)] .$$

We start by proving (1.19). From Theorem 1.5, we have the local convergence of preferential attachment models (A-F) to the $\text{RPPT}(M, \delta)$. Define for fixed $k \ge 1$,

(7.2)
$$h(H,u) = \sum_{\substack{v,j:\\ u \xrightarrow{\downarrow} v}} \frac{\mathbb{1}_{\{d_v(H)=k\}}}{m_u} ,$$

where $d_v(H)$ is the degree of the vertex v in the graph H. Clearly h is a bounded continuous function on \mathcal{G}_{\star} . Therefore by (7.1),

(7.3)
$$\frac{1}{n} \sum_{\substack{u,v,j:\\u \stackrel{j}{\to} v}} \frac{\mathbb{I}_{\{d_v(n)=k\}}}{m_u} \xrightarrow{\mathbb{P}} \mathbb{E}_{\mu_\star} \left[h(G, \emptyset) \right],$$

where (G, \emptyset) is the rooted RPPT (M, δ) with law μ_{\star} . Now h(H, o) is the fraction of older neighbours of o in H having degree k. Therefore, $\mathbb{E}_{\mu_{\star}}[h(G, \emptyset)]$ is the probability that a random 0 labelled neighbour of the root of RPPT (M, δ) has degree k. From the definition of RPPT (M, δ) , the degree distribution of an 0 labelled node with age a_{ω} is $1 + M^{(\delta)} +$ $Y(M^{(\delta)} + 1, a_{\omega})$, where $Y(M^{(\delta)} + 1, a_{\omega})$ is a mixed Poisson random variable with mixing distribution $\Gamma_{in}(M^{(\delta)} + 1)\lambda(a_{\omega})$ and $\Gamma_{in}(M^{(\delta)} + 1)$ is as defined in Section 1.4. On the other hand, a random 0 labelled neighbour of the root \emptyset has age distributed as $U_{\emptyset}U_1^{1/\chi}$. Therefore a random 0 labelled neighbour of the root has the degree distribution $1 + M^{(\delta)} + Y(M^{(\delta)} + 1, A_{\circ})$, where $Y(M^{(\delta)} + 1, A_{\circ})$ is as defined previously in Corollary 1.6 and

(7.4)
$$\mathbb{E}_{\mu_{\star}}[h(G,o)] = \mathbb{P}(1 + M^{(\delta)} + Y(M^{(\delta)} + 1, A_{o}) = k) = \tilde{p}_{k}^{(o)}$$

We prove (1.20) in a similar way with a few adaptations. Instead of considering all vertices, we consider the vertices that have at least one younger neighbour. Next we choose the bounded continuous function as follows

(7.5)
$$h_1(H,v) = \sum_{\substack{u,j:\\u \xrightarrow{j} v v}} \frac{\mathbb{I}_{\{d_v(H) > m_v\}} \mathbb{I}_{\{d_u(H) = k\}}}{d_v(H) - m_v} ,$$

and

 $h_2(H,v) = \mathbb{1}_{\{v \text{ has at least one younger neighbour}\}}$.

Similarly by (7.1),

(7.6)
$$\frac{1}{n} \sum_{\substack{u,v,j:\\u \stackrel{\rightarrow}{\rightarrow} v}} \frac{\mathbb{1}_{\{d_v(n) > m_v\}} \mathbb{1}_{\{d_u(n) = k\}}}{d_v(n) - m_v} \stackrel{\mathbb{P}}{\rightarrow} \mathbb{E}_{\mu_\star} \left[h_1(G, \emptyset) \right],$$

and $\frac{1}{n} \sum_{v \in [n]} \mathbb{1}_{\{v \text{ has at least one younger neighbour}\}} \xrightarrow{\mathbb{P}} \mathbb{E}_{\mu_{\star}} \left[h_2(G, \emptyset) \right]$.

where (G, \emptyset) is the rooted RPPT (M, δ) with law μ_{\star} . Now $h_1(H, o)$ is the fraction of younger neighbours of o in H having degree k. If o has no younger neighbour then define $h_1(H, o) = 0$. Therefore

(7.7)
$$\mathbb{E}_{\mu_{\star}}\left[h_{1}(G, \varnothing)\right] = \mathbb{E}_{\mu_{\star}}\left[h_{1}(G, \varnothing) \left| d_{\varnothing}^{(\mathrm{in})} > 0\right] \mathbb{P}(d_{\varnothing}^{(\mathrm{in})} > 0)\right]$$

and $\mathbb{E}_{\mu_*}[h_1(G, \emptyset) | d_{\emptyset}^{(\text{in})} > 0]$ is the probability that a random γ labelled neighbour of the root of RPPT (M, δ) has degree k, conditionally on the event that the root has at least one younger neighbour. From the definition of RPPT (M, δ) , the degree distribution of a γ labelled node of age a_{ω} is $M^{(0)} + Y(M^{(0)}, a_{\omega})$ where $Y(M^{(0)}, a_{\omega})$ is a mixed Poisson random variable with mixing distribution $\Gamma_{in}(M^{(0)})\lambda(a_{\omega})$.

Using the fact that $d_{\varnothing}^{(in)}$ is the total number of points in a Poisson point process, the ages of the γ labelled neighbours of \varnothing , conditioned on $d_{\varnothing}^{(in)} = n$, are i.i.d. random variables with density (1.21) [36, Exercise 4.34]. Hence conditioned on $d_{\varnothing}^{(in)} = n$, a uniformly chosen younger neighbour of the root \varnothing has age distribution given by A_1 with density (1.21). Therefore conditionally on the root having at least one younger neighbour, a random γ labelled neighbour of the root has the degree distribution $M^{(0)} + Y(M^{(0)}, A_{\gamma})$ where $Y(M^{(0)}, A_{\gamma})$ is as defined earlier in Corollary 1.6 and

(7.8)
$$\mathbb{E}_{\mu_{\star}}\left[h_{1}(G, \varnothing)\right] = \mathbb{P}\left(d_{\varnothing}^{(\mathrm{in})} > 0\right) \mathbb{P}\left(M^{(0)} + Y\left(M^{(0)}, A_{\mathsf{Y}}\right) = k\right) = \mathbb{P}\left(d_{\varnothing}^{(\mathrm{in})} > 0\right) \tilde{p}_{k}^{(\mathsf{Y})}$$

On the other hand, $\mathbb{E}_{\mu_*}[h_2(G, \emptyset)]$ is the probability of the root \emptyset having at least one younger neighbour and it is given by $\mathbb{P}(d_{\emptyset}^{(in)} > 0)$. Therefore,

(7.9)
$$\mathbb{E}_{\mu_{\star}}\left[h_{2}(G, \varnothing)\right] = \mathbb{P}\left(d_{\varnothing}^{(\mathrm{in})} > 0\right).$$

Since $\mathbb{P}(d_{\emptyset}^{(in)} > 0)$ is non-zero, (1.20) follows immediately from (7.8) and (7.9). This completes the proof of (a).

The proof of (b) makes use of similar calculations in [8, Lemma 5.2] and [18, Proposition 1.4] and we defer this proof to Appendix D. \Box

Acknowledgments. The authors would like to thank Shankar Bhamidi, Tiffany Y. Y. Lo and Delphin Sénizergues for their insightful suggestions on the initial version of this paper.

Funding. The work of AG, RSH, RvdH and RR is supported in part by the Netherlands Organisation for Scientific Research (NWO) through the Gravitation NETWORKS grant 024.002.003. The work of RvdH is further supported by the Netherlands Organisation for Scientific Research (NWO) through VICI grant 639.033.806. The work of RR is further supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 945045.



REFERENCES

- ALBERT, R. and BARABÁSI, A.-L. (2002). Statistical mechanics of complex networks. *Rev. Modern Phys.* 74 47–97. https://doi.org/10.1103/RevModPhys.74.47 MR1895096
- [2] ALDOUS, D. and STEELE, J. M. (2004). The objective method: probabilistic combinatorial optimization and local weak convergence. In *Probability on discrete structures. Encyclopaedia Math. Sci.* 110 1–72. Springer, Berlin. https://doi.org/10.1007/978-3-662-09444-0_1 MR2023650
- [3] BANERJEE, S., DEKA, P. and OLVERA-CRAVIOTO, M. (2023). Local weak limits for collapsed branching processes with random out-degrees. https://doi.org/10.48550/ARXIV.2302.00562
- BARABÁSI, A.-L. and ALBERT, R. (1999). Emergence of scaling in random networks. Science 286 509– 512. https://doi.org/10.1126/science.286.5439.509 MR2091634
- [5] BASU, A., SHANNIGRAHI, S., CHHABRA, S. S. and BRUNDAVANAM, A. (2014). On the Rise and Fall of Online Social Networks. https://doi.org/10.48550/ARXIV.1403.5617
- [6] BENJAMINI, I. and SCHRAMM, O. (2001). Recurrence of distributional limits of finite planar graphs. *Electron. J. Probab.* 6 no. 23, 13. https://doi.org/10.1214/EJP.v6-96 MR1873300
- [7] BERGER, N., BORGS, C., CHAYES, J. T. and SABERI, A. (2005). On the spread of viruses on the internet. In *Proceedings of the Sixteenth Annual ACM-SIAM Symposium on Discrete Algorithms* 301–310. ACM, New York. MR2298278
- [8] BERGER, N., BORGS, C., CHAYES, J. T. and SABERI, A. (2014). Asymptotic behavior and distributional limits of preferential attachment graphs. Ann. Probab. 42 1–40. https://doi.org/10.1214/12-AOP755 MR3161480
- [9] BHAMIDI, S. (2007). Universal techniques to analyze preferential attachment trees: Global and local analysis.
- [10] BIANCONI, G. and BARABÁSI, A.-L. (2011). Competition and multiscaling in evolving networks. In *The Structure and Dynamics of Networks* 361–367. Princeton University Press.
- [11] BINGHAM, N. H., GOLDIE, C. M. and TEUGELS, J. L. (1987). Regular variation. Encyclopedia of Mathematics and its Applications 27. Cambridge University Press, Cambridge. https://doi.org/10.1017/CBO9780511721434 MR898871
- [12] BOLLOBÁS, B. and RIORDAN, O. (2004). The diameter of a scale-free random graph. Combinatorica 24 5–34. https://doi.org/10.1007/s00493-004-0002-2 MR2057681
- [13] BOLLOBÁS, B., RIORDAN, O., SPENCER, J. and TUSNÁDY, G. (2001). The degree sequence of a scale-free random graph process. *Random Structures Algorithms* 18 279–290. https://doi.org/10.1002/rsa.1009 MR1824277
- [14] BORGS, C., CHAYES, J., DASKALAKIS, C. and ROCH, S. (2007). First to market is not everything: an analysis of preferential attachment with fitness. In STOC'07—Proceedings of the 39th Annual ACM Symposium on Theory of Computing 135–144. ACM, New York. https://doi.org/10.1145/1250790.1250812 MR2402437
- [15] CARAVENNA, F., GARAVAGLIA, A. and VAN DER HOFSTAD, R. (2019). Diameter in ultra-small scalefree random graphs. *Random Structures Algorithms* 54 444–498. https://doi.org/10.1002/rsa.20798 MR3938775
- [16] CHOI, B. D. and SUNG, S. H. (1987). Almost sure convergence theorems of weighted sums of random variables. *Stochastic Anal. Appl.* 5 365–377. https://doi.org/10.1080/07362998708809124 MR912863
- [17] COOPER, C. and FRIEZE, A. (2003). A general model of web graphs. *Random Structures Algorithms* 22 311–335. https://doi.org/10.1002/rsa.10084 MR1966545
- [18] DEIJFEN, M., VAN DEN ESKER, H., VAN DER HOFSTAD, R. and HOOGHIEMSTRA, G. (2009). A preferential attachment model with random initial degrees. Ark. Mat. 47 41–72. https://doi.org/10.1007/s11512-007-0067-4 MR2480915

48

- [19] DEREICH, S. (2016). Preferential attachment with fitness: unfolding the condensate. *Electron. J. Probab.* 21 Paper No. 3, 38. https://doi.org/10.1214/16-EJP3801 MR3485345
- [20] DEREICH, S., MÖNCH, C. and MÖRTERS, P. (2012). Typical distances in ultrasmall random networks. Adv. in Appl. Probab. 44 583–601. https://doi.org/10.1239/aap/1339878725 MR2977409
- [21] DEREICH, S. and MÖRTERS, P. (2009). Random networks with sublinear preferential attachment: degree evolutions. *Electron. J. Probab.* 14 no. 43, 1222–1267. https://doi.org/10.1214/EJP.v14-647 MR2511283
- [22] DEREICH, S. and MÖRTERS, P. (2011). Random networks with concave preferential attachment rule. Jahresber. Dtsch. Math.-Ver. 113 21–40. https://doi.org/10.1365/s13291-010-0011-6 MR2760002
- [23] DEREICH, S. and MÖRTERS, P. (2013). Random networks with sublinear preferential attachment: the giant component. Ann. Probab. 41 329–384. https://doi.org/10.1214/11-AOP697 MR3059201
- [24] DEREICH, S. and ORTGIESE, M. (2014). Robust analysis of preferential attachment models with fitness. Combin. Probab. Comput. 23 386–411. https://doi.org/10.1017/S0963548314000157 MR3189418
- [25] DOMMERS, S., VAN DER HOFSTAD, R. and HOOGHIEMSTRA, G. (2010). Diameters in preferential attachment models. J. Stat. Phys. 139 72–107. https://doi.org/10.1007/s10955-010-9921-z MR2602984
- [26] DOROGOVTSEV, S. N., GOLTSEV, A. V. and MENDES, J. F. F. (2008). Critical phenomena in complex networks. *Rev. Mod. Phys.* 80 1275–1335. https://doi.org/10.1103/RevModPhys.80.1275
- [27] DOROGOVTSEV, S. N., MENDES, J. F. F. and SAMUKHIN, A. N. (2000). Structure of Growing Networks with Preferential Linking. *Phys. Rev. Lett.* 85 4633–4636. https://doi.org/10.1103/PhysRevLett.85.4633
- [28] GAO, F. and VAN DER VAART, A. (2017). On the asymptotic normality of estimating the affine preferential attachment network models with random initial degrees. *Stochastic Process. Appl.* 127 3754–3775. https://doi.org/10.1016/j.spa.2017.03.008 MR3707244
- [29] GARAVAGLIA, A. (2019). Preferential attachment models for dynamic networks, PhD thesis, Mathematics and Computer Science.
- [30] GARAVAGLIA, A. and VAN DER HOFSTAD, R. (2018). From trees to graphs: collapsing continuous-time branching processes. J. Appl. Probab. 55 900–919. https://doi.org/10.1017/jpr.2018.57 MR3877316
- [31] GUT, A. (2013). Probability: a graduate course, second ed. Springer Texts in Statistics. Springer, New York. https://doi.org/10.1007/978-1-4614-4708-5 MR2977961
- [32] JAGERS, P. and NERMAN, O. (1984). The Growth and Composition of Branching Populations. Advances in Applied Probability 16 221–259.
- [33] JORDAN, J. (2018). Preferential attachment graphs with co-existing types of different fitnesses. J. Appl. Probab. 55 1211–1227. https://doi.org/10.1017/jpr.2018.80 MR3899937
- [34] LO, T. Y. Y. (2021). Weak local limit of preferential attachment random trees with additive fitness. https://doi.org/10.48550/ARXIV.2103.00900
- [35] MÓRI, T. F. (2005). The maximum degree of the Barabási-Albert random tree. *Combin. Probab. Comput.* 14 339–348. https://doi.org/10.1017/S0963548304006133 MR2138118
- [36] RESNICK, S. (1992). Adventures in stochastic processes. Birkhäuser Boston, Inc., Boston, MA. MR1181423
- [37] RUDAS, A., TÓTH, B. and VALKÓ, B. (2007). Random trees and general branching processes. Random Structures Algorithms 31 186–202. https://doi.org/10.1002/rsa.20137 MR2343718
- [38] SÉNIZERGUES, D. (2021). Geometry of weighted recursive and affine preferential attachment trees. *Electron. J. Probab.* 26 Paper No. 80, 56. https://doi.org/10.1214/21-ejp640 MR4269210
- [39] VAN DER HOFSTAD, R. (2017). Random graphs and complex networks. Vol. 1. Cambridge Series in Statistical and Probabilistic Mathematics, [43]. Cambridge University Press, Cambridge. https://doi.org/10.1017/9781316779422 MR3617364
- [40] VAN DER HOFSTAD, R. (2023). *Random graphs and complex networks. Vol.* 2. To appear in: Cambridge Series in Statistical and Probabilistic Mathematics.
- [41] VOITALOV, I., VAN DER HOORN, P., VAN DER HOFSTAD, R. and KRIOUKOV, D. (2019). Scale-free networks well done. *Phys. Rev. Research* 1 033034. https://doi.org/10.1103/PhysRevResearch.1.033034
- [42] WACLAW, B. and SOKOLOV, I. M. (2007). Finite-size effects in Barabási-Albert growing networks. *Phys. Rev. E (3)* 75 056114, 10. https://doi.org/10.1103/PhysRevE.75.056114 MR2361828

APPENDIX A: ADDITIONAL PROOFS

In this section we prove Theorems 3.9 and 3.10. These proofs follow that of Theorem 3.1, except for a few minor adaptations. Here we show that models (D) and (B) are equal in distribution with $PU^{(NSL)}$ and $CPU^{(NSL)}$ respectively. Recall that, $CPU^{(NSL)}$ could be obtained

by collapsing a special case $PU^{(NSL)}$. So, we prove Lemma A.1-A.2 and Corollary A.3 for general choice of m and ψ . Later, while proving Theorem 3.9, we shall use these lemmas and corollary with m = 1, whereas for proving Theorem 3.10, we continue with the m of model (D). The following lemmas and propositions are the $PU^{(NSL)}$ analogues of the lemmas and propositions proved in Section 3.1. The conditional edge-connection probabilities for $PU^{(NSL)}$ are given as follows:

LEMMA A.1 (Conditional edge-connection probabilities of $\mathrm{PU}^{(\mathrm{NSL})}$). Conditionally on \boldsymbol{m} and $(\psi_k)_{k\geq 1}$, the probability of connecting an edge from v to u in $\mathrm{PU}_n^{(\mathrm{NSL})}(\boldsymbol{m}, \boldsymbol{\psi})$, is given by $\psi_v(1-\psi)_{(v,u)}$.

The proof to this lemma is identical to that of Lemma 3.4, using the definition of $PU^{(NSL)}$. Similarly as Lemma 3.5 the above lemma allows us to compute the conditional law of $PU^{(NSL)}$:

LEMMA A.2 (Conditional graph probability of $PU^{(NSL)}$). For any graph $H \in \mathcal{H}_n$,

(A.1)
$$\mathbb{P}_{m,\psi}\left(\mathrm{PU}_{n}^{(\mathrm{NSL})}(\boldsymbol{m},\boldsymbol{\psi})\overset{\star}{\simeq}H\right) = \prod_{s\in[2,n]}\psi_{s}^{p_{s}}(1-\psi_{s})^{q_{s}},$$

where

$$p_s = d_s(H) - f_s,$$
 and $q_s = \sum_{u \in (2,n]} \sum_{j=1}^{m_u} \mathbb{1}_{\{s \in (v(u,j),u)\}}$

where v(u, j) is the vertex to which the *j*-th out-edge of *u* connects and $f_s = m_s$ for all $s \ge 3$ and $f_1 = a_1$ and $f_2 = a_2$ denote the degrees of the vertex 1 and 2 in the initial graph G_0 .

Since the proof strategy of this lemma is identical to that of Lemma 3.5, we omit the proof to this lemma also. Using Lemma 3.6, conditionally on m, the graph probability of $PU^{(NSL)}$ is computed as follows:

COROLLARY A.3. For $H \in \mathcal{H}_n$,

(A.2)
$$\mathbb{P}_m\left(\mathrm{PU}_n^{(\mathrm{NSL})}(\boldsymbol{m}, \boldsymbol{\psi}) \stackrel{\star}{\simeq} H\right) = \prod_{s \in [2,n)} \frac{(\alpha_s + p_s - 1)_{p_s}(\beta_s + q_s - 1)_{q_s}}{(\alpha_s + \beta_s + p_s + q_s - 1)_{p_s + q_s}}$$

where α_s and β_s are the first and second parameters of the Beta random variables and p_s, q_s are defined in Lemma A.2.

Following the steps in (3.18)-(3.20), we can simplify q_s as

(A.3)
$$q_s = d_{[s-1]}(H) - a_{[2]} - 2(m_{[s-1]} - 2) - m_s \quad \text{for } s \ge 3,$$

with $q_2 = d_1(H) - a_1$, and q_s satisfies

(A.4)
$$p_s + q_s = q_{s+1} + m_{s+1}$$
 for $s \ge 3$.

Now, we have all tools to adapt Proposition 3.2 to $PU^{(NSL)}$:

PROPOSITION A.4 (Equivalence of pre-collapsed model (B) and $PU^{(NSL)}$). For any graph $H \in \mathcal{H}_{m_{[n]}}$,

(A.5)
$$\mathbb{P}_m\left(\mathrm{PA}_n^{\scriptscriptstyle(\mathrm{B})}(\boldsymbol{m},1,\delta)\overset{\star}{\simeq} H\right) = \mathbb{P}_m\left(\mathrm{PU}_{m_{[n]}}^{\scriptscriptstyle(\mathrm{NSL})}(\boldsymbol{1},\boldsymbol{\psi})\overset{\star}{\simeq} H\right).$$

PROOF. Following a similar calculation as the one leading to (3.10),

$$\mathbb{P}_{m}\left(\mathrm{PA}_{n}^{(\mathrm{B})}(\boldsymbol{m},1,\delta) \stackrel{\star}{\simeq} H\right) = \prod_{u \in [n]} \prod_{j \in [m_{u}]} \prod_{i=a_{m_{[u-1]}+j}}^{d_{m_{[u-1]}+j}(H)-1} \left(i + \frac{\delta}{m_{u}}\right)$$
(A.6)
$$\times \prod_{u \in [3,n]} \prod_{j \in [m_{u}]} \frac{1}{a_{[2]} + 2(m_{[u-1]} + j - 3) + \left((u-1) + \frac{j}{m_{u}}\right)\delta}.$$

For model (B), we consider $\mathrm{PU}_{m_{[n]}}^{(\mathrm{NSL})}(\mathbf{1}, \boldsymbol{\psi})$, where $\boldsymbol{\psi}$ is the sequence of Beta variables defined in (3.1) and (3.2). Therefore, $p_s = d_s(H) - 1$ for $s \ge 1$, and

$$q_s = d_{[s-1]}(H) - a_{[2]} - 2(s-3) - 1$$
 for $s \ge 3$

By Corollary A.3,

(A.7)
$$\mathbb{P}_{m}\left(\mathrm{PU}_{m_{[n]}}^{(\mathrm{NSL})}(\mathbf{1}, \psi) \stackrel{\star}{\simeq} H\right) = \prod_{s \in [2, m_{[n]})} \frac{(\alpha_{s} + p_{s} - 1)_{p_{s}}(\beta_{s} + q_{s} - 1)_{q_{s}}}{(\alpha_{s} + \beta_{s} + p_{s} + q_{s} - 1)_{p_{s} + q_{s}}},$$

where α_s and β_s are the first and second parameters of the Beta variable ψ_s defined in (3.1) and (3.2) respectively. Then, by (A.4), the recursion (3.24) again holds. Now, following the calculations in (3.25)–(3.27), and substituting the values of $\alpha_s, \beta_s, p_s, q_s$, it follows immediately that

$$\mathbb{P}_{m}\left(\mathrm{PU}_{m_{[n]}}^{(\mathrm{NSL})}(\mathbf{1}, \boldsymbol{\psi}) \stackrel{\star}{\simeq} H\right) = \prod_{u \in [n]} \prod_{j \in [m_{u]}} \prod_{i=a_{m_{[u-1]}+j}}^{d_{m_{[u-1]}+j}(H)-1} \left(i + \frac{\delta}{m_{u}}\right)$$
(A.8)
$$\times \prod_{u \in [3,n]} \prod_{j \in [m_{u]}} \frac{1}{a_{[2]} + 2(m_{[u-1]} + j - 3) + \left((u - 1) + \frac{j}{m_{u}}\right)\delta},$$
as required.

as required.

Proof of Theorem 3.9. Theorem 3.9 follows immediately from Proposition A.4 in exactly the same way as Theorem 3.1 follows from Proposition 3.2.

Proof of Theorem 3.10. Conditionally on m, the distribution of model (D) is

$$\mathbb{P}_{m}\left(\mathrm{PA}_{n}^{(\mathrm{D})}(\boldsymbol{m},\delta) \stackrel{\star}{\simeq} H\right) = \prod_{u \in [3,n]} \prod_{j \in [m_{u}]} \frac{d_{v(u,j)}(u,j-1) + \delta}{a_{[2]} + 2(m_{[u-1]} - 2) + (j-1) + (u-1)\delta} ,$$

where v(u,j) is the vertex in [u-1] to which u connects with its j-th edge and $d_v(u,j)$ denotes the degree of the vertex v in $PA_{u,j}^{(D)}(\boldsymbol{m},\delta)$. Rearranging the numerators of RHS of (A.9),

(A.10)
$$\mathbb{P}_{m}\left(\mathrm{PA}_{n}^{(\mathrm{D})}(\boldsymbol{m},\delta) \stackrel{\star}{\simeq} H\right)$$
$$= \prod_{u \in [n]} \prod_{i=f_{u}}^{d_{u}(H)-1} (i+\delta) \prod_{u \in [3,n]} \prod_{j=1}^{m_{u}} \frac{1}{a_{[2]} + 2(m_{[u-1]} - 2) + (j-1) + (u-1)\delta}$$

For model (B), we use the ψ defined in (3.29). Using Corollary A.3, we calculate the conditional distribution of $\mathrm{PU}_n^{\scriptscriptstyle\mathrm{(SL)}}(\boldsymbol{m},\psi)$ as

(A.11)
$$\mathbb{P}_m\left(\mathrm{PU}_n^{(\mathrm{NSL})}(\boldsymbol{m}, \boldsymbol{\psi}) \stackrel{\star}{\simeq} H\right) = \prod_{s \in [2,n)} \frac{(\alpha_s + p_s - 1)_{p_s}(\beta_s + q_s - 1)_{q_s}}{(\alpha_s + \beta_s + p_s + q_s - 1)_{p_s + q_s}},$$

where, for s > 3,

(A.12)
$$\begin{aligned} \alpha_s &= m_s + \delta, & \text{and} & \beta_s &= a_{[2]} + 2\left(m_{[s]} - 2\right) - m_s + (s - 1)\delta, \\ p_s &= d_s(H) - f_s, & \text{and} & q_s &= d_{[s-1]}(H) - \left(a_{[2]} + 2\left(m_{[s]} - 2\right)\right) + m_s. \end{aligned}$$

Therefore, for $s \ge 3$, the recursion relation in (3.24) becomes

(A.13)
$$\alpha_s + \beta_s = \beta_{s+1} - m_{s+1}, \qquad p_s + q_s = q_{s+1} + m_{s+1}$$

This gives us all the necessary tools to prove Theorem 3.10. From the recursion relation in (A.13), it follows that

(A.14)
$$\frac{(\beta_s + q_s - 1)_{q_s}}{(\alpha_s + \beta_s + p_s + q_s - 1)_{q_s + p_s}} = \frac{(\beta_s + q_s - 1)_{q_s}}{(\beta_{s+1} + q_{s+1} - 1)_{q_{s+1} + m_{s+1}}} = \frac{1}{(\beta_{s+1} - 1)_{m_{s+1}}} \frac{(\beta_s + q_s - 1)_{q_s}}{(\beta_{s+1} + q_{s+1} - 1)_{q_{s+1}}}$$

On the other hand, the first factor in RHS of (A.11) can be rewritten as

$$\frac{(\alpha_2 + p_2 - 1)_{p_2}(\beta_2 + q_2 - 1)_{q_2}}{(\alpha_2 + \beta_2 + p_2 + q_2)_{p_2 + q_2}} = \frac{1}{(\beta_3 + q_3 - 1)_{q_3}} \prod_{i=0}^{m_3 - 1} \frac{1}{a_{[2]} + i + 2\delta} \prod_{i \in [2]} \prod_{j=f_i}^{d_i(H) - 1} (j + \delta).$$

Hence substituting the simplifications obtained from (A.14) and (A.15) in (A.11),

$$\mathbb{P}_{m}\left(\mathrm{PU}_{n}^{(\mathrm{NSL})}(\boldsymbol{m},\boldsymbol{\psi})\overset{\star}{\simeq}H\right)$$
(A.16)
$$=\prod_{u\in[n]}\prod_{i=f_{u}}^{d_{u}(H)-1}(i+\delta)\prod_{u\in[3,n]}\prod_{j=1}^{m_{u}}\frac{1}{a_{[2]}+2(m_{[u-1]}-2)+(j-1)+(u-1)\delta},$$
as required.

as required.

APPENDIX B: ADAPTED PROOFS FOR PRELIMINARY RESULTS

In this appendix, we prove Proposition 4.2, and the regularity of the Random Pólya point tree in Lemma 4.7. Versions of these results were proved in [8], and we have adapted those proofs to our settings of PAMs with random out-degrees. Before proving the position concentration result in Proposition 4.2, we prove an auxiliary lemma that we will use several times, and which is a direct application of the dominated convergence theorem and strong law of large numbers:

LEMMA B.1. Let X_1, X_2, \ldots be a sequence of i.i.d. random variables with $X_1 > c$ for some c > 0 a.s. and finite mean. Then, with $X_{[n]} = X_1 + \cdots + X_n$,

(B.1)
$$\mathbb{E}\left[\frac{1}{X_{[n]}}\right] = (1+o(1))\frac{1}{n\mathbb{E}[X_1]}.$$

PROOF. Note that $X_i \ge c > 0$, hence both $1/\mathbb{E}[X]$ and $n/X_{[n]}$ have upper bounds 1/c. By the strong law of large numbers,

$$\frac{X_{[n]}}{n} \stackrel{a.s.}{\to} \mathbb{E}[X_1].$$

Since $\mathbb{E}[X_1] > 0$ and $n/X_{[n]} \leq 1/c$, by the dominated convergence theorem,

$$\mathbb{E}\left[\frac{n}{X_{[n]}}\right] = (1 + o(1))\frac{1}{\mathbb{E}[X_1]}$$

Hence the lemma follows immediately.

Proof of Proposition 4.2. Proof for CPU: We follow the line of proof provided in [8, Lemma 3.1] with the adaptations as required for our case of i.i.d. out-degrees.

Fix $\omega, \varepsilon > 0$, and let $\bar{\omega} = \log(1 + \omega)$. We use the definition of $\mathcal{S}_k^{(n)}$ to bound the error in estimating $\mathcal{S}_k^{(n)}$ by $\left(\frac{k}{n}\right)^{\chi}$. For all $k \in [n-1]$,

(B.2)
$$S_k^{(n)} = \prod_{l=m_{[k]}+1}^{m_{[n]}} (1-\psi_l) = \exp\left[\sum_{l=m_{[k]}+1}^{m_{[n]}} \log(1-\psi_l)\right],$$

with $S_n^{(n)} \equiv 1$. We concentrate on the argument of the exponential in (B.2). Note that

(B.3)
$$\operatorname{Var}\left(\log(1-\psi_{l})\right) \leq \mathbb{E}\left[\log^{2}(1-\psi_{l})\right] \leq \mathbb{E}\left[\frac{\psi_{l}^{2}}{(1-\psi_{l})^{2}}\right]$$

By (B.3) and Kolmogorov's maximal inequality, (B.4)

$$\mathbb{P}\Big(\max_{l\in[m_{[n]}-1]}\Big|\sum_{k=l+1}^{m_{[n]}}\log(1-\psi_k)-\mathbb{E}\Big[\sum_{k=l+1}^{m_{[n]}}\log(1-\psi_k)\Big]\Big| \ge \frac{\bar{\omega}}{2}\Big) \le \frac{4}{\bar{\omega}^2}\mathbb{E}\left[\sum_{i=2}^{m_{[n]}}\mathbb{E}_m\left[\frac{\psi_i^2}{(1-\psi_i)^2}\right]\right]$$

Equation (B.4) shows that the maximum of the fluctuations of the argument in (B.2) can be bounded by the variances of the singles terms. By properties of the Beta distribution, and recalling that, for u = 3, ..., n and $j \in [m_u]$,

$$\psi_{m_{[u-1]}+j} \sim \mathsf{Beta}\left(1 + \frac{\delta}{m_u}, a_{[2]} + 2\left(m_{[u-1]} + j - 2\right) + (u-1)\delta + \frac{j-1}{m_u}\delta - 1\right),$$

we can bound, for u > 2 and $j \in [m_u]$,

(B.5)
$$\mathbb{E}_{m}\left[\frac{\psi_{m_{[u-1]}+j}^{2}}{\left(1-\psi_{m_{[u-1]}+j}\right)^{2}}\right] = \mathcal{O}\left(m_{[u-1]+j}^{-2}\right)$$

Notice that $m_{[n]} \ge n$ for all $n \ge 1$, and hence $m_{[n]} \to \infty$ as $n \to \infty$. Equation (B.5) assures us that the sum on the RHS of (B.4) is finite as $n \to \infty$. Therefore, we can fix $N_1(\bar{\omega}) \in \mathbb{N}$ such that $\mathbb{E}\left[\sum_{i=N_1}^{\infty} \frac{\psi_i^2}{(1-\psi_i)^2}\right] \le \varepsilon \bar{\omega}^2/4$. As a consequence, bounding the sum on the RHS of (B.4) by the tail of the series, for $n > N_1$,

(B.6)
$$\mathbb{P}\left(\max_{i\in[m_{[n]}]\setminus[m_{[N_1]}]}\left|\sum_{l=i+1}^{m_{[n]}}\log(1-\psi_l)-\mathbb{E}\left[\sum_{l=i+1}^{m_{[n]}}\log(1-\psi_l)\right]\right|\geq\frac{\bar{\omega}}{2}\right)$$
$$\leq\frac{4}{\bar{\omega}^2}\mathbb{E}\left[\sum_{i=N_1}^{\infty}\frac{\psi_i^2}{(1-\psi_i)^2}\right]\leq\varepsilon.$$

Next, we wish to compare the expectations of $\sum_{k=i}^{m_{[n]}} \log(1-\psi_k)$ and $\sum_{k=i}^{m_{[n]}} \psi_k$ for n large enough. Using the fact that, for $x \in (0, 1)$,

$$|\log(1-x) - x| \le x^2/(1-x),$$

and using (B.5) we bound, for $m_{[N_1]} \leq i \leq m_{[n]}$,

(B.7)
$$\left| \mathbb{E} \Big[\sum_{l=i+1}^{m_{[n]}} \log(1-\psi_l) \Big] - \mathbb{E} \Big[\sum_{l=i+1}^{m_{[n]}} \psi_l \Big] \right| \le \mathbb{E} \left[\sum_{l=i+1}^{m_{[n]}} \frac{\psi_l^2}{(1-\psi_l)} \right] < \infty.$$

Similarly, there exists $N_2(\omega) \in \mathbb{N}$ such that $\mathbb{E}\left[\sum_{l=N_2}^{\infty} \psi_l^2/(1-\psi_l)\right] \leq \bar{\omega}/3$ and $1/\sqrt{N_2} \leq \bar{\omega}/6$. On the other hand, for u > 2 and $j \in [m_u]$,

(B.8)
$$\mathbb{E}_{m}\left[\psi_{m_{[u-1]}+j}\right] = \frac{1 + \frac{\delta}{m_{u}}}{a_{[2]} + 2\left[m_{[u-1]} + j - 2\right] + (u - 1)\delta + \frac{j}{m_{u}}\delta} = \frac{1 + \frac{\delta}{m_{u}}}{2m_{[u-1]} + (u - 1)\delta} \left(1 + o\left(1\right)\right).$$

Therefore, for all u > 2,

$$\mathbb{E}\Big[\sum_{j=1}^{m_u} \psi_{m_{[u-1]}+j}\Big] = \mathbb{E}\Big[\mathbb{E}_m\Big[\sum_{j=1}^{m_u} \psi_{m_{[u-1]}+j}\Big]\Big] = \mathbb{E}\left[\frac{m_u+\delta}{2m_{[u-1]}+\delta}\right](1+o(1))$$
(B.9)
$$= (\mathbb{E}[M]+\delta)\mathbb{E}\left[\frac{1}{2m_{[u-1]}+\delta}\right](1+o(1)) = \frac{\mathbb{E}[M]+\delta}{(u-1)(2\mathbb{E}[M]+\delta)}(1+o(1))$$

$$= \frac{\chi}{u-1}(1+o(1)),$$

by Lemma B.1. Now, using the bounds in (B.7) and (B.9), for all $N_2 \le k \le n$,

(B.10)
$$\left| \mathbb{E} \left[\sum_{i=m_{[k]}+1}^{m_{[n]}} \log(1-\psi_i) \right] - \chi \log\left(\frac{k}{n}\right) \right| \le o\left(k^{-1/2}\right) + \bar{\omega}/3 \le \frac{\bar{\omega}}{2}$$

As a consequence, for $n > N_2$,

(B.11)
$$\max_{k \in (N_2, n]} \left| \mathbb{E} \left[\sum_{i=m_{[k]}+1}^{m_{[n]}} \log(1-\psi_i) \right] - \chi \log\left(\frac{k}{n}\right) \right| \le \frac{\bar{\omega}}{2}.$$

Let $N_0 = \max\{N_1, N_2\}$. By (B.11) and (B.6), for $n > N_0$,

(B.12)
$$\mathbb{P}\left(\max_{k\in(N_0,n]}\left|\sum_{i=m_{[k]}+1}^{m_{[n]}}\log(1-\psi_i)-\chi\log\left(\frac{k}{n}\right)\right|\geq\bar{\omega}\right)\leq\varepsilon.$$

Recalling that $\log S_k^{(n)} = \sum_{i=m_{[k]}+1}^{m_{[n]}} \log(1-\psi_i)$ and $\bar{\omega} = \log(1+\omega)$, (B.12) implies that with probability at least $1-\varepsilon$, for every $i = (N_0, n]$,

(B.13)
$$\frac{1}{1+\omega} \left(\frac{k}{n}\right)^{\chi} \le \mathcal{S}_k^{(n)} \le (1+\omega) \left(\frac{k}{n}\right)^{\chi}.$$

Since, $\frac{1}{1+\omega} \ge 1 - \omega$, we obtain from (B.13) that

(B.14)
$$\mathbb{P}\Big(\bigcap_{u\in(N_0,n]}\Big\{\Big|\mathcal{S}_u^{(n)}-\Big(\frac{u}{n}\Big)^{\chi}\Big|\leq\omega\left(\frac{u}{n}\right)^{\chi}\Big\}\Big)\geq 1-\varepsilon,$$

which proves (4.8).

Finally, to prove (4.7), we observe that, for fixed $\omega > 0$ and $\varepsilon > 0$, $\left(\frac{N_0}{n}\right)^{\chi} \le \omega/3$ for large enough n. Now,

(B.15)
$$\max_{u \in [N_0]} \left| \mathcal{S}_u^{(n)} - \left(\frac{u}{n}\right)^{\chi} \right| \le \mathcal{S}_{N_0}^{(n)} + \left(\frac{N_0}{n}\right)^{\chi} \le \omega,$$

as required.

Proof for PU: In Pólya urn graphs, conditionally on m, for $u \ge 2$,

(B.16)
$$\psi_u \sim \text{Beta}\left(m_u + \delta, a_{[2]} + 2(m_{[u-1]} - 2) + m_u + (u-1)\delta\right).$$

We prove the position concentration lemma for PU following the proof for CPU. Using Kolmogorov's inequality as in (B.4), (B.17)

$$\mathbb{P}\Big(\max_{l\in[n-1]\setminus[n_1]}\Big|\sum_{k=l+1}^n\log(1-\psi_k)-\mathbb{E}\Big[\sum_{k=l+1}^n\log(1-\psi_k)\Big]\Big| \ge \frac{\bar{\omega}}{2}\Big) \le \frac{4}{\bar{\omega}^2}\mathbb{E}\Big[\sum_{i=n_1}^n\frac{\psi_i^2}{(1-\psi_i)^2}\Big]$$

Now similarly as in (B.5) we first bound $\mathbb{E}\left[\frac{\psi_i^2}{(1-\psi_i)^2}\right]$ by i^{-p} and thus we can choose n_1 large enough such that the RHS of (B.17) is smaller than ε . Then,

(B.18)
$$\mathbb{E}\left[\frac{\psi_u^2}{(1-\psi_u)^2}\right] \le \mathbb{E}\left[\left(\frac{m_u+\delta}{a_{[2]}+2(m_{[u-1]}-3)+m_u+(u-1)\delta}\right)^2\right]$$

Since the random variable in the RHS of (B.18) is bounded above by 1, we can bound its second moment by its (1 + p)-th moment and obtain the following bound on the LHS of (B.18):

(B.19)
$$\mathbb{E}\left[\frac{\psi_{u}^{2}}{(1-\psi_{u})^{2}}\right] \leq \mathbb{E}\left[\left(\frac{m_{u}+\delta}{a_{[2]}+2(m_{[u-1]}-3)+m_{u}+(u-1)\delta}\right)^{1+p}\right] \\ \leq \mathbb{E}\left[\left(\frac{m_{u}+\delta}{a_{[2]}+2(m_{[u-1]}-3)+(u-1)\delta}\right)^{1+p}\right].$$

Now the numerator and the denominator are independent of each other. Using the law of large numbers and Lemma B.1,

(B.20)
$$\mathbb{E}\left[\left(\frac{1}{a_{[2]}+2(m_{[u-1]}-3)+(u-1)\delta}\right)^{1+p}\right] = (1+o(1))\frac{1}{u^{1+p}(2\mathbb{E}[M]+\delta)^{1+p}}$$

Since m_u has finite (1+p)-th moment, there exists a constant $\xi_0 > 0$ such that

(B.21)
$$\mathbb{E}\left[\frac{\psi_u^2}{(1-\psi_u)^2}\right] \le \xi_0 u^{-(1+p)} ,$$

which replaces (B.5). Now it remains to adapt a similar result to (B.9) to complete the proof for PU. By (B.16),

(B.22)
$$\mathbb{E}[\psi_u] = \mathbb{E}\left[\frac{m_u + \delta}{a_{[2]} + 2(m_{[u]} - 2) + u\delta}\right].$$

Note that the numerator and the denominator are not independent here. We can bound the RHS from above as

$$\mathbb{E}\left[\frac{m_u + \delta}{a_{[2]} + 2(m_{[u]} - 2) + u\delta}\right] \le \mathbb{E}\left[\frac{m_u + \delta}{a_{[2]} + 2(m_{[u-1]} - 2) + (u-1)\delta}\right] = \frac{\chi}{u-1}(1+o(1)).$$

On the other hand for the lower bound, we truncate m_u at $\log(u)$ as

(B.24)
$$\mathbb{E}\left[\frac{m_u + \delta}{a_{[2]} + 2(m_{[u]} - 2) + u\delta}\right] \ge \mathbb{E}\left[\frac{m_u \mathbb{1}_{\{m_u \le \log u\}} + \delta}{a_{[2]} + 2(m_{[u-1]} - 2) + 2\log(u) + u\delta}\right]$$

Again we can split the numerator and the denominator since they are now independent. Since m_u has finite mean, $\mathbb{E}[m_u \mathbb{1}_{\{m_u \le \log u\}} + \delta] = (\mathbb{E}[M] + \delta)(1 + o(1))$. By Lemma B.1,

$$\mathbb{E}\left[\frac{m_u + \delta}{a_{[2]} + 2(m_{[u]} - 2) + u\delta}\right] \ge \frac{\mathbb{E}[M] + \delta}{(u - 1)(2\mathbb{E}[M] + \delta)}(1 + o(1)) = \frac{\chi}{u - 1}(1 + o(1)).$$

Hence from (B.23) and (B.25), we obtain a similar result as in (B.8) as

(B.26)
$$\mathbb{E}[\psi_u] = \mathbb{E}\left[\frac{m_u + \delta}{a_{[2]} + 2(m_{[u]} - 2) + u\delta}\right] = \frac{\chi}{u - 1}(1 + o(1)).$$

Proof of Lemma 4.7. The lemma holds trivially for r = 0 and $\varepsilon/4$, i.e., with probability at least $1 - \varepsilon/4$,

- (1) $A_{\emptyset} \ge \eta\left(\frac{\varepsilon}{4}, 0\right)$, where $\eta\left(\frac{\varepsilon}{4}, 0\right)$ is a positive value depending on ε and r = 0;
- (2) $|B_0^{(G)}(\emptyset)| = 1;$
- (3) $\Gamma_{\emptyset} \leq K\left(\frac{\varepsilon}{4}, 0\right)$, where $K\left(\frac{\varepsilon}{4}, 0\right)$ is a natural number depending on ε and r = 0.

We thus prove the lemma for r = 1. The proof for $r \ge 2$ then easily follows by induction on r.

Let $U_{(M)}$ be the smallest order statistic of U_1, \ldots, U_M , i.e., $U_{(M)} = \min\{U_1, \ldots, U_M\}$ and $\eta \leq \eta \left(\frac{\varepsilon}{4}, 0\right)$. To prove (1), it is enough to show that

$$\mathbb{P}\left(\left|A_{\varnothing}U_{\scriptscriptstyle (M)}^{1/\chi} < \eta \right| \left|A_{\varnothing} > \eta(\varepsilon/4, 0)\right) \le \frac{\varepsilon}{4}.\right.$$

First, we condition on M and find a lower bound on the LHS. Next, we take an expectation over M. Using the density of $U_{(M)}$ and Taylor's inequality,

$$\mathbb{P}\left(\left.A_{\varnothing}U_{\scriptscriptstyle(M)}^{1/\chi} < \eta \right| \ M = m, A_{\varnothing} \ge \eta(\varepsilon/4, 0)\right) = \int_{\eta(\varepsilon/4, 0)}^{1} \left(1 - \left(1 - \left(\frac{\eta}{x}\right)^{\chi}\right)^{m}\right) \, dx$$
(B.27)
$$\leq \int_{\eta(\varepsilon/4, 0)}^{1} m\left(\frac{\eta}{x}\right)^{\chi} \, dx \le \frac{m\eta^{\chi}}{1 - \chi}.$$

Integrating over m and choosing η suitably,

(B.28)
$$\mathbb{P}\left(\left|A_{\varnothing}U_{(M)}^{1/\chi} < \eta \right| \left|A_{\varnothing} \ge \eta(\varepsilon/4, 0)\right| \le \frac{\eta^{\chi}}{2\mathbb{E}[M] + \eta} = \frac{\varepsilon}{4}$$

Denote the η in (B.28) by $\eta(\varepsilon, 1)$.

Next, we prove (2). The number of children of the root is distributed as $(M + \Lambda)$, where Λ is the total number of points in an inhomogeneous Poisson process with intensity $\rho_{\emptyset}(x)$. Since M is uniformly integrable, there exists m_0 such that $\mathbb{P}(M \le m_0) \ge 1 - \frac{\varepsilon}{8}$. Moreover, it can also be shown that

$$\mathbb{P}(M^{(0)} > m_0) = \frac{1}{\mathbb{E}[M]} \sum_{k \ge m_0 + 1} k \mathbb{P}(M = k) = \frac{\mathbb{E}\left[M \mathbb{1}_{\{M > m_0\}}\right]}{\mathbb{E}[M]} \le \frac{\varepsilon}{8}.$$

Similar results also hold for $M^{(\delta)}$, and are useful for the induction step when r > 1. Now, we have the parameter of Λ bounded above and hence there exists $C(\varepsilon, 1)$ such that

$$M + \Lambda \le C(\varepsilon, 1)$$

We can choose $K(\varepsilon, 1) < \infty$ such that $\mathbb{P}(\Gamma_{\omega} \leq K(\varepsilon, 1) \text{ for all } \omega \in B_1^{(G)}(\emptyset)) \geq 1 - \frac{\varepsilon}{4}$. \Box

APPENDIX C: COUPLING BETWEEN MODEL (D) AND (F)

The proof to Proposition 6.4 follows exactly the same line as that of Proposition 6.1. To prove Proposition 6.1, we needed Lemma 6.2 and 6.3. Lemma 6.3 proves a property of model (D), and thus we do not need to adapt this lemma. Additionally, subject to Lemma 6.2, the proof of Proposition 6.1 does not use any property of model (E). Thus, if we manage to adapt Lemma 6.2 to model (F), the proof of Proposition 6.4 follows immediately.

For the proof of Lemma 6.2 for model (F), we construct a similar coupling between model (D) and (F) as we have constructed for model (D) and (E) in Section 6. Then we claim that Lemma 6.2 also holds for model (F).

Proof of Lemma 6.2 for model (F). Note that for model (F), there cannot be more than one edge between two vertices. Let p'_l denote the conditional edge-connection probability of connecting the *n*-th vertex to vertex *v* with its *l*-th edge and Q'_k denote the probability of connecting *k* edges from *n* to *v* in model (F). Then the total variation distance in (6.20) can be rewritten for model (F) as

(C.1)
$$d_{\text{TV}}(D_1^{(n)}, D_2^{(n)}) = \frac{1}{2} \sum_{k=0}^{m_n} |Q_k' - T_k| = \frac{1}{2} \Big[|Q_0' - T_0| + |Q_1' - T_1| + \sum_{k=2}^{m_n} T_k \Big] .$$

Since $\sum_{k=0}^{1} Q'_k = \sum_{k=0}^{m_n} T_k = 1$, we bound $|Q'_1 - T_1|$ as

(C.2)
$$|Q'_1 - T_1| = \left| (1 - Q'_0) - (1 - T_0 - \sum_{k=2}^{m_n} T_k) \right| \le |Q'_0 - T_0| + \sum_{k=2}^{m_n} T_k ,$$

and the total variation distance can be bounded by

(C.3)
$$d_{\scriptscriptstyle \mathrm{TV}}\left(D_1^{\scriptscriptstyle (n)}, D_2^{\scriptscriptstyle (n)}\right) \le |Q_0' - T_0| + \sum_{k=2}^{m_n} T_k$$

Therefore we do not need to bound $|Q'_1 - T_1|$, if we manage to bound the RHS of (C.3). From (6.23), we already have the required bound for $\sum_{k=2}^{m_n} T_k$. For the other term in the RHS of (C.3), we use a triangle inequality argument to obtain

(C.4)
$$|Q'_0 - T_0| \le |Q_0 - T_0| + |Q'_0 - Q_0|,$$

where Q_k is defined in (6.18). Therefore, again from (6.22), we have the desired bound for the first term in the RHS of (C.4). Now we are left to show that the other term also has the same upper bound. Note that $p' \le p'_l$ for all $l \ge 1$ and

(C.5)
$$p'_{m_n} \le \frac{d+\delta}{\sum\limits_{n \ne u} (d_v(n-1)+\delta)} ,$$

where $n \not\sim u$ implies that u and n are not connected by an edge. Let $\zeta_0 = \delta + \inf \operatorname{supp}(M) > 0$. Then for all $u \in [n-1]$, $d_v(n-1) + \delta \geq \zeta_0$ and the denominator of the RHS of (C.5) can be further lower bounded by $(n - m_n)\zeta_0$. On the event space \mathcal{E} , $m_n \leq n^{1-\varrho/2}$. Therefore there exists another constant $\zeta_1 > 0$, such that on \mathcal{E} , the denominator can be further lower bounded by $n\zeta_1$ and the LHS of (C.5) can be upper bounded by $(d + \delta)/(n\zeta_1)$.

Therefore using the telescoping sum bound in (6.21), we have upper bound for $|Q'_0 - Q_0|$.

APPENDIX D: TAIL DEGREE DISTRIBUTION OF THE NEIGHBOURS

Even though Corollary 1.6(b) is about the tails of the 0 and Y neighbours, since the proof follows the same way, we prove the tail of the root also. In Corollary 1.6(a), we have proved that the degree of a randomly chosen 0 child of the root is given by $1 + M^{(\delta)} + Y(M^{(\delta)} + 1, A_{\circ})$, where A_{\circ} is the age of a randomly chosen 0 neighbour of the root; the degree of a randomly chosen Y child of the root is given by $M^{(0)} + Y(M^{(0)}, A_{\gamma})$, where A_{\circ} is the age of a randomly chosen Y neighbour of the root. Lastly for the root, the degree is given by $M + Y(M, U_{\emptyset})$ where U_{\emptyset} is a uniform random variable on [0, 1]. Therefore for all $t \in \mathbb{N}$,

$$\mathbb{P}(1 + M^{(\delta)} + Y(M^{(\delta)} + 1, A_{\circ}) = t) = \sum_{m=1}^{t-1} \mathbb{P}(M^{(\delta)} = m)$$

(D.1)

$$\int_0^1 \mathbb{P}(Y(m+1,a) = t - m - 1) f_0(a) \, da \,,$$

$$\mathbb{P}(M^{(0)} + Y(M^{(0)}, A_{\mathsf{Y}}) = t) = \sum_{m=1}^{t} \mathbb{P}(M^{(0)} = m) \int_{0}^{1} \mathbb{P}(Y(m, a) = t - m) f_{\mathsf{Y}}(a) \, da$$

and
$$\mathbb{P}(M + Y(M, U_{\varnothing}) = t) = \sum_{m=1}^{t} \mathbb{P}(M = m) \int_{0}^{1} \mathbb{P}(Y(m, a) = t - m) da$$
,

where f_{\circ} and f_{\vee} are the age distribution functions of a random \circ and \vee child of the root. If we manage to explicitly calculate the densities f_{\circ} and f_{\vee} , then we are left to find out $\mathbb{P}(Y(m,a) = t - m)$. Y(m,a) is a mixed Poisson random variable with mixing parameter $\Gamma_{in}(m)\lambda(a)$. Therefore

$$\mathbb{P}(Y(m,a) = t - m) = \int_{0}^{\infty} \frac{(\gamma\lambda(a))^{t-m}}{(t-m)!} \exp\left(-\gamma\lambda(a)\right) \frac{1}{\Gamma(m+\delta)} \gamma^{(m+\delta)-1} \exp\left(-\gamma\right) d\gamma$$
(D.2)
$$= \frac{\lambda(a)^{t-m}}{(t-m)!\Gamma(m+\delta)} \int_{0}^{\infty} \gamma^{(t+\delta)-1} \exp\left(-\gamma(\lambda(a)+1)\right) d\gamma$$

$$= \frac{\Gamma(t+\delta)}{(t-m)!\Gamma(m+\delta)} \lambda(a)^{t-m} (1+\lambda(a))^{t+\delta}.$$

Substituting the value of $\lambda(a) = a^{-(1-\chi)}(1-a^{1-\chi})$ we obtain a compact form of the probability on the LHS of (D.2) as

(D.3)
$$\mathbb{P}(Y(m,a) = t - m) = \frac{\Gamma(t+\delta)}{\Gamma(m+\delta)(t-m)!} (1 - a^{1-\chi})^{t-m} (a^{1-\chi})^{m+\delta}.$$

This is a similar result as [8, Lemma 5.2].

D.1. The age densities. We proceed to explicitly calculate the densities f_0 and f_y :

LEMMA D.1 (Age of random 0). The density of the age of a randomly chosen 0 child of the root is given by

(D.4)
$$f_{o}(a) = \frac{\chi}{1-\chi} a^{-(1-\chi)} (1-a^{1-\chi}).$$

PROOF. From the definition of the RPPT (M, δ) we have that A_0 is distributed as $U_1 U_2^{1/\chi}$ where U_1 and U_2 are independent uniform random variables on [0, 1]. Therefore,

(D.5)
$$\mathbb{P}(A_{o} \le a) = \mathbb{P}(U_{1}U_{2}^{1/\chi} \le a) = \int_{0}^{1} \mathbb{P}(U_{2} \le a^{\chi}x^{-\chi}) dx$$

Note that for $x \in [0, a]$ the probability in the RHS of (D.5) is always 1. For $x \in (a, 1]$ the probability is $a^{\chi}x^{-\chi}$. Therefore the RHS of (D.5) simplifies as

(D.6)
$$\mathbb{P}(A_{\circ} \le a) = a + a^{\chi} \int_{a}^{1} x^{-\chi} dx = a + \frac{a^{\chi}}{1 - \chi} (1 - a^{1 - \chi})$$

Now differentiating the RHS of (D.6) with respect to a,

(D.7)
$$f_{o}(a) = 1 + \frac{\chi}{1-\chi} a^{-(1-\chi)} (1-a^{1-\chi}) - \frac{1-\chi}{1-\chi} a^{\chi} a^{-\chi} = \frac{\chi}{1-\chi} a^{-(1-\chi)} (1-a^{1-\chi}).$$

Similarly, we have seen that conditionally on the existence of at least one γ child its age distribution is given in equation (1.21) of the main paper. We simplify the density here:

LEMMA D.2 (Age of random Y). The density of the age of a randomly chosen Y child of the root is given by

(D.8)
$$f_{\mathsf{Y}}(a) = a^{-\chi} \int_0^{a^{1-\chi}} x^{\tau_e - 2} (1-x)^{-1} dx ,$$

where $\tau_e = 3 + \delta/\mathbb{E}(M)$.

PROOF. From equation (1.21), if U_1 is a uniform random variable on [0, 1], then

(D.9)
$$f_{\mathsf{Y}}(a) = \mathbb{E}\left[(1-\chi) \frac{a^{-\chi} \mathbb{1}_{\{a \ge U_1\}}}{1-U_1^{1-\chi}} \right] = (1-\chi)a^{-\chi} \int_0^a (1-y^{1-\chi})^{-1} dy$$

Now we perform a change of variable operation by substituting $y^{1-\chi} = x$ in the RHS of (D.9).

(D.10)
$$f_{\mathsf{r}}(a) = a^{-\chi} \int_0^{a^{1-\chi}} x^{1/(1-\chi)-1} (1-x)^{-1} dx \, .$$

Substituting $\tau_e = 1 + \frac{1}{1-\chi}$ in the RHS of (D.10) we obtain the desired form of the density of a randomly chosen γ neighbour of the root.

D.2. Tail calculations. With Lemma D.1 and D.2 and equation (D.3) in hand, we evaluate the expressions in the RHS of (D.1) for the root, O and γ neighbours. Before proceeding with the calculation, note that

(D.11)
$$\begin{split} \mathbb{P}(M=t) &= L(t)t^{-\tau_M} ,\\ \mathbb{P}(M^{(0)}=t) &= L_1(t)t^{-(\tau_M-1)} ,\\ \text{and} \quad \mathbb{P}(M^{(\delta)}=t) &= L_2(t)t^{-(\tau_M-1)} . \end{split}$$

for some slowly varying functions L(t), $L_1(t)$ and $L_2(t)$. On the other hand, using Stirling's approximation,

(D.12)
$$\frac{\Gamma(m+\delta+k)}{\Gamma(m+\delta)} = m^k (1+\mathcal{O}(1/m))$$
$$\frac{\Gamma(t+\delta)}{\Gamma(t+\delta+k)} = t^{-k} (1+\mathcal{O}(1/m)).$$

(D.11) and (D.12) will be useful in several steps in the tail calculation of the degree distributions.

The root. First, we carry out the simplest of all these calculations. As we can see from (D.1), we need to integrate the RHS of (D.3). (D.13)

$$\int_0^1 \mathbb{P}(Y(m,a) = t - m) \, da = \frac{\Gamma(t+\delta)}{\Gamma(m+\delta)(t-m)!} \int_0^1 (1 - a^{1-\chi})^{t-m} (a^{1-\chi})^{m+\delta} \, da \, .$$

Now we do the same change of variable as we did in (D.9) and obtain

$$\int_0^1 \mathbb{P}(Y(m,a) = t - m) \, da = \frac{\Gamma(t+\delta)}{(1-\chi)\Gamma(m+\delta)(t-m)!} \int_0^1 u^{(m+\delta+\tau_e-1)-1} (1-u)^{t-m} \, du$$
(D.14)
$$= \frac{(\tau_e - 1)\Gamma(t+\delta)\Gamma(t-m+1)\Gamma(m+\delta+\tau_e-1)}{\Gamma(m+\delta)(t-m)!\Gamma(t+\delta+\tau_e)}$$

$$= \frac{(\tau_e - 1)\Gamma(t+\delta)\Gamma(m+\delta+\tau_e-1)}{\Gamma(m+\delta)\Gamma(t+\delta+\tau_e)}.$$

This matches with the expression presented in [18]. Now plugging this integral value in (D.1) and using (D.11) and (D.12),

$$\mathbb{P}(M + Y(M, U_{\varnothing}) = t) = \frac{(\tau_e - 1)\Gamma(t + \delta)}{\Gamma(t + \delta + \tau_e)} \sum_{m=1}^{t} \mathbb{P}(M = m) \frac{\Gamma(m + \delta + \tau_e - 1)}{\Gamma(m + \delta)}$$
(D.15)
$$= (\tau_e - 1)t^{-\tau_e} (1 + \mathcal{O}(1/t)) \sum_{m=1}^{t} L(m)m^{-(\tau_M - \tau_e + 1)} (1 + \mathcal{O}(1/m))$$

For $\tau_M > \tau_e$, the sum on the RHS of (D.15) is finite. On the other hand, for $\tau_M \le \tau_e$ the sum varies regularly as $t^{-(\tau_M - \tau_e)}$ and hence

(D.16)
$$\mathbb{P}(M+Y(M,U_{\varnothing})=t)=L^{(\varnothing)}(t)t^{-\tau},$$

where $\tau = \min\{\tau_M, \tau_e\}$ and $L^{(\emptyset)}(\cdot)$ is a slowly varying function.

The \circ **child**. From Lemma D.1, we have the expression for $f_{\circ}(a)$ and therefore the integral in (D.1) can be simplified as

$$\begin{aligned} \int_0^1 \mathbb{P}(Y(m,a) &= t - m) f_0(a) \, da \\ \text{(D.17)} \quad &= \frac{\Gamma(t+\delta)}{\Gamma(m+\delta)(t-m)!} \int_0^1 (1 - a^{1-\chi})^{t-m} (a^{1-\chi})^{m+\delta} \frac{\chi}{1-\chi} a^{-(1-\chi)} (1 - a^{1-\chi}) \, da \\ &= (\tau_e - 2) \frac{\Gamma(t+\delta)}{\Gamma(m+\delta)(t-m)!} \int_0^1 (1 - a^{1-\chi})^{t-m+1} (a^{1-\chi})^{m+\delta-1} \, da \, . \end{aligned}$$

Again we do the same change of variable and simplify the RHS of (D.17) as

$$\int_{0}^{1} \mathbb{P}(Y(m,a) = t - m) f_{0}(a) da$$

$$= (\tau_{e} - 2)(\tau_{e} - 1) \frac{\Gamma(t + \delta)}{\Gamma(m + \delta)(t - m)!} \int_{0}^{1} (1 - u)^{t - m + 1} u^{m + \delta + \tau_{e} - 3} da$$

$$= (\tau_{e} - 2)(\tau_{e} - 1) \frac{\Gamma(t + \delta)}{\Gamma(m + \delta)(t - m)!} \frac{\Gamma(m + \delta + \tau_{e} - 2)\Gamma(t - m + 2)}{\Gamma(t + \delta + \tau_{e})}$$

$$= (\tau_{e} - 2)(\tau_{e} - 1)(t - m + 1) \frac{\Gamma(t + \delta)\Gamma(m + \delta + \tau_{e} - 2)}{\Gamma(m + \delta)\Gamma(t + \delta + \tau_{e})}.$$

Therefore,

(D.19)
$$\int_0^1 \mathbb{P}(Y(m+1,a) = t - m - 1) f_o(a) \, da$$
$$= (\tau_e - 2)(\tau_e - 1)(t - m) \frac{\Gamma(t+\delta)\Gamma(m+\delta+\tau_e - 1)}{\Gamma(m+1+\delta)\Gamma(t+\delta+\tau_e)}.$$

Now substituting this integral value in (D.1), the sum in the RHS could be simplified as

$$\mathbb{P}(1 + M^{(\delta)} + Y(M^{(\delta)} + 1, A_{o}) = t)$$

(D.20)

$$=(\tau_e - 2)(\tau_e - 1)\frac{\Gamma(t+\delta)}{\Gamma(t+\delta+\tau_e)}\sum_{m=1}^{t-1}\mathbb{P}(M^{(\delta)} = m)(t-m)\frac{\Gamma(m+\delta+\tau_e - 1)}{\Gamma(m+1+\delta)}$$
$$=(\tau_e - 2)(\tau_e - 1)t^{-\tau_e}(1 + \mathcal{O}(1/t))\sum_{m=1}^{t-1}L_1(m)m^{-(\tau_M - \tau_e + 1)}(t-m)(1 + \mathcal{O}(1/m))$$

Now if $\tau_M \leq \tau_e$, the sum in (D.20) varies regularly as $t^{(\tau_e - \tau_M)+1}$. On the other hand for $\tau_M > \tau_e$, the sum varies regularly as t. Therefore there exists a slowly varying function $L^{(o)}(t)$ such that

(D.21)
$$\mathbb{P}(1 + M^{(\delta)} + Y(M^{(\delta)} + 1, A_{o}) = t) = L^{(o)}(t)t^{-\tau_{(o)}}$$

where $\tau_{(0)} = \min{\{\tau_M, \tau_e\}} - 1$. For $\tau_M > \tau_e$, $L^{(0)}$ turns out to be a constant. On the other hand, for $\tau_M \leq \tau_e$, using Karamata's Theorem [11, Proposition 1.5.8], $L^{(0)}$ can be shown to be asymptotically equal to $L_1/\tau_{(0)}$.

The γ *child*. The calculation for γ neighbours of the root is more involved. We first substitute the $f_{\gamma}(a)$ in (D.1)

$$\int_{0}^{1} \mathbb{P}(Y(m,a) = t - m) f_{\mathsf{Y}}(a) \, da$$

(D.22)
$$= \frac{\Gamma(t+\delta)}{\Gamma(m+\delta)(t-m)!} \int_{0}^{1} (1 - a^{1-\chi})^{t-m} (a^{1-\chi})^{(m+\delta)} a^{-\chi} \times \int_{0}^{a^{1-\chi}} x^{\tau_{e}-2} (1-x)^{-1} \, dx \, da \, .$$

We perform the usual change of variable $u = a^{1-\chi}$ and simplify the above equation as

(D.23)
$$\int_{0}^{1} \mathbb{P}(Y(m,a) = t - m) f_{\mathsf{v}}(a) da$$
$$= (\tau_{e} - 1) \frac{\Gamma(t+\delta)}{\Gamma(m+\delta)(t-m)!} \int_{0}^{1} (1-u)^{t-m} u^{m+\delta} \int_{0}^{u} x^{\tau_{e}-2} (1-x)^{-1} dx du$$

Using the fact that $(1-x)^{-1} = \sum\limits_{i \geq 0} x^i$ we simplify the inner integral as

(D.24)
$$\int_0^u x^{\tau_e - 2} (1 - x)^{-1} dx = \sum_{i \ge 0} \int_0^u x^{\tau_e + i - 2} dx = \sum_{i \ge 0} \frac{u^{\tau_e + i - 1}}{\tau_e + i - 1} \, .$$

Substituting the RHS of (D.24) in (D.23),

$$\begin{split} &\int_0^1 \mathbb{P}(Y(m,a)=t-m)f_{\mathsf{v}}(a)\,da \\ &= \frac{(\tau_e-1)\Gamma(t+\delta)}{\Gamma(m+\delta)(t-m)!}\sum_{i\geq 0}\frac{1}{\tau_e+i-1}\int_0^1(1-u)^{(t-m+1)-1}u^{(m+\delta+\tau_e+i)-1}\,du \\ (\mathbf{D.25}) &= (\tau_e-1)\frac{\Gamma(t+\delta)\Gamma(t-m+1)}{\Gamma(m+\delta)(t-m)!}\sum_{i\geq 0}\frac{1}{(\tau_e+i-1)}\frac{\Gamma(m+\delta+\tau_e+i)}{\Gamma(t+\delta+\tau_e+i+1)} \\ &= (\tau_e-1)\frac{\Gamma(t+\delta)}{\Gamma(m+\delta)}\sum_{i\geq 0}\frac{1}{(\tau_e+i-1)}\frac{\Gamma(m+\delta+\tau_e+i)}{\Gamma(t+\delta+\tau_e+i+1)} \\ &= \frac{\Gamma(t+\delta)\Gamma(m+\delta+\tau_e)}{\Gamma(m+\delta)\Gamma(t+\delta+\tau_e+1)} \\ &+ (\tau_e-1)\frac{\Gamma(t+\delta)}{\Gamma(m+\delta)}\sum_{i\geq 1}\frac{1}{(\tau_e+i-1)}\frac{\Gamma(m+\delta+\tau_e+i)}{\Gamma(t+\delta+\tau_e+i+1)} \,. \end{split}$$

Now the RHS of (D.25) is lower bounded by the first term and hence the RHS of (D.1) can be lower bounded as

$$\mathbb{P}(M^{(0)} + Y(M^{(0)}, A_{\mathsf{v}}) = t) \ge \sum_{m=1}^{t} \mathbb{P}(M^{(0)} = m) \frac{\Gamma(t+\delta)\Gamma(m+\delta+\tau_e)}{\Gamma(m+\delta)\Gamma(t+\delta+\tau_e+1)}$$

(D.26)
$$= t^{-(\tau_e+1)}(1 + \mathcal{O}(1/t)) \sum_{m=1}^{t} L_2(m)m^{-(\tau_M-\tau_e-1)}(1 + \mathcal{O}(1/m))$$

It can easily be shown that the RHS of (D.26) varies regularly with $t^{-(\tau_e+1)}$ if $\tau_M > \tau_e + 2$. When $\tau_M \leq \tau_e + 2$, it varies regularly with $t^{-(\tau_M-1)}$. Therefore the lower bound varies regularly with $t^{-\tau_{(Y)}}$, where $\tau_{(Y)} = \min\{\tau_M - 1, \tau_e + 1\}$ and some slowly varying function $L'_2(t)$. For $\tau_M > \tau_e + 2$, $L'_2(t)$ turns out to be a constant and for $\tau_M = \tau_e + 2$,

(D.27)
$$L_2'(t) = \sum_{t=1}^m L_2(m)/m$$

and lastly for $\tau_M < \tau_e + 2$, we use the same Karamata's Theorem as before to obtain that $L'_2(t) = \Theta(L_2(t))$. Now we move on to analyse the second term of RHS of (D.25).

$$\begin{split} &(\mathrm{D.28})\\ &(\tau_e-1)\frac{\Gamma(t+\delta)}{\Gamma(m+\delta)}\sum_{i\geq 1}\frac{1}{(\tau_e+i-1)}\frac{\Gamma(m+\delta+\tau_e+i)}{\Gamma(t+\delta+\tau_e+i+1)}\\ &=(\tau_e-1)\frac{\Gamma(t+\delta)}{\Gamma(m+\delta)}\sum_{i\geq 1}\frac{1}{(\tau_e+i-1)(m+\delta+\tau_e+i)}\frac{\Gamma(m+\delta+\tau_e+i+1)}{\Gamma(t+\delta+\tau_e+i+1)}\\ &=(\tau_e-1)\frac{\Gamma(t+\delta)\Gamma(m+\delta+\tau_e+2)}{\Gamma(m+\delta)\Gamma(t+\delta+\tau_e+2)}\sum_{i\geq 0}\frac{1}{(\tau_e+i)(m+\delta+\tau_e+i+1)}\prod_{j=0}^{i-1}\frac{m+\delta+\tau_e+j+2}{t+\delta+\tau_e+j+2}\,. \end{split}$$

LEMMA D.3. For any $m \ge 1$,

(D.29)
$$\sum_{i\geq 0} \frac{1}{(\tau_e+i)(m+\delta+\tau_e+i+1)} \prod_{j=0}^{i-1} \frac{m+\delta+\tau_e+j+2}{t+\delta+\tau_e+j+2} = \mathcal{O}(m^{-1}(2-\log(1-2m/(t+m+1))).$$

Subject to Lemma D.3, there exists $J_0 > 0$ such that the RHS of (D.28) can be upper bounded by $J_0(1 + \log(1 - 2m/(t + m + 1)))t^{-(\tau_e+2)}m^{(\tau_e+1)}(1 + \mathcal{O}(1/t))(1 + \mathcal{O}(1/m))$. Therefore, using (D.25), (D.26), (D.28) and Lemma D.3, the RHS of (D.1) can be upper bounded as

$$\mathbb{P}(M^{(0)} + Y(M^{(0)}, A_{\mathsf{v}}) = t)$$

$$\leq L_2'(t)t^{-\tau_{(\mathsf{v})}} + J_0 t^{-(\tau_e+2)}(1 + \mathcal{O}(1/t))$$
(D.30)
$$\times \sum_{k=1}^{t} L_2(m)(2 - \log(1 - 2m/(t + m + 1)))m^{-(\tau_M - \tau_e - 2)}(1 + \mathcal{O}(1/m))$$

$$\sum_{m=1}^{\infty} L_2(m)(2 - \log(1 - 2m/(t + m + 1)))m \quad (m + 0, (1 + O(1/m))).$$

When $\tau_M > \tau_e + 2$, the second sum is o(t) and hence the second term is $o(t^{-(\tau_e+1)})$. Therefore the tail degree distribution of the τ child varies regularly with $t^{-\tau_{(Y)}}$ when $\tau_M > \tau_e + 2$.

On the other hand, when $\tau_M \leq \tau_e + 2$, the upper bound can be shown to vary regularly with $t^{-\tau_{(Y)}}$ and the slowly varying function is again $\Theta(L'_2(t))$. Therefore considering $L'_2(t)$ as $L^{(Y)}(t)$, we can say that

(D.31)
$$\mathbb{P}(M^{(0)} + Y(M^{(0)}, A_{\mathsf{Y}}) = t) = \Theta(L^{(\mathsf{Y})}(t))t^{-\tau_{(\mathsf{Y})}}$$

From our calculation here we could not the exact slowly varying function in t when $\tau_M \leq \tau_e + 2$. We think this could also be shown by tweaking the sum in (D.25) properly.

We now complete the argument. Equations (D.31) and (D.21) complete the proof of Corollary 1.6(b) and (D.16) proves the claims in (1.17)-(1.18) which matches with [18, Proposition 1.4]. It remains to prove Lemma D.3:

Proof of Lemma D.3. For $i \ge m - \delta - \tau_e - 1$, we bound

(D.32)
$$\prod_{j=0}^{i-1} \frac{m+\delta+\tau_e+j+2}{t+\delta+\tau_e+j+2} \le 1,$$

so that (D.33)

$$\sum_{i\geq m-\delta-\tau_e-1}^{1} \frac{1}{(\tau_e+i)(m+\tau_e+i+\lfloor\delta\rfloor)} \leq \sum_{i\geq m-\delta-\tau_e-1}^{1} \frac{1}{(\tau_e+i)(\tau_e+i+\lfloor\delta\rfloor)} = \Theta(1/m),$$

as required.

For $i < m - \delta - \tau_e - 1$, instead, we rewrite the summands in (D.29) as

(D.34)
$$\frac{1}{(\tau_e+i)(m+\delta+\tau_e+i+1)} \prod_{j=0}^{i-1} \frac{m+\delta+\tau_e+j+2}{t+\delta+\tau_e+j+2}$$
$$= \frac{1}{(\tau_e+i)(m+\delta+\tau_e+1)} \prod_{j=0}^{i-1} \frac{m+\delta+\tau_e+j+1}{t+\delta+\tau_e+j+2} \,.$$

Now we bound the product terms as

(D.35)
$$\frac{m+\delta+\tau_e+j+1}{t+\delta+\tau_e+j+2} \le \frac{m+\delta+\tau_e+i-1+1}{t+\delta+\tau_e+i-1+2} \le \frac{2m}{t+m+1}.$$

Therefore,

(D.36)
$$\prod_{j=0}^{i-1} \frac{m+\delta+\tau_e+j+1}{t+\delta+\tau_e+j+2} \le \left(\frac{2m}{t+m+1}\right)^i.$$

We conclude that, also using that $\tau_e \geq 1$,

(D.37)
$$\sum_{i=0}^{m-\delta-\tau_e-1} \frac{1}{(\tau_e+i)(m+\delta+\tau_e+i+1)} \prod_{j=0}^{i-1} \frac{m+\delta+\tau_e+j+2}{t+\delta+\tau_e+j+2}$$
$$\leq \frac{1}{m+\delta+\tau_e+1} + \frac{1}{m} \sum_{i=1}^{m-\delta-\tau_e-1} \frac{1}{i} \left(\frac{2m}{t+m+1}\right)^i$$
$$\leq \Theta(1/m) + \frac{1}{m} \sum_{i\geq 1} \frac{1}{i} \left(\frac{2m}{t+m+1}\right)^i$$
$$= \Theta(m^{-1}(1-\log(1-2m/(t+m+1)))),$$

as required.

64