

On approximation for time-fractional stochastic diffusion equations on the unit sphere

Tareq Alodat, Quoc T. Le Gia* and Ian H. Sloan

La Trobe University, Melbourne, VIC 3086, Australia

The University of New South Wales, Sydney, NSW 2052, Australia

[math.PR] 4 Mar 2024

arXiv:2212.05690v3

ARTICLE INFO

Keywords:

Fractional derivative
Random field
Angular power spectrum
Spherical harmonics
Time-delayed Brownian motion

ABSTRACT

This paper develops a two-stage stochastic model to investigate the evolution of random fields on the unit sphere \mathbb{S}^2 in \mathbb{R}^3 . The model is defined by a time-fractional stochastic diffusion equation on \mathbb{S}^2 governed by a diffusion operator with a time-fractional derivative defined in the Riemann-Liouville sense. In the first stage, the model is characterized by a homogeneous problem with an isotropic Gaussian random field on \mathbb{S}^2 as an initial condition. In the second stage, the model becomes an inhomogeneous problem driven by a time-delayed Brownian motion on \mathbb{S}^2 . The solution to the model is given in the form of an expansion in terms of complex spherical harmonics. An approximation to the solution is given by truncating the expansion of the solution at degree $L \geq 1$. The rate of convergence of the truncation errors as a function of L and the mean square errors as a function of time are also derived. It is shown that the convergence rates depend not only on the decay of the angular power spectrum of the driving noise and the initial condition, but also on the order of the fractional derivative. We study sample properties of the stochastic solution and show that the solution is an isotropic Hölder continuous random field. Numerical examples and simulations inspired by the cosmic microwave background (CMB) are given to illustrate the theoretical findings.

1. Introduction

Let $\mathbb{S}^2 \subset \mathbb{R}^3$ be the 2-dimensional unit sphere in the Euclidean space \mathbb{R}^3 . In this paper, we develop a two-stage stochastic model defined by time-fractional stochastic diffusion equations on \mathbb{S}^2 . The stochastic model can be used to model the evolution of spatio-temporal stochastic systems such as climate changes and the density fluctuations in the primeval universe (see [2, 4, 5, 28]). For general stochastic diffusion models (or Itô diffusion) we refer the reader to [27]).

The model is constructed as follows. Let ξ be a strongly isotropic Gaussian random field on \mathbb{S}^2 and let $U(t)$, $t \in (0, \infty)$, be a random field on \mathbb{S}^2 , which solves the following time-fractional diffusion equation:

$$dU(t) - D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U(t) dt = \begin{cases} 0, & t \in (0, \tau], \\ dW_\tau(t), & t \in [\tau, \infty), \end{cases} \quad (1)$$

with the initial condition $U(0) = \xi$, where $\Delta_{\mathbb{S}^2}$ is the Laplace-Beltrami operator, α is a constant satisfying $0 < \alpha \leq 1$, D_t^β is the Riemann-Liouville time fractional derivative of order $0 \leq \beta < 1$ defined, see [31], as

$$D_t^\beta g(t) := \frac{d}{dt} \int_0^t \frac{1}{\Gamma(1-\beta)} \frac{1}{(t-s)^\beta} g(s) ds, \quad t > 0, \quad (2)$$

and W_τ , $\tau > 0$, is a time-delayed Brownian motion defined in $L_2(\Omega \times \mathbb{S}^2)$ (the product space of the probability space Ω and the sphere \mathbb{S}^2). The structure of the noise W_τ will be given in Section 2.3.

In words, the model in (1) describes an initial isotropic random field evolving by a time-fractional diffusion process, with no external input until time τ , at which time external noise is switched on.

*Quoc T. Le Gia

✉ T.Alodat@latrobe.edu.au (T. Alodat); qlegia@unsw.edu.au (Q.T.L. Gia); I.sloan@unsw.edu.au (I.H. Sloan)

🌐 <https://research.unsw.edu.au/people/dr-quoc-thong-le-gia> (Q.T.L. Gia)

ORCID(s):

Note that in the model (1), it is assumed that the random field ξ is independent of the noise W_τ . Since equation (1) is linear, it is easy to see that U can be decomposed as

$$U(t) := U^H(t) + U^I(t), \quad t \in (0, \infty), \quad (3)$$

where U^H satisfies the homogeneous equation

$$dU^H(t) - D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U^H(t) dt = 0, \quad t \in (0, \infty), \quad (4)$$

with the initial condition $U^H(0) = \xi$, and U^I satisfies the inhomogeneous equation

$$dU^I(t) - D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U^I(t) dt = \begin{cases} 0, & t \in (0, \tau], \\ dW_\tau(t), & t \in [\tau, \infty), \end{cases} \quad (5)$$

with the condition $U^I(t) = 0, t \in (0, \tau]$, from which it follows that $U^I(t)$ is zero until time τ .

We write the equation (5) in the integral form, for $t > \tau$, as

$$U^I(t) - \frac{1}{\Gamma(\alpha)} \int_\tau^t \frac{\Delta_{\mathbb{S}^2} U^I(s)}{(t-s)^{1-\alpha}} ds = W_\tau(t). \quad (6)$$

Note that in (1) there are two sources of randomness, namely the initial condition ξ and the driving noise W_τ . The split of the stochastic solution U in equation (3) into two independent parts (due to linearity) separates the two sources of randomness: the homogeneous solution U^H is random only through the initial condition, while the inhomogeneous solution U^I is random only through the time-delayed Brownian motion W_τ .

We shall derive the stochastic solution U to the equation (1), which represents a random field defined in $L_2(\Omega \times \mathbb{S}^2)$. The solution U is given in the form of an expansion in terms of complex orthonormal spherical harmonics $\{Y_{\ell,m} : \ell \in \mathbb{N}_0, m = -\ell, \dots, \ell\}$, i.e.,

$$U(\omega, t) := \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \hat{U}_{\ell,m}(\omega, t) Y_{\ell,m}, \quad t \in (0, \infty), \quad (7)$$

where the random variables $\hat{U}_{\ell,m}$ represent the Fourier coefficients of U and are given by

$$\hat{U}_{\ell,m}(\omega, t) := \int_{\mathbb{S}^2} U(\omega, \mathbf{x}, t) \overline{Y_{\ell,m}(\mathbf{x})} \mu(d\mathbf{x}), \quad t \in (0, \infty),$$

where $\overline{Y_{\ell,m}}$ denotes the complex conjugate of $Y_{\ell,m}$ and μ is the normalized Riemann surface measure on the sphere \mathbb{S}^2 (details are given in Section 2).

Given that the initial field ξ is a strongly isotropic Gaussian random field on \mathbb{S}^2 , we shall show that the stochastic solution U is also a strongly isotropic Gaussian random field on \mathbb{S}^2 .

We truncate the expansion (7) at a level $\ell = L \geq 1$ to obtain an approximate version U_L of the solution U , i.e.,

$$U_L(\omega, t) := \sum_{\ell=0}^L \sum_{m=-\ell}^{\ell} \hat{U}_{\ell,m}(\omega, t) Y_{\ell,m}, \quad t \in (0, \infty).$$

We investigate the approximate solution U_L and its rate of convergence as $L \rightarrow \infty$. We shall show that the convergence rate (in L_2 -norm on $L_2(\Omega \times \mathbb{S}^2)$) depends not only on the decay of the power spectrum of the driving noise W_τ and the initial condition ξ , but also on the order of the fractional derivative α . Also, we show that the stochastic solution $U(\omega, t), t > \tau$, evolves continuously with time by deriving an upper bound in the L_2 -norm for its temporal increments from time t to $t+h$ of the form $q(t)h^{\frac{1}{2}}$, $h > 0, t \in (\tau, \infty)$, where the continuous function $q(\cdot)$ does not depend on h . Finally, under some conditions, we show the existence of a locally Hölder continuous modification of U .

Note that the proposed model, defined in (1), describes the evolution (in time) of a two-stage stochastic system. The homogeneous equation given in (4) determines the evolution for the system with no external input while the

inhomogeneous equation (5) provides the perturbation produced by the Brownian motion starting at time τ . The cosmic microwave background (CMB) provides motivation for considering multi-stage stochastic systems. Roughly speaking, CMB is electromagnetic radiation that has been emitted across the cosmos since ionised atoms and electrons recombined roughly 370,000 years after the Big Bang [9, 29, 30]. Since CMB has passed through several stages called formation epochs or phases (such as Planck, Grand Unification, Inflation and recombination), it can be considered as an example of a multi-stage stochastic system.

The literature shows a variety of studies where models using stochastic partial differential equations were developed, see for example [4, 6, 18, 21, 22, 25]. However, little attention has been paid to two-stage stochastic models of the kind considered here. A fractional SPDE, governed by a fractional derivative in time and a fractional diffusion operator in space, was developed in [1] to model evolution of tangent vector random fields on the unit sphere. In a recent paper, Anh et.al. [2] considered a two-stage stochastic model defined by SPDEs on the sphere governed by a fractional diffusion operator in space and a fractional Brownian motion as driving noise. The case where a random initial condition given by a fractional stochastic Cauchy problem was considered. They showed that the truncation errors of the stochastic solution have, in the L_2 -norm, the convergence rates $C_t L^{-r}$, $r > 1$, under the assumption that the variances of the driving noise satisfy some smoothness condition. Also, as a function of time t , it was demonstrated that the ‘‘constant’’ C_t blows up when t is sufficiently small. In a recent article [16], a time-fractional stochastic heat equation driven by time-space white noise in \mathbb{R}^d was considered. The model is defined in the distribution sense, and an explicit solution is derived within the space of tempered distributions. The stochastic solution is demonstrated to exhibit mild behavior exclusively when the dimension is either $d = 1$ or $d = 2$, and the order of the fractional derivative falls within the range $\alpha \in (1, 2)$, while for $\alpha < 1$, the solution is not mild for any d .

The paper is structured as follows. Section 2 presents necessary material from the theory of functions on the sphere \mathbb{S}^2 , spherical harmonics, Gaussian random fields, $L_2(\mathbb{S}^2)$ -valued time-delayed Brownian motions, and some necessary tools from the theory of stochastic integrals. Section 3 derives properties of the solution to the homogeneous equation (4). In Section 4 we study the solution to the inhomogeneous equation (5). In Section 5 we give results for the combined equation (1). Section 6 derives an approximation to the solution of equation (1), the rate of convergence for the truncation errors. In Section 7 we study the temporal increments, in the L_2 -norm, of the stochastic solution of (1) and its sample Hölder continuity property. In Section 8 we provide some numerical examples to explain the theoretical findings. In particular, Section 8.1 illustrates the evolution of the stochastic solution of (1) using simulated data inspired by the CMB map. Section 8.2 explains the convergence rates of the truncation errors, in the L_2 -norm, of the stochastic solution of (1). Finally, Section 8.3 explores the convergence rates of the temporal increments, in the L_2 -norm, of the solution of (1).

2. Preliminaries

Let \mathbb{R}^3 be the 3-dimensional Euclidean space. Let $\mathbf{x}, \mathbf{y} \in \mathbb{R}^3$, i.e. $\mathbf{x} = (x_1, x_2, x_3)$ and $\mathbf{y} = (y_1, y_2, y_3)$, then the inner product of \mathbf{x} and \mathbf{y} is $\mathbf{x} \cdot \mathbf{y} := \sum_{i=1}^3 x_i y_i$ and the Euclidean norm of \mathbf{x} is $\|\mathbf{x}\| := \sqrt{\mathbf{x} \cdot \mathbf{x}}$. We denote by $\mathbb{S}^2 \subset \mathbb{R}^3$ the 2-dimensional unit sphere in \mathbb{R}^3 , i.e., $\mathbb{S}^2 := \{\mathbf{x} \in \mathbb{R}^3 : \|\mathbf{x}\| = 1\}$. The pair $(\mathbb{S}^2, \tilde{d})$ forms a compact metric space, where \tilde{d} is the geodesic metric defined by $\tilde{d}(\mathbf{x}, \mathbf{y}) := \arccos(\mathbf{x} \cdot \mathbf{y})$, $\mathbf{x}, \mathbf{y} \in \mathbb{S}^2$.

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $L_2(\Omega, \mathbb{P})$ be the L_2 -space on Ω with respect to the probability measure \mathbb{P} , endowed with the norm $\|\cdot\|_{L_2(\Omega)}$. For two random variables X and Y on $(\Omega, \mathcal{F}, \mathbb{P})$, we denote by $\mathbb{E}[X]$ the expected value of X and by $Cov(X, Y) := \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])]$ the covariance between X and Y . For two random variables X and Y , we use $X \simeq Y$ to denote the equality in distribution. Note that when $X \simeq Y$, $Cov(X, X)$ is the variance of X , which is denoted by $Var[X]$. For a random variable X on $(\Omega, \mathcal{F}, \mathbb{P})$, we denote by $\varphi_X(\cdot)$ the characteristic function of X , i.e. $\varphi_X(r) := \mathbb{E}[e^{irX}]$.

2.1. Functions on the sphere

Let μ be the normalized Riemann surface measure on the sphere \mathbb{S}^2 (i.e. $\int_{\mathbb{S}^2} d\mu = 1$). We denote by $L_2(\mathbb{S}^2) := L_2(\mathbb{S}^2, \mu)$ the space of complex-valued square μ -integrable functions on \mathbb{S}^2 , i.e.,

$$L_2(\mathbb{S}^2) = \left\{ f : \int_{\mathbb{S}^2} |f(\mathbf{x})|^2 d\mu(\mathbf{x}) < \infty \right\}.$$

Let f, g be two arbitrary functions in $L_2(\mathbb{S}^2)$, the inner product $\langle f, g \rangle_{L_2(\mathbb{S}^2)}$ is defined as

$$\langle f, g \rangle_{L_2(\mathbb{S}^2)} := \int_{\mathbb{S}^2} f(\mathbf{x}) \overline{g(\mathbf{x})} d\mu(\mathbf{x})$$

and the norm $\|f\|_{L_2(\mathbb{S}^2)}$ of f is

$$\|f\|_{L_2(\mathbb{S}^2)} := \left(\int_{\mathbb{S}^2} |f(\mathbf{x})|^2 d\mu(\mathbf{x}) \right)^{\frac{1}{2}}.$$

Let $\mathbb{N}_0 = \{0, 1, 2, \dots\}$ and let $P_\ell, \ell \in \mathbb{N}_0$, be the Legendre polynomials of degree ℓ , i.e.,

$$P_\ell(t) := 2^{-\ell} \frac{1}{\ell!} \frac{d^\ell}{dt^\ell} (t^2 - 1)^\ell, \quad \ell \in \mathbb{N}_0, \quad t \in [-1, 1].$$

The Legendre polynomials define the associated Legendre functions $P_{\ell,m}, \ell \in \mathbb{N}_0, m = 0, \dots, \ell$, by

$$P_{\ell,m}(t) := (-1)^m (1 - t^2)^{\frac{m}{2}} \frac{d^m}{dt^m} P_\ell(t), \quad \ell \in \mathbb{N}_0, \quad t \in [-1, 1].$$

Let $\{Y_{\ell,m} : \ell \in \mathbb{N}_0, m = -\ell, \dots, \ell\}$ be an orthonormal basis of complex-valued spherical harmonics of $L_2(\mathbb{S}^2)$. Using spherical coordinates $\mathbf{x} := (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta) \in \mathbb{S}^2, \theta \in [0, \pi], \varphi \in [0, 2\pi)$, the spherical harmonic functions can be written as, for $\ell \in \mathbb{N}_0$,

$$Y_{\ell,m}(\theta, \varphi) := \sqrt{\frac{(2\ell + 1)(\ell - m)!}{(\ell + m)!}} P_{\ell,m}(\cos \theta) e^{im\varphi}, \quad m = 0, \dots, \ell,$$

$$Y_{\ell,m} := (-1)^m \overline{Y_{\ell,-m}}, \quad m = -\ell, \dots, -1.$$

In what follows, we will adopt the notation $Y_{\ell,m}$ to represent spherical harmonics as a function of both spherical and Euclidean coordinates. Specifically, for all $\mathbf{x} \in \mathbb{S}^2$, we shall write (with a slight abuse of notation) $Y_{\ell,m}(\mathbf{x}) := Y_{\ell,m}(\theta, \varphi)$, where $\mathbf{x} = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta)$ and $(\theta, \varphi) \in [0, \pi] \times [0, 2\pi)$.

The basis $Y_{\ell,m}$ and the Legendre polynomial P_ℓ satisfy the addition theorem (see [26]), that is for $\mathbf{x}, \mathbf{y} \in \mathbb{S}^2$, there holds

$$\sum_{m=-\ell}^{\ell} Y_{\ell,m}(\mathbf{x}) \overline{Y_{\ell,m}(\mathbf{y})} = (2\ell + 1) P_\ell(\mathbf{x} \cdot \mathbf{y}). \quad (8)$$

The Laplace-Beltrami operator $\Delta_{\mathbb{S}^2}$ (or the spherical Laplacian) on the sphere \mathbb{S}^2 at \mathbf{x} is given in terms of the spherical coordinates by (see [8, 26])

$$\Delta_{\mathbb{S}^2} := \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \left(\frac{1}{\sin \theta} \right)^2 \frac{\partial^2}{\partial \varphi^2}.$$

It is well-known that the spherical harmonic functions $\{Y_{\ell,m} : \ell \in \mathbb{N}_0, m = -\ell, \dots, \ell\}$ are the eigenfunctions of the negative Laplace-Beltrami operator $-\Delta_{\mathbb{S}^2}$ on the sphere \mathbb{S}^2 with eigenvalues

$$\lambda_\ell := \ell(\ell + 1), \quad \ell \in \mathbb{N}_0, \quad (9)$$

that is for $\ell \in \mathbb{N}_0, m = -\ell, \dots, \ell$,

$$-\Delta_{\mathbb{S}^2} Y_{\ell,m} = \lambda_\ell Y_{\ell,m}. \quad (10)$$

An arbitrary complex-valued function $f \in L_2(\mathbb{S}^2)$ can be expanded in terms of a Fourier-Laplace series, in the $L_2(\mathbb{S}^2)$ sense,

$$f = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \widehat{f}_{\ell,m} Y_{\ell,m}, \quad \text{with } \widehat{f}_{\ell,m} := \int_{\mathbb{S}^2} f(\mathbf{x}) \overline{Y_{\ell,m}(\mathbf{x})} d\mu(\mathbf{x}). \quad (11)$$

The $\widehat{f}_{\ell,m}$, for $\ell \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$, are known as the Fourier coefficients for the function f under the Fourier basis $Y_{\ell,m}$.

By Parseval's theorem, for $f \in L_2(\mathbb{S}^2)$, there holds

$$\|f\|_{L_2(\mathbb{S}^2)}^2 = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} |\widehat{f}_{\ell,m}|^2.$$

Note that by the properties of the spherical harmonic functions, the Fourier-Laplace series (11) of any real-valued $f \in L_2(\mathbb{S}^2)$ takes the form

$$f = \sum_{\ell=0}^{\infty} \left(\widehat{f}_{\ell,0} Y_{\ell,0} + 2 \sum_{m=1}^{\ell} \left(\operatorname{Re} \widehat{f}_{\ell,m} \operatorname{Re} Y_{\ell,m} - \operatorname{Im} \widehat{f}_{\ell,m} \operatorname{Im} Y_{\ell,m} \right) \right).$$

2.2. Isotropic random fields on the sphere

This subsection introduces isotropic Gaussian random fields on the sphere \mathbb{S}^2 and their expansions in terms of complex spherical harmonics [12, 21].

We denote by $\mathfrak{B}(\mathbb{S}^2)$ the Borel σ -algebra on the sphere \mathbb{S}^2 and $SO(3)$ the rotation group on \mathbb{R}^3 . An $\mathcal{F} \otimes \mathfrak{B}(\mathbb{S}^2)$ -measurable function $\xi : \Omega \times \mathbb{S}^2 \rightarrow \mathbb{R}$ is called a (jointly measurable) real-valued random field on \mathbb{S}^2 . In the paper, we assume that the random field $\xi \in L_2(\Omega \times \mathbb{S}^2)$. Let $L_2(\Omega \times \mathbb{S}^2, \mathbb{P} \otimes \mu) =: L_2(\Omega \times \mathbb{S}^2)$ be the real-valued L_2 -space of random fields on $\Omega \times \mathbb{S}^2$, with the product measure $\mathbb{P} \otimes \mu$. Using Fubini's theorem, the inner product of $\xi_1, \xi_2 \in L_2(\Omega \times \mathbb{S}^2)$ can be written as

$$\begin{aligned} \langle \xi_1, \xi_2 \rangle_{L_2(\Omega \times \mathbb{S}^2)} &:= \int_{\Omega \times \mathbb{S}^2} \xi_1(\omega, \mathbf{x}) \xi_2(\omega, \mathbf{x}) d(\mathbb{P} \otimes \mu)(\omega, \mathbf{x}) \\ &= \mathbb{E} \left[\langle \xi_1, \xi_2 \rangle_{L_2(\mathbb{S}^2)} \right], \end{aligned}$$

and the norm of $\xi_1 \in L_2(\Omega \times \mathbb{S}^2)$ is

$$\|\xi_1\|_{L_2(\Omega \times \mathbb{S}^2)} := \sqrt{\langle \xi_1, \xi_1 \rangle_{L_2(\Omega \times \mathbb{S}^2)}}.$$

In particular, $\xi(\omega, \cdot) \in L_2(\mathbb{S}^2)$ \mathbb{P} -a.s., since $\|\xi\|_{L_2(\Omega \times \mathbb{S}^2)}^2 = \mathbb{E}[\|\xi\|_{L_2(\mathbb{S}^2)}^2]$. Note that an arbitrary random field $\xi(\omega, \cdot) \in L_2(\mathbb{S}^2)$ \mathbb{P} -a.s., admits an expansion in terms of spherical harmonics, \mathbb{P} -a.s.,

$$\xi = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \widehat{\xi}_{\ell,m} Y_{\ell,m}, \quad \widehat{\xi}_{\ell,m} = \int_{\mathbb{S}^2} \xi(\mathbf{x}) \overline{Y_{\ell,m}(\mathbf{x})} d\mu(\mathbf{x}), \quad (12)$$

where the convergence is in the $L_2(\Omega \times \mathbb{S}^2)$ sense.

In this paper, we consider a particular class of random fields called isotropic random fields. Following [24], the random field ξ is strongly isotropic if for any $k \in \mathbb{N}$ and for all sets of k points $\mathbf{x}_1, \dots, \mathbf{x}_k \in \mathbb{S}^2$, and for any rotation $R \in SO(3)$, the joint distributions of $\xi(\mathbf{x}_1), \dots, \xi(\mathbf{x}_k)$ and $\xi(R\mathbf{x}_1), \dots, \xi(R\mathbf{x}_k)$ coincide.

The random field ξ is called a 2-weakly, isotropic random field if for all $\mathbf{x} \in \mathbb{S}^2$, the second moment of $\xi(\mathbf{x})$ is finite, that is $\mathbb{E}[|\xi(\mathbf{x})|^2] < \infty$, and for all $\mathbf{x}_1, \mathbf{x}_2 \in \mathbb{S}^2$, and for any rotation $R \in SO(3)$, (see [24]), there holds

$$\mathbb{E}[\xi(\mathbf{x}_1)] = \mathbb{E}[\xi(R\mathbf{x}_1)], \quad \mathbb{E}[\xi(\mathbf{x}_1)\xi(\mathbf{x}_2)] = \mathbb{E}[\xi(R\mathbf{x}_1)\xi(R\mathbf{x}_2)].$$

If ξ is a centered, 2-weakly isotropic random field, then the Fourier coefficients $\widehat{\xi}_{\ell,m}$ of ξ in (12) are uncorrelated mean-zero complex-valued random variables (see Remark 5.15 in [24]), i.e., for $\ell \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$,

$$\mathbb{E} \left[\widehat{\xi}_{\ell,m} \right] = 0, \quad \mathbb{E} \left[\widehat{\xi}_{\ell,m} \overline{\widehat{\xi}_{\ell',m'}} \right] = C_{\ell,m} \delta_{\ell\ell'} \delta_{mm'},$$

where the $C_{\ell,m}$ are non-negative numbers and $\delta_{\ell\ell'}$ is the Kronecker delta function. The sequence $\{C_{\ell,m} : \ell \in \mathbb{N}_0, m = -\ell, \dots, \ell\}$ is called the angular power spectrum of the random field ξ . It follows that for each $\mathbf{x} \in \mathbb{S}^2$, $\xi(\mathbf{x})$ is a centered random variable and its covariance function takes the form

$$\mathbb{E}[\xi(\mathbf{x})\xi(\mathbf{y})] = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} C_{\ell,m} Y_{\ell,m}(\mathbf{x}) \overline{Y_{\ell,m}(\mathbf{y})}, \quad \mathbf{x}, \mathbf{y} \in \mathbb{S}^2.$$

For a centered, 2-weakly isotropic random field ξ it holds, for $\mathbf{x}, \mathbf{y} \in \mathbb{S}^2$, that $\mathbb{E}[\xi(\mathbf{x})] = 0$ and $\mathbb{E}[\xi(\mathbf{x})\xi(\mathbf{y})]$ is rotationally invariant. In this case the angular power spectrum is independent of m , that is $C_{\ell,m} = C_{\ell}$, $\ell \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$, and can be written as

$$C_{\ell} = \mathbb{E}[|\widehat{\xi}_{\ell,m}|^2]. \quad (13)$$

Thus, the covariance function $\mathbb{E}[\xi(\mathbf{x})\xi(\mathbf{y})]$ takes the form, using the addition theorem for spherical harmonics (see equation (8)),

$$\mathbb{E}[\xi(\mathbf{x})\xi(\mathbf{y})] = \sum_{\ell=0}^{\infty} (2\ell + 1) C_{\ell} P_{\ell}(\mathbf{x} \cdot \mathbf{y}). \quad (14)$$

Since $|P_{\ell}(\cdot)| \leq 1$, for $\ell \rightarrow \infty$, the series (14) is convergent provided that

$$\sum_{\ell=0}^{\infty} (2\ell + 1) C_{\ell} < \infty. \quad (15)$$

Throughout all subsequent sections, we will assume that the condition given in equation (15) remains valid.

A random field ξ on \mathbb{S}^2 is Gaussian if for each $k \in \mathbb{N} = \{1, 2, 3, \dots\}$ and each k points $\mathbf{x}_1, \dots, \mathbf{x}_k \in \mathbb{S}^2$, the vector $(\xi(\mathbf{x}_1), \dots, \xi(\mathbf{x}_k))$ has a multivariate Gaussian distribution.

Note that a Gaussian random field ξ is strongly isotropic if and only if it is 2-weakly isotropic (see [24]). In this paper, we consider random fields that are Gaussian and strongly isotropic.

We recall the following result (see [21], Corollary 2.5).

Proposition 2.1. [21] *Let $\{C_{\ell} : \ell \in \mathbb{N}_0\}$ be the angular power spectrum of a centered, 2-weakly isotropic Gaussian random field ξ on \mathbb{S}^2 . Then ξ admits the expansion*

$$\xi = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \widehat{\xi}_{\ell,m} Y_{\ell,m},$$

which is convergent in the $L_2(\Omega \times \mathbb{S}^2)$ sense, where $Y_{\ell,m}$, $\ell \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$, are spherical harmonic functions and the elements of the set $\mathcal{V} := \{\widehat{\xi}_{\ell,m} : \ell \in \mathbb{N}_0, m = -\ell, \dots, \ell\}$ are complex-valued, centered Gaussian random variables satisfying:

- The subset $\mathcal{V}' := \{\widehat{\xi}_{\ell,m} : \ell \in \mathbb{N}_0, m = 0, \dots, \ell\} \subset \mathcal{V}$ consists of **independent**, complex-valued Gaussian random variables.
- The elements of \mathcal{V}' with $m > 0$ have the property that $\text{Re} \widehat{\xi}_{\ell,m}$ and $\text{Im} \widehat{\xi}_{\ell,m}$ are independent and $\mathcal{N}(0, C_{\ell}/2)$ distributed.
- The elements of \mathcal{V}' with $m = 0$ are real-valued, and the elements $\text{Re} \widehat{\xi}_{\ell,0}$ are $\mathcal{N}(0, C_{\ell})$ distributed.
- The elements of \mathcal{V} with $m < 0$ can be obtained from those of \mathcal{V}' by applying the relations

$$\text{Re} \widehat{\xi}_{\ell,m} = (-1)^m \text{Re} \widehat{\xi}_{\ell,-m} \text{ and } \text{Im} \widehat{\xi}_{\ell,m} = (-1)^{m+1} \text{Im} \widehat{\xi}_{\ell,-m}.$$

Remark 2.1. Let $\{C_\ell : \ell \in \mathbb{N}_0\}$ be the angular power spectrum of a centered, strongly isotropic Gaussian random field ξ on \mathbb{S}^2 . Let for $\ell \in \mathbb{N}_0$,

$$\mathcal{Z}_\ell := \left\{ Z_{\ell,0}^{(1)}, Z_{\ell,1}^{(1)}, \dots, Z_{\ell,\ell}^{(1)}, Z_{\ell,1}^{(2)}, Z_{\ell,2}^{(2)}, \dots, Z_{\ell,\ell}^{(2)} \right\}, \quad (16)$$

be a set of independent, real-valued, standard normally distributed random variables. Then, by Proposition 2.1, the field ξ can be represented, in the $L_2(\Omega \times \mathbb{S}^2)$ sense, as

$$\begin{aligned} \xi : & \simeq \sum_{\ell=0}^{\infty} \left(\sqrt{C_\ell} Z_{\ell,0}^{(1)} Y_{\ell,0} \right. \\ & \left. + \sqrt{2C_\ell} \sum_{m=1}^{\ell} \left(Z_{\ell,m}^{(1)} \operatorname{Re} Y_{\ell,m} + Z_{\ell,m}^{(2)} \operatorname{Im} Y_{\ell,m} \right) \right), \end{aligned}$$

where $Z_{\ell,m_1}^{(1)}, Z_{\ell,m_2}^{(2)} \in \mathcal{Z}_\ell$, $\ell \in \mathbb{N}_0$, $m_1 = 0, \dots, \ell$, $m_2 = 1, \dots, \ell$.

2.3. Brownian motion

The standard Brownian motion (or the Wiener process) $B(t)$, $t \geq 0$, with variance 1 at $t = 1$, is a centered Gaussian process on $\mathbb{R}_+ = [0, \infty)$ satisfying $B(0) = 0$ and

$$\mathbb{E} \left[\left| B(t) - B(s) \right|^2 \right] = |t - s|, \quad s, t \geq 0.$$

Let $B_\tau(t)$, $t \geq \tau \geq 0$, be a real-valued, time-delayed Brownian motion with $B_\tau(\tau) = 0$ and variance 1 at $t = \tau + 1$. Note that it is easy to show that $B_\tau(t)$, $t \geq \tau \geq 0$, is a centered Gaussian process on $[\tau, \infty)$ satisfying

$$\mathbb{E} \left[\left| B_\tau(t) - B_\tau(s) \right|^2 \right] = |t - s|, \quad t, s \geq \tau. \quad (17)$$

Putting $s = \tau$ in (17) we have

$$\mathbb{E} \left[\left(B_\tau(t) \right)^2 \right] = |t - \tau|, \quad t \geq \tau \geq 0. \quad (18)$$

The case when $\tau = 0$, $B_0(t)$, $t \geq 0$, represents the standard Brownian motion. We will for brevity write $B_0(t)$ as $B(t)$ if no confusion arises.

We define a time-delayed Brownian motion on the unit sphere \mathbb{S}^2 as in [7, Sections 4.1.1 and 4.1.2]. Let $W_\tau(t)$, $\tau \geq 0$, be an $L_2(\mathbb{S}^2)$ -valued time-delayed Brownian motion. Then, for each real-valued f in $L_2(\mathbb{S}^2)$, the stochastic process $\langle W_\tau(t), f \rangle_{L_2(\mathbb{S}^2)}$ is a real-valued, time-delayed Brownian motion.

For arbitrary real-valued $f, g \in L_2(\mathbb{S}^2)$, $t, s \geq \tau$,

$$\mathbb{E} \left[\langle W_\tau(t), f \rangle_{L_2(\mathbb{S}^2)} \langle W_\tau(s), f \rangle_{L_2(\mathbb{S}^2)} \right] = (t \wedge s) \mathbb{E} \left[\langle W_\tau(\tau + 1), f \rangle_{L_2(\mathbb{S}^2)}^2 \right]$$

and

$$\mathbb{E} \left[\langle W_\tau(t), f \rangle_{L_2(\mathbb{S}^2)} \langle W_\tau(s), g \rangle_{L_2(\mathbb{S}^2)} \right] = (t \wedge s) \mathbb{E} \left[\langle Qf, g \rangle_{L_2(\mathbb{S}^2)} \right],$$

where Q is the covariance operator for the law of $W_\tau(1 + \tau)$, see [7, Section 2.3.1].

Definition 2.1. An $L_2(\mathbb{S}^2)$ -valued stochastic process $W_\tau(t)$, for $t \geq \tau$, is called a Q -Wiener process if

1. $W_\tau(\tau) = 0$,
2. W_τ has continuous trajectories,
3. W_τ has independent increments,
4. $W_\tau(t) - W_\tau(s)$ follows the normal distribution $\mathcal{N}(0, (t - s)Q)$, $t \geq s \geq \tau$.

For the complete orthonormal basis $Y_{\ell,m}$, there is a sequence of non-negative numbers \mathcal{A}_ℓ ,

$$QY_{\ell,m} = \mathcal{A}_\ell Y_{\ell,m}, \quad \ell = 0, 1, 2, \dots; m = -\ell, \dots, \ell.$$

Proposition 2.2. *If Q is of trace class, i.e., $\text{Tr}(Q) < \infty$, then the following condition holds true*

$$\sum_{\ell=0}^{\infty} (2\ell + 1) \mathcal{A}_\ell < \infty. \quad (19)$$

PROOF. By definition,

$$\text{Tr}(Q) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \langle QY_{\ell,m}, Y_{\ell,m} \rangle_{L_2(\mathbb{S}^2)} = \sum_{\ell=0}^{\infty} (2\ell + 1) \mathcal{A}_\ell < \infty. \quad (20)$$

In what follows we will assume that the condition (19) remains valid.

Let for $\ell \in \mathbb{N}_0$, $t \geq \tau \geq 0$,

$$\mathcal{B}_{\ell,\tau}(t) := \left\{ \beta_{\ell,0,\tau}^{(1)}(t), \beta_{\ell,1,\tau}^{(1)}(t), \dots, \beta_{\ell,\ell,\tau}^{(1)}(t), \beta_{\ell,1,\tau}^{(2)}(t), \beta_{\ell,2,\tau}^{(2)}(t), \dots, \beta_{\ell,\ell,\tau}^{(2)}(t) \right\}, \quad (21)$$

be a set of independent, real-valued time-delayed Brownian motions with variances 1 when $t = \tau + 1$.

Using the above setting, we have the following proposition. Its proof is similar to the one from [7, Proposition 4.3].

Proposition 2.3. *The noise $W_\tau(t)$, $t \geq \tau \geq 0$, admits the following representation*

$$W_\tau(t) := \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \widehat{W}_{\ell,m,\tau}(t) Y_{\ell,m}, \quad t \geq \tau \geq 0, \quad (22)$$

where $\widehat{W}_{\ell,m,\tau}(t)$, $t \geq \tau \geq 0$, $\ell \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$, are defined by

$$\widehat{W}_{\ell,m,\tau} := \begin{cases} \sqrt{\mathcal{A}_\ell} \beta_{\ell,0,\tau}^{(1)}, & m = 0, \\ \sqrt{\frac{\mathcal{A}_\ell}{2}} \left(\beta_{\ell,m,\tau}^{(1)} - i \beta_{\ell,m,\tau}^{(2)} \right), & m = 1, \dots, \ell, \\ (-1)^m \sqrt{\frac{\mathcal{A}_\ell}{2}} \left(\beta_{\ell,|m|,\tau}^{(1)} + i \beta_{\ell,|m|,\tau}^{(2)} \right), & m = -\ell, \dots, -1, \end{cases} \quad (23)$$

where for $\ell \in \mathbb{N}_0$, $m_1 = 0, \dots, \ell$, $m_2 = 1, 2, \dots, \ell$, $\beta_{\ell,m_1,\tau}^{(1)}, \beta_{\ell,m_2,\tau}^{(2)} \in \mathcal{B}_{\ell,\tau}$ and $\mathcal{B}_{\ell,\tau}$ is defined by (21).

It follows, using (23), that the expansion (22) can be written, in the $L_2(\Omega \times \mathbb{S}^2)$ sense, as

$$\begin{aligned} W_\tau(t) &\simeq \sum_{\ell=0}^{\infty} \left(\sqrt{\mathcal{A}_\ell} \beta_{\ell,0,\tau}^{(1)}(t) Y_{\ell,0} \right. \\ &\quad \left. + \sqrt{2\mathcal{A}_\ell} \sum_{m=1}^{\ell} \left(\beta_{\ell,m,\tau}^{(1)}(t) \text{Re } Y_{\ell,m} + \beta_{\ell,m,\tau}^{(2)}(t) \text{Im } Y_{\ell,m} \right) \right), \quad t \geq \tau \geq 0. \end{aligned} \quad (24)$$

Since by (19), $\sum_{\ell=0}^{\infty} (2\ell + 1) \mathcal{A}_\ell < \infty$, we obtain, using (24),

$$\begin{aligned} \|W_\tau(t)\|_{L_2(\Omega \times \mathbb{S}^2)}^2 &= \sum_{\ell=0}^{\infty} \left(\mathcal{A}_\ell \mathbb{E}[(\beta_{\ell,0,\tau}^{(1)}(t))^2] \|Y_{\ell,0}\|_{L_2(\mathbb{S}^2)}^2 \right. \\ &\quad \left. + 2\mathcal{A}_\ell \sum_{m=1}^{\ell} \left(\mathbb{E}[(\beta_{\ell,m,\tau}^{(1)}(t))^2] \|\text{Re } Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 + \mathbb{E}[(\beta_{\ell,m,\tau}^{(2)}(t))^2] \|\text{Im } Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 \right) \right) \\ &= (t - \tau) \sum_{\ell=0}^{\infty} (2\ell + 1) \mathcal{A}_\ell < \infty, \end{aligned}$$

where the second step uses (18). Thus the noise W_τ is well-defined in $L_2(\Omega \times \mathbb{S}^2)$.

Note that the sequence $\{\mathcal{A}_\ell : \ell \in \mathbb{N}_0\}$ is called the angular power spectrum of W_τ .

2.4. Stochastic integrals of Mittag-Leffler functions

In this subsection, we provide some tools and technical results from the theory of stochastic integrals.

It is well-known (see [19]) that for a non-random, measurable function g in $L_2([s, t])$, and a real-valued Brownian motion $B(t)$, stochastic integrals $\int_s^t g(u)dB(u)$, are defined as Wiener integrals. We can similarly define stochastic integrals, in the $L_2(\Omega \times \mathbb{S}^2)$ sense, of the form $\int_s^t g(u)dW_\tau(u)$, $t > s \geq \tau$.

The following definition states that an $L_2(\mathbb{S}^2)$ -valued stochastic integral $\int_s^t g(u)dW_\tau(u)$ can be defined as an expansion in spherical harmonics with random coefficients of the form $\int_s^t g(u)d\widehat{W}_{\ell,m,\tau}(u)$.

Definition 2.2. Let W_τ be an $L_2(\mathbb{S}^2)$ -valued time-delayed Brownian motion on \mathbb{S}^2 . Let $\widehat{W}_{\ell,m,\tau}(t)$, $t \geq \tau$, $\ell \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$, be the Fourier coefficients for W_τ defined by (23). For $t > s \geq \tau$, the stochastic integral $\int_s^t g(u)dW_\tau(u)$ for a measurable function g in $L_2([s, t])$ is defined, in the $L_2(\Omega \times \mathbb{S}^2)$ sense, by

$$\int_s^t g(u)dW_\tau(u) := \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \left(\int_s^t g(u)d\widehat{W}_{\ell,m,\tau}(u) \right) Y_{\ell,m}.$$

The following result gives expressions for stochastic integrals of the form $\int_s^t g(u)dW_\tau(u)$ in terms of the angular power spectrum of W_τ .

Proposition 2.4. Let W_τ be an $L_2(\mathbb{S}^2)$ -valued time-delayed Brownian motion. Let $\{\mathcal{A}_\ell : \ell \in \mathbb{N}_0\}$ be the angular power spectrum of W_τ . Then, for $t > s \geq \tau$, the stochastic integral $\int_s^t g(u)dW_\tau(u)$ given by Definition 2.2 satisfies

$$\mathbb{E} \left[\left\| \int_s^t g(u)dW_\tau(u) \right\|_{L_2(\mathbb{S}^2)}^2 \right] = \sum_{\ell=0}^{\infty} (2\ell + 1) \mathcal{A}_\ell \int_s^t |g(u)|^2 du.$$

PROOF. By Definition 2.2 and Parseval's identity, we get

$$\begin{aligned} \mathbb{E} \left[\left\| \int_s^t g(u)dW_\tau(u) \right\|_{L_2(\mathbb{S}^2)}^2 \right] &= \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \mathbb{E} \left[\left\| \int_s^t g(u)d\widehat{W}_{\ell,m,\tau}(u) \right\|^2 \right] \\ &= \left(\int_s^t |g(u)|^2 du \right) \sum_{\ell=0}^{\infty} \mathcal{A}_\ell \sum_{m=-\ell}^{\ell} 1 \\ &= \left(\int_s^t |g(u)|^2 du \right) \sum_{\ell=0}^{\infty} (2\ell + 1) \mathcal{A}_\ell, \end{aligned}$$

where the second step uses Itô's isometry (see [19]) and the last step uses the properties of $Y_{\ell,m}$, which completes the proof.

Let

$$E_{\alpha,\beta}(z) := \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)}, \quad \alpha, \beta > 0, z \in \mathbb{C}, \quad (25)$$

be the two-parametric Mittag-Leffler function [13]. For $\beta = 1$, it reduces to the classical Mittag-Leffler function, i.e., $E_{\alpha,1}(z) := E_\alpha(z)$. Here $\Gamma(\cdot)$ stands for the gamma function and \mathbb{C} is the set of complex numbers.

The following upper bound holds true [31, Theorem 1.6]

$$E_{\alpha,\beta}(-z) \leq \frac{C}{1+z}, \quad \alpha < 2, \beta \in \mathbb{R}, C > 0, z \geq 0. \quad (26)$$

For $\alpha \in (0, 1]$, let M_α be defined as

$$M_\alpha := \begin{cases} (\Gamma(1+\alpha))^2 |2\alpha - 1|^{-1}, & \alpha \neq \frac{1}{2}, \\ (\Gamma(3/2))^2, & \alpha = \frac{1}{2}. \end{cases} \quad (27)$$

For $\ell \in \mathbb{N}_0$, $t \geq 0$, and $\alpha \in (0, 1]$, let $\sigma_{\ell,t,\alpha}^2$ be defined as

$$\sigma_{\ell,t,\alpha}^2 := \int_0^t (E_\alpha(-\lambda_\ell(t-u)^\alpha))^2 du = \int_0^t (E_\alpha(-\lambda_\ell r^\alpha))^2 dr, \quad (28)$$

where $E_\alpha(\cdot)$ is defined in (25) and λ_ℓ is given by (9).

Remark 2.2. It is well-known that the Mittag-Leffler function $E_\alpha(-\lambda_\ell t^\alpha)$, $\alpha \in (0, 1]$, $t > 0$, is positive and decreasing, with $E_\alpha(0) = 1$. Thus, for $t \geq 0$, $\ell \in \mathbb{N}_0$, there holds $E_\alpha(-\lambda_\ell t^\alpha) \in (0, 1]$, $(E_\alpha(-\lambda_\ell t^\alpha))^2 \leq E_\alpha(-\lambda_\ell t^\alpha)$ and the integral defined in (28) is finite, i.e.,

$$\sigma_{\ell,t,\alpha}^2 = \int_0^t (E_\alpha(-\lambda_\ell r^\alpha))^2 dr \leq \int_0^t E_\alpha(-\lambda_\ell r^\alpha) dr \leq t < \infty.$$

Proposition 2.5. Let $\alpha \in (0, 1]$ and $\mathcal{I}_{\ell,m,\alpha}^{(j)}(t)$, $j = 1, 2$, $\ell \in \mathbb{N}_0$, $m = 0, \dots, \ell$, be stochastic integrals defined as

$$\mathcal{I}_{\ell,m,\alpha}^{(j)}(t) := \int_0^t E_\alpha(-\lambda_\ell(t-u)^\alpha) d\beta_{\ell,m}^{(j)}(u), \quad t > 0, \quad (29)$$

where for each $j = 1, 2$, the process $\beta_{\ell,m}^{(j)}(u)$ is an independent real-valued Brownian motion with variance 1 at $u = 1$. Then, for $m = 0, \dots, \ell$, $\ell \in \mathbb{N}_0$, each stochastic integral in (29) is normally distributed with mean zero and variance $\sigma_{\ell,t,\alpha}^2$, where $\sigma_{\ell,t,\alpha}^2$ is given by (28).

PROOF. Note that for $t > 0$ and $\lambda_\ell \geq 0$, $E_\alpha(-\lambda_\ell t^\alpha)$ is a bounded measurable function on \mathbb{R}_+ . Then, by Itô's isometry (see [19]), for $j = 1, 2$, there holds

$$\begin{aligned} \mathbb{E} \left[(\mathcal{I}_{\ell,m,\alpha}^{(j)}(t))^2 \right] &= \mathbb{E} \left[\left| \int_0^t E_\alpha(-\lambda_\ell(t-u)^\alpha) d\beta_{\ell,m}^{(j)}(u) \right|^2 \right] \\ &= \mathbb{E} \left[\int_0^t |E_\alpha(-\lambda_\ell(t-u)^\alpha)|^2 du \right] \\ &= \int_0^t (E_\alpha(-\lambda_\ell r^\alpha))^2 dr = \sigma_{\ell,t,\alpha}^2. \end{aligned}$$

By [19] (see Theorem 2.3.4), we conclude that each stochastic integral defined in (29) is a Gaussian random variable with mean zero and variance $\sigma_{\ell,t,\alpha}^2$ given by (28), thus completing the proof.

Proposition 2.6. For $t > 0$ and $\ell \in \mathbb{N}$ we have

$$\sigma_{\ell,t,\alpha}^2 \leq \begin{cases} \lambda_\ell^{-\frac{1}{\alpha}} + M_\alpha t^{1-2\alpha} \lambda_\ell^{-2}, & \alpha \in (0, \frac{1}{2}), \\ \lambda_\ell^{-\frac{1}{\alpha}} (1 + M_\alpha), & \alpha \in (\frac{1}{2}, 1], \\ \lambda_\ell^{-2} (1 + M_{\frac{1}{2}} \ln(\lambda_\ell^2 t)), & \alpha = \frac{1}{2}, \end{cases} \quad (30)$$

where M_α is given by (27).

PROOF. Using (28) with the substitution $s = \lambda_\ell^{\frac{1}{\alpha}} r$ we get

$$\sigma_{\ell,t,\alpha}^2 = \lambda_\ell^{-\frac{1}{\alpha}} \int_0^{\lambda_\ell^{\frac{1}{\alpha}} t} (E_\alpha(-s^\alpha))^2 ds. \quad (31)$$

Suppose that $t\lambda_\ell^{\frac{1}{\alpha}} > 1$ and write, using (31),

$$\begin{aligned}\sigma_{\ell,t,\alpha}^2 &= \lambda_\ell^{-\frac{1}{\alpha}} \left(\int_0^1 (E_\alpha(-s^\alpha))^2 ds + \int_1^{\lambda_\ell^{\frac{1}{\alpha}} t} (E_\alpha(-s^\alpha))^2 ds \right) \\ &\leq \lambda_\ell^{-\frac{1}{\alpha}} \left(1 + (\Gamma(1+\alpha))^2 \int_1^{\lambda_\ell^{\frac{1}{\alpha}} t} s^{-2\alpha} ds \right),\end{aligned}\quad (32)$$

where the second step uses Remark 2.2 and the upper bound (see [33], Theorem 4),

$$E_\alpha(-s^\alpha) \leq \frac{1}{1 + (\Gamma(1+\alpha))^{-1} s^\alpha} \leq \Gamma(1+\alpha) s^{-\alpha}, \quad s > 0. \quad (33)$$

Note that if $\alpha \in (0, \frac{1}{2})$, then

$$\begin{aligned}\int_1^{\lambda_\ell^{\frac{1}{\alpha}} t} s^{-2\alpha} ds &= \frac{1}{1-2\alpha} \left(\left(\lambda_\ell^{\frac{1}{\alpha}} t \right)^{1-2\alpha} - 1 \right) \\ &\leq \frac{1}{1-2\alpha} \left(\lambda_\ell^{\frac{1}{\alpha}} t \right)^{1-2\alpha} \\ &\leq \frac{1}{1-2\alpha} t^{1-2\alpha} \lambda_\ell^{\frac{1}{\alpha}-2}.\end{aligned}\quad (34)$$

Similarly if $\alpha \in (\frac{1}{2}, 1]$, then

$$\begin{aligned}\int_1^{\lambda_\ell^{\frac{1}{\alpha}} t} s^{-2\alpha} ds &= \frac{1}{2\alpha-1} \left(1 - \left(\lambda_\ell^{\frac{1}{\alpha}} t \right)^{1-2\alpha} \right) \\ &\leq \frac{1}{2\alpha-1}.\end{aligned}\quad (35)$$

For the critical case $\alpha = \frac{1}{2}$,

$$\int_1^{\lambda_\ell^{\frac{1}{\alpha}} t} s^{-2\alpha} ds = \ln(\lambda_\ell^{\frac{1}{\alpha}} t). \quad (36)$$

Thus the result follows by using the estimates (34), (35), and (36) with (32).

In subsequent sections, some results are required for establishing the pathwise solution of (5). In particular, a formal justification for taking the fractional derivative of stochastic integrals defined by (29) is required. Since there is currently no existing result in the literature on this matter, we embark on proving the following theorem.

Theorem 2.1. *Let $B(t)$ be a real-valued Brownian motion with variance 1 at $t = 1$. Let $Y(t, s) : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$ be a deterministic function such that both $Y(t, s)$ and ${}_s D_t^{1-\alpha} Y(t, s)$, $\alpha \in (0, 1)$, are continuous in t and s in some region of the ts -plane. Suppose that for $\alpha \in (0, 1)$ the following stochastic integrals are well-defined*

$$\int_0^t Y(s, t) dB(s) \text{ and } \int_0^t ({}_s D_t^{1-\alpha} Y(t, s)) dB(s).$$

Assume moreover that

$$\int_0^t (t-s)^{\alpha-1} \left(\int_0^s |Y(s, r)|^2 dr \right)^{\frac{1}{2}} ds < \infty, \quad (37)$$

then for all $t > 0$ there holds

$$D_t^{1-\alpha} \left(\int_0^t \Upsilon(t, r) d\mathbf{B}(r) \right) dt \simeq \left(\int_0^t {}_r D_t^{1-\alpha} \Upsilon(t, r) d\mathbf{B}(r) \right) dt + \left(\lim_{r \rightarrow t} {}_r D_t^{-\alpha} \Upsilon(t, r) \right) d\mathbf{B}(t), \quad (38)$$

where ${}_x D_t^{-p} \Upsilon(t, x) = \frac{1}{\Gamma(p)} \int_x^t (t-s)^{p-1} \Upsilon(s, x) ds$, $p > 0$, is a p -fold integral, see [31, Section 2.3.2].

PROOF. Using (2) and (37) with the stochastic Fubini theorem (see [7, Section 4.5]) we get

$$\begin{aligned} D_t^{1-\alpha} \left(\int_0^t \Upsilon(t, r) d\mathbf{B}(r) \right) dt &\simeq \frac{1}{\Gamma(\alpha)} d \left(\int_0^t \left(\frac{1}{(t-s)^{1-\alpha}} \int_0^s \Upsilon(s, r) d\mathbf{B}(r) \right) ds \right) \\ &\simeq d \left(\int_0^t \left(\frac{1}{\Gamma(\alpha)} \int_r^t \frac{\Upsilon(s, r)}{(t-s)^{1-\alpha}} ds \right) d\mathbf{B}(r) \right) \\ &\simeq d \left(\int_0^t \tilde{\Upsilon}(t, r) d\mathbf{B}(r) \right), \end{aligned}$$

where d denotes the t -differential and

$$\tilde{\Upsilon}(t, x) := \frac{1}{\Gamma(\alpha)} \int_x^t \frac{\Upsilon(s, x)}{(t-s)^{1-\alpha}} ds.$$

By Proposition 1.3 in [15], we could write

$$\begin{aligned} D_t^{1-\alpha} \left(\int_0^t \Upsilon(t, r) d\mathbf{B}(r) \right) dt &\simeq \left(\int_0^t \frac{\partial}{\partial t} \tilde{\Upsilon}(t, r) d\mathbf{B}(r) \right) dt + \lim_{r \rightarrow t} \tilde{\Upsilon}(t, r) d\mathbf{B}(t) \\ &\simeq \int_0^t {}_r D_t^{1-\alpha} \Upsilon(t, r) d\mathbf{B}(r) dt + \left(\lim_{r \rightarrow t} {}_r D_t^{-\alpha} \Upsilon(t, r) \right) d\mathbf{B}(t), \end{aligned}$$

which completes the proof.

It is worth noting that if we have $\Upsilon(t-r)$ instead of the function $\Upsilon(t, r)$, then (38) takes the form

$$D_t^{1-\alpha} \left(\int_0^t \Upsilon(t-r) d\mathbf{B}(r) \right) dt \simeq \left(\int_0^t D_r^{1-\alpha} \Upsilon(r) d\mathbf{B}(r) \right) dt + \left(\lim_{r \rightarrow 0} D_r^{-\alpha} \Upsilon(r) \right) d\mathbf{B}(t). \quad (39)$$

Lemma 2.1. *Let $t > \tau > 0$. For $\ell \geq 1$ and $\alpha \in (0, 1)$, there holds*

$$D_t^{1-\alpha} \int_0^{t-\tau} E_\alpha(-\lambda_\ell (t-\tau-r)^\alpha) d\mathbf{B}(r) \simeq \int_\tau^t D_r^{1-\alpha} E_\alpha(-\lambda_\ell r^\alpha) d\mathbf{B}_\tau(r), \quad (40)$$

where λ_ℓ is given by (9) and \mathbf{B}_τ is a real-valued, time-delayed Brownian motion starting at time $\tau > 0$.

PROOF. By Remark (2.2),

$$\begin{aligned} \int_\tau^t (t-s)^{\alpha-1} \left(\int_0^s |E_\alpha(-\lambda_\ell r^\alpha)|^2 dr \right)^{\frac{1}{2}} ds &\leq \sqrt{t} \int_\tau^t (t-s)^{\alpha-1} ds \\ &\leq \sqrt{t} (t-\tau)^\alpha / \alpha \\ &< \infty, \end{aligned}$$

and by Proposition 2.5, the stochastic integral

$$\int_0^{t-\tau} E_\alpha(-\lambda_\ell (t-\tau-r)^\alpha) d\mathbf{B}(r)$$

is well-defined.

Let us show that the stochastic integral

$$\int_{\tau}^t D_r^{1-\alpha} E_{\alpha}(-\lambda_{\ell} r^{\alpha}) dB_{\tau}(r)$$

is well-defined. Using Itô's isometry and Lemma 3.10 in [31] we obtain

$$\begin{aligned} \mathbb{E} \left[\left| \int_{\tau}^t D_r^{1-\alpha} E_{\alpha}(-\lambda_{\ell} r^{\alpha}) dB_{\tau}(r) \right|^2 \right] &= \int_{\tau}^t \left| D_r^{1-\alpha} E_{\alpha}(-\lambda_{\ell} r^{\alpha}) \right|^2 dr \\ &= \frac{1}{\lambda_{\ell}^2} \int_{\tau}^t \left(\frac{-\lambda_{\ell}}{\Gamma(\alpha)} \frac{d}{dr} \int_0^r \frac{E_{\alpha}(-\lambda_{\ell} s^{\alpha})}{(r-s)^{1-\alpha}} \right)^2 dr \\ &= \frac{1}{\lambda_{\ell}^2} \int_{\tau}^t \left(\frac{d}{dr} (E_{\alpha}(-\lambda_{\ell} r^{\alpha}) - 1) \right)^2 dr \\ &\leq C \lambda_{\ell}^{-2} \tau^{-1} < \infty, \end{aligned}$$

where the last second step used the relations (4.3.1) and (5.1.14) in [13] and the upper bound (26).

By the relation (39) we obtain

$$\begin{aligned} D_t^{1-\alpha} \left(\int_0^{t-\tau} E_{\alpha}(-\lambda_{\ell} (t-\tau-r)^{\alpha}) dB(r) \right) dt &\simeq \left(\int_{\tau}^t D_r^{1-\alpha} E_{\alpha}(-\lambda_{\ell} r^{\alpha}) dB_{\tau}(r) \right) dt \\ &\quad + \lim_{r \rightarrow 0} (D_r^{-\alpha} E_{\alpha}(-\lambda_{\ell} r^{\alpha})) dB_{\tau}(t). \end{aligned} \quad (41)$$

Using equations (2.88) and (1.100) in [31],

$$\begin{aligned} \lim_{r \rightarrow 0} D_r^{-\alpha} E_{\alpha}(-\lambda_{\ell} r^{\alpha}) &= \lim_{r \rightarrow 0} \frac{1}{\Gamma(\alpha)} \int_0^r (r-s)^{\alpha-1} E_{\alpha}(-\lambda_{\ell} s^{\alpha}) ds \\ &= \lim_{r \rightarrow 0} r^{\alpha} E_{\alpha,1+\alpha}(-\lambda_{\ell} r^{\alpha}) = 0. \end{aligned}$$

Thus, equation (41) becomes

$$D_t^{1-\alpha} \int_0^{t-\tau} E_{\alpha}(-\lambda_{\ell} (t-\tau-r)^{\alpha}) dB(r) \simeq \int_{\tau}^t D_r^{1-\alpha} E_{\alpha}(-\lambda_{\ell} r^{\alpha}) dB_{\tau}(r),$$

which completes the proof.

3. Solution of the homogeneous problem

Consider a random field $U^H(t) \in L_2(\Omega \times \mathbb{S}^2)$, $t \in (0, \infty)$. We say that the field U^H satisfies the equation (4), in the $L_2(\Omega \times \mathbb{S}^2)$ sense, if for a given $t > 0$, there holds

$$\sup_{t>0} \left\| \frac{d}{dt} U^H(t) - D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U^H(t) \right\|_{L_2(\Omega \times \mathbb{S}^2)}^2 = 0.$$

This section derives the solution $U^H(t)$, $t \in (0, \infty)$, to the homogeneous equation (4) under the random initial condition $U^H(0) = \xi$, where ξ is a centered, strongly isotropic Gaussian random field on \mathbb{S}^2 .

Note that since $U^H \in L_2(\Omega \times \mathbb{S}^2)$, it follows that the field U^H takes the series expansion

$$U^H(t) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \widehat{U}_{\ell,m}^H(t) Y_{\ell,m}, \quad t \in (0, \infty),$$

where the random coefficients $\widehat{U}^H_{\ell,m}$, $\ell \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$, are given by

$$\widehat{U}^H_{\ell,m}(t) = \int_{\mathbb{S}^2} U^H(\mathbf{x}, t) \overline{Y_{\ell,m}(\mathbf{x})} \mu(d\mathbf{x}), \quad t \in (0, \infty).$$

By multiplying both sides of (4) by $\overline{Y_{\ell,m}}$ and integrating over \mathbb{S}^2 , we obtain, with the help of (10), the following set of ordinary differential equations

$$d\widehat{U}^H_{\ell,m} + \lambda_\ell D_t^{1-\alpha} \widehat{U}^H_{\ell,m} dt = 0, \quad \widehat{U}^H_{\ell,m}(0) = \widehat{\xi}_{\ell,m}. \quad (42)$$

We denote by $\widetilde{U}^H_{\ell,m}(z)$ the Laplace transform of $\widehat{U}^H_{\ell,m}$ with respect to t , i.e.,

$$\widetilde{U}^H_{\ell,m}(z) = \mathcal{L}(\widehat{U}^H_{\ell,m}(t)) := \int_0^\infty e^{-tz} \widehat{U}^H_{\ell,m}(t) dt, \quad (43)$$

assuming that the integral converges. The condition for its convergence will be given in Proposition 3.1. Since $\mathcal{L}(D_t^{1-\alpha} \widehat{U}^H_{\ell,m}(t)) = z^{1-\alpha} \widetilde{U}^H_{\ell,m}(z)$ and $\mathcal{L}(\frac{d}{dt} \widehat{U}^H_{\ell,m}(t)) = z \widetilde{U}^H_{\ell,m}(z) - \widehat{U}^H_{\ell,m}(0)$, then by taking the Laplace transform of both sides of (42), we get

$$z \widetilde{U}^H_{\ell,m}(z) - \widehat{\xi}_{\ell,m} + \lambda_\ell z^{1-\alpha} \widetilde{U}^H_{\ell,m}(z) = 0.$$

Now solving for $\widetilde{U}^H_{\ell,m}(z)$, we arrive at

$$\widetilde{U}^H_{\ell,m}(z) = \frac{\widehat{\xi}_{\ell,m}}{z + \lambda_\ell z^{1-\alpha}}. \quad (44)$$

Recall that (see [31], equation 1.80)

$$\mathcal{L}\{E_\alpha(-\lambda_\ell t^\alpha)\} = \frac{1}{z + \lambda_\ell z^{1-\alpha}},$$

where $E_\alpha(\cdot)$ is the Mittag-Leffler function (see equation (25)).

By taking the inverse Laplace transform in (44), the solution of (4) becomes

$$U^H(t) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} E_\alpha(-\lambda_\ell t^\alpha) \widehat{\xi}_{\ell,m} Y_{\ell,m}, \quad t \in [0, \infty). \quad (45)$$

In order to prove that U^H is the solution of (4) we need to prove a truncation field U_L^H of U^H is a solution and then pass it to the limit as the truncation degree $L \rightarrow \infty$. Also, we need the uniform convergence of $\frac{d}{dt} U_L^H$ to $\frac{d}{dt} U^H$ in some sense. We begin with the following lemma.

Let $V_L^H(t)$ be defined as

$$V_L^H(t) := \sum_{\ell=1}^L \sum_{m=-\ell}^{\ell} (-\lambda_\ell) t^{\alpha-1} E_{\alpha,\alpha}(-\lambda_\ell t^\alpha) \widehat{\xi}_{\ell,m} Y_{\ell,m} = \frac{d}{dt} U_L^H(t), \quad (46)$$

where $\widehat{\xi}_{\ell,m}$ are the Fourier coefficients of ξ .

Lemma 3.1. *Let ξ , be a centered, 2-weakly isotropic Gaussian random field on \mathbb{S}^2 . Let $\{C_\ell : \ell \in \mathbb{N}_0\}$, the angular power spectrum of ξ , satisfy (15). Let $t_0 > 0$ be given. For $t \in [t_0, \infty)$, then V_L^H is convergent to*

$$V^H(t) := \sum_{\ell=1}^{\infty} \sum_{m=-\ell}^{\ell} (-\lambda_\ell) t^{\alpha-1} E_{\alpha,\alpha}(-\lambda_\ell t^\alpha) \widehat{\xi}_{\ell,m} Y_{\ell,m} \quad (47)$$

in the following sense

$$\sup_{t \geq t_0} \mathbb{E} \left[\left\| V_L^H(t) - V^H(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] \rightarrow 0, \quad \text{as } L \rightarrow \infty.$$

PROOF. For $t \geq t_0$, $L < M$, by Parseval's formula, we have

$$\begin{aligned}
 \mathbb{E} \left[\|V_L^H(t) - V_M^H(t)\|_{L_2(\mathbb{S}^2)}^2 \right] &= \mathbb{E} \left[\sum_{\ell=L+1}^M \sum_{m=-\ell}^{\ell} \lambda_{\ell}^2 t^{2\alpha-2} (E_{\alpha,\alpha}(-\lambda_{\ell} t^{\alpha}))^2 |\widehat{\xi}_{\ell,m}|^2 \right] \\
 &= \sum_{\ell=L+1}^M \sum_{m=-\ell}^{\ell} \lambda_{\ell}^2 t^{2\alpha-2} (E_{\alpha,\alpha}(-\lambda_{\ell} t^{\alpha}))^2 \mathbb{E} [|\widehat{\xi}_{\ell,m}|^2] \\
 &= \sum_{\ell=L+1}^M \lambda_{\ell}^2 t^{2\alpha-2} (E_{\alpha,\alpha}(-\lambda_{\ell} t^{\alpha}))^2 (2\ell + 1) C_{\ell} \text{ by using (13)} \\
 &\leq C^2 t^{-2} \sum_{\ell=L+1}^M (2\ell + 1) C_{\ell} \text{ by using (26)} \\
 &\leq C^2 t_0^{-2} \sum_{\ell=L+1}^M (2\ell + 1) C_{\ell}. \tag{48}
 \end{aligned}$$

For a given $\epsilon > 0$, using the condition (15), there is L_0 independent of t such that for $L, M \geq L_0$, the RHS of (48) is smaller than ϵ . So $\{V_L^H\}$ is a Cauchy sequence in $L_2(\Omega \times \mathbb{S}^2)$ and hence it is convergent. We define the limit of V_L^H as $L \rightarrow \infty$ as in (47). Note that by the condition (15) we obtain

$$\mathbb{E} \left[\|V^H(t)\|_{L_2(\mathbb{S}^2)}^2 \right] \leq C^2 t_0^{-2} \sum_{\ell=1}^{\infty} (2\ell + 1) C_{\ell} < \infty,$$

which guarantees that V^H is well-defined in the $L_2(\Omega \times \mathbb{S}^2)$ sense.

Now we prove the following lemma in which we adapt the techniques from the proofs of [32, Theorems 7.11 and 7.17].

Lemma 3.2. *Let $T > t_0 > 0$ be fixed. For $\hat{t} \in [t_0, T]$, we have*

$$\lim_{L \rightarrow \infty} \mathbb{E} \left[\left\| \frac{dU^H}{dt}(\hat{t}) - V_L^H(\hat{t}) \right\|_{L_2(\mathbb{S}^2)}^2 \right] = 0.$$

PROOF. Let us define for $t \neq \hat{t}$

$$\phi(t) := \frac{U^H(t) - U^H(\hat{t})}{t - \hat{t}}$$

and

$$\phi_L(t) := \frac{U_L^H(t) - U_L^H(\hat{t})}{t - \hat{t}}.$$

Then we have

$$\lim_{t \rightarrow \hat{t}} \phi(t) = \frac{dU^H}{dt}(\hat{t}) = V(\hat{t}). \tag{49}$$

We also have

$$\begin{aligned}
 \lim_{t \rightarrow \hat{t}} \phi_L(t) &= \sum_{\ell=0}^L \lim_{t \rightarrow \hat{t}} \frac{E_{\alpha}(-\lambda_{\ell} t^{\alpha}) - E_{\alpha}(-\lambda_{\ell} \hat{t}^{\alpha})}{t - \hat{t}} \sum_{m=-\ell}^{\ell} \widehat{\xi}_{\ell,m} Y_{\ell,m} \\
 &= \sum_{\ell=0}^L \frac{d}{d\hat{t}} E_{\alpha}(-\lambda_{\ell} \hat{t}^{\alpha}) \sum_{m=-\ell}^{\ell} \widehat{\xi}_{\ell,m} Y_{\ell,m}
 \end{aligned}$$

$$= \sum_{\ell=1}^L (-\lambda_\ell) \hat{t}^{\alpha-1} E_{\alpha,\alpha}(-\lambda_\ell \hat{t}^\alpha) \sum_{m=-\ell}^{\ell} \hat{\xi}_{\ell,m} Y_{\ell,m} = V_L^H(\hat{t}). \quad (50)$$

We remark, for a fixed $\omega \in \Omega$, that $V_L^H(\hat{t}) < \infty$ for $\hat{t} \geq t_0 > 0$.

Since by Lemma 3.1, $V_L^H(t)$ and $V_L^H(\hat{t})$ converge uniformly to $V^H(t)$ and $V^H(\hat{t})$ respectively in $L_2(\Omega \times \mathbb{S}^2)$, for any given $\epsilon > 0$, we can choose L_0 independent of t such that for all $L, M \geq L_0$,

$$\mathbb{E} \left[\|U_L^H(\hat{t}) - U_M^H(\hat{t}) - U_L^H(t) + U_M^H(t)\|_{L_2(\mathbb{S}^2)}^2 \right] \leq \frac{|\hat{t} - t|^2}{(T - t_0)^2} \epsilon. \quad (51)$$

Therefore,

$$\mathbb{E} \left[\|\phi_L(t) - \phi_M(t)\|_{L_2(\mathbb{S}^2)}^2 \right] \leq \epsilon. \quad (52)$$

This shows that $\{\phi_L\}$ is a Cauchy sequence in $L_2(\Omega \times \mathbb{S}^2)$ and hence for $t \in [t_0, T]$ and $t \neq \hat{t}$,

$$\mathbb{E} \left[\|\phi(t) - \phi_L(t)\|_{L_2(\mathbb{S}^2)}^2 \right] \leq \epsilon. \quad (53)$$

Now, by taking the limit as $t \rightarrow \hat{t}$ in (53), with the help of (49) and (50) we obtain

$$\mathbb{E} \left[\left\| \frac{dU^H}{dt}(\hat{t}) - V_L^H(\hat{t}) \right\|_{L_2(\mathbb{S}^2)}^2 \right] \leq \epsilon,$$

which completes the proof.

Let G_L be defined as

$$G_L(t) := \sum_{\ell=1}^L (-\lambda_\ell) E_\alpha(-\lambda_\ell t^\alpha) \sum_{m=-\ell}^{\ell} \hat{\xi}_{\ell,m} Y_{\ell,m}. \quad (54)$$

Using (54) with the Parseval formula, (13) and the upper bound (33) then for $L < M$ we have

$$\begin{aligned} \mathbb{E} \|G_L(t) - G_M(t)\|_{L_2(\mathbb{S}^2)}^2 &= \sum_{\ell=L+1}^M (\lambda_\ell)^2 (E_\alpha(-\lambda_\ell t^\alpha))^2 (2\ell + 1) C_\ell \\ &\leq (\Gamma(1 + \alpha) t^{-\alpha})^2 \sum_{\ell=L+1}^M (2\ell + 1) C_\ell \\ &\leq (\Gamma(1 + \alpha) t_0^{-\alpha})^2 \sum_{\ell=L+1}^M (2\ell + 1) C_\ell. \end{aligned} \quad (55)$$

For a given $\epsilon > 0$, using the condition (15), there is L_0 independent of t such that for $L, M \geq L_0$, the RHS of (55) can be smaller than ϵ . We then let

$$G(t) := \sum_{\ell=1}^{\infty} (-\lambda_\ell) E_\alpha(-\lambda_\ell t^\alpha) \sum_{m=-\ell}^{\ell} \hat{\xi}_{\ell,m} Y_{\ell,m}, \quad (56)$$

which is convergent in $L_2(\Omega \times \mathbb{S}^2)$.

Lemma 3.3. *Let ξ be a centered, 2-weakly isotropic Gaussian random field on \mathbb{S}^2 . Let $\{C_\ell : \ell \in \mathbb{N}_0\}$, the angular power spectrum of ξ , satisfy (15). Let $t_0 > 0$ be given. For $t \geq t_0$ there holds*

$$\lim_{L \rightarrow \infty} \mathbb{E} \left[\|D_t^{1-\alpha} G_L(t) - D_t^{1-\alpha} G(t)\|_{L_2(\mathbb{S}^2)}^2 \right] = 0,$$

where G_L and G are given by (54) and (56) respectively.

PROOF. Using Lemma 3.10 in [13] we have

$$D_t^{1-\alpha} G_L(t) = V_L^H(t),$$

which implies, by Lemma 3.1, that $\{D_t^{1-\alpha} G_L\}$ is a Cauchy sequence in $L_2(\Omega \times \mathbb{S}^2)$, i.e., we can choose L_0 independent of t such that for all $L, M \geq L_0$, there holds

$$\mathbb{E} \left[\|D_t^{1-\alpha} G_L(t) - D_t^{1-\alpha} G_M(t)\|_{L_2(\mathbb{S}^2)}^2 \right] \leq \epsilon.$$

Hence,

$$\mathbb{E} \left[\|D_t^{1-\alpha} G_L(t) - D_t^{1-\alpha} G(t)\|_{L_2(\mathbb{S}^2)}^2 \right] \leq \epsilon,$$

which completes the proof.

Lemma 3.4. For $t \in [t_0, T]$, we have

$$\lim_{L \rightarrow \infty} \mathbb{E} \left[\left\| V_L^H(t) - D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U^H(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] = 0.$$

PROOF. Using Theorem 4.1.2 (see [3, Satz 4.1.2]), we can show that

$$\lim_{L \rightarrow \infty} \mathbb{E} \left[\|\Delta_{\mathbb{S}^2} U^H(t) - G_L(t)\|_{L_2(\mathbb{S}^2)}^2 \right] = 0.$$

By Lemma 3.3, noting that $D_t^{1-\alpha} G_L(t) = V_L^H(t)$, we obtain the conclusion of the lemma.

Proposition 3.1. Let the angular power spectrum $\{C_\ell : \ell \in \mathbb{N}_0\}$ of the isotropic Gaussian random field ξ satisfy assumption (15). Then the random field U^H defined by (45) satisfies (4), in the $L_2(\Omega \times \mathbb{S}^2)$ sense, under the initial condition $U^H(0) = \xi$. In particular, for a given $t_0 > 0$, we have

$$\sup_{t \geq t_0} \mathbb{E} \left[\left\| \frac{d}{dt} U^H(t) - D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U^H(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] = 0.$$

PROOF. Let us fix $L \geq 1$, then we have

$$\begin{aligned} \frac{d}{dt} U_L^H(t) &= \sum_{\ell=1}^L \sum_{m=-\ell}^{\ell} \frac{d}{dt} E_\alpha(-\lambda_\ell t^\alpha) \widehat{\xi}_{\ell,m} Y_{\ell,m} \\ &= \sum_{\ell=1}^L \sum_{m=-\ell}^{\ell} (-\lambda_\ell) t^{\alpha-1} E_{\alpha,\alpha}(-\lambda_\ell t^\alpha) \widehat{\xi}_{\ell,m} Y_{\ell,m} = V_L^H(t). \end{aligned}$$

Using Lemma 3.10 in [13] we have

$$\begin{aligned} D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U_L^H(t) &= \sum_{\ell=0}^L \sum_{m=-\ell}^{\ell} (-\lambda_\ell) D_t^{1-\alpha} E_\alpha(-\lambda_\ell t^\alpha) \widehat{\xi}_{\ell,m} Y_{\ell,m} \\ &= \sum_{\ell=1}^L \sum_{m=-\ell}^{\ell} \frac{d}{dt} (E_\alpha(-\lambda_\ell t^\alpha) - 1) \widehat{\xi}_{\ell,m} Y_{\ell,m} \\ &= \sum_{\ell=1}^L \sum_{m=-\ell}^{\ell} (-\lambda_\ell) t^{\alpha-1} E_{\alpha,\alpha}(-\lambda_\ell t^\alpha) \widehat{\xi}_{\ell,m} Y_{\ell,m} = V_L^H(t). \end{aligned}$$

Combining the above two equations, then for any fixed $\omega \in \Omega$ we obtain

$$\frac{d}{dt} U_L^H(t) - D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U_L^H(t) = V_L^H(t) - V_L^H(t) = 0. \quad (57)$$

It follows from (57) that

$$\mathbb{E} \left[\left\| \frac{d}{dt} U_L^H(t) - D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U_L^H(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] = 0.$$

By the triangle inequality

$$\begin{aligned} \mathbb{E} \left[\left\| \frac{d}{dt} U^H(t) - D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U^H(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] &\leq 2\mathbb{E} \left[\left\| \frac{d}{dt} U^H(t) - \frac{d}{dt} U_L^H(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] \\ &+ 2\mathbb{E} \left[\left\| D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U_L^H(t) - D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U^H(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right]. \end{aligned}$$

By Lemma 3.2, we can choose L so that

$$\mathbb{E} \left[\left\| \frac{d}{dt} U^H(t) - \frac{d}{dt} U_L^H(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] \leq \frac{\epsilon}{4}. \quad (58)$$

By Lemma 3.4, we can choose L so that

$$\mathbb{E} \left[\left\| D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U_L^H(t) - D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U^H(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] \leq \frac{\epsilon}{4}. \quad (59)$$

So the proposition is proved.

The following result shows that the solution U^H to the equation (4) is a centered, 2-weakly isotropic Gaussian random field on \mathbb{S}^2 . Here we assume that the angular power spectrum $\{C_\ell : \ell \in \mathbb{N}_0\}$ of ξ decays algebraically with order $\kappa_1 > 2$, i.e., there exist constants $\tilde{D}, \tilde{C} > 0$ such that

$$C_\ell \leq \begin{cases} \tilde{D}, & \ell = 0, \\ \tilde{C}\ell^{-\kappa_1}, & \ell \geq 1, \kappa_1 > 2. \end{cases} \quad (60)$$

For $\kappa_1 > 2$ and $\tilde{C} > 0$, we let

$$\tilde{C}_{\kappa_1} := \left(\tilde{C} \left(\frac{2}{\kappa_1 - 2} + \frac{1}{\kappa_1 - 1} \right) \right)^{1/2}. \quad (61)$$

Proposition 3.2. *Let the field U^H , given in (45), be the solution to the equation (4) under the initial condition $U^H(0) = \xi$, where ξ is a centered, 2-weakly isotropic Gaussian random field on \mathbb{S}^2 . Let $\{C_\ell : \ell \in \mathbb{N}_0\}$, the angular power spectrum of ξ , satisfy (60). For a fixed $t \in (0, \infty)$, $U^H(t)$ is a centred, 2-weakly isotropic Gaussian random field on \mathbb{S}^2 , and its random coefficients*

$$\widehat{U}_{\ell,m}^H(t) = E_\alpha(-\lambda_\ell t^\alpha) \widehat{\xi}_{\ell,m}, \quad (62)$$

satisfy for $\ell, \ell' \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$ and $m' = -\ell', \dots, \ell'$,

$$\mathbb{E} \left[\widehat{U}_{\ell,m}^H(t) \overline{\widehat{U}_{\ell',m'}^H(t)} \right] = (E_\alpha(-\lambda_\ell t^\alpha))^2 C_\ell \delta_{\ell\ell'} \delta_{mm'}, \quad (63)$$

where λ_ℓ is given in (9) and $\delta_{\ell\ell'}$ is the Kronecker delta function.

PROOF. Consider the field U^H defined by (45). Since ξ is centred we have $\mathbb{E}[U^H(t)] = 0$. Let $t \in (0, \infty)$ and $\mathbf{x}, \mathbf{y} \in \mathbb{S}^2$, then by (45), we can write

$$\mathbb{E} [U^H(\mathbf{x}, t) U^H(\mathbf{y}, t)] = \sum_{\ell=0}^{\infty} \sum_{\ell'=0}^{\infty} \sum_{m=-\ell}^{\ell} \sum_{m'=-\ell'}^{\ell'} E_\alpha(-\lambda_\ell t^\alpha) E_\alpha(-\lambda_{\ell'} t^\alpha)$$

$$\times \mathbb{E} \left[\widehat{\xi}_{\ell,m} \widehat{\xi}_{\ell',m'} \right] Y_{\ell,m}(\mathbf{x}) \overline{Y_{\ell',m'}(\mathbf{y})}.$$

Since ξ is a centred, 2-weak isotropic Gaussian random field we have $\mathbb{E} \left[\widehat{\xi}_{\ell,m} \widehat{\xi}_{\ell',m'} \right] = C_\ell \delta_{\ell\ell'} \delta_{mm'}$, and by the addition theorem (see equation (8)) and Proposition 2.1, we have

$$\begin{aligned} \mathbb{E} [U^H(\mathbf{x}, t) U^H(\mathbf{y}, t)] &= \sum_{\ell=0}^{\infty} (E_\alpha(-\lambda_\ell t^\alpha))^2 C_\ell \sum_{m=-\ell}^{\ell} Y_{\ell,m}(\mathbf{x}) \overline{Y_{\ell,m}(\mathbf{y})} \\ &= \sum_{\ell=0}^{\infty} (E_\alpha(-\lambda_\ell t^\alpha))^2 (2\ell + 1) C_\ell P_\ell(\mathbf{x} \cdot \mathbf{y}), \end{aligned}$$

where $P_\ell(\cdot)$, $\ell \in \mathbb{N}_0$, is the Legendre polynomial of degree ℓ . As $P_\ell(\mathbf{x} \cdot \mathbf{y})$ depends only on the inner product of \mathbf{x} and \mathbf{y} , we conclude that the covariance function $\mathbb{E} [U^H(\mathbf{x}, t) U^H(\mathbf{y}, t)]$ is rotationally invariant. Note that for $\ell, \ell' \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$ and $m' = -\ell', \dots, \ell'$, by the 2-weak isotropy of ξ , and Proposition 2.1,

$$\begin{aligned} \mathbb{E} \left[\widehat{U}_{\ell,m}^H(t) \overline{\widehat{U}_{\ell',m'}^H(t)} \right] &= E_\alpha(-\lambda_\ell t^\alpha) E_\alpha(-\lambda_{\ell'} t^\alpha) \mathbb{E} \left[\widehat{\xi}_{\ell,m} \widehat{\xi}_{\ell',m'} \right] \\ &= (E_\alpha(-\lambda_\ell t^\alpha))^2 C_\ell \delta_{\ell\ell'} \delta_{mm'}. \end{aligned}$$

To prove that $U^H(t)$ is Gaussian, first note that its variance can be bounded, using (60) and that $0 < E_\alpha(\cdot) \leq 1$, by

$$\begin{aligned} \text{Var} [U^H(t)] &= \sum_{\ell=0}^{\infty} (E_\alpha(-\lambda_\ell t^\alpha))^2 (2\ell + 1) C_\ell \\ &\leq \tilde{D} + \tilde{C} \sum_{\ell=1}^{\infty} (2\ell + 1) \ell^{-\kappa_1} \\ &\leq \tilde{D} + \tilde{C} \int_1^{\infty} (2x^{1-\kappa_1} + x^{-\kappa_1}) dx \\ &\leq \tilde{D} + \tilde{C} \left(\frac{2}{\kappa_1 - 2} + \frac{1}{\kappa_1 - 1} \right) \\ &\leq \tilde{D} + \tilde{C}_{\kappa_1}^2 < \infty. \end{aligned} \tag{64}$$

Now let $T_\ell(t)$, $\ell \in \mathbb{N}_0$, be defined as

$$T_\ell(t) := \sum_{m=-\ell}^{\ell} \widehat{U}_{\ell,m}^H(t) Y_{\ell,m} = \sum_{m=-\ell}^{\ell} E_\alpha(-\lambda_\ell t^\alpha) \widehat{\xi}_{\ell,m} Y_{\ell,m},$$

and note that since $\widehat{\xi}_{\ell,m}$ are centred, independent Gaussian random variables (by the 2-weak isotropy of ξ), then T_ℓ is a Gaussian random variable with mean zero and variance, using (63), (8), and properties of spherical harmonics, given by

$$\text{Var}[T_\ell] = (E_\alpha(-\lambda_\ell t^\alpha))^2 (2\ell + 1) C_\ell. \tag{65}$$

Let $U_N^H(t)$, $N \geq 1$, be defined as

$$U_N^H(t) := \sum_{\ell=0}^N T_\ell(t).$$

It is easy to see, using (65), that

$$\text{Var}[U_N^H(t)] = \sum_{\ell=0}^N \text{Var}[T_\ell(t)] = \sum_{\ell=0}^N (E_\alpha(-\lambda_\ell t^\alpha))^2 (2\ell + 1) C_\ell, \tag{66}$$

and, using (64),

$$\lim_{N \rightarrow \infty} \text{Var}[U_N^H(t)] = \sum_{\ell=0}^{\infty} (E_{\alpha}(-\lambda_{\ell} t^{\alpha}))^2 (2\ell + 1) C_{\ell} < \infty. \quad (67)$$

The characteristic function $\varphi_{U_N^H(t)}(r)$ of $U_N^H(t)$ can be written, using (65) and (66), as

$$\begin{aligned} \varphi_{U_N^H(t)}(r) &= \mathbb{E} \left[e^{irU_N^H(t)} \right] = \prod_{\ell=0}^N \mathbb{E} \left[e^{irT_{\ell}(t)} \right] = \prod_{\ell=0}^N \varphi_{T_{\ell}(t)}(r) \\ &= e^{-\frac{1}{2} r^2 \sum_{\ell=0}^N (E_{\alpha}(-\lambda_{\ell} t^{\alpha}))^2 (2\ell + 1) C_{\ell}}, \end{aligned}$$

since T_{ℓ} , $\ell \geq 0$, are independent Gaussian random variables.

Hence,

$$\lim_{N \rightarrow \infty} \varphi_{U_N^H(t)}(r) = e^{-\frac{1}{2} r^2 \sum_{\ell=0}^{\infty} (E_{\alpha}(-\lambda_{\ell} t^{\alpha}))^2 (2\ell + 1) C_{\ell}},$$

which is the characteristic function of some Gaussian random variable with mean zero and variance $\sum_{\ell=0}^{\infty} (E_{\alpha}(-\lambda_{\ell} t^{\alpha}))^2 (2\ell + 1) C_{\ell} < \infty$ (by (67)). Then by the continuity theorem (see [10], Theorem 3.3.6), we conclude that the field U^H is Gaussian, thus completing the proof.

Remark 3.1. Let $\{C_{\ell} : \ell \in \mathbb{N}_0\}$ be the angular power spectrum of ξ . Then the Fourier coefficients $\widehat{U}^H_{\ell,m}$ of U^H can be written, using (62) and Proposition 2.1, as

$$\widehat{U}^H_{\ell,m}(t) := \begin{cases} \sqrt{C_{\ell}} E_{\alpha}(-\lambda_{\ell} t^{\alpha}) Z_{\ell,0}^{(1)}, & m = 0, \\ \sqrt{\frac{C_{\ell}}{2}} E_{\alpha}(-\lambda_{\ell} t^{\alpha}) \left(Z_{\ell,m}^{(1)} - i Z_{\ell,m}^{(2)} \right), & m = 1, \dots, \ell, \\ (-1)^m \sqrt{\frac{C_{\ell}}{2}} E_{\alpha}(-\lambda_{\ell} t^{\alpha}) \left(Z_{\ell,|m|}^{(1)} + i Z_{\ell,|m|}^{(2)} \right), & m = -\ell, \dots, -1, \end{cases} \quad (68)$$

where $Z_{\ell,m_1}^{(1)}, Z_{\ell,m_2}^{(2)} \in \mathcal{Z}_{\ell}$, $\ell \in \mathbb{N}_0, m_1 = 0, \dots, \ell, m_2 = 1, 2, \dots, \ell$ and \mathcal{Z}_{ℓ} is defined by (16).

Moreover, the solution $U^H(t)$, $t \in (0, \infty)$, can be represented, using (68), as

$$\begin{aligned} U^H(t) &= \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \widehat{U}^H_{\ell,m}(t) Y_{\ell,m} \\ &= \sum_{\ell=0}^{\infty} \left(E_{\alpha}(-\lambda_{\ell} t^{\alpha}) \left(\sqrt{C_{\ell}} Z_{\ell,0}^{(1)} Y_{\ell,0} \right. \right. \\ &\quad \left. \left. + \sqrt{2C_{\ell}} \sum_{m=1}^{\ell} \left(Z_{\ell,m}^{(1)} \text{Re} Y_{\ell,m}(\mathbf{x}) + Z_{\ell,m}^{(2)} \text{Im} Y_{\ell,m} \right) \right) \right), \end{aligned} \quad (69)$$

where the above expansion is convergent in $L_2(\Omega \times \mathbb{S}^2)$.

4. Solution of the inhomogeneous problem

Consider a random field $U^I(t) \in L_2(\Omega \times \mathbb{S}^2)$, $t \in (0, \infty)$. We say that U^I is a solution to (5) if the equality in (6) holds in the sense of $L_2(\Omega \times \mathbb{S}^2)$, that is

$$\sup_{t \geq \tau} \mathbb{E} \left\| U^I(t) - \frac{1}{\Gamma(\alpha)} \int_{\tau}^t \frac{\Delta_{\mathbb{S}^2} U^I(s)}{(t-s)^{1-\alpha}} ds - W_{\tau}(t) \right\|_{L_2(\mathbb{S}^2)}^2 = 0.$$

This section derives the solution $U^I(t) \in L_2(\Omega \times \mathbb{S}^2)$, $t \in (0, \infty)$, to the inhomogeneous equation (5), under the condition $U^I(t) = 0$ for $t \in (0, \tau]$.

Note that since $U^I \in L_2(\Omega \times \mathbb{S}^2)$, we can write

$$U^I(t) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \widehat{U}^I_{\ell,m}(t) Y_{\ell,m}, \quad t \in (0, \infty),$$

where the random coefficients $\widehat{U}^I_{\ell,m}$, $\ell \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$, are given by

$$\widehat{U}^I_{\ell,m}(t) = \int_{\mathbb{S}^2} U^I(\mathbf{x}, t) \overline{Y_{\ell,m}(\mathbf{x})} \mu(d\mathbf{x}), \quad t \in (0, \infty).$$

By multiplying both sides of equation (5) by $\overline{Y_{\ell,m}}$ and integrating over \mathbb{S}^2 , we obtain a set of ordinary differential equations

$$d\widehat{U}^I_{\ell,m} + \lambda_{\ell} D_t^{1-\alpha} \widehat{U}^I_{\ell,m} dt = d\widehat{W}_{\ell,m,\tau}, \quad \widehat{U}^I_{\ell,m}(0) = 0, \quad (70)$$

where $\widehat{W}_{\ell,m,\tau}$ are the Fourier coefficients of W_{τ} .

If $\widehat{W}_{\ell,m,\tau}$ were deterministic differentiable functions in time, then we could write (70) as

$$\frac{d}{dt} \widehat{U}^I_{\ell,m} + \lambda_{\ell} D_t^{1-\alpha} \widehat{U}^I_{\ell,m} = \frac{d}{dt} \widehat{W}_{\ell,m,\tau}, \quad \widehat{U}^I_{\ell,m}(0) = 0. \quad (71)$$

By taking the Laplace transform of (71) (see equation (43)), we get

$$z \widetilde{U}^I_{\ell,m}(z) - \widehat{U}^I_{\ell,m}(0) + \lambda_{\ell} z^{1-\alpha} \widetilde{U}^I_{\ell,m}(z) = e^{-z\tau} \widetilde{W}_{\ell,m,0}(z), \quad (72)$$

where $\widetilde{U}^I_{\ell,m}(z)$ is the Laplace transform of $\widehat{U}^I_{\ell,m}(t)$ and $e^{-z\tau} \widetilde{W}_{\ell,m,0}(z)$ is the Laplace transform of $\frac{d}{dt} \widehat{W}_{\ell,m,\tau}(t)$.

Now solving equation (72) for $\widetilde{U}^I_{\ell,m}(z)$, we obtain

$$\widetilde{U}^I_{\ell,m}(z) = e^{-z\tau} \frac{\widetilde{W}_{\ell,m,0}(z)}{z + \lambda_{\ell} z^{1-\alpha}}. \quad (73)$$

Using the convolution theorem (see [11], Theorem 3.2), we could write

$$\mathcal{L}^{-1} \left\{ \frac{\widetilde{W}_{\ell,m,0}(z)}{z + \lambda_{\ell} z^{1-\alpha}} \right\} = \int_0^t E_{\alpha}(-\lambda_{\ell}(t-s)^{\alpha}) \frac{d}{ds} \widehat{W}_{\ell,m,0}(s) ds,$$

where $\mathcal{L}^{-1}(\cdot)$ denotes the inverse Laplace transform.

Now by taking the inverse Laplace transform in (73) with the help of the second shift theorem (see [11], Theorem 2.4) we obtain

$$\widehat{U}^I_{\ell,m}(t) = \mathcal{H}(t - \tau) \int_0^{t-\tau} E_{\alpha}(-\lambda_{\ell}(t - \tau - s)^{\alpha}) \frac{d}{ds} \widehat{W}_{\ell,m,0}(s) ds,$$

where $\mathcal{H}(\cdot)$ is the Heaviside unit step function (i.e., $\mathcal{H}(t) = 1$, for $t \geq 0$, and zero for $t < 0$).

Since $\widehat{W}_{\ell,m,0}(\cdot)$ are 1-dimensional, real-valued Brownian motions which are continuous but nowhere differentiable, we have to write instead

$$\widehat{U}^I_{\ell,m}(t) = \mathcal{H}(t - \tau) \int_0^{t-\tau} E_{\alpha}(-\lambda_{\ell}(t - \tau - s)^{\alpha}) d\widehat{W}_{\ell,m,0}(s),$$

where the integral is in the Itô sense.

Thus the solution of equation (5) takes the form, for $t > \tau$,

$$U^I(t) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \int_0^{t-\tau} E_{\alpha}(-\lambda_{\ell}(t-\tau-s)^{\alpha}) d\widehat{W}_{\ell,m,0}(s) Y_{\ell,m}, \quad (74)$$

and $U^I(t) = 0$ for $t \leq \tau$.

By the condition (19) and Remark 2.2, for every $\ell \in \mathbb{N}_0$, we have $\sigma_{\ell,t,\alpha}^2 \leq t$, which guarantees the convergence of (74) in the $L_2(\Omega \times \mathbb{S}^2)$ space.

For $L \in \mathbb{N}$, let

$$V_L^I(t) := D_t^{1-\alpha} \Delta_{\mathbb{S}^2} U_L^I(t) = \frac{1}{\Gamma(\alpha)} \frac{d}{dt} \int_{\tau}^t \frac{\Delta_{\mathbb{S}^2} U_L^I(s)}{(t-s)^{1-\alpha}} ds. \quad (75)$$

Remark 4.1. By Lemma 2.1, we can express V_L^I in the form

$$\begin{aligned} V_L^I(t) &= \sum_{\ell=0}^L \sum_{m=-\ell}^{\ell} -\lambda_{\ell} D_t^{1-\alpha} \int_{\tau}^t E_{\alpha}(-\lambda_{\ell}(t-\tau-r)^{\alpha}) d\widehat{W}_{\ell,m,\tau}(r) Y_{\ell,m} \\ &= \sum_{\ell=0}^L \sum_{m=-\ell}^{\ell} -\lambda_{\ell} \int_{\tau}^t D_r^{1-\alpha} E_{\alpha}(-\lambda_{\ell} r^{\alpha}) d\widehat{W}_{\ell,m,\tau}(r) Y_{\ell,m} \\ &= \sum_{\ell=0}^L \sum_{m=-\ell}^{\ell} \int_{\tau}^t \left(\frac{-\lambda_{\ell}}{\Gamma(\alpha)} \frac{d}{dr} \int_0^r \frac{E_{\alpha}(-\lambda_{\ell} s^{\alpha})}{(r-s)^{1-\alpha}} ds \right) d\widehat{W}_{\ell,m,\tau}(r) Y_{\ell,m} \\ &= \sum_{\ell=0}^L \sum_{m=-\ell}^{\ell} \int_{\tau}^t \frac{d}{dr} \left(E_{\alpha}(-\lambda_{\ell} r^{\alpha}) - 1 \right) d\widehat{W}_{\ell,m,\tau}(r) Y_{\ell,m} \\ &= \sum_{\ell=0}^L \sum_{m=-\ell}^{\ell} \int_{\tau}^t (-\lambda_{\ell}) r^{\alpha-1} E_{\alpha,\alpha}(-\lambda_{\ell} r^{\alpha}) d\widehat{W}_{\ell,m,\tau}(r) Y_{\ell,m}, \end{aligned} \quad (76)$$

where the third step used Lemma 3.10 in [13].

Lemma 4.1. Let W_{τ} be an $L_2(\mathbb{S}^2)$ -valued time-delayed Brownian motion. Let $\{\mathcal{A}_{\ell} : \ell \in \mathbb{N}_0\}$, the angular power spectrum of W_{τ} , satisfy (19). Then, as $L \rightarrow \infty$, V_L^I is convergent to

$$V^I(t) := \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \int_{\tau}^t (-\lambda_{\ell}) s^{\alpha-1} E_{\alpha,\alpha}(-\lambda_{\ell} s^{\alpha}) d\widehat{W}_{\ell,m,\tau}(s) Y_{\ell,m}, \quad t \geq \tau, \quad (77)$$

in the following sense

$$\sup_{t \geq \tau} \mathbb{E} \left\| V_L^I(t) - V^I(t) \right\|_{L_2(\mathbb{S}^2)}^2 \rightarrow 0, \quad \text{as } L \rightarrow \infty.$$

PROOF. For $t \geq \tau$, $L < M$, by Parseval's formula and Itô's isometry, we have

$$\begin{aligned} \mathbb{E} \left\| V_L^I(t) - V_M^I(t) \right\|_{L_2(\mathbb{S}^2)}^2 &= \mathbb{E} \left[\sum_{\ell=L+1}^M \sum_{m=-\ell}^{\ell} \lambda_{\ell}^2 \int_{\tau}^t (-\lambda_{\ell}) s^{\alpha-1} E_{\alpha,\alpha}(-\lambda_{\ell} s^{\alpha}) d\widehat{W}_{\ell,m,\tau}(s) Y_{\ell,m} \right] \\ &= \sum_{\ell=L+1}^M \lambda_{\ell}^2 (2\ell+1) \mathcal{A}_{\ell} \int_{\tau}^t s^{2\alpha-2} (E_{\alpha,\alpha}(-\lambda_{\ell} s^{\alpha}))^2 ds \text{ by using (13)} \\ &\leq C^2 \int_{\tau}^t s^{-2} ds \sum_{\ell=L+1}^M (2\ell+1) \mathcal{A}_{\ell} \end{aligned}$$

$$\leq C^2 \tau^{-1} \sum_{\ell=L+1}^M (2\ell + 1) \mathcal{A}_\ell, \quad (78)$$

where the third step used (26).

For a given $\epsilon > 0$, using the condition (19), there is L_0 independent of t such that for $L, M \geq L_0$, the RHS of (78) is smaller than ϵ . So $\{V_L^I\}$ is a Cauchy sequence in $L_2(\Omega \times \mathbb{S}^2)$ and hence it is convergent. We define the limit of V_L^I as $L \rightarrow \infty$ as in (77). Note that by the condition (19) we obtain

$$\mathbb{E} \left[\|V^I(t)\|_{L_2(\mathbb{S}^2)}^2 \right] \leq C^2 \tau^{-1} \sum_{\ell=0}^{\infty} (2\ell + 1) \mathcal{A}_\ell < \infty, \quad (79)$$

which guarantees that V^I is well-defined in the $L_2(\Omega \times \mathbb{S}^2)$ sense.

Remark 4.2. Let $T > \tau > 0$, then by (78) we obtain

$$\int_{\tau}^T \mathbb{E} \left[\|V_L^I(s)\|_{L_2(\mathbb{S}^2)}^2 \right] ds \leq C^2 \tau^{-1} T \sum_{\ell=0}^{\infty} (2\ell + 1) \mathcal{A}_\ell < \infty,$$

which implies, by Lemma 4.1 and the dominated convergence theorem, that $\int_{\tau}^T V_L^I(s) ds$ is convergent as $L \rightarrow \infty$, to $\int_{\tau}^T V^I(s) ds$, in the $L_2(\Omega \times \mathbb{S}^2)$ sense.

For $L \in \mathbb{N}$, $t > \tau$, we define

$$U_L^I(t) := \int_{\tau}^t V_L^I(s) ds + W_{\tau}^{(L)}(t), \quad (80)$$

where

$$W_{\tau}^{(L)}(t) := \sum_{\ell=0}^L \sum_{m=-\ell}^{\ell} \widehat{W}_{\ell,m,\tau}(t) Y_{\ell,m}, \quad (81)$$

and V_L^I is defined by (76).

Lemma 4.2. Let W_{τ} be an $L_2(\mathbb{S}^2)$ -valued time-delayed Brownian motion. Let $\{\mathcal{A}_\ell : \ell \in \mathbb{N}_0\}$, the angular power spectrum of W_{τ} , satisfy the condition (19). For $t \in [\tau, T]$, then U_L^I is convergent as $L \rightarrow \infty$, to $U^I(t) := \int_{\tau}^t V^I(s) ds + W_{\tau}(t)$, in the following sense

$$\sup_{t \in [\tau, T]} \mathbb{E} \left[\left\| U_L^I(t) - U^I(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] \rightarrow 0, \quad \text{as } L \rightarrow \infty.$$

PROOF. For $t \geq \tau$, $L < M$, we have, using (80),

$$\begin{aligned} \mathbb{E} \left[\left\| U_L^I(t) - U_M^I(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] &= \mathbb{E} \left[\left\| \int_{\tau}^t V_L^I(s) ds - \int_{\tau}^t V_M^I(s) ds \right. \right. \\ &\quad \left. \left. + \sum_{\ell=L+1}^M \sum_{m=-\ell}^{\ell} \widehat{W}_{\ell,m,\tau}(s) Y_{\ell,m} \right\|_{L_2(\mathbb{S}^2)}^2 \right] \\ &\leq \mathbb{E} \left[\left\| \int_{\tau}^t V_L^I(s) ds - \int_{\tau}^t V_M^I(s) ds \right\|_{L_2(\mathbb{S}^2)}^2 \right] \\ &\quad + (t - \tau) \sum_{\ell=L+1}^M (2\ell + 1) \mathcal{A}_\ell, \end{aligned} \quad (82)$$

where the second step used Parseval's formula.

By Remark 4.2, there is L_0 independent of t such that for $L, M \geq L_0$, the term

$$\mathbb{E} \left[\left\| \int_{\tau}^t V_L^I(s) ds - \int_{\tau}^t V_M^I(s) ds \right\|_{L_2(\mathbb{S}^2)}^2 \right] \leq \epsilon/2,$$

for $\epsilon > 0$. Similarly, using the condition (19), the second term of (82) can be smaller than $\epsilon/2$. So $\{U_L^I\}$ is a Cauchy sequence in $L_2(\Omega \times \mathbb{S}^2)$ and hence it is convergent. Thus the proof is complete.

Proposition 4.1. *Let the angular power spectrum $\{\mathcal{A}_{\ell} : \ell \in \mathbb{N}_0\}$ of W_{τ} satisfy assumption (19). Then the random field $U^I(t)$, $t > \tau$, defined by (74), satisfies (5) in the $L_2(\Omega \times \mathbb{S}^2)$ sense. In particular, for $t \in [\tau, T]$, there holds*

$$\sup_{t \in [\tau, T]} \mathbb{E} \left\| U^I(t) - \frac{1}{\Gamma(\alpha)} \int_{\tau}^t \frac{\Delta_{\mathbb{S}^2} U^I(s)}{(t-s)^{1-\alpha}} ds - W_{\tau}(t) \right\|_{L_2(\mathbb{S}^2)}^2 = 0. \quad (83)$$

PROOF. By Lemma 4.2 there holds

$$\sup_{t \in [\tau, T]} \mathbb{E} \left\| U^I(t) - \int_{\tau}^t V^I(s) ds - W_{\tau}(t) \right\|_{L_2(\mathbb{S}^2)}^2 = 0.$$

Also, using (75) we obtain

$$\int_{\tau}^t V^I(s) ds = \frac{1}{\Gamma(\alpha)} \int_{\tau}^t \frac{\Delta_{\mathbb{S}^2} U^I(s)}{(t-s)^{1-\alpha}} ds,$$

which is convergent in $L_2(\Omega \times \mathbb{S}^2)$ due to (79). Thus the proof is complete.

The following result shows that the inhomogeneous solution $U^I(t)$, $t \in (\tau, \infty)$, to the equation (5) is a centered, 2-weakly isotropic Gaussian random field. Here we assume that the angular power spectrum $\{\mathcal{A}_{\ell} : \ell \in \mathbb{N}_0\}$ of W_{τ} , decays algebraically with order $\kappa_2 > 2$, i.e., there exist constants $\tilde{K}, \tilde{A} > 0$ such that

$$\mathcal{A}_{\ell} \leq \begin{cases} \tilde{K}, & \ell = 0, \\ \tilde{A} \ell^{-\kappa_2}, & \ell \geq 1, \kappa_2 > 2. \end{cases} \quad (84)$$

Let \tilde{A}_{κ_2} and $\gamma_{\alpha}(\kappa_2)$ be defined as

$$\tilde{A}_{\kappa_2} := \left(\tilde{A} \left(\frac{2}{\kappa_2 - 2} + \frac{1}{\kappa_2 - 1} \right) \right)^{1/2} \quad (85)$$

and

$$\gamma_{\alpha}(\kappa_2) := \begin{cases} \kappa_2 + 2, & \alpha \in (0, \frac{1}{2}), \\ \kappa_2 + \frac{2}{\alpha} - 2, & \alpha \in (\frac{1}{2}, 1], \\ \kappa_2, & \alpha = \frac{1}{2}. \end{cases} \quad (86)$$

Proposition 4.2. *Let the field U^I be defined as in (74), with the condition $U^I(t) = 0$, $t \in (0, \tau]$. Let $\widehat{W}_{\ell, m, 0}$ be the Fourier coefficients of W_0 with variances $\{\mathcal{A}_{\ell} : \ell \in \mathbb{N}_0\}$ given in (84). For a fixed $t \in (\tau, \infty)$, $U^I(t)$ is a centred, 2-weakly isotropic Gaussian random field on \mathbb{S}^2 , and its random coefficients*

$$\widehat{U}_{\ell, m}^I(t) = \int_0^{t-\tau} E_{\alpha}(-\lambda_{\ell}(t-\tau-s)^{\alpha}) d\widehat{W}_{\ell, m, 0}(s), \quad (87)$$

satisfy for $\ell, \ell' \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$ and $m' = -\ell', \dots, \ell'$,

$$\mathbb{E} \left[\widehat{U}_{\ell, m}^I(t) \overline{\widehat{U}_{\ell', m'}^I(t)} \right] = \mathcal{A}_{\ell} \sigma_{\ell, t-\tau, \alpha}^2 \delta_{\ell \ell'} \delta_{m m'}, \quad (88)$$

where λ_{ℓ} is given in (9), $\delta_{\ell \ell'}$ is the Kronecker delta function, and $\sigma_{\ell, t, \alpha}^2$ is given by (28).

PROOF. Consider the field U^I defined by (74). Since W_0 is centred we have $\mathbb{E}[U^I(t)] = 0$. Let $t \in (\tau, \infty)$ and $\mathbf{x}, \mathbf{y} \in \mathbb{S}^2$, then by using (74) we write

$$\begin{aligned} \mathbb{E}[U^I(\mathbf{x}, t)U^I(\mathbf{y}, t)] &= \sum_{\ell=0}^{\infty} \sum_{\ell'=0}^{\infty} \sum_{m=-\ell}^{\ell} \sum_{m'=-\ell'}^{\ell'} Y_{\ell,m}(\mathbf{x}) \overline{Y_{\ell',m'}(\mathbf{y})} \\ &\times \mathbb{E} \left[\left(\int_0^{t-\tau} E_{\alpha}(-\lambda_{\ell}(t-\tau-s)^{\alpha}) d\widehat{W}_{\ell,m,0}(s) \right) \overline{\left(\int_0^{t-\tau} E_{\alpha}(-\lambda_{\ell'}(t-\tau-v)^{\alpha}) d\widehat{W}_{\ell',m',0}(v) \right)} \right]. \end{aligned}$$

Using Itô's isometry (see [19]) and the addition theorem (see (8)) we can write

$$\begin{aligned} \mathbb{E}[U^I(\mathbf{x}, t)U^I(\mathbf{y}, t)] &= \sum_{\ell=0}^{\infty} \mathcal{A}_{\ell} \int_0^{t-\tau} (E_{\alpha}(-\lambda_{\ell}(t-\tau-s)^{\alpha}))^2 ds \sum_{m=-\ell}^{\ell} Y_{\ell,m}(\mathbf{x}) \overline{Y_{\ell,m}(\mathbf{y})} \\ &= \sum_{\ell=0}^{\infty} (2\ell+1) \mathcal{A}_{\ell} P_{\ell}(\mathbf{x} \cdot \mathbf{y}) \int_0^{t-\tau} (E_{\alpha}(-\lambda_{\ell}(t-\tau-s)^{\alpha}))^2 ds \\ &= \sum_{\ell=0}^{\infty} (2\ell+1) \mathcal{A}_{\ell} \sigma_{\ell,t-\tau,\alpha}^2 P_{\ell}(\mathbf{x} \cdot \mathbf{y}), \end{aligned}$$

where $P_{\ell}(\cdot)$, $\ell \in \mathbb{N}_0$, is the Legendre polynomial of degree ℓ . As $P_{\ell}(\mathbf{x} \cdot \mathbf{y})$ depends only on the inner product of \mathbf{x} and \mathbf{y} , we conclude that the covariance function $\mathbb{E}[U^I(\mathbf{x}, t)U^I(\mathbf{y}, t)]$ is rotationally invariant. Note that for $\ell, \ell' \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$ and $m' = -\ell', \dots, \ell'$, by the 2-weak isotropy of W_0 and Itô's isometry,

$$\mathbb{E} \left[\widehat{U}_{\ell,m}^I(t) \overline{\widehat{U}_{\ell',m'}^I(t)} \right] = \mathcal{A}_{\ell} \sigma_{\ell,t-\tau,\alpha}^2 \delta_{\ell\ell'} \delta_{mm'}.$$

To prove that $U^I(\mathbf{x}, t)$ is Gaussian, we note that its variance can be written, using (84), as

$$\begin{aligned} \text{Var}[U^I(\mathbf{x}, t)] &= \sum_{\ell=0}^{\infty} (2\ell+1) \mathcal{A}_{\ell} \sigma_{\ell,t-\tau,\alpha}^2 \\ &= \widetilde{K} \sigma_{0,t-\tau,\alpha}^2 + \sum_{\ell=1}^{\infty} (2\ell+1) \mathcal{A}_{\ell} \sigma_{\ell,t-\tau,\alpha}^2 \\ &\leq (t-\tau) \left(\widetilde{K} + \sum_{\ell=1}^{\infty} (2\ell+1) \mathcal{A}_{\ell} \right) \\ &\leq (t-\tau) \left(\widetilde{K} + \widetilde{A}_{k_2}^2 \right) < \infty, \end{aligned} \tag{89}$$

where \widetilde{A}_{k_2} is given by (85) and the last step uses (28) and steps similar to (64).

Note that $R_{\ell}(t) := \sum_{m=-\ell}^{\ell} \widehat{U}_{\ell,m}^I(t) Y_{\ell,m}$ is a centred, Gaussian random variable with mean zero and variance given, using (88), by $(2\ell+1) \mathcal{A}_{\ell} \sigma_{\ell,t-\tau,\alpha}^2$. Define $U_N^I(t) := \sum_{\ell=0}^N R_{\ell}(t)$, $N \geq 1$, then by (89) and similar steps as in (66) and (67), with direct application of the continuity theorem (see [10], Theorem 3.3.6), we conclude that the field U^I is Gaussian, thus completing the proof.

Remark 4.3. Let $\{\mathcal{A}_{\ell} : \ell \in \mathbb{N}_0\}$ be the angular power spectrum of W_0 . Then for $t > \tau$ the Fourier coefficients $\widehat{U}_{\ell,m}^I(t)$ of the field $U^I(t)$ can be written, using (23) and (74), as

$$\widehat{U}_{\ell,m}^I(t) := \begin{cases} \sqrt{\mathcal{A}_{\ell}} \mathcal{I}_{\ell,0,\alpha}^{(1)}(t-\tau), & m = 0, \\ \sqrt{\mathcal{A}_{\ell}/2} \left(\mathcal{I}_{\ell,m,\alpha}^{(1)}(t-\tau) - i \mathcal{I}_{\ell,m,\alpha}^{(2)}(t-\tau) \right), & m = 1, \dots, \ell, \\ (-1)^m \sqrt{\mathcal{A}_{\ell}/2} \left(\mathcal{I}_{\ell,|m|,\alpha}^{(1)}(t-\tau) + i \mathcal{I}_{\ell,|m|,\alpha}^{(2)}(t-\tau) \right), & m = -\ell, \dots, -1, \end{cases} \tag{90}$$

where $\mathcal{I}_{\ell,m,\alpha}^{(j)}(\cdot)$, $j = 1, 2$, are given in (29).

Moreover, for $t > \tau$, $U^I(t)$ can be expressed, using (90), as

$$\begin{aligned} U^I(t) &= \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \widehat{U}^I_{\ell,m}(t) Y_{\ell,m} \\ &= \sum_{\ell=0}^{\infty} \left(\sqrt{\mathcal{A}_{\ell}} \mathcal{I}_{\ell,0,\alpha}^{(1)}(t-\tau) Y_{\ell,0} \right. \\ &\quad \left. + \sqrt{2\mathcal{A}_{\ell}} \sum_{m=1}^{\ell} \left(\mathcal{I}_{\ell,m,\alpha}^{(1)}(t-\tau) \operatorname{Re} Y_{\ell,m} + \mathcal{I}_{\ell,m,\alpha}^{(2)}(t-\tau) \operatorname{Im} Y_{\ell,m} \right) \right), \end{aligned} \quad (91)$$

where the above expansion is convergent in $L_2(\Omega \times \mathbb{S}^2)$.

5. Solution of the time-fractional diffusion equation

This section demonstrates the solution $U(t)$, $t \in (0, \infty)$, to the time-fractional diffusion equation given in (1).

Let the conditions of Propositions 3.1 be satisfied. Then, the solution $U(t)$, $t \in (0, \infty)$, of equation (1) is given by

$$U(t) := U^H(t) + U^I(t), \quad t \in (0, \infty), \quad (92)$$

where U^H is given by (69) and $U^I(t) = 0$ for $t \leq \tau$, while for $t > \tau$, $U^I(t)$ is given by (91).

The following result shows that the solution $U(t)$, $t \in (0, \infty)$, to the equation (1) is a centred, 2-weakly isotropic Gaussian random field.

Proposition 5.1. *Let the field U , given by (92), be the solution to the equation (1). Then, for a fixed $t \in (0, \infty)$, $U(t)$ is a centred, 2-weakly isotropic Gaussian random field with random coefficients*

$$\widehat{U}_{\ell,m}(t) := \begin{cases} \widehat{U}^H_{\ell,m}(t), & t \leq \tau, \\ \widehat{U}^H_{\ell,m}(t) + \widehat{U}^I_{\ell,m}(t), & t > \tau, \end{cases} \quad (93)$$

where $\widehat{U}^H_{\ell,m}(t)$ and $\widehat{U}^I_{\ell,m}(t)$ are given by (62) and (87) respectively.

Moreover, for $\ell, \ell' \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$ and $m' = -\ell', \dots, \ell'$, there holds

$$\mathbb{E} \left[\widehat{U}_{\ell,m}(t) \overline{\widehat{U}_{\ell',m'}(t)} \right] = \delta_{\ell\ell'} \delta_{mm'} \begin{cases} C_{\ell}(E_{\alpha}(-\lambda_{\ell} t^{\alpha}))^2, & t \leq \tau, \\ C_{\ell}(E_{\alpha}(-\lambda_{\ell} t^{\alpha}))^2 + \mathcal{A}_{\ell} \sigma_{\ell,t-\tau,\alpha}^2, & t > \tau. \end{cases}$$

PROOF. For $t \leq \tau$ the result follows by Proposition 3.2. Let $t > \tau$, then by Propositions 3.2, 4.2, and the uncorrelatedness of U^H and U^I , we obtain

$$\begin{aligned} \mathbb{E} [U(\mathbf{x}, t) U(\mathbf{y}, t)] &= \mathbb{E} [U^H(\mathbf{x}, t) U^H(\mathbf{y}, t)] + \mathbb{E} [U^I(\mathbf{x}, t) U^I(\mathbf{y}, t)] \\ &= \sum_{\ell=0}^{\infty} (2\ell + 1) \left(C_{\ell}(E_{\alpha}(-\lambda_{\ell} t^{\alpha}))^2 + \mathcal{A}_{\ell} \sigma_{\ell,t-\tau,\alpha}^2 \right) P_{\ell}(\mathbf{x} \cdot \mathbf{y}). \end{aligned}$$

As $P_{\ell}(\mathbf{x} \cdot \mathbf{y})$ depends only on the inner product of \mathbf{x} and \mathbf{y} , we conclude that the covariance function $\mathbb{E} [U^I(\mathbf{x}, t) U^I(\mathbf{y}, t)]$ is rotationally invariant. Since $t > \tau$, we note that for $\ell, \ell' \in \mathbb{N}_0$, $m = -\ell, \dots, \ell$ and $m' = -\ell', \dots, \ell'$, by Propositions 3.2 and 4.2,

$$\mathbb{E} \left[\widehat{U}_{\ell,m}(t) \overline{\widehat{U}_{\ell',m'}(t)} \right] = \left((E_{\alpha}(-\lambda_{\ell} t^{\alpha}))^2 C_{\ell} + \mathcal{A}_{\ell} \sigma_{\ell,t-\tau,\alpha}^2 \right) \delta_{\ell\ell'} \delta_{mm'}.$$

Now using the results of Propositions 4.2 and 3.2, we conclude that the field U is Gaussian, thus completing the proof.

Remark 5.1. The solution U given by (92), can be expressed, using (69) and (91), as

$$U(t) = \sum_{\ell=0}^{\infty} (U_{\ell}^H(t) + U_{\ell}^I(t)), \quad t \in (0, \infty),$$

which is convergent in $L_2(\Omega \times \mathbb{S}^2)$, where $U_{\ell}^H(t)$ is defined as

$$\begin{aligned} U_{\ell}^H(t) &= E_{\alpha}(-\lambda_{\ell} t^{\alpha}) \left(\sqrt{C_{\ell}} Z_{\ell,0}^{(1)} Y_{\ell,0} \right. \\ &\quad \left. + \sqrt{2C_{\ell}} \sum_{m=1}^{\ell} \left(Z_{\ell,m}^{(1)} \operatorname{Re} Y_{\ell,m} + Z_{\ell,m}^{(2)} \operatorname{Im} Y_{\ell,m} \right) \right), \end{aligned} \quad (94)$$

and $U_{\ell}^I(t) = 0$ for $t \leq \tau$, while for $t > \tau$,

$$\begin{aligned} U_{\ell}^I(t) &= \sqrt{A_{\ell}} \mathcal{I}_{\ell,0,\alpha}^{(1)}(t - \tau) Y_{\ell,0} \\ &\quad + \sqrt{2A_{\ell}} \sum_{m=1}^{\ell} \left(\mathcal{I}_{\ell,m,\alpha}^{(1)}(t - \tau) \operatorname{Re} Y_{\ell,m} + \mathcal{I}_{\ell,m,\alpha}^{(2)}(t - \tau) \operatorname{Im} Y_{\ell,m} \right), \end{aligned} \quad (95)$$

where $\mathcal{I}_{\ell,m,\alpha}^{(j)}(\cdot)$, $j = 1, 2$, are given in (29).

6. Approximation to the solution

The results in the previous sections give a series representation of the solution $U(t)$, $t \in (0, \infty)$, of the time-fractional diffusion equation (1). This section provides an approximation to the solution $U(t)$ by truncating its expansion at a degree (truncation level) $L \in \mathbb{N}$. Then we give an upper bound for the approximation error of U_L .

Definition 6.1. The approximation U_L of degree $L \in \mathbb{N}$ to the solution U given in (92) is defined as

$$\begin{aligned} U_L(t) &:= U_L^H(t) + U_L^I(t) \\ &= \sum_{\ell=0}^L (U_{\ell}^H(t) + U_{\ell}^I(t)), \quad t \in (0, \infty), \end{aligned} \quad (96)$$

where $U_{\ell}^H(t)$ is given by (94) and for $t \leq \tau$, $U_{\ell}^I(t) = 0$, while for $t > \tau$, $U_{\ell}^I(t)$ is given by (95).

Proposition 6.1. Let $\{C_{\ell} : \ell \in \mathbb{N}_0\}$, the angular power spectrum of ξ , satisfy (60). Let $Q_L^H(t)$, $t \in (0, \infty)$, be defined as

$$Q_L^H(t) := \left\| U^H(t) - U_L^H(t) \right\|_{L_2(\Omega \times \mathbb{S}^2)} = \left\| \sum_{\ell=L+1}^{\infty} U_{\ell}^H(t) \right\|_{L_2(\Omega \times \mathbb{S}^2)},$$

where U_{ℓ}^H is given in (94). Then, the following estimates hold true:

a) for $0 < t \leq \lambda_L^{-\frac{1}{\alpha}}$, $L \geq \ell_0$,

$$Q_L^H(t) \leq \tilde{C}_{\kappa_1} L^{-(\kappa_1-2)/2},$$

where \tilde{C}_{κ_1} is given by (61).

b) for $t > \lambda_L^{-\frac{1}{\alpha}}$, $L \geq \ell_0$,

$$Q_L^H(t) \leq \psi_{\alpha}^H(t) \tilde{C}_{\kappa_1} L^{-(2+\kappa_1)/2},$$

where $\psi_\alpha^H(\cdot)$ is defined as

$$\psi_\alpha^H(t) := \Gamma(1 + \alpha)t^{-\alpha} \quad \alpha \in (0, 1]. \quad (97)$$

PROOF. By (94) we can write

$$\begin{aligned} (Q_L^H(t))^2 &= \mathbb{E} \left[\left\| \sum_{\ell=L+1}^{\infty} \left[E_\alpha(-\lambda_\ell t^\alpha) \left(\sqrt{C_\ell} Z_{\ell,0}^{(1)} Y_{\ell,0} \right. \right. \right. \right. \\ &\quad \left. \left. \left. + \sqrt{2C_\ell} \sum_{m=1}^{\ell} \left(Z_{\ell,m}^{(1)} \operatorname{Re} Y_{\ell,m} + Z_{\ell,m}^{(2)} \operatorname{Im} Y_{\ell,m} \right) \right) \right] \right\|_{L_2(\mathbb{S}^2)}^2 \right]. \end{aligned}$$

Since $Z_{\ell,m_1}^{(1)}, Z_{\ell,m_2}^{(2)} \in \mathcal{Z}_\ell, \ell \in \mathbb{N}_0, m_1 = 0, 1, \dots, \ell, m_2 = 1, \dots, \ell$, we have

$$\begin{aligned} (Q_L^H(t))^2 &= \sum_{\ell=L+1}^{\infty} \left[(E_\alpha(-\lambda_\ell t^\alpha))^2 \left(C_\ell \mathbb{E} \left[(Z_{\ell,0}^{(1)})^2 \right] \|Y_{\ell,0}\|_{L_2(\mathbb{S}^2)}^2 \right. \right. \\ &\quad \left. \left. + 2C_\ell \sum_{m=1}^{\ell} \left(\mathbb{E} \left[(Z_{\ell,m}^{(1)})^2 \right] \|\operatorname{Re} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 + \mathbb{E} \left[(Z_{\ell,m}^{(2)})^2 \right] \|\operatorname{Im} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 \right) \right) \right] \\ &= \sum_{\ell=L+1}^{\infty} (E_\alpha(-\lambda_\ell t^\alpha))^2 \left[C_\ell \|Y_{\ell,0}\|_{L_2(\mathbb{S}^2)}^2 \right. \\ &\quad \left. + 2C_\ell \sum_{m=1}^{\ell} \left(\|\operatorname{Re} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 + \|\operatorname{Im} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 \right) \right]. \end{aligned}$$

By the properties of the spherical harmonics we have

$$\begin{aligned} \|Y_{\ell,0}\|_{L_2(\mathbb{S}^2)}^2 + 2 \sum_{m=1}^{\ell} \left(\|\operatorname{Re} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 + \|\operatorname{Im} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 \right) &= \sum_{m=-\ell}^{\ell} \|Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 \\ &= (2\ell + 1)P_\ell(1) = (2\ell + 1), \end{aligned} \quad (98)$$

and hence

$$(Q_L^H(t))^2 = \sum_{\ell=L+1}^{\infty} (E_\alpha(-\lambda_\ell t^\alpha))^2 (2\ell + 1)C_\ell. \quad (99)$$

Now, assume $0 < t \leq \lambda_L^{-\frac{1}{\alpha}}$ for $L \geq \ell_0$, then since $E_\alpha(-s^\alpha) \leq 1$, equation (99) can be bounded by

$$\begin{aligned} (Q_L^H(t))^2 &\leq \sum_{\ell=L+1}^{\infty} (2\ell + 1)C_\ell \\ &\leq \tilde{C} \sum_{\ell=L+1}^{\infty} (2\ell + 1)\ell^{-\kappa_1} \\ &\leq \tilde{C} \int_L^{\infty} (2x + 1)x^{-\kappa_1} dx \\ &= \tilde{C} \left(\frac{2}{\kappa_1 - 2} + \frac{L^{-1}}{\kappa_1 - 1} \right) L^{2-\kappa_1} \\ &\leq \tilde{C} \left(\frac{2}{\kappa_1 - 2} + \frac{1}{\kappa_1 - 1} \right) L^{2-\kappa_1}. \end{aligned} \quad (100)$$

Now, let us consider part b. Since $E_\alpha(-\lambda_\ell t^\alpha)$ is positive and decreasing for $t > 0$, then for $t > \lambda_L^{-\frac{1}{\alpha}}$, using (33), (99) can be bounded by

$$\begin{aligned} (Q_L^H(t))^2 &\leq (E_\alpha(-\lambda_L t^\alpha))^2 \sum_{\ell=L+1}^{\infty} (2\ell+1)C_\ell \\ &\leq (\Gamma(1+\alpha))^2 t^{-2\alpha} \lambda_L^{-2} \sum_{\ell=L+1}^{\infty} (2\ell+1)C_\ell \\ &\leq (\psi_\alpha^H(t))^2 \lambda_L^{-2} \tilde{C}_{\kappa_1}^2 L^{2-\kappa_1}, \end{aligned} \quad (101)$$

thus completing the proof.

In the next proposition, we derive an upper bound for the approximation errors of the truncated solution $U_L^I(t)$, $t > \tau$.

Let $\psi_\alpha^I(t)$, $t > 0$, be defined as

$$\psi_\alpha^I(t) := \begin{cases} \sqrt{1 + M_\alpha t^{1-2\alpha}}, & \alpha \in (0, \frac{1}{2}), \\ \sqrt{1 + M_\alpha}, & \alpha \in (\frac{1}{2}, 1], \\ K(t), & \alpha = \frac{1}{2}, \end{cases} \quad (102)$$

where M_α is given by (27) and

$$K(t) := \begin{cases} \sqrt{1 + M_{\frac{1}{2}}(2 + \ln(t))}, & t > 1, \\ \sqrt{1 + 2M_{\frac{1}{2}}}, & t \leq 1. \end{cases} \quad (103)$$

Proposition 6.2. Let $\{\mathcal{A}_\ell : \ell \in \mathbb{N}_0\}$, the angular power spectrum of W_τ , satisfy (84). Let $Q_L^I(t)$, $t > \tau$, be defined as

$$Q_L^I(t) := \left\| U^I(t) - U_L^I(t) \right\|_{L_2(\Omega \times \mathbb{S}^2)} = \left\| \sum_{\ell=L+1}^{\infty} U_\ell^I(t) \right\|_{L_2(\Omega \times \mathbb{S}^2)}, \quad (104)$$

where for $t > \tau$, $U_\ell^I(t)$ is given by (95). Then, the following estimates hold true:

i) for $\tau < t \leq \tau + \lambda_L^{-\frac{1}{\alpha}}$, $L \geq \ell_0$,

$$Q_L^I(t) \leq \tilde{A}_{\kappa_2} L^{-\frac{(\kappa_2 + \frac{2}{\alpha} - 2)}{2}},$$

where \tilde{A}_{κ_2} is given by (85).

ii) For $t > \tau + \lambda_L^{-\frac{1}{\alpha}}$, $L \geq \ell_0$,

$$Q_L^I(t) \leq \psi_\alpha^I(t - \tau) \tilde{A}_{\kappa_2} L^{-\frac{\gamma_\alpha(\kappa_2)}{2}},$$

where $\gamma_\alpha(\kappa_2)$ is given in (86) and $\psi_\alpha^I(\cdot)$ is given in (102).

PROOF. By (95) the term $(Q_L^I(t))^2$ can be written, for $t > \tau$, as

$$(Q_L^I(t))^2 = \left\| \sum_{\ell=L+1}^{\infty} U_\ell^I(t) \right\|_{L_2(\Omega \times \mathbb{S}^2)}^2 = \left\| \sum_{\ell=L+1}^{\infty} \left[\sqrt{\mathcal{A}_\ell} \mathcal{I}_{\ell,0,\alpha}^{(1)}(t - \tau) Y_{\ell,0} \right] \right\|_{L_2(\Omega \times \mathbb{S}^2)}^2$$

$$+ \sqrt{2\mathcal{A}_\ell} \sum_{m=1}^{\ell} \left(\mathcal{I}_{\ell,m,\alpha}^{(1)}(t-\tau) \operatorname{Re} Y_{\ell,m} + \mathcal{I}_{\ell,m,\alpha}^{(2)}(t-\tau) \operatorname{Im} Y_{\ell,m} \right) \Big\|_{L_2(\Omega \times \mathbb{S}^2)}^2.$$

By Proposition 2.5, for $\ell \in \mathbb{N}_0, m_1 = 0, \dots, \ell, m_2 = 1, \dots, \ell$, we know that $\{(\mathcal{I}_{\ell,m_1,\alpha}^{(1)}(\cdot), \mathcal{I}_{\ell,m_2,\alpha}^{(2)}(\cdot))\}$ is a sequence of independent, real-valued Gaussian random variables with mean zero and variances $\sigma_{\ell,t-\tau,\alpha}^2, t > \tau$. It follows that

$$\begin{aligned} (Q_L^I(t))^2 &= \sum_{\ell=L+1}^{\infty} \left[\left(\mathcal{A}_\ell \mathbb{E} \left[(\mathcal{I}_{\ell,0,\alpha}^{(1)}(t-\tau))^2 \right] \|Y_{\ell,0}\|_{L_2(\mathbb{S}^2)}^2 \right. \right. \\ &\quad + 2\mathcal{A}_\ell \sum_{m=1}^{\ell} \left(\mathbb{E} \left[(\mathcal{I}_{\ell,m,\alpha}^{(1)}(t-\tau))^2 \right] \|\operatorname{Re} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 \right. \\ &\quad \left. \left. + \mathbb{E} \left[(\mathcal{I}_{\ell,m,\alpha}^{(2)}(t-\tau))^2 \right] \|\operatorname{Im} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 \right) \right] \\ &= \sum_{\ell=L+1}^{\infty} \sigma_{\ell,t-\tau,\alpha}^2 \left[\mathcal{A}_\ell \|Y_{\ell,0}\|_{L_2(\mathbb{S}^2)}^2 \right. \\ &\quad \left. + 2\mathcal{A}_\ell \sum_{m=1}^{\ell} \left(\|\operatorname{Re} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 + \|\operatorname{Im} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 \right) \right] \\ &= \sum_{\ell=L+1}^{\infty} \mathcal{A}_\ell (2\ell + 1) \sigma_{\ell,t-\tau,\alpha}^2 \\ &\leq \sigma_{L,t-\tau,\alpha}^2 \sum_{\ell=L+1}^{\infty} \mathcal{A}_\ell (2\ell + 1), \end{aligned} \tag{105}$$

since by (28) $\sigma_{\ell,t,\alpha}^2$ is decreasing in ℓ .

Now let us consider part i. Since $0 < (t-\tau)\lambda_L^{\frac{1}{\alpha}} \leq 1$, then by (31), $\sigma_{L,t-\tau,\alpha}^2$ is bounded by $\lambda_L^{-\frac{1}{\alpha}}$. Similar to (100), the upper bound (105) can be bounded, for $\tau < t \leq \tau + \lambda_L^{-\frac{1}{\alpha}}, L \geq \ell_0$, by

$$(Q_L^I(t))^2 \leq \lambda_L^{-\frac{1}{\alpha}} \sum_{\ell=L+1}^{\infty} \mathcal{A}_\ell (2\ell + 1) \leq \tilde{A}_{\kappa_2}^2 L^{2-(\kappa_2 + \frac{2}{\alpha})},$$

which completes the proof of i.

To prove ii, note that by using the upper bound (31) with the estimates (34), (35), and (36), we obtain

$$\sigma_{L,t-\tau,\alpha}^2 \leq \begin{cases} \lambda_L^{-\frac{1}{\alpha}} + M_\alpha (t-\tau)^{1-2\alpha} \lambda_L^{-2}, & \alpha \in (0, \frac{1}{2}), \\ \lambda_L^{-\frac{1}{\alpha}} (1 + M_\alpha), & \alpha \in (\frac{1}{2}, 1], \\ \lambda_L^{-2} (1 + M_{\frac{1}{2}} \ln(\lambda_L^2 (t-\tau))), & \alpha = \frac{1}{2}, \end{cases} \tag{106}$$

where M_α is given in (27).

By using (105) with the estimate (106) and the properties of \mathcal{A}_ℓ given in (84), we get

$$\begin{aligned} (Q_L^I(t))^2 &\leq \tilde{A}_{L,t-\tau,\alpha}^2 \sum_{\ell=L+1}^{\infty} (2\ell + 1) \ell^{-\kappa_2} \\ &\leq (\psi_\alpha^I(t-\tau) \tilde{A}_{\kappa_2})^2 L^{-\kappa_\alpha(\kappa_2)}, \end{aligned} \tag{107}$$

where $\gamma_\alpha(\kappa_2)$ is given in (86) and $\psi_\alpha^I(\cdot)$ is given in (102), thus completing the proof.

By combining the results of Propositions 6.1 and 6.2, we present the following theorem.

Theorem 6.1. *Let $U(t)$, $t \in (0, \infty)$, be the solution (92) to the equation (1), and the conditions of Propositions 6.1 and 6.2 be satisfied. Let $\tilde{\kappa}_\alpha := \min(\kappa_1 - 2, \kappa_2 + \frac{2}{\alpha} - 2)$, $\hat{\kappa}_\alpha := \min(\kappa_1 + 2, \kappa_2 + \frac{2}{\alpha} - 2)$, and $\kappa_\alpha := \min(\kappa_1 + 2, \gamma_\alpha(\kappa_2))$, where $\gamma_\alpha(\kappa_2)$ is given in (86). Let $\mathcal{Q}_L(t)$, $t \in (0, \infty)$, be defined as*

$$\mathcal{Q}_L(t) := \left\| U(t) - U_L(t) \right\|_{L_2(\Omega \times \mathbb{S}^2)}, \quad L \geq 1, \quad (108)$$

where U_L is given by (96). Then, the following estimates hold true:

I) For $0 < t \leq \lambda_L^{-\frac{1}{\alpha}}$, $\tau \geq \lambda_L^{-\frac{1}{\alpha}}$, $L \geq \ell_0$,

$$\mathcal{Q}_L(t) \leq \tilde{C}_{\kappa_1} L^{-(\kappa_1 - 2)/2},$$

where \tilde{C}_{κ_1} is given by (61).

II) For $\lambda_L^{-\frac{1}{\alpha}} < t \leq \tau + \lambda_L^{-\frac{1}{\alpha}}$, $L \geq \ell_0$,

$$\mathcal{Q}_L(t) \leq \left((\psi_\alpha^H(t) \tilde{C}_{\kappa_1})^2 + \tilde{A}_{\kappa_2}^2 \right)^{\frac{1}{2}} L^{-\hat{\kappa}_\alpha/2},$$

where \tilde{A}_{κ_2} is given by (85).

III) For $t > \tau + \lambda_L^{-\frac{1}{\alpha}}$, $L \geq \ell_0$,

$$\mathcal{Q}_L(t) \leq \left((\psi_\alpha^H(t) \tilde{C}_{\kappa_1})^2 + (\psi_\alpha^I(t - \tau) \tilde{A}_{\kappa_2})^2 \right)^{\frac{1}{2}} L^{-\kappa_\alpha/2},$$

where $\psi_\alpha^H(\cdot)$ is given by (97) and $\psi_\alpha^I(\cdot)$ is given by (102).

Remark 6.1. *In view of Theorem 6.1 and the functions $\psi_\alpha^H(t)$ and $\psi_\alpha^I(t)$ given by (97) and (102) respectively, it is easy to see that for $\alpha \in (\frac{1}{2}, 1]$, both functions $\psi_\alpha^H(t)$ and $\psi_\alpha^I(t)$ are bounded when $t \rightarrow \infty$ and hence the truncation errors $\mathcal{Q}_L(t)$ are bounded when $t \rightarrow \infty$ (see case III). Also, it is worth noting that when $\alpha \in (\frac{1}{2}, 1]$, the truncation errors $\mathcal{Q}_L(t)$ as functions of t are bounded as well when $t \rightarrow 0$ (see case I), which improves the upper bound for the truncation errors obtained in [2] when $\alpha = 1$ and t is sufficiently small.*

7. Temporal increments and sample Hölder continuity of solution

In this section we study some properties of the stochastic solution of equation (1). In particular, we derive an upper bound, in L_2 -norm, of the temporal increments of the stochastic solution $U(t)$, $t \in (\tau, \infty)$, to equation (1). Then we study the sample properties of the stochastic solution U . Under some conditions, we show the existence of a locally Hölder continuous modification of the random field U . We demonstrate how the \mathbb{P} -almost sure Hölder continuity of the solution U depends on the decay of the angular power spectra of the initial condition ξ and the driving noise W_τ .

Theorem 7.1. *Let U be the solution (92) to the equation (1). Let J_h , $h > 0$, be defined as*

$$J_h(t) := \left\| U(t+h) - U(t) \right\|_{L_2(\Omega \times \mathbb{S}^2)}, \quad t > \tau. \quad (109)$$

Then, for $h > 0$, there holds

$$J_h(t) \leq q(t)h^{\frac{1}{2}},$$

where $q(t)$, $t > \tau$, is given by

$$q(t) = \sqrt{C \tilde{C}_{\kappa_1}^2 t^{-1} + (1+C) \tilde{A}_{\kappa_2}^2}, \quad C > 0,$$

\tilde{C}_{κ_1} , and \tilde{A}_{κ_2} are defined in (61) and (85) respectively.

PROOF. Let $t \in (\tau, \infty)$, then by the triangle inequality for $L_2(\Omega \times \mathbb{S}^2)$, we can write

$$(\mathcal{J}_h(t))^2 \leq \mathcal{J}_h^H(t) + \mathcal{J}_h^I(t),$$

where $\mathcal{J}_h^H(t)$ and $\mathcal{J}_h^I(t)$ are defined as

$$\mathcal{J}_h^H(t) := \left\| \sum_{\ell=0}^{\infty} (U_{\ell}^H(t+h) - U_{\ell}^H(t)) \right\|_{L_2(\Omega \times \mathbb{S}^2)}^2,$$

$$\mathcal{J}_h^I(t) := \left\| \sum_{\ell=0}^{\infty} (U_{\ell}^I(t+h) - U_{\ell}^I(t)) \right\|_{L_2(\Omega \times \mathbb{S}^2)}^2,$$

where U_{ℓ}^H and U_{ℓ}^I are given by (94) and (95) respectively.

Using (94), the term $\mathcal{J}_h^H(t)$ can be written as

$$\begin{aligned} \mathcal{J}_h^H(t) = \mathbb{E} \left[\left\| \sum_{\ell=0}^{\infty} \left[(E_{\alpha}(-\lambda_{\ell}(t+h)^{\alpha}) - E_{\alpha}(-\lambda_{\ell}t^{\alpha}))^2 \left(\sqrt{C_{\ell}} Z_{\ell,0}^{(1)} Y_{\ell,0} \right. \right. \right. \right. \\ \left. \left. \left. + \sqrt{2C_{\ell}} \sum_{m=1}^{\ell} \left(Z_{\ell,m}^{(1)} \operatorname{Re} Y_{\ell,m} + Z_{\ell,m}^{(2)} \operatorname{Im} Y_{\ell,m} \right) \right) \right] \right\|_{L_2(\mathbb{S}^2)}^2 \right]. \end{aligned}$$

Since $Z_{\ell,m_1}^{(1)}, Z_{\ell,m_2}^{(2)} \in \mathcal{Z}_{\ell}$, $\ell \in \mathbb{N}_0$, $m_1 = 0, \dots, \ell$, $m_2 = 1, 2, \dots, \ell$, there holds, using (98),

$$\begin{aligned} \mathcal{J}_h^H(t) &= \sum_{\ell=0}^{\infty} \left[\left| E_{\alpha}(-\lambda_{\ell}(t+h)^{\alpha}) - E_{\alpha}(-\lambda_{\ell}t^{\alpha}) \right|^2 \left(C_{\ell} \mathbb{E} \left[(Z_{\ell,0}^{(1)})^2 \right] \|Y_{\ell,0}\|_{L_2(\mathbb{S}^2)}^2 \right. \right. \\ &\quad \left. \left. + 2C_{\ell} \sum_{m=1}^{\ell} \left(\mathbb{E} \left[(Z_{\ell,m}^{(1)})^2 \right] \|\operatorname{Re} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 + \mathbb{E} \left[(Z_{\ell,m}^{(2)})^2 \right] \|\operatorname{Im} Y_{\ell,m}\|_{L_2(\mathbb{S}^2)}^2 \right) \right) \right] \\ &= \sum_{\ell=1}^{\infty} \left| E_{\alpha}(-\lambda_{\ell}(t+h)^{\alpha}) - E_{\alpha}(-\lambda_{\ell}t^{\alpha}) \right|^2 (2\ell + 1) C_{\ell}. \end{aligned} \quad (110)$$

Since $0 < E_{\alpha}(\cdot) \leq 1$, (110) becomes bounded by

$$\mathcal{J}_h^H(t) \leq \sum_{\ell=1}^{\infty} \left| E_{\alpha}(-\lambda_{\ell}(t+h)^{\alpha}) - E_{\alpha}(-\lambda_{\ell}t^{\alpha}) \right| (2\ell + 1) C_{\ell}. \quad (111)$$

Now using the mean-value theorem for the function $E_{\alpha}(-\lambda_{\ell}t^{\alpha})$ (see [13], equation 4.3.1), then (111) becomes

$$\begin{aligned} \mathcal{J}_h^H(t) &\leq h \sum_{\ell=1}^{\infty} \left| -\alpha \lambda_{\ell} t_2^{\alpha-1} E_{\alpha,\alpha+1}^2(-\lambda_{\ell}t_2^{\alpha}) \right| (2\ell + 1) C_{\ell} \\ &\leq h t^{\alpha-1} \sum_{\ell=1}^{\infty} \lambda_{\ell} \alpha E_{\alpha,\alpha+1}^2(-\lambda_{\ell}t^{\alpha}) (2\ell + 1) C_{\ell}, \end{aligned} \quad (112)$$

where $t_2 \in (t, t+h)$ and $E_{\alpha,\alpha+1}^2(\cdot)$ is the 3-parameter Mittag-Leffler function (see [13]) and the second step uses that $E_{\alpha,\alpha+1}^2(\cdot)$ is decreasing in t .

Using the relations (see [13], equation 5.1.14)

$$\alpha E_{\alpha,\beta}^2 = E_{\alpha,\beta-1} - (1 + \alpha - \beta) E_{\alpha,\beta}, \quad \alpha > 0, \beta > 1, \quad (113)$$

and (26) we estimate (112) as

$$\begin{aligned}
 \mathcal{J}_h^H(t) &\leq ht^{\alpha-1} \sum_{\ell=1}^{\infty} \lambda_{\ell} E_{\alpha,\alpha}(-\lambda_{\ell} t^{\alpha})(2\ell+1)C_{\ell} \\
 &\leq Cht^{-1} \sum_{\ell=1}^{\infty} (2\ell+1)C_{\ell} \left(\frac{\lambda_{\ell} t^{\alpha}}{1+\lambda_{\ell} t^{\alpha}} \right) \\
 &\leq Cht^{-1} \sum_{\ell=1}^{\infty} (2\ell+1)C_{\ell} \\
 &\leq Cht^{-1} \tilde{C}_{\kappa_1}^2.
 \end{aligned}$$

Now we treat the term $\mathcal{J}_h^I(t)$. By using (95) we have

$$\begin{aligned}
 \mathcal{J}_h^I(t) &= \sum_{\ell=0}^{\infty} \left[\left(\mathcal{A}_{\ell} \mathbb{E} \left[\left(\mathcal{I}_{\ell,0,\alpha}^{(1)}(t-\tau+h) - \mathcal{I}_{\ell,0,\alpha}^{(1)}(t-\tau) \right)^2 \right] \|Y_{\ell,0}\|_{L_2(\mathbb{S}^2)}^2 \right. \right. \\
 &\quad + 2\mathcal{A}_{\ell} \sum_{m=1}^{\ell} \left(\mathbb{E} \left[\left(\mathcal{I}_{\ell,m,\alpha}^{(1)}(t-\tau+h) - \mathcal{I}_{\ell,m,\alpha}^{(1)}(t-\tau) \right)^2 \right] \| \operatorname{Re} Y_{\ell,m} \|_{L_2(\mathbb{S}^2)}^2 \right. \\
 &\quad \left. \left. + \mathbb{E} \left[\left(\mathcal{I}_{\ell,m,\alpha}^{(2)}(t-\tau+h) - \mathcal{I}_{\ell,m,\alpha}^{(2)}(t-\tau) \right)^2 \right] \| \operatorname{Im} Y_{\ell,m} \|_{L_2(\mathbb{S}^2)}^2 \right) \right], \tag{114}
 \end{aligned}$$

since by Proposition 2.5, $\mathcal{I}_{\ell,m_1,\alpha}^{(1)}, \mathcal{I}_{\ell,m_2,\alpha}^{(2)}$, $m_1 = 0, 1, \dots, \ell$, $m_2 = 1, \dots, \ell$, are independent.

Let $\Xi_{\ell,m_1,\alpha}^{(1)}(h)$ and $\Xi_{\ell,m_2,\alpha}^{(2)}(h)$, $m_1 = 0, 1, \dots, \ell$, $m_2 = 1, \dots, \ell$, be defined as

$$\Xi_{\ell,m_1,\alpha}^{(1)}(h) := \mathbb{E} \left[\left(\mathcal{I}_{\ell,m_1,\alpha}^{(1)}(t-\tau+h) - \mathcal{I}_{\ell,m_1,\alpha}^{(1)}(t-\tau) \right)^2 \right]$$

and

$$\Xi_{\ell,m_2,\alpha}^{(2)}(h) := \mathbb{E} \left[\left(\mathcal{I}_{\ell,m_2,\alpha}^{(2)}(t-\tau+h) - \mathcal{I}_{\ell,m_2,\alpha}^{(2)}(t-\tau) \right)^2 \right].$$

Since by Proposition 2.5, $\beta_{\ell,m_1}^{(1)} \in \mathcal{B}_{\ell,0}$, $\ell \in \mathbb{N}_0$, $m_1 = 0, \dots, \ell$, there holds

$$\begin{aligned}
 \Xi_{\ell,m_1,\alpha}^{(1)}(h) &= \mathbb{E} \left[\left(\int_{t-\tau}^{t-\tau+h} E_{\alpha}(-\lambda_{\ell}(t-\tau+h-u)^{\alpha}) d\beta_{\ell,m_1}^{(1)}(u) \right. \right. \\
 &\quad \left. \left. + \int_0^{t-\tau} \left(E_{\alpha}(-\lambda_{\ell}(t+h-\tau-u)^{\alpha}) - E_{\alpha}(-\lambda_{\ell}(t-\tau-u)^{\alpha}) \right) d\beta_{\ell,m_1}^{(1)}(u) \right)^2 \right] \\
 &\leq \mathbb{E} \left[\left(\int_{t-\tau}^{t-\tau+h} E_{\alpha}(-\lambda_{\ell}(t-\tau+h-u)^{\alpha}) d\beta_{\ell,m_1}^{(1)}(u) \right)^2 \right] \\
 &\quad + \mathbb{E} \left[\left(\int_0^{t-\tau} \left(E_{\alpha}(-\lambda_{\ell}(t+h-\tau-u)^{\alpha}) - E_{\alpha}(-\lambda_{\ell}(t-\tau-u)^{\alpha}) \right) d\beta_{\ell,m_1}^{(1)}(u) \right)^2 \right] \\
 &\leq \int_{t-\tau}^{t-\tau+h} \left(E_{\alpha}(-\lambda_{\ell}(t-\tau+h-u)^{\alpha}) \right)^2 du \\
 &\quad + \int_0^{t-\tau} \left(E_{\alpha}(-\lambda_{\ell}(t+h-\tau-u)^{\alpha}) - E_{\alpha}(-\lambda_{\ell}(t-\tau-u)^{\alpha}) \right)^2 du, \tag{115}
 \end{aligned}$$

where the last step uses Itô's isometry (see [19]).

Since $0 < E_\alpha(\cdot) \leq 1$, we can write (115), using Remark 2.2, as

$$\begin{aligned} \Xi_{\ell, m_1, \alpha}^{(1)}(h) &\leq \sigma_{\ell, h, \alpha}^2 + \int_0^{t-\tau} \left| E_\alpha(-\lambda_\ell(t+h-\tau-u)^\alpha) - E_\alpha(-\lambda_\ell(t-\tau-u)^\alpha) \right| du \\ &\leq \sigma_{\ell, h, \alpha}^2 + h \int_0^{t-\tau} \left| -\alpha \lambda_\ell(t-\tau-u)^{\alpha-1} E_{\alpha, \alpha+1}^2(-\lambda_\ell(t-\tau-u)^\alpha) \right| du \\ &\leq \sigma_{\ell, h, \alpha}^2 + h \lambda_\ell \int_0^{t-\tau} \left| (t-\tau-u)^{\alpha-1} E_{\alpha, \alpha}(-\lambda_\ell(t-\tau-u)^\alpha) \right| du, \end{aligned} \quad (116)$$

where the second step uses the mean-value theorem for the function $E_\alpha(-\lambda_\ell(t-\tau)^\alpha)$ (see [13], equation 4.3.1) and the last step uses (113).

Using Remark 2.2 and relation (4.4.4) in [13], then (116) becomes

$$\begin{aligned} \Xi_{\ell, m_1, \alpha}^{(1)}(h) &\leq h + h \lambda_\ell (t-\tau)^\alpha E_{\alpha, \alpha}(-\lambda_\ell (t-\tau)^\alpha) \\ &\leq h + hC \frac{\lambda_\ell (t-\tau)^\alpha}{1 + \lambda_\ell (t-\tau)^\alpha} \\ &\leq h(1 + C), \end{aligned} \quad (117)$$

where the second step uses (26).

Similarly, we obtain, for $m_2 = 1, \dots, \ell$, that

$$\Xi_{\ell, m_2, \alpha}^{(2)}(h) \leq h(1 + C). \quad (118)$$

Using (117), (118), and (98) with (114), we get

$$\begin{aligned} J_h^I(t) &\leq \sum_{\ell=0}^{\infty} h(1 + C)(2\ell + 1) \mathcal{A}_\ell \\ &\leq h(1 + C) \sum_{\ell=0}^{\infty} (2\ell + 1) \mathcal{A}_\ell \leq h(1 + C) \tilde{A}_{\kappa_2}^2. \end{aligned}$$

Thus the result follows by combining the upper bounds $J_h^H(\cdot)$ and $J_h^I(\cdot)$.

Now we demonstrate continuity properties of the stochastic solution U at a given time t with respect to the geodesic distance on \mathbb{S}^2 , i.e. $\tilde{d}(\mathbf{x}, \mathbf{y})$, $\mathbf{x}, \mathbf{y} \in \mathbb{S}^2$.

Notice that by Proposition 5.1, the stochastic solution U is known to be mean square continuous (see [23]). However, to obtain sample Hölder continuity of the stochastic solution U we need stronger assumptions on the angular power spectra \mathcal{C}_ℓ and \mathcal{A}_ℓ , $\ell \in \mathbb{N}_0$.

Assumption 7.1. Assume that the angular power spectra $\{\mathcal{C}_\ell, \ell \in \mathbb{N}_0\}$ and $\{\mathcal{A}_\ell, \ell \in \mathbb{N}_0\}$ satisfy (60) and (84) with $\kappa_1, \kappa_2 > 2(1 + \beta^*)$ for some $\beta^* \in (0, 1]$.

It follows from Assumption 7.1 that for $\beta^* \in (0, 1]$ there exist constants $K_{\beta^*}^{(1)}, K_{\beta^*}^{(2)} > 0$ such that

$$K_{\beta^*}^{(1)} := \sum_{\ell=1}^{\infty} \ell^{1+2\beta^*} \mathcal{C}_\ell < \infty \text{ and } K_{\beta^*}^{(2)} := \sum_{\ell=1}^{\infty} \ell^{1+2\beta^*} \mathcal{A}_\ell < \infty.$$

The following theorem provides bounds on moments of the variance $\text{Var}[U(\mathbf{x}, t) - U(\mathbf{y}, t)]$, for $\mathbf{x}, \mathbf{y} \in \mathbb{S}^2$ in terms of the geodesic distance $\tilde{d}(\mathbf{x}, \mathbf{y})$.

Theorem 7.2. Let U , $t \in (0, \infty)$, be the solution given by (92) to the equation (1), and the angular power spectra \mathcal{C}_ℓ and \mathcal{A}_ℓ satisfy the Assumption 7.1. Then there exists a constant $K_{\beta^*} > 0$ such that

$$\text{Var}[U(\mathbf{x}, t) - U(\mathbf{y}, t)] \leq K_{\beta^*} (\tilde{d}(\mathbf{x}, \mathbf{y}))^{2\beta^*}, \quad \beta^* \in (0, 1].$$

PROOF. Let $t \leq \tau$. Then by (92) and (93) there holds

$$\begin{aligned} \mathbb{E}[U(\mathbf{x}, t)U(\mathbf{y}, t)] &= \mathbb{E}[U^H(\mathbf{x}, t)U^H(\mathbf{y}, t)] \\ &= \sum_{\ell=0}^{\infty} (2\ell + 1)C_{\ell}(E_{\alpha}(-\lambda_{\ell}t^{\alpha}))^2 P_{\ell}(\mathbf{x} \cdot \mathbf{y}), \end{aligned}$$

where $P_{\ell}(\cdot)$, $\ell \in \mathbb{N}_0$, is the Legendre polynomial of degree ℓ .

Thus for $t \leq \tau$ there holds

$$\begin{aligned} \text{Var}[U(\mathbf{x}, t) - U(\mathbf{y}, t)] &= \text{Var}[U^H(\mathbf{x}, t)] + \text{Var}[U^H(\mathbf{y}, t)] - 2\mathbb{E}[U^H(\mathbf{x}, t)U^H(\mathbf{y}, t)] \\ &= 2(\text{Var}[U^H(\mathbf{x}, t)] - \mathbb{E}[U^H(\mathbf{x}, t)U^H(\mathbf{y}, t)]) \\ &= 2 \sum_{\ell=0}^{\infty} (2\ell + 1)C_{\ell}(E_{\alpha}(-\lambda_{\ell}t^{\alpha}))^2 (1 - P_{\ell}(\mathbf{x} \cdot \mathbf{y})) \\ &\leq 2 \sum_{\ell=0}^{\infty} (2\ell + 1)C_{\ell} |1 - P_{\ell}(\cos \tilde{d}(\mathbf{x}, \mathbf{y}))|, \end{aligned} \quad (119)$$

where the last step uses that $0 < E_{\alpha}(-\lambda_{\ell}t^{\alpha}) \leq 1$.

Since $P_{\ell}(1) = 1$ for all $\ell \in \mathbb{N}_0$ and $P'_{\ell}(x) \leq P'_{\ell}(1)$ for $x \in [-1, 1]$, we obtain

$$|1 - P_{\ell}(x)| = \left| \int_x^1 P'_{\ell}(y) dy \right| \leq |1 - x| \left(\frac{\ell(\ell + 1)}{2} \right), \quad x \in [-1, 1]. \quad (120)$$

Also, we observe that

$$|1 - P_{\ell}(x)| \leq 2, \quad x \in [-1, 1]. \quad (121)$$

Then, for some $\beta^* \in (0, 1]$ we can write, using (120) and (121),

$$\begin{aligned} |1 - P_{\ell}(x)| &= |1 - P_{\ell}(x)|^{\beta^*} |1 - P_{\ell}(x)|^{1-\beta^*} \\ &\leq \left(|1 - x| \frac{\ell(\ell + 1)}{2} \right)^{\beta^*} 2^{1-\beta^*} \\ &\leq 2^{1-2\beta^*} |1 - x|^{\beta^*} (\ell(\ell + 1))^{\beta^*}. \end{aligned} \quad (122)$$

Thus (119) becomes bounded, using (122), by

$$\begin{aligned} \text{Var}[U(\mathbf{x}, t) - U(\mathbf{y}, t)] &\leq 2^{2-2\beta^*} |1 - \cos \tilde{d}(\mathbf{x}, \mathbf{y})|^{\beta^*} \sum_{\ell=0}^{\infty} (2\ell + 1)C_{\ell} \lambda_{\ell}^{\beta^*} \\ &\leq 2^{2-2\beta^*} (\tilde{d}(\mathbf{x}, \mathbf{y}))^{2\beta^*} \sum_{\ell=1}^{\infty} (2\ell + 1)C_{\ell} (\ell(\ell + 1))^{\beta^*} \\ &\leq 2^{4-\beta^*} (\tilde{d}(\mathbf{x}, \mathbf{y}))^{2\beta^*} K_{\beta^*}^{(1)}, \end{aligned}$$

where the second step uses the inequality

$$\begin{aligned} |1 - \cos r| &= \left| \int_0^r \sin x dx \right| \leq r \sin r \\ &\leq r \left| \int_0^r \cos x dx \right| \leq r^2. \end{aligned}$$

Similarly, for $t > \tau$, we have

$$\mathbb{E}[U(\mathbf{x}, t)U(\mathbf{y}, t)] = \sum_{\ell=0}^{\infty} (2\ell + 1) \left(C_{\ell}(E_{\alpha}(-\lambda_{\ell}t^{\alpha}))^2 + \mathcal{A}_{\ell} \sigma_{\ell, t-\tau, \alpha}^2 \right) P_{\ell}(\mathbf{x} \cdot \mathbf{y})$$

and by Assumption 7.1 and (122) we obtain

$$\begin{aligned} \text{Var}[U(\mathbf{x}, t) - U(\mathbf{y}, t)] &= 2 \sum_{\ell=0}^{\infty} (2\ell + 1) \left(C_{\ell} (E_{\alpha}(-\lambda_{\ell} t^{\alpha}))^2 + \mathcal{A}_{\ell} \sigma_{\ell, t-\tau, \alpha}^2 \right) (1 - P_{\ell}(\mathbf{x} \cdot \mathbf{y})) \\ &\leq 2 \sum_{\ell=0}^{\infty} (2\ell + 1) (C_{\ell} + (t - \tau) \mathcal{A}_{\ell}) (1 - P_{\ell}(\mathbf{x} \cdot \mathbf{y})) \\ &\leq 2^{4-\beta^*} (\tilde{d}(\mathbf{x}, \mathbf{y}))^{2\beta^*} (K_{\beta^*}^{(1)} + (t - \tau) K_{\beta^*}^{(2)}), \end{aligned}$$

where the second step uses Remark 2.2.

Thus for $t > 0$, there exists a constant K_{β^*} such that

$$\text{Var}[U(\mathbf{x}, t) - U(\mathbf{y}, t)] \leq K_{\beta^*} (\tilde{d}(\mathbf{x}, \mathbf{y}))^{2\beta^*},$$

where

$$K_{\beta^*} := 2^{4-\beta^*} \begin{cases} K_{\beta^*}^{(1)}, & t \leq \tau, \\ K_{\beta^*}^{(1)} + (t - \tau) K_{\beta^*}^{(2)}, & t > \tau, \end{cases}$$

which completes the proof.

Under the assumptions of Theorem 7.2, applying Kolmogorov's continuity criterion (see Corollary 4.5 in [20]), for any fixed $t > 0$, we deduce the existence of a locally Hölder continuous version (a modification) for the random field U of order $\gamma^* \in (0, \beta^*)$.

8. Numerical studies

In this section, we present some numerical examples for the solution U of equation (1). In particular, we show the evolution of the stochastic solution U of equation (1) using simulated data inspired by the CMB map. Also, we explain the convergence rates of the truncation errors of the approximations $U_L(t)$, $t \in (0, \infty)$, and the L_2 -norm of the temporal increments of the stochastic solution $U(t)$, $t \in (\tau, \infty)$, to the equation (1).

8.1. Evolution of the solution

In this subsection, we illustrate the evolution of the stochastic solution U of the equation (1) using simulated data inspired by the CMB map.

First, in Figure 1 we display a realization of the truncated initial condition at time $t = 0$ of degree $L = 600$ (i.e. $\xi = U_{600}(0)$) whose angular power spectrum C_{ℓ} has the form given by (60) with $\tilde{D} = \tilde{C} = 1$, $\kappa_1 = 2.3$. The Python HEALPy package (see [14]) was used to generate such a realization. Then, we use the obtained initial realization $U_{600}(0)$ to generate two realizations for truncated solution $U_{600}(t)$ given by (96) at different times, namely $t = \tau$ and $t = 10\tau$ with $\tau = 10^{-5}$ for the case $\alpha = 0.5$, $\kappa_1 = 2.3$, and $\kappa_2 = 2.5$. We use the equation (84) with $\tilde{K} = \tilde{A} = 10^4$ to compute the angular power spectrum \mathcal{A}_{ℓ} for the time-delayed Brownian motion W_{τ} . The Fourier coefficients $\hat{U}_{\ell, m}(t)$ for the solution $U_L(t)$ are computed, using (93) with (68) and (90), by

$$V_{\ell, 0}(t) := \begin{cases} E_{\alpha}(-\lambda_{\ell} t^{\alpha}) \sqrt{C_{\ell}} Z_{\ell, 0}^{(1)}, & t \leq \tau, \\ E_{\alpha}(-\lambda_{\ell} t^{\alpha}) \sqrt{C_{\ell}} Z_{\ell, 0}^{(1)} + \sqrt{\mathcal{A}_{\ell}} \mathcal{I}_{\ell, 0, \alpha}^{(1)}(t - \tau), & t > \tau, \end{cases} \quad (123)$$

and for $m = 1, \dots, \ell$,

$$V_{\ell, m}(t) := \begin{cases} E_{\alpha}(-\lambda_{\ell} t^{\alpha}) \sqrt{C_{\ell}/2} Z_{\ell, m}, & t \leq \tau, \\ E_{\alpha}(-\lambda_{\ell} t^{\alpha}) \sqrt{C_{\ell}/2} Z_{\ell, m} + \sqrt{\mathcal{A}_{\ell}/2} \mathcal{I}_{\ell, m, \alpha}(t - \tau), & t > \tau, \end{cases} \quad (124)$$

where $Z_{\ell, m} := Z_{\ell, m}^{(1)} - iZ_{\ell, m}^{(2)}$ and $\mathcal{I}_{\ell, m, \alpha}(t) := \mathcal{I}_{\ell, m, \alpha}^{(1)}(t) - i\mathcal{I}_{\ell, m, \alpha}^{(2)}(t)$, $t > 0$.




Figure 1: A realization of the truncated initial condition $U_{600}(0)$ with $L = 600$, $\alpha = 0.5$ and $\kappa_1 = 2.30$.

Note that for each realization and given time $t > \tau$, we simulate the stochastic integrals $\mathcal{I}_{\ell,m,\alpha}^{(j)}(t - \tau)$, $j = 1, 2$, using Proposition 2.5, as independent, normally distributed random variables with mean zero and variances $\sigma_{\ell,t-\tau,\alpha}^2$ where $\sigma_{\ell,t,\alpha}^2$ is given by (28). Figures 2a and 2b, respectively, show two realizations of the truncated solutions $U_L(t)$ and $U_L^H(t)$ with $L = 600$, using the initial realization obtained in Figure 1, at times $t = \tau$ and $t = 10\tau$ with $\tau = 10^{-5}$. Figure 3a shows a realization of the inhomogeneous solution $U_L^I(t)$ at time $t = 10\tau$ with $\tau = 10^{-5}$ and $\alpha = 0.5$, while Figure 3b shows a realization of the combined solution $U_L(t)$ using the realizations obtained in Figures 2b and 3a.




Figure 2: Truncated homogeneous solutions $U_{600}(\tau) = U_{600}^H(\tau)$ and $U_{600}^H(10\tau)$, using the initial realization in Figure 1, with $\alpha = 0.5$, $\kappa_1 = 2.3$, and $\kappa_2 = 2.5$.

The Figures 2a and 2b reflect the time evolution of the truncated homogeneous stochastic solution $U_{600}^H(t)$ at time $t = \tau$ and $t = 10\tau$ respectively. In this case, the randomness comes only from the initial condition. Figure 3a reflects the time evolution of the truncated inhomogeneous stochastic solution $U_{600}^I(t)$ at time $t = 10\tau$ (in this case the randomness comes only from W_τ). Figure 3b reflects the evolution of the same realization obtained in Figure 2b at $t = 10\tau$ but for which the noise W_τ is switched on (in this case the randomness comes from both the initial condition and W_τ , i.e. $U_{600}(10\tau) = U_{600}^H(10\tau) + U_{600}^I(10\tau)$).

8.2. Simulations of truncation errors of solution

This subsection presents some numerical examples for the truncation errors of the combined stochastic solution U of the equation (1). In particular, the numerical examples explain the convergence rates of the truncation errors $Q_L(t)$, given by (108), of the stochastic solution of (1).







Figure 3: (a) The truncated inhomogeneous solution $U_L^I(10\tau)$ with $\tau = 10^{-5}$, $\tilde{A} = 10^4$, $\alpha = 0.5$ and $\kappa_2 = 2.5$, (b) the truncated combined solution $U_{600}(10\tau)$, using the truncated homogeneous and inhomogeneous solutions in Figures 2b and 3a.

To produce numerical results, we use the angular power spectra \mathcal{A}_ℓ and \mathcal{C}_ℓ given by (60) and (84) respectively with $\tilde{A} = \tilde{K} = 10^4$, $\tilde{C} = \tilde{D} = 1$, $\kappa_1 = 2.3$, and $\kappa_2 = 2.5$. Also, we use $U_{\tilde{L}}$ with $\tilde{L} = 1500$ as a substitution of the solution U to the equation (1) which is given by (92). Then the (squared) mean L_2 -errors are computed, using Parseval's identity, by

$$\begin{aligned}
 (Q_{L,\tilde{L}}(t))^2 &:= \left\| U_{\tilde{L}}(t) - U_L(t) \right\|_{L_2(\Omega \times \mathbb{S}^2)}^2 \\
 &= \mathbb{E} \left[\left\| U_{\tilde{L}}(t) - U_L(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] \\
 &= \mathbb{E} \left[\left\| \sum_{\ell=L+1}^{\tilde{L}} \sum_{m=-\ell}^{\ell} \hat{U}_{\ell,m}(\omega, t) Y_{\ell,m} \right\|_{L_2(\mathbb{S}^2)}^2 \right] \\
 &\approx \frac{1}{\hat{N}} \sum_{j=1}^{\hat{N}} \sum_{\ell=L+1}^{\tilde{L}} \sum_{m=-\ell}^{\ell} \left| \hat{U}_{\ell,m}(\omega_j, t) \right|^2 \\
 &= \frac{1}{\hat{N}} \sum_{j=1}^{\hat{N}} \sum_{\ell=L+1}^{\tilde{L}} \sum_{m=0}^{\ell} \left| V_{\ell,m}(\omega_j, t) \right|^2, \tag{125}
 \end{aligned}$$

where the fourth line approximates the expectation by the mean of $\hat{N} = 100$ realizations, and $\{V_{\ell,m} : \ell = L+1, \dots, \tilde{L}; m = 0, \dots, \ell\}$ are computed, using (123) and (124).

To illustrate the results of Theorem 6.1, we consider two cases, namely $\alpha = 0.5$ and $\alpha = 0.75$. Then we compute the root mean L_2 -errors using equation (125) with degree up to $L = 800$ for different values of t . Figure 4 shows numerical errors $Q_{L,\tilde{L}}(t)$ of $\hat{N} = 100$ realizations and the corresponding theoretical errors for different values of t and τ with $\kappa_1 = 2.3$ and $\kappa_2 = 2.5$ for the case $\alpha = 0.5$. In particular, in Figure 4a we illustrate the case I ($0 < t \leq \lambda_L^{-2}$) in Theorem 6.1 with $t = 10^{-12}$. In Figure 4b we illustrate the case II with $t = \tau + \lambda_L^{-2}$ for $\tau = 10^{-5}$, while Figure 4c illustrates the case III with $t = 10\tau$ when $\tau = 10^{-5}$. Similarly, we illustrate the results of Theorem 6.1 using $\alpha = 0.75$. The corresponding results are displayed in Figures 5a-5c. The blue points in each picture in Figures 4a-4c and 5a-5c show the (sample) mean square approximation errors of $\hat{N} = 100$ realizations of $Q_{L,\tilde{L}}(t)$. The red line in each figure shows the corresponding theoretical approximation upper bound. The numerical results show that the convergence rates of the mean L_2 -error of U_L are consistent with the corresponding theoretical results in Theorem 6.1.


 (a) case I with $t = 10^{-12}$

 (b) case II with $t = \tau + \lambda_{800}^{-2}$, $\tau = 10^{-5}$

 (c) case III with $t = 10\tau$, $\tau = 10^{-5}$
Figure 4: Numerical errors for the combined solution $U(t)$ with $\kappa_1 = 2.3$, $\kappa_2 = 2.5$, $\alpha = 0.5$.

8.3. Simulations of temporal increments




This subsection presents a numerical example for the stochastic solution U of the equation (1) in time. In particular, the numerical example explains the L_2 -norm temporal increments $\mathcal{J}_h(t)$, given by (109), of equation (1).

To produce numerical results, we use the angular power spectra given by (60) and (84). Then, we use the truncated version of the L_2 -norm temporal increments of the solution $U(t)$ for $t \in (\tau, \infty)$ which can be estimated as


$$\begin{aligned}
 (\mathcal{J}_{h,L}(t))^2 &:= \left\| U_L(t+h) - U_L(t) \right\|_{L_2(\Omega \times \mathbb{S}^2)}^2 \\
 &= \mathbb{E} \left[\left\| U_L(t+h) - U_L(t) \right\|_{L_2(\mathbb{S}^2)}^2 \right] \\
 &= \mathbb{E} \left[\left\| \sum_{\ell=0}^{\tilde{L}} \sum_{m=-\ell}^{\ell} (\hat{U}_{\ell,m}(\omega, t+h) - \hat{U}_{\ell,m}(\omega, t)) Y_{\ell,m} \right\|_{L_2(\mathbb{S}^2)}^2 \right] \\
 &\approx \frac{1}{\hat{N}} \sum_{j=1}^{\hat{N}} \sum_{\ell=0}^{\tilde{L}} \sum_{m=0}^{\ell} \left| (V_{\ell,m}(\omega_j, t+h) - V_{\ell,m}(\omega_j, t)) \right|^2, \tag{126}
 \end{aligned}$$

where $\{V_{\ell,m} : \ell = 0, \dots, \tilde{L}; m = 0, \dots, \ell\}$ are computed using (123) and (124).

To illustrate the results of Theorem 7.1, we use $U_{\tilde{L}}$ with $\tilde{L} = 1500$ as a substitution of the solution U to the equation (1) which is given by (92). Then, the L_2 -norm temporal increments $\mathcal{J}_{h,L}(t)$ are computed using the estimate (126) with $\alpha = 0.5$, $t = \tau + \delta$, with $\tau = 10^{-5}$ and $\delta = 10^{-6}$. Figure 6 shows numerical temporal increments


 (a) case I with $t = 10^{-12}$

 (b) case II with $t = \tau + \lambda_{800}^{-2}$, $\tau = 10^{-5}$

 (c) case III with $t = 10\tau$, $\tau = 10^{-5}$
Figure 5: Numerical errors for the combined solution $U(t)$ with $\kappa_1 = 2.3$, $\kappa_2 = 2.5$, $\alpha = 0.75$.

$\mathcal{J}_{h,L}(t)$ of $\hat{N} = 100$ realizations and the corresponding theoretical upper bound for the temporal increments with $t = \tau + \delta$, $\alpha = 0.5$, $\kappa_1 = 2.3$, and $\kappa_2 = 2.5$, and time increment h ranging from δ to 11δ . The blue points in Figure 6


Figure 6: Numerical values of $\mathcal{J}_{h,L}(t) = \|U_L(t+h) - U_L(t)\|_{L_2(\Omega \times \mathbb{S}^2)}$ with $L = 1500$ and $t = \tau + \delta$, $\tau = 10^{-5}$, $\delta = 10^{-6}$ and $h = k\delta$, $k = 1, \dots, 11$.

show the (sample) mean square approximation errors of $\hat{N} = 100$ realizations of \mathcal{J}_h . The red straight line shows the theoretical approximation upper bound with time increment h . The numerical results show that the convergence rates

of the L_2 -norm temporal increments of the solution U of the equation (1) are consistent with the theoretical results in Theorem 7.1.

Acknowledgements

This research was supported under the Australian Research Council's Discovery Project funding scheme (Discovery Project number DP180100506). This research includes extensive computations using the Linux computational cluster Katana [17] supported by the Faculty of Science, The University of New South Wales, Sydney. The authors are also grateful to Bill McLean for discussions on some properties of the Mittag-Leffler function and to an anonymous referee's comments to improve the presentation.

Declarations

- All authors declare that they have no conflicts of interest.
- Availability of data and code: The Python code used for numerical simulations in the paper can be found at <https://github.com/qlegia/Stochastic-diffusion-on-sphere>.

References

- [1] V. V. Anh, A. Olenko, and G. Y. Wang. Fractional stochastic partial differential equation for random tangent fields on the sphere. *Theor. Probability and Math. Statist.*, 104:3-22, 2021.
- [2] V. V. Anh, P. Broadbridge, A. Olenko, and G. Y. Wang. On approximation for fractional stochastic partial differential equations on the sphere. *Stoch Environ Res Risk Assess.*, 32:2585–2603, 2018.
- [3] H. Berens, P.L. Butzer and S.Pawelke. Limitierungsverfahren von Reihen mehrdimensionaler Kugelfunktionen und deren Saturationsverhalten, Publ. RIMS, Kyoto Univ. Ser. A, Vol. 4 (1968), pp. 201-268.
- [4] P. Broadbridge, A. Kolesnik, N. Leonenko, and A. Olenko. Random spherical hyperbolic diffusion. *J. Stat. Phys.*, 177(5):889–916, 2019.
- [5] S. Castruccio and M. L. Stein. Global space-time models for climate ensembles. *Ann Appl Stat.*, 7(3):1593-1611, 2013.
- [6] D., Cohen, and A. Lang. Numerical approximation and simulation of the stochastic wave equation on the sphere. *Calcolo*, 59(3), 2022.
- [7] G. Da Prato and J. Zabczyk. *Stochastic Equations in Infinite Dimensions. Encyclopedia of Mathematics and Its Applications*, vol. 152. Cambridge University Press, Cambridge, 2014.
- [8] F. Dai and Y. Xu. *Approximation Theory and Harmonic Analysis on Spheres and Balls*. Springer, New York, 2013.
- [9] S. Dodelson. *Modern Cosmology*. Academic Press, New York, 2003.
- [10] R. Durrett. *Probability: Theory and Examples, volume 31 of Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2010.
- [11] P. Dyke. *An Introduction to Laplace Transforms and Fourier Series*. Springer, New York, 2014.
- [12] Q. L. Gia, I. H. Sloan, R. S. Womersley, and Y. G. Wang. Sparse isotropic regularization for spherical harmonic representations of random fields on the sphere. *Appl. Comput. Harmon. Anal.*, 49(1):257–278, 2020.
- [13] R. Gorenflo, A. A. Kilbas, F. Mainardi, and S. V. Rogosin. *Mittag-Leffler Functions: Related Topics and Applications*. Springer, Berlin, 2014.
- [14] K. M. Górski, E. Hivon, A. J. Banday, B. D. Wandelt, F. K. Hansen, M. Reinecke, and M. Bartelmann. HEALPix: a framework for high-resolution discretization and fast analysis of data distributed on the sphere. *Astrophys. J.*, 622(2):759–771, 2005.
- [15] M. Hess. *The stochastic Leibniz formula for Volterra integrals under enlarged filtrations*. *Stoch. Model.*, 39(4), 823–850.
- [16] R. Hachemi and B. Øksendal. *The fractional stochastic heat equation driven by time-space white noise*. *Fract. Calc. Appl. Anal.*, 26:513–532, 2023.
- [17] Katana. Published online 2010. doi:10.26190/669X-A286
- [18] Y. Kazashi and Q. L. Gia. A non-uniform discretization of stochastic heat equations with multiplicative noise on the unit sphere. *J. Complex.*, 50:43–65, 2019.
- [19] H-H. Kuo. *Introduction to Stochastic Integration*. Springer, New York, 2006.
- [20] A. Lang, J. Potthoff, M. Schlather, and D. Schwab. Continuity of random fields on Riemannian manifolds. *Commun. Stoch. Anal.*, 10(2):185–193, 2016.
- [21] A. Lang and C. Schwab. Isotropic Gaussian random field sphere: regularity, fast simulation and stochastic partial differential equations. *Ann. Appl. Probab.*, 25(6):3047–3094, 2015.
- [22] N. Leonenko and J. Vaz Jr. Spectral analysis of fractional hyperbolic diffusion equations with random Data. *J. Stat. Phys.*, 179:155–175, 2020.
- [23] D. Marinucci and G. Peccati. *Mean-square continuity on homogeneous spaces of compact groups*. *Electron. Commun. Probab.*, 18:1–10, 2013.
- [24] D. Marinucci and G. Peccati. *Random Fields on the Sphere: Representation, Limit Theorems and Cosmological Applications*. Cambridge University Press, Cambridge, 2011.
- [25] W. McLean and V. Thomée. Numerical solution via Laplace transforms of a fractional order evolution equation. *J. Integr. Equ. Appl.*, 22(1):57–94, 2010.
- [26] C. Müller. *Spherical Harmonics*. Springer, Berlin, New York, 1966.

- [27] B. Øksendal. *Stochastic Differential Equations. An Introduction with Applications*, 5th edn. Springer-Verlag Heidelberg, New York, 2023.
- [28] Planck Collaboration and Adam, R. et al. Planck 2015 results - I. Overview of products and scientific results. *Astron. Astrophys.*, 594:A1, 2016.
- [29] Planck Collaboration and Adam, R. et al. Planck 2015 results - IX. Diffuse component separation: CMB maps. *Astron. Astrophys.*, 594:A9, 2016.
- [30] Planck Collaboration and Adam, R. et al. Planck 2015 results - XVI. Isotropy and statistics of the CMB. *Astron. Astrophys.*, 594:A16, 2016.
- [31] I. Podlubny. *Fractional Differential Equations*. Academic Press, Inc., San Diego, CA, 1999.
- [32] W. Rudin. *Principles of Mathematical Analysis*. International Series in Pure and Applied Mathematics, McGraw-Hill, 1976.
- [33] T. Simon. Comparing Fréchet and positive stable laws. *Electron. J. Probab.* 19:1–25, 2014.