A CLT FOR THE LSS OF LARGE DIMENSIONAL SAMPLE COVARIANCE MATRICES WITH DIVERGING SPIKES

BY ZHIJUN LIU^{*}, JIANG HU[†], ZHIDONG BAI[‡], AND HAIYAN SONG[§]

KLASMOE and School of Mathematics and Statistics, Northeast Normal University, China. *liuzj037@nenu.edu.cn; †huj156@nenu.edu.cn; ‡baizd@nenu.edu.cn; §songhy716@nenu.edu.cn

> In this paper, we establish the central limit theorem (CLT) for linear spectral statistics (LSSs) of a large-dimensional sample covariance matrix when the population covariance matrices are involved with diverging spikes. This constitutes a nontrivial extension of the Bai-Silverstein theorem (BST) (Ann Probab 32(1):553-605, 2004), a theorem that has strongly influenced the development of high-dimensional statistics, especially in the applications of random matrix theory to statistics. Recently, there has been a growing realization that the assumption of uniform boundedness of the population covariance matrices in the BST is not satisfied in some fields, such as economics, where the variances of principal components may diverge as the dimension tends to infinity. Therefore, in this paper, we aim to eliminate this obstacle to applications of the BST. Our new CLT accommodates spiked eigenvalues, which may either be bounded or tend to infinity. A distinguishing feature of our result is that the variance in the new CLT is related to both spiked eigenvalues and bulk eigenvalues, with dominance being determined by the divergence rate of the largest spiked eigenvalues. The new CLT for LSS is then applied to test the hypothesis that the population covariance matrix is the identity matrix or a generalized spiked model. The asymptotic distributions of the corrected likelihood ratio test statistic and the corrected Nagao's trace test statistic are derived under the alternative hypothesis. Moreover, we present power comparisons between these two LSSs and Roy's largest root test. In particular, we demonstrate that except for the case in which the number of spikes is equal to one, the LSSs could exhibit higher asymptotic power than Roy's largest root test.

1. Introduction. We consider the general sample covariance matrix $\mathbf{B}_n = \frac{1}{n} \mathbf{T}_p \mathbf{X}_n \mathbf{X}_n^* \mathbf{T}_p^*$, where \mathbf{X}_n is a $p \times n$ matrix with independent and identically distributed (i.i.d.) standardized entries $\{x_{ij}\}_{1 \le i \le p, 1 \le j \le n}$, \mathbf{T}_p is a $p \times p$ deterministic matrix, $\mathbf{T}_p \mathbf{X}_n$ is considered a random sample from the population with the population covariance matrix $\mathbf{T}_p \mathbf{T}_p^* = \mathbf{\Sigma}$, and * represents the complex conjugate transpose. In the sequel, we simply write $\mathbf{B} \equiv \mathbf{B}_n$, $\mathbf{T} \equiv \mathbf{T}_p$ and $\mathbf{X} \equiv \mathbf{X}_n$ when there is no confusion. Let $\lambda_1 \ge \cdots \ge \lambda_p$ be the eigenvalues of \mathbf{B} . For a known test function f, we call $\sum_{j=1}^p f(\lambda_j)$ a linear spectral statistic (LSS) of \mathbf{B} . Because most of the classical test statistics in multivariate statistical analysis are associated with the eigenvalues of sample covariance matrices, LSSs are remarkable tools in many statistical problems (see Anderson (2003); Yao et al. (2015) for details). Through extensive study of high-dimensional data, it has been discovered that the distributions of LSSs significantly differ between lowdimensional and high-dimensional data. For example, in the low-dimensional setting, Wilks' theorem (see Wilks (1938)) provides the χ^2 approximation for the likelihood ratio test (LRT) statistic, which is a kind of LSS. However, when p is large compared with the sample size n, the LRT statistic exhibits Gaussian fluctuations (see Bai et al. (2009); Jiang and Yang (2013)). More generally, Bai and Silverstein (2004) established the central limit theorem

MSC2020 subject classifications: Primary 60B20; secondary 60F05.

Keywords and phrases: Empirical spectral distribution, linear spectral statistic, random matrix, Stieltjes transform.

(CLT) for the LSSs of a high-dimensional B under Gaussian-like moments condition by employing random matrix theory (RMT). Here the term 'Gaussian-like moments' refers to the population second-order and fourth-order moments are the same as those of real or complex Gaussian population. We refer to this CLT as the Bai–Silverstein theorem (BST) for brevity. Following the work of Bai and Silverstein (2004), many extensions have been developed under different settings. Pan and Zhou (2008) generalized the BST by relaxing the Gaussianlike moments condition of x_{ij} , which at the price of adding a structural condition on T. Zheng (2012), Yang and Pan (2015) and Bao et al. (2022) extended the BST to multivariate F matrices, canonical correlation matrices and block correlation matrices, respectively. Pan (2014) presented the CLT for the LSS of noncentered sample covariance matrices, and Zheng et al. (2015) studied the case of an unbiased sample covariance matrix when the population mean is unknown. Chen and Pan (2015) focused on the ultrahigh dimensional case in which the dimension p is much larger than the sample size n. Gao et al. (2017) and Li et al. (2021) studied the CLTs for the LSSs of high-dimensional Spearman and Kendall's rank correlation matrices, respectively. Without attempting to be comprehensive, we also refer readers to other extensions (Bai et al., 2007, 2015, 2019; Zheng et al., 2019; Banna et al., 2020; Najim and Yao, 2016; Baik et al., 2018; Hu et al., 2019; Jiang and Bai, 2021).

Almost all the literature mentioned above have traditionally assumed that the spectral norms of Σ are bounded in n. This assumption limits the applications in data analysis because in many fields, such as economics and wireless communication networks, the leading eigenvalues may tend to infinity. We present two examples here.

• Signal detection (Johnstone and Nadler (2017)): We consider a single signal model:

$$\boldsymbol{x} = \chi_s^{1/2} \boldsymbol{u} \boldsymbol{h} + \sigma \boldsymbol{v},$$

where h is an unknown *p*-dimensional unit vector, u is a random variable distributed as N(0,1), χ_s is the signal strength, σ is the noise level, and v is a random noise vector that is independent of u and follows a multivariate Gaussian distribution $N_p(0, \Sigma_v)$. It is easy to check that the covariance matrix of x is $\Sigma_x = \sigma^2 \Sigma_v + \chi_s h h^{\top}$. When the noise level is low, but the signal strength is large and sometimes tends to infinity, it is illogical to assume the boundedness of Σ_x .

• *Factor model* (Bai and Ng (2002)): Many economic analyses, such as arbitrage pricing theory and analyses of the rank of a demand system, align naturally within the framework of the factor model:

$$\boldsymbol{x}_t = \boldsymbol{\Lambda} \quad \boldsymbol{f}_t + \boldsymbol{\varepsilon}_t \quad t = 1, \dots, T.$$

where x_t is the observed data, N represents the number of cross-sections, T is a large time dimension, and f_t , Λ and ε_t represent the common factors, the factor loadings and the idiosyncratic error term, respectively. To ensure the identification of the model, several conventional assumptions are needed, such as $\mathbb{E}f_t = 0$, $\mathbb{E}(f_t f_t^{\top}) = \mathbf{I}_r$, ε_t is independent of f_t with $\mathbb{E}\varepsilon_t = 0$ and $\mathbb{E}\varepsilon_t \varepsilon_t^{\top} = \Sigma_{\varepsilon} > 0$. Then the covariance matrices of x_t can be expressed as $\Sigma_x = \Lambda \Lambda^{\top} + \Sigma_{\varepsilon}$. A pervasive assumption is that the variances of the principal components Λf_t can diverge as N increases to infinity (see Assumption B of Bai and Ng (2002)). Therefore, the spectral norms of Σ_x are unbounded.

For these reasons, it is of practical value to obtain the asymptotic properties of the LSS when Σ is unbounded. Therefore, in this paper, we focus on a generalized CLT for the LSSs of a spiked covariance matrix with the following structure:

(1.1)
$$\boldsymbol{\Sigma} = \mathbf{V} \begin{pmatrix} \mathbf{D}_1 & 0\\ 0 & \mathbf{D}_2 \end{pmatrix} \mathbf{V}^*,$$

where V is a unitary matrix, D_1 is a diagonal matrix consisting of the descending unbounded eigenvalues, and D_2 is the diagonal matrix of the bounded eigenvalues. As an application, the established CLT is employed to study the asymptotic behaviors of two special LSSs, i.e., the likelihood ratio (LR) statistic and Nagao's trace (NT) statistic, under the hypothesis

(1.2)
$$H_0: \mathbf{\Sigma} = \mathbf{I}_p \quad \text{vs.} \quad H_1: \mathbf{\Sigma} = \mathbf{V} \begin{pmatrix} \mathbf{D}_1 & 0\\ 0 & \mathbf{I}_{p-M} \end{pmatrix} \mathbf{V}^*,$$

where M is a constant. We also derive the asymptotic power of Roy's largest root test to detect the above hypothesis and make a comparison with these two LSSs.

The setting (1.1) is attributed to the famous spiked model in which a few large eigenvalues of the population covariance matrix are assumed to be well separated from the remaining eigenvalues (Johnstone, 2001). The spiked model has served as the foundation for a rich theory of principal component analysis through the performance of extreme eigenvalues, as discussed in Baik and Silverstein (2006); Paul (2007); Bai and Yao (2008); Nadler (2008); Jung and Marron (2009); Bai and Yao (2012); Onatski et al. (2014); Bloemendal et al. (2016); Wang and Yao (2017); Donoho et al. (2018); Perry et al. (2018); Johnstone and Paul (2018); Yang and Johnstone (2018); Yao et al. (2018); Dobriban (2020); Johnstone and Onatski (2020); Cai et al. (2020); Jiang and Bai (2021). There are also several works that have considered the asymptotic behaviors of various quantities as the spike strengths tend to infinity. Specifically, Zhou and Marron (2015) focused on the consistency of the sample eigenvector, corresponding to the largest eigenvalue of the sample covariance matrix, under high dimension and low sample size settings, when Σ is unbounded and the data set is Gaussian. Wang and Fan (2017) derived the asymptotic distributions of the spiked eigenvalues and eigenvectors when Σ is diagonal and unbounded, and the data set is sub-Gaussian. Recently, Li et al. (2020), Yin (2021) and Zhang et al. (2022) investigated the trace of a large sample covariance matrix under the spiked model assumption.

To summarize, the contributions of this paper are as follows.

- We demonstrate a nontrivial extension of the BST to the situation in which the spectral norms of the population covariance matrices are allowed to diverge as min{p, n} → ∞. In particular, we show how the test function f and the divergence rate of the population spectral norm affect the new CLT.
- 2. It was previously reported that Gaussian-like moments or diagonality of the population covariance matrix are necessary for the CLT of the LSS (e.g., Zheng et al. (2015)). Nevertheless, we prove that these restrictions can be completely removed by normalizing the LSS. More importantly, even if no limit exists on the variance of the LSS, the new CLT could still hold.
- 3. The entire technical part of this paper is built on the decomposition of the LSS $\sum_{j=1}^{p} f(\lambda_j) = \sum_{j=1}^{M} f(\lambda_j) + \sum_{j=M+1}^{p} f(\lambda_j)$. Because the classical delta method cannot be applied to the unbounded part $\sum_{j=1}^{M} f(\lambda_j)$ and the bounded part $\sum_{j=M+1}^{p} f(\lambda_j)$ is not a strict LSS of a sample covariance matrix, the results of Bai and Silverstein (2004) and Jiang and Bai (2021) cannot be adopted directly. In this paper, we leverage a 'generalized delta method' and employ skillful transformations to prove the CLTs for the unbounded parts are asymptotically independent, which leads to the establishment of the new CLT.
- 4. We verify that Roy's largest root test is most powerful among the common tests when the alternative (1.2) has only one spiked eigenvalue, which has also been mentioned by Olson (1974); Johnstone and Nadler (2017). Furthermore, we demonstrate that when the number of spikes is larger than one, the LSSs could exhibit higher asymptotic power than Roy's largest root test when the divergence rates of spiked eigenvalues are higher than \sqrt{n} .

The remaining sections are organized as follows: Section 2 presents a detailed description of our notations and assumptions. The main results for the CLT for the LSSs of sample covariance matrices are stated in Section 3. In Section 4, we explore an application of our main results. We also report the results of numerical studies in Section 5. Technical proofs are presented in Section 6. This paper is also accompanied by an online supplementary file that includes the following materials: (i) some postponed proofs for Theorems 3.1–4.5; (ii) some additional simulation results, and (iii) some useful lemmas.

2. Notations and assumptions. Throughout the paper, we use bold capital letters and bold italic lowercase letters to represent matrices and vectors, respectively. Scalars are represented by regular letters. e_i denotes a standard basis vector whose components are all zero, except the *i*-th component, which is equal to 1. We use $tr(\mathbf{A})$, \mathbf{A}^{\top} and \mathbf{A}^* to denote the trace, transpose and conjugate transpose of matrix \mathbf{A} , respectively. We also use f' to denote the derivative of function f, and we use $\frac{\partial}{\partial z_1} f(z_1, z_2)$ to denote the partial derivative of function f with respect to z_1 . Let $[\mathbf{A}]_{ij}$ denote the (i, j)-th entry of the matrix \mathbf{A} and $\oint_C f(z)dz$ denote the contour integral of f(z) on the contour C. Let $\lambda_i^{\mathbf{A}}$ be the *i*th largest eigenvalue of matrix \mathbf{A} . Weak convergence is denoted by $\stackrel{d}{\rightarrow}$. Throughout this paper, we use o(1) (resp. $o_p(1)$) to denote a negligible scalar (resp. in probability), and the notation C represents a generic constant that may vary from line to line.

We adopt the notation $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_n) = (x_{ij}), 1 \le i \le p, 1 \le j \le n$. Let $\rho_1 \ge \dots \ge \rho_p$ be the eigenvalues of Σ and the singular value decomposition of \mathbf{T} be

(2.3)
$$\mathbf{T} = \mathbf{V}\mathbf{D}^{1/2}\mathbf{U}^* = (\mathbf{V}_1, \mathbf{V}_2) \begin{pmatrix} \mathbf{D}_1^{\frac{1}{2}} & 0\\ 0 & \mathbf{D}_2^{\frac{1}{2}} \end{pmatrix} (\mathbf{U}_1, \mathbf{U}_2)^*$$

Here U and V are unitary matrices, and $\mathbf{D}_1 = \text{diag}(\underbrace{\alpha_1, \dots, \alpha_1}_{d_1}, \underbrace{\alpha_2, \dots, \alpha_2}_{d_2}, \dots, \underbrace{\alpha_K, \dots, \alpha_K}_{d_K})$

is a diagonal matrix whose diagonal elements tend to infinity. To avoid confusion, we refer to $\{\alpha_i, i = 1, ..., K\}$ as the diverging spikes in the following. Assume $d_1 + \cdots + d_K = M$. \mathbf{D}_2 is the diagonal matrix of the eigenvalues with bounded components, including bounded spiked eigenvalues and bulk eigenvalues. Moreover, let $d_0 = 0$ and $J_k = \left\{\sum_{i=0}^{k-1} d_i + 1, \ldots, \sum_{i=0}^{k} d_i\right\}$, thus $\rho_i = \alpha_k$ if $i \in J_k$. Then, the corresponding sample covariance matrix. Corresponding to the decomposition of \mathbf{D} , we decompose $\mathbf{V} = (\mathbf{V}_1, \mathbf{V}_2)$, $\mathbf{U} = (\mathbf{U}_1, \mathbf{U}_2)$, and denote $\mathbf{\Gamma} = \mathbf{V}_2 \mathbf{D}_2^{1/2} \mathbf{U}_2^*$, $\mathbf{r}_j = \frac{1}{\sqrt{n}} \mathbf{\Gamma} \mathbf{x}_j$, and $\mathbf{A}_j = \frac{1}{n} \mathbf{\Gamma} \mathbf{X} \mathbf{X}^* \mathbf{\Gamma}^* - z\mathbf{I} - \mathbf{r}_j \mathbf{r}_j^*$. Let \mathbb{E}_j be the conditional expectation with respect to the σ -field generated by $\mathbf{r}_1, \ldots, \mathbf{r}_j$. For any matrix \mathbf{A} with real eigenvalues, the empirical spectral distribution of \mathbf{A} is denoted by

$$F^{\mathbf{A}}(x) = \frac{1}{p}$$
 (number of eigenvalues of $\mathbf{A} \le x$).

For any function of bounded variation F on the real line, its Stieltjes transform is defined by

$$m_F(z) = \int \frac{1}{\lambda - z} \, \mathrm{d}F(\lambda), \quad z \in \mathbb{C}^+ := \{ z \in \mathbb{C} : \Im z > 0 \}.$$

The assumptions used to obtain the results in this paper are as follows:

ASSUMPTION 1. $\{x_{ij}, 1 \le i \le p, 1 \le j \le n\}$ are i.i.d. random variables with common moments

$$\mathbb{E}x_{ij} = 0, \quad \mathbb{E}|x_{ij}|^2 = 1, \quad \beta_x = \mathbb{E}|x_{ij}|^4 - |\mathbb{E}x_{ij}^2|^2 - 2, \quad \alpha_x = |\mathbb{E}x_{ij}^2|^2.$$

ASSUMPTION 2. As $\min\{p, n\} \to \infty$, the ratio of the dimension-to-sample size (RDS) $c_n := p/n \to c > 0$.

REMARK 2.1. Assumptions 1 and 2 are standard in RMT. If $\mathbb{E}x_{ij} \neq 0$, we can use the centralized sample covariance matrices and n-1 instead of \mathbf{B}_n and n, respectively, and the following results also hold. Details can be found in (Zheng et al., 2015). Therefore, in the sequel, we assume that $\mathbb{E}x_{ij} = 0$ without loss of generality.

ASSUMPTION 3. **T** is nonrandom. As $\min\{p, n\} \to \infty$, $\alpha_K \to \infty$ and $H_n := F^{\Gamma\Gamma^*} \xrightarrow{d} H$, where *H* is a distribution function on the real line. *M* is fixed.

REMARK 2.2. Similar to Silverstein (1995) that under Assumptions 1-3, $F^{\mathbf{B}} \xrightarrow{d} F^{c,H}$ almost surely, where $F^{c,H}$ is a nonrandom distribution function whose Stieltjes transform $m := m_{F^{c,H}}(z)$ satisfies the following equation:

(2.4)
$$m = \int \frac{1}{t(1 - c - czm) - z} dH(t).$$

In the sequel, we call $F^{c,H}$ the limiting spectral distribution (LSD) of **B**. Moreover, because the matrix $\underline{\mathbf{B}} = \frac{1}{n} \mathbf{X}^* \mathbf{T}^* \mathbf{T} \mathbf{X}$ shares the same nonzero eigenvalues as **B**, equation (2.4) can be rewritten as

$$\underline{m} = -\left(z - c\int \frac{t}{1 + t\underline{m}}dH(t)\right)^{-1},$$

where $\underline{m} := m_{F^{c,H}}(z)$ represents the Stieltjes transform of the LSD of $\underline{\mathbf{B}}$.

ASSUMPTION 4. Test functions f_1, \ldots, f_h are analytic on a connected open region of the complex plane containing the support of F^{c_n,H_n} for almost all n. Moreover, we suppose that for any $l = 1, \ldots, h$,

$$\lim_{\substack{\{x_n, y_n\}\to\infty\\x_n/y_n\to 1}}\frac{f_l'(x_n)}{f_l'(y_n)} = 1.$$

REMARK 2.3. In fact, Assumption 4 is not highly restrictive in practice, as many common functions such as logarithmic and polynomial functions satisfy it. However, it is worth noting that the exponential function does not satisfy this assumption.

For convenience of description, we introduce some notations before presenting the main results in the next section. Let $\underline{F}^{c,H}$ denote the LSD of matrix $n^{-1}\mathbf{X}^*\mathbf{U}_2\mathbf{D}_2\mathbf{U}_2^*\mathbf{X}$, $\mathbf{U}_1 = (u_{ij})_{i=1,\dots,p;j=1,\dots,M}$, $\mathcal{U}_{i_1j_1i_2j_2} = \sum_{t=1}^p \overline{u}_{ti_1}u_{tj_1}u_{ti_2}\overline{u}_{tj_2}$, $\phi_n(x) = x\left(1+c_n\int \frac{t}{x-t}dH_n(t)\right)$, $\phi_k = \phi(x)|_{x=\alpha_k} = \alpha_k\left(1+c\int \frac{t}{\alpha_k-t}dH(t)\right)$, $\theta_k = \phi_k^2\underline{m}_2(\phi_k)$, $\nu_k = \phi_k^2\underline{m}^2(\phi_k)$, $\underline{m}(\lambda) = \int \frac{1}{x-\lambda}d\underline{F}^{c,H}(x)$, $\underline{m}_2(\lambda) = \int \frac{1}{(\lambda-x)^2}d\underline{F}^{c,H}(x)$,

 $c_{nM} = \frac{p - M}{n}, \ H_{2n} = F^{\mathbf{D}_2}, \ \mathbf{P}_n(z) = ((1 - c_{nM})\mathbf{\Gamma}\mathbf{\Gamma}^* - zc_{nM}m_{2n0}(z)\mathbf{\Gamma}\mathbf{\Gamma}^* - z\mathbf{I}_p)^{-1},$

$$\begin{split} \varpi_{nkl} &= \frac{\phi_n(\alpha_k)}{\sqrt{n}} f_l'(\phi_n(\alpha_k)), \ s_k^2 = \frac{(\alpha_x + 1) d_k}{\theta_k} + \frac{\beta_x \nu_k \sum_{j_1, j_2 \in J_k} \mathcal{U}_{j_1 j_2 j_2}}{\theta_k^2}, \\ &\qquad \vartheta_n^2 = \Theta_{0,n}(z_1, z_2) + \alpha_x \Theta_{1,n}(z_1, z_2) + \beta_x \Theta_{2,n}(z_1, z_2), \\ &\qquad \Theta_{0,n}(z_1, z_2) = \frac{\underline{m}'_{2n0}(z_1)\underline{m}'_{2n0}(z_2)}{(\underline{m}_{2n0}(z_1) - \underline{m}_{2n0}(z_2))^2} - \frac{1}{(z_1 - z_2)^2}, \\ &\qquad \Theta_{1,n}(z_1, z_2) = \frac{\partial}{\partial z_2} \left\{ \frac{\partial \mathcal{A}_n(z_1, z_2)}{\partial z_1} \frac{1}{1 - \alpha_x \mathcal{A}_n(z_1, z_2)} \right\}, \\ &\qquad \mathcal{A}_n(z_1, z_2) = \frac{z_1^2 z_2^2 \underline{m}'_{2n0}(z_1) \underline{m}_{2n0}(z_2) \mathrm{tr} \mathbf{\Gamma}^* \mathbf{P}_n(z_1) \mathbf{\Gamma} \mathbf{\Gamma}^\top \mathbf{P}_n(z_2)^\top \mathbf{\bar{\Gamma}}, \\ &\qquad \Theta_{2,n}(z_1, z_2) = \frac{z_1^2 z_2^2 \underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{n} \sum_{i=1}^p \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_1) \mathbf{\Gamma} \right]_{ii} \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_2) \mathbf{\Gamma} \right]_{ii}, \\ &\mu_l = -\frac{\alpha_x}{2\pi i} \cdot \oint_{\mathcal{C}} \frac{c_{nM} f_l(z) \int \underline{m}_{2n0}^3(z) t^2 \left(1 + t\underline{m}_{2n0}(z)\right)^{-3} dH_{2n}(t)}{\left(1 - \alpha_n M \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t\underline{m}_{2n0}(z))^2} dH_{2n}(t)\right) \left(1 - \alpha_x c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t\underline{m}_{2n0}(z))^2} dH_{2n}(t)\right)} dz \end{split}$$

$$-\frac{\beta_x}{2\pi i} \cdot \oint_{\mathcal{C}} \frac{c_{nM} f_l(z) \int \underline{m}_{2n0}^3(z) t^2 \left(1 + t \underline{m}_{2n0}(z)\right)^{-3} dH_{2n}(t)}{1 - c_{nM} \int \underline{m}_{2n0}^2(z) t^2 \left(1 + t \underline{m}_{2n0}(z)\right)^{-2} dH_{2n}(t)} dz, \quad l = 1, \dots, h.$$

Here, $m_{2n0}(z)$ is the Stieltjes transform of $F^{c_{nM},H_{2n}}$, where $F^{c_{nM},H_{2n}}$ is the LSD $F^{c,H}$ with $\{c, H\}$ replaced by $\{c_{nM}, H_{2n}\}$, $\underline{m}_{2n0}(z) = -\frac{1-c_{nM}}{z} + c_{nM}m_{2n0}(z)$ and C is a closed contour in the complex plane enclosing the support of F^{c_n,H_n} and it is also enclosed in the analytic area of f_l . For clarity, $m_{1n0}(z)$ denotes the Stieltjes transform of F^{c_n,H_n} , $m_n = \frac{1}{p} \operatorname{tr} (\mathbf{B} - z\mathbf{I}_p)^{-1}$, and $m_{2n} = \frac{1}{p-M} \operatorname{tr} (\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-1}$.

Note that

$$\sum_{j=1}^{p} f(\lambda_j) = p \int f(x) \, dF^{\mathbf{B}}(x).$$

Thus, for brevity, we define the normalized LSSs as

$$Y_{l} = \int f_{l}(x) \, dG_{n}(x) - \sum_{k=1}^{K} d_{k} f_{l}(\phi_{n}(\alpha_{k})) - \frac{M}{2\pi i} \oint_{\mathcal{C}} f_{l}(z) \, \frac{\underline{m}'_{2n0}(z)}{\underline{m}_{2n0}(z)} dz, \quad l = 1, 2, \dots, h,$$

where

$$G_n(x) = p[F^{\mathbf{B}}(x) - F^{c_n, H_n}(x)].$$

3. Main results. Now, we are in a position to present our main theorems and their proofs are provided in Section 6 and the supplementary material. We first establish a CLT for an LSS without any restrictions imposed on the Gaussian moments or on the structures of the population covariance matrix by normalizing the LSS.

THEOREM 3.1. Under Assumptions 1–4, we have

$$\frac{Y_1 - \mu_1}{\varsigma_1} \xrightarrow{d} N\left(0, 1\right),$$

where

(3.5)
$$\varsigma_1^2 = \sum_{k=1}^K \varpi_{nk1}^2 s_k^2 - \frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_1(z_1) f_1(z_2) \vartheta_n^2 dz_1 dz_2,$$

 C_1 and C_2 are nonoverlapping and closed contours in the complex plane enclosing the support of F^{c_n,H_n} . C_1 and C_2 are also enclosed in the analytic area of f_1 .

REMARK 3.1. Recall the definitions $s_k^2 = \frac{(\alpha_k+1)d_k}{\theta_k} + \frac{\beta_x \nu_k \sum_{j_1, j_2 \in J_k} \mathcal{U}_{j_1 j_1 j_2 j_2}}{\theta_k^2}$ and $\vartheta_n^2 = \Theta_{0,n}(z_1, z_2) + \alpha_x \Theta_{1,n}(z_1, z_2) + \beta_x \Theta_{2,n}(z_1, z_2)$. Notably, the term $\Theta_{0,n}(z_1, z_2)$ has a limitation under Assumptions 1-4, which has already been discussed in Bai and Silverstein (2004). Additionally, if Σ is complex, the convergence of $\Theta_{1,n}(z_1, z_2)$ is not guaranteed. The term $\Theta_{2,n}(z_1, z_2)$ involves the quantities $[\Gamma^* \mathbf{P}_n^2(z_i) \Gamma]_{ii}$, which depend not only on the eigenvalues of \mathbf{D}_2 but also on their associated eigenvectors. Furthermore, the term s_k^2 indicates that the variance is influenced by the second and fourth moments of x_{ij} , spiked eigenvalues, and their associated eigenvectors. The limit of ϖ_{nk1}^2 is allowed to be infinite.

REMARK 3.2. After a closer look at the variance (3.5), we can also find that the first part of the formula (3.5) is the variance containing diverging spikes and the second part is the variance containing the bounded eigenvalues. When $\phi_n(\alpha_1) f'_1(\phi_n(\alpha_1)) = o(\sqrt{n})$, the first term in formula (3.5) tends to 0, and the variance is mainly affected by the second part. When $\phi_n(\alpha_1) f'_1(\phi_n(\alpha_1))$ is of order \sqrt{n} , the two parts of formula (3.5) are of the same order, and the variance is affected by both. When the order of $\phi_n(\alpha_1) f'_1(\phi_n(\alpha_1))$ is higher than \sqrt{n} , the first part of formula (3.5) is much larger than the second part; therefore, the spiked part dominates the variance value.

As a minor price for the removal of the bounded spectrum condition, the new CLT described above applies only to a single LSS. To guarantee that the new CLT will apply to multiple normalized LSSs, structural assumptions about the population covariance matrices are needed.

ASSUMPTION 5. T is real or the variables x_{ij} are complex satisfying $\alpha_x = 0$.

ASSUMPTION 6. $\mathbf{T}^*\mathbf{T}$ is diagonal or $\beta_x = 0$.

REMARK 3.3. Assumptions 5 and 6 are used as a replacement for the Gaussian-like moments condition. It has been proved by Zheng et al. (2015) that these two structural assumptions regarding the population matrices are necessary for their results when the Gaussianlike moments condition in the BST does not hold.

The following theorem is a nontrivial extension of the BST:

THEOREM 3.2. Under Assumptions 1–6, the random vector

$$\left(\frac{Y_1-\mu_1}{\sigma_1},\ldots,\frac{Y_h-\mu_h}{\sigma_h}\right)^{\top} \stackrel{d}{\to} N_h\left(0,\Psi\right),$$

with variance

$$\sigma_l^2 = \sum_{k=1}^K \varpi_{nkl}^2 s_k^2 - \kappa_{nll}, \quad l = 1, \dots, h,$$

and covariance matrix $\Psi = (\psi_{st})_{h \times h}$, where $\psi_{st} = \lim_{n \to \infty} \psi_{nst}$,

$$\begin{split} \psi_{nst} &= \frac{\sum_{k=1}^{K} \overline{\omega}_{nks} \overline{\omega}_{nkt} s_{k}^{2} - \kappa_{nst}}{\sqrt{\sum_{k=1}^{K} \overline{\omega}_{nks}^{2} s_{k}^{2} - \kappa_{nss}} \sqrt{\sum_{k=1}^{K} \overline{\omega}_{nkt}^{2} s_{k}^{2} - \kappa_{ntt}}}, \\ \kappa_{nst} &= \frac{1}{4\pi^{2}} \oint_{\mathcal{C}_{1}} \oint_{\mathcal{C}_{2}} \frac{f_{s}(z_{1}) f_{t}(z_{2})}{(\underline{m}_{2n0}(z_{1}) - \underline{m}_{2n0}(z_{2}))^{2}} d\underline{m}_{2n0}(z_{1}) d\underline{m}_{2n0}(z_{2}) \\ &+ \frac{c_{nM} \beta_{x}}{4\pi^{2}} \oint_{\mathcal{C}_{1}} \oint_{\mathcal{C}_{2}} \int \frac{f_{s}(z_{1}) f_{t}(z_{2}) t^{2}}{(\underline{m}_{2n0}(z_{1}) t + 1)^{2} (\underline{m}_{2n0}(z_{2}) t + 1)^{2}} dH_{2n}(t) d\underline{m}_{2n0}(z_{1}) d\underline{m}_{2n0}(z_{2}) \\ &+ \frac{1}{4\pi^{2}} \oint_{\mathcal{C}_{1}} \oint_{\mathcal{C}_{2}} f_{s}(z_{1}) f_{t}(z_{2}) \left[\frac{\partial^{2}}{\partial z_{1} \partial z_{2}} \log\left(1 - a_{n}(z_{1}, z_{2})\right) \right] dz_{1} dz_{2}, \end{split}$$

and

$$a_{n}(z_{1}, z_{2}) = \alpha_{x} \left(1 + \frac{\underline{m}_{2n0}(z_{1}) \underline{m}_{2n0}(z_{2})(z_{1} - z_{2})}{\underline{m}_{2n0}(z_{2}) - \underline{m}_{2n0}(z_{1})} \right).$$

 $\operatorname{REMARK} 3.4. \quad \operatorname{Define} \tilde{\psi}_{nst} = \frac{\sum_{k=1}^{K} \varpi_{nks} \varpi_{nkt} s_k^2 - \frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_s(z_1) f_t(z_2) \vartheta_n^2 dz_1 dz_2}{\varsigma_s \varsigma_t}.$

If $\tilde{\Psi}_n = (\tilde{\psi}_{nst})_{h \times h}$ is invertible for all sufficiently large *n*, we conjecture that, similar to Theorem 3.1, the convergence

$$\tilde{\boldsymbol{\Psi}}_{n}^{-1/2} \left(\frac{Y_{1} - \mu_{1}}{\sigma_{1}}, \dots, \frac{Y_{h} - \mu_{h}}{\sigma_{h}} \right)^{\top} \stackrel{d}{\to} N_{h} \left(0, \mathbf{I}_{h} \right)$$

holds without requiring Assumptions 5 and 6. It should be noted that $\tilde{\Psi}_n$ is singular if the set of test functions is linearly dependent. However, determining the invertibility of $\tilde{\Psi}_n$ becomes challenging when the test functions are completely linearly independent. Hence, the extension to the removal of Assumptions 5 and 6 in Theorem 3.2 is left for future work.

REMARK 3.5. If $\varpi_{nkl} \to 0$ as $n \to \infty$, Theorem 3.2 coincides with Theorem 2.1 of Zheng et al. (2015). If the test functions $f_l = x$ and x^2 , then Theorem 3.2 reduces to Theorem 2.1 of Yin (2021). Notably, the results in Yin (2021) required higher-order moment conditions.

4. Application. In this section, we focus on a hypothesis test concerning whether the population covariance matrix Σ is equal to the identity matrix or a spiked model, i.e.,

(4.6)
$$H_0: \mathbf{\Sigma} = \mathbf{I}_p \quad \text{vs.} \quad H_1: \mathbf{\Sigma} = \mathbf{V} \begin{pmatrix} \mathbf{D}_1 & 0 \\ 0 & \mathbf{I}_{p-M} \end{pmatrix} \mathbf{V}^*,$$

where D_1 is a diagonal matrix of the diverging spiked eigenvalues of Σ . There are several classical test statistics for this problem, but due to the limited length of this paper, we only consider the likelihood ratio (LR) test statistic (Wilks, 1938) and the Nagao's trace (NT) test statistic (Nagao, 1973) in this section. Specifically, the LR and NT statistics can be formulated as

$$L = \operatorname{tr} \mathbf{B} - \log |\mathbf{B}| - p \text{ and } W = \operatorname{tr} (\mathbf{B} - \mathbf{I}_p)^2,$$

respectively. Under the null hypothesis, the asymptotic properties of the LR and NT statistics for high-dimensional settings have been investigated extensively in the literature; here, we refer to Bai et al. (2009); Jiang and Yang (2013); Ledoit and Wolf (2002); Wang and Yao (2013); Onatski et al. (2013) for more details. Thus, in this section, we mainly focus on the alternative hypothesis. However, to provide better comparisons, we also present the asymptotic distributions under the null hypothesis in the following theorems.

4.1. Asymptotic results for the LR and NT statistics. In this subsection, we present the asymptotic results for LR and NT test statistics for the testing problem (4.6).

THEOREM 4.1 (CLT for the LR statistic). Under Assumptions 1–4 with $c_n = p/n \rightarrow c \in (0, 1)$, we have

• (Under H_0)

$$\frac{L - p\ell_l - \mu_l}{\varsigma_l} \xrightarrow{d} N(0, 1),$$

where

$$\ell_l = 1 - \frac{c_n - 1}{c_n} \log(1 - c_n), \quad \mu_l = -\frac{\log(1 - c_n)}{2} \alpha_x + \frac{c_n}{2} \beta_x$$

and

$$\varsigma_l^2 = (\alpha_x + 1)(-\log(1 - c_n) - c_n).$$

• (Under H_1)

$$\frac{L - (p - M)\check{\ell}_l - \check{\mu}_l}{\check{q}} \longrightarrow N(0, 1),$$

where

$$\check{\ell}_{l} = 1 - \frac{c_{nM} - 1}{c_{nM}} \log \left(1 - c_{nM} \right), \ \check{\mu}_{l} = -\frac{\log \left(1 - c_{nM} \right)}{2} \alpha_{x} + \frac{c_{nM}}{2} \beta_{x}$$

$$+ \sum_{k=1}^{K} d_{k} \left(\phi_{n} \left(\alpha_{k} \right) - \log \phi_{n} \left(\alpha_{k} \right) - 1 \right) - M(c_{nM} + \log(1 - c_{nM}))$$

and

$$\vec{\varsigma}_l^2 = \sum_{k=1}^K \frac{(\phi_n(\alpha_k) - 1)^2}{n} s_k^2 + (\alpha_x + 1) \left(-\log(1 - c_{nM}) - c_{nM}\right)$$

REMARK 4.1. If $c \ge 1$, then \mathbf{B}_n could be singular for large n, which would give rise to an undefined LR statistic L. Thus, the additional restriction c < 1 is added in Theorem 4.1.

REMARK 4.2. Note that $\phi_n(\alpha_k)$ and s_k are defined in Section 2. Under the alternative hypothesis H_1 in (4.6), we can adopt simplification $\phi_n(\alpha_k) = \alpha_k + c_{nM} + o(1)$, and

$$s_k^2 = (\alpha_x + 1) d_k + \beta_x \sum_{j_1, j_2 \in J_k} \mathcal{U}_{j_1 j_1 j_2 j_2} + o(1).$$

THEOREM 4.2 (CLT for the NT statistic). Under Assumptions 1–4, we have • (Under H_0)

$$\frac{W - pc_n - \mu_w}{\varsigma_w} \xrightarrow{d} N(0, 1),$$

where

$$\mu_w = c_n(\alpha_x + \beta_x)$$
 and $\varsigma_w^2 = (\alpha_x + 1)(4c_n^3 + 2c_n^2) + 4\beta_x c_n^3$.

• (Under H_1)

$$\frac{W - (p - M)c_{nM} - \breve{\mu}_w}{\breve{\varsigma}_w} \xrightarrow{d} N(0, 1),$$

where

$$\breve{\mu}_{w} = c_{nM}(\alpha_{x} + \beta_{x}) + \sum_{k=1}^{K} d_{k} (\phi_{n}(\alpha_{k}) - 1)^{2} - Mc_{nM}^{2}$$

and

$$\vec{\varsigma}_{w}^{2} = \sum_{k=1}^{K} \frac{4\phi_{n}^{2}\left(\alpha_{k}\right)\left(\phi_{n}\left(\alpha_{k}\right)-1\right)^{2}}{n} s_{k}^{2} + \left(\alpha_{x}+1\right)\left(4c_{nM}^{3}+2c_{nM}^{2}\right) + 4\beta_{x}c_{nM}^{3}.$$

REMARK 4.3. From the covariance terms ξ_l^2 and ξ_w^2 , one can find that these CLTs are related to the components of the right singular vectors **U**, but not to the left singular vectors **V**. Furthermore, since **D**₂ is an identity matrix, **U**₂ does not affect the asymptotic CLTs. Therefore, the only singular vectors of **T** affecting the results are **U**₁, which are involved in s_k^2 .

The proofs of Theorems 4.1 and 4.2 are given in the supplementary material. To avoid confusion with the classical distributions of the LR test and NT test, we refer to the CLTs above as the corrected LR test (CLRT) and corrected NT test (CNTT) in the sequel. From Theorems 4.1 and 4.2, we reject the null hypothesis H_0 in (4.6) if

$$L > z_{\xi}\varsigma_l + p\ell_l + \mu_l$$

and

$$W > z_{\xi}\varsigma_w + c_n(p + \alpha_x + \beta_x),$$

where ξ is the significance level of the test and z_{ξ} is the $1 - \xi$ quantile of the standard Gaussian distribution Φ . For the power functions of CLRT and CNTT, we have the following theorems.

THEOREM 4.3 (Power function of CLRT). Under Assumptions 1–4 with $c_n = p/n \rightarrow c \in (0,1)$ and H_1 in (4.6), we have that the power function of the CLRT $P_L = P(L > z_{\xi}\varsigma_l + p\ell_l + \mu_l)$ satisfies

$$(4.7) P_L - \Phi\left(\frac{\sum_{k=1}^K d_k \left(\phi_n\left(\alpha_k\right) - \log \phi_n\left(\alpha_k\right)\right) - M(1+c) - z_{\xi}\varsigma_l}{\sqrt{\sum_{k=1}^K \frac{\left(\phi_n\left(\alpha_k\right) - 1\right)^2}{n}} s_k^2 + \left(\alpha_x + 1\right)\left(-\log(1-c) - c\right)}}\right) \to 0,$$

as $n \to \infty$.

THEOREM 4.4 (Power function of CNTT). Under Assumptions 1–4 and H_1 in (4.6), we have that the power function of the CNTT $P_W = P(W > z_{\xi}\varsigma_w + c_n(p + \alpha_x + \beta_x))$ satisfies

$$(4.8) \quad P_W - \Phi\left(\frac{\sum_{k=1}^K d_k \left(\phi_n(\alpha_k) - 1\right)^2 - Mc^2 - 2Mc - z_{\xi}\varsigma_w}{\sqrt{\sum_{k=1}^K \frac{4\phi_n^2(\alpha_k)(\phi_n(\alpha_k) - 1)^2}{n}} s_k^2 + (\alpha_x + 1)\left(4c^3 + 2c^2\right) + 4\beta_x c^3}\right) \to 0,$$

as $n \to \infty$.

REMARK 4.4. Since s_k^2 is nonrandom and of order O(1), P_L and P_W tend to 1 as $\alpha_1 \rightarrow \infty$. The detailed analysis of the power functions of P_L and P_W is discussed in the next subsection.

10

4.2. *Power analysis.* This subsection discusses the power functions of P_L and P_W . For simplicity, in this subsection, we assume that $\{x_{ij}\}$ are real, i.e., $\alpha_x = 1$. We first derive the asymptotic power of Roy's largest root test (RLRT) to detect H_1 in (4.6) for comparison. Recall RLRT statistic λ_1 . Under Assumptions 1–4 and H_0 in (4.6), it follows from Theorem 2.7 of Ding and Yang (2018) that

$$\frac{\lambda_1 - \mu_r}{\varsigma_r} \stackrel{d}{\to} F_{TW},$$

where $\mu_r = (1 + \sqrt{c_n})^2$, $\varsigma_r = n^{-2/3} (1 + \sqrt{c_n}) (1 + \sqrt{c_n^{-1}})^{1/3}$ and F_{TW} is the Type 1 Tracy-Widom (TW) distribution. Let t_{ξ} be the $1 - \xi$ quantile of TW distribution with significance level ξ . Then we have the following theorem regarding the power function of RLRT.

THEOREM 4.5 (Power function of RLRT). Under Assumptions 1–4 and H_1 in (4.6), if the multiplicity of α_1 is one, then the power function of the RLRT $P_R = \mathbb{P}(\lambda_1 > t_{\xi}\varsigma_r + \mu_r)$ satisfies

(4.9)
$$P_R - \Phi\left(-\frac{t_{\xi}\varsigma_r + \mu_r - \phi_n\left(\alpha_1\right)}{s_1\phi_n\left(\alpha_1\right)/\sqrt{n}}\right) \to 0,$$

as $n \to \infty$.

The proof of this theorem is postponed to the supplementary material. It is clear that if $\alpha_1 > 1 + \sqrt{c}$ uniformly, then $P_R \rightarrow 1$ as $n \rightarrow \infty$. According to Anderson (2003), compared with the classical LSSs, RLRT has the highest asymptotic power to detect rank-one alternatives and under low dimensional settings. This property has also been demonstrated by Olson (1974) and Johnstone and Nadler (2017). In the following, we discuss the asymptotic power functions of CLRT, CNTT and RLRT. In particular, we will show that except for the case in which the number of spikes is equal to 1, CLRT and CNTT may exhibit higher asymptotic power than RLRT in some scenarios.

Define

(4.10)

$$\varkappa_{L} = \frac{\sum_{k=1}^{K} d_{k} \left(\phi_{n} \left(\alpha_{k}\right) - \log \phi_{n} \left(\alpha_{k}\right)\right) - M(1+c) - z_{\xi}\varsigma_{l}}{\sqrt{\sum_{k=1}^{K} \frac{(\phi_{n} \left(\alpha_{k}\right) - 1)^{2}}{n}} s_{k}^{2} - 2 \left(\log(1-c) + c\right)}}$$

$$\varkappa_{W} = \frac{\sum_{k=1}^{K} d_{k} \left(\phi_{n} \left(\alpha_{k}\right) - 1\right)^{2} - Mc^{2} - 2Mc - z_{\xi}\varsigma_{w}}{\sqrt{\sum_{k=1}^{K} \frac{4\phi_{n}^{2}\left(\alpha_{k}\right)\left(\phi_{n} \left(\alpha_{k}\right) - 1\right)^{2}}{n}} s_{k}^{2} + 2 \left(4c^{3} + 2c^{2}\right) + 4\beta_{x}c^{3}}}$$

$$\varkappa_{R} = \frac{\phi_{n} \left(\alpha_{1}\right) - \mu_{r} - t_{\xi}\varsigma_{r}}{s_{1}\phi_{n} \left(\alpha_{1}\right)/\sqrt{n}}.$$

Since CLRT, CNTT and RLRT statistics are all asymptotically normally distributed under the alternative hypothesis, according to formulas (4.7)–(4.9), comparing the convergence rates of power functions P_L , P_W , and P_R is equivalent to comparing the divergence rates \varkappa_L , \varkappa_W and \varkappa_R tend to infinity. Note that $\{z_{\xi}\varsigma_l, z_{\xi}\varsigma_w, t_{\xi}\varsigma_r\}$ are all of order O(1), $\{K, M\}$ are fixed, 0 < c < 1, $\phi_n(\alpha_k) = \alpha_k + c + o(1)$ and $s_k^2 = 2d_k + \beta_x \sum_{j_1, j_2 \in J_k} \mathcal{U}_{j_1 j_1 j_2 j_2} + o(1)$. In the sequel, we use the notations $A_n = \Omega(B_n)$, $A_n \simeq B_n$ and $A_n \asymp B_n$ to denote $B_n = O(A_n)$, $A_n = B_n + o(B_n)$ and $C^{-1}A_n < B_n < CA_n$, respectively, for some constant C > 1. Then, we have the following conclusions.

• (M = 1) For M = 1, i.e., there is only one diverging spike, we report the divergence rates of \varkappa_L , \varkappa_W and \varkappa_R in Table 1. From these results, we can conclude that the RLRT is

asymptotically more powerful than CLRT and CNTT whenever $\alpha_1 \to \infty$. Here, one should note that $1 + 2c + \beta_x c > 0$ and $\log(1 - c) + c < 0$ provided that c < 1. Moreover, if $\alpha_1 = o(n^{1/2})$, then the divergence rate of \varkappa_W is higher than \varkappa_L . However, when $\alpha_1 = \Omega(n^{1/2})$, \varkappa_L could be larger than \varkappa_W , such as $n = o(\alpha_1^2)$.

	\varkappa_L	\varkappa_W	\varkappa_R
$\alpha_1 = o(n^{\frac{1}{4}})$	$\asymp \alpha_1$	$\asymp \alpha_1^2$	$\simeq \frac{\sqrt{n}}{s_1}$
$\Omega(n^{\frac{1}{4}}) = \alpha_1 = o(n^{\frac{1}{2}})$	$\asymp \alpha_1$	$\simeq \frac{\sqrt{n}}{2\sqrt{s_1^2 + c^2(1 + 2c + \beta_x c)n/\alpha_1^4}}$	$\simeq \frac{\sqrt{n}}{s_1}$
$\alpha_1 = \Omega(n^{\frac{1}{2}})$	$\simeq \frac{\sqrt{n}}{\sqrt{s_1^2 - 2(\log(1-c) + c)n/\alpha_1^2}}$	$\simeq rac{\sqrt{n}}{2s_1}$	$\simeq \frac{\sqrt{n}}{s_1}$

TABLE 1 Divergence rates of \varkappa_L , \varkappa_W and \varkappa_R when M = 1

• (M = 2) For M = 2, we assume that the two diverging spikes are not equal, i.e., $d_1 = d_2 = 1$. In addition, for convenience of analysis, we assume that the two spikes have the same divergence rate, i.e., $\alpha_2 = k_2 \alpha_1$ with some $k_2 < 1$. The results are presented in Table 2. Since the power function of RLRT is relevant only to the largest spikes and not to the other spikes, \varkappa_R has the same result for M = 2 as it does for M = 1, that is $\varkappa_R \simeq \sqrt{n}/s_1$. Thus, we omit the results of \varkappa_R in Table 2. We can conclude from Table 2 that RLRT is asymptotically more powerful than CNTT whenever $\alpha_1 \to \infty$, because $1 + k_2^2 < 2$ and $s_2 \ge 0$. However, for CLRT, if $n = o(\alpha_1^2)$ and $(s_1^2 + k_2^2 s_2^2)/(1 + k_2)^2 < s_1^2$, then \varkappa_L could be larger than \varkappa_R . Since $s_k^2 = 2 + \beta_k \sum_{t=1}^p |u_{tk}|^4 + o(1)$, with suitable values of $\beta_x \ge -2$, $\sum_{t=1}^p |u_{tk}|^4 \in [1/p, 1]$ and $k_2 < 1$, the inequality $(s_1^2 + k_2^2 s_2^2)/(1 + k_2)^2 < s_1^2$ can be satisfied, such as choosing $\beta_x = 0$. This property indicates that the LSSs could exhibit higher asymptotic power than RLRT statistic in some special scenarios.

TABLE 2Divergence rates of \varkappa_L and \varkappa_W when M = 2 and $\alpha_2 = k_2 \alpha_1$

	3 8 E W	
	\varkappa_L	\varkappa_W
$\alpha_1 = o(n^{\frac{1}{4}})$	$\asymp \alpha_1$	$\asymp \alpha_1^2$
$\Omega(n^{\frac{1}{4}}) = \alpha_1 = o(n^{\frac{1}{2}})$	$\asymp \alpha_1$	$\simeq \frac{\sqrt{n}(1+k_2^2)}{2\sqrt{s_1^2 + k_2^4 s_2^2 + c^2(1+2c+\beta_x c)n/\alpha_1^4}}$
$\alpha_1 = \Omega(n^{\frac{1}{2}})$	$\simeq \frac{(1+k_2)\sqrt{n}}{\sqrt{s_1^2 + k_2^2 s_2^2 - 2(\log(1-c) + c)n/\alpha_1^2}}$	$\simeq \frac{\sqrt{n}(1+k_2^2)}{2\sqrt{s_1^2+k_2^4s_2^2}}$

• $(M \ge 3)$ For $M \ge 3$, the discussion is analogous to the cases of M = 1 and M = 2; thus, we omit the details because of space limitations. We merely wish to emphasize here that when $M \ge 3$, CNTT is also potentially asymptotically more powerful than RLRT. Suppose that $\alpha_t = k_t \alpha_1$, $t = 1, \ldots, M$ and $1 = k_1 > k_2 > \cdots > k_M > 0$. It is not difficult to find that if $n = o(\alpha_1^4)$, then $\varkappa_W \simeq \sqrt{n} \sum_{t=1}^M k_t^2 / \sqrt{4 \sum_{t=1}^M k_t^4 s_t^2}$, which can be larger than $\varkappa_R \simeq \sqrt{n}/s_1$ with suitable values of k_t and s_t , $t = 1, \ldots, M$. For example, if M = 3, we can choose k_2 and k_3 close to 1, while choosing s_2 and s_3 close to 0. Notably, if $\beta_x = 0$, e.g., $\{x_{ij}\}$ are Gaussian, then M must be at least 5 for $\varkappa_W > \varkappa_R$ to asymptotically hold.

For illustration, we present some graphs of the functions \varkappa_L , \varkappa_W and \varkappa_R in (4.10) with different numbers of diverging spikes in Figure 1.




Fig 1: Graphs of the functions \varkappa_L , \varkappa_W and \varkappa_R . We fix $(p, n, \xi, \beta_x) = (100, 300, 0.05, 0)$. The left panel shows the curves for M = 1. The middle panel shows the curves for M = 2 with $\alpha_2 = 0.9\alpha_1$. The right panel shows the curves for M = 5 with $\alpha_2 = 0.9\alpha_1$, $\alpha_3 = 0.85\alpha_1$, $\alpha_4 = 0.8\alpha_1$ and $\alpha_5 = 0.75\alpha_1$.

5. Numerical studies . In this section, we report short numerical studies as an illustration of our results. Our objective in the simulations is to examine the power analysis in subsection 4.2.

We examine the following three different distributions of x_{ij} :

 Dt_1 : $\{x_{ij}\}\$ are i.i.d. samples from a standard Gaussian population. Dt_2 : $\{x_{ij}\}\$ are i.i.d. samples from Gamma(4, 0.5) - 2. Dt_3 : $\{x_{ij}\}\$ are i.i.d. samples from Uniform population distribution $U[-\sqrt{3}, \sqrt{3}]$.

Note that in above settings, $\beta_x = 0, \frac{3}{2}, -\frac{6}{5}$, respectively.

In the current numerical studies, the null hypothesis is defined as $H_0: \Sigma = \mathbf{I}_p$. For the alternative hypothesis, we adopt the following six population covariance matrix structures:

$$H_{1}: \Sigma = \Lambda_{1} = diag(\alpha_{1}, \underbrace{1, 1, \dots, 1}_{p-1}).$$

$$H_{2}: \Sigma = \Lambda_{2} = diag(\alpha_{1}, \alpha_{2}, \underbrace{1, 1, \dots, 1}_{p-2}), \alpha_{2} = 0.9\alpha_{1}.$$

$$H_{3}: \Sigma = \Lambda_{3} = diag(\alpha_{1}, \dots, \alpha_{5}, \underbrace{1, 1, \dots, 1}_{p-5}), \alpha_{2} = 0.9\alpha_{1}, \alpha_{3} = 0.85\alpha_{1}, \alpha_{4} = 0.8\alpha_{1}, \alpha_{5} = 0.9\alpha_{1}.$$

 $0.75\alpha_1.$

*H*₄: $\Sigma = U_0 \Lambda_1 U_0^*$, where U_0 is the left singular vectors of a $p \times p$ random matrix with i.i.d. N(0, 1) entries

 $H_5: \Sigma = \mathbf{U}_0 \mathbf{\Lambda}_2 \mathbf{U}_0^*$, and \mathbf{U}_0 is defined in H_4 .

 $H_6: \Sigma = \mathbf{U}_0 \mathbf{\Lambda}_3 \mathbf{U}_0^*$, and \mathbf{U}_0 is defined in H_4 .

Note that in above settings, Σ is diagonal and $\mathcal{U}_{j_1j_1j_2j_2} = 1$ under H_1-H_3 , whereas Σ is nondiagonal and $\mathcal{U}_{j_1j_1j_2j_2} \approx 1/p$ under H_4-H_6 . The settings for the significance level ξ are constructed as follows: 0.05, 0.01, and 1 ×

The settings for the significance level ξ are constructed as follows: 0.05, 0.01, and 1×10^{-4} . The empirical results are obtained based on 10,000 replications with dimension p = 50,100,200, respectively. We set the RDS p/n = 1/3.

In Tables 3–6, we list the empirical sizes and powers of CLRT, CNTT, and RLRT under different settings. In the captions of these tables, " (Dt_*, H_*) " stands for the setting Dt_*, H_* . For the alternative hypothesis, due to space limitations, we present only some selected tables with significant properties in the paper, and the tables for other cases are provided in the supplementary material. Below are our conclusions based on our simulation studies:

- (1) For the null hypothesis, the empirical sizes of CLRT and CNTT are closer to significance levels than that of RLRT overall. This is because the rate of convergence for the largest eigenvalue distributions is slow. This property has been discussed extensively in the literature and we omit the details here. We refer interested readers to Johnstone (2001, 2008).
- (2) For the alternative hypothesis:
 - From Table 4, it is easy to find that RLRT has the highest asymptotic power under H₁. When there are two spikes, as seen from Table 5, CNTT and RLRT have higher asymptotic power than CLRT. As seen from Table 6, when there are five spikes, CNTT seems to have the highest asymptotic empirical power. This is consistent with our analysis in subsection 4.2 that when the number of spikes increases, CLRT and CNTT may exhibit higher asymptotic power than RLRT in some scenarios. To be noticed that, in Table 6, we only list the powers when ξ = 1 × 10⁻⁴ and consider smaller values of α₁ since the powers of three tests are all equal to 1 when ξ = 0.05 and ξ = 0.01, or when α₁ is not small enough, making comparisons infeasible.
 - From Tables 4–6, we can also find that in each row, the asymptotic power derived under (Dt_2, H_*) is smaller than that under (Dt_1, H_*) , and power derived under (Dt_3, H_*) is larger than that under (Dt_1, H_*) . This is because when Σ is diagonal, a smaller β_x may result in higher asymptotic power, corresponding to Theorems 4.3 and 4.4.

TABLE 3
Empirical probability of rejecting H_0 at significance levels 0.05 and 0.01 under assumptions of Gaussian,
Gamma, and Uniform distributions

	Dt_1 Dt_2 Dt_3											
		D	-	<i>D</i>		Dt_3						
test	(p,n)	$\xi = 0.05$	$\xi = 0.01$	$\xi = 0.05$	$\xi = 0.01$	$\xi = 0.05$	$\xi = 0.01$					
CLRT	(50,150)	0.0563	0.0132	0.0545	0.0123	0.0538	0.0128					
	(100,300)	0.0517	0.0105	0.0567	0.0120	0.0525	0.0102					
	(200,600)	0.0543	0.0117	0.0502	0.0110	0.0527	0.0101					
CNTT	(50,150)	0.0591	0.0138	0.0784	0.0219	0.0522	0.0107					
	(100,300)	0.0530	0.0106	0.0633	0.0175	0.0513	0.0112					
	(200,600)	0.0530	0.0122	0.0542	0.0142	0.0493	0.0117					
RLRT	(50,150)	0.0426	0.0089	0.1104	0.0328	0.0215	0.0033					
	(100,300)	0.0459	0.0082	0.0969	0.0260	0.0276	0.0046					
	(200,600)	0.0495	0.0095	0.0779	0.0170	0.0279	0.0068					

TABLE 4

Empirical probability of rejecting H_1 at significance level $\xi = 0.05$ under assumptions of Gaussian, Gamma, and Uniform distributions

			(Dt_1, H_1)			(Dt_2, H_1)		(Dt_3, H_1)		
test	(p,n)	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$
CLRT	(50,150)	0.7236	0.9989	1	0.6920	0.9981	1	0.7511	1	1
	(100,300)	0.7439	0.9999	1	0.7251	0.9998	1	0.7499	1	1
	(200,600)	0.7544	0.9999	1	0.7468	0.9998	1	0.7632	1	1
CNTT	(50,150)	0.9919	1	1	0.9702	1	1	0.9996	1	1
	(100,300)	0.9990	1	1	0.9898	1	1	1	1	1
	(200,600)	0.9998	1	1	0.9973	1	1	1	1	1
RLRT	(50,150)	0.9998	1	1	0.9988	1	1	1	1	1
	(100,300)	1	1	1	1	1	1	1	1	1
	(200,600)	1	1	1	1	1	1	1	1	1

CLT FOR LSS WITH DIVERGING SPIKES

1	F ,	, .jj	-	and Unifor	m distribut					
		(Dt_1, H_2)				(Dt_2, H_2)		(Dt_3, H_2)		
test	(p,n)	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$
\overline{CLRT}	(50,150)	0.9811	1	1	0.9633	1	1	0.9899	1	1
	(100,300)	0.9894	1	1	0.9835	1	1	0.9929	1	1
	(200,600)	0.9928	1	1	0.9894	1	1	0.9948	1	1
CNTT	(50,150)	1	1	1	1	1	1	1	1	1
	(100,300)	1	1	1	1	1	1	1	1	1
	(200,600)	1	1	1	1	1	1	1	1	1
RLRT	(50,150)	1	1	1	1	1	1	1	1	1
	(100,300)	1	1	1	1	1	1	1	1	1
	(200,600)	1	1	1	1	1	1	1	1	1

TABLE 5 Empirical probability of rejecting H_2 at significance level $\xi = 0.05$ under assumptions of Gaussian, Gamma,

TABLE 6

Empirical probability of rejecting H_3 at significance level $\xi = 1 \times 10^{-4}$ under assumptions of Gaussian, Gamma, and Uniform distributions

			(Dt_1, H_3))		(Dt_2, H_3))	(Dt_3, H_3)		
						α_1				
test	(p,n)	2.2	2.5	2.8	2.2	2.5	2.8	2.2	2.5	2.8
\overline{CLRT}	(50,150)	0.3911	0.8636	0.9918	0.3883	0.8232	0.9834	0.3825	0.8945	0.9976
	(100,300)	0.3876	0.8946	0.9985	0.3804	0.8705	0.9951	0.3861	0.9110	0.9985
	(200,600)	0.3779	0.9032	0.9980	0.3846	0.8946	0.9976	0.3825	0.9134	0.9997
CNTT	(50,150)	0.9972	1	1	0.9972	0.9998	1	0.9999	1	1
	(100,300)	0.9997	1	1	0.9917	1	1	1	1	1
	(200,600)	1	1	1	0.9964	1	1	1	1	1
RLRT	(50,150)	0.8933	0.9968	1	0.8954	0.9943	0.9999	0.8909	0.9994	1
	(100,300)	0.9821	1	1	0.9756	0.9999	1	0.9904	1	1
	(200,600)	0.9996	1	1	0.9989	1	1	1	1	1

6. Technical proofs. In this section, we present some lemmas that are needed in the proofs of the main results. The truncation and renormalization are postponed to the end of this paper.

6.1. Some primary definitions and lemmas. In this section, we provide some useful results that are used later in the proofs of Theorems 3.1 and 3.2. For the population covariance matrix $\Sigma = TT^*$, we consider the corresponding sample covariance matrix $B = TS_xT^*$, where $\mathbf{S}_x = \frac{1}{n} \mathbf{X} \mathbf{X}^*$. By singular value decomposition of T (see (2.3)),

$$\mathbf{B} = \mathbf{V} \begin{pmatrix} \mathbf{D}_{1}^{\frac{1}{2}} \mathbf{U}_{1}^{*} \mathbf{S}_{x} \mathbf{U}_{1} \mathbf{D}_{1}^{\frac{1}{2}}, \mathbf{D}_{1}^{\frac{1}{2}} \mathbf{U}_{1}^{*} \mathbf{S}_{x} \mathbf{U}_{2} \mathbf{D}_{2}^{\frac{1}{2}} \\ \mathbf{D}_{2}^{\frac{1}{2}} \mathbf{U}_{2}^{*} \mathbf{S}_{x} \mathbf{U}_{1} \mathbf{D}_{1}^{\frac{1}{2}}, \mathbf{D}_{2}^{\frac{1}{2}} \mathbf{U}_{2}^{*} \mathbf{S}_{x} \mathbf{U}_{2} \mathbf{D}_{2}^{\frac{1}{2}} \end{pmatrix} \mathbf{V}^{*}.$$

Note that

$$\mathbf{S} = \begin{pmatrix} \mathbf{D}_{1}^{\frac{1}{2}} \mathbf{U}_{1}^{*} \mathbf{S}_{x} \mathbf{U}_{1} \mathbf{D}_{1}^{\frac{1}{2}}, \mathbf{D}_{1}^{\frac{1}{2}} \mathbf{U}_{1}^{*} \mathbf{S}_{x} \mathbf{U}_{2} \mathbf{D}_{2}^{\frac{1}{2}} \\ \mathbf{D}_{2}^{\frac{1}{2}} \mathbf{U}_{2}^{*} \mathbf{S}_{x} \mathbf{U}_{1} \mathbf{D}_{1}^{\frac{1}{2}}, \mathbf{D}_{2}^{\frac{1}{2}} \mathbf{U}_{2}^{*} \mathbf{S}_{x} \mathbf{U}_{2} \mathbf{D}_{2}^{\frac{1}{2}} \end{pmatrix} \triangleq \begin{pmatrix} \mathbf{S}_{11}, \mathbf{S}_{12} \\ \mathbf{S}_{21}, \mathbf{S}_{22} \end{pmatrix}$$

Moreover, B and S have the same eigenvalues.

Recall that $\underline{\mathbf{B}} = \frac{1}{n} \mathbf{X}^* \mathbf{T}^* \mathbf{T} \mathbf{X}$ (the spectrum of which differs from that of \mathbf{B} by |n-p| zeros). Its LSD is $\underline{F}^{c,H}$, $\underline{F}^{c,H} \equiv (1-c) \mathbb{1}_{[0,\infty)} + cF^{c,H}$, and its Stieltjes transform is $\underline{m}(z)$. Let $\widetilde{\lambda}_j$ be the eigenvalues of \mathbf{S}_{22} so that the LSS of \mathbf{S}_{22} is $\sum_{j=1}^{p-M} f(\widetilde{\lambda}_j)$. Correspondingly, recall that $c_{nM} := \frac{p-M}{n}$, $H_{2n} := F^{\mathbf{D}_2}$ and $m_{2n0} := m_{2n0}(z)$ is the Stieltjes transform of $F^{c_{nM},H_{2n}}$. First, in Lemma 6.1 we derive that the difference between the two centers is 0.

LEMMA 6.1. Under Assumptions 1–4,

$$(p-M)\int f(x) dF^{c_{nM},H_{2n}} = p\int f(x) dF^{c_n,H_n}(x)$$

PROOF. By the Cauchy integral formula,

$$p\int f(x)dF^{c_n,H_n} = -\frac{p}{2\pi i}\oint_{\mathcal{C}} f(z)m_{1n0}dz = -\frac{n}{2\pi i}\oint_{\mathcal{C}} f(z)\underline{m}_{1n0}dz,$$
$$(p-M)\int f(x)dF^{c_{nM},H_{2n}} = -\frac{p-M}{2\pi i}\oint_{\mathcal{C}} f(z)m_{2n0}dz = -\frac{n}{2\pi i}\oint_{\mathcal{C}} f(z)\underline{m}_{2n0}dz.$$

where \underline{m}_{1n0} and \underline{m}_{2n0} are the Stieltjes transforms of \underline{F}^{c_n,H_n} and $\underline{F}^{c_{nM},H_{2n}}$, respectively. Then,

$$(p-M)\int f(x)dF^{c_{nM},H_{2n}} - p\int f(x)dF^{c_n,H_n} = \frac{n}{2\pi i}\oint_{\mathcal{C}} f(z)\left(\underline{m}_{1n0} - \underline{m}_{2n0}\right)dz.$$

Next, we prove that $\underline{m}_{1n0} = \underline{m}_{2n0}$.

Note that m_{1n0} and m_{2n0} are the unique solutions to

(6.11)
$$z = -\frac{1}{\underline{m}_{1n0}} + c_n \int \frac{t dH_n(t)}{1 + t\underline{m}_{1n0}}$$

(6.12)
$$z = -\frac{1}{\underline{m}_{2n0}} + c_{nM} \int \frac{t dH_{2n}(t)}{1 + t \underline{m}_{2n0}}$$

respectively, where $\underline{m}_{1n0} = -\frac{1-c_n}{z} + c_n m_{1n0}$ and $\underline{m}_{2n0} = -\frac{1-c_{nM}}{z} + c_{nM} m_{2n0}$. Since

$$H_n(t) = \frac{1}{p} \left[\sum_{i=1}^M \mathbb{1}_{\{0 \le t\}} + \sum_{i=M+1}^p \mathbb{1}_{\{\alpha_i \le t\}} \right] = \frac{M}{p} + \frac{1}{p} \sum_{i=M+1}^p \mathbb{1}_{\{\alpha_i \le t\}}$$

and $H_{2n}(t) = \frac{1}{p-M} \sum_{i=M+1}^{p} \mathbb{1}_{\{\alpha_i \le t\}}$, (6.11) can be written as

$$z = -\frac{1}{\underline{m}_{1n0}} + \frac{p}{n} \int \frac{td\left(\frac{M}{p} + \frac{1}{p}\sum_{i=M+1}^{p} \mathbb{1}_{\{\alpha_i \le t\}}(t)\right)}{1 + t\underline{m}_{1n0}} = -\frac{1}{\underline{m}_{1n0}} + \frac{p}{n} \int \frac{td\left(\frac{1}{p}\sum_{i=M+1}^{p} \mathbb{1}_{\{\alpha_i \le t\}}(t)\right)}{1 + t\underline{m}_{1n0}}$$

(6.13)

$$= -\frac{1}{\underline{m}_{1n0}} + \frac{1}{n} \sum_{i=M+1}^{p} \frac{\alpha_i}{1 + \alpha_i \underline{m}_{1n0}}.$$

Similarly, equation (6.12) can be written as $z = -\frac{1}{\underline{m}_{2n0}} + \frac{1}{n} \sum_{i=M+1}^{p} \frac{\alpha_i}{1 + \alpha_i \underline{m}_{2n0}}$. Thus, according to the fact that m_{1n0} and m_{2n0} are the unique solutions of (6.13) and (6.1), respectively, we have $m_{1n0} = m_{2n0}$, which completes the proof of this lemma.

Note that for the bounded part of the LSS, $\sum_{j=M+1}^{p} f(\lambda_j)$, the BST cannot be used directly since it is not an LSS of a sample covariance matrix. In fact, it approximates the LSS of \mathbf{S}_{22} , that is $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$, but they are not equal since the off-diagonal blocks of the sample covariance matrix are not null. The following lemma measures the difference between $\sum_{j=M+1}^{p} f(\lambda_j)$ and $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$.

16

LEMMA 6.2. Under Assumptions 1–4,

$$\sum_{j=M+1}^{p} f(\lambda_{j}) - \sum_{j=1}^{p-M} f\left(\tilde{\lambda}_{j}\right) - \frac{M}{2\pi i} \oint_{\mathcal{C}} f(z) \frac{\underline{m}_{2n0}'(z)}{\underline{m}_{2n0}(z)} dz = o_{p}(1).$$

PROOF. Note that $L_1 := \sum_{j=M+1}^p f(\lambda_j)$. By the Cauchy integral formula, we have $L_1 = -\frac{p}{2\pi i} \oint_{\mathcal{C}} f(z) m_n(z) dz$, where $m_n = \frac{1}{p} \operatorname{tr} (\mathbf{S} - z\mathbf{I}_p)^{-1} = \frac{1}{p} \operatorname{tr} (\mathbf{B} - z\mathbf{I}_p)^{-1}$. Analogously, we have $L_2 := \sum_{j=1}^{p-M} f(\widetilde{\lambda}_j) = -\frac{p-M}{2\pi i} \oint_{\mathcal{C}} f(z) m_{2n}(z) dz$, where $m_{2n} = \frac{1}{p-M} \operatorname{tr} (\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-1}$. By applying the block matrix inversion formula to m_n , we can obtain

(6.14)
$$L_1 - L_2 = -\frac{1}{2\pi i} \oint_{\mathcal{C}} f(z) \left(T_1 - T_2\right) dz,$$

where

$$T_{1} = \operatorname{tr} \left(\mathbf{S}_{11} - z \mathbf{I}_{M} - \mathbf{S}_{12} \left(\mathbf{S}_{22} - z \mathbf{I}_{p-M} \right)^{-1} \mathbf{S}_{21} \right)^{-1},$$

$$T_{2} = -\operatorname{tr} \left[\left(\mathbf{S}_{11} - z \mathbf{I}_{M} - \mathbf{S}_{12} \left(\mathbf{S}_{22} - z \mathbf{I}_{p-M} \right)^{-1} \mathbf{S}_{21} \right)^{-1} \mathbf{S}_{12} \left(\mathbf{S}_{22} - z \mathbf{I}_{p-M} \right)^{-2} \mathbf{S}_{21} \right].$$

Note that for any matrix **Z**,

$$\mathbf{Z} \left(\mathbf{Z}^* \mathbf{Z} - \lambda \mathbf{I} \right)^{-1} \mathbf{Z}^* = \mathbf{I} + \lambda \left(\mathbf{Z} \mathbf{Z}^* - \lambda \mathbf{I} \right)^{-1}$$

which, together with the notation $\Upsilon_n := \frac{1}{n} \mathbf{D}_1^{\frac{1}{2}} \mathbf{U}_1^* \mathbf{X} \left(\frac{1}{n} \mathbf{X}^* \mathbf{U}_2 \mathbf{D}_2 \mathbf{U}_2^* \mathbf{X} - z \mathbf{I}_n \right)^{-1} \mathbf{X}^* \mathbf{U}_1 \mathbf{D}_1^{\frac{1}{2}}$, implies that

$$T_{1} = -z^{-1} \operatorname{tr} \left(\mathbf{I}_{M} + \boldsymbol{\Upsilon}_{n} \right)^{-1}$$
$$T_{2} = z^{-1} \operatorname{tr} \left[\left(\mathbf{I}_{M} + \boldsymbol{\Upsilon}_{n} \right)^{-1} \mathbf{S}_{12} \left(\mathbf{S}_{22} - z \mathbf{I}_{p-M} \right)^{-2} \mathbf{S}_{21} \right]$$

 $\underline{m}_{2n} = \underline{m}_{2n}(z)$ denotes the Stieltjes transform of $F_n^{\frac{1}{n}\mathbf{X}^*\mathbf{U}_2\mathbf{D}_2\mathbf{U}_2^*\mathbf{X}}$. Thus, we have that $\underline{m}_{2n}(z) - \underline{m}(z) = o_p(1)$ for any $z \in \mathcal{C}$. From Theorem 3.1 of (Jiang and Bai, 2021), we know that

(6.15)
$$\frac{1}{n}\mathbf{U}_{1}^{*}\mathbf{X}\left(\frac{1}{n}\mathbf{X}^{*}\mathbf{U}_{2}\mathbf{D}_{2}\mathbf{U}_{2}^{*}\mathbf{X}-z\mathbf{I}_{n}\right)^{-1}\mathbf{X}^{*}\mathbf{U}_{1}=\underline{m}_{2n}\left(z\right)\mathbf{I}_{M}+O_{p}\left(n^{-\frac{1}{2}}\right).$$

Thus, under Assumption 3, we find that

(6.16)
$$\mathbf{D}_{1}^{1/2} \left(\mathbf{I}_{M} + \boldsymbol{\Upsilon}_{n} \right)^{-1} \mathbf{D}_{1}^{1/2} = \frac{1}{\underline{m}(z)} \mathbf{I}_{M} + o_{p}(1)$$

which yields

(6.17)
$$T_1 = o_p(1).$$

It follows that

$$\mathbf{D}_{1}^{-1/2} \mathbf{S}_{12} (\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-2} \mathbf{S}_{21} \mathbf{D}_{1}^{-1/2}$$

$$= \frac{1}{n} \operatorname{tr} \left[(\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-2} \mathbf{S}_{22} \right] \mathbf{I}_{M} + O_{p} (n^{-\frac{1}{2}})$$

$$= \frac{1}{n} \operatorname{tr} (\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-1} \mathbf{I}_{M} + \frac{z}{n} \operatorname{tr} (\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-2} \mathbf{I}_{M} + O_{p} (n^{-\frac{1}{2}})$$

$$= cm_{2n} (z) \mathbf{I}_{M} + zcm'_{2n} (z) \mathbf{I}_{M} + o_{p} (1)$$

$$= cm_{2n0} (z) \mathbf{I}_{M} + zcm'_{2n0} (z) \mathbf{I}_{M} + o_{p} (1)$$
(6.18)
$$= \underline{m}_{2n0} (z) \mathbf{I}_{M} + z\underline{m}'_{2n0} (z) \mathbf{I}_{M} + o_{p} (1),$$

where the last equality is derived from $\underline{m}_{2n0}(z) = -\frac{1-c}{z} + cm_{2n0}(z)$. Therefore, according to (6.16) and (6.18), we obtain

$$T_{2} = M \frac{\underline{m}_{2n0}(z) + z\underline{m}'_{2n0}(z)}{z\underline{m}_{2n0}(z)} + o_{p}(1),$$

which, together with (6.14) and (6.17), implies that

$$L_{1} - L_{2} = \frac{M}{2\pi i} \oint_{\mathcal{C}} f(z) \frac{\underline{m}_{2n0}(z) + z\underline{m}_{2n0}'(z)}{z\underline{m}_{2n0}(z)} dz + o_{p}(1) = \frac{M}{2\pi i} \oint_{\mathcal{C}} f(z) \frac{\underline{m}_{2n0}'(z)}{\underline{m}_{2n0}(z)} dz + o_{p}(1).$$

Therefore, the proof of this lemma is complete.

Therefore, the proof of this lemma is complete.

Define random vector $\boldsymbol{\gamma}_k = (\gamma_{kj})^\top = \left(\sqrt{n}\frac{\lambda_j - \phi_n(\alpha_k)}{\phi_n(\alpha_k)}, j \in J_k\right)^\top$, where J_k is the indicator set of a packet of d_k consecutive sample eigenvalues. Then, we present the following lemma, which is borrowed from Jiang and Bai (2021) and characterizes the limiting distribution of the spiked eigenvalues of the sample covariance matrix.

LEMMA 6.3. (Jiang and Bai (2021)) Under Assumptions 1–4, random vector γ_k converges weakly to the joint distribution of d_k eigenvalues of a Gaussian random matrix

$$-rac{1}{ heta_k}\left[oldsymbol{\Omega}_{\phi_k}
ight]_{kk},$$

where

$$\theta_k = \phi_k^2 \underline{m}_2(\phi_k), \ \underline{m}_2(\lambda) = \int \frac{1}{(\lambda - x)^2} d\underline{F}^{c,H}(x)$$

with $\underline{F}^{c,H}$ being the LSD of matrix $n^{-1}\mathbf{X}^*\mathbf{U}_2\mathbf{D}_2\mathbf{U}_2^*\mathbf{X}$, $\phi_k = \alpha_k \left(1 + c\int \frac{t}{\alpha_k - t} dH(t)\right)$. $[\mathbf{\Omega}_{\phi_k}]_{kk}$ is the kth diagonal block of matrix $\mathbf{\Omega}_{\phi_k}$. The variances and covariances of the elements ω_{ij} of Ω_{ϕ_k} are:

$$\operatorname{Cov}(\omega_{i_{1},j_{1}},\omega_{i_{2},j_{2}}) = \begin{cases} (\alpha_{x}+1)\theta_{k} + \beta_{x}\mathcal{U}_{iiii}\nu_{k}, & i_{1}=j_{1}=i_{2}=j_{2}=i\\ \theta_{k} + \beta_{x}\mathcal{U}_{ijj}\nu_{k}, & i_{1}=i_{2}=i \neq j_{1}=j_{2}=j\\ \beta_{x}\mathcal{U}_{i_{1}j_{1}i_{2}j_{2}}\nu_{k}, & other \ cases \end{cases}$$

where $\beta_x \mathcal{U}_{i_1 j_1 i_2 j_2} = \sum_{t=1}^p \bar{u}_{t i_1} u_{t j_1} u_{t i_2} \bar{u}_{t j_2} \beta_x$, $\boldsymbol{u}_i = (u_{1i}, \dots, u_{pi})^\top$ are the *i* th column of the matrix \mathbf{U}_1 , $\nu_k = \phi_k^2 \underline{m}^2(\phi_k)$.

Recall that λ_j are the eigenvalues of **B**, and $\tilde{\lambda}_j$ are the eigenvalues of **S**₂₂. The following lemma shows the asymptotic independence between $\sum_{j=1}^{M} f(\lambda_j)$ and $\sum_{j=1}^{p-M} f(\widetilde{\lambda}_j)$.

LEMMA 6.4. Under Assumptions 1–4, $\sum_{j=1}^{M} f(\lambda_j)$ and $\sum_{j=1}^{p-M} f(\widetilde{\lambda}_j)$ are asymptotically independent.

PROOF. It is sufficient to prove that for a given $\sum_{j=1}^{p-M} f(\widetilde{\lambda}_j)$, the asymptotic limiting distribution of $\sum_{j=1}^{M} f(\lambda_j)$ does not depend on the random part of $\sum_{j=1}^{p-M} f(\widetilde{\lambda}_j)$, that is, it only depends on its limit.

First, we consider f(x) = x. From the proof of Theorem 3.1 in Jiang and Bai (2021), we have the following determinant equation

$$0 = \left| \left[\mathbf{\Omega}_M \left(\phi_k \right) \right]_{kk} + \lim \gamma_{kj} \left\{ \phi_k^2 \underline{m}_2(\phi_k) \right\} \mathbf{I}_{d_k} \right|,$$

where $\Omega_M(\phi_k)$

(6.

$$=\frac{\phi_k}{\sqrt{n}}\left[\operatorname{tr}\left\{\left(\phi_k\mathbf{I}-\frac{1}{n}\mathbf{X}^*\mathbf{U}_2\mathbf{D}_2\mathbf{U}_2^*\mathbf{X}\right)^{-1}\right\}\mathbf{I}-\mathbf{U}_1^*\mathbf{X}\left(\phi_k\mathbf{I}-\frac{1}{n}\mathbf{X}^*\mathbf{U}_2\mathbf{D}_2\mathbf{U}_2^*\mathbf{X}\right)^{-1}\mathbf{X}^*\mathbf{U}_1\right],$$

and $\underline{m}_2(\phi_k)$ is the limit of tr $(\phi_k \mathbf{I} - \frac{1}{n} \mathbf{X}^* \mathbf{U}_2 \mathbf{D}_2 \mathbf{U}_2^* \mathbf{X})^{-2}$. Then, we know that γ_{kj} has the same asymptotic distribution with eigenvalues of $-\frac{[\mathbf{\Omega}_M(\phi_k)]_{kk}}{\phi_k^2 \underline{m}_2(\phi_k)}$ in order. From Jiang and Bai (2021), we can obtain that the limiting distribution of $\mathbf{\Omega}_M(\phi_k)$ does not change if $\mathbf{U}_2^* \mathbf{X}$ is replaced by $\mathbf{U}_2^* \mathbf{Y}$ while $\mathbf{U}_1^* \mathbf{X}$ remains unchanged. Here \mathbf{Y} and \mathbf{X} are i.i.d.. Therefore in $\mathbf{\Omega}_M(\phi_k)$, we can assume that $\mathbf{U}_1^* \mathbf{X}$ and $\mathbf{U}_2^* \mathbf{X}$ are independent without loss of generality. Then, given $\mathbf{U}_2^* \mathbf{X}$, the limiting distribution of γ_{kj} only depends on the limit of tr $(\phi_k \mathbf{I} - \frac{1}{n} \mathbf{X}^* \mathbf{U}_2 \mathbf{D}_2 \mathbf{U}_2^* \mathbf{X})^{-1}$, that is, $\underline{m}_2(\phi_k)$, and has nothing to do with the random part. Therefore, it is found that $\sum_{j=1}^M f(\lambda_j)$ and $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$ are asymptotically independent when f(x) = x.

When $f(x) \neq x$, by using the Newton-Leibniz formula, we have

$$\sum_{j=1}^{M} f(\lambda_{j}) - \sum_{k=1}^{K} d_{k}f(\phi_{n}(\alpha_{k})) = \sum_{k=1}^{K} \sum_{j \in J_{k}} (f(\lambda_{j}) - f(\phi_{n}(\alpha_{k})))$$

$$= \sum_{k=1}^{K} \sum_{j \in J_{k}} \int_{0}^{\frac{\phi_{n}(\alpha_{k})}{\sqrt{n}}} \gamma_{kj} f'(t + \phi_{n}(\alpha_{k})) dt$$

$$= \sum_{k=1}^{K} \sum_{j \in J_{k}} \int_{0}^{1} \frac{\phi_{n}(\alpha_{k})}{\sqrt{n}} \gamma_{kj} \frac{f'\left(\phi_{n}(\alpha_{k})\left(1 + \frac{\gamma_{kj}}{\sqrt{n}}s\right)\right)}{f'(\phi_{n}(\alpha_{k}))} f'(\phi_{n}(\alpha_{k})) ds$$

$$= \sum_{k=1}^{K} \sum_{j \in J_{k}} \int_{0}^{1} \varpi_{nk} \gamma_{kj} \frac{f'\left(\phi_{n}(\alpha_{k})\left(1 + \frac{\gamma_{kj}}{\sqrt{n}}s\right)\right)}{f'(\phi_{n}(\alpha_{k}))} ds$$

$$19) \qquad \rightarrow \sum_{k=1}^{K} \sum_{j \in J_{k}} \int_{0}^{1} \gamma_{kj} \varpi_{nk} ds = \sum_{k=1}^{K} \sum_{j \in J_{k}} \varpi_{nk} \gamma_{kj},$$

where (6.19) is true due to Assumption 4, and we denote $\varpi_{nk} = \frac{\phi_n(\alpha_k)}{\sqrt{n}} f'(\phi_n(\alpha_k))$. Thus, we convert the above equation into a function of γ_{kj} . The above calculations represent the underlying idea of the generalized delta method we mentioned in the Introduction. Since we have proven above that given $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$, the limiting distribution of γ_{kj} is concerned only with the limit of $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$, as is $\sum_{k=1}^{K} \sum_{j \in J_k} \varpi_{nk} \gamma_{kj}$, accordingly, we can conclude that $\sum_{j=1}^{M} f(\lambda_j)$ and $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$ are asymptotically independent. The proof is complete.

In the following lemma, we derive the asymptotic distribution of the LSS generated from submatrix S_{22} .

LEMMA 6.5. Define $Q_1 = \sum_{j=1}^{p-M} f_1(\widetilde{\lambda}_j) - (p-M) \int f_1(x) dF^{c_{nM},H_{2n}}$; then, under Assumptions 1–4, we have

$$\kappa_1^{-1}(Q_1 - \mu_1) \xrightarrow{d} N(0, 1)$$

with mean function

$$\begin{split} \mu_1 &= -\frac{\alpha_x}{2\pi i} \cdot \oint_{\mathcal{C}} f_1(z) \frac{c_{nM} \int \underline{m}_{2n0}^3(z) t^2 \left(1 + t\underline{m}_{2n0}(z)\right)^{-3} dH_{2n}(t)}{\left(1 - c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t\underline{m}_{2n0}(z))^2} dH_{2n}(t)\right) \left(1 - \alpha_x c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t\underline{m}_{2n0}(z))^2} dH_{2n}(t)\right)} dz \\ &- \frac{\beta_x}{2\pi i} \cdot \oint_{\mathcal{C}} f_1(z) \frac{c_{nM} \int \underline{m}_{2n0}^3(z) t^2 \left(1 + t\underline{m}_{2n0}(z)\right)^{-3} dH_{2n}(t)}{1 - c_{nM} \int \underline{m}_{2n0}^2(z) t^2 \left(1 + t\underline{m}_{2n0}(z)\right)^{-3} dH_{2n}(t)} dz, \end{split}$$

and the covariance function is

$$\kappa_{1}^{2} = -\frac{1}{4\pi^{2}} \oint_{\mathcal{C}_{1}} \oint_{\mathcal{C}_{2}} f_{1}(z_{1}) f_{1}(z_{2}) \vartheta_{n}^{2} dz_{1} dz_{2},$$

where $\vartheta_n^2 = \Theta_{0,n}(z_1, z_2) + \alpha_x \Theta_{1,n}(z_1, z_2) + \beta_x \Theta_{2,n}(z_1, z_2),$ $\Theta_{0,n}(z_1, z_2) = \frac{\underline{m}'_{2n0}(z_1)\underline{m}'_{2n0}(z_2)}{(\underline{m}_1 - (\underline{z}_1) - \underline{m}_2 - (\underline{z}_1))^2} - \frac{1}{(\underline{m}_1 - \underline{z}_2)^2},$

$$\Theta_{0,n}(z_1, z_2) = \frac{\partial}{\partial z_2} \left\{ \frac{\partial \mathcal{A}_n(z_1, z_2)}{\partial z_1} \frac{1}{1 - \alpha_x \mathcal{A}_n(z_1, z_2)} \right\},$$

$$\mathcal{A}_n(z_1, z_2) = \frac{z_1 z_2}{n} \underline{m}_{2n0}(z_1) \underline{m}_{2n0}(z_2) \operatorname{tr} \mathbf{\Gamma}^* \mathbf{P}_n(z_1) \mathbf{\Gamma} \mathbf{\Gamma}^\top \mathbf{P}_n(z_2)^\top \bar{\mathbf{\Gamma}},$$

$$\Theta_{2,n}(z_1, z_2) = \frac{z_1^2 z_2^2 \underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{n} \sum_{i=1}^p \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_1) \mathbf{\Gamma} \right]_{ii} \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_2) \mathbf{\Gamma} \right]_{ii},$$

and the definitions of \mathbf{P}_n , Γ , and \underline{m}_{2n0} are defined in Section 3.

PROOF. From Zheng et al. (2015), we have that under Assumptions 1–4, the random variable $(\kappa_1^0)^{-1} (Q_1 - \mu_1) \xrightarrow{d} N(0, 1)$, with mean function

$$\begin{split} \mu_{1} &= -\frac{\alpha_{x}}{2\pi i} \cdot \oint_{\mathcal{C}} \frac{f_{1}(z)c_{nM} \int \underline{m}_{2n0}^{3}(z)t^{2} \left(1 + t\underline{m}_{2n0}(z)\right)^{-3} dH_{2n}(t)}{\left(1 - c_{nM} \int \frac{\underline{m}_{2n0}^{2}(z)t^{2}}{(1 + t\underline{m}_{2n0}(z))^{2}} dH_{2n}(t)\right) \left(1 - \alpha_{x}c_{nM} \int \frac{\underline{m}_{2n0}^{2}(z)t^{2}}{(1 + t\underline{m}_{2n0}(z))^{2}} dH_{2n}(t)\right)} dz \\ &- \frac{\beta_{x}}{2\pi i} \cdot \oint_{\mathcal{C}} \frac{f_{1}(z)c_{nM} \int \underline{m}_{2n0}^{3}(z)t^{2} \left(1 + t\underline{m}_{2n0}(z)\right)^{-3} dH_{2n}(t)}{1 - c_{nM} \int \underline{m}_{2n0}^{2}(z)t^{2} \left(1 + t\underline{m}_{2n0}(z)\right)^{-2} dH_{2n}(t)} dz, \end{split}$$

and the covariance function is

$$\left(\kappa_{1}^{0}\right)^{2} = -\frac{1}{4\pi^{2}} \oint_{\mathcal{C}_{1}} \oint_{\mathcal{C}_{2}} f_{1}\left(z_{1}\right) f_{1}\left(z_{2}\right) \left(\vartheta_{n}^{0}\right)^{2} dz_{1} dz_{2},$$

where

$$\begin{aligned} (\vartheta_n^0)^2 &= \frac{b_n\left(z_1\right)b_n\left(z_2\right)}{n^2} \sum_{j=1}^n \operatorname{tr} \mathbb{E}_j \Gamma \Gamma^* \mathbf{A}_j^{-1}\left(z_1\right) \mathbb{E}_j \left(\Gamma \Gamma^* \mathbf{A}_j^{-1}\left(z_2\right) \right) \\ &+ \frac{\alpha_x b_n\left(z_1\right)b_n\left(z_2\right)}{n^2} \sum_{j=1}^n \operatorname{tr} \mathbb{E}_j \Gamma^* \mathbf{A}_j^{-1}\left(z_1\right) \Gamma \mathbb{E}_j \left(\Gamma^\top \left(\mathbf{A}_j^\top\right)^{-1}\left(z_2\right) \bar{\Gamma} \right) \\ &+ \frac{\beta_x b_n\left(z_1\right)b_n\left(z_2\right)}{n^2} \sum_{j=1}^n \sum_{i=1}^p \boldsymbol{e}_i^\top \Gamma^* \mathbf{A}_j^{-1}\left(z_1\right) \Gamma \boldsymbol{e}_i \cdot \boldsymbol{e}_i^\top \Gamma^* \mathbf{A}_j^{-1}\left(z_2\right) \Gamma \boldsymbol{e}_i, \end{aligned}$$

where $b_n(z) = \frac{1}{1+n^{-1}\mathbb{E}\mathrm{tr}\mathbf{\Gamma}\mathbf{\Gamma}^*\mathbf{A}_j^{-1}(z)}$. The symbols $\mathbf{A}_j, \mathbf{e}_i$ are defined in Section 2. Moreover Najim and Yao (2016) provided an estimation ϑ_n^2 for $(\vartheta_n^0)^2$ and proved that $(\vartheta_n^0)^2$ is close to ϑ_n^2 in the Lévy–Prohorov distance, where $\vartheta_n^2 = \Theta_{0,n}(z_1, z_2) + \alpha_x \Theta_{1,n}(z_1, z_2) + \beta_x \Theta_{2,n}(z_1, z_2)$,

$$\begin{split} \Theta_{0,n}(z_1, z_2) &= \frac{\underline{m}'_{2n0}(z_1)\underline{m}'_{2n0}(z_2)}{(\underline{m}_{2n0}(z_1) - \underline{m}_{2n0}(z_2))^2} - \frac{1}{(z_1 - z_2)^2}, \\ \Theta_{1,n}(z_1, z_2) &= \frac{\partial}{\partial z_2} \left\{ \frac{\partial \mathcal{A}_n(z_1, z_2)}{\partial z_1} \frac{1}{1 - \alpha_x \mathcal{A}_n(z_1, z_2)} \right\}, \\ \mathcal{A}_n(z_1, z_2) &= \frac{z_1 z_2}{n} \underline{m}_{2n0}(z_1) \underline{m}_{2n0}(z_2) \mathrm{tr} \mathbf{\Gamma}^* \mathbf{P}_n(z_1) \mathbf{\Gamma} \mathbf{\Gamma}^\top \mathbf{P}_n(z_2)^\top \overline{\mathbf{\Gamma}}, \\ \Theta_{2,n}(z_1, z_2) &= \frac{z_1^2 z_2^2 \underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{n} \sum_{i=1}^p \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_1) \mathbf{\Gamma} \right]_{ii} \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_2) \mathbf{\Gamma} \right]_{ii}, \end{split}$$

and the definitions of \mathbf{P}_n , $\mathbf{\Gamma}$, and \underline{m}_{2n0} are given in Section 3. Notably, if $\mathbf{\Gamma}$ is not real, then the convergence of $\Theta_{1,n}(z_1, z_2)$ is not guaranteed. However, if $\mathbf{\Gamma}$ and entries x_{ij} are real, that is, $\alpha_x = 1$, then it can be easily proven that $\Theta_{0,n}(z_1, z_2) = \Theta_{1,n}(z_1, z_2)$. Similarly, the convergence of $\Theta_{2,n}(z_1, z_2)$ depends on the assumption that $\mathbf{\Gamma}^*\mathbf{\Gamma}$ is diagonal; thus, under Assumptions 1–4, $\Theta_{1,n}(z_1, z_2)$ and $\Theta_{2,n}(z_1, z_2)$ may have no limits.

Therefore, the covariance term $(\kappa_1^0)^2$ is estimable, and the estimate is κ_1^2 , with

$$\kappa_1^2 = -\frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_1(z_1) f_1(z_2) \vartheta_n^2 dz_1 dz_2.$$

blete.

Thus, the proof is complete.

6.2. *Truncation and renormalization*. In this subsection, we truncate and renormalize the random variables to ensure the existence of their higher-order moments.

Similar to Jiang and Bai (2021), we may select $\eta_n \to 0$ such that $\eta_n^{-4}n^2 P(|x_{ij}| \ge \eta_n \sqrt{n}) \to 0$. Let $\hat{x}_{ij} = x_{ij} \mathbb{1}_{\{|x_{ij}| < \eta_n \sqrt{n}\}}$ and $\tilde{x}_{ij} = \frac{\hat{x}_{ij} - \mathbb{E}\hat{x}_{ij}}{\hat{\sigma}_n}$, where $\hat{\sigma}_n^2 = \mathbb{E} |\hat{x}_{ij} - \mathbb{E}\hat{x}_{ij}|^2$. Analogous to the discussion in Li and Bai (2015), the sequence $\eta_n \to 0$ can be made arbitrarily slow, therefore, we may require it satisfying $\eta_n n^t \to \infty$ for any fixed t > 0. Correspondingly, we define $\hat{\mathbf{B}} = \frac{1}{n} \mathbf{T} \hat{\mathbf{X}} \hat{\mathbf{X}}^* \mathbf{T}^*$ and $\tilde{\mathbf{B}} = \frac{1}{n} \mathbf{T} \mathbf{X} \tilde{\mathbf{X}}^* \mathbf{T}^*$, where $\hat{\mathbf{X}} = (\hat{x}_{ij})$ and $\tilde{\mathbf{X}} = (\tilde{x}_{ij})$. \hat{G}_n and \tilde{G}_n denote the analogs of G_n with the matrix \mathbf{B} replaced by $\hat{\mathbf{B}}$ and $\tilde{\mathbf{B}}$, respectively. Next, we demonstrate that the limiting distribution of the LSS is unchanged when the entries of \mathbf{X} are replaced by the truncated and renormalized entries.

From Supplement B in Jiang and Bai (2021), we have

$$P(\mathbf{B} \neq \hat{\mathbf{B}}) \le \sum_{i,j} P\left(|x_{ij}| \ge \eta_n \sqrt{n}\right) \le np P\left(|x_{11}| \ge \eta_n \sqrt{n}\right) \to 0, \quad \text{as } n, p \to \infty.$$

It follows from the definition of LSS and Lemma 6.2, for any l = 1, ..., h,

$$\int f_{l}(x)d\hat{G}_{n} - \int f_{l}(x)d\tilde{G}_{n}(x) = \sum_{i=1}^{p} (f_{l}(\lambda_{i}^{\hat{\mathbf{B}}}) - f_{l}(\lambda_{i}^{\tilde{\mathbf{B}}}))$$

$$= \sum_{i=1}^{M} (f_{l}(\lambda_{i}^{\hat{\mathbf{B}}}) - f_{l}(\lambda_{i}^{\tilde{\mathbf{B}}})) + \sum_{i=M+1}^{p} (f_{l}(\lambda_{i}^{\hat{\mathbf{B}}}) - f_{l}(\lambda_{i}^{\tilde{\mathbf{B}}}))$$

$$= \sum_{i=1}^{M} (f_{l}(\lambda_{i}^{\hat{\mathbf{B}}}) - f_{l}(\lambda_{i}^{\tilde{\mathbf{B}}})) + \sum_{i=M+1}^{p} f_{l}(\lambda_{i}^{\hat{\mathbf{B}}}) - \sum_{i=1}^{p-M} f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}}) + \sum_{i=1}^{p-M} (f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}}) - f_{l}(\lambda_{i}^{\tilde{\mathbf{S}}_{22}}))$$

$$+ \sum_{i=1}^{p-M} f_l(\lambda_i^{\tilde{\mathbf{S}}_{22}}) - \sum_{i=M+1}^{p} f_l(\lambda_i^{\tilde{\mathbf{B}}})$$

$$= \sum_{i=1}^{M} (f_l(\lambda_i^{\hat{\mathbf{B}}}) - f_l(\lambda_i^{\tilde{\mathbf{B}}})) + \sum_{i=1}^{p-M} (f_l(\lambda_i^{\hat{\mathbf{S}}_{22}}) - f_l(\lambda_i^{\tilde{\mathbf{S}}_{22}}))$$

$$+ \frac{M}{2\pi i} \oint_{\mathcal{C}} f_l(z) \frac{\hat{m}'_{2n0}(z)}{\hat{m}_{2n0}(z)} dz - \frac{M}{2\pi i} \oint_{\mathcal{C}} f_l(z) \frac{\hat{m}'_{2n0}(z)}{\hat{m}_{2n0}(z)} dz$$

Here $\hat{\mathbf{S}}_{22}$ and $\tilde{\mathbf{S}}_{22}$ denote the analogs of \mathbf{S}_{22} with the matrix \mathbf{X} replaced by $\hat{\mathbf{X}}$ and $\tilde{\mathbf{X}}$, respectively. $\underline{\hat{m}}_{2n0}(z)$ and $\underline{\tilde{m}}_{2n0}(z)$ denote the analogs of $\underline{m}_{2n0}(z)$ with the matrix \mathbf{X} replaced by $\hat{\mathbf{X}}$ and $\tilde{\mathbf{X}}$, respectively.

$$\begin{split} \text{Therefore, } \left| \hat{Y}_{l} - \tilde{Y}_{l} \right| &= \left| \sum_{i=1}^{M} (f_{l}(\lambda_{i}^{\hat{\mathbf{B}}}) - f_{l}(\lambda_{i}^{\hat{\mathbf{B}}})) + \sum_{i=1}^{p-M} (f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}}) - f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}})) \right| &\leq \\ \left| \sum_{i=1}^{M} (f_{l}(\lambda_{i}^{\hat{\mathbf{B}}}) - f_{l}(\lambda_{i}^{\hat{\mathbf{B}}})) \right| + \left| \sum_{i=1}^{p-M} (f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}}) - f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}})) \right|. \\ \text{For } \left| \sum_{i=1}^{p-M} (f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}}) - f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}})) \right|, \text{ we have} \\ &\left| \sum_{i=1}^{p-M} (f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}}) - f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}})) \right| &\leq \sum_{i=1}^{p-M} \left| f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}}) - f_{l}(\lambda_{i}^{\hat{\mathbf{S}}_{22}}) \right| \leq C \sum_{i=1}^{p-M} \left| \lambda_{i}^{\hat{\mathbf{S}}_{22}} - \lambda_{i}^{\hat{\mathbf{S}}_{22}} \right|. \end{split}$$

Similar to Lemma 2.7 in Bai (1999), we have

$$\begin{split} &\sum_{i=1}^{p-M} \left| \lambda_i^{\hat{\mathbf{S}}_{22}} - \lambda_i^{\tilde{\mathbf{S}}_{22}} \right| \leq \left(\frac{2}{n} \operatorname{tr}(\hat{\mathbf{S}}_{22} + \tilde{\mathbf{S}}_{22}) \operatorname{tr} \mathbf{D}_2^{\frac{1}{2}} \mathbf{U}_2^* (\hat{\mathbf{X}} - \tilde{\mathbf{X}}) (\hat{\mathbf{X}} - \tilde{\mathbf{X}})^* \mathbf{U}_2 \mathbf{D}_2^{\frac{1}{2}} \right)^{1/2} \\ &= \left(\frac{2}{n} \operatorname{tr}(\hat{\mathbf{S}}_{22} + \tilde{\mathbf{S}}_{22}) \operatorname{tr}(\hat{\mathbf{X}} - \tilde{\mathbf{X}}) (\hat{\mathbf{X}} - \tilde{\mathbf{X}})^* \mathbf{U}_2 \mathbf{D}_2 \mathbf{U}_2^* \right)^{1/2} \\ &\leq \left(\frac{2C}{n} \operatorname{tr}(\hat{\mathbf{S}}_{22} + \tilde{\mathbf{S}}_{22}) \operatorname{tr}(\hat{\mathbf{X}} - \tilde{\mathbf{X}}) (\hat{\mathbf{X}} - \tilde{\mathbf{X}})^* \right)^{1/2}, \end{split}$$

where the last inequality is because $\mathbf{U}_2\mathbf{D}_2\mathbf{U}_2^*$ is bounded. It is easy to prove $\frac{1}{n}\mathbb{E}\mathrm{tr}(\hat{\mathbf{S}}_{22} + \tilde{\mathbf{S}}_{22}) < C$, and

(6.20)
$$\mathbb{E}\operatorname{tr}(\hat{\mathbf{X}} - \tilde{\mathbf{X}})(\hat{\mathbf{X}} - \tilde{\mathbf{X}})^* = \sum_{i,j} \mathbb{E} \left| (1 - \frac{1}{\hat{\sigma}_n}) \hat{x}_{ij} + \frac{\mathbb{E}\hat{x}_{ij}}{\hat{\sigma}_n} \right|^2$$
$$= \sum_{i,j} \left(\operatorname{var}((1 - \frac{1}{\hat{\sigma}_n}) \hat{x}_{ij} + \frac{\mathbb{E}\hat{x}_{ij}}{\hat{\sigma}_n}) + |\mathbb{E}\hat{x}_{ij}|^2 \right).$$

Since $\operatorname{var}\left((1-\frac{1}{\hat{\sigma}_n})\hat{x}_{ij}+\frac{\mathbb{E}\hat{x}_{ij}}{\hat{\sigma}_n}\right) = (\hat{\sigma}_n-1)^2 \leq \frac{(\hat{\sigma}_n^2-1)^2}{\hat{\sigma}_n^2+1} \leq 2\left(\mathbb{E}|x_{ij}|^2 \mathbb{1}_{\left\{|x_{ij}| \geq \eta_n \sqrt{n}\right\}}\right)^2$, and by the selection of η_n , $\mathbb{E}|x_{ij}|^2 \mathbb{1}_{\left\{|x_{ij}| \geq \eta_n \sqrt{n}\right\}} = o(\eta_n^2 n^{-1})$, $\mathbb{E}\hat{x}_{ij} = o(\eta_n n^{-3/2})$, then we can obtain (6.20) is o(1). Therefore, $\sum_{i=1}^{p-M} \left|\lambda_i^{\hat{\mathbf{S}}_{22}} - \lambda_i^{\hat{\mathbf{S}}_{22}}\right|$ is $o_p(1)$.

For the first term $\left|\sum_{i=1}^{M} (f_l(\lambda_i^{\hat{\mathbf{B}}}) - f_l(\lambda_i^{\hat{\mathbf{B}}}))\right|$, from the arguments in Supplement B of Jiang and Bai (2021), we know that

(6.21)
$$\left|\lambda_i^{\hat{\mathbf{B}}} - \lambda_i^{\tilde{\mathbf{B}}}\right| = o_p(n^{-\frac{1}{2}}\rho_i).$$

We recall that $\rho_i = \alpha_k$ if $i \in J_k$. Then, for brevity, we denote $\beta_i = (\lambda_i^{\hat{\mathbf{B}}} - \lambda_i^{\hat{\mathbf{B}}})/\rho_i$ and obtain that

(6.22)
$$f_l(\lambda_i^{\tilde{\mathbf{B}}}) - f_l(\lambda_i^{\tilde{\mathbf{B}}}) = \int_0^{\beta_i \rho_i} f_l'(t + \lambda_i^{\tilde{\mathbf{B}}}) dt$$
$$= \int_0^1 \beta_i \rho_i f_l'(\beta_i \rho_i s + \lambda_i^{\tilde{\mathbf{B}}}) ds = \beta_i \rho_i f_l'(\rho_i) \int_0^1 \frac{f_l'(\rho_i(\beta_i s + \frac{\lambda_i^{\tilde{\mathbf{B}}}}{\rho_i}))}{f_l'(\rho_i)} ds.$$

Since $\beta_i = o_p(n^{-\frac{1}{2}}), \frac{\lambda_i^{\hat{\mathbf{B}}}}{\rho_i}$ tends to 1, then from Assumption 4 we obtain

(6.23)
$$\sqrt{n} \sum_{i=1}^{M} \frac{\left| f_l(\lambda_i^{\mathbf{B}}) - f_l(\lambda_i^{\mathbf{B}}) \right|}{f_l'(\rho_i)\rho_i} = o_p(1).$$

Then we have $\frac{\left|\sum_{i=1}^{M} (f_{l}(\lambda_{i}^{\hat{\mathbf{B}}}) - f_{l}(\lambda_{i}^{\hat{\mathbf{B}}}))\right|}{\varsigma_{l}} \leq \frac{\sum_{i=1}^{M} \left|f_{l}(\lambda_{i}^{\hat{\mathbf{B}}}) - f_{l}(\lambda_{i}^{\hat{\mathbf{B}}})\right|}{\varsigma_{l}} \leq \sum_{i=1}^{M} \frac{\left|f_{l}(\lambda_{i}^{\hat{\mathbf{B}}}) - f_{l}(\lambda_{i}^{\hat{\mathbf{B}}})\right|}{f_{l}'(\rho_{i})\rho_{i}/\sqrt{n}} = o_{p}(1).$ Thus, it is concluded that the procedure of truncation does not affect the limiting distribution

Thus, it is concluded that the procedure of truncation does not affect the limiting distribution of LSS.

Therefore, in the following proofs, we can safely assume that $|x_{ij}| < \eta_n \sqrt{n}$.

Acknowledgments. The authors would like to thank Professor Jeff Yao for many helpful suggestions and discussions. Zhidong Bai was partially supported by NSFC Grants No.12171198, No.12271536, and Team Project of Jilin Provincial Department of Science and Technology No.20210101147JC. Jiang Hu was partially supported by NSFC Grant No.12171078.

SUPPLEMENTARY MATERIAL

Supplement to "A CLT for the LSS of large dimensional sample covariance matrices with diverging spikes"

This supplementary document contains proofs of Theorems 3.1–4.5, and some useful lemmas. We also report some additional simulation results in this document.

REFERENCES

Theodore Wilbur Anderson. An Introduction to Multivariate Statistical Analysis. Third Edition. Wiley New York, 2003.

- Jushan Bai and Serena Ng. Determining the number of factors in approximate factor models. *Econometrica*, 70 (1):191–221, 2002.
- Zhidong Bai Methodologies in spectral analysis of large dimensional random matrices, A review. *Statistica Sinica*, 9:611–677, 1999.
- Zhidong Bai, Baiqi Miao, and Guangming Pan. On asymptotics of eigenvectors of large sample covariance matrix. *The Annals of Probability*, 35(4):1532–1572, 2007.
- Zhidong Bai, Jiang Hu, Guangming Pan, and Wang Zhou. Convergence of the empirical spectral distribution function of beta matrices. *Bernoulli*, 21(3):1538–1574, 2015.
- Zhidong Bai, Huiqin Li, and Guangming Pan. Central limit theorem for linear spectral statistics of large dimensional separable sample covariance matrices. *Bernoulli*, 25(3):1838–1869, 2019.
- Zhidong Bai and Jack W. Silverstein. CLT for linear spectral statistics of large-dimensional sample covariance matrices. *The Annals of Probability*, 32(1A):553 605, 2004.
- Zhidong Bai and Jianfeng Yao. Central limit theorems for eigenvalues in a spiked population model. *Annales de l'Institut Henri Poincaré, Probabilités et Statistiques*, 44(3):447 474, 2008.
- Zhidong Bai and Jianfeng Yao. On sample eigenvalues in a generalized spiked population model. *Journal of Multivariate Analysis*, 106:167–177, 2012.
- Zhidong Bai, Dandan Jiang, Jianfeng Yao, and Shurong Zheng. Corrections to LRT on large-dimensional covariance matrix by RMT. *The Annals of Statistics*, 37(6B):3822 – 3840, 2009.

- Jinho Baik and Jack W. Silverstein. Eigenvalues of large sample covariance matrices of spiked population models. Journal of Multivariate Analysis, 97(6):1382–1408, 2006.
- Jinho Baik, Ji Oon Lee, and Hao Wu. Ferromagnetic to paramagnetic transition in spherical spin glass. Journal of Statistical Physics, 173(5):1484–1522, 2018.
- Marwa Banna, Jamal Najim, and Jianfeng Yao. A clt for linear spectral statistics of large random informationplus-noise matrices. *Stochastic Processes and their Applications*, 130(4):2250–2281, 2020.
- Zhigang Bao, Jiang Hu, Xiaocong Xu, and Xiaozhuo Zhang. Spectral statistics of sample block correlation matrices. arXiv:2207.06107, 2022.
- Alex Bloemendal, Antti Knowles, Horng-Tzer Yau, and Jun Yin. On the principal components of sample covariance matrices. *Probability Theory and Related Fields*, 164(1-2):459–552, 2016.
- T. Tony Cai, Xiao Han, and Guangming Pan. Limiting laws for divergent spiked eigenvalues and largest nonspiked eigenvalue of sample covariance matrices. *The Annals of Statistics*, 48(3):1255–1280, 2020.
- Binbin Chen and Guangming Pan. Clt for linear spectral statistics of normalized sample covariance matrices with the dimension much larger than the sample size. *Bernoulli*, 21(2):1089–1133, 2015.
- Xiucai Ding and Fan Yang A necessary and sufficient condition for edge universality at the largest singular values of covariance matrices. *The Annals of Applied Probability*, 28(3):1679–1738, 2018.
- Edgar Dobriban. Permutation methods for factor analysis and pca. *The Annals of Statistics*, 48(5):2824–2847, 2020.
- David Donoho, Matan Gavish, and Iain Johnstone. Optimal shrinkage of eigenvalues in the spiked covariance model. *The Annals of Statistics*, 46(4):1742–1778, 2018.
- Jiti Gao, Xiao Han, Guangming Pan, and Yanrong Yang. High dimensional correlation matrices: The central limit theorem and its applications. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79 (3):677–693, 2017.
- Jiang Hu, Weiming Li, Zhi Liu, and Wang Zhou. High-dimensional covariance matrices in elliptical distributions with application to spherical test. *The Annals of Statistics*, 47(1):527–555, 2019.
- Dandan Jiang and Zhidong Bai. Generalized four moment theorem and an application to clt for spiked eigenvalues of high-dimensional covariance matrices. *Bernoulli*, 27(1):274–294, 2021.
- Tiefeng Jiang and Fan Yang. Central limit theorems for classical likelihood ratio tests for high-dimensional normal distributions. *The Annals of Statistics*, 41(4):2029–2074, 2013.
- Iain M. Johnstone. On the distribution of the largest eigenvalue in principal components analysis. *The Annals of Statistics*, 29(2):295 327, 2001.
- Iain M. Johnstone. Multivariate analysis and Jacobi ensembles: largest eigenvalue, Tracy-Widom limits and rates of convergence. *The Annals of Statistics*, 36(6):2638 – 2716, 2008.
- Iain M. Johnstone and Boaz Nadler. Roy's largest root test under rank-one alternatives. *Biometrika*, 104(1): 181–193, 2017.
- Iain M. Johnstone and Alexei Onatski. Testing in high-dimensional spiked models. *The Annals of Statistics*, 48 (3):1231–1254, 2020.
- Iain M. Johnstone and Debashis Paul. PCA in high dimensions: An orientation. *Proceedings of the IEEE*, 106(8): 1277–1292, 2018.
- Sungkyu Jung and J. S. Marron. PCA consistency in high dimension, low sample size context. The Annals of Statistics, 37(6B):4104–4130, 2009.
- Olivier Ledoit and Michael Wolf. Some hypothesis tests for the covariance matrix when the dimension is large compared to the sample size. *The Annals of Statistics*, 30(4):1081–1102, 2002.
- Huiqin Li and Zhidong Bai. Extreme eigenvalues of large dimensional quaternion sample covariance matrices. *Journal of Statistical Planning and Inference*, 159:1–14, 2015.
- Zeng Li, Fang Han, and Jianfeng Yao. Asymptotic joint distribution of extreme eigenvalues and trace of large sample covariance matrix in a generalized spiked population model. *The Annals of Statistics*, 48(6):3138–3160, 2020.
- Zeng Li, Qinwen Wang, and Runze Li. Central limit theorem for linear spectral statistics of large dimensional kendall's rank correlation matrices and its applications. *The Annals of Statistics*, 49(3):1569–1593, 2021.
- Boaz Nadler. Finite sample approximation results for principal component analysis: A matrix perturbation approach. *The Annals of Statistics*, 36(6):2791–2817, 2008.
- Hisao Nagao. On Some Test Criteria for Covariance Matrix The Annals of Statistics 1(4):700-709, 1973.
- Jamal Najim and Jianfeng Yao. Gaussian fluctuations for linear spectral statistics of large random covariance matrices. *The Annals of Applied Probability*, 26(3):1837–1887, 2016.
- Chester L. Olson. Comparative robustness of six tests in multivariate analysis of variance *Journal of the American Statistical Association*, 69, 894–908, 1974.
- Alexei Onatski, Marcelo J. Moreira, and Marc Hallin. Asymptotic power of sphericity tests for high-dimensional data. *The Annals of Statistics*, 41(3):1204–1231, 2013.

- Alexei Onatski, Marcelo J. Moreira, and Marc Hallin. Signal detection in high dimension: The multispiked case. *The Annals of Statistics*, 42(1):225–254, 2014.
- Guangming Pan. Comparison between two types of large sample covariance matrices. Annales de l'Institut Henri Poincaré, Probabilités et Statistiques, 50(2):655–677, 2014.
- Guangming Pan and Wang Zhou. Central limit theorem for signal-to-interference ratio of reduced rank linear receiver. *The Annals of Applied Probability*, 18(3):1232–1270, 2008.
- Debashis Paul. Asymptotics of sample eigenstructure for a large dimensional spiked covariance model. *Statistica Sinica*, 17(4):1617–1642, 2007.
- Amelia Perry, Alexander S. Wein, Afonso S. Bandeira, and Ankur Moitra. Optimality and sub-optimality of pca i: Spiked random matrix models. *The Annals of Statistics*, 46(5):2416–2451, 2018.
- Jack W. Silverstein. Strong convergence of the empirical distribution of eigenvalues of large dimensional random matrices. *Journal of Multivariate Analysis*, 55(2):331–339, 1995.
- Qinwen Wang and Jianfeng Yao. On the sphericity test with large-dimensional observations. *Electronic Journal* of Statistics, 7:2164–2192, 2013.
- Qinwen Wang and Jianfeng Yao. Extreme eigenvalues of large-dimensional spiked fisher matrices with application. *The Annals of Statistics*, 45(1):415–460, 2017.
- Weichen Wang and Jianqing Fan. Asymptotics of empirical eigenstructure for high dimensional spiked covariance. *The Annals of Statistics*, 45(3):1342–1374, 2017.
- Samuel S. Wilks. The Large-Sample Distribution of the Likelihood Ratio for Testing Composite Hypotheses. *The Annals of Mathematical Statistics*, 9(1):60–62, 1938.
- Jeha Yang and Iain M. Johnstone. Edgeworth correction for the largest eigenvalue in a spiked pca model. *Statistica Sinica*, 2018.
- Yanrong Yang and Guangming Pan. Independence test for high dimensional data based on regularized canonical correlation coefficients. *The Annals of Statistics*, 43(2):467–500, 2015.
- Jianfeng Yao, Shurong Zheng, and Zhidong Bai. Large Sample Covariance Matrices and High-Dimensional Data Analysis. Cambridge University Press, 2015.
- Zhigang Yao, Ye Zhang, Zhidong Bai, and William F. Eddy. Estimating the number of sources in magnetoencephalography using spiked population eigenvalues. *Journal of the American Statistical Association*, 113(522): 505–518, 2018.
- Yanqing Yin. Spectral statistics of high dimensional sample covariance matrix with unbounded population spectral norm. *Bernoulli*, 28(3):1729–1756, 2022.
- Zhixiang Zhang, Shurong Zheng, Guangming Pan, and Pingshou Zhong. Asymptotic independence of spiked eigenvalues and linear spectral statistics for large sample covariance matrices. *The Annals of Statistics*, 50(4): 2205–2230, 2022.
- Shurong Zheng. Central limit theorems for linear spectral statistics of large dimensional F-matrices. Annales de l'Institut Henri Poincaré, Probabilités et Statistiques, 48(2):444–476, 2012.
- Shurong Zheng, Zhidong Bai, and Jianfeng Yao. Substitution principle for CLT of linear spectral statistics of high-dimensional sample covariance matrices with applications to hypothesis testing. *The Annals of Statistics*, 43(2):546 – 591, 2015.
- Shurong Zheng, Guanghui Cheng, Jianhua Guo, and Hongtu Zhu. Test for high-dimensional correlation matrices. *The Annals of Statistics*, 47(5):2887–2921, 2019.
- Yihui Zhou and James Stephen Marron. High dimension low sample size asymptotics of robust PCA. *Electronic Journal of Statistics*, 9(1):204–218, 2015.

SUPPLEMENT TO "A CLT FOR THE LSS OF LARGE DIMENSIONAL SAMPLE COVARIANCE MATRICES WITH DIVERGING SPIKES"

BY ZHIJUN LIU^{*}, JIANG HU[†], ZHIDONG BAI[‡], AND HAIYAN SONG[§]

KLASMOE and School of Mathematics and Statistics, Northeast Normal University, China. *liuzj037@nenu.edu.cn; †huj156@nenu.edu.cn; ‡baizd@nenu.edu.cn; § songhy716@nenu.edu.cn

In this document we present some technical details involved in Liu et al. (2022). More precisely, in Section 7, we prove Theorems 3.1–4.5 of the main file. Some derivations and calculations in Section 7 are postponed to Section 8. In Section 9 we provide some useful lemmas. Finally, in Section 10, we report some additional simulation results in this part.

The number of scheme(equations, theorems, lemmas, etc.) is shared with the main document so that there are no misunderstandings with the use of references.

7. Proofs of Theorems 3.1–4.5.

7.1. *Proof of Theorem 3.1.* The proof of Theorem 3.1 builds on the decomposition analysis of the LSSs and it is divided into part (I) $\sum_{j=1}^{M} f(\lambda_j)$ and part (II) $\sum_{j=M+1}^{p} f(\lambda_j)$. Enlightened by the BST in Bai and Silverstein (2004), we have

$$\begin{split} &\sum_{j=1}^{p} f(\lambda_{j}) - p \int f(x) dF^{c_{n},H_{n}} \\ &= \sum_{j=1}^{M} f(\lambda_{j}) + \sum_{j=M+1}^{p} f(\lambda_{j}) - p \int f(x) dF^{c_{n},H_{n}} \\ &= \sum_{j=1}^{M} f(\lambda_{j}) + \sum_{j=1}^{p-M} f\left(\tilde{\lambda}_{j}\right) - (p-M) \int f(x) dF^{c_{nM},H_{2n}} + \sum_{j=M+1}^{p} f(\lambda_{j}) - \sum_{j=1}^{p-M} f\left(\tilde{\lambda}_{j}\right) \\ &+ (p-M) \int f(x) dF^{c_{nM},H_{2n}} - p \int f(x) dF^{c_{n},H_{n}}. \end{split}$$

Since Lemma 6.1 has shown the difference between $(p - M) \int f(x) dF^{c_{nM}, H_{2n}}$ and $p \int f(x) dF^{c_{n,H_n}}$ is 0. Moreover, in Lemma 6.2 we have proved

$$\sum_{j=M+1}^{p} f(\lambda_j) - \sum_{j=1}^{p-M} f\left(\tilde{\lambda}_j\right) = \frac{M}{2\pi i} \oint_C f(z) \frac{\underline{m}'_{2n0}(z)}{\underline{m}_{2n0}(z)} dz + o_p(1).$$

It follows that

$$\sum_{j=1}^{p} f(\lambda_{j}) - p \int f(x) dF^{c_{n},H_{n}}$$

$$= \sum_{j=1}^{M} f(\lambda_{j}) + \sum_{j=1}^{p-M} f\left(\widetilde{\lambda}_{j}\right) - (p-M) \int f(x) dF^{c_{nM},H_{2n}} + \frac{M}{2\pi i} \oint_{C} f(z) \frac{\underline{m}_{2n0}^{'}(z)}{\underline{m}_{2n0}(z)} dz + o_{p}(1),$$

MSC2020 subject classifications: Primary 60B20; secondary 60F05.

Keywords and phrases: Empirical spectral distribution, linear spectral statistic, random matrix, Stieltjes transform.

which yields

$$\sum_{j=1}^{p} f(\lambda_{j}) - p \int f(x) dF^{c_{n},H_{n}} - \sum_{k=1}^{K} d_{k} f(\phi_{n}(\alpha_{k})) - \frac{M}{2\pi i} \oint_{C} f(z) \frac{\underline{m}'_{2n0}(z)}{\underline{m}_{2n0}(z)} dz$$
(7.24)
$$= \sum_{j=1}^{M} f(\lambda_{j}) - \sum_{k=1}^{K} d_{k} f(\phi_{n}(\alpha_{k})) + \sum_{j=1}^{p-M} f(\widetilde{\lambda}_{j}) - (p-M) \int f(x) dF^{c_{nM},H_{2n}} + o_{p}(1)$$

The analysis below is executed by dividing (7.24) into two parts: (I) $\sum_{j=1}^{M} f(\lambda_j)$ – $\sum_{k=1}^{K} d_k f(\phi_n(\alpha_k)) \text{ and (II)} \sum_{j=1}^{p-M} f(\widetilde{\lambda}_j) - (p-M) \int f(x) dF^{c_{nM}, H_{2n}}, \text{ where we ignore the impact of } o_p(1) \text{ on the asymptotic distribution. Since we have derived the asymptotic dis$ tribution of part (II) in Lemma 6.5, we only need to consider the asymptotic distribution of part (I) $\sum_{j=1}^{M} f(\lambda_j) - \sum_{k=1}^{K} d_k f(\phi_n(\alpha_k))$. From the proof of Lemma 6.4, $\sum_{j=1}^{M} f(\lambda_j) - \sum_{k=1}^{K} d_k f(\phi_n(\alpha_k))$ has the same limiting distribution as $\sum_{k=1}^{K} \varpi_{nk} \sum_{j \in J_k} \gamma_{kj}$. From Lemma 6.3, we have $(\gamma_{kj}, j \in J_k)^{\top}$ converges weakly to the joint distribution of the d_k eigenvalues of Gaussian random matrix $-\frac{1}{\theta_k} [\Omega_{\phi_k}]_{kk}$, so $\sum_{j \in J_k} \gamma_{kj} \stackrel{d}{\to} -\frac{1}{\theta_k} \operatorname{tr} [\Omega_{\phi_k}]_{kk}$. Recall that ω_{ij} is the element of Ω_{ϕ_k} , and $\operatorname{tr} [\Omega_{\phi_k}]_{kk}$ is the summation of the diagonal element, that is, $\sum_{j \in J_k} \omega_{jj}$. Since the diagonal elements are i.i.d., then $\mathbb{E}\left(\sum_{j \in J_k} \omega_{jj}\right) = 0$, $\operatorname{Var}\left(\sum_{j\in J_k}\omega_{jj}\right) = \sum_{j\in J_k}\operatorname{Var}\left(\omega_{jj}\right) + \sum_{j_1\neq j_2}\operatorname{cov}\left(\omega_{j_1j_1},\omega_{j_2j_2}\right) = \sum_{j\in J_k}\left(\left(\alpha_x+1\right)\theta_k + \beta_x\mathcal{U}_{j_1j_j}\nu_k\right) + \sum_{j_1\neq j_2}\beta_x\mathcal{U}_{j_1j_1j_2j_2}\nu_k = \left(\alpha_x+1\right)\theta_kd_k + \sum_{j_1,j_2\in J_k}\mathcal{U}_{j_1j_1j_2j_2}\beta_x\nu_k$ Therefore, from Lemma 6.3, we have that the asymptotic distribution of $\sum_{j\in J_k}\gamma_{kj}$ is a

Gaussian distribution with

$$\mathbb{E}\sum_{j\in J_k}\gamma_{kj}=0,$$

$$s_k^2 \triangleq \operatorname{Var}\left(\sum_{j \in J_k} \gamma_{kj}\right) = \frac{(\alpha_x + 1)\,\theta_k d_k + \sum_{j_1, j_2 \in J_k} \mathcal{U}_{j_1 j_1 j_2 j_2} \beta_x \nu_k}{\theta_k^2}$$

and then, we directly derive that the mean function of $\sum_{k=1}^{K} \varpi_{nk} \sum_{i \in J_k} \gamma_{kj}$ is 0 and that its covariance function is

$$\operatorname{Var}(Y_{f_1}) = \sum_{k=1}^{K} \varpi_{nk1}^2 s_k^2.$$

Finally, we focus on the asymptotic distribution of equation (7.24). Because of Lemma 6.4, the two LSSs are asymptotically independent; thus, the random variable

$$\varsigma_1^{-1}(Y_1 - \mathbb{E}Y_1) \xrightarrow{d} N(0, 1)$$

with mean function $\mathbb{E}Y_1 = \mu_1$ being

$$-\frac{\alpha_x}{2\pi i} \cdot \oint_{\mathcal{C}} f_1(z) \frac{c_{nM} \int \underline{m}_{2n0}^3(z) t^2 \left(1 + t\underline{m}_{2n0}(z)\right)^{-3} dH_{2n}(t)}{\left(1 - c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t\underline{m}_{2n0}(z))^2} dH_{2n}(t)\right) \left(1 - \alpha_x c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t\underline{m}_{2n0}(z))^2} dH_{2n}(t)\right)} dz$$
$$-\frac{\beta_x}{2\pi i} \cdot \oint_{\mathcal{C}} f_1(z) \frac{c_{nM} \int \underline{m}_{2n0}^3(z) t^2 \left(1 + t\underline{m}_{2n0}(z)\right)^{-3} dH_{2n}(t)}{1 - c_{nM} \int \underline{m}_{2n0}^2(z) t^2 \left(1 + t\underline{m}_{2n0}(z)\right)^{-2} dH_{2n}(t)} dz,$$

and covariance function ς_1^2 being

$$\sum_{k=1}^{K} \varpi_{nk1}^2 s_k^2 - \frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_1(z_1) f_1(z_2) \vartheta_n^2 dz_1 dz_2,$$

where ϑ_n^2 is defined in Lemma 6.5. Therefore, the proof is finished.

7.2. *Proof of Theorem 3.2.* Similar to the proof of Theorem 3.1, we divide the LSSs into two parts. Different from the above analysis, in this section, we focus on the multidimensional case under Assumptions 1-6. Recall that we defined

$$G_{n}(x) = p\left[F^{\mathbf{B}}(x) - F^{c_{n},H_{n}}(x)\right],$$

$$Y_{l} = \int f_{l}(x) \, dG_{n}(x) - \sum_{k=1}^{K} d_{k} f_{l}(\phi_{n}(\alpha_{k})) - \frac{M}{2\pi i} \oint_{\mathcal{C}} f_{l}(z) \, \frac{\underline{m}'_{2n0}(z)}{\underline{m}_{2n0}(z)} dz.$$

Because of equation (7.24), the random vector (Y_1, \ldots, Y_h) shares the same asymptotic distribution with the summation of two random vectors

$$\left(\sum_{k=1}^{K} \varpi_{nk1} \sum_{j \in J_k} \gamma_{kj}, \dots, \sum_{k=1}^{K} \varpi_{nkh} \sum_{j \in J_k} \gamma_{kj}\right)$$

and

$$\left(\sum_{j=1}^{p-M} f_1\left(\tilde{\lambda}_j\right) - (p-M) \int f_1(x) dF^{c_{nM}, H_{2n}}, \dots, \sum_{j=1}^{p-M} f_h\left(\tilde{\lambda}_j\right) - (p-M) \int f_h(x) dF^{c_{nM}, H_{2n}}\right).$$

First, we focus on the first random vector. Similar to the proof of Theorem 3.1, we derive that the mean function of the first random vector is 0 and that the covariance function is

$$\operatorname{Cov}\left(Y_{f_{s}}, Y_{f_{t}}\right) = \sum_{k=1}^{K} \frac{\phi_{n}^{2}\left(\alpha_{k}\right)}{n} f_{s}'\left(\phi_{n}\left(\alpha_{k}\right)\right) f_{t}'\left(\phi_{n}\left(\alpha_{k}\right)\right) s_{k}^{2} = \sum_{k=1}^{K} \varpi_{nks} \varpi_{nkt} s_{k}^{2},$$

Moreover, the asymptotic distribution of the second random vector is derived in Zheng et al. (2015). Because of Lemma 6.4, two random vectors are asymptotically independent; thus, the random vector

$$(Y_1 - \mathbb{E}Y_1, \dots, Y_h - \mathbb{E}Y_h)^\top \xrightarrow{d} N_h(0, \mathbf{\Omega})$$

with mean function $\mathbb{E}Y_l$ is the same as μ_l , and the covariance matrix is Ω with its entries

$$\omega_{st} = \sum_{k=1}^{K} \varpi_{nks} \varpi_{nkt} s_k^2 - \kappa_{nst},$$

where

$$\begin{split} \kappa_{nst} &= \frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} \frac{f_s\left(z_1\right) f_t\left(z_2\right)}{\left(\underline{m}_{2n0}\left(z_1\right) - \underline{m}_{2n0}\left(z_2\right)\right)^2} d\underline{m}_{2n0}\left(z_1\right) d\underline{m}_{2n0}\left(z_2\right) + \frac{c_{nM}\beta_x}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_s\left(z_1\right) \\ f_t\left(z_2\right) \left[\int \frac{t}{\left(\underline{m}_{2n0}\left(z_1\right) t + 1\right)^2} \times \frac{t}{\left(\underline{m}_{2n0}\left(z_2\right) t + 1\right)^2} dH_{2n}(t) \right] d\underline{m}_{2n0}\left(z_1\right) d\underline{m}_{2n0}\left(z_2\right) \\ &+ \frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_s\left(z_1\right) f_t\left(z_2\right) \left[\frac{\partial^2}{\partial z_1 \partial z_2} \log\left(1 - a_n\left(z_1, z_2\right)\right) \right] dz_1 dz_2, \\ a_n\left(z_1, z_2\right) &= \alpha_x \left(1 + \frac{\underline{m}_{2n0}\left(z_1\right) \underline{m}_{2n0}\left(z_2\right) \left(z_1 - z_2\right)}{\underline{m}_{2n0}\left(z_2\right) - \underline{m}_{2n0}\left(z_1\right)} \right). \end{split}$$

28

Then, we obtain the random vector

$$\left(\frac{Y_1 - \mathbb{E}Y_1}{\sigma_1}, \dots, \frac{Y_h - \mathbb{E}Y_h}{\sigma_h}\right)^\top \stackrel{d}{\to} N_h(0, \boldsymbol{\Psi}),$$

which has a mean function that is the same as that in Theorem 3.1, and variance function

$$\sigma_l^2 = \sum_{k=1}^K \varpi_{nkl}^2 s_k^2 - \kappa_{nll}, \quad l = 1, \dots, h,$$

and the covariance matrix $\Psi = (\psi_{st})_{h \times h}$ is the correlation coefficient matrix of random vector $(Y_1, \ldots, Y_h)^\top$ with its entries $\psi_{st} = \lim_{n \to \infty} \psi_{nst}$,

$$\psi_{nst} = \frac{\sum_{k=1}^{K} \overline{\omega}_{nks} \overline{\omega}_{nkt} s_k^2 - \kappa_{nst}}{\sqrt{\sum_{k=1}^{K} \overline{\omega}_{nks}^2 s_k^2 - \kappa_{nss}} \sqrt{\sum_{k=1}^{K} \overline{\omega}_{nkt}^2 s_k^2 - \kappa_{ntt}}},$$

Note that renormalization is necessary to guarantee that elements in the correlation coefficient matrix Ψ are limited. Therefore, the proof is finished.

7.3. *Proof of Theorem 4.1.* The result under H_0 is a direct result of Theorem 4.1 in Zheng et al. (2015) using the substitution principle. Therefore, we omit the proof here. Next, we focus on the result under H_1 . Recall that

$$G_{n}(x) = p\left[F^{\mathbf{B}}(x) - F^{c_{n},H_{n}}(x)\right],$$

$$Y = \int f_L(x) \, dG_n(x) - \sum_{k=1}^K d_k f_L(\phi_n(\alpha_k)) - \frac{M}{2\pi i} \oint_{\mathcal{C}} f_L(z) \, \frac{\underline{m}'_{2n0}(z)}{\underline{m}_{2n0}(z)} dz,$$

when $f_L(x) = x - \log x - 1$. After some calculations, we obtain

$$\int f_{L}(x) dG_{n}(x) = \operatorname{tr} \mathbf{B} - \log |\mathbf{B}| - p - p \int f_{L}(x) dF^{c_{n},H_{n}}(x) = L - p \int f_{L}(x) dF^{c_{n},H_{n}}(x),$$
(7.25)
$$p \int f_{L}(x) dF^{c_{n},H_{n}}(x) = (p - M)(1 - \frac{c_{nM} - 1}{c_{nM}} \log (1 - c_{nM})),$$

$$\sum_{k=1}^{K} d_{k} f_{L}(\phi_{n}(\alpha_{k})) = \sum_{k=1}^{K} d_{k}(\phi_{n}(\alpha_{k}) - \log \phi_{n}(\alpha_{k}) - 1),$$
(7.26)
$$\frac{M}{2\pi i} \oint_{\mathcal{C}} f_{L}(z) \frac{\underline{m}'_{2n0}(z)}{\underline{m}_{2n0}(z)} dz = -M(c_{nM} + \log(1 - c_{nM})),$$

where (7.25) is obtained from Lemma 6.1 and Bai et al. (2009). For consistency, we present the proof of (7.26) in Section 8. According to Theorem 3.1, since $f_L(x) = x - \log x - 1$, $\mathbf{D}_2 = \mathbf{I}_{p-M}$, $\mathbf{\Gamma} = \mathbf{V}_2 \mathbf{U}_2^*$, then we have

$$\frac{L - p \int f_L(x) dF^{c_n, H_n}(x) - \breve{\mu}_l}{\breve{\varsigma}_l} \xrightarrow{d} N(0, 1).$$

where the mean function is $\breve{\mu}_l = -\frac{\log(1-c_{nM})}{2}\alpha_x + \frac{c_{nM}}{2}\beta_x + \sum_{k=1}^K d_k(\phi_n(\alpha_k) - \log\phi_n(\alpha_k) - 1) - M(c_{nM} + \log(1-c_{nM}))$. For covariance term, $\breve{\varsigma}_l^2$ equals $\sum_{k=1}^K \frac{(\phi_n(\alpha_k)-1)^2}{n}s_k^2 - \frac{1}{4\pi^2}\oint_{\mathcal{C}_1}\oint_{\mathcal{C}_2} (z_1 - \log(z_1) - 1)(z_2 - \log(z_2) - 1)\vartheta_n^2 dz_1 dz_2, \text{ where } t_1 = 0$

$$s_k^2 = \frac{(\alpha_x + 1)d_k}{\theta_k} + \frac{\sum_{j_1, j_2 \in J_k} \mathcal{U}_{j_1 j_1 j_2 j_2} \beta_x \nu_k}{\theta_k^2}, \ \vartheta_n^2 = \Theta_{0,n}(z_1, z_2) + \alpha_x \Theta_{1,n}(z_1, z_2) + \beta_x \Theta_{2,n}(z_1, z_2),$$
 and

$$\Theta_{0,n}(z_1, z_2) = \frac{\underline{m}'_{2n0}(z_1)\underline{m}'_{2n0}(z_2)}{(\underline{m}_{2n0}(z_1) - \underline{m}_{2n0}(z_2))^2} - \frac{1}{(z_1 - z_2)^2}.$$

For $\Theta_{1,n}(z_1, z_2)$, since

$$\mathbf{P}_{n}(z) = ((1 - c_{nM}) \mathbf{V}_{2} \mathbf{V}_{2}^{*} - z c_{nM} m_{2n0}(z) \mathbf{V}_{2} \mathbf{V}_{2}^{*} - z \mathbf{I}_{p})^{-1}$$

then

$$\begin{aligned} \mathcal{A}_{n}(z_{1},z_{2}) &= \frac{z_{1}z_{2}}{n} \underline{m}_{2n0}(z_{1}) \underline{m}_{2n0}(z_{2}) \operatorname{tr} \mathbf{U}_{2} \mathbf{V}_{2}^{*} \mathbf{P}_{n}(z_{1}) \, \mathbf{V}_{2} \mathbf{U}_{2}^{*} \overline{\mathbf{U}}_{2} \mathbf{V}_{2}^{\top} \mathbf{P}_{n}^{\top}(z_{2}) \, \overline{\mathbf{V}}_{2} \mathbf{U}_{2}^{\top}, \\ &= \frac{z_{1}z_{2} \underline{m}_{2n0}(z_{1}) \underline{m}_{2n0}(z_{2}) \operatorname{tr} \mathbf{U}_{2} \mathbf{U}_{2}^{*} \overline{\mathbf{U}}_{2} \mathbf{U}_{2}^{\top}}{n \left(1 - c_{nM} - z_{1} c_{nM} m_{2n0}(z_{1}) - z_{1}\right) \left(1 - c_{nM} - z_{2} c_{nM} m_{2n0}(z_{2}) - z_{2}\right)}, \\ &= \frac{\underline{m}_{2n0}(z_{1}) \underline{m}_{2n0}(z_{2})}{n \left(1 + \underline{m}_{2n0}(z_{1})\right) \left(1 + \underline{m}_{2n0}(z_{2})\right)} \operatorname{tr} \mathbf{U}_{2} \mathbf{U}_{2}^{*} \overline{\mathbf{U}}_{2} \mathbf{U}_{2}^{\top}. \end{aligned}$$

For tr $\mathbf{U}_{2}\mathbf{U}_{2}^{*}\mathbf{\bar{U}}_{2}\mathbf{U}_{2}^{\top}$, since $\mathbf{U}\mathbf{U}^{*} = \mathbf{I}_{p}$, therefore tr $\mathbf{U}_{2}\mathbf{U}_{2}^{*}\mathbf{\bar{U}}_{2}\mathbf{U}_{2}^{\top} = \text{tr}(\mathbf{I}_{p} - \mathbf{U}_{1}\mathbf{U}_{1}^{*})(\mathbf{I}_{p} - \mathbf{U}_{1}\mathbf{U}_{1}^{*})^{\top} = p - \text{tr}(\mathbf{U}_{1}\mathbf{U}_{1}^{*})^{\top} - \text{tr}\mathbf{U}_{1}\mathbf{U}_{1}^{*} + \text{tr}\mathbf{U}_{1}\mathbf{U}_{1}^{*}(\mathbf{U}_{1}\mathbf{U}_{1}^{*})^{\top}$. Moreover, since tr $\mathbf{U}_{1}\mathbf{U}_{1}^{*} = M$, tr $\mathbf{U}_{1}\mathbf{U}_{1}^{*}(\mathbf{U}_{1}\mathbf{U}_{1}^{*})^{\top} = \sum_{s,t=1}^{p} \left(\sum_{i=1}^{M} u_{si}\bar{u}_{ti}\right)^{2}$. Therefore, $\mathcal{A}_{n(z_{1},z_{2})} = \frac{p-2M+\sum_{s,t=1}^{p} \left(\sum_{i=1}^{M} u_{si}\bar{u}_{ti}\right)^{2}}{n} \frac{\underline{m}_{2n0}(z_{1})\underline{m}_{2n0}(z_{2})}{(1+\underline{m}_{2n0}(z_{1}))(1+\underline{m}_{2n0}(z_{2}))}$. Denote $\tilde{c} = \frac{p-2M+\sum_{s,t=1}^{p} \left(\sum_{i=1}^{M} u_{si}\bar{u}_{ti}\right)^{2}}{n}$, then $\mathcal{A}_{n}(z_{1},z_{2}) = \frac{\tilde{c}\underline{m}_{2n0}(z_{1})\underline{m}_{2n0}(z_{2})}{(1+\underline{m}_{2n0}(z_{1}))(1+\underline{m}_{2n0}(z_{2}))}$. Therefore fore

$$\begin{split} \Theta_{1,n}(z_1, z_2) = & \frac{\partial}{\partial z_2} \left\{ \frac{\partial \mathcal{A}_n(z_1, z_2)}{\partial z_1} \frac{1}{1 - \alpha_x \mathcal{A}_n(z_1, z_2)} \right\}, \\ = & \frac{\tilde{c}\underline{m}'_{2n0}(z_1)\underline{m}'_{2n0}(z_2)}{\left(\left(1 + \underline{m}_{2n0}(z_1)\right)\left(1 + \underline{m}_{2n0}(z_2)\right) - \alpha_x \tilde{c}\underline{m}_{2n0}(z_1)\underline{m}_{2n0}(z_2)\right)^2}. \end{split}$$

For $\Theta_{2,n}(z_1, z_2)$, since $\Theta_{2,n}(z_1, z_2) = \frac{z_1^2 z_2^2 \underline{m}'_{2n0}(z_1)\underline{m}'_{2n0}(z_2)}{n} \sum_{i=1}^p \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_1)\mathbf{\Gamma}\right]_{ii} \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_2)\mathbf{\Gamma}\right]_{ii}, \end{split}$
and

$$\mathbf{\Gamma}^* \mathbf{P}_n^2(z_1) \mathbf{\Gamma} = \mathbf{U}_2 \mathbf{V}_2^* \left((1 - c_{nM}) \mathbf{V}_2 \mathbf{V}_2^* - z_1 c_{nM} m_{2n0}(z_1) \mathbf{V}_2 \mathbf{V}_2^* - z \mathbf{I}_p \right)^{-2} \mathbf{V}_2 \mathbf{U}_2^*,$$

by using lemma 9.2, we have

$$\mathbf{P}_{n}(z_{1}) = \frac{\underline{m}_{2n0}(z_{1})}{z(1+\underline{m}_{2n0}(z_{1}))} \mathbf{V}_{2} \mathbf{V}_{2}^{*} - \frac{1}{z_{1}} \mathbf{I}_{p},$$

then

and

$$\begin{split} \mathbf{\Gamma}^* \mathbf{P}_n^2(z_1) \mathbf{\Gamma} = & \mathbf{U}_2 \mathbf{V}_2^* (\frac{\underline{m}_{2n0}(z_1)}{z_1 \left(1 + \underline{m}_{2n0}(z_1)\right)} \mathbf{V}_2 \mathbf{V}_2^* - \frac{1}{z_1} \mathbf{I}_p) (\frac{\underline{m}_{2n0}(z_1)}{z_1 \left(1 + \underline{m}_{2n0}(z_1)\right)} \mathbf{V}_2 \mathbf{V}_2^* - \frac{1}{z_1} \mathbf{I}_p) \mathbf{V}_2 \mathbf{U}_2^*, \\ = & \mathbf{U}_2 \mathbf{V}_2^* (\frac{\underline{m}_{2n0}^2(z_1)}{z_1^2 \left(1 + \underline{m}_{2n0}(z_1)\right)^2} \mathbf{V}_2 \mathbf{V}_2^* - \frac{2\underline{m}_{2n0}(z_1)}{z_1^2 \left(1 + \underline{m}_{2n0}(z_1)\right)} \mathbf{V}_2 \mathbf{V}_2^* + \frac{1}{z_1^2} \mathbf{I}_p) \mathbf{V}_2 \mathbf{U}_2^*, \\ = & \mathbf{U}_2 (\frac{\underline{m}_{2n0}^2(z_1)}{z_1^2 \left(1 + \underline{m}_{2n0}(z_1)\right)^2} \mathbf{I}_{p-M} - \frac{2\underline{m}_{2n0}(z_1)}{z_1^2 \left(1 + \underline{m}_{2n0}(z_1)\right)} \mathbf{I}_{p-M} + \frac{1}{z_1^2} \mathbf{I}_{p-M}) \mathbf{U}_2^*, \\ = & \frac{1}{z_1^2 \left(1 + \underline{m}_{2n0}(z_1)\right)^2} \mathbf{U}_2 \mathbf{U}_2^*. \end{split}$$

Therefore,

$$\begin{split} \sum_{i=1}^{p} [\mathbf{\Gamma}^{*} \mathbf{P}_{n}^{2}(z_{1}) \mathbf{\Gamma}]_{ii} [\mathbf{\Gamma}^{*} \mathbf{P}_{n}^{2}(z_{2}) \mathbf{\Gamma}]_{ii} &= \sum_{i=1}^{p} [\frac{1}{z_{1}^{2} \left(1 + \underline{m}_{2n0}(z_{1})\right)^{2}} \mathbf{U}_{2} \mathbf{U}_{2}^{*}]_{ii} [\frac{1}{z_{2}^{2} \left(1 + \underline{m}_{2n0}(z_{2})\right)^{2}} \mathbf{U}_{2} \mathbf{U}_{2}^{*}]_{ii}. \end{split}$$
Since
$$[\frac{1}{z_{1}^{2} \left(1 + \underline{m}_{2n0}(z_{1})\right)^{2}} \mathbf{U}_{2} \mathbf{U}_{2}^{*}]_{ii} &= [\frac{1}{z_{1}^{2} \left(1 + \underline{m}_{2n0}(z_{1})\right)^{2}} \mathbf{I}_{p} - \frac{1}{z_{1}^{2} \left(1 + \underline{m}_{2n0}(z_{1})\right)^{2}} \mathbf{U}_{1} \mathbf{U}_{1}^{*}]_{ii} \\ &= \frac{1 - \sum_{j=1}^{M} |u_{ij}|^{2}}{z_{1}^{2} \left(1 + \underline{m}_{2n0}(z_{1})\right)^{2}}, \end{split}$$

then

$$\sum_{i=1}^{p} \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_1) \mathbf{\Gamma} \right]_{ii} \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_2) \mathbf{\Gamma} \right]_{ii} = \sum_{i=1}^{p} \frac{\left(1 - \sum_{j=1}^{M} |u_{ij}|^2 \right)^2}{z_1^2 z_2^2 \left(1 + \underline{m}_{2n0}(z_1) \right)^2 \left(1 + \underline{m}_{2n0}(z_2) \right)^2},$$
$$= \frac{p - 2M + \sum_{i=1}^{p} \left(\sum_{j=1}^{M} |u_{ij}| \right)^2}{z_1^2 z_2^2 \left(1 + \underline{m}_{2n0}(z_1) \right)^2 \left(1 + \underline{m}_{2n0}(z_2) \right)^2} = \frac{p - 2M + \sum_{j_1, j_2 = 1}^{M} \mathcal{U}_{j_1 j_1 j_2 j_2}}{z_1^2 z_2^2 \left(1 + \underline{m}_{2n0}(z_1) \right)^2 \left(1 + \underline{m}_{2n0}(z_2) \right)^2}.$$

Then

$$\Theta_{2,n}(z_1, z_2) = \frac{z_1^2 z_2^2 \underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{n} \sum_{i=1}^{p} \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_1) \mathbf{\Gamma} \right]_{ii} \left[\mathbf{\Gamma}^* \mathbf{P}_n^2(z_2) \mathbf{\Gamma} \right]_{ii},$$

$$= \frac{z_1^2 z_2^2 \underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{n} \frac{p - 2M + \sum_{j_1, j_2=1}^{M} \mathcal{U}_{j_1 j_1 j_2 j_2}}{z_1^2 z_2^2 \left(1 + \underline{m}_{2n0}(z_1)\right)^2 \left(1 + \underline{m}_{2n0}(z_2)\right)^2}$$

$$= \frac{p - 2M + \sum_{j_1, j_2=1}^{M} \mathcal{U}_{j_1 j_1 j_2 j_2}}{n} \frac{\underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{\left(1 + \underline{m}_{2n0}(z_1)\right)^2 \left(1 + \underline{m}_{2n0}(z_2)\right)^2}.$$

Since the covariance of bulk part is $-\frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} (z_1 - \log(z_1) - 1) (z_2 - \log(z_2) - 1) \vartheta_n^2 dz_1 dz_2$, where $\vartheta_n^2 = \Theta_{0,n}(z_1, z_2) + \alpha_x \Theta_{1,n}(z_1, z_2) + \beta_x \Theta_{2,n}(z_1, z_2)$. By contour integral calculations, we obtain the covariance equals $-\log(1 - c_{nM}) - c_{nM} + \alpha_x (-\log(1 - \tilde{c}) - \tilde{c})$, where $\tilde{c} = \frac{p - 2M + \sum_{s,t=1}^p \left(\sum_{i=1}^M u_{si} \bar{u}_{ti}\right)^2}{n}$. Since $\tilde{c} - c_{nM} \to 0$ as $n \to \infty$, therefore $\vec{\varsigma}_{l}^{2} = \sum_{k=1}^{K} \frac{(\phi_{n}(\alpha_{k}) - 1)^{2}}{n} s_{k}^{2} + (\alpha_{x} + 1) \left(-\log(1 - c_{nM}) - c_{nM}\right),$ and

$$\frac{L - p \int f_L(x) dF^{c_n, H_n}(x) - \breve{\mu}_l}{\breve{\varsigma}_l} \xrightarrow{d} N(0, 1).$$

The proof of Theorem 4.1 is finished.

7.4. *Proof of Theorem* 4.2. First, we focus on the results under H_0 . From Lemma 9.1, we have

$$I_1(f_W) = c,$$

$$I_2(f_W) = c,$$

32

$$J_1(f_W, f_W) = 4c^3 + 2c^2,$$

$$J_2(f_W, f_W) = 4c^3,$$

which then yields

$$\mu_w = \alpha_x I_1(f_W) + \beta_x I_2(f_W) = \alpha_x c + \beta_x c,$$

$$\varsigma_w^2 = (\alpha_x + 1) J_1(f_W, f_W) + \beta_x J_2(f_W, f_W) = (\alpha_x + 1) (4c^3 + 2c^2) + 4\beta_x c^3.$$

The results are still valid if c is replaced by c_n . Moreover, the center term

(7.27)
$$\int f_W(x)dF^{c_n,H_n} = c_n,$$

is a direct result of Lemma 2.2 in Wang and Yao (2013). Therefore, from Zheng et al. (2015) or Wang and Yao (2013), we have

$$\frac{W - p \int f_W(x) dF^{c_n, H_n} - \mu_w}{\varsigma_w} \stackrel{d}{\longrightarrow} N(0, 1)$$

Then, we focus on the results under H_1 . Note that

$$Y = \int f_W(x) \, dG_n(x) - \sum_{k=1}^K d_k f_W(\phi_n(\alpha_k)) - \frac{M}{2\pi i} \oint_{\mathcal{C}} f_W(z) \, \frac{\underline{m}'_{2n0}(z)}{\underline{m}_{2n0}(z)} dz$$

After some calculations, we obtain

(7.28)

$$\int f_{W}(x) dG_{n}(x) = \operatorname{tr}(\mathbf{B} - \mathbf{I}_{p})^{2} - p \int f_{W}(x) dF^{c_{n}, H_{n}} = W - p \int f_{W}(x) dF^{c_{n}, H_{n}},$$

$$p \int f_{W}(x) dF^{c_{n}, H_{n}} = (p - M) \int f_{W}(x) dF^{c_{nM}, H_{2n}} = (p - M)c_{nM},$$

$$\sum_{k=1}^{K} d_{k} f_{W}(\phi_{n}(\alpha_{k})) = \sum_{k=1}^{K} d_{k} \left(\phi_{n}^{2}(\alpha_{k}) - 2\phi_{n}(\alpha_{k}) + 1\right),$$

$$\frac{M}{2\pi i} \oint_{\mathcal{C}} f_{W}(z) \frac{\underline{m}'_{2n0}(z)}{\underline{m}_{2n0}(z)} dz = -Mc_{nM}^{2}.$$

For consistency, we present the proof of (7.28) in Section 8. Therefore, from Theorem 3.1, we have

$$\frac{W - (p - M)\tilde{\ell}_w - \breve{\mu}_w}{\breve{\varsigma}_w} \xrightarrow{d} N(0, 1),$$

where

$$\check{\ell}_w = c_{nM}, \quad \check{\mu}_w = \alpha_x c_{nM} + \beta_x c_{nM} + \sum_{k=1}^K d_k \left(\phi_n^2 \left(\alpha_k\right) - 2\phi_n \left(\alpha_k\right) + 1\right) - M c_{nM}^2,$$
$$\check{\varsigma}_w^2 = -\frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} (z_1 - 1)^2 (z_2 - 1)^2 \vartheta_n^2 dz_1 dz_2 + \sum_{k=1}^K \frac{4\phi_n^2 \left(\alpha_k\right) \left(\phi_n \left(\alpha_k\right) - 1\right)^2}{n} s_k^2.$$

Since $\vartheta_n^2 = \Theta_{0,n}(z_1, z_2) + \alpha_x \Theta_{1,n}(z_1, z_2) + \beta_x \Theta_{2,n}(z_1, z_2)$, and $\Theta_{0,n}(z_1, z_2)$, $\Theta_{1,n}(z_1, z_2)$, and $\Theta_{2,n}(z_1, z_2)$ are calculated in the proof of Theorem 4.1, then by some calculations we obtain $-\frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} (z_1 - 1)^2 (z_2 - 1)^2 \vartheta_n^2 dz_1 dz_2$ equals $4c_{nM}^3 + 2c_{nM}^2 + \alpha_x \left(4\tilde{c}^3 + 2\tilde{c}^2\right) + 2\tilde{c}^2$

 $4\beta_x\check{c}^3, \text{ where } \check{c} = \frac{p-2M + \sum_{j_1, j_2=1}^M \mathcal{U}_{j_1 j_1 j_2 j_2}}{n}. \text{ Since } \tilde{c} - c_{nM} \to 0, \check{c} - c_{nM} \to 0 \text{ as } n \to \infty, \text{ therefore}$

$$\vec{\varsigma}_w^2 = \sum_{k=1}^K \frac{4\phi_n^2 \left(\alpha_k\right) \left(\phi_n \left(\alpha_k\right) - 1\right)^2}{n} s_k^2 + \left(\alpha_x + 1\right) \left(4c_{nM}^3 + 2c_{nM}^2\right) + 4\beta_x c_{nM}^3,$$

and

$$\frac{W - (p - M)\breve{\ell}_W - \breve{\mu}_w}{\breve{\varsigma}_w} \longrightarrow N(0, 1),$$

then the proof is finished.

7.5. *Proof of Theorem* 4.3. Let ξ be the significance level, and z_{ξ} is the $1 - \xi$ quantile of the standard Gaussian distribution Φ . Since

$$\xi = P\left(L > z_{\xi}\varsigma_l + p\ell_l + \mu_l\right),$$

for brevity, we denote $L_0 = p\ell_l + \mu_l$, $L_1 = (p - M)\check{\ell}_l + \check{\mu}_l$. Therefore, the power to detect the hypothesis is

$$P_L = P\left(L > z_{\xi\varsigma l} + L_0\right) = P\left(\frac{L - L_1}{\breve{\varsigma}_l} > \frac{z_{\xi\varsigma l} + L_0 - L_1}{\breve{\varsigma}_l}\right)$$

Since $\frac{L-L_1}{\zeta_l}$ is asymptotically normal distributed, then P_L is approximate to $\Phi\left(\frac{L_1-L_0}{\zeta_l}-z_{\xi}\frac{\zeta_l}{\zeta_l}\right)$. After some elementary calculations, we obtain as $n \to \infty$,

$$L_1 - L_0 \to -Mc + \sum_{k=1}^K d_k (\phi_k - \log \phi_k - 1),$$
$$\varsigma_l \to \sqrt{(\alpha_x + 1)(-\log(1 - c) - c)},$$

$$\breve{\varsigma}_l - \sqrt{(\alpha_x + 1)(-\log(1 - c) - c) + \sum_{k=1}^K \frac{(\phi_n(\alpha_k) - 1)^2}{n} s_k^2} \to 0.$$

Therefore, we have as n tends to infinity,

(7.29)
$$P_L - \Phi\left(\frac{\sum_{k=1}^K d_k \left(\phi_n(\alpha_k) - \log \phi_n(\alpha_k)\right) - M - Mc - z_{\xi}\varsigma_l}{\sqrt{(\alpha_x + 1)(-\log(1 - c) - c) + \sum_{k=1}^K \frac{\left(\phi_n(\alpha_k) - 1\right)^2}{n}s_k^2}}\right) \to 0,$$

then the proof of Theorem 4.3 is finished.

7.6. Proof of Theorem 4.4. Since $\xi = P(W > z_{\xi}\varsigma_w + c_n(p + \alpha_x + \beta_x))$, for brevity, we use the notation $W_0 = pc_n + c_n(\alpha_x + \beta_x)$, $W_1 = (p - M) \int f_W(x) dF^{c_{nM}, H_{2n}} + \breve{\mu}_w$. Therefore, the power to detect the hypothesis is

$$P_W = P\left(W > z_{\xi}\varsigma_w + W_0\right) = P\left(\frac{W - W_1}{\breve{\varsigma}_w} > \frac{z_{\xi}\varsigma_w + W_0 - W_1}{\breve{\varsigma}_w}\right)$$

Since $\frac{W-W_1}{\zeta_w}$ is asymptotically normal distributed, then P_W is approximated to $\Phi(\frac{W_1-W_0}{\zeta_w} - z_{\xi}\frac{\zeta_w}{\zeta_w})$. Here $W_1 - W_0 = (p-M)c_{nM} - pc_n + (\beta_x + \alpha_x)c_{nM} - (\beta_x + \alpha_x)c_n + \sum_{k=1}^{K} d_k(\phi_n(\alpha_k) - 1)^2 - Mc_{nM}^2$. After some elementary calculations, we obtain, as $n \to \infty$,

$$W_1 - W_0 \to \sum_{k=1}^{K} d_k \, (\phi_n \, (\alpha_k) - 1)^2 - Mc^2 - 2Mc,$$

$$\varsigma_w \to \sqrt{(\alpha_x + 1)(4c^3 + 2c^2) + 4\beta_x c^3},$$

$$\breve{\varsigma}_w - \sqrt{(\alpha_x + 1)(4c^3 + 2c^2) + 4\beta_x c^3 + \sum_{k=1}^K \frac{4\phi_n^2(\alpha_k) (\phi_n(\alpha_k) - 1)^2}{n} s_k^2} \to 0.$$

Then we have as n tends to infinity,

$$P_{W} - \Phi\left(\frac{\sum_{k=1}^{K} d_{k} (\phi_{k} - 1)^{2} - Mc^{2} - 2Mc - z_{\xi}\varsigma_{w}}{\sqrt{(\alpha_{x} + 1)(4c^{3} + 2c^{2}) + 4\beta_{x}c^{3} + \sum_{k=1}^{K} \frac{4\phi_{n}^{2} (\alpha_{k}) (\phi_{n} (\alpha_{k}) - 1)^{2}}{n}s_{k}^{2}}\right) \to 0,$$

then the proof is finished.

7.7. *Proof of Theorem 4.5.* From Jiang and Bai (2021), for spike α_1 , we eliminate the multiplicity of it and then we have

$$\sqrt{\frac{n\theta_1^2}{2\theta_1 + \sum_{t=1}^p |u_{t1}|^4 \beta_x \nu_1}} \frac{\lambda_1 - \phi_n(\alpha_1)}{\phi_n(\alpha_1)} \xrightarrow{d} N(0, 1)$$

Then the power of test R equals

$$P_{R} = P\left(\lambda_{1} > t_{\xi}\varsigma_{r} + \mu_{r}\right)$$

$$= P\left(\sqrt{\frac{n\theta_{1}^{2}}{2\theta_{1} + \sum_{t=1}^{p}|u_{t1}|^{4}\beta_{x}\nu_{1}}}\frac{\lambda_{1} - \phi_{n}\left(\alpha_{1}\right)}{\phi_{n}\left(\alpha_{1}\right)} > \sqrt{\frac{n\theta_{1}^{2}}{2\theta_{1} + \sum_{t=1}^{p}|u_{t1}|^{4}\beta_{x}\nu_{1}}}\frac{t_{\xi}\varsigma_{r} + \mu_{r} - \phi_{n}\left(\alpha_{1}\right)}{\phi_{n}\left(\alpha_{1}\right)}\right)$$

Since $\sqrt{\frac{n\theta_1^2}{2\theta_1 + \sum_{t=1}^{p} |u_{t1}|^4 \beta_x \nu_1}} \frac{\lambda_1 - \phi_n(\alpha_1)}{\phi_n(\alpha_1)}$ is asymptotically standard normal distributed, then P_R is approximate to $1 - \Phi\left(\sqrt{\frac{n\theta_1^2}{2\theta_1 + \sum_{t=1}^{p} |u_{t1}|^4 \beta_x \nu_1}} \frac{t_{\xi}\varsigma_r + \mu_r - \phi_n(\alpha_1)}{\phi_n(\alpha_1)}\right)$, and it equals $\Phi\left(-\sqrt{n\frac{t_{\xi}\varsigma_r + \mu_r - \phi_n(\alpha_1)}{s_1\phi_n(\alpha_1)}}\right)$, then the proof is finished.

8. Some deviations and calculations. This section contains proof of formulas stated in the proof of Theorems 4.1 and 4.2. We begin by deriving formula (7.26). First, we consider $\oint_{\mathcal{C}} f_L(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz$.

$$\oint_{\mathcal{C}} f_L(z) \, \frac{\underline{m}'(z)}{\underline{m}(z)} dz = \oint_{\mathcal{C}} f_L(z) \, d\log \underline{m}(z) = -\oint_{\mathcal{C}} f'_L(z) \log \underline{m}(z) \, dz$$

(8.31)
$$= \int_{a(c)}^{b(c)} f'_{L}(z) \left[\log \underline{m}(x+i\varepsilon) - \log \underline{m}(x-i\varepsilon) \right] dx$$
$$= 2i \int_{a(c)}^{b(c)} f'_{L}(z) \Im \log \underline{m}(x+i\varepsilon) dx$$

Here, $a(c) = (1 - \sqrt{c})^2$ and $b(c) = (1 + \sqrt{c})^2$. Since

$$\underline{m}(z) = -\frac{1-c}{z} + cm(z),$$

under H_1 , we have

$$\underline{m}(z) = \frac{-(z+1-c) + \sqrt{(z-1-c)^2 - 4c}}{2z}.$$

As $z \to x \in [a(c), b(c)]$, we obtain

$$\underline{m}(x) = \frac{-(x+1-c) + \sqrt{4c - (x-1-c)^2}i}{2x}.$$

Therefore,

$$\begin{split} &\int_{a(c)}^{b(c)} f'_{L}(z) \Im \log \underline{m}(x+i\varepsilon) dx \\ &= \int_{a(c)}^{b(c)} f'_{L}(x) \tan^{-1} \left(\frac{\sqrt{4c - (x-1-c)^{2}}}{-(x+1-c)} \right) dx \\ &= \left[\tan^{-1} \left(\frac{\sqrt{4c - (x-1-c)^{2}}}{-(x+1-c)} \right) f_{L}(x) \Big|_{a(c)}^{b(c)} - \int_{a(c)}^{b(c)} f_{L}(x) d \tan^{-1} \left(\frac{\sqrt{4c - (x-1-c)^{2}}}{-(x+1-c)} \right) \right] . \end{split}$$

It is easy to verify that the first term is 0, and we now focus on the second term,

(8.32)
$$\int_{a(c)}^{b(c)} f_L(x) d \tan^{-1} \left(\frac{\sqrt{4c - (x - 1 - c)^2}}{-(x + 1 - c)} \right) = \int_{a(c)}^{b(c)} \frac{(x - \log x - 1)}{1 + \frac{4c - (x - 1 - c)^2}{(x + 1 - c)^2}} \cdot \frac{\sqrt{4c - (x - 1 - c)^2} + \frac{(x - 1 - c)(x + 1 - c)}{\sqrt{4c - (x - 1 - c)^2}}}{(x + 1 - c)^2} dx.$$

By substituting $x = 1 + c - 2\sqrt{c}\cos(\theta)$, we obtain

$$(8.32) = \frac{1}{2} \int_0^{2\pi} \left(1 + c - 2\sqrt{c}\cos(\theta) - \log\left(1 + c - 2\sqrt{c}\cos(\theta)\right) - 1 \right) \frac{c - \sqrt{c}\cos(\theta)}{1 + c - 2\sqrt{c}\cos(\theta)} d\theta$$
$$= \frac{1}{2} \int_0^{2\pi} \left[1 - \frac{\log\left(1 + c - 2\sqrt{c}\cos(\theta)\right) + 1}{1 + c - 2\sqrt{c}\cos(\theta)} \right] \left(c - \sqrt{c}\cos(\theta) \right) d\theta$$

(8.33)

$$= \frac{1}{2} \int_0^{2\pi} \left(c - \sqrt{c} \cos(\theta) \right) d\theta - \frac{1}{2} \int_0^{2\pi} \frac{\log\left(1 + c - 2\sqrt{c} \cos(\theta)\right)}{1 + c - 2\sqrt{c} \cos(\theta)} \left(c - \sqrt{c} \cos(\theta) \right) d\theta - \frac{1}{2} \int_0^{2\pi} \frac{c - \sqrt{c} \cos(\theta)}{1 + c - 2\sqrt{c} \cos(\theta)} d\theta$$

It is easy to obtain that the first term of (8.33) is πc ; then, we consider the second term. By substituting $\cos \theta = \frac{z+z^{-1}}{2}$, we turn it into a contour integral on |z| = 1

$$\begin{split} &\frac{1}{2} \int_{0}^{2\pi} \frac{\log\left(1+c-2\sqrt{c}\cos(\theta)\right)}{1+c-2\sqrt{c}\cos(\theta)} \left(c-\sqrt{c}\cos\theta\right) d\theta \\ &= \frac{1}{2} \oint_{|z|=1} \log|1-\sqrt{c}z|^2 \cdot \frac{c-\sqrt{c}\frac{z+z^{-1}}{2}}{1+c-2\sqrt{c}\cdot\frac{z+z^{-1}}{2}} \frac{dz}{iz} \\ &= \frac{1}{4i} \oint_{|z|=1} \log|1-\sqrt{c}z|^2 \cdot \frac{2cz-\sqrt{c}\left(z^2+1\right)}{(z-\sqrt{c})(-\sqrt{c}z+1)z} dz \end{split}$$

When c < 1, 0 and \sqrt{c} are poles, by using the residue theorem, the integral is $-\pi \log(1-c)$. The same argument also holds for the third term, and the integral is 0 after some calculation. Therefore,

$$\frac{M}{2\pi i} \oint_{\mathcal{C}} f_L(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz = -M(c + \log(1-c)),$$

and the result is still valid if c is replaces c_{nM} ; therefore, formula (7.26) holds. Now, we prove (7.28). Since $z = -\frac{1}{\underline{m}} + \frac{c}{1+\underline{m}}$, we have, for c > 1,

$$\oint_{\mathcal{C}} f_W(z) \, \frac{\underline{m}'(z)}{\underline{m}(z)} dz = \oint_{\mathcal{C}_1} f_W(z) \, \frac{\underline{m}'(z)}{\underline{m}(z)} dz + \oint_{\mathcal{C}_2} f_W(z) \, \frac{\underline{m}'(z)}{\underline{m}(z)} dz,$$

where C_1 is a contour that includes the interval $((1 - \sqrt{c})^2, (1 + \sqrt{c})^2)$, and C_2 is a contour that includes the origin. Using $C_{\underline{m}}$ to denote the contour of \underline{m} , we obtain

$$\oint_{\mathcal{C}_1} f_W(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz = \oint_{\mathcal{C}_{\underline{m}}} (-\frac{1}{\underline{m}} + \frac{c}{1+\underline{m}} - 1)^2 \frac{\underline{m}'(z)}{\underline{m}(z)} \frac{dz}{d\underline{m}} d\underline{m}$$
$$= \oint_{\mathcal{C}_{\underline{m}}} (-\frac{1+\underline{m}}{\underline{m}} + \frac{c}{1+\underline{m}})^2 \frac{1}{\underline{m}} d\underline{m} = \oint_{\mathcal{C}_{\underline{m}}} (\frac{(1+\underline{m})^2}{\underline{m}^3} + \frac{c^2}{(1+\underline{m})^2\underline{m}} - \frac{2c}{\underline{m}^2}) d\underline{m}$$

Since the z contour cannot enclose the origin, neither can the resulting m contour. Thus, the only pole is -1, the residue is $-c^2$ by residue theorem, and we obtain the integral as $-2\pi i c^2$.

Then, we focus on the second integral $\oint_{\mathcal{C}_2} f_W(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz$. When z = 0, we obtain $\underline{m} = \frac{1}{c-1}$; since c > 1, $\frac{1}{c-1} > 0$. Both $\underline{m} = 0$ and $\underline{m} = -1$ are not in the contour. Thus, the integrand $(\frac{(1+\underline{m})^2}{\underline{m}^3} + \frac{c^2}{(1+\underline{m})^2\underline{m}} - \frac{2c}{\underline{m}^2})$ is analytic in the contour. The integral is 0. Therefore, when c > 1, $\frac{M}{2\pi i} \oint_{\mathcal{C}} f_W(z) \frac{\overline{m'(z)}}{m(z)} dz = -Mc^2$. When c < 1, the contour integral $\oint_{\mathcal{C}} f_W(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz$ reduces to $\oint_{\mathcal{C}_1} f_W(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz$, and the result is also the same as above. When c = 1, the result is still true by continuity in c. The results above are still valid if c is replaced by c_{nM} . Therefore, the proof of (7.28) is complete.

CLT FOR LSS WITH DIVERGING SPIKES

			C	and Unifor	m distribut	ions				
			(Dt_1, H_1))		(Dt_2, H_1)		(Dt_3, H_1)		
test	(p,n)	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$
\overline{CLRT}	(50,150)	0.5142	0.9954	1	0.4896	0.9896	1	0.5130	0.9995	1
	(100,300)	0.5056	0.9992	1	0.4954	0.9972	1	0.5134	0.9999	1
	(200,600)	0.5178	0.9996	1	0.5085	0.9990	1	0.5210	1	1
CNTT	(50,150)	0.9814	1	1	0.9287	1	1	0.9984	1	1
	(100,300)	0.9938	1	1	0.9662	1	1	0.9995	1	1
	(200,600)	0.9985	1	1	0.9873	1	1	1	1	1
RLRT	(50,150)	0.9983	1	1	0.9947	1	1	1	1	1
	(100,300)	1	1	1	1	1	1	1	1	1
	(200,600)	1	1	1	1	1	1	1	1	1

 TABLE 7

 Empirical probability of rejecting H_1 at significance level $\xi = 0.01$ under assumptions of Gaussian, Gamma, and Uniform distributions

9. Some useful lemmas.

LEMMA 9.1. If $\mathbf{D}_2 = \mathbf{I}_{p-M}$, then the mean function μ_1 and κ_{nst} in the covariance function of Theorem 3.2 can be simplified from the results in Wang and Yao (2013) and Zheng et al. (2015), i.e.,

$$\mu_1 = \alpha_x I_1(f_1) + \beta_x I_2(f_1),$$

$$\kappa_{nst} = (\alpha_x + 1)J_1(f_s, f_t) + \beta_x J_2(f_s, f_t),$$

$$\begin{split} I_{1}(f_{1}) &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} f_{1} \left(|1 + \sqrt{c_{nM}} z|^{2} \right) \left[\frac{z}{z^{2} - r^{-2}} - \frac{1}{z} \right] dz, \\ I_{2}(f_{1}) &= \frac{1}{2\pi i} \oint_{|z|=1} f_{1} \left(|1 + \sqrt{c_{nM}} z|^{2} \right) \frac{1}{z^{3}} dz, \\ J_{1}(f_{s}, f_{t}) &= \lim_{r \downarrow 1} \frac{-1}{4\pi^{2}} \oint_{|z_{1}|=1} \oint_{|z_{2}|=1} \frac{f_{s} \left(|1 + \sqrt{c_{nM}} z_{1}|^{2} \right) f_{t} \left(|1 + \sqrt{c_{nM}} z_{2}|^{2} \right)}{(z_{1} - rz_{2})^{2}} dz_{1} dz_{2}, \\ J_{2}(f_{s}, f_{t}) &= -\frac{1}{4\pi^{2}} \oint_{|z_{1}|=1} \frac{f_{s} \left(|1 + \sqrt{c_{nM}} z_{1}|^{2} \right)}{z_{1}^{2}} dz_{1} \oint_{|z_{2}|=1} \frac{f_{t} \left(|1 + \sqrt{c_{nM}} z_{2}|^{2} \right)}{z_{2}^{2}} dz_{2}. \end{split}$$

LEMMA 9.2. Note that for any matrix **Z**,

$$\mathbf{Z} \left(\mathbf{Z}^* \mathbf{Z} - \lambda \mathbf{I} \right)^{-1} \mathbf{Z}^* = \mathbf{I} + \lambda \left(\mathbf{Z} \mathbf{Z}^* - \lambda \mathbf{I} \right)^{-1}.$$

10. Tables for simulation studies. In this section, we present additional simulation tables regarding empirical probability of rejecting alternative hypotheses in Section 5 in the main file.

			6	ind Unifor	m distribut	tions								
	(Dt_1, H_2) (Dt_2, H_2) (Dt_3, H_2)													
test	(p,n)	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$				
\overline{CLRT}	(50,150)	0.9387	1	1	0.9045	1	1	0.9574	1	1				
	(100,300)	0.9496	1	1	0.9330	1	1	0.9666	1	1				
	(200,600)	0.9624	1	1	0.9553	1	1	0.9710	1	1				
CNTT	(50,150)	1	1	1	0.9998	1	1	1	1	1				
	(100,300)	1	1	1	1	1	1	1	1	1				
	(200,600)	1	1	1	1	1	1	1	1	1				
RLRT	(50,150)	1	1	1	1	1	1	1	1	1				
	(100,300)	1	1	1	1	1	1	1	1	1				
	(200,600)	1	1	1	1	1	1	1	1	1				

TABLE 8Empirical probability of rejecting H_2 at significance level $\xi = 0.01$ under assumptions of Gaussian, Gamma,and Uniform distributions

TABLE 9Empirical probability of rejecting H_4 at significance level $\xi = 0.05$ under assumptions of Gaussian, Gamma,
and Uniform distributions

			(Dt_1, H_4)			(Dt_2, H_4)		(Dt_3, H_4)		
test	(p,n)	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$
\overline{CLRT}	(50,150)	0.7288	0.9991	1	0.7074	0.9989	1	0.7269	0.9993	1
	(100,300)	0.7397	0.9999	1	0.7364	0.9997	1	0.7411	0.9997	1
	(200,600)	0.7568	1	1	0.7484	1	1	0.7649	1	1
CNTT	(50,150)	0.9917	1	1	0.9821	1	1	0.9974	1	1
	(100,300)	0.9985	1	1	0.9951	1	1	0.9997	1	1
	(200,600)	0.9997	1	1	0.9983	1	1	1	1	1
RLRT	(50,150)	0.9997	1	1	0.9992	1	1	0.9996	1	1
	(100,300)	1	1	1	1	1	1	1	1	1
	(200,600)	1	1	1	1	1	1	1	1	1

TABLE 10Empirical probability of rejecting H_4 at significance level $\xi = 0.01$ under assumptions of Gaussian, Gamma,
and Uniform distributions

			c	and Onigon	n aisin ioni	10115				
			(Dt_1, H_4))		(Dt_2, H_4))	(Dt_3, H_4)		
test	(p,n)	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$
\overline{CLRT}	(50,150)	0.5068	0.9967	1	0.5043	0.9952	1	0.5101	0.9966	1
	(100,300)	0.5137	0.9996	1	0.5082	0.9987	1	0.5193	0.9996	1
	(200,600)	0.5155	0.9995	1	0.5192	0.9997	1	0.5100	0.9998	1
CNTT	(50,150)	0.9782	1	1	0.9469	1	1	0.9934	1	1
	(100,300)	0.9935	1	1	0.9803	1	1	0.9988	1	1
	(200,600)	0.9984	1	1	0.9921	1	1	0.9997	1	1
RLRT	(50,150)	0.9978	1	1	0.9991	1	1	0.9981	1	1
	(100,300)	1	1	1	1	1	1	1	1	1
	(200,600)	1	1	1	1	1	1	1	1	1

CLT FOR LSS WITH DIVERGING SPIKES

and Uniform distributions											
		(Dt_1, H_5)			(Dt_2, H_5)			(Dt_3, H_5)			
test	(p,n)	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	
\overline{CLRT}	(50,150)	0.9803	1	1	0.9778	1	1	0.9819	1	1	
	(100,300)	0.9889	1	1	0.9864	1	1	0.9861	1	1	
	(200,600)	0.9928	1	1	0.9913	1	1	0.9908	1	1	
CNTT	(50,150)	1	1	1	1	1	1	1	1	1	
	(100,300)	1	1	1	1	1	1	1	1	1	
	(200,600)	1	1	1	1	1	1	1	1	1	
RLRT	(50,150)	1	1	1	1	1	1	1	1	1	
	(100,300)	1	1	1	1	1	1	1	1	1	
	(200,600)	1	1	1	1	1	1	1	1	1	

 TABLE 11

 Empirical probability of rejecting H_5 at significance level $\xi = 0.05$ under assumptions of Gaussian, Gamma, and Uniform distributions

TABLE 12Empirical probability of rejecting H_5 at significance level $\xi = 0.01$ under assumptions of Gaussian, Gamma,
and Uniform distributions

		(Dt_1, H_5)			(Dt_2, H_5)			(Dt_3, H_5)		
test	(p,n)	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$	$\alpha_1 = 3$	$\alpha_1 = 5$	$\alpha_1 = 7$
CLRT	(50,150)	0.9321	1	1	0.9218	1	1	0.9418	1	1
	(100,300)	0.9530	1	1	0.9521	1	1	0.9528	1	1
	(200,600)	0.9610	1	1	0.9586	1	1	0.9638	1	1
CNTT	(50,150)	1	1	1	0.9998	1	1	1	1	1
	(100,300)	1	1	1	1	1	1	1	1	1
	(200,600)	1	1	1	1	1	1	1	1	1
RLRT	(50,150)	1	1	1	1	1	1	1	1	1
	(100,300)	1	1	1	1	1	1	1	1	1
	(200,600)	1	1	1	1	1	1	1	1	1

TABLE 13Empirical probability of rejecting H_6 at significance level $\xi = 1 \times 10^{-4}$ under assumptions of Gaussian,
Gamma, and Uniform distributions

		(Dt_1, H_6)			(Dt_2, H_6)			(Dt_3, H_6)		
		α_1								
test	(p,n)	2.2	2.5	2.8	2.2	2.5	2.8	2.2	2.5	2.8
\overline{CLRT}	(50,150)	0.3876	0.8702	0.9943	0.6033	0.9347	0.9971	0.3881	0.8639	0.9946
	(100,300)	0.3849	0.8870	0.9980	0.6069	0.9616	0.9993	0.3856	0.8917	0.9985
	(200,600)	0.3761	0.9038	0.9978	0.6272	0.9698	0.9996	0.3842	0.9026	0.9990
CNTT	(50,150)	0.9974	1	1	0.9969	0.9999	1	0.9998	1	1
	(100,300)	0.9996	1	1	0.9994	1	1	1	1	1
	(200,600)	0.9998	1	1	1	1	1	1	1	1
RLRT	(50,150)	0.8845	0.9972	1	0.9111	0.9979	1	0.8718	0.9962	1
	(100,300)	0.9833	1	1	0.9858	1	1	0.9802	1	1
	(200,600)	0.9994	1	1	0.9995	1	1	0.9995	1	1

- Zhidong Bai and Jack W. Silverstein. CLT for linear spectral statistics of large-dimensional sample covariance matrices. *The Annals of Probability*, 32(1A):553 605, 2004.
- Zhidong Bai, Dandan Jiang, Jianfeng Yao, and Shurong Zheng. Corrections to LRT on large-dimensional covariance matrix by RMT. *The Annals of Statistics*, 37(6B):3822 – 3840, 2009.
- Dandan Jiang and Zhidong Bai. Generalized four moment theorem and an application to clt for spiked eigenvalues of high-dimensional covariance matrices. *Bernoulli*, 27(1):274–294, 2021.
- Zhijun Liu, Zhidong Bai, Jiang Hu and Haiyan Song. A CLT for the LSS of large dimensional sample covariance matrices with diverging spikes.
- Qinwen Wang and Jianfeng Yao. On the sphericity test with large-dimensional observations. *Electronic Journal* of Statistics, 7:2164–2192, 2013.
- Shurong Zheng, Zhidong Bai, and Jianfeng Yao. Substitution principle for CLT of linear spectral statistics of high-dimensional sample covariance matrices with applications to hypothesis testing. *The Annals of Statistics*, 43(2):546 – 591, 2015.