LOCALITY OF CRITICAL PERCOLATION ON EXPANDING GRAPH SEQUENCES

MICHAEL REN^{**} AND NIKE SUN^{*}

ABSTRACT. We study the locality of critical percolation on finite graphs: let G_n be a sequence of finite graphs, converging locally weakly to a (random, rooted) infinite graph G. Consider Bernoulli edge percolation: does the critical probability for the emergence of an infinite component on G coincide with the critical probability for the emergence of a linear-sized component on G_n ? In this short article we give a positive answer provided the graphs G_n satisfy an expansion condition, and the limiting graph G has finite expected root degree. The main result of Benjamini, Nachmias, and Peres (2011), where this question was first formulated, showed the result assuming the G_n satisfy a uniform degree bound and uniform expansion condition, and converge to a deterministic limit G. Later work of Sarkar (2021) extended the result to allow for a random limit G, but still required a uniform degree bound and uniform expansion for G_n . Our result replaces the degree bound on G_n with the (milder) requirement that G must have finite expected root degree. Our proof is a modification of the previous results, using a pruning procedure and the second moment method to control unbounded degrees.

1. INTRODUCTION

In this article we study the question of locality of critical percolation, as formulated by [\[BNP11\]](#page-11-0). Informally, if G_n is a graph sequence converging locally weakly to the random graph (G, ρ) , then does Bernoulli bond percolation have the same critical probability on G as on G_n ? The main result of [\[BNP11\]](#page-11-0) gives a positive answer in the case that the G_n have a uniform Cheeger constant $h > 0$ and a uniform degree bound $d < \infty$, and converges locally weakly to a deterministic limiting graph (G, ρ) . A later work [\[Sar21\]](#page-12-0) extends this result to the more general case of a random limiting graph (G, ρ) , but still requires that the graphs G_n satisfy uniform expansion and a uniform degree bound. In this article we extend the result of [\[Sar21\]](#page-12-0) by further relaxing the degree condition: more precisely, we show that the degree restriction on G_n can be replaced with the requirement that the limit (G, ρ) must have finite expected root degree.

1.1. Statement of main result. To formally state our main result, we set some notations and review some standard definitions (see [\[AL07\]](#page-11-1)). If v is a vertex in graph H , we let $B_R(v; H)$ be the R-neighborhood of v in H; we can regard $B_R(v; H)$ as a graph rooted at v.

Definition 1 (space of rooted graphs). Let \mathcal{C}_\star denote the space of connected locally finite rooted graphs *

$$
(G, \rho), modulo\ rooted\ isomorphism. A\ metric\ on\ \mathcal{G}_\star\ is\ given\ by
$$

$$
d\Big((G_1, \rho_1), (G_2, \rho_2)\Big) = \inf \left\{\frac{1}{1+R} : R \geqslant 0, B_R(\rho_1; G_1) \cong B_R(\rho_2; G_2) \right\} \in [0, 1]. \tag{1.1}
$$

The space \mathcal{G}_{\star} is separable and complete in this metric.

Definition 2 (local weak convergence). Suppose $G_n \equiv (V_n, E_n)$ is a sequence of finite graphs, and let U_n be a uniformly random vertex in G_n . We say G_n converges locally weakly to the (random) element $(G, \rho) \in$ \mathscr{G}_{\star} if $B_R(U_n; G_n)$ converges in law to (G, ρ) in the topology of \mathscr{G}_{\star} . This will be denoted $G_n \to_{lw} (G, \rho)$; we will often write simply $G_n \rightarrow_{lw} G$ with the understanding that G is a (random) rooted graph.

Date: December 13, 2022.

^{*}Massachusetts Institute of Technology. *University of Cambridge.

Definition 3 (expanding graph sequences). Given a finite graph $G = (V, E)$ and $0 < \delta \leq 1/2$, define

$$
h_{\delta}(G) \equiv \min \left\{ \frac{|E_G(A, V \setminus A)|}{|A|} : A \subseteq V, \delta \le \frac{|A|}{|V|} \le \frac{1}{2} \right\},\,
$$

where $E_G(A, B)$ denotes the set of all edges in G between the vertex sets A and B. We say that a sequence G_n
of finite graphs is **expanding** if for all $\delta > 0$ the hevelim inful $h(G) > 0$. That is to see, for all $\delta > 0$ of finite graphs is **expanding** if for all $\delta > 0$ we have lim inf_n $h_\delta(G_n) > 0$. That is to say, for all $\delta > 0$ there exist $n_{\delta} < \infty$ and $c_{\delta} > 0$ such that $h_{\delta}(G_n) \geq c_{\delta}$ for all $n \geq n_{\delta}$.

Definition 4 (critical percolation). Let G be an infinite graph, and let \mathbb{P}_p denote the law of a Bernoulli bond percolation (or edge percolation) configuration $\omega = \omega(G)$ on G. Then *

$$
p_c(G) = \sup_p \left\{ p \in [0, 1] : \mathbb{P}_p \left(\omega \text{ has no infinite component} \right) = 1 \right\}
$$

is the critical probability for Bernoulli bond percolation on G .

Theorem 1. Suppose that $G_n = (V_n, E_n)$ is an expanding sequence of finite graphs such that $G_n \to_{lw} (G, \rho)$ for some locally finite random infinite rooted graph (G, ρ) with $\mathbb{E}[\deg \rho] < \infty$. If $\omega(G_n)$ denotes edge percolation on G_n with probability p, then the following hold:

- (i) The critical edge percolation probability $p_c(G)$ is almost surely constant.
- (ii) If $p < p_c(G)$, then for any $\alpha > 0$, the probability that the largest component in $\omega(G_n)$ has size more than $\alpha |V_n|$ tends to 0 as $n \to \infty$.
- (iii) If $p > p_c(G)$, then there exists $\alpha > 0$ such that the probability that the largest component in $\omega(G_n)$ has size more than $\alpha|V_n|$ tends to 1 as $n \to \infty$.

Items (i)–(iii) can be summarized by saying that " $p_c((G_n)_{n\geq 0}) = p_c(G)$."

Theorem [1](#page-1-0) removes the uniform degree bound assumption from [\[BNP11\]](#page-11-0) and [\[Sar21\]](#page-12-0). Further, [\[BNP11,](#page-11-0) [Sar21\]](#page-12-0) assume **uniform expansion** ($h_{\delta}(G_n) \geq c_{\delta}$ for all $n \geq 1$ $n \geq 1$), while Theorem 1 uses only the weaker expansion assumption of Definition [3](#page-1-1) ($h_\delta(G_n) \geq c_\delta$ for all $n \geq n_\delta$). In fact, our argument for the unbounded degree setting involves pruning high-degree vertices, which can break a uniform expansion assumption.

Additionally, we remark that the locality result can fail without some kind of expansion condition. As an example,^{[1](#page-1-2)} let $h_n \to 0$, and let H be a graph on d/h_n vertices, uniformly random subject to the requirement
that all vertices have degree 2d event for one vertex $T_c(H)$ of degree d_c Let H_c H be disjoint copies that all vertices have degree 2d except for one vertex $v_*(H)$ of degree d. Let H_1, \ldots, H_n be disjoint copies of H, and let *J* be a random *d*-regular graph on the vertices $\{v(H_1), \ldots, v(H_n)\}\$. Let G_n be the union of H_1, \ldots, H_n , J, so it is a $(2d)$ -regular graph. It has poor expansion: if we take $A = H_1 \cup \ldots \cup H_k$, then there are at most kd edges between A and its complement, so

$$
\frac{|E_G(A, V \setminus A)|}{|A|} \leqslant \frac{kd}{kd/h_n} = h_n.
$$

Then G_n converges locally weakly to the (2*d*)-regular tree G , which has critical probability $p_c(G) = 1/(2d-1)$. On the other hand, the sequence of finite graphs has critical probability $p_c(G) \rightarrow \infty$ $1/(2d - 1)$. On the other hand, the sequence of finite graphs has critical probability $p_c((G_n)_{n\geqslant 0})$ $1/(d-1)$, as the emergence of a giant component on G_n is determined by percolation on J.

¹We learned of this example from Elchanan Mossel.

1.2. **Related work.** The locality question, as formulated by $[BNP11]$, was motivated by a related conjec-ture of Oded Schramm (see [\[BNP11,](#page-11-0) Conjecture 1.2]) for the setting of infinite, vertex transitive graphs G_n . This conjecture has since been proven in a few special cases, e.g., for uniformly nonemenable transitive graphs [\[SXZ14\]](#page-12-1), for abelian Cayley graphs [\[MT17\]](#page-11-2), for transitive graphs with exponential growth [\[Hut20\]](#page-11-3), and for transitive graphs with exponential growth [\[CMT22\]](#page-11-4).

Percolation on general finite graphs, initiated by [\[ABS04\]](#page-11-5), has also been well-studied. Beyond the existence of a percolation threshold for sequences of graphs [\[Fou07,](#page-11-6) [Sha13\]](#page-12-2), work has also been done to address the uniqueness of the giant component [\[ABS04,](#page-11-5) [EH21\]](#page-11-7) and scaling windows around the critical probability [\[NP10,](#page-11-8) [BCvdH](#page-11-9)⁺05]. We remark that many of these results which give detailed descriptions of behavior above, below, and near the threshold are about specific graph sequences such as random regular graphs or rely on strong assumptions about the graphs such as having uniformly bounded degrees. On the other hand, the results from [\[BNP11\]](#page-11-0) and our results apply more generally but only show the existence of a threshold for locally weakly convergent graph sequences.

In the case that G_n converges locally weakly to a (random, rooted) tree G , more can be said about the percolation threshold: it is well known that $p_c(G)$ is the reciprocal of the tree **branching number** br(G) [\[Lyo90\]](#page-11-10). More recently, physicists have proposed [\[KNZ14\]](#page-11-11) that $p_c((G_n)_{n\geq 0})$ can be estimated by the reciprocal of $\lambda^{NB}(G_n)$, the top eigenvalue of the nonbacktracking matrix of G_n . Our main result
Theorem 1 for the gase the local limit G is a tree implies $n((G_n)_{n-1}) = 1/\text{hr}(G)$. This paturelly suggests Theorem [1,](#page-1-0) for the case the local limit G is a tree, implies $p_c((G_n)_{n\geq 0}) = 1/\text{br}(G)$. This naturally suggests that one approach for proving the conjecture of [\[KNZ14\]](#page-11-11) would be to relate the asympotics of $\lambda(B_n)$ to the branching number of G_n .

1.3. Organization. In Section [2](#page-2-0) we prove some preliminary consequences of the local weak convergence assumption, and give our pruning argument. In Section [3](#page-5-0) we complete the proof of Theorem [1.](#page-1-0)

Acknowledgements. We thank Elchanan Mossel, Lenka Zdeborová, Jiaoyang Huang, and Fan Wei for helpful conversations. Research supported in part by NSF CAREER grant DMS-1940092.

2. Preliminaries

We fix some basic notation to be used throughout. In a graph H, let $d_H(u, v)$ be the distance between vertices u and v; we will sometimes write simply $d(u, v)$ if H is clear from context. Recall that $B_R(v; H)$ denotes the R-neighborhood of v in H , i.e., the subgraph of H induced by all vertices within distance R of v. More generally, if S is any subset of vertices in H, we write $B_R(S; H)$ for the subgraph of H induced by all vertices within distance R of S. We write $\partial B_R(v; H)$ for the set of vertices in H at exactly distance R from v .

We will also be working with a few different probability measures throughout the proof. Let \mathbb{P}_p and $\mathbb{P}_{n,p}$ denote edge percolation on G and G_n , respectively, with probability p. We use $\omega(G)$ and $\omega(G_n)$ to denote the percolation configurations. Let \mathbf{L}_n be the law of a uniform random vertex U_n of G_n , and let μ be the law of (G, ρ) .

2.1. **Continuity lemma.** It follows directly from the definition of local weak convergence (Definition [2\)](#page-0-0)

that if *f* is any bounded continuous function on the space
$$
\mathcal{G}_{\star}
$$
 (Definition 1), and $G_n \to_{lw} (G, \rho)$, then
\n
$$
\mathbb{E}_{\mathbf{L}_n} f(G_n, U_n) \xrightarrow{n \to \infty} \int f(G, \rho) d\mu.
$$
\n(2.1)

An immediate consequence is the following:

Lemma 2. Suppose $f(G, \rho)$ is a bounded function on \mathcal{G}_* that depends only on $B_R(\rho; G)$ for some finite R. Then f is continuous on \mathcal{G}_\star , so [\(2.1\)](#page-2-1) holds whenever $G_n \to_{lw} (G, \rho)$.

Proof. Recall from Definition [1](#page-0-1) that a metric on \mathcal{G}_\star is given by d from [\(1.1\)](#page-0-2). If

$$
d\Big((G_1,\rho_1),(G_2,\rho_2)\Big) \leq \frac{1}{1+R}
$$

then $B_R(\rho_1; G_1)$ and $B_R(\rho_2; G_2)$ agree, in which case the assumption on f implies $f(G_1, \rho_1) = f(G_2, \rho_2)$.
This shows that f is (uniformly) continuous on $\mathscr C$ This shows that f is (uniformly) continuous on \mathcal{G}_\star .

Corollary 3. If
$$
G_n \to_{lwc} (G, \rho)
$$
, then for any measurable event E we have

$$
\mathbf{L}_n \otimes \mathbb{P}_{n,p} \bigg(\omega(B_R(U_n; G_n)) \in E \bigg) \stackrel{n \to \infty}{\longrightarrow} \mu \otimes \mathbb{P}_p \bigg(\omega(B_R(\rho; G)) \in E \bigg).
$$

Proof. Define a function f on \mathscr{G}_{\star} by setting

$$
f(G,\rho) = \mathbb{P}_p\bigg(\omega(B_R(\rho;G)) \in E\bigg)
$$

ˆ

Then f clearly depends only on $B_R(\rho; G)$, so the claim follows from Lemma [2.](#page-2-2)

Lemma [2](#page-2-2) and Corollary [3](#page-3-0) will be used repeatedly throughout the proofs below.

2.2. **Pruning.** We will assume that the average degree of G_n is uniformly bounded, i.e., that there exists a constant d such that the average degree of G_n is at most d for all n . In the case that this does not hold, the result can be recovered from the former case via a pruning argument, which we now give. The main result of this subsection is the following:

Proposition 4. Suppose (G, ρ) is a (random) infinite rooted graph, locally finite, with $E(\deg \rho) < \infty$. Suppose G_n is an expanding sequence of finite graphs with $G_n \to_{loc} (G, \rho)$. Then, for any $\varepsilon > 0$, we can choose a sequence of subgraphs $\overline{\overline{G}}_n$ of \overline{G}_n such that (i) each $\overline{\overline{G}}_n$ has average degree at most $\overline{E}(\deg \rho) + \varepsilon$; and
(ii) the \overline{G} also form an apparenting asquange with $\overline{\overline{G}}$ and $(C, \alpha$ (ii) the \bar{G}_n also form an expanding sequence, with $\bar{G}_n \to_{lwc} (G, \rho)$.

Lemma 5. Suppose (G, ρ) is a (random) infinite rooted graph, locally finite, with $\mathbb{E}(\deg \rho) < \infty$. Suppose G_n is an expanding sequence of finite graphs with $G_n \to_{lw} (G, \rho)$. Let k_n be any sequence of integers with $k_n \to \infty$, and let $\overline{\tilde{G}}_n$ be the subgraph of G_n formed by removing all edges incident to vertices of degree at least k_n . Then \bar{G}_n is also an expanding sequence with $\bar{G}_n \to_{lwc} (G, \rho)$.

Proof. Let S_n denote the subset of all vertices in G_n with degree at least k_n . We first claim that for any fixed $R < \infty$,

$$
\lim_{n \to \infty} \frac{|B_R(S_n; G_n)|}{|V_n|} = 0.
$$
\n(2.2)

Indeed, if a vertex is within distance R of S_n , then it must have at least k_n vertices in its $(R + 1)$ neighborhood, so ˆˇ

$$
\frac{|B_R(S_n; G_n)|}{|V_n|} \leqslant \mathbf{L}_n\left(\left|B_{R+1}(U_n; G_n)\right| \geqslant k_n\right).
$$
\nif follows that for any finite k we have

Since $k_n \to \infty$ by assumption, it follows that for any finite k we have

$$
\limsup_{n\to\infty} \mathbf{L}_n\left(\left|B_{R+1}(U_n;G_n)\right|\geqslant k_n\right)\leqslant \limsup_{n\to\infty} \mathbf{L}_n\left(\left|B_{R+1}(U_n;G_n)\right|\geqslant k\right)
$$

Finally, by the assumption $G_n \to_{lw} (G, \rho) \sim \mu$ together with Lemma [2,](#page-2-2) we have

$$
\lim_{n\to\infty} \mathbf{L}_n\bigg(\Big|B_{R+1}(U_n;G_n)\Big| \geq k\bigg) = \mu\bigg(\Big|B_{R+1}(\rho,G)\Big| \geq k\bigg).
$$

Combining the above calculations gives

$$
\limsup_{n \to \infty} \frac{|B_R(S_n; G_n)|}{|V_n|} \le \mu\bigg(\Big|B_{R+1}(\rho, G)\Big| \ge k\bigg)
$$

for any finite k. In the above, the right-hand side can be made arbitrarily small by taking $k \to \infty$, while
the left hand side does not depend on k, so this proves the alsim (2.2) the left-hand side does not depend on k , so this proves the claim [\(2.2\)](#page-3-1).

We next argue that $\bar{G}_n \to_{lw} (G, \rho)$. The graphs G_n and \bar{G}_n have the same vertex set, so if U_n is a if \bar{G}_n is a set of \bar{G}_n is a set of \bar{G}_n is a set of \bar{G}_n . uniformly random vertex in G_n , then it is also a uniformly random vertex in \bar{G}_n . Moreover, it is clear that the subgraphs $B_R(U_n; G_n)$ and $B_R(U_n; \bar{G}_n)$ are the same as long as U_n does not belong to $B_R(S_n; G_n)$.
We see from (2.2) that U_n lise in $B_n(S_n; G_n)$ with probability $\Omega_n(1)$ as we senaly that $B_n(U_n; \bar{G}_n)$ also We see from [\(2.2\)](#page-3-1) that U_n lies in $B_R(S_n; G_n)$ with probability $o_n(1)$, so we conclude that $B_R(U_n; \bar{G}_n)$ also converges in law to $B_R(\rho; G)$. This proves that $\bar{G}_n \to_{lwc} (G, \rho)$, as claimed.

We now argue that \overline{G}_n is an expanding sequence. Given $\delta > 0$, let $A \subseteq V_n$ with $\delta \le |A|/|V_n| \le 1/2$. Since the original sequence G_n is assumed to be expanding, there exists $c_{\delta} > 0$ such that

$$
\frac{|E_{G_n}(A, V_n \backslash A)|}{|A|} \geq c_{\delta}
$$

as long as $n \geq n_\delta$. The graph \bar{G}_n differs from G_n only in the deletion of edges incident to S_n , and the total number of deleted edges can be upper bounded by $|B_1(S_n; G_n)|$. It follows using [\(2.2\)](#page-3-1) that

$$
\frac{|E_{\tilde{G}_n}(A, V_n \setminus A)|}{|A|} \geqslant \frac{|E_{G_n}(A, V_n \setminus A)|}{|A|} - \frac{|B_1(S_n; G_n)|}{|V_n|} \frac{|V_n|}{|A|} \geqslant \frac{|E_{G_n}(A, V_n \setminus A)|}{|A|} - \frac{o_n(1)}{\delta} \,,
$$

which is at least $c_{\delta}/2$ for *n* large enough. This shows that \bar{G}_n is also an expanding sequence, and this concludes the preaf of the large concludes the proof of the lemma.

Lemma 6. Let D_n be a nonnegative integer-valued random variable converging in law to an integrable random variable D. Then, for any $\varepsilon > 0$, it is possible to choose a sequence $k_n \to \infty$ such that $\mathbb{E}(D_n; D_n <$ k_n) \leq $ED + \varepsilon$ $< \infty$.

Proof. First we note that for any finite integer k ,

$$
\mathbb{E}(D; D < k) = \sum_{\ell \geq 1} \mathbb{P}\bigg(D\mathbf{1}\{D < k\} \geq \ell\bigg) = \sum_{\ell=1}^{k-1} \mathbb{P}(\ell \leq D < k) \leq \sum_{\ell=1}^{k-1} \mathbb{P}(D \geq \ell).
$$

It follows from the assumption that for any finite ℓ we have $\mathbb{P}(D_n \geq \ell) \to \mathbb{P}(D \geq \ell)$ as $n \to \infty$, and consequently $\mathbb{P}(D_n \geq \ell) \leq \mathbb{P}(D \geq \ell) + \varepsilon/2^{\ell}$ for all $n \geq m_{\ell}$, where m_{ℓ} is a finite integer depending on ℓ
and ε . We then set $k = n$ if $\mathbb{P}(D_n > \ell) \leq \mathbb{P}(D_n > \ell) + \varepsilon/2^{\ell}$ for all $\ell > 1$ an *

and
$$
\varepsilon
$$
. We then set $k_n \equiv n$ if $\mathbb{P}(D_n \ge \ell) \le \mathbb{P}(D \ge \ell) + \varepsilon/2^{\ell}$ for all $\ell \ge 1$ and otherwise
\n
$$
k_n \equiv \min \left\{ \ell \ge 1 : \mathbb{P}(D_n \ge \ell) > \mathbb{P}(D \ge \ell) + \frac{\varepsilon}{2^{\ell}} \right\} \ge \min \left\{ \ell \ge 1 : m_{\ell} > n \right\}.
$$

Our chosen k_n must tend to infinity as $n \to \infty$ since the m_ℓ are finite. The definition of k_n implies

$$
\mathbb{E}(D_n; D_n < k_n) \leq \sum_{\ell=1}^{k_n-1} \mathbb{P}(D_n \geq \ell) \leq \sum_{\ell=1}^{k_n-1} \left\{ \mathbb{P}(D \geq \ell) + \frac{\varepsilon}{2^{\ell}} \right\} \leq \mathbb{E}D + \varepsilon,
$$

as required. This finishes the proof. \Box

Proof of Proposition [4.](#page-3-2) This follows immediately by combining Lemmas [5](#page-3-3) and [6.](#page-4-0) \Box

3. Proof of Theorem [1](#page-1-0)

In this section we give the proof of Theorem [1.](#page-1-0) We prove parts (i), (ii), and (iii) respectively in $\S 3.1$, $\S 3.2$, and \S 3.3. As noted above, our proofs are based on arguments of [\[BNP11,](#page-11-0) [Sar21\]](#page-12-0), with modifications to handle unbounded degrees. With the aim of being self-contained, we present a full proof below, but will repeatedly point out similarities and differences with the prior works.

3.[1](#page-1-0). **Rerooting invariant functions.** The proof of Theorem 1 part (i), after applying Proposition [4,](#page-3-2) is very similar to the proof of the corresponding result in [\[Sar21\]](#page-12-0), but we have made some simplications to the main lemma. As in [\[Sar21\]](#page-12-0), the proof shows that any rerooting invariant function of G is almost surely constant, so the local weak limit of an expanding sequence of finite graphs with finite expected root degree is in fact an extremal unimodular graph (see Corollary [11](#page-7-1) below).

Proposition 7. Suppose that G_1 and G_2 are disjoint subsets of \mathcal{C}_\star such that each G_i is closed under rerooting. Then, for any compact subsets $H_i \subseteq G_i$ and any finite integer K, there exists a finite integer R such that for any $(G_1, \rho_1) \in H_1$ and $(G_2, \rho_2) \in H_2$, we have that $B_R(u_1, G_1)$ is not isomorphic to $B_R(u_2, G_2)$ for all $u_1 \in B_K(\rho_1, G_1), u_2 \in B_K(\rho_2, G_2).$

In other words, given a rooted graph from one of G_1 and G_2 , we can determine which set it came from by looking at the R -neighborhood of an arbitrary vertex in the K -neighborhood of the root. This was proven in [\[Sar21\]](#page-12-0) by considering finite open covers of the H_i . We now give a conceptually cleaner proof
of this feet of this fact.

Lemma 8. Recall the metric $d((G_1,\rho_1),(G_2,\rho_2))$ on \mathcal{G}_* defined by [\(1.1\)](#page-0-2). For any finite K, let $D_K:\mathcal{G}_*\times\mathcal{G}_*\to$ $\lceil 0, 1 \rceil$ be defined by *

$$
D_K((G_1,\rho_1),(G_2,\rho_2)) = \min \left\{d\Big((G_1,u_1),(G_2,u_2)\Big) : u_i \in B_K(\rho_i;G_i)\right\},\,
$$

where (G_i, u_i) denotes the graph (G_i, ρ_i) rerooted at u_i . The function D_K is continuous.

Proof. Since the graphs G_i are locally finite, the definition of D_K is a minimum over finitely many quantities. Therefore we can choose $u_{i\star} \in B_K(\rho_i; G_i)$ such that

$$
D_K\Big((G_1,\rho_1),(G_2,\rho_2)\Big)=\boldsymbol{d}\Big((G_1,u_{1\star}),(G_2,u_{2\star})\Big)\,.
$$
\n(3.1)

Now suppose that (H, σ) is any element of \mathcal{G}_* that lies within distance $\delta = 1/(1 + R)$ of (G_2, ρ_2) , where we assume $R \geq K$. From the definition [\(1.1\)](#page-0-2), it implies that there is a rooted isomorphism φ that maps $B_R(\rho_2;G_2)$ to $B_R(\sigma;H)$. Let $v_{2\star}\equiv \varphi(u_{2\star})$. We then have

$$
D_K\Big((G_1,\rho_1),(H,\sigma)\Big) \leq d\Big((G_1,u_{1\star}),(H,v_{2\star})\Big) \leq d\Big((G_1,u_{1\star}),(G_2,u_{2\star})\Big) + d\Big((G_2,u_{2\star}),(H,v_{2\star})\Big) \leq D_K\Big((G_1,\rho_1),(G_2,\rho_2)\Big) + \frac{1}{1+(R-K)}
$$

where the first step is by the definition of D_K , the second step is by the triangle inequality, and the last $\cos \theta = \sin(\theta + B)$. step is by [\(3.1\)](#page-5-2) together with the observation that $B_{R-K}(u_{2*}; G_2)$ must be isomorphic to $B_{R-K}(u_{2*}; H)$. By exchanging the roles of (G_2, ρ_2) and (H, σ) we must also have

$$
D_K\Big((G_1,\rho_1),(G_2,\rho_2)\Big)\leq D_K\Big((G_1,\rho_1),(H,\sigma)\Big)+\frac{1}{1+(R-K)},
$$

and combining the last two inequalities gives

$$
\left| D_K\Big((G_1,\rho_1),(G_2,\rho_2)\Big) - D_K\Big((G_1,\rho_1),(H,\sigma)\Big) \right| \leq \frac{1}{1 + (R - K)}.
$$

This shows that $D_K : \mathcal{G}_\star \times \mathcal{G}_\star \to [0, 1]$ is uniformly continuous in the second coordinate. By symmetry it is also uniformly continuous in the first coordinate which proves the eleim is also uniformly continuous in the first coordinate, which proves the claim. \Box

Proof of Proposition [7.](#page-5-3) By the assumption that the G_i are disjoint and closed under rerooting, the function D_K must be strictly positive on $\mathcal{L}_1 \times \mathcal{L}_2$, and hence also on $H_1 \times H_2$. Since D_K is continuous by Lemma [8,](#page-5-4) we conclude that the minimum of D_K on the compact set $H_1 \times H_2$ must be lower bounded by some $\delta > 0$. Then, as long as we have $\delta > 1/(1 + R)$, it follows from the definition of D_K that for any $(G_i, \rho_i) \in H_i$, the graphs $R_2(u \circ C_i)$ and $R_2(u \circ C_i)$ must be non-isomorphic for all $u \in R_2(\alpha \circ C_i)$. the graphs $B_R(u_1; G_1)$ and $B_R(u_2; G_2)$ must be non-isomorphic for all $u_i \in B_K(\rho_i; G_i)$.

As in [\[Sar21\]](#page-12-0), the rest of the proof proceeds by showing that if $p_c(G)$ can take on different values, then the root vertices in G_n that produce these different values in the limit can be connected by a short path, contradicting Proposition [7.](#page-5-3) This relies crucially on the expansion condition, which will be used in the application of the following classical result:

Theorem 9 (Menger's theorem). In a finite graph $G = (V, E)$, for any two disjoint subsets $A, B \subseteq V$, the minimum size of an A-B edge cut equals the maximum number of pairwise edge-disjoint paths from A to B.

Corollary 10. Suppose G_n is an expanding sequence of finite graphs with $G_n \to_{lw} (G, \rho)$, and such that each G_n has average degree at most $d < \infty$. Then $p_c(G)$ is almost surely constant.

Proof. This proof is essentially identical to that of $\lceil \text{Sar21}, \text{Theorem 1.1} \rceil$. The uniform degree bound is only used to bound the number of edges in the graph, so the argument can be directly adapted to when there is a uniform average degree bound.

Suppose to the contrary that $p_c(G)$ is not almost surely constant. Then there exist $0 \le a < b \le 1$ such that, if we define !
!) !
. \mathbf{r}

$$
\mathcal{G}^1 = \left\{ G : p_c(G) \leq a \right\}, \quad \mathcal{G}^2 = \left\{ G : p_c(G) \geq b \right\},
$$

then we will have a positive constant p_0 such that $\min \Big\{ \mu(\mathcal{G}^1) \Big\}$

$$
\min\left\{\mu(\mathcal{G}^1),\mu(\mathcal{G}^2)\right\}\geq p_0.
$$

)

It is clear that G^1 and G^2 are disjoint, and invariant under rerooting. Recall that \mathscr{G}_\star is a Polish space (see Definition [1\)](#page-0-1), and that all Borel measures on Polish spaces are inner regular. It follows that there exist compact subsets $H^i \subseteq \mathcal{G}^i$ such that)

$$
\min\left\{\mu(H^1), \mu(H^2)\right\} \geqslant \frac{p_0}{2} \,. \tag{3.2}
$$

Recall that G_n is assumed to be an expanding sequence in the sense of Definition [3,](#page-1-1) and let h be a positive constant such that

$$
\frac{1}{2}\liminf_{n\to\infty} h_{p_0/4}(G_n) \geq h.
$$
\n(3.3)

Recall that G_n is also assumed to have average degree at most d , and define

$$
K=\frac{4d}{hp_0}.
$$

Now apply Proposition [7](#page-5-3) with H^1 , H^2 , and K as defined above, and let R be the integer that results from
the conclusion of the proposition. If we define the greate the conclusion of the proposition. If we define the events *

$$
E^i = \left\{ (G, \rho) : B_{R+K}(\rho; G) \cong B_{R+K}(\rho^i; G^i) \text{ for some } (G^i, \rho^i) \in H^i \right\},\
$$

then E^1 and E^2 are disjoint, because Proposition [7](#page-5-3) guarantees in particular that $B_R(\rho^1; G^1) \not\cong B_R(\rho^2; G^2)$
for all $(G^i, g^i) \in H^i$. Let A^i be the set of vertices in G , such that (G, g^n) belongs to symptoge for all $(G^t, \rho^t) \in H^t$. Let A^t be the set of vertices in G_n such that (G_n, v) belongs to event E^t . Since the events E^1 and E^2 are disjoint, the vertex sets A^1 and A^1 are disjoint. Since membership in E^i depends only on the $(R + K)$ -neighborhood of the root vertex, Lemma [2](#page-2-2) gives

$$
\lim_{n\to\infty}\frac{|A^i|}{|V_n|}=\lim_{n\to\infty}\mathbf{L}_n\left((G_n,U_n)\in E^i\right)=\mu\Big((G,\rho)\in E^i\Big)\geqslant\mu(H^i)\stackrel{(3.2)}{\geqslant}\frac{p_0}{2}>0.
$$

Thus, for large enough *n* we have $|A^i|/|V_n| \ge p_0/4$. It follows from the expansion assumption [\(3.3\)](#page-6-1) that, again for *n* large enough, the minimum edge cut separating A^1 from A^2 has at least $\kappa_n = h p_0 |V_n|/4$ edges.
It follows by Mangar's theorem (Theorem 0) that there are at least κ , adge disjoint paths ioining A^1 It follows by Menger's theorem (Theorem [9\)](#page-6-2) that there are at least κ_n edge-disjoint paths joining A^1 to A^2 .
Since C , hes at most $|V|d/2$ edges in total, at least helf of the naths must have length at most Since G_n has at most $|V_n|d/2$ edges in total, at least half of the paths must have length at most

$$
\frac{|V_n|d/2}{\kappa_n/2}=\frac{4d}{hp_0}=K.
$$

However, even the existence of a single such path joining $v^1 \in A^1$ to $v^2 \in A^2$ results in a contradiction: it
implies that the *P* poishborhood of v^2 lies inside the $(P_+ + K)$ poishborhood of v^1 in C. The definit implies that the R-neighborhood of v^2 lies inside the $(R + K)$ -neighborhood of v^1 in G_n . The definition of R of A^i implies $B_{R+K}(v^i; G_n) \cong B_{R+K}(p^i; G^i)$ for some $(G^i, \rho^i) \in H_i$. This contradicts the definition of R
and concludes the proof and concludes the proof.

We remark that the above argument implies the result of $\lceil \text{Sar21}, \text{Theorem 1.1} \rceil$ under the milder assump-tions of Theorem [1.](#page-1-0) Any local weak limit of a sequence of finite graphs is a random rooted graph whose measure is unimodular (see [\[AL07,](#page-11-1) Definition 2.1]). The space of unimodular probability measures on \mathcal{E}_\star is convex, and its extreme points are said to be **extremal** or **ergodic**. A unimodular measure μ on \mathcal{G}_* is extremal if and only if $\mu(E) \in \{0, 1\}$ for any event E that is invariant under non-rooted isomorphisms $($ [\[AL07,](#page-11-1) Theorem 4.7]). It then follows from the above that the local weak limit of an expanding sequence of finite graphs is extremal if its root has finite expected degree:

Corollary 11. Under the conditions of Theorem [1,](#page-1-0) G is an extremal unimodular random graph.

Proof of Theorem [1](#page-1-0) part (i). By Proposition [4,](#page-3-2) there exists a expanding sequence of subgraphs \bar{G}_n of G_n and that each \bar{G}_n has every degree at most $\mathbb{F}(dog \Omega)$ + 1 and $\bar{G}_n \to (G, g)$. Anglying Carollary 10 such that each \bar{G}_n has average degree at most $\mathbb{E}(\deg \rho) + 1$ and $\bar{G}_n \to_{lw} (G, \rho)$. Applying Corollary [10](#page-6-3) to \bar{G}_n gives the result. \Box gives the result.

3.2. Subcritical percolation regime. In this subsection we prove Theorem [1](#page-1-0) part (ii), which says in short that $p_c(G_n) \geqslant p_c(G)$. The result is a straightforward consequence of Lemmas [2](#page-2-2) and Corollary [3.](#page-3-0) Let us point out that for this part of the proof we work with the original sequence G_n , without pruning, so we do not assume that G_n has bounded average degree. The reason for this is that (a priori) we only know that $p_c(\bar{G}_n) \geq p_c(G_n)$, so it is not sufficient to only prove $p_c(\bar{G}_n) \geq p_c(G)$.

Recall that we let ζ $= \zeta(G)$ denote an adso percelation configuration

Recall that we let $\omega = \omega(G)$ denote an edge percolation configuration on graph G; we also view ω as a (random) subgraph of G. Let $C(v) \equiv C(v; \omega)$ denote the connected component of ω that contains v. For any vertex subsets A and B, let $A \leftrightarrow_{\omega} B$ indicate that A and B are connected by a path using only edges in ω . We write simply $A \leftrightarrow B$ when ω is clear from context.

Proof of Theorem [1](#page-1-0) part (ii). The proof of $\lceil \text{Sar21}, \text{Corollary 1.2} \rceil$ (which is based on the proof of $\lceil \text{BNP11}, \rceil$ Theorem 1.3]) applies here verbatim. To be self-contained, we also give the argument here. Under the assumptions of Theorem [1,](#page-1-0) we will show that for any $p < p_c(G)$ and any $\alpha > 0$, the probability that the largest component in $\omega(G_n)$ contains more than $\alpha|V_n|$ vertices tends to zero as $n \to \infty$. Let C_n denote

the vertices in the largest component of
$$
\omega(G_n)
$$
. For any $R > 0$, we have the chain of inequalities
\n
$$
\alpha \mathbb{P}_{n,p}(|C_n| \ge \alpha |V_n|)
$$
\n
$$
\le \mathbf{L}_n \otimes \mathbb{P}_{n,p}(|C_n| \ge \alpha |V_n|, U_n \in C_n)
$$
\n
$$
\le \mathbf{L}_n \otimes \mathbb{P}_{n,p}(|C(U_n)| \ge \alpha |V_n|)
$$
\n
$$
\le \mathbf{L}_n \otimes \mathbb{P}_{n,p}(|C(U_n)| \ge |B_R(U_n, G_n)|) + \mathbf{L}_n(|B_R(U_n, G_n)| \ge \alpha |V_n|)
$$
\n
$$
\le \mathbf{L}_n \otimes \mathbb{P}_{n,p}(|U_n \leftrightarrow \partial B_R(U_n, G_n)|) + \mathbf{L}_n(|B_R(U_n, G_n)| \ge \alpha |V_n|).
$$
\n(3.4)

We will consider separately the two terms on the right-hand side of (3.4) . For the first term, Corollary [3](#page-3-0) directly implies

$$
\lim_{n\to\infty} \mathbf{L}_n \otimes \mathbf{P}_{n,p} \Big(U_n \leftrightarrow \partial B_R (U_n, G_n) \Big) = \mu \otimes \mathbf{P}_p \Big(\rho \leftrightarrow \partial B_R (\rho, G) \Big)
$$

For the second term on the right-hand side of (3.4) , for any finite K we have

$$
\limsup_{n\to\infty} \mathbf{L}_n\Big(|B_R(U_n;G_n)|\geq \alpha|V_n|\Big)\leq \lim_{n\to\infty} \mathbf{L}_n\Big(|B_R(U_n;G_n)|\geq K\Big)=\mu\Big(|B_R(\rho;G)|\geq K\Big)\,,
$$

where the last step is by Lemma [2.](#page-2-2) Since $|B_R(\rho; G)|$ is almost surely finite, the right-hand side tends to zero as $K \to \infty$. The left-hand side does not depend on K , so we conclude that in fact

$$
\lim_{n\to\infty} \mathbf{L}_n\Big(|B_R(U_n;G_n)|\geqslant \alpha|V_n|\Big)=0\,.
$$

Substituting these limits back into [\(3.4\)](#page-8-1) gives

$$
\limsup_{n\to\infty} \alpha \mathbb{P}_{n,p}\Big(|C_n| \geq \alpha |V_n|\Big) \leq \mu \otimes \mathbb{P}_p\Big(\rho \leftrightarrow \partial B_R(\rho, G)\Big)
$$

Since $p < p_c(G)$, the right-hand side tends to zero as $R \to \infty$. The left-hand side does not depend on R , so we finally conslude that so we finally conclude that

so we finally conclude that
$$
\lim_{n\to\infty} \mathbb{P}_{n,p}\left(|C_n| \geq \alpha |V_n|\right) = 0,
$$
 as claimed.

3.3. Supercritical percolation regime. In this subsection we prove Theorem [1](#page-1-0) part (iii), which says in short that $p_c(G_n) \leq p_c(G)$. We make use of Proposition [4](#page-3-2) to reduce to the case where the G_n have uniformly bounded average degree. The rest of the proof is adapted from that of [\[BNP11,](#page-11-0) Theorem 1.3], where the idea is to first percolate with a smaller probability $p - \varepsilon > p_c(G)$, then connect the resulting clusters into a linear-sized component with the remaining probability. While [\[BNP11\]](#page-11-0) used the bounded degree assumption to obtain a strong concentration bound for the number of percolation clusters, we will use Chebychev's inequality to obtain a weaker (but sufficient) bound for our more general setting. This is based on the following lemma:

Lemma 12. Under the assumptions of Theorem [1,](#page-1-0) we have for any fixed R that

$$
\lim_{n\to\infty}\frac{1}{|V_n|^2}\bigg|\Big\{(u,v)\in (V_n)^2:d(u,v)\leq R\Big\}\bigg|=0.
$$

Proof. We can bound the quantity of interest by

$$
r(n) = \frac{1}{|V_n|^2} \Biggl\{ (u, v) \in (V_n)^2 : d(u, v) \le R \Biggr\} = \sum_{v \in V_n} \frac{|B_R(v; G_n)|}{|V_n|^2}
$$

= $\frac{\mathbb{E}_{\mathbf{L}_n} |B_R(U_n; G_n)|}{|V_n|} \le \frac{K}{|V_n|} + \mathbf{L}_n \Bigl(|B_R(U_n, G_n)| \ge K \Bigr),$

where the last inequality holds for any finite K . It follows using Lemma [2](#page-2-2) that

$$
\limsup_{n\to\infty} r(n) \leq \mu\Big(|B_R(\rho, G)| \geq K\Big).
$$

Since $|B_R(\rho, G)|$ is finite almost surely, the right-hand side tends to zero as $K \to \infty$. The left-hand side does not denoted as K so we conclude $r(n) \to 0$ or claimed does not depend on *K*, so we conclude $r(n) \to 0$ as claimed.

The rest of the proof closely follows that of $[BNP11, Theorem 1.3]$ $[BNP11, Theorem 1.3]$, which uses a sprinkling argument (see also [\[AKS82,](#page-11-12) [ABS04\]](#page-11-5)). Given $p > p_c(G)$, let $p_1 > p_c(G)$ and $\varepsilon > 0$ such that

$$
1-p=(1-p_1)(1-\varepsilon).
$$

Let ω_1 denote an edge percolation configuration with edge probability p_1 , and let ω_{ε} denote an edge percolation configuration with edge probability ε . Then the union of ω_1 and ω_ε is equidistributed as a p-percolation. The first step is to show that a positive proportion of the vertices in ω_1 percolate far and hence are contained in sizable components.

Lemma 13. If
$$
p > p_c(G)
$$
, then there exists $\delta > 0$ such that for any $R > 0$,
\n
$$
\lim_{n \to \infty} \mathbb{P}_{n,p}\left(\frac{1}{|V_n|}\Big|\Big\{v \in G_n : v \leftrightarrow \partial B_R(v;G_n)\Big\}\Big| \leq \delta\right) = 0.
$$

Proof. The first part of the proof establishes a positive lower bound on the expectation of the quantity of interest and is essentially identical to the argument in the proof of [\[Sar21,](#page-12-0) Corollary 1.2]. Given (G, ρ) , let $f(G,\rho)$ be the probability under a p -percolation that the root percolates:
 $f(G,\rho)\equiv \mathbb{P}_p\Big(\Big|C(\rho;\omega(G))\Big|=\infty\Big)$,

$$
f(G,\rho) \equiv \mathbb{P}_p\bigg(\bigg|C(\rho;\omega(G))\bigg| = \infty\bigg),\,
$$

where $C(\rho;\omega(G))$ is the connected component of $\omega(G)$ that contains ρ . This is a measurable function on \mathscr{G}_\star , since it is the decreasing limit of the local (hence measurable) functions

$$
f_R(G,\rho) \equiv \mathbb{P}_p\bigg(\rho \leftrightarrow \partial B_R(\rho;G)\bigg).
$$

Since
$$
p > p_c(G)
$$
, the quantity $f(G, \rho)$ must be positive, μ -almost surely. Averaging over μ gives

$$
\mu \otimes \mathbb{P}_p\left(\rho \leftrightarrow \partial B(\rho, R)\right) \ge \mu \otimes \mathbb{P}_p\left(\left|C(\rho; \omega(G))\right| = \infty\right) \ge \delta > 0
$$

uniformly over all *R*. It follows by Corollary 3 that
\n
$$
\lim_{n \to \infty} \mathbf{L}_n \otimes \mathbb{P}_{n,p} \left(U_n \leftrightarrow \partial B_R(U_n; G_n) \right) = \mu \otimes \mathbb{P}_p \left(\rho \leftrightarrow \partial B(\rho, R) \right) \ge \delta.
$$

Let $Y_v = Y_v(R)$ be the indicator that v percolates at least distance R under the p -percolation,

$$
Y_v = \mathbf{1}\Big\{v \leftrightarrow \partial B_R(v;G_n)\Big\}\,.
$$

Let $X_n = X_n(R)$ be the sum of Y_v over all $v \in V_n$, so X_n is the quantity of interest. If $\mathbb{E}_{n,v}$ denotes expectation with respect to $\mathbb{P}_{n,p}$, then for all n large enough we have

$$
\frac{\mathbb{E}_{n,p} X_n}{|V_n|} = \mathbf{L}_n \otimes \mathbb{P}_{n,p} \left(U_n \leftrightarrow \partial B_R (U_n; G_n) \right) \geq \frac{\delta}{2}
$$

We next apply the second moment method to show that X_n concentrates, as the method used in [\[BNP11\]](#page-11-0)
no longer applies due to petential unboundedness of the degrees. Note that $Cov(Y, Y) < 1$ always (since no longer applies due to potential unboundedness of the degrees. Note that $Cov(Y_u, Y_v) \leq 1$ always (since Y_u , Y_v are indicators). If u and v lie at distance more than 2R in G_n , then Y_u and Y_v are clearly independent, so in this case $\text{Cov}(Y_u, Y_v) = 0$. Hence, we have that

$$
\frac{\text{Var}_{n,p} X_n}{|V_n|^2} \leq \frac{1}{|V_n|^2} \left| \left\{ (u,v) \in (V_n)^2 : d(u,v) \leq 2R \right\} \right| = o_n(1),
$$

by Lemma [12.](#page-8-2) It follows by Chebyshev's inequality that

$$
\mathbb{P}_{n,p_1}\bigg(X_n\leqslant \frac{\delta|V_n|}{4}\bigg)\to 0\,.
$$

The claim follows by redefining δ .

Once we prune the G_n , the remainder of the proof is essentially the same as in [\[BNP11\]](#page-11-0), where the uniform degree bound is only used to bound the number of edges in the graph. For the sake of being self-contained, we reproduce the following lemma from the end of the proof of [\[BNP11,](#page-11-0) Theorem 1.3]:

Lemma 14. Suppose G_n is an expanding sequence of finite graphs, each with average degree at most d. Let ω_1 , ω_ε be two independent edge percolation configurations on G_n , with edge probabilities p_1 and ε respectively. Let $X_n(R)$ be the number of vertices in G_n that percolate to distance R under ω_1 . Let C_n be the largest

connected component in
$$
\omega_1 \cup \omega
$$
 and $\delta \in (0, 1)$. Then there is a large enough constant R such that
\n
$$
\lim_{n \to \infty} \mathbb{P}_{n,p}\left(|C_n| \geq \frac{\delta |V_n|}{3} \middle| X_n(R) > \delta |V_n| \right) = 1.
$$

The value of R depends on d, δ , ε , and the expansion rate, but not on n.

Proof. We condition on ω_1 and use the randomness of ω_ε only. Consider the connected components of ω_1 of size at least R, of which there are at most $m = |V_n|/R$. By the bound on $X_n(R)$, these components cover at least a δ fraction of the vertices in G_n . Let A, B be any partition of these components into two sets, such that *

$$
\min\left\{|A|, |B|\right\} \geqslant \frac{\delta |V_n|}{3} \,. \tag{3.5}
$$

Let h be a positive constant such that

$$
\frac{1}{2}\liminf_{n\to\infty}h_{\delta/3}(G_n)\geq h.
$$

By the same reasoning as in the proof of Corollary [10,](#page-6-3) A and B must be joined by at least

$$
\lambda_n = \frac{h\delta|V_n|}{3}
$$

edge-disjoint paths for sufficiently large n , at least half of which must be of length at most

$$
\frac{|V_n|d/2}{\lambda_n/2} = \frac{3d}{h\delta} \equiv L.
$$

Thus, the probability that ω_{ε} fails to have a path from A to B is at most
 $(1 - \varepsilon^L)^{\lambda_n/2} \leqslant \exp\left\{-\frac{\lambda_n}{2} \varepsilon^L\right\}$

$$
(1 - \varepsilon^{L})^{\lambda_{n}/2} \leqslant \exp\left\{-\frac{\lambda_{n}}{2}\varepsilon^{L}\right\}
$$

Let *E* be the event that
$$
\omega_{\varepsilon}
$$
 has a path from *A* to *B* for every partition (A, B) satisfying (3.5). Then
\n
$$
1 - \mathbb{P}(E) \le 2^m \exp\left\{-\frac{\lambda_n}{2} \varepsilon^L\right\} \le \exp\left\{|V_n|\left(\frac{1}{R} - \frac{h\delta}{6} \varepsilon^L\right)\right\},
$$

which tends to zero by taking R a large enough constant. On the event E, the configuration $\omega = \omega_1 \cup \omega_\varepsilon$
contains a component that equipment a set a $\delta/3$ fraction of the vertices in C , as desired contains a component that occupies at least a $\delta/3$ fraction of the vertices in G_n , as desired.

Proof of Theorem [1](#page-1-0) part (iii). By Proposition [4,](#page-3-2) there exists a expanding sequence of subgraphs \bar{G}_n of G_n
such that each \bar{G}_n has average degree at most $\mathbb{E}(dge, \alpha)$, $\mathbb{E}(dge, \alpha)$, \bar{G}_n , α). Since such that each \bar{G}_n has average degree at most $\mathbb{E}(\deg \rho) + 1$ and $\bar{G}_n \to_{lw} (G, \rho)$. Since \bar{G}_n is a subgraph of G_n , the probability that $\omega(G_n)$ contains a component of linear size is at least the probability that $\omega(\bar{G}_n)$ contains a component of linear size, i.e. $p_c(\bar{G}_n) \geq p_c(G_n)$. Thus, it suffices to prove the statement for \bar{G}_n .

Let $X_n(R)$ be the number of vertices in \overline{G}_n that percolate to distance R under ω_1 and C_n be the largest connected component in $\omega(\bar{G}_n)$. We have that

$$
\mathbb{P}_{n,p}\left(|C_n| \geq \frac{\delta|V_n|}{3}\right) \geq \mathbb{P}_{n,p_1}(X_n(R) > \delta|V_n|) \times \mathbb{P}_{n,p}\left(|C_n| \geq \frac{\delta|V_n|}{3}\middle|X_n(R) > \delta|V_n|\right)
$$

for any R , $\delta > 0$. Taking δ to be the constant that results from applying Lemma [13](#page-9-0) to $\omega_1(\bar{G}_n)$ and R to be the constant that results from applying I amma 14 to \bar{G} and δ we have that

the constant that results from applying Lemma 14 to
$$
\bar{G}_n
$$
 and δ , we have that\n
$$
\lim_{n \to \infty} \mathbb{P}_{n,p_1}(X_n(R) > \delta |V_n|) = \lim_{n \to \infty} \mathbb{P}_{n,p} \left(|C_n| \geq \frac{\delta |V_n|}{3} \, \middle| \, X_n(R) > \delta |V_n| \right) = 1.
$$

It follows that

$$
\lim_{n\to\infty}\mathbb{P}_{n,p}\bigg(|C_n|\geqslant \frac{\delta|V_n|}{3}\bigg)=1,
$$

so we may take $\alpha = \delta/3$.

REFERENCES

[ABS04] N. Alon, I. Benjamini, and A. Stacey. Percolation on finite graphs and isoperimetric inequalities. Ann. Probab., 32(3A):1727–1745, 2004. [AKS82] M. Ajtai, J. Komlós, and E. Szemerédi. Largest random component of a *k*-cube. *Combinatorica*, 2(1):1–7, 1982.
[AL07] D. Aldous and R. Lyons. Processes on unimodular random networks. *Electron. J. Probab.*, 12:no. D. Aldous and R. Lyons. Processes on unimodular random networks. Electron. J. Probab., 12:no. 54, 1454–1508, 2007. [BCvdH`05] C. Borgs, J. T. Chayes, R. van der Hofstad, G. Slade, and J. Spencer. Random subgraphs of nite graphs. I. The scaling window under the triangle condition. Random Structures Algorithms, 27(2):137–184, 2005. [BNP11] I. Benjamini, A. Nachmias, and Y. Peres. Is the critical percolation probability local? Probab. Theory Related Fields, 149(1-2):261–269, 2011. [CMT22] D. Contreras, S. Martineau, and V. Tassion. Locality of percolation for graphs with polynomial growth. arXiv:2205.10253, 2022. [EH21] P. Easo and T. Hutchcroft. Supercritical percolation on finite transitive graphs I: Uniqueness of the giant component. arXiv:2112.12778, 2021. [Fou07] N. Fountoulakis. Percolation on sparse random graphs with given degree sequence. Internet Math., 4(4):329–356, 2007. [Hut20] T. Hutchcroft. Locality of the critical probability for transitive graphs of exponential growth. Ann. Probab., 48(3):1352–1371, 2020. [KNZ14] B. Karrer, M. E. J. Newman, and L. Zdeborová. Percolation on sparse networks. Phys. Rev. Lett., 113:208702, Nov 2014. [Lyo90] R. Lyons. Random walks and percolation on trees. Ann. Probab., 18(3):931–958, 1990. [MT17] S. Martineau and V. Tassion. Locality of percolation for Abelian Cayley graphs. Ann. Probab., 45(2):1247–1277, 2017. [NP10] A. Nachmias and Y. Peres. Critical percolation on random regular graphs. Random Structures Algorithms, 36(2):111–148, 2010.

- [Sar21] S. Sarkar. A note on the local weak limit of a sequence of expander graphs. Electron. Commun. Probab., 26:Paper No. 32, 6, 2021.
- [Sha13] Y. Shang. The sharp threshold for percolation on expander graphs. Math. Slovaca, 63(5):1141-1152, 2013.
- [SXZ14] H. Song, K.-N. Xiang, and S.-C.-H. Zhu. Locality of percolation critical probabilities: uniformly nonamenable case. arXiv:1410.2453, 2014.