

# Entropic information of a particle trapped by the Aharonov-Bohm-type effect

N. Ali,<sup>1,2,\*</sup> C. A. S. Almeida,<sup>3,†</sup> C. O. Edet,<sup>2,4,5,‡</sup> F. C. E. Lima,<sup>3,§</sup> and A. R. P. Moreira<sup>3,¶</sup>

<sup>1</sup>*Advanced Communication Engineering (ACE) Centre of Excellence,*

*Universiti Malaysia Perlis, 01000 Kangar, Perlis, Malaysia*

<sup>2</sup>*Faculty of Electronic Engineering Technology,*

*Universiti Malaysia Perlis, 02600 Arau, Perlis, Malaysia.*

<sup>3</sup>*Universidade Federal do Ceará (UFC),*

*Departamento de Física - Campus do Pici,*

*Fortaleza, CE, C. P. 6030, 60455-760, Brazil.*

<sup>4</sup>*Institute of Engineering Mathematics,*

*Universiti Malaysia Perlis, 02600 Arau, Perlis, Malaysia*

<sup>5</sup>*Department of Physics, Cross River University of Technology, Calabar, Nigeria.*

In this research article, we use the Shannon's formalism to investigate the quantum information entropy of a particle trapped by the Aharonov-Bohm-type effect. For quantum information study, it is necessary to investigate the eigenstates of the quantum system, i. e., the wave functions and energies of the quantum states. We assumed that the particle is in principle, confined in a cylindrical box in the presence of Aharonov-Bohm-type effect due to dislocation defect. Analysis of the quantum information entropy, reveals that the dislocation influences the eigenstates and, consequently, the quantum information of the system.

---

\*Electronic address: E-mail: norshamsuri@unimap.edu.my

†Electronic address: E-mail: carlos@fisica.ufc.br

‡Electronic address: E-mail: collinsokonedet@gmail.com

§Electronic address: E-mail: cleiton.estevao@fisica.ufc.br

¶Electronic address: E-mail: allan.moreira@fisica.ufc.br

## I. INTRODUCTION

Vector and scalar potentials appear in physics as auxiliary tools describing the electric and magnetic fields [1, 2]. Furthermore, potentials play a fundamental role in the canonical description of a physical system at the classical [3] and quantum levels [4, 5]. In quantum theory, potentials are responsible for the emergence of the Aharonov-Bohm (AB) effect [1].

The AB effect is a quantum phenomenon in which an electrically charged particle is affected by an electromagnetic potential [4, 5]. In principle, the AB effect tells us that the particle will feel the influence of the field even when the particle is at a null-field region [1, 4, 5]. This effect was described in 1959 by Yakir Aharonov and David Bohm in their seminal paper *Significance of Electromagnetic Potentials in Quantum Theory* [1].

After the publication of the paper of Aharonov and Bohm, several applications discussing AB effects have appeared in the literature [6–8]. For example, the AB effect arises when studying carbon nanotubes [9], in the scattering process [10], and in photonic systems [11]. The fact is that there is a growing interest in the study of the AB effect [12–14]. In particular, this interest motivates us to understand the quantum information of a particle trapped by an AB-type effect due to the presence of a topological defect, i. e., the dislocation defect.

The idea of quantum information arises in search of a general communication theory [15]. Claude E. Shannon sought an extension of communication theory to include new factors. In particular, Shannon sought to implement in his study the noise effect on the channel and the possible savings in the propagation process of the message [15]. This relevant study made Shannon one of the forerunners of communication theory.

Shannon's theory has been used in quantum mechanics to study information from physical systems [16–18]. In quantum theory, Shannon's theory is a theoretical measuring of the communication responsible for informing us of the measure of uncertainty in the position and momentum of the particle [17, 18]. In other words, Shannon's information is information theoretical-measuring associated with the position space and reciprocal. Indeed, this informational analysis plays a relevant role in the foundations of information theory [15] and quantum mechanics problems [4, 5, 19] (see Refs. [20–22]). Furthermore, Shannon's entropy is a useful approach used in the study of cryptography [23], noise theory [24], harmonic oscillators [25–27], potential wells [28–36], and theories of effective mass [37–40].

Some studies are found in the literature discussing the influence of the AB effect on the

system eigenstates [41]. Furthermore, some studies were also performed on the information of quantum-mechanical systems with AB effect with a specific interaction [16]. However, as far as we know, no studies have been carried out purely studying the influence of the AB-type effect on the quantum information of a spinless particle. Therefore, the main purpose is to study how quantum information changes as the AB-type effect is modified.

We organize the paper as follows: In Sec. II, we build the quantum theory and present the analytical solution of the eigenstates of the system. Posteriorly, in Sec. III, the quantum information of the system is studied considering an alteration of the AB-type effect. Finally, in Sec. IV, we announce our findings.

## II. QUANTUM DESCRIPTION OF THE SYSTEM

Our purpose in this section is the quantum description of a spinless particle subjected to the AB effect in the presence of dislocation. So, the eigenstates depend on the AB effect and the dislocation defect. In summary, the dislocation defect corresponds to the distortion of a vertical line into a spiral [42, 43]. Seeking to understand how the AB effect and dislocation modify the eigenstates, let us consider a quantum system composed of a spinless particle in the presence of a dislocation [44], i. e.,

$$ds^2 = dr^2 + 2\beta dr d\theta + r^2 d\theta^2 + dz^2, \quad (1)$$

where the parameter  $\beta$  describes the dislocation and obeys the constraint  $0 < \beta < 1$ . Furthermore, the cylindrical symmetry considered (1) requires that  $0 < r < \infty$ ,  $0 \leq \theta \leq 2\pi$  and  $-\infty \leq z \leq \infty$ .

Considering a nonrelativistic particle in the presence of dislocation [44], we write the Schrödinger equation as

$$-\frac{1}{2m} \frac{1}{\sqrt{g}} \partial_i (g^{ij} \sqrt{g} \partial_j) \psi(r, \theta, z) = \mathcal{E} \psi(r, \theta, z). \quad (2)$$

Here, we use the  $\hbar = c = 1$ .

Now using the line element (1) and considering the Eq. (2), one obtains

$$-\frac{1}{2m} \left\{ \frac{\partial^2}{\partial r^2} + \frac{\partial}{\partial z^2} + \frac{1}{(r^2 - \beta^2)} \left[ r \frac{\partial}{\partial r} + \left( \frac{\partial}{\partial \theta} - \beta \frac{\partial}{\partial z} \right)^2 \right] \right\} \psi(r, \theta, z) = \mathcal{E} \psi(r, \theta, z). \quad (3)$$

To solve Schrödinger's equation (3), allow us assume

$$\psi(r, \theta, z) = e^{i(l\theta + kz)}R(r). \quad (4)$$

Here,  $l = 0, \pm 1, \pm 2, \pm 3, \dots$  and  $k \in \mathbb{R}$ . Besides, note that the parameters  $l$  and  $k$  are the eigenvalues associated with the operators  $\hat{p}_z = -i\partial_z$  and  $\hat{L}_z = -i\partial_\theta$ , respectively.

Assuming the transformation (4) where  $\theta$  and  $z$  are cyclic coordinates, we write Schrödinger's equation as

$$R''(r) + \frac{1}{(r^2 - \beta^2)} \left[ rR'(r) - (l - \beta k)^2 R(r) \right] + (2m\mathcal{E} - k^2)R(r) = 0. \quad (5)$$

Here, prime notation represents the derivative concerning the radial coordinate.

Seeking to describe the confinement of the spinless particle, let us assume the coordinate change

$$r = \frac{x}{\sqrt{2m\mathcal{E} - k^2}}. \quad (6)$$

This coordinate change appears in Refs. [44, 45] to study a particle confinement by a hard wall.

Considering the Eq. (6), we rewrite the Eq. (5) as

$$\left(1 - \frac{4\varsigma^2}{x^2}\right)\ddot{R}(x) + \frac{1}{x}\dot{R}(x) + \left(1 - \frac{\lambda}{x^2} - \frac{2\varsigma^2}{x^2}\right)R(x) = 0, \quad (7)$$

where

$$\lambda = (l - \beta k)^2 + \varsigma^2, \quad (8)$$

and

$$\varsigma^2 = \frac{\beta^2(2m\mathcal{E} - k^2)}{4}. \quad (9)$$

Here, the point notation represents the derivative concerning the variable  $x$ .

Let us highlight that the parameter that describes the dislocation (or distortion of the metric) has a value restrained in the range  $[0, 1]$  [42, 43]. So, one can assume the parameter  $\beta$  is small. Therefore, the higher order terms of  $\beta$  will be negligible. In this way, we reformulate the Eq. (7) as follows:

$$\ddot{R}(x) + \frac{1}{x}\dot{R}(x) + \left[1 - \left(\frac{l - \beta k}{x}\right)^2\right]R(x) = 0. \quad (10)$$

Here, allow us to remember that every equation written in the form

$$f''(x) + \frac{1}{x}f'(x) + \left(k^2 - \frac{\alpha^2}{x^2}\right)f(x) = 0 \quad (11)$$

is known as Bessel's equation [2, 46, 47]. Thus, the general solution of the Bessel equation will be

$$f(x) = c_1 J_\alpha(x) + c_2 J_{-\alpha}(x), \quad (12)$$

where

$$J_\alpha(x) = \sum_{k=0}^{\infty} (-1)^k \frac{1}{k! \Gamma(\alpha + k + 1)} \left(\frac{x}{2}\right)^{\alpha+2k}. \quad (13)$$

For more details on the Bessel equation, see Refs. [2, 46, 47].

Comparing Eq. (11) and its solution (12) with Eq. (10) and requiring the wave function to be normalizable, one obtains that the solution of Eq. (10) is

$$R(x) = A_0 J_{|l-\beta k|}(x). \quad (14)$$

Here  $A_0$  is the normalization constant, and  $J_{|l-\beta k|}(x)$  is Bessel's function of first kind [2, 46, 47].

Building the general wave function that describes spinless particles in the presence of disclination, one obtains

$$\psi(r, \theta, z) = A_0 e^{i(l\theta + kz)} J_{|l-\beta k|}\left(\frac{2\zeta}{\beta^2} r\right). \quad (15)$$

As discussed by Silva and Bakke [44], let us analyze the confinement of the spinless particle by a hard-wall confining potential in the elastic medium. For this, allow us to assume that

$$\lim_{x \rightarrow x_0} R(x) = 0 \quad \text{with} \quad x_0 = \beta \sqrt{2m\mathcal{E} - k^2}. \quad (16)$$

Applying the constraint (16), one obtains that

$$\mathcal{E}_{n,l,k} = \frac{1}{2m} \left(\frac{\Theta_{n,|l-\beta k|}}{r_0}\right)^2 + \frac{k^2}{2m}. \quad (17)$$

Here  $\Theta_{n,|l-\beta k|}$  is the  $n$ -th zero of the  $|l-\beta k|$ -th Bessel's function [4]. Furthermore,  $n = 0, 1, 2, \dots$ , and the quantum number associated to the angular momentum is  $l = 0, \pm 1, \pm 2, \dots$ .

The results of the Eqs. (15) and (17) expose how eigenstates change with disclination. So, we expect the existence of a dependency on the quantum information with the disclination defect. In the next section, we present a study of the variation of Shannon information as the disclination changes.

Note that the parameters  $n$  and  $|l - \beta k|$  play the role of quantum numbers associated with the radial and angular variables, respectively. Therefore, this result allows us to perceive the influence of the defect on the energy eigenvalues. Moreover, one notes this influence by a change in the quantum number of the angular momentum. It is relevant to mention that this change arises even if there is no interaction between the particle and the defect. Indeed, this effect is called the Aharonov-Bohm-type effect and appears due to the presence of the topological defect [44, 48].

### III. INFORMATION THEORETICAL MEASURE: SHANNON'S ENTROPY

The growing interest in studies of information-theoretical measurements of quantum-mechanical systems is evident [23–40]. Initially, the entropic measure of information comes up with an alternative proposal to the well-known Heisenberg uncertainty relation [39, 49]. Among different approaches used to measure information entropies is the so-called Shannon's entropy [39]. In quantum theories, Shannon's entropy plays an essential role in uncertainty measurement. In particular, this entropy gives us a new version of Heisenberg's uncertainty relationship between the position and momentum spaces [49]. This relation is

$$S_r + S_p \geq D(1 + \ln \pi). \quad (18)$$

Here,  $D$  is the spatial dimension (in the one-dimensional case  $D=1$ ),  $S_r$  is the information entropy in the position space, and  $S_p$  is the information entropy in the momentum space.

Using the definition of Shannon's entropy, let us write the quantum information entropy as

$$S_r = - \int_{-\infty}^{\infty} |\psi(x)|^2 \ln |\psi(x)|^2 dx, \quad (19)$$

and

$$S_p = - \int_{-\infty}^{\infty} |\psi(p)|^2 \ln |\psi(p)|^2 dp, \quad (20)$$

where

$$\psi(p) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \psi(x) e^{ipx} dx. \quad (21)$$

Shannon's entropy helps us to calculate the entropic information measures. In this context, Shannon's entropy has performed a relevant role in several studies. For example, one can apply this approach in investigating Aharonov-Bohm rings [16], quantum systems with double-well potentials [21], hyperbolic interaction [29], position-dependent mass theories [38, 40], and the Aharonov-Bohm effect [50].

Motivated by these applications, we will use this approach to study quantum information theoretical measurements of a spinless particle. That particle is in the presence of a screw displacement, which produces the Aharonov-Bohm-type effect. So, to achieve our purpose, let us consider the wave eigenfunctions (15) and substitute in Eq. (19). This substitution leads us to Shannon's entropy in the position space. Subsequently, we perform Fourier's transform (21) of the eigenfunctions (15) and calculate Shannon's entropy in the reciprocal space using Eq. (20). The numerical results shown in Tab. III are obtained numerically solving the integrals (19) and (20) for the first eigenstates. In Fig. 1, the probability densities at the position space for the ground state and the first two excited states used in Shannon's entropy calculation are displayed. On the other hand, in Fig. 2, probability densities at the reciprocal space used in Shannon's entropy calculation are exposed.

Analyzing Fig. 1, i. e., the probability density for the first three energy levels, one notes that the more energetic states have greater amplitudes and oscillations. This behavior restricts the regions most likely that the particle can be. In addition, the results presented in Fig. 1 show that by increasing the influence of the topological defect responsible for the Aharonov-Bohm type effect, the amplitudes of the probability density increase. That suggests that the quantum information of the system increases as the Aharonov-Bohm-type effect increases (i.e., as the parameter  $\beta$  increases).

Looking at Fig. 2, one can see that the probability density of the particle at the reciprocal space is a Dirac delta-like distribution. This probability density profile suggests minimal uncertainty in the momentum of the spinless particle. Furthermore, with the probability densities shown in Figs. 1 and 2, we obtain the numerical results of Tab. III. This result tells us that the quantum information decreases as the disclination increases. This decreasing in information suggests an increase in uncertainty measurements associated with




Figure 1: Probability density at the position space. (a) The case:  $n = 0$  and  $l = 0$ . (b) The case:  $n = 1$  and  $l = -1$ . (c) The case:  $n = 2$  and  $l = -2$ .

particle position measurements. Besides, the quantum information at the reciprocal space increases when the disclination is larger. Thus, suggesting less uncertainty in the momentum measurement of the particle. Furthermore, one notes that these results remain valid for quantum states with higher energy. It is worth mentioning that in all cases, we have

$$S_r + S_p \geq 6.43419, \quad (22)$$

and therefore, the Bialynicki-Birula and Mycielski [49] relationship is preserved.

#### IV. FINAL REMARKS

In this research, we have employed Shannon's entropy to study the quantum information entropy of a non-relativistic particle trapped by the Aharonov-Bohm-type effect. The Schrodinger equation in cylindrical coordinate is solved in the presence of topological defect and the resulting Aharonov-Bohm effect to obtain the wave functions and eigenvalues of quantum system. These eigensolutions are employed to evaluate the Shannon entropy






Figure 2: Probability density at the reciprocal space. (a) The case:  $n = 0$  and  $l = 0$ . (b) The case:  $n = 1$  and  $l = -1$ . (c) The case:  $n = 2$  and  $l = -2$ .

numerically. From our analysis of the quantum information, it is possible to see that the dislocation influences the eigenstates and, consequently, the quantum information of the system. It is interesting to point out that the famous Bialynicki-Birula and Mycielski (BBM) relationship is satisfied as well. Going forward, the results of this study will find applications in quantum information processing and related areas.

### Authors Declaration

### Funding

The research has been carried out under Long Term Research Grant Scheme project LRGS/1/2020/UM/01/5/2 provided by Ministry of Higher Education of Malaysia (MOHE).

$n$	$l$	$\beta$	$S_r$	$S_p$	$S_r + S_p$	$3(1 + \ln \pi)$
0	0	0.2	9.74631	0.06678	9.81309	6.43419
		0.4	9.74262	0.07558	9.81821	
		0.8	9.74040	0.12158	9.86199	
	-1	0.2	9.74435	0.05526	9.79961	6.43419
		0.4	9.74424	0.10408	9.84832	
		0.8	9.74387	0.29596	10.03984	
1	0	0.2	9.74483	0.00770	9.75254	6.43419
		0.4	9.74439	0.10381	9.84821	
		0.8	9.74353	0.29641	10.03991	
1	0.4	0.2	9.74410	0.01640	9.76050	6.43419
		0.4	9.74377	0.13777	9.88154	
		0.8	9.74312	0.29938	10.04251	
	-2	0.2	9.74461	0.02251	9.76713	6.43419
		0.4	9.74329	0.22803	9.97133	
		0.8	9.74277	0.35949	10.10231	
	-1	0.2	9.74518	0.02720	9.77238	6.43419
		0.4	9.74361	0.25583	9.99944	
		0.8	9.74301	0.48609	10.22910	
2	0	0.2	9.74482	0.04111	9.78593	6.43419
		0.4	9.74370	0.25717	10.00091	
		0.8	9.74317	0.52424	10.26742	
1	0.4	0.2	9.74435	0.04188	9.78623	6.43419
		0.4	9.74335	0.35605	10.09942	
		0.8	9.74291	0.85920	10.60214	
2	0.4	0.2	9.74399	0.07462	9.81861	6.43419
		0.4	9.74311	0.44406	10.18722	
		0.8	9.74272	0.91082	10.65351	

Table I: Numerical result of Shannon's entropy.

### Conflicts of interest/Competing interest

All the authors declared that there is no conflict of interest in this manuscript

### Acknowledgements

This research has been carried out under LRGS Grant LRGS/1/2020/UM/01/5/2 (9012-00009) Fault-tolerant Photonic Quantum States for Quantum Key Distribution provided by Ministry of Higher Education of Malaysia (MOHE). C. O. Edet acknowledges eJDS (ICTP).

C. A. S. Almeida thanks to Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), nº 309553/2021-0. F. C. E. Lima is grateful to Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), nº 88887.372425/2019-00.

### Data Availability

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request

- 
- [1] Y. Aharonov and D. Bohm, *Phys. Rev.* **115** (1959) 485.
  - [2] G. B. Arfken and H. J. Weber, *Mathematical methods for physicists*, (6th ed., Elsevier Academic Press, New York, 2005).
  - [3] J. B. Marion, *Classical dynamics of particles and systems*, (5th Revised ed., Cengage Learning, USA, 2003).
  - [4] D. J. Griffiths, *Introduction to quantum mechanics*, (Prentice Hall, USA, New Jersey, 1994).
  - [5] J. J. Sakurai and J. Napolitano, *Modern quantum mechanics*, (Addison-Wesley Publishing Company, USA, 1994).
  - [6] C. Overstreet, P. Asenbaum, J. Curti, M. Kim and M. A. Kasevich, *Science* **375** (2022) 226.
  - [7] Y. Ronen, T. Werkmeister, D. H. Najafabadi, A. T. Pierce, L. E. Anderson, Y. J. Shin, S. Y. Lee, Y. H. Lee, B. Johnson, K. Watanabe, T. Taniguchi, A. Yacoby and P. Kim, *Nature* **16** (2021) 563.
  - [8] E. Mefford and K. Suzuki, *JHEP* **05** (2021) 026.
  - [9] H. Ajiki and T. Ando, *Physica B* **201** (1994) 349.
  - [10] S. N. M. Ruijsenaars, *Ann. Phys.* **146** (1983) 1-34.
  - [11] K. Fang, Z. Yu and S. Fan, *Phys. Rev. Lett.* **108** (2012) 153901.
  - [12] L. Vaidman, *Phys. Rev. A* **86** (2012) 040101(R).
  - [13] A. Caprez, B. Barwick and H. Batelaan, *Phys. Rev. Lett.* **99** (2007) 210401.
  - [14] M. Chaichian, P. Prešnajdera, M. M. Sheikh-Jabbaric and A. Tureanu, *Phys. Lett. B* **527** (2002) 149.
  - [15] C. E. Shannon, *Bell Syst. Tech. J.* **27** (1948) 623.

- [16] C. O. Edet, F. C. E. Lima, C. A. S. Almeida, N. Ali and M. Asjad, *Entropy*, **24** (2022) 1059.
- [17] F. C. E. Lima, A. R. P. Moreira and C. A. S. Almeida, *Int. J. Quant. Chem.* **121** (2021) e26645.
- [18] F. C. E. Lima, A. R. P. Moreira, L. E. S. Machado and C. A. S. Ameida, *Int. J. Quant. Chem.* **121** (2021) e26749.
- [19] R. K. Pathria, *Statistical Mechanics*, (2nd ed., Butterworth Heinemann, Jordan Hill, Oxford, 1996).
- [20] C. A. Gil-Barrera, R. S. Carrillo, G.-H. Sun and S. -H. Dong, *Entropy* **24** (2022) 604.
- [21] G. -H. Sun, S. -H. Dong, K. D. Launey, T. Dytrych and J. P. Draayer, *Int. J. Quant. Chem.* **115** (2015) 891.
- [22] G. -H. Sun, S. -H. Dong and N. Saad, *Int. J. Quant. Chem.* **525** (2013) 934.
- [23] F. Grosshans, N. J. Cerf, *Phys. Rev. Lett.* **92** (2004) 047905.
- [24] A. D. Wyner, S. Shamai, *Proc. IEEE* **86** (1998) 442.
- [25] A. Boumali, M. Labidi, *Mod. Phys. Lett. A* **33** (2018) 1850033.
- [26] R. J. Yañez, W. V. Assche and J. S. Dehesa, *Phys. Rev. A* **50** (1994) 3065.
- [27] J. S. Dehesa, E. D. Belega, I. V. Toranzo and A. I. Aptekarev, *Int. J. Quantum Chem.* **119** (2019) e25977.
- [28] P. A. Bouvrie, J. C. Angulo, J. S. Dehesa, *Phys. A* **390** (2011) 2215.
- [29] R. Valencia-Torres, G. H. Sun and S. H. Dong, *Phys. Scr.* **90** (2015) 035205.
- [30] J. S. Dehesa, A. Martínez-Finkelshtein, V. N. Sorokin, *Mol. Phys.* **104** (2006) 613.
- [31] G. -H. Sun, M. A. Aoki, S. -H. Dong, *Chin. Phys. B* **22** (2013) 050302.
- [32] V. Majernik, R. Charvot, E. Majernikova, *J. Phys. A: Math. Gen.* **32** (1999) 2207.
- [33] S. T. Tserkis, C. C. Moustakidis, S. E. Massen, C. P. Panos, *Phys. Lett. A* **378** (2014) 497.
- [34] N. Mukherjee, A. K. Roy, *Int. J. Quantum Chem.* **11**, 2018, e25596.
- [35] P. R. Kumar, G. Kumar, R. Kumar, A. Kumar, *Int. J. Quantum Chem.* **116** (2016) 1413.
- [36] P. O. Amadi, A. N. Ikot, A. T. Ngiangia, U. S. Okorie, G. J. Rampho, H. Y. Abdullah, *Int. J. Quantum Chem.* **120** (2020) e26246.
- [37] F. A. Serrano, B. J. Falaye, S. H. Dong, *Phys. A Stat. Mech. Appl.* **446** (2016) 152.
- [38] S. Guo-Hua, D. Popov, O. Camacho-Nieto, S. H. Dong, *Chin. Phys. B* **24** (2015) 100303.
- [39] G. Yañez-Navarro, G. H. Sun, T. Dytrych, K. D. Launey, S. H. Dong, J. P. Draayer, *Ann. Phys.* **348** (2014) 153.

- [40] F. C. E. Lima, Ann. Phys. **442** (2022) 168906.
- [41] A. N. Ikot, C. O. Edet, P. O. Amadi, U. S. Okorie, G. J. Rampho and H. Y. Abdullah, Eur. Phys. J. D **74** (2020) 159.
- [42] K. S. Valanis and V. P. Panoskaltsis, Acta Mech. **175** (2005), 77-103.
- [43] R. A. Puntigam and H. H. Soleng, Class. Quant. Grav. **14** (1997), 1129-1149.
- [44] W. C. F. da Silva and K. Bakke, Eur. Phys. J. C **79** (2019) no.7, 559.
- [45] H.J.W. Müller-Kirsten, Introduction to quantum mechanics: Schrödinger equation and path integral (Word Scientific, Singapore, 2006).
- [46] M. Abramowitz, I.A. Stegun, Handbook of mathematical functions (Dover Publications Inc., New York, 1965).
- [47] E. Butkov, *Mathematical Physics*, (1st ed., Addison-Wesley, USA, 1973).
- [48] A. L. Silva Netto, C. Chesman and C. Furtado, Phys. Lett. A **372** (2008) 3894.
- [49] I. Białynicki-Birula and J. Mycielski, Comm. Math. Phys. **44**, (1975) 129.
- [50] C. O Edet and A. N. Ikot, Eur. Phys. J. C **136** (2021) 432.
- [51] V. B. Bezerra, J. Math. Phys. **38** (1997), 2553-2564.
- [52] A. I. Aptekarev, J. S. Dehesa, R. J. Yáñez, J. Math. Phys. **35** (1994) 4423.
- [53] W. Van Assche, R. J. Yáñez, J. S. Dehesa, J. Math. Phys. **36** (1995) 4106.
- [54] I.I. Hirschmann Jr., Amer. J. Math. **79** (1957) 152.