

Sound propagation in cigar-shaped Bose liquids in the Thomas-Fermi approximation: A comparative study between Gross-Pitaevskii and logarithmic models

Konstantin G. Zloshchastiev*

*Institute of Systems Science, Durban University of Technology, P.O. Box 1334,
Durban 4000, South Africa; kostiantynz@dut.ac.za, kostya@u.nus.edu*

(Dated: received: 30 September 2022)

Abstract: A comparative study is done of the propagation of sound pulses in elongated Bose liquids and Bose-Einstein condensates in Gross-Pitaevskii and logarithmic models, by means of the Thomas-Fermi approximation. It is shown that in the linear regime the propagation of small density fluctuations is essentially one-dimensional in both models, in the direction perpendicular to the cross section of a liquid's lump. Under these approximations, it is shown that the speed of sound scales as a square root of particle density in the case of the Gross-Pitaevskii liquid/condensate, but it is constant in a case of the homogeneous logarithmic liquid.

Keywords: quantum fluid; Bose-Einstein condensate; logarithmic Schrödinger equation; cigar-shaped Bose-Einstein condensate; Thomas-Fermi approximation; Gross-Pitaevskii equation

PACS numbers: 03.75.Kk, 47.37.+q

1. INTRODUCTION

The remarkable experiments with cooling magnetically trapped alkali atoms down to nanokelvin temperatures demonstrated a steep narrowing of the velocity and density distribution profiles [1–3]. This was explained by the occurrence of the Bose-Einstein condensation (BEC) phenomenon, earlier known to exist in noble gases, such as helium-4 [4], and excitonic semiconductors at low temperatures [5].

From the modern point of view, Bose-Einstein condensates fall under a wide category of quantum Bose liquid. This special kind of matter includes hydrodynamic media formed not only by integer-spin particles but also by the “bosonized” combinations of fermions. The standard theory of quantum Bose liquids is far from its completion, due to the obvious complexity of this matter and large number of phenomena involved, as we shall discuss below.

Historically, it was the Gross-Pitaevskii (GP) model, which was first proposed for describing Bose-Einstein condensation [6, 7]. This is essentially a hydrodynamic version of a quantum many-body model whose condensate particles' interaction potential was truncated at the two-body contact term, within the perturbation approach's paradigm [8, 9]. In this model, the cubic nonlinear Schrödinger equation naturally occurs as an equation for the condensate's wavefunction. This approach thus assumes that the pair interaction is largest in magnitude when it comes to the Bose-Einstein condensation. This assumption does seem acceptable for the diluted condensates, such as vapours of alkali atoms, where interparticle distances are sufficiently large to neglect the multi-body (three or more) collisions. However, as the density grows,

multi-body interactions should come into play, even for laboratory quantum liquids [10–12], let alone quantum fluids in high-energy physics and astrophysics. On top of that, effects of lower-dimensionality and nontrivial vacuum's presence can come into play and somewhat narrow an applicability range of the Gross-Pitaevskii model.

This inspired further corrections to the Gross-Pitaevskii model, *e.g.*, in the form of adding higher-order polynomial terms to the (formerly) cubic Schrödinger equation, to mention just some literature examples [13–15]. Therefore, the question remains whether these polynomial terms are exact, or they actually represent an infinite series expansion of some non-polynomial function of condensate density.

The conceptually different model of the Bose-Einstein condensate, defined by a wave equation with the non-polynomial (logarithmic) nonlinearity, was proposed in [16]. Subsequently, it was shown that the logarithmic nonlinearity should universally occur in the systems with the following properties: (i) they allow a hydrodynamic description in terms of the fluid wavefunction (cf. [17]), and (ii) their particles' characteristic interaction potentials are substantially larger than kinetic energies [18]. A profound relation between logarithmic Schrödinger equations and quantum information entropy should be mentioned as well [19, 20]. Fruitful applications of the logarithmic models were found in various physical systems [21–32]. Extensive mathematical studies of logarithmically nonlinear systems were performed as well, to mention only very recent results [33–41].

In this paper, we perform a comparative study of the propagation of sound pulses in cigar-shaped (effectively one-dimensional) Bose-Einstein condensates in Gross-Pitaevskii and logarithmic models. We shall use the Thomas-Fermi approximation as well as consider a linear regime at some stage. This is usually believed to deliver a clear analytical picture of the underlying physics, while avoiding technical complications. In the next section we

*Electronic address: <https://orcid.org/0000-0002-9960-2874>

give a brief description of both models. In section 3, we consider their thermodynamical properties, and in section 4 we study the propagation of small fluctuations of density. Some discussion and conclusions are presented in section 5.

2. THE MODELS

Both Gross-Pitaevskii and logarithmic Bose-Einstein condensates are described by a condensate wavefunction $\Psi = \Psi(\mathbf{x}, t)$, which is normalized to a number N of condensate particles of mass m :

$$\int_{\mathcal{V}} |\Psi|^2 d\mathcal{V} = \int_{\mathcal{V}} n d\mathcal{V} = N, \quad (1)$$

where $n = \rho/m = |\Psi|^2$ is particle density and \mathcal{V} is the volume of the condensate. This wavefunction obeys a minimal $U(1)$ -symmetric nonlinear Schrödinger equation

$$i\hbar\partial_t\Psi = \left[-\frac{1}{2}\hbar\mathcal{D}\nabla^2 + V(\mathbf{x}) - F(|\Psi|^2) \right] \Psi, \quad (2)$$

where $\mathcal{D} = \hbar/m$, $V(\mathbf{x})$ is an external field potential which usually describes trapping fields and confining vessel's boundaries in BEC experimental setups. Furthermore, $F(n)$ is a preselected function of particle density n , which defines the condensate's type; in our case:

$$F(n) = \begin{cases} -U_0 n, & \text{(Gross-Pitaevskii BEC)} \\ b \ln(n/n_c), & \text{(Logarithmic BEC)} \end{cases} \quad (3)$$

where U_0 is the strength of the two-body interaction, and b is the logarithmic nonlinear coupling, and n_c is a critical density value (at which the logarithmic term is nil). These parameters have different physical meaning and origins. The parameter U_0 is a fixed parameter of the GP model, which can be expressed in terms of atom-atom collisions:

$$U_0 = 4\pi\hbar a\mathcal{D}, \quad (4)$$

where a is the scattering length.

The parameter b of the logarithmic model is linearly related to the wave-mechanical temperature T_Ψ , which is defined as a thermodynamical conjugate of the Everett-Hirschman entropy function $S_\Psi = -\int_{\mathcal{V}} |\Psi|^2 \ln(|\Psi|^2/n_c) d\mathcal{V}$. In case of Bose-Einstein condensates, this is quantum information entropy which emerges from the logarithmic term when one averages the logarithmic nonlinear wave equation in a Hilbert space of wavefunctions Ψ [19, 20]. It becomes the thermodynamical entropy of the condensate $S = -\int_{\mathcal{V}} n \ln(n/n_c) d\mathcal{V}$. Correspondingly, it is conjectured that T_Ψ is linearly related to the conventional (thermal) temperature T [18], thus we can assume here that

$$b \sim T_\Psi \sim T, \quad (5)$$

or

$$b = \chi(T - T_c), \quad (6)$$

where T_c is a critical temperature at which the logarithmic term vanishes, and χ would be a scale constant dimensionless in units where the Boltzmann constant is one. This formula indicates that b is not a fixed parameter of the model, but its value depends on the environment of the logarithmic condensate.

One can see that the attractive/repulsive behaviour for the Gross-Pitaevskii condensate is fixed by a sign of the strength U_0 , while the logarithmic condensate can switch between attractive and repulsive regimes as density goes across the value n_c . This leads to many profound effects which are only pertinent to the logarithmic Bose liquids, including the enhanced stability [20, 29, 42].

3. THERMODYNAMICS AND FLUID ANALOGY

Let us first derive an expression for the chemical potential μ , which is defined as the Lagrange multiplier that ensures constancy of number N under arbitrary variations of wavefunction [8]. Following the standard procedure, we use the stationary ansatz $\Psi(\mathbf{x}, t) = \psi(\mathbf{x}) \exp(-i\mu t/\hbar)$, to obtain from (2) the general expression for the chemical potential:

$$\mu = -\frac{1}{2}\hbar\mathcal{D} \frac{\nabla^2\psi}{\psi} + V(\mathbf{x}) - F(|\psi|^2), \quad (7)$$

where $F(n)$ would be given by equations (3) for the models in question. For example, for a free condensate, $V(\mathbf{x}) \equiv 0$, we obtain

$$\mu \approx \begin{cases} U_0|\psi|^2 = U_0 n, \\ -b \ln(|\psi|^2/n_c) = -b \ln(n/n_c), \end{cases} \quad (8)$$

where the approximation indicates that we neglected the kinetic energy term, trapping fields and boundary effects. Here and in equations below, we shall omit the model titles, assuming that top and bottom lines after a curly bracket refer to the Gross-Pitaevskii and logarithmic condensates, respectively.

One can see that, similarly to the attractive/repulsive behaviour discussed earlier, the chemical potential's sign in the case the Gross-Pitaevskii condensate is fixed by a sign of the strength U_0 , while the logarithmic condensate's μ switches its sign as density goes across the value n_c . Thus, adding or removing particles to the logarithmic condensate is energetically favorable in two of the four regions $\{n \geq n_c, T \geq T_c\}$, and unfavorable in others. This, of course, has a crucial influence upon a stability of the free logarithmic condensate, as mentioned above. In presence of an external potential, the picture gets slightly more complicated but the analysis can be easily done by analogy.

Furthermore, let us deduce an equation of state from equation (2). To do that, we use the analogy between Schrödinger-type equations and inviscid isentropic flows. This analogy has been known for a long time - since works by de Broglie and Madelung [43, 44], and subsequently it was rediscovered few times [45–48], to mention only examples known to the author. For the logarithmic Schrödinger equation specifically, the fluid-Schrödinger analogy was derived in [22], although with some error in the equation of state; the correct equation of state and derivation can be found in [20, 23]. For the equation (2) with a general function $F(n)$, the fluid-Schrödinger analogy was derived in [49].

According to the analogy, the wave equation (2) can be rewritten in the hydrodynamic form by using the Madelung ansatz [43, 44], from which one can deduce the equation of state and related values. In the leading order approximation with respect to Planck constant, we can neglect second derivatives of density (which is robust as long as we do not consider shock waves and other fluctuations with a non-smooth density profile), the hydrodynamic equations have the perfect-fluid form, from which we obtain the equation of state $p = p(n)$ and speed of sound c , respectively, as:

$$p \approx - \int nF'(n) dn, \quad c \approx \sqrt{-nF'(n)/m}, \quad (9)$$

where prime denotes an ordinary derivative. Therefore, using the definitions (3), we obtain for pressure:

$$p \approx \begin{cases} n^2 U_0/2, \\ -bn, \end{cases} \quad (10)$$

and velocity of sound:

$$c \approx \begin{cases} \sqrt{nU_0/m}, \\ \sqrt{-b/m}, \end{cases} \quad (11)$$

where we omitted the approximation sign from now on. One can see for that the squared velocity of sound is proportional to density in the Gross-Pitaevskii case, whereas for the logarithmic fluid it is independent of density.

A direct comparison of the formulae above indicates that the logarithmic condensate is actually closer to the ideal gas (whose pressure is a linear function of density) than the Gross-Pitaevskii condensate is. Therefore, if one does a Taylor series expansion of the general barotropic fluid's equation of state with respect to density, then the first-order term (apart from a constant) would represent the logarithmic liquid, whereas the Gross-Pitaevskii term would be only a second-order (quadratic) correction.

4. SOUND IN CIGAR-SHAPED CONDENSATES

Let us consider a cloud or lump of BEC whose transverse dimensions R being significantly less than its axial

dimension H aligned along z axis. In this setup, we assume that sound pulses have characteristic length scale ℓ in the axial direction, $R \ll \ell \ll H$, which is usually what takes place in experiments like [1, 3]. Along the lines of those experimental setups, we expect that vertical variations of the confining potential are small compared to the length ℓ , as well as that the particles density value being sufficient for the Thomas-Fermi approximation to be robust.

Since the problem becomes essentially one-dimensional, the cloud can be characterized by its velocity $v = v(z)$ and linear particle density

$$\sigma = \sigma(z) = \int n(\mathbf{x}) dx dy, \quad (12)$$

where (x, y) are coordinates in transverse dimensions. Because in the previous section we established that the nonlinear Schrödinger equations for both models can be approximately rewritten in a hydrodynamic perfect-fluid form, we can derive hydrodynamic equations for v and σ by integrating over condensate lump's transverse coordinates [50]. From the continuity and Euler equations, we obtain, respectively:

$$\frac{\partial \sigma}{\partial t} + \frac{\partial(\sigma v)}{\partial z} = 0, \quad (13)$$

$$m\sigma \frac{dv}{dt} + \sigma \frac{\partial V}{\partial z} = - \int \frac{\partial p}{\partial z} dx dy, \quad (14)$$

where the external potential V occurs due to the trap and electromagnetic interactions, and pressure p is given by either of the formulae (10).

While the continuity equation remains common for both models, Euler equations must be derived separately. Using expressions (10), we obtain

$$m\sigma \frac{dv}{dt} + \sigma \frac{\partial V}{\partial z} = \begin{cases} -U_0 \int n \frac{\partial n}{\partial z} dx dy, \\ b \int \frac{\partial n}{\partial z} dx dy, \end{cases} \quad (15)$$

where we also assumed a constant b in the logarithmic case, remembering its thermal interpretation (6).

In order to simplify these formulae, let us recall that in 1D case the change of density n can be expressed as $dn = d\sigma/\mathcal{A}$, where \mathcal{A} the area of lump's cross section. Therefore, the Euler equations (15) become

$$m\sigma \frac{dv}{dt} + \sigma \frac{\partial V}{\partial z} = \begin{cases} -\frac{U_0}{\mathcal{A}} \sigma \frac{\partial \sigma}{\partial z}, \\ b \frac{\partial \sigma}{\partial z}, \end{cases} \quad (16)$$

where we also recalled the definition (12).

Let us consider the linear approximation, where we also assume that the variation of the external potential in the z direction is small compared to other terms. In case

of the Gross-Pitaevskii condensate, this would also mean that one can neglect spatial variations of the particle density averaged over the lump's cross section. Altogether, from (15) one can derive

$$\frac{\partial}{\partial z} \left(\sigma \frac{dv}{dt} \right) \approx \begin{cases} -\frac{U_0}{m} \frac{\partial}{\partial z} \left(\frac{\sigma}{\mathcal{A}} \frac{\partial \sigma}{\partial z} \right) \approx -\frac{\bar{n}U_0}{m} \frac{\partial^2 \sigma}{\partial z^2}, \\ \frac{b}{m} \frac{\partial^2 \sigma}{\partial z^2}, \end{cases}$$

where $\bar{n} = \sigma/\mathcal{A}$ is the particle density averaged over the lump's cross section. Finally, taking the total time derivative of the continuity equation and using the last formula, we obtain

$$\frac{\partial^2 \sigma}{\partial t^2} \approx \begin{cases} \frac{\bar{n}U_0}{m} \frac{\partial^2 \sigma}{\partial z^2}, \\ -\frac{b}{m} \frac{\partial^2 \sigma}{\partial z^2}, \end{cases} \quad (17)$$

where we are left only with terms which are linear with respect to σ .

If we assume that U_0 is positive, and b is negative, $b = -|b|$ (*i.e.*, $\chi > 0$ and $T < T_c$), then we obtain the hyperbolic-type partial differential equation in both cases. This indicates that small density fluctuations propagate in the direction perpendicular to the lump's cross section with the velocities

$$\bar{c} = \begin{cases} \sqrt{\bar{n}U_0/m}, \\ \sqrt{|b|/m}, \end{cases} \quad (18)$$

which are similar to the corresponding expressions for the homogeneous gas, except for the Gross-Pitaevskii condensate density must be replaced by the average density over the cross section of the lump. It also reaffirms the one-dimensional character of sound propagation in Bose-Einstein condensates of both types.

5. CONCLUSION

In this paper, we performed a comparative study of the propagation of small fluctuations of density in cigar-shaped Bose-Einstein condensates in Gross-Pitaevskii and logarithmic models, using the Thomas-Fermi and linear approximations.

It is shown that the propagation of sound pulses is effectively one-dimensional in both cases. The difference between them is that speed of sound scales as square root of particle density in a case of the Gross-Pitaevskii condensate, whereas this velocity is constant in a case of the logarithmic condensate. It should be emphasized that these results were obtained assuming a number of approximations, which do not necessarily hold in a presence of large density inhomogeneities. For example, logarithmic condensate, being essentially nonlinear, is known to form such inhomogeneities, because it behaves more like quantum liquid than gas [20, 21, 26]. Apart from the nonlinear phenomena, shock waves and other objects with effectively non-smooth density profiles can violate the above-mentioned approximations too.

Study of nonlinear and shock-wave effects in one-dimensional quantum Bose liquids under more general conditions goes beyond a scope of this report, and thus it will be a subject of future work.

Acknowledgments

Author gratefully acknowledges an invitation and a full waiver on this article's processing charges by Multidisciplinary Digital Publishing Institute to get it published in the Open Access form. This work is based on the research supported by the Department of Higher Education and Training of South Africa and in part by the National Research Foundation of South Africa (Grants Nos. 95965, 131604 and 132202).

-
- [1] Anderson, M.H.; Ensher, J.R.; Matthews, M.R.; Wieman, C.E.; Cornell, E.A. Observation of Bose-Einstein Condensation in a Dilute Atomic Vapor. *Science* **1995**, *269*, 198-201.
 - [2] Bradley, C.C.; Sackett, C.A.; Tollett, J.J.; Hulet, R.G. Evidence of Bose-Einstein Condensation in an Atomic Gas with Attractive Interactions. *Phys. Rev. Lett.* **1995**, *75*, 1687-1690.
 - [3] Andrews, M.R.; Kurn, D.M.; Miesner, H.-J.; Durfee, D.S.; Townsend, C.G.; Inouye, S.; Ketterle, W. Propagation of Sound in a Bose-Einstein Condensate. *Phys. Rev. Lett.* **1997**, *79*, 553-556.
Andrews, M.R.; Stamper-Kurn, D.M.; Miesner, H.-J.; Durfee, D.S.; Townsend, C.G.; Inouye, S.; Ketterle, W. Erratum: Propagation of Sound in a Bose-Einstein Condensate. *Phys. Rev. Lett.* **1998**, *80*, 2967.
 - [4] London, F. The λ -Phenomenon of Liquid Helium and the Bose-Einstein Degeneracy. *Nature* **1938**, *141*, 643-644.
 - [5] Snoke, D.W.; Wolfe, J.P.; Mysyrowicz, A. Evidence for Bose-Einstein Condensation of Excitons in Cu₂O. *Phys. Rev. B* **1990**, *41*, 11171-11184.
 - [6] Gross, E.P. Structure of a Quantized Vortex in Boson Systems. *Nuov. Cim.* **1961**, *20*, 454.
 - [7] Pitaevskii, L.P. Vortex Lines in an Imperfect Bose Gas. *Sov. Phys. JETP* **1961**, *13*, 451.
 - [8] Pethick, C.J.; Smith, H. *Bose-Einstein Condensation in Dilute Gases*; Cambridge University Press: New York, NY, USA, 2008.
 - [9] Van Gorder, R.A. Perturbation Theory for Bose-Einstein Condensates on Bounded Space Domains. *Proc. R. Soc. A* **2020**, *476*, 20200674.
 - [10] Efimov, V. Weakly-bound States of Three Resonantly-

- interacting Particles. *Sov. J. Nucl. Phys.* **1971**, *12*, 589.
- [11] Efimov, V. Energy Levels of Three Resonantly Interacting Particles. *Nucl. Phys. A* **1973**, *210*, 157.
- [12] Kolganova, E.A.; Motovilov, A.K.; Sandhas, W. The ^4He Trimer as an Efimov System. *Few-Body Syst.* **2011**, *51*, 249-257.
- [13] Kolomeisky, E.B.; Straley, J.P. Low-Dimensional Bose Liquids: Beyond the Gross-Pitaevskii Approximation. *Phys. Rev. B* **1992**, *46*, 11749.
- [14] Chui, S.T.; Ryzhov, V.N. Collapse Transition in Mixtures of Bosons and Fermions. *Phys. Rev. A* **2004**, *69*, 043607.
- [15] Chui, S.T.; Ryzhov, V.N.; Tareyeva, E.E. Stability of Bose system in Bose-Fermi Mixture with Attraction Between Bosons and Fermions. *JETP Lett.* **2004**, *80*, 274-279.
- [16] Zloshchastiev, K.G. Spontaneous Symmetry Breaking and Mass Generation as Built-in Phenomena in Logarithmic Nonlinear Quantum Theory. *Acta Phys. Polon.* **2011**, *42*, 261-292.
- [17] Rylov, Yu. A. Spin and Wave Function as Attributes of Ideal Fluid. *J. Math. Phys.* **1999**, *40*, 256-278.
- [18] Zloshchastiev, K.G. On the Dynamical Nature of Nonlinear Coupling of Logarithmic Quantum Wave Equation, Everett-Hirschman Entropy and Temperature. *Z. Naturforsch. A* **2018**, *73*, 619-628.
- [19] Brasher, J.D. Nonlinear Wave Mechanics, Information Theory, and Thermodynamics. *Int. J. Theor. Phys.* **1991**, *30*, 979-984.
- [20] Avdeenkov, A.V.; Zloshchastiev, K.G. Quantum Bose Liquids with Logarithmic Nonlinearity: Self-sustainability and Emergence of Spatial Extent. *J. Phys. B: At. Mol. Opt. Phys.* **2011**, *44*, 195303.
- [21] Zloshchastiev, K.G. Volume Element Structure and Roton-maxon-phonon Excitations in Superfluid Helium Beyond the Gross-Pitaevskii Approximation. *Eur. Phys. J. B* **2012**, *85*, 273.
- [22] De Martino, S.; Falanga, M.; Godano, C.; Lauro, G. Logarithmic Schrödinger-like Equation in Magma. *Europhys. Lett. (EPL)* **2003**, *63*, 472-475.
- [23] Bottiglieri, M.; Godano, C.; Lauro, G. Volcanic Eruptions: Initial State of Magma Melt Pulse Unloading. *Europhys. Lett. (EPL)* **2012**, *97*, 29001.
- [24] Zloshchastiev, K.G. Nonlinear Wave-mechanical Effects in Korteweg Fluid Magma Transport. *Europhys. Lett. (EPL)* **2018**, *122*, 39001.
- [25] Carles, R.; Carrapatoso, K.; Hillairet, M. Rigidity Results in Generalized Isothermal Fluids. *Ann. H. Lebesgue* **2018**, *1*, 47-85.
- [26] Zloshchastiev, K.G. Temperature-driven Dynamics of Quantum Liquids: Logarithmic Nonlinearity, Phase Structure and Rising Force. *Int. J. Mod. Phys. B* **2019**, *33*, 1950184.
- [27] Scott, T.C.; Zloshchastiev, K.G. Resolving the Puzzle of Sound Propagation in Liquid Helium at Low Temperatures. *Low Temp. Phys.* **2019**, *45*, 1231.
- [28] Zloshchastiev, K.G. An Alternative to Dark Matter and Dark Energy: Scale-dependent Gravity in Superfluid Vacuum Theory. *Universe* **2020**, *6*, 180.
- [29] Zloshchastiev, K.G. Superfluid Stars and Q-balls in Curved Spacetime. *Low Temp. Phys.* **2021**, *47*, 89.
- [30] Lasich, M.; Zloshchastiev, K.G. Particle Size and Phase Equilibria in Classical Logarithmic Fluid. *J. Phys.: Conf. Ser.* **2021**, *1740*, 012042.
- [31] Zloshchastiev, K.G. Resolving the Puzzle of Sound Propagation in a Dilute Bose-Einstein Condensate. *Int. J. Mod. Phys. B* **2022**, *36*, 2250121.
- [32] Zueva, T.I.; Sokolov, S.S. Self-localized Electron State in a Dense Helium Gas: Variational and Self-consistent Approaches. *Low Temp. Phys.* **2022**, *48*, 674-683.
- [33] Carles, R.; Ferriere, G. Logarithmic Schrödinger Equation with Quadratic Potential. *Nonlinearity* **2021**, *34*, 8283-8310.
- [34] Fan, H.; Feng, Z.; Yan, X. Positive Solutions for the Fractional Schrödinger Equations with Logarithmic and Critical Non-linearities. *Trans. London Math. Soc.* **2021**, *8*, 206-242.
- [35] Ji, C.; Xue, Y. Existence and Concentration of Positive Solution for a Fractional Logarithmic Schrödinger equation. *Differ. Integral Equ.* **2022**, *35*, 677-704.
- [36] Lima, F.C.E.; Petrov, A.Yu.; Almeida, C.A.S. Vortex Solutions in Nonpolynomial Scalar QED. *Phys. Rev. D* **2021**, *103*, 096019.
- [37] Lima, F.C.E.; Almeida, C.A.S. Differential Configurational Complexity and Phase Transitions of the BPS Solutions in the O(3)-sigma Model. *Ann. Phys.* **2022**, *442*, 168904.
- [38] Kai, Y.; Yin, Z. On the Gaussian Traveling Wave Solution to a Special Kind of Schrödinger Equation with logarithmic nonlinearity. *Mod. Phys. Lett. B* **2022**, *36*, 2150543.
- [39] Read, L.; Zegarliński, B.; Zhang, M. Logarithmic Schrödinger Equations in Infinite Dimensions. *J. Math. Phys.* **2022**, *63*, 111502.
- [40] Carles, R.; Su, C. Numerical Study of the Logarithmic Schrödinger Equation with Repulsive Harmonic Potential. *Discrete Contin. Dyn. Syst. B* **2022**, *Online*, DOI: 10.3934/dcdsb.2022206.
- [41] Wang, K.; Zhao, D.; Feng, B. Optimal Bilinear Control of the Logarithmic Schrödinger Equation. *Math. Meth. Appl. Sci.* **2022**, *Online*, DOI: 10.1002/mma.8840.
- [42] Zloshchastiev, K.G. Stability and Metastability of Trapless Bose-Einstein Condensates and Quantum Liquids. *Z. Naturforsch. A* **2017**, *72*, 677-687.
- [43] de Broglie, L. Sur la Possibilité de Relier les Phénomènes d'Interférence et de Diffraction à la Théorie des Quanta de Lumière. *Comptes Rendus* **1926**, *183*, 447-448.
- [44] Madelung, E. Quantentheorie in Hydrodynamischer Form. *Z. Phys.* **1927**, *40*, 322-326.
- [45] Halbwachs, F. *Théorie Relativiste des Fluides à Spin*; Gauthier-Villars: Paris, France, 1960.
- [46] Fröhlich, H. Macroscopic Wave Functions in Superconductors. *Proc. Phys. Soc.* **1966**, *87*, 330.
- [47] Spiegel, E.A. Fluid Dynamical Form of the Linear and Nonlinear Schrödinger Equations. *Phys. D: Nonlinear Phenom.* **1980**, *1*, 236-240.
- [48] Rylov, Yu.A. The Equations for Isentropic Motion of Inviscid Fluid in Terms of Wave Function. *J. Math. Phys.* **1989**, *30*, 2516-2520.
- [49] Zloshchastiev, K.G. Matrix Logarithmic Wave Equation and Multi-channel Systems in Fluid Mechanics. *J. Theor. Appl. Mech.* **2019**, *57*, 843-852.
- [50] Kavoulakis, G.M.; Pethick, C.J. Quasi-one-dimensional Character of Sound Propagation in Elongated Bose-Einstein Condensed Clouds. *Phys. Rev. A* **1997**, *58*, 1563-1566.