Page time and the order parameter for a consciousness state

Alexander Gorsky^{a,b}

^aInstitute for Information Transmission Problems, Moscow 127994, Russia ^bLaboratory of Complex Networks, Brain and Consciousness Research Center, Moscow

March 7, 2023

Abstract

In this Letter using the analogy with the recent resolution of the black hole information paradox we conjecture the order parameter for the state of consciousness based on the notion of the Page curve and the Page time. The entanglement between the state of the brain and time series of neuronal firing as well as the non-orthogonality of the functional connectomes play a key role.

1 Introduction and motivation

The origin of the consciousness certainly is the interesting scientific problem. It is desirable to identify the guidance principle behind the phenomenon and there are several ideas formulated to handle with this issue. In the integrated information theory(IIT) [1, 2] it is assumed that the entanglement of the clusters in the times series data for the neuronal firing could play the role of the order parameter of the consciousness state. The identification of the clusters can be simplified via the spectral analysis of the network formed from the time series [3]. The idea of the global neuronal workspace is based on the version of the self-organized criticality [4, 5]. It is assumed that interaction between the neuronal groups in nonlinear and transition to the consciousness state is a kind of phase transition. Another combination of ideas yields the model of competing cellular assemblies [6]. The neuronal group selection theory focuses at the evolutionary aspects of the formation of consciousness [7].

More physically motivated thermodynamical approach via a kind of the free energy principle was suggested in [8]. Some ideas based on the quantum decoherence were discussed in [9]. The review of the measures of consciousness can be found in [10] while the recent review of the most popular theories of consciousness can be found in [11].

Several features are quite common for all models

- The state of consciousness is related to consolidation of effective degrees of freedom in the brain functional connectomes
- There is specific timescale attributed to the formation of consciousness state from the non-conscious state

- Some version of entanglement serves as the order parameter for the consciousness state
- There is the interplay between the deterministic component and the ensemble average in the formation of consciousness state
- The cluster structure in the time series is important in some models
- The evolution of the system is unitary

We shall argue that these features can be naturally unified within the framework of the Page curve and Page time [12, 13] assuming the entanglement between the brain states and time series of neuron firing.

The Page curve has been invented to keep unitarity in a two-component system in a pure random state taking into account properly the entanglement of subsystems. The entanglement is evaluated via the density matrix projected into the subsystem [13]. The spectrum of projected density matrix before the Page time obeys the Marchenko-Pastur (MP) law but gets modified after the Page time. This approach has been applied for the entanglement of a black hole (BH) and its radiation [12] however the BH example is not unique and in principle one can consider any other "black box" entangled with its own radiation or time series of some activity.

The BH information paradox has been known since 70-ths [14] and it claims that the entropy production of the Hawking BH radiation violates a unitarity at large times, see [15, 16] for the recent reviews. The resolution of the information paradox has been found very recently and the island conjecture has been formulated. It was argued [17, 18] that the state of BH radiation is encoded in some island subregions behind the BH horizon. This has been quantified in the simplified model via end of world branes(EOW) [19], each of them has large number of internal degrees of freedom [20, 21]. The number of EOW branes grows with time and BH with EOW branes get entangled with the BH radiation. This is the first ingredient important for the resolution of the information paradox.

The second crucial ingredient goes as follows. The EOW branes interact and can develop the wormhole non-perturbative configurations connecting the pairs of the EOW branes. At the Page time the EOW branes get consolidated into the single disc geometry since the multiple non-perturbative wormholes start to dominate [21]. After the Page time the entropy of radiation goes down and the unitarity gets restored. It is important that after the Page time the Petz map known in the information theory [22, 23] allows to map the operations inside the BH Hilbert space into the operations in the Hilbert space of BH radiation [21]. Such mapping is impossible before the Page time. If the deterministic component is introduced into the system [24, 25] a kind of perturbative interaction between the EOW branes is induced.

We conjecture that somewhat similar picture emerges statistically during formation of consciousness state from the non-conscious one. The Hilbert space of brain and the Hilbert space of time series together are considered as the two-component entangled system. We assume that the functional connectomes are analogue of the EOW branes and encode the information concerning the time series of neuronal activity. The functional connectomes are non-orthogonal and interact in two ways. First they interact as multilayer networks and secondly there is the "contact interaction" since some neurons belong to several functional connectomes simultaneously. The state of brain with ensemble of functional networks is entangled with the time series and the number of functional networks involved grows in time. At time of formation of a consciousness state the functional networks get consolidated similar to the consolidation of EOW branes and the entropy attributed to time series stops to grow.

Hence we assume that the Page curve describes the time evolution of the entropy production in the time series of neuronal firing and at the Page time the functional connectomes get consolidated which is necessary condition for the establishing of a consciousness state. We conjecture that the sign of time derivative of the entropy production in the time series and clustering of the networks formed from the time series can serve as the indicators for a state of consciousness.

2 BH toy model

2.1 The entanglement of the BH and radiation

Let us recall the notion of the Page curve for a entangled two-component system. It was shown in [13] that the entanglement spectrum of the system in pure random state involving two subsystems and projected to one of them provides the important information concerning unitarity. Let rectangular matrix X describes uniformly random pure quantum state in $C^n \otimes C^m$. Then $Y = XX^+$ - reduced density matrix on C^n whose eigenvalue distribution depends on the evolution time. At small times spectral density of entanglement matrix Y obeys MP distribution [13]. However at some time, called the Page time the spectrum of the entanglement matrix gets modified and the entropy production in the subsystem which we have projected on starts to decrease. The entropy behaves according to Page curve as follows

$$S(n,m) = \left(\log n - \frac{n}{2m}\right)\theta(m-n) + \left(\log m - \frac{m}{2n}\right)\theta(n-m)$$
(1)

If the number of degrees of freedom in one subsystem grows with time the change of regime occurs exactly at the Page time,

The Page curve has been recognized recently in the context of the resolution of the BH information paradox. In this case the entangled two-component system is as follows. The one subsystem involves e_{BH}^S degrees of freedom where $S_{BH} = \frac{A}{4G_N}$ is the Bekenstein-Hawking BH entropy and A is the area of horizon. The second system describes the ongoing radiation of the BH and dimension of the corresponding Hilbert space k grows in time. These two subsystems are entangled and it was argued [17] that there is quantum extremal surface Q behind the BH horizon [26] encoding the BH radiation. The island conjecture assumes that the entropy of BH radiation reads as

$$S(\rho_R) = \min\left[\frac{Area(Q)}{4G_N} + S_{\text{bulk}}\right]$$
(2)

Here ρ_R is the density matrix of the BH radiation in the full theory coupled to quantum gravity. The second term corresponds to the von Neumann entropy of the region in the interior of the BH bounded by the quantum extremal surface Q. At small times the second contribution is small however at large times the leading saddle in the semi-classical path integral on the Euclidean black hole gets modified and the different extremal surface starts to dominate. The key point is that in this new phase the entropy of the radiation decreases in agreement with a unitarity. The corresponding behaviour with the Page time is presented in Fig.1

In the simplified version instead of the extremal surface one considers multiple EOW branes [20, 21] and analyses what is the saddle point configuration at the different times.




Figure 1: Page curve for the entropy of BH radiation(red)

Naively one considers the states

$$\sum_{i=1}^{k} |\Psi_i\rangle \otimes |i\rangle \tag{3}$$

where Ψ_i is the state of BH with the non-dynamical EOW branes behind the horizon while the state *i* belongs to the auxiliary basis of ongoing Hawking modes [21]. The states Ψ_i are not orthogonal which is the second important point behind the explanation of the BH information paradox

$$\langle \Psi_i | \Psi_j \rangle = e^{2S_{BH}} \delta_{ij} + O\left(e^{S_{BH}}\right). \tag{4}$$

The second term corresponds to the contribution from non-perturbative wormholes or effect of more direct EOW brane interactions and generically is subject to the ensemble averaging with some measure. Quantitatively the density matrix projected onto the radiation reads as

$$\rho_R = \sum_{i,j}^k \langle \Psi_j | \Psi_i \rangle | i \rangle \langle j | = \sum_{i,j}^k \sum_{a=1}^{e^S} C_{aj}^* C_{ai} | i \rangle \langle j |$$
(5)

In the simplest case the random rectangular $e^{S_{BH}} \times k$ matrices C are averaged with the Gauss measure with unit covariance

$$\int dC dC^+ \exp(-\operatorname{Tr} C^+ C) \tag{6}$$

The spectral density of ρ_R has been evaluated in [21] via solution to the Schwinger-Dyson equation for the resolvent in the planar approximation which takes into account the replica wormholes between the EOW branes. In microcanonical ensemble with unit Gaussian distribution it reads as

$$\rho_R(x) = \frac{ke_{BH}^S}{2\pi x} \sqrt{[x - (k^{-1/2} - e^{S_{BH}/2})^2][(k^{-1/2} + e^{S_{BH}/2})^2 - x)} + \delta(x)(k - e_{BH}^S)\theta(k - e_{BH}^S)$$
(7)

We see that its smooth part coincides with the MP distribution.

At some value of k the entropy of the BH and the entropy evaluated from the radiation density matrix ρ_R coincide. This happens at the Page time and the interpretation from the viewpoint of internal degrees of freedom behind the horizon has been developed in [21]. It has been argued that the semiclassical saddle point for EOW branes before the Page time implies that they can be considered as the separated degrees of freedom which nonperturbatively interact via wormhole solutions. However upon the summation of the replica wormholes at large times another saddle point dominates and the EOW branes form the




Figure 2: Islands behind the horizon are entangled with BH radiation. At the Page time the multiple islands get consolidated into the disc-like geometry.

single disk-like geometry which should be considered as single consolidated state of multiple EOW branes see Fig 2. The off-diagonal terms in the matrix elements start to dominate. After the Page time the Petz map [22, 23] allows to map the operations in the radiation Hilbert space into the operations in the BH Hilbert space [21].

2.2 BBP transition and entanglement spectrum

We have argued above that at the Page time the entanglement spectrum of the radiation gets modified and some number of the eigenvalues get separated from continuum after the Page time. This follows from the effect of non-orthogonality of the internal states of the BH. Recently it has been suggested [25] that the admixture of the determinism in the averaging over ensemble can induce the "perturbative" interaction of the EOW branes. Moreover it properly induces the eigenvalue separation phenomenon in the MP distribution necessary to get the Page curve.

The phenomenon of the eigenvalue separation is familiar in the probability theory when we perturb the Gaussian measure for random variables. We are interested in the deformation of the measure involved into the projected density matrix in the pure random states that is in the Wishart ensemble. Two types of measure deformation are known; the shifted Gaussian measure $\exp(-\operatorname{Tr}(H - H_0)^2)$ when the expansion around the nontrivial saddle takes place [27] and the measure with non-null covariance $\exp(-\operatorname{Tr} X^+ X\Sigma)$ [28,29]. In both cases there is the critical value of parameters involved in H_0 or Σ when one or several eigenvalues get decoupled from the bulk of the spectrum. It is a kind of phase transition (BBP transition) with own universality class and the distribution of the largest eigenvalue obeys the so-called spiked Tracy-Widom(TW) law exactly at the transition point.

To fit with our conjecture we modify the Gaussian measure not by the shift of mean value as in [25] but by introducing non-null complex covariance which does the same job [28,29]. This goes as follows. Let us consider M vectors $(\vec{y_1} \dots \vec{y_M})$ each of dimension N. The density of all vectors is Gaussian with covariance Σ , where Σ is positive $N \times N$ matrix

$$P(\vec{y}) = \frac{1}{(2\pi)^{N/2} (\det \Sigma)^{1/2}} \exp\left(-\frac{1}{2} \left< \vec{y}, \Sigma^{-1}, \vec{y} \right>\right)$$
(8)

Consider the $N \times M$ matrix $X = [\mathbf{y}_1 - \mathbf{Y}, \dots, \mathbf{y}_M - \mathbf{Y}]$ and combine it into $N \times N$ covariance matrix $S = XX^t$. Let us assume that $M \to \infty$ and $N \to \infty$ such that their ratio is finite $\frac{M}{N} = \gamma$. The matrix S belongs to the Laguerre or Wishart ensemble whose

asymptotic properties at $\Sigma = Id$ are known. The spectral density obeys the MP law

$$\rho(x) = \frac{\gamma^2}{2x} \sqrt{(b-x)(x-a)}, \qquad a < x < b \tag{9}$$

where

$$a = \left(\frac{\gamma - 1}{\gamma}\right)^2; \qquad b = \left(\frac{\gamma + 1}{\gamma}\right)^2 \tag{10}$$

The largest eigenvalue fixing the spectral edge is $\lambda_{max} = b$ and the spectral fluctuations at the spectral edge obey the TW distribution

$$P\left((\lambda_{max} - b)\frac{\gamma}{(1+\gamma)^{4/3}}M^{2/3} \le s\right) \to F_{GOE}(s) \tag{11}$$

where F_{GOE} is the TW distribution for orthogonal ensemble. Now departure from the identity matrix is introduced and some number of degenerate non-unit eigenvalues of the covariance matrix Σ are selected $l_1 \dots l_k \neq 1$. The value l_1 is considered as the control parameter for the BBP phase transition.

For complex Gaussian samples [28] the critical value of the non-unit covariance eigenvalue for perturbed case has been found $l_{crit} = 1 + \gamma^{-1}$ where γ is a parameter of the ensemble.

$$P\left((\lambda_{max} - l_{crit}^2)\frac{\gamma}{(1+\gamma)^{4/3}}N^{2/3} \le s\right) \to F_k(s) \tag{12}$$

The function $F_k(s)$ is the limiting distribution, where the largest eigenvalue l_1 of the covariance matrix has the multiplicity k and is equal to the critical value, hence we have single perturbation parameter. If k non-unit eigenvalues of the covariance matrix are above the critical value l_{crit} , i.e. $l_i > l_{crit}$, distribution of λ_{max} obeys the distribution in $k \times k$ Gaussian unitary ensemble, $G_k(s)$.

$$P\left(\left(\lambda_{max} - \left(l_1 + \frac{l_1\gamma^{-2}}{l_1 - 1}\right)\right) N^{1/2} \sqrt{l_1^2 - \frac{l_1^2\gamma^{-2}}{(l_1 - 1)^2}} \le s\right) \to G_k(s) \tag{13}$$

The function $F_k(s)$ can be generalized to the function $F_k(s, w_1, \ldots, w_k)$ if all perturbation parameters are different. Similarly to the pure TW case, the distribution obeys some solution to the Painleve II equation with particular monodromy properties [29]. In general case the determinantal representation of $F_k(s, w_1, \ldots, w_k)$ is available in terms of the single function of two variables f(s, w) which satisfy the pair of differential equations depending on the Painleve II solutions – see [29] for details.

Summarizing, there are three regimes for the spectral density depending on the relation between k and $e^{S_{BH}}$. The transition at the Page time indicates that the new saddle in the gravitational path integral corresponding to connected geometry with multiple wormholes starts to dominate at $k \propto e^{S_{BH}}$. In terms of the eigenvalues exactly at the Page transition $k - e^{S_{BH}}$ eigenvalues get escaped from the bulk of distribution and settle at $\lambda = 0$. If somebody would be interested in the question if all EOW branes get consolidated or not one should look at the sign of the time derivative for the radiation entropy and at the number of isolated eigenvalues. This is the order parameter separating the phases.

3 Towards the order parameter for a consciousness state

3.1 Entropy for the time series

Since we will assume that the time series of the brain activity is the part of the whole two-component system it is necessary to provide the tool to evaluate the entropy of the time series similarly to the entropy of the BH radiation. The effective way goes as follows, first we represent the time series data via the network and than evaluate the entropy of this network.

There are at least three ways to generate the network from the time series (see, [30] for the review). The first way involves the introduction of the effective parameter which quantifies the vicinity in the data set and the nodes within this effective radius are connected by links. In the second approach the notion of visability is introduced and the horizontally visable nodes are connected by links. Finally, in the third approach a kind of Markov dynamics for the probability is formulated and its Markov operator yields the network adjacency matrix.

Once we find the network representation for the time series some standard entropy measured can be applied. The simplest one is the conventional thermodynamic entropy defined through the density metrics for the network

$$\rho = \frac{\exp(-\beta L)}{Z} \tag{14}$$

where L is the graph Laplacian or normalized graph Laplacian, and β is auxiliary parameter which allows to scan the spectrum. The thermodynamical entropy is defined as

$$S = \operatorname{Tr} \rho \log \rho \tag{15}$$

and is useful to determine the characteristic time scales for the diffusion in the network [31].

However we are interested in the entropy production hence we have to consider the coarse-grained entropy instead of the fine-grained entropy which is constant in time. The useful coarse-grained information entropy production measure has been discussed in [32] which is suitable for our purpose,

$$\dot{S} = \sum_{ij} P_{ij} log(\frac{P_{ij}}{P_{ji}}) \tag{16}$$

where $P_{ij} = Prob[x_{t-1} = i, x_t = j]$ and x_t corresponds to the state of the system at time t. It is nothing but the Kullback-Leibler divergence between the forward and reverse transition probabilities. This measure for the entropy production for different tasks has been investigated experimentally [32]. It was shown that the entropy production vanishes or is very small in non-conscious state, is quite substantial for the cognitive task and maximal at the physical activity. Remark that the entropy production provides the thermodynamical time arrow and it was noted recently that in the interacting many-body system the time arrow emerges via the interaction only [33].

3.2 The conjecture

To formulate our conjecture remind that the brain involves structural and multiple functional networks. The structural network just reflects the connections between neurons and is completely known for C'Elegance with 302 nodes only. Each functional network or functional connectome involves all neurons specialized in some function. They are connected to each other and can be located in the very different parts of the brain. The functional networks are assumed to interact and the individual neurons can belong to the several functional networks simultaneously. It is generally believed that in the state of consciousness the functional connectomes get consolidated via some mechanism.

Let us apply the Page's ideas to the brain and assume that the functional connectomes play the similar role as EOW branes in the BH example while the spiking time series play




Figure 3: The functional connectome represents the network of neurons with the same specialization. There are multiple functional connectomes in the brain which can overlap due to the multispecialized neurons. We conjecture that functional connectomes are entangled with the time series of neuron activity and at the effective Page time when the state of consciousness gets established they get consolidated.

the role of BH radiation. Hence we have the "black box" with some degrees of freedom inside and its radiation and would like to investigate the phase of internal state of the brane looking at the properties of time series only.

Let us follow the same logic as before and assume that the state of the system involves the state Ξ_i which corresponds to the state of the brain with the functional connectomes in the state i and the states $|i\rangle$ which are the eigenfunctions of the Laplacian of the graph representing the time series. Hence we consider the states

$$\sum_{i} |\Xi_i\rangle\langle i| \tag{17}$$

The density matrix projected to the time series state ρ_{TS} is

$$\rho_{TS} = \sum_{i,j}^{k} \langle \Xi_j | \Xi_i \rangle |i\rangle \langle j| = \sum_{i,j}^{k} \sum_{a=1}^{S_{brain}} C_{aj}^* C_{ai} |i\rangle \langle j|$$
(18)

where S_{brane} is the entropy of the brain in the non-conscious state. The value of k measures the number of functional connectomes which are switched on.

The projected density matrix yields the entanglement entropy of the spiking time series S_{TS} . To keep the unitarity we assume the existence of the Page curve and Page time when the internal functional networks get consolidated, see Fig.3, providing the state of consciousness at the Page time. After the Page time presumably the Petz map yields the mapping of the operations within the time series to the operations between the functional connectomes in the brain.

As in the BH toy example the functional connectomes overlap and are non-orthogonal. The interaction of the functional connectomes can have perturbative and non-perturbative contributions. The simplest one is generated by " contact interaction" we have described above. In the case of a brain this interaction can be attributed to the neurons with several specializations. Due to the overlap of functional connectomes the effective brain entropy and the entropy of the time series become of the same order and the "common space" of the functional connectomes gets formed at the Page time. Note that the contact interaction amounts to the non-unit covariance matrix which induces the isolated eigenvalues in the

spectrum corresponding to the clusters in the time series network [34]. Such clusters in the time series networks are the essential ingredients of IIT.

Our conjecture fits with the general expected features of the consciousness state

- The consolidation of the functional networks occurs at the Page time
- There is the timescale when consciousness state gets established
- Order parameter indeed is related to some specific entanglement
- Deterministic component is important. It induces the effective interactions and simultaneously generates the clusters in the time series

Summarizing the arguments above we suggest that the sign of the time derivative of the entropy production in the time series serves as the order parameter R for a consciousness state

$$R = \operatorname{sign}\left(\frac{dS_{TS}}{dt}\right). \tag{19}$$

Another important indicator is the increasing of the clusters in the network formed from the time series. We can not exclude that the entropy production stop raising at the Page and decreases very slowly at the late times.

Hence it is natural to question how the predictions following from our conjecture can be measured experimentally and if there are any contradictions with the existing experimental data. The simplest possibility concerns the investigation of the entropy production and clusterization of the time series during the waking up process and the exit from the state of anesthesia. Hopefully some relevant results concerning the anesthesia state are available. It was found in [35] that at exit from anesthesia two effects are observed both for mice and humans (see also [36–39]). First, the number of clusters in the networks formed by time series (microstates in notations of [35]) increases substantially and secondly, consolidation of the functional connectomes gets increased as well. Moreover both features are reversible while depth of consolidation depends on the type of anesthesia applied [40]. This is consistent with our conjecture.

Note also that the establishing time of consciousness state is different for the waking up and the anesthesia exit processes. That is we can assume that the number of functional networks switched off at anesthesia is larger then at sleep state. It would be very interesting to measure the entropy production at the anesthesia process to compare with our conjecture.

4 Conclusion

In this Letter we have conjectured using very general statistical arguments for the unitary two-component entangled many-body system the new order parameter for the consciousness state which measures if the functional connectomes are consolidated or not. Certainly it is necessary but insufficient criteria. Two points are crucial for this conjecture. First, it is the entanglement between the time series of the neuron activity and the brain state involving multiple overlapping functional connectomes. Secondly, the non-orthogonality of the states of functional connectomes which is small at early time but starts to dominate at the effective Page time. The mechanism behind the resolution of the information paradox for a BH radiation provides the proper framework to analyze this problem. The clustering of the time series density matrix occurs naturally within this approach and is another indication of the state of consciousness.

The conjecture is a bit speculative but we believe that this framework could be useful for this research area and suggest the very clear-cut check. We do not provide the microscopic explanation of the consciousness state restricting ourselves by the order parameter and do not present the hypothetical functional whose extremization presumably yields two different saddle points. However we expect that some notions of the information geometry could be useful in this respect. In particular the recent observation [41] concerning the relation between the singularity of the information metric in the simplest all-to-all disordered Kuramoto model and the synchronization phase transition provides the example of such relation. Indeed microscopically the consolidation of the functional connectomes implies the synchronization phenomena in the multiplex networks. Another approach concerning the quantum information geometry approach can be found in [42]. Note also that some analogy between the brain and the BH in a different setting has been mentioned in [43]. It would be interesting to combine the ideas from that paper with our study.

I am grateful to A. Alexandrov, K. Anokhin, A. Milekhin, S. Nechaev and N. Pospelov for the useful discussions and comments. I want to thank IHES and Nordita where the parts of the work have been done for the hospitality and support. The work was supported by grant №075-15-2020-801 by Ministry of Science and Higher Education of Russian Federation.

References

- Giulio Tononi and Gerald M Edelman. Consciousness and complexity. science, 282(5395):1846–1851, 1998.
- [2] Giulio Tononi. An information integration theory of consciousness. BMC neuroscience, 5(1):1–22, 2004.
- [3] Daniel Toker and Friedrich T Sommer. Information integration in large brain networks. PLoS computational biology, 15(2):e1006807, 2019.
- [4] Stanislas Dehaene and Jean-Pierre Changeux. Experimental and theoretical approaches to conscious processing. *Neuron*, 70(2):200–227, 2011.
- [5] George A Mashour, Pieter Roelfsema, Jean-Pierre Changeux, and Stanislas Dehaene. Conscious processing and the global neuronal workspace hypothesis. *Neuron*, 105(5):776–798, 2020.
- [6] Francis Crick and Christof Koch. A framework for consciousness. Nature neuroscience, 6(2):119–126, 2003.
- [7] Gerald M Edelman. Wider than the sky: The phenomenal gift of consciousness. Yale University Press, 2004.
- [8] Karl Friston. The free-energy principle: a unified brain theory? Nature reviews neuroscience, 11(2):127–138, 2010.
- [9] Stuart Hameroff and Roger Penrose. Consciousness in the universe: A review of the 'orch or'theory. *Physics of life reviews*, 11(1):39–78, 2014.
- [10] Anil K Seth, Eugene Izhikevich, George N Reeke, and Gerald M Edelman. Theories and measures of consciousness: an extended framework. *Proceedings of the National Academy of Sciences*, 103(28):10799–10804, 2006.

- [11] KV Anokhin. The cognitome: Seeking the fundamental neuroscience of a theory of consciousness. Neuroscience and Behavioral Physiology, 51(7):915-937, 2021.
- [12] Don N Page. Information in black hole radiation. Physical review letters, 71(23):3743, 1993.
- [13] Don N Page. Average entropy of a subsystem. *Physical review letters*, 71(9):1291, 1993.
- [14] Stephen W Hawking. Breakdown of predictability in gravitational collapse. Physical Review D, 14(10):2460, 1976.
- [15] Ahmed Almheiri, Thomas Hartman, Juan Maldacena, Edgar Shaghoulian, and Amirhossein Tajdini. The entropy of hawking radiation. *Reviews of Modern Physics*, 93(3):035002, 2021.
- [16] Suvrat Raju. Lessons from the information paradox. *Physics Reports*, 943:1–80, 2022.
- [17] Geoffrey Penington. Entanglement wedge reconstruction and the information paradox. Journal of High Energy Physics, 2020(9):1–84, 2020.
- [18] Ahmed Almheiri, Netta Engelhardt, Donald Marolf, and Henry Maxfield. The entropy of bulk quantum fields and the entanglement wedge of an evaporating black hole. *Journal of High Energy Physics*, 2019(12):1–47, 2019.
- [19] Ioanna Kourkoulou and Juan Maldacena. Pure states in the syk model and nearlyads_2 gravity. arXiv preprint arXiv:1707.02325, 2017.
- [20] Ahmed Almheiri, Thomas Hartman, Juan Maldacena, Edgar Shaghoulian, and Amirhossein Tajdini. Replica wormholes and the entropy of hawking radiation. *Jour*nal of High Energy Physics, 2020(5), may 2020.
- [21] Geoff Penington, Stephen H Shenker, Douglas Stanford, and Zhenbin Yang. Replica wormholes and the black hole interior. *Journal of High Energy Physics*, 2022(3):1–87, 2022.
- [22] Dénes Petz. Sufficient subalgebras and the relative entropy of states of a von neumann algebra. Communications in mathematical physics, 105(1):123–131, 1986.
- [23] Mark M Wilde. Quantum information theory. Cambridge University Press, 2013.
- [24] Andreas Blommaert and Jorrit Kruthoff. Gravity without averaging. SciPost Physics, 12(2), feb 2022.
- [25] Andreas Blommaert and Mykhaylo Usatyuk. Microstructure in matrix elements. Journal of High Energy Physics, 2022(9):1–42, 2022.
- [26] Netta Engelhardt and Aron C Wall. Quantum extremal surfaces: holographic entanglement entropy beyond the classical regime. *Journal of High Energy Physics*, 2015(1):1–27, 2015.
- [27] Kevin E Bassler, Peter J Forrester, and Norman E Frankel. Eigenvalue separation in some random matrix models. *Journal of mathematical physics*, 50(3):033302, 2009.

- [28] Jinho Baik, Gérard Ben Arous, and Sandrine Péché. Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices. *The Annals of Probability*, 33(5):1643–1697, 2005.
- [29] Jinho Baik. Painlevé formulas of the limiting distributions for nonnull complex sample covariance matrices. Duke Mathematical Journal, 133(2):205–235, 2006.
- [30] Thomas F Varley and Olaf Sporns. Network analysis of time series: Novel approaches to network neuroscience. *Frontiers in Neuroscience*, 15, 2021.
- [31] Manlio De Domenico and Jacob Biamonte. Spectral entropies as information-theoretic tools for complex network comparison. *Physical Review X*, 6(4):041062, 2016.
- [32] Christopher W Lynn, Eli J Cornblath, Lia Papadopoulos, Maxwell A Bertolero, and Danielle S Bassett. Broken detailed balance and entropy production in the human brain. *Proceedings of the National Academy of Sciences*, 118(47):e2109889118, 2021.
- [33] Christopher W Lynn, Caroline M Holmes, William Bialek, and David J Schwab. Decomposing the local arrow of time in interacting systems. *Physical review letters*, 129(11):118101, 2022.
- [34] Raj Rao Nadakuditi and Mark EJ Newman. Spectra of random graphs with arbitrary expected degrees. *Physical Review E*, 87(1):012803, 2013.
- [35] Michael Wenzel, Shuting Han, Elliot H Smith, Erik Hoel, Bradley Greger, Paul A House, and Rafael Yuste. Reduced repertoire of cortical microstates and neuronal ensembles in medically induced loss of consciousness. *Cell systems*, 8(5):467–474, 2019.
- [36] Andrea I Luppi, Daniel Golkowski, Andreas Ranft, Rüdiger Ilg, Denis Jordan, David K Menon, and Emmanuel A Stamatakis. Brain network integration dynamics are associated with loss and recovery of consciousness induced by sevoflurane. *Human Brain* Mapping, 42(9):2802–2822, 2021.
- [37] Richard Boyce, Robin F Dard, and Rosa Cossart. Cortical neuronal assemblies coordinate with eeg microstate dynamics during resting wakefulness. *Cell Reports*, 42(2), 2023.
- [38] Arjun Bharioke, Martin Munz, Alexandra Brignall, Georg Kosche, Max Ferdinand Eizinger, Nicole Ledergerber, Daniel Hillier, Brigitte Gross-Scherf, Karl-Klaus Conzelmann, Emilie Macé, et al. General anesthesia globally synchronizes activity selectively in layer 5 cortical pyramidal neurons. *Neuron*, 110(12):2024–2040, 2022.
- [39] André S Nilsen, Bjørn Juel, Benjamin Thürer, and Johan Frederik Storm. Proposed eeg measures of consciousness: a systematic, comparative review. 2020.
- [40] Thomas F Varley, Olaf Sporns, Aina Puce, and John Beggs. Differential effects of propofol and ketamine on critical brain dynamics. *PLoS computational biology*, 16(12):e1008418, 2020.
- [41] Artem Alexandrov and Alexander Gorsky. Information geometry and synchronization phase transition in kuramoto model. *arXiv preprint arXiv:2211.15617*, 2022.
- [42] Danko D Georgiev. Quantum information theoretic approach to the mind-brain problem. Progress in Biophysics and Molecular Biology, 158:16–32, 2020.

[43] Gia Dvali. Black holes as brains: neural networks with area law entropy. *Fortschritte* der Physik, 66(4):1800007, 2018.