arXiv:2212.13433v2 [astro-ph.CO] 25 Apr 2024

FRB dark sirens: Measuring the Hubble constant with unlocalized fast radio bursts

Ze-Wei Zhao,¹ Ji-Guo Zhang,¹ Yichao Li,¹ Jing-Fei Zhang,¹ and Xin Zhang^{1,2,3,*}

¹Key Laboratory of Cosmology and Astrophysics (Liaoning Province) & Department of Physics,

College of Sciences, Northeastern University, Shenyang 110819, China

²Key Laboratory of Data Analytics and Optimization for Smart Industry (Ministry of Education),

Northeastern University, Shenyang 110819, China

³National Frontiers Science Center for Industrial Intelligence and Systems Optimization,

Northeastern University, Shenyang 110819, China

Fast radio bursts (FRBs) can be used to measure cosmological parameters by employing the Macquart relation. However, at present, only a small number of FRB events are localized to host galaxies with known redshifts. Inspired by the dark siren method in gravitational wave cosmology, we develop a Bayesian method to statistically measure the Hubble constant using unlocalized FRBs and galaxy catalog data, which makes it possible to constrain cosmological parameters from a large number of FRB data without known redshifts, meanwhile including the real galaxy information. We assume that the probability for a galaxy to host an FRB is proportional to the luminosity of this galaxy and use the results from the IllustrisTNG simulation as the priors of FRB host galaxy parameters. Ignoring some systematic errors, we obtain the first statistical H_0 measurement only using twelve unlocalized FRB events combined with the big bang nucleosynthesis result, i.e., $H_0 = 80.4^{+24.1}_{-19.4}$ km s⁻¹ Mpc⁻¹ (68% highest-density interval). This method can also be refined to constrain other cosmological and FRB parameters. It is applicable to well-localized FRBs that still have several potential hosts.

I. INTRODUCTION

The Hubble tension has now been one of the most important issues in current cosmology. It shows that the values of the Hubble constant H_0 estimated by the earlyand late-universe observations are apparently inconsistent. Actually, the disagreement of the measurements of H_0 has reached 4.8 σ . The constraint from the Planck 2018 data of the cosmic microwave background (CMB) anisotropies [1] gives $H_0 = 67.36 \pm 0.54 \,\mathrm{km \, s^{-1} \, Mpc^{-1}}$, based on the Λ cold dark matter (Λ CDM) model, and the direct measurement by the SH0ES team [2] shows $H_0 = 73.04 \pm 1.04 \,\mathrm{km \, s^{-1} \, Mpc^{-1}}$, using the "distance ladder" method. In order to solve the Hubble tension, some extensions of the Λ CDM model have been proposed, but a new concordance model has not yet been widely accepted [3–7].

Another possible explanation for the Hubble tension may be the unknown systematic errors in the observations. However, the current reexaminations of uncertainties are unable to fully solve the tension [6]. Therefore, it is necessary to develop new precise cosmological probes to provide a cross-check. A key factor in developing a precise probe is that there should be a vast number of data that can be used to reduce the random error. Fast radio bursts (FRBs) are a kind of new astronomical phenomenon and it is possible for FRBs to accumulate a lot of data in the future (see Refs. [8–14] for reviews), due to their high event rate [15].

FRBs are luminous millisecond pulses detected in the radio band. When FRB photons propagate through plasma, it will interact with free electrons and generate dispersion between different frequencies. The observed high dispersion measures (DMs) greatly exceed the expectation value of the Milky Way, indicating that most FRBs are likely of extragalactic origin, except one FRB event associated with a Galactic magnetar [16, 17]. If an FRB can be localized to a unique galaxy, the redshift of this FRB can be inferred from its host galaxy [18–20].

Although the sources and radiation mechanisms of most FRBs have not been generally identified [21–23], one can still use them as a cosmological probe. Since DMs contain the baryonic information along the cosmological distances FRBs' photons travelled, localized FRBs can be used to constrain cosmological parameters through the Macquart relation (i.e., the DM-z relation) [24-42] and to constrain related parameters [43-50]. An outstanding application is shown in solving the "missing baryon" problem. Macquart et al. [51] used five localized FRBs to derive a cosmic baryon density constraint which is consistent with the big bang nucleosynthesis (BBN) and CMB measurements. To study the Hubble tension, the H_0 measurements from localized FRBs were obtained by several independent groups [52-59], with a precision of around 10%.

However, despite hundreds of FRB events having been detected, little more than forty FRBs are currently localized to their host galaxies [60]. The application of FRBs in cosmology is greatly limited because it is difficult to get the redshift information of FRBs. In order to exploit the potential of FRB data in cosmology research, it is necessary to find a way of using unlocalized FRBs to constrain cosmological parameters. In Ref. [61], authors used the information from sky coordinates, galaxy fluxes, and angular sizes to estimate the probability that an FRB is associated with a candidate host.

^{*} Corresponding author.

zhangxin@mail.neu.edu.cn

In this work, we extend the dark siren method from gravitational wave (GW) cosmology [62–67] to the FRB field. We statistically measure the Hubble constant using the DM data of unlocalized FRBs in conjunction with galaxy catalogs.

In the dark siren method, we utilize GW events, such as binary black hole (BBH) mergers, which lack electromagnetic (EM) counterparts. By integrating these events with galaxy catalog data, we can constrain cosmological parameters by associating each GW event with its potential host galaxies. This method operates within a Bayesian framework and has been validated with mock data [68]. Similarly, the Hubble constant can be constrained using unlocalized FRBs by identifying all potential host galaxies within their localization region and subsequently marginalizing over these possibilities.

Most previous studies have focused on using localized FRBs to constrain the Hubble constant [52–54, 57–59], while another approach for unlocalized FRBs involves marginalizing the likelihood across the entire redshift range [0.01, 5] [55, 56]. The advantage of our method is that it allows for the constraint of cosmological parameters using a large dataset of unlocalized FRBs, while also incorporating actual galaxy information (including host data) into the analysis. This enhancement potentially improves the utility of FRBs as a new cosmological probe in two significant ways: it increases the volume of data utilized and tightens the constraints derived from each unlocalized FRB event.

This paper is organized as follows. In Sec. II, we briefly describe the galaxy catalog and FRB data. In Sec. III, the model of the likelihood and the Bayesian method are introduced. The constraints and relevant discussion are given in Sec. IV. Conclusions are given in Sec. V.

II. DATA

A. Galaxy catalog data

The Dark Energy Spectroscopic Instrument (DESI) project is a stage IV dark energy measurement aimed at measuring the expansion of the universe and studying the physics of dark energy, by constructing a threedimensional map of the large-scale structure of the universe. The DESI Legacy Imaging Surveys have observed the sky in three optical (g, r, and z) bands [69], to provide the targets for the DESI survey. The optical imaging data are collected by three independent programs (the Dark Energy Camera Legacy Survey, the Beijing-Arizona Sky Survey, and the Mayall z-band Legacy Survey). The 5σ detection thresholds are 24.0, 23.4, and 22.5 AB magnitudes for the q, r, and z bands, respectively, for a fiducial galaxy with an exponential light profile and half-light radius 0.45 arcsec. The optical data are combined with the infrared data at two bands, i.e., W1 (3.4 μ m) and W2 (4.6 μ m), observed from the Wide-field Infrared Survey Explorer (WISE) satellite [70]. We use the processed DESI galaxy catalog data published in Ref. [71], in which the DESI Legacy Imaging Surveys DR8 data are considered. In total, 129.35 million galaxies are selected based on the selection criteria in Ref. [71].

B. FRB data

The Australian Square Kilometre Array Pathfinder (ASKAP) project can provide precise localization for FRB events [72]. The Commensal Real-time ASKAP Fast Transients (CRAFT) group [73] has performed FRB surveys in two modes: single-antenna ("Flye's Eye", or "FE") mode during the "lat50" survey, i.e. observing at Galactic latitudes $|b| \sim 50^{\circ}$ [74, 75], and incoherent sum (ICS) mode, incoherently adding the spectra from all antennas to localize FRBs at sub-arcsecond precision [19]. The FRBs observed by the ICS mode are mainly localized, so in this work we only use the FRB data observed by the FE mode.

We then select the FRB events whose localization region is covered by the sky coverage of the catalog data of the DESI Legacy Imaging Surveys. We also follow the "gold-standard sample" criteria of FRBs for cosmological study [51]¹, and finally 12 FRB data are selected. The selected FRB data and their properties [75–77] are listed in Table I, and their positions in the galaxy catalog are shown in Fig. 1.

III. METHODOLOGY

A. Dispersion measure distribution

We use a DM model similar to Ref. [51]. The observed DM of an FRB consists of contributions from the Milky Way's interstellar medium (ISM), our Galactic halo, the cosmological distribution of the intergalactic medium (IGM) and galaxy haloes, and the FRB host galaxy,

$$DM = DM_{MW,ISM} + DM_{MW,halo} + DM_{cosmic} + DM_{host}.$$
(1)

We use the NE2001 model [78] to estimate $DM_{MW,ISM}$. The contribution from our Galactic halo with a hot, diffuse gas is still uncertain. We will discuss it below Eq. (8).

The average value of DM_{cosmic} at redshift z is given by the Macquart relation [24, 79, 80],

$$\langle \mathrm{DM}_{\mathrm{cosmic}} \rangle = \int_{0}^{z} \frac{c \bar{n}_{e}(z') dz'}{H_{0}(1+z')^{2} E(z')} \\ = \frac{3c f_{\mathrm{d}} \Omega_{\mathrm{b}} H_{0}^{2}}{8\pi G m_{\mathrm{p}} H_{0}} \int_{0}^{z} \frac{\chi(z')(1+z') dz'}{E(z')}, \qquad (2)$$

¹ The first criterion [51] only applies to localized FRBs, so we use the rest of criteria.



FIG. 1. The sky positions of the FRB data (black circles) and the footprints of the galaxy catalog (the blue area) used in this paper. The left and right panels show the data in the south Galactic cap (SGC) and north Galactic cap (NGC), respectively.

TABLE I. Properties of unlocalized FRBs observed by ASKAP and the numbers of potential host galaxies (N_{gal}) for each FRB. The names and properties of FRBs are taken from the public Transient Name Server website^a.

FRB event	Right ascension	Declination	$DM (pc cm^{-3})$	$N_{\rm gal}$
20170107A	$11h23m18s\pm10.5$	$-05^{\circ}00'00'' \pm 10'$	609.5	911
20170416A	$22h13m00s{\pm}15'$	$-10^{\circ}56'00"\pm 9'$	523.2	375
20170712A	$22h36m00s{\pm}15'$	$-60^{\circ}57'00"\pm10'$	312.8	551
20171116A	$03h31m00s\pm10$ '	$-17^{\circ}14'00"\pm10'$	618.5	596
20171213A	$03h39m00s\pm30'$	$-10^{\circ}56'00'' \pm 20'$	158.6	4363
20180119A	$03h29m18s\pm8'$	$-12^{\circ}44'00"\pm 8'$	402.7	427
20180128A	$13h56m00s\pm20'$	$-06^{\circ}43'00"\pm15'$	441.4	2425
20180131A	$21\mathrm{h}49\mathrm{m}54\mathrm{s}{\pm}12^{\prime}$	$-40^{\circ}41'00"\pm8'$	657.7	557
20180212A	$14h21m00s\pm30'$	-03°35'00"±30'	167.5	6811
20180417 A	$12h24m56s\pm7'$	$+14^{\circ}13'00"\pm7'$	474.8	485
20180515A	$23h13m12s\pm7$	$-42^{\circ}14'46''\pm7'$	355.2	270
$20180525 \mathrm{A}$	$14h40m00s\pm30'$	$-02^{\circ}12'00"\pm6'$	388.1	1257

^a https://www.wis-tns.org/

where \bar{n}_e is the mean free electron density, Ω_b is the present-day baryon density parameter, f_d is the fraction of cosmic baryons in diffuse ionized gas, $\chi(z)$ represents the fraction of ionized electrons in hydrogen and helium atoms, and m_p is the mass of a proton. The ionization fraction is

$$\chi(z) = Y_{\rm H}\chi_{\rm e,H}(z) + \frac{1}{2}Y_{\rm He}\chi_{\rm e,He}(z).$$
 (3)

where $Y_{\rm H} = 3/4$ and $Y_{\rm He} = 1/4$ are the mass fractions of hydrogen and helium, respectively, and $\chi_{\rm e,H}$ and $\chi_{\rm e,He}$ are the ionization fractions for hydrogen and helium, respectively. We take $\chi_{\rm e,H} = \chi_{\rm e,He} = 1$, assuming both hydrogen and helium are fully ionized at z < 3 [81]. For $f_{\rm d}$, we use a redshift-dependent form [51] calculated from the public FRB code².

In the flat Λ CDM model, the dimensionless Hubble parameter is

$$E(z) = \sqrt{\Omega_{\rm m}(1+z)^3 + (1-\Omega_{\rm m})},$$
 (4)

where $\Omega_{\rm m}$ is the present-day matter density parameter. We focus on the Hubble constant measurement and set $\Omega_{\rm m} = 0.315$ [1]. With more FRB data in future works, the influence of cosmological models and other cosmological parameters could be further studied.

Equation (2) reflects that $\langle \text{DM}_{\text{IGM}} \rangle$ is proportional to $\Omega_{\text{b}}h$, with $h = H_0/(100 \,\text{km}\,\text{s}^{-1}\text{Mpc}^{-1})$. However, $\Omega_{\text{b}}h^2$ is usually the directly observed quantity in other cosmological observations, so we write as $\langle \text{DM}_{\text{IGM}} \rangle \propto \Omega_{\text{b}}h^2/h$,

² https://github.com/FRBs/FRB

which is also a straightforward consequence of \bar{n}_e/H_0 . Recently, a constraint $\Omega_b h^2 = 0.02233 \pm 0.00036$ is obtained only based on the big bang nucleosynthesis theory with an improved rate of deuterium burning [82]. We use its mean value as default value on baryon density, and thus the observed DM can be used to measure H_0 .

Due to the existence of the cosmic web of filaments, voids, and other substructures, the values of DM_{cosmic} have variances along different sightlines. This variation is mainly affected by the galactic feedback, based on cosmological simulations [83]. The probability distribution of DM_{cosmic} can be described by [51, 84],

$$p_{\text{cosmic}}(\Delta) = A\Delta^{-\beta} \exp\left[-\frac{(\Delta^{-\alpha} - C_0)^2}{2\alpha^2 \sigma_{\text{DM}}^2}\right], \quad \Delta > 0,$$
(5)

where $\Delta \equiv \text{DM}_{\text{cosmic}}/\langle \text{DM}_{\text{cosmic}} \rangle$ and A is a normalization factor. The parameters $\alpha = 3$ and $\beta = 3$ provide the best match to the models [51]. The effective standard deviation equals approximately $Fz^{-0.5}$ for z < 1 where the parameter F describes the strength of the baryon feedback. The parameter C_0 can be fixed by the requirement $\langle \Delta \rangle = 1$. Combining Eqs. (2)–(5), we can obtain the probability distribution $p_{\text{cosmic}}(\text{DM}_{\text{cosmic}}|z, H_0, F)$.

The DM contribution from host galaxy includes host

galaxy halo and FRB's local environment. A log-normal distribution shows a fine fit to the results from the IllustrisTNG simulation [85], and its asymmetric long tail also can account for the contribution from the gas nearby FRB sources. Hence, DM_{host} can be modeled by a log-normal distribution,

$$p_{\rm host}(\rm DM'_{\rm host}) = \frac{1}{\rm DM'_{\rm host}} \frac{1}{\sigma_{\rm host}\sqrt{2\pi}} \exp\left[-\frac{(\ln \rm DM'_{\rm host} - \mu)^2}{2\sigma_{\rm host}^2}\right]$$
(6)

where DM'_{host} is the value of DM_{host} referenced to the rest frame of the host galaxy. The mean value and variance are e^{μ} and $e^{2\mu+\sigma^2_{\text{host}}}(e^{\sigma^2_{\text{host}}}-1)$, respectively. We apply a redshift correction $\text{DM}_{\text{host}} =$ $\text{DM}'_{\text{host}}/(1+z)$ and then acquire the probability distribution $p_{\text{host}}(\text{DM}_{\text{host}}|z,e^{\mu},\sigma_{\text{host}})$.

We rewrite Eq. (1) as

$$DM = DM_{MW,ISM} + DM', \tag{7}$$

where $DM' = DM_{MW,halo} + DM_{cosmic} + DM_{host} = DM_{MW,halo} + DM_E$ and $DM_E = DM_{cosmic} + DM_{host}$ is the extragalactic contribution. We ignore measurement error on DM and subtract $DM_{MW,ISM}$ from the observed DM value to obtain DM' for each FRB. Hence, the DM likelihood is

$$p(\mathrm{DM}'|z, H_0, e^{\mu}, \sigma_{\mathrm{host}}, F) = \int p_{\mathrm{halo}}(\mathrm{DM}_{\mathrm{MW,halo}}) \int_0^{\mathrm{DM}' - \mathrm{DM}_{\mathrm{MW,halo}}} p_{\mathrm{host}}(\mathrm{DM}_{\mathrm{host}}|z, e^{\mu}, \sigma_{\mathrm{host}})$$
$$p_{\mathrm{cosmic}}(\mathrm{DM}' - \mathrm{DM}_{\mathrm{MW,halo}} - \mathrm{DM}_{\mathrm{host}}|z, H_0, F) \, d\mathrm{DM}_{\mathrm{host}} \, d\mathrm{DM}_{\mathrm{MW,halo}}, \tag{8}$$

where p_{host} and p_{cosmic} can be calculated from Eqs. (2)–(6). The $p_{\text{halo}}(\text{DM}_{\text{MW,halo}})$ term is the model of $\text{DM}_{\text{MW,halo}}$. The simulation [86] showed that a representative halo electron contribution is about 30 pc cm⁻³. By using several observations, the value of $\text{DM}_{\text{MW,halo}}$ was estimated in the range of [50, 80] pc cm⁻³ [87]. According to these estimates, we use a Gaussian distribution $p_{\text{halo}}(\text{DM}_{\text{MW,halo}})$ with the mean value of 55 pc cm⁻³ and standard deviation of 25 pc cm⁻³ to describe the distribution of $\text{DM}_{\text{MW,halo}}$ for all FRBs [53]. In principle, all the unknown parameters, i.e. e^{μ} , σ_{host} , and F, should be treated as free parameters and be simultaneously constrained by data. However, the purpose of this work is to demonstrate the feasibility of using unlocalized FRBs and the real galaxy catalog to measure the Hubble constant. Thus, for convenience, we use the results from other works as the priors of the parameters e^{μ} , σ_{host} , and F. We first assume F = 0.31, which is a common assumption in the literature [55, 88], based on the constraint results in Refs. [51, 56].

For the FRB host galaxy parameters e^{μ} and σ_{host} , Zhang *et al.* [85] gave the best-fitting parameters $e^{\mu} \sim 32.97(1 + z)^{0.84} \text{ pc cm}^{-3}$ and $\sigma_{\text{host}} \sim 1.27$ for one-off FRBs, based on the state-of-the-art IllustrisTNG simulation, which can be used as priors in the simulation-based case. Using DM model from cosmological simulations to help constrain cosmological parameters is regarded as one of the approach to avoid uncontrolled systematic errors [89]. They also gave that the standard deviation of e^{μ} in the simulation is about 15 pc cm⁻³ [85], so we assume a Gaussian distribution $p_{\mu}(e^{\mu})$ on e^{μ} and integrate over it. Compared to Ref. [53], we further take the deviation of the simulation into account. Macquart *et al.* [51] also provided the median values of e^{μ} and σ_{host} around 68 pc cm⁻³ and 0.88, respectively, using localized FRBs while treating $\Omega_{\text{b}}h$ and F as free parameters. Similarly, the observation-based case assumes a Gaussian distribution $p_{\mu}(e^{\mu})$ on e^{μ} with a standard deviation 45 pc cm⁻³ [51]. We fix the values of σ_{host} in two cases. Then Eq. (8) becomes

$$p(\mathrm{DM}'|z, H_0) = \int_{30 \,\mathrm{pc} \,\mathrm{cm}^{-3}}^{80 \,\mathrm{pc} \,\mathrm{cm}^{-3}} p_{\mathrm{halo}}(\mathrm{DM}_{\mathrm{MW,halo}}) \int_{e_{\mathrm{min}}^{\mu}}^{e_{\mathrm{max}}^{\mu}} p_{\mu}(e^{\mu}) \int_{0}^{\mathrm{DM}' - \mathrm{DM}_{\mathrm{MW,halo}}} p_{\mathrm{host}}(\mathrm{DM}_{\mathrm{host}}|z, e^{\mu}) p_{\mathrm{cosmic}}(\mathrm{DM}' - \mathrm{DM}_{\mathrm{MW,halo}} - \mathrm{DM}_{\mathrm{host}}|z, H_0) \, d\mathrm{DM}_{\mathrm{host}} \, de^{\mu} \, d\mathrm{DM}_{\mathrm{MW,halo}},$$
(9)

where $e_{\text{max}}^{\mu} = 18(1+z)^{0.84} \text{ pc cm}^{-3}$ and $e_{\text{min}}^{\mu} = 48(1+z)^{0.84} \text{ pc cm}^{-3}$ in the simulation-based case [85] and $e_{\text{max}}^{\mu} = 23 \text{ pc cm}^{-3}$ and $e_{\text{min}}^{\mu} = 113 \text{ pc cm}^{-3}$ in the observation-based case [51]. This likelihood is normalized to all possible DM realizations. We again note that Eq. (9) used in this paper is an approximation of Eq. (8) and we just use it in order to facilitate the calculations.

B. Bayesian framework

We present an overview of the Bayesian framework to estimate H_0 using unlocalized FRBs. By using Bayes' theorem, the posterior of H_0 given the FRB data from a single detection, d_{FRB} , should be

$$p(H_0|\mathbf{d}_{\mathrm{FRB}}) \propto p(\mathbf{d}_{\mathrm{FRB}}|H_0)p(H_0),\tag{10}$$

where $p(d_{\text{FRB}}|H_0)$ is the FRB likelihood, and $p(H_0)$ is the prior on H_0 . We use the DM likelihood, Eq. (9), as a proxy of the FRB likelihood. By marginalizing over redshift and sky location Ω , the FRB likelihood can be calculated by

$$p(\mathbf{d}_{\mathrm{FRB}}|H_0) \propto \frac{\iint p(\mathrm{DM}'|z, H_0)p(z, \Omega) \, d\Omega \, dz}{\beta(H_0)},\tag{11}$$

where $\beta(H_0)$ is a normalization term, and the probability $p(z, \Omega)$ represents an FRB occurring at redshift z and sky location Ω . From the galaxy catalog data, we can select the galaxies within the angular localization errors of each FRB. The photometric redshift errors of the galaxy data are about 0.01 + 0.015z. But we ignore the errors and expect the final DESI data could provide precise spectroscopic redshift measurements. Then, the $p(z, \Omega)$ term can be written as the sum of the Dirac delta functions,

$$p(z,\Omega) = \sum_{i}^{N_{\text{gal}}} w_i \frac{R(z)}{1+z} \delta(z-z_i) \delta(\Omega-\Omega_i), \qquad (12)$$

where N_{gal} is the number of host galaxy candidates, (z_i, Ω_i) represent the redshift and sky location of the *i*-th host galaxy candidate, and w_i is the probability that the *i*-th galaxy hosts a FRB source. We consider two scenarios for this weight. The first one, called the "equal weight" scenario, assumes equal probability for each galaxy to host an FRB source, $w_i = 1/N_{\text{gal}}$, and the second one, called the "luminosity weight" scenario, assumes that this probability is proportional to the luminosity of this galaxy, $w_i \propto L_i$, meanwhile ensuring that the weights are normalized. The R(z) term represents the intrinsic redshift evolution of the FRB rate, but we assume that it is redshift-independent, R(z) = constant. This term may also be a function of the FRB luminosity function [90, 91], because the FRB luminosity could determine how far it can be detected and then affect the redshift distribution of the FRB rate. The factor 1/(1+z) converts the FRB rate from the source frame to the detector frame. Inserting Eq. (12) into Eq. (11), we have

$$p(\mathbf{d}_{\text{FRB}}|H_0) \propto \frac{\sum_{i=1}^{N_{\text{gal}}} w_i p(\mathrm{DM}'|z_i, H_0) R(z_i) / (1+z_i)}{\beta(H_0)}.$$
 (13)

The normalization term $\beta(H_0)$ accounts for selection effects to make sure the results are unbiased. Because we have followed the "gold-standard sample" criterion of FRBs [51] and then selected a part of the FRB data which are enough bright to avoid the selection effect induced by detection threshold, this procedure may discard the FRBs from higher redshift, which are more likely with higher DM_E value. So the used FRB data are below a cut-off value DM_{cutoff} (in this paper DM_{cutoff} = 570 pc cm⁻³). Integrating over all possible FRB realizations below DM_{cutoff} leads to

$$\beta(H_0) \propto \int \int_0^{\mathrm{DM}_{\mathrm{cutoff}}} p(\mathrm{DM}_{\mathrm{E}}|z, H_0) p(z) \, d\mathrm{DM}_{\mathrm{E}} \, dz$$
$$\propto \sum_{i=1}^{N_{\mathrm{gal}}} \int_0^{\mathrm{DM}_{\mathrm{cutoff}}} p(\mathrm{DM}_{\mathrm{E}}|z_i, H_0) w_i \frac{R(z_i)}{1+z_i} \, d\mathrm{DM}_{\mathrm{E}}, \tag{14}$$

where p(z) is identical to Eq. (12).

For multiple FRB events $\{d_{FRB,j}\}$, we assume that they are independent of each other and rewrite Eq. (10) as

$$p(H_0|\{\mathrm{d}_{\mathrm{FRB},j}\}) \propto p(H_0) \prod_j p(\mathrm{d}_{\mathrm{FRB},j}|H_0).$$

$$(15)$$

We take the prior on H_0 to be uniform over the range [20, 140] km s⁻¹ Mpc⁻¹. We use the Markov chain Monte Carlo method to sample the posterior and estimate constraints, which is implemented by the emcee package [92] with 32 walkers of 20000 samples after discarding 200 burn-in steps.

IV. RESULTS AND DISCUSSION

The core of the analysis is the probability $p(z_i|\text{DM}') \propto p(\text{DM}'|z_i)p(z_i)$ calculated at the redshift of the *i*-th potential host, z_i . As an example, the discrete probability $p(z_i|\text{DM}')$ in the "equal weight" scenario is shown in Fig. 2, with the assumption $\text{DM}_{\text{MW,halo}} = 50 \,\text{pc cm}^{-3}$, $H_0 = 70 \,\text{km s}^{-1} \,\text{Mpc}^{-1}$, $e^{\mu} = 50 \,\text{pc cm}^{-3}$, and $\sigma_{\text{host}} = 1$. It demonstrates that the galaxies above the maximum redshift of the galaxy catalog (z > 1) could almost contribute little to the total likelihood.

In Fig. 3, we show the constraints on H_0 assuming different host galaxy parameters from 12 unlocalized ASKAP FRBs in the "equal weight" scenario. The maximum a posteriori (MAP) values with the minimal 68.3% credible intervals are $H_0 = 58.7^{+19.8}_{-15.2}$ km s⁻¹ Mpc⁻¹ in the simulation-based case and $H_0 = 62.3^{+19.8}_{-16.6}$ km s⁻¹ Mpc⁻¹ in the observation-based case. Two estimates are basically consistent, and the difference can be regarded as a systematic error until we know more about the host galaxy parameters. The results of Planck and SH0ES are both within the 1 σ credible intervals of these two estimates. At current precision, our measurement cannot help resolve the Hubble tension.

It should be noted that actually, above $z \sim 0.5$, the galaxy catalog is incomplete [71], thus some high-redshift galaxies are absent in our analysis. This leads to the constraints lower compared to the reality (i.e., with a complete galaxy catalog), because of the positive correlation between z and H_0 [Eq. (2)]. The "luminosity weight" scenario is a reasonable way to reduce this effect. Because the absent high-redshift galaxies are with low luminosity, they would only contribute a little to the likelihood and final results. We should admit that the incompleteness of galaxy catalog needs a full analysis, which can refer to the calculations in the dark siren method [68], but this calculation is extremely time-consuming for FRBs due to the multiple integral in the DM likelihood, Eq. (9).

Figure 4 is the same as Fig. 3 but in the "luminosity weight" scenario. We can clearly see that the H_0 constraints are obviously increased in this scenario. The simulation-based and observation-based cases provide constraints, $H_0 = 80.4^{+24.1}_{-19.4} \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $H_0 = 81.9^{+23.4}_{-17.1} \text{ km s}^{-1} \text{ Mpc}^{-1}$, respectively. In Fig. 5, we plot the posterior distributions of H_0 for each FRB event in the simulation-based case and the "luminosity weight" scenario. We can see that although most of the single events only provide very weak constraints, the combined constraint can provide an informative constraint around 80 km s⁻¹ Mpc⁻¹.

We compare our H_0 constraints with the ones in the literature. The constraints, $H_0 = 62.3 \pm 9.1$ km s⁻¹ Mpc⁻¹ from 9 localized FRBs [52], $H_0 =$ $68.81^{+4.99}_{-4.33}~{\rm km\,s^{-1}\,Mpc^{-1}}$ from 18 localized FRBs [53], H_0 = 70.60 \pm 2.11 ${\rm km\,s^{-1}\,Mpc^{-1}}$ from a cosmologicalmodel-independent method [54], $H_0 = 69.4 \pm 4.7$ $\mathrm{km}\,\mathrm{s}^{-1}\,\mathrm{Mpc}^{-1}$ from 23 well-localized FRBs [57], and $H_0 = 65.5^{+6.4}_{-5.4}\,\mathrm{km}\,\mathrm{s}^{-1}\,\mathrm{Mpc}^{-1}$ from Supernova Ia combined with 18 localized FRBs [58] are all lower than our results in the "luminosity weight" scenario, but still consistent with our results at the 1σ credible level, due to the current loose constraints. Another work setting a flat prior on $DM_{MW,halo}$ provided $H_0 = 95.8^{+7.8}_{-9.2}$ $\rm km \, s^{-1} \, Mpc^{-1}$ [59], which is obviously higher than the estimates above but consistent with our results. The other two works [55, 56] using both localized and unlocalized FRBs gave $H_0 = 73^{+12}_{-8} \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $H_0 = 85.3^{+9.4}_{-8.1} \text{ km s}^{-1} \text{ Mpc}^{-1}$, respectively, which are closer to our MAP value, $80 \text{ km s}^{-1} \text{ Mpc}^{-1}$, of the "luminosity weight" constraint. Note that their analysis just marginalized the likelihood over a broad redshift range for unlocalized FRBs, but without the real galaxy information.

We also compare the relative errors of our results with the one using dark sirens in GW astronomy. 46 GW events combined with the GLADE+ galaxy catalog data could give about 19% precision constraint on H_0 [93], yet our method achieves about 27% precision constraint using 12 FRB events. By approximating a Gaussian distribution and scaling our result as $1/\sqrt{N}$ behaviour, where N is the number of FRB events, we see that our method can provide similar constraint ability to the dark siren method.

The 12 FRB events account for about a half of the total FRB events observed by ASKAP FE mode, because the catalog footprint covers about a half of the total sky. Applying this proportion to the total current FRB data, we would have about 400 events to perform this analysis. Again scaling our result as $1/\sqrt{N}$ behaviour, the constraint on H_0 is able to reach 8% precision using those 400 unlocalized FRBs. However, at that point, the impacts of the host galaxy parameters, the choice of priors and cosmological models, and other systematic errors would become more important and should be carefully treated. We will study the systematic errors of this method in the next paper.

We expect that this method can be further developed to measure other cosmological parameters and FRB parameters [94], benefiting by the high detection rates of FRBs. Actually, even for some of the well-localized FRBs (such as being localized to arcsecond), it is still possible more than one single obvious host galaxy in the FRB localization region. Therefore, cosmological parameter estimations still have to be marginalized over multiple potential hosts for these FRBs.



FIG. 2. The discrete probabilities $p(\{z_i\}|\text{DM}')$ calculated at the redshifts of potential hosts, $\{z_i\}$, for each FRB event in the "equal weight" scenario. The DM' value of FRB events (from low to high) are indicated by the colors of lines (from purple to yellow). Note that we assume $\text{DM}_{\text{MW,halo}} = 50 \,\text{pc}\,\text{cm}^{-3}$, $H_0 = 70 \,\text{km}\,\text{s}^{-1}\,\text{Mpc}^{-1}$, $e^{\mu} = 50 \,\text{pc}\,\text{cm}^{-3}$, and $\sigma_{\text{host}} = 1$ as an example.



FIG. 3. The posteriors for H_0 from 12 unlocalized ASKAP FRBs in the "equal weight" scenario. The constraints for the simulation-based case and the observation-based case are shown in blue and red solid lines, respectively, and their 1σ credible intervals are shown in vertical dotted lines. The constraints from Planck [1] and SH0ES [2] are also shown in orange and grey regions, respectively.

V. CONCLUSIONS

In this paper, we use the statistical galaxy catalog method similar to the dark siren method in GW cosmology and obtain the measurement of the Hubble constant H_0 using twelve unlocalized ASKAP FRB data as an example. We find all potential host galaxies in the FRB localization region and then marginalize the FRB likelihood over them. For the weights on the probability for a galaxy to host an FRB source, two different scenarios are considered, i.e. the "equal weight" scenario and the "luminosity weight" scenario. We also use the results from the IllustrisTNG simulation and the constraints from the localized FRBs as two priors of the DM_{host}



FIG. 4. The posteriors for H_0 from 12 unlocalized ASKAP FRBs in the "luminosity weight" scenario. The constraints for the simulation-based case and the observation-based case are shown in blue and red solid lines, respectively, and their 1σ credible intervals are shown in vertical dotted lines. The constraints from Planck [1] and SH0ES [2] are also shown in orange and grey regions, respectively.



FIG. 5. The posteriors for H_0 from each unlocalized ASKAP FRB. The constraints for individual FRB are shown in dotted lines, and the combined constraint is shown in solid line.

model, denoted by the simulation-based and observationbased cases, respectively. In the "equal weight" scenario, the constraints are $H_0 = 58.7^{+19.8}_{-15.2} \text{ km s}^{-1} \text{ Mpc}^{-1}$ in the simulation-based case and $H_0 = 62.3^{+19.8}_{-16.6} \text{ km s}^{-1} \text{ Mpc}^{-1}$ in the observation-based case, combined with the BBN result on $\Omega_{\rm b}h^2$. To reduce the effect of the galaxy catalog incompleteness, we obtain the constraints in the "luminosity weight" scenario, $H_0 = 80.4^{+24.1}_{-19.4} \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $H_0 = 81.9^{+23.4}_{-17.1} \text{ km s}^{-1} \text{ Mpc}^{-1}$, in the simulation-based and observation-based cases, respectively. These results are very rough estimates, but serve to demonstrate the feasibility of this method.

We should point out one disadvantage of this work that we used the priors for the host galaxy parameters, i.e. e^{μ} and σ_{host} . In principle, they should be regarded as free parameters. The results obtained by fixing these parameters may be biased [55]. Actually, this problem also exists in early study of the dark siren method in GW cosmology, in which the underlying source mass distribution is fixed, and the systematic errors are studied by a series following papers. Nonetheless, the deviation between the cases of different host galaxy parameters could be regarded as a systematic error, until more accurate information about FRB hosts is known. It should be noted that other emerging H_0 measurements (such as standard sirens [95–116]) also have different systematic errors, e.g., systematic errors of bright sirens from the viewing angle of binary neutron stars [117], of dark sirens from the assumed BBH mass distribution [93], of localized FRBs from tentative host galaxy [51, 118]. So it is still im-

- N. Aghanim *et al.* (Planck), Planck 2018 results. VI. Cosmological parameters, Astron. Astrophys. **641**, A6 (2020), [Erratum: Astron.Astrophys. 652, C4 (2021)], arXiv:1807.06209 [astro-ph.CO].
- [2] A. G. Riess *et al.*, A Comprehensive Measurement of the Local Value of the Hubble Constant with 1 km s⁻¹ Mpc⁻¹ Uncertainty from the Hubble Space Telescope and the SH0ES Team, Astrophys. J. Lett. **934**, L7 (2022), arXiv:2112.04510 [astro-ph.CO].
- [3] R.-Y. Guo, J.-F. Zhang, and X. Zhang, Can the H₀ tension be resolved in extensions to ΛCDM cosmology?, JCAP 02, 054, arXiv:1809.02340 [astro-ph.CO].
- [4] R.-G. Cai, Z.-K. Guo, S.-J. Wang, W.-W. Yu, and Y. Zhou, No-go guide for the Hubble tension: Latetime solutions, Phys. Rev. D 105, L021301 (2022), arXiv:2107.13286 [astro-ph.CO].
- [5] E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri, D. F. Mota, A. G. Riess, and J. Silk, In the realm of the Hubble tension—a review of solutions, Class. Quant. Grav. 38, 153001 (2021), arXiv:2103.01183 [astro-ph.CO].
- [6] E. Abdalla *et al.*, Cosmology intertwined: A review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies, JHEAp **34**, 49 (2022), arXiv:2203.06142 [astro-ph.CO].
- [7] J.-P. Hu and F.-Y. Wang, Hubble Tension: The Evidence of New Physics, Universe 9, 94 (2023), arXiv:2302.05709 [astro-ph.CO].
- [8] S. Bhandari and C. Flynn, Probing the Universe with Fast Radio Bursts, Universe 7, 85 (2021).
- [9] D. Xiao, F. Wang, and Z. Dai, The physics of fast radio bursts, Sci. China Phys. Mech. Astron. 64, 249501 (2021), arXiv:2101.04907 [astro-ph.HE].
- [10] L. Nicastro, C. Guidorzi, E. Palazzi, L. Zampieri, M. Turatto, and A. Gardini, Multiwavelength observations of Fast Radio Bursts, Universe 7, 76 (2021), arXiv:2103.07786 [astro-ph.HE].
- [11] E. Petroff, J. W. T. Hessels, and D. R. Lorimer, Fast radio bursts at the dawn of the 2020s, Astron. Astrophys.

portant to develop new H_0 measurements to provide a cross-check for each other.

ACKNOWLEDGMENTS

We are very grateful to Jia-Ming Zou, J. X. Prochaska, Zheng-Xiang Li, and Yougang Wang for fruitful discussions. This work was supported by the National SKA Program of China (Grants Nos. 2022SKA0110200 and 2022SKA0110203), the National Natural Science Foundation of China (Grants Nos. 11975072, 11875102, and 11835009), the science research grants from the China Manned Space Project (Grant No. CMS-CSST-2021-B01), the Liaoning Revitalization Talents Program (Grant No. XLYC1905011), the National Program for Support of Top-Notch Young Professionals (Grant No. W02070050), and the National 111 Project of China (Grant No. B16009).

Rev. 30, 2 (2022), arXiv:2107.10113 [astro-ph.HE].

- [12] M. Caleb and E. Keane, A Decade and a Half of Fast Radio Burst Observations, Universe 7, 453 (2021).
- [13] B. Zhang, The physics of fast radio bursts, Rev. Mod. Phys. 95, 035005 (2023), arXiv:2212.03972 [astroph.HE].
- [14] M. Bailes, The discovery and scientific potential of fast radio bursts, Science 378, abj3043 (2022), arXiv:2211.06048 [astro-ph.HE].
- [15] M. Amiri *et al.* (CHIME/FRB), The First CHIME/FRB Fast Radio Burst Catalog, Astrophys. J. Supp. **257**, 59 (2021), arXiv:2106.04352 [astro-ph.HE].
- [16] C. D. Bochenek, V. Ravi, K. V. Belov, G. Hallinan, J. Kocz, S. R. Kulkarni, and D. L. McKenna, A fast radio burst associated with a Galactic magnetar, Nature 587, 59 (2020), arXiv:2005.10828 [astro-ph.HE].
- [17] B. C. Andersen *et al.* (CHIME/FRB), A bright millisecond-duration radio burst from a Galactic magnetar, Nature 587, 54 (2020), arXiv:2005.10324 [astroph.HE].
- [18] S. Chatterjee *et al.*, The direct localization of a fast radio burst and its host, Nature **541**, 58 (2017), arXiv:1701.01098 [astro-ph.HE].
- [19] K. W. t. Bannister, A single fast radio burst localized to a massive galaxy at cosmological distance, Science 365, 565 (2019), arXiv:1906.11476 [astro-ph.HE].
- [20] C. H. Niu *et al.*, A repeating fast radio burst associated with a persistent radio source, Nature **606**, 873 (2022), [Erratum: Nature 611, E10 (2022)], arXiv:2110.07418 [astro-ph.HE].
- [21] J.-S. Wang, Y.-P. Yang, X.-F. Wu, Z.-G. Dai, and F.-Y. Wang, Fast Radio Bursts from the Inspiral of Double Neutron Stars, Astrophys. J. Lett. 822, L7 (2016), arXiv:1603.02014 [astro-ph.HE].
- [22] Y. Lyubarsky, Emission Mechanisms of Fast Radio Bursts, Universe 7, 56 (2021), arXiv:2103.00470 [astroph.HE].
- [23] J.-J. Geng, B. Li, and Y.-F. Huang, Repeating Fast Ra-

dio Bursts from Collapses of the Crust of a Strange Star, The Innovation **2**, 100152 (2021), arXiv:2103.04165 [astro-ph.HE].

- [24] W. Deng and B. Zhang, Cosmological Implications of Fast Radio Burst/Gamma-Ray Burst Associations, Astrophys. J. Lett. 783, L35 (2014), arXiv:1401.0059 [astro-ph.HE].
- [25] B. Zhou, X. Li, T. Wang, Y.-Z. Fan, and D.-M. Wei, Fast radio bursts as a cosmic probe?, Phys. Rev. D 89, 107303 (2014), arXiv:1401.2927 [astro-ph.CO].
- [26] H. Gao, Z. Li, and B. Zhang, Fast Radio Burst/Gamma-Ray Burst Cosmography, Astrophys. J. 788, 189 (2014), arXiv:1402.2498 [astro-ph.CO].
- [27] Y.-P. Yang and B. Zhang, Extracting host galaxy dispersion measure and constraining cosmological parameters using fast radio burst data, Astrophys. J. Lett. 830, L31 (2016), arXiv:1608.08154 [astro-ph.HE].
- [28] Z.-X. Li, H. Gao, X.-H. Ding, G.-J. Wang, and B. Zhang, Strongly lensed repeating fast radio bursts as precision probes of the universe, Nature Commun. 9, 3833 (2018), arXiv:1708.06357 [astro-ph.CO].
- [29] A. Walters, A. Weltman, B. M. Gaensler, Y.-Z. Ma, and A. Witzemann, Future Cosmological Constraints from Fast Radio Bursts, Astrophys. J. 856, 65 (2018), arXiv:1711.11277 [astro-ph.CO].
- [30] J.-J. Wei, X.-F. Wu, and H. Gao, Cosmology with Gravitational Wave/Fast Radio Burst Associations, Astrophys. J. Lett. 860, L7 (2018), arXiv:1805.12265 [astroph.CO].
- [31] B. Liu, Z. Li, H. Gao, and Z.-H. Zhu, Prospects of strongly lensed repeating fast radio bursts: Complementary constraints on dark energy evolution, Phys. Rev. D 99, 123517 (2019), arXiv:1907.10488 [astro-ph.CO].
- [32] L. Zhang and Z. Li, Combinations of Standard Pings and Standard Candles: An Effective and Hubble Constant-free Probe of Dark Energy Evolution, Astrophys. J. 901, 130 (2020).
- [33] Z.-W. Zhao, Z.-X. Li, J.-Z. Qi, H. Gao, J.-F. Zhang, and X. Zhang, Cosmological parameter estimation for dynamical dark energy models with future fast radio burst observations, Astrophys. J. 903, 83 (2020), arXiv:2006.01450 [astro-ph.CO].
- [34] X.-W. Qiu, Z.-W. Zhao, L.-F. Wang, J.-F. Zhang, and X. Zhang, A forecast of using fast radio burst observations to constrain holographic dark energy, JCAP 02 (02), 006, arXiv:2108.04127 [astro-ph.CO].
- [35] D.-C. Qiang and H. Wei, Effect of Redshift Distributions of Fast Radio Bursts on Cosmological Constraints, Phys. Rev. D 103, 083536 (2021), arXiv:2102.00579 [astro-ph.CO].
- [36] J.-P. Dai and J.-Q. Xia, Reconstruction of baryon fraction in intergalactic medium through dispersion measurements of fast radio bursts, Mon. Not. Roy. Astron. Soc. 503, 4576 (2021), arXiv:2103.08479 [astro-ph.CO].
- [37] S. Zhao, B. Liu, Z. Li, and H. Gao, Model-independent Estimation of H_0 and Ω_K from Strongly Lensed Fast Radio Bursts, Astrophys. J. **916**, 70 (2021).
- [38] C. Zhu and J. Zhang, Improvement of cosmological constraints with the cross-correlation between line-of-sight optical galaxy and FRB dispersion measures, Phys. Rev. D 106, 023513 (2022), arXiv:2205.03867 [astro-ph.CO].
- [39] Z.-W. Zhao, L.-F. Wang, J.-G. Zhang, J.-F. Zhang, and X. Zhang, Probing the interaction between dark energy and dark matter with future fast radio burst observa-

tions, JCAP 04, 022, arXiv:2210.07162 [astro-ph.CO].

- [40] P.-J. Wu, Y. Shao, S.-J. Jin, and X. Zhang, A path to precision cosmology: synergy between four promising late-universe cosmological probes, JCAP 06, 052, arXiv:2202.09726 [astro-ph.CO].
- [41] K. B. Yang, Q. Wu, and F. Y. Wang, Finding the Missing Baryons in the Intergalactic Medium with Localized Fast Radio Bursts, Astrophys. J. Lett. 940, L29 (2022), arXiv:2211.04058 [astro-ph.HE].
- [42] J.-G. Zhang, Z.-W. Zhao, Y. Li, J.-F. Zhang, D. Li, and X. Zhang, Cosmology with fast radio bursts in the era of SKA, Sci. China Phys. Mech. Astron. 66, 120412 (2023), arXiv:2307.01605 [astro-ph.CO].
- [43] Z. Li, H. Gao, J.-J. Wei, Y.-P. Yang, B. Zhang, and Z.-H. Zhu, Cosmology-independent estimate of the fraction of baryon mass in the IGM from fast radio burst observations, Astrophys. J. 876, 146 (2019), arXiv:1904.08927 [astro-ph.CO].
- [44] R.-G. Cai, T.-B. Liu, S.-J. Wang, and W.-T. Xu, Probing cosmic anisotropy with GW/FRB as upgraded standard sirens, JCAP 09, 016, arXiv:1905.01803 [astroph.CO].
- [45] J.-J. Wei, Z. Li, H. Gao, and X.-F. Wu, Constraining the Evolution of the Baryon Fraction in the IGM with FRB and H(z) data, JCAP 09, 039, arXiv:1907.09772 [astro-ph.CO].
- [46] Q. Wu, H. Yu, and F. Y. Wang, A New Method to Measure Hubble Parameter H(z) using Fast Radio Bursts, Astrophys. J. 895, 33 (2020), arXiv:2004.12649 [astro-ph.CO].
- [47] K.-G. Lee, M. Ata, I. S. Khrykin, Y. Huang, J. X. Prochaska, J. Cooke, J. Zhang, and A. Batten, Constraining the Cosmic Baryon Distribution with Fast Radio Burst Foreground Mapping, Astrophys. J. 928, 9 (2022), arXiv:2109.00386 [astro-ph.CO].
- [48] R. Gao, Z. Li, and H. Gao, Prospects of strongly lensed fast radio bursts: simultaneous measurement of post-Newtonian parameter and Hubble constant, Mon. Not. Roy. Astron. Soc. **516**, 1977 (2022), arXiv:2208.10175 [astro-ph.CO].
- [49] R. Reischke and S. Hagstotz, Consistent constraints on the equivalence principle from localized fast radio bursts, Mon. Not. Roy. Astron. Soc. **523**, 6264 (2023), arXiv:2302.10072 [astro-ph.CO].
- [50] B. Wang, J.-J. Wei, X.-F. Wu, and M. López-Corredoira, Revisiting constraints on the photon rest mass with cosmological fast radio bursts, JCAP 09, 025, arXiv:2304.14784 [astro-ph.HE].
- [51] J. P. Macquart *et al.*, A census of baryons in the Universe from localized fast radio bursts, Nature **581**, 391 (2020), arXiv:2005.13161 [astro-ph.CO].
- [52] S. Hagstotz, R. Reischke, and R. Lilow, A new measurement of the Hubble constant using fast radio bursts, Mon. Not. Roy. Astron. Soc. 511, 662 (2022), arXiv:2104.04538 [astro-ph.CO].
- [53] Q. Wu, G.-Q. Zhang, and F.-Y. Wang, An 8 per cent determination of the Hubble constant from localized fast radio bursts, Mon. Not. Roy. Astron. Soc. 515, L1 (2022), arXiv:2108.00581 [astro-ph.CO].
- [54] Y. Liu, H. Yu, and P. Wu, Cosmological-modelindependent Determination of Hubble Constant from Fast Radio Bursts and Hubble Parameter Measurements, Astrophys. J. Lett. **946**, L49 (2023), arXiv:2210.05202 [astro-ph.CO].

- [55] C. W. James *et al.*, A measurement of Hubble's Constant using Fast Radio Bursts, Mon. Not. Roy. Astron. Soc. **516**, 4862 (2022), arXiv:2208.00819 [astro-ph.CO].
- [56] J. Baptista, J. X. Prochaska, A. G. Mannings, C. W. James, R. M. Shannon, S. D. Ryder, A. T. Deller, D. R. Scott, M. Glowacki, and N. Tejos, Measuring the Variance of the Macquart Relation in Redshift–Extragalactic Dispersion Measure Modeling, Astrophys. J. 965, 57 (2024), arXiv:2305.07022 [astro-ph.CO].
- [57] J. A. S. Fortunato, W. S. Hipólito-Ricaldi, and M. V. dos Santos, Cosmography from well-localized fast radio bursts, Mon. Not. Roy. Astron. Soc. **526**, 1773 (2023), arXiv:2307.04711 [astro-ph.CO].
- [58] J. Gao, Z. Zhou, M. Du, R. Zou, J. Hu, and L. Xu, A Measurement of Hubble Constant Using Cosmographic Approach from Fast Radio Bursts and SNe Ia, (2023), arXiv:2307.08285 [astro-ph.CO].
- [59] J.-J. Wei and F. Melia, Investigating Cosmological Models and the Hubble Tension Using Localized Fast Radio Bursts, Astrophys. J. 955, 101 (2023), arXiv:2308.05918 [astro-ph.CO].
- [60] S. Paine, T. Hawkins, D. R. Lorimer, J. Stanley, J. Kania, F. Crawford, and N. Fairfield, A search for millisecond radio bursts from Messier 82, Mon. Not. Roy. Astron. Soc. **528**, 6340 (2024), arXiv:2401.17242 [astroph.HE].
- [61] K. Aggarwal, T. Budavári, A. T. Deller, T. Eftekhari, C. W. James, J. X. Prochaska, and S. P. Tendulkar (FRB Associations Team), Probabilistic Association of Transients to their Hosts (PATH), Astrophys. J. 911, 95 (2021), arXiv:2102.10627 [astro-ph.HE].
- [62] W. Del Pozzo, Inference of the cosmological parameters from gravitational waves: application to second generation interferometers, Phys. Rev. D 86, 043011 (2012), arXiv:1108.1317 [astro-ph.CO].
- [63] H.-Y. Chen, M. Fishbach, and D. E. Holz, A two per cent Hubble constant measurement from standard sirens within five years, Nature 562, 545 (2018), arXiv:1712.06531 [astro-ph.CO].
- [64] M. Soares-Santos et al. (DES, LIGO Scientific, Virgo), First Measurement of the Hubble Constant from a Dark Standard Siren using the Dark Energy Survey Galaxies and the LIGO/Virgo Binary–Black-hole Merger GW170814, Astrophys. J. Lett. 876, L7 (2019), arXiv:1901.01540 [astro-ph.CO].
- [65] B. P. Abbott *et al.* (LIGO Scientific, Virgo, VIRGO), A Gravitational-wave Measurement of the Hubble Constant Following the Second Observing Run of Advanced LIGO and Virgo, Astrophys. J. **909**, 218 (2021), [Erratum: Astrophys.J. 923, 279 (2021)], arXiv:1908.06060 [astro-ph.CO].
- [66] A. Palmese, C. R. Bom, S. Mucesh, and W. G. Hartley, A Standard Siren Measurement of the Hubble Constant Using Gravitational-wave Events from the First Three LIGO/Virgo Observing Runs and the DESI Legacy Survey, Astrophys. J. 943, 56 (2023), arXiv:2111.06445 [astro-ph.CO].
- [67] J.-Y. Song, L.-F. Wang, Y. Li, Z.-W. Zhao, J.-F. Zhang, W. Zhao, and X. Zhang, Synergy between CSST galaxy survey and gravitational-wave observation: Inferring the Hubble constant from dark standard sirens, Sci. China Phys. Mech. Astron. 67, 230411 (2024), arXiv:2212.00531 [astro-ph.CO].

- [68] R. Gray *et al.*, Cosmological inference using gravitational wave standard sirens: A mock data analysis, Phys. Rev. D **101**, 122001 (2020), arXiv:1908.06050 [grqc].
- [69] A. Dey et al. (DESI), Overview of the DESI Legacy Imaging Surveys, Astron. J. 157, 168 (2019), arXiv:1804.08657 [astro-ph.IM].
- [70] E. L. Wright *et al.*, The Wide-field Infrared Survey Explorer (WISE): Mission Description and Initial On-orbit Performance, Astron. J. **140**, 1868 (2010), arXiv:1008.0031 [astro-ph.IM].
- [71] X. Yang *et al.*, An Extended Halo-based Group/Cluster finder: application to the DESI legacy imaging surveys DR8, Astrophys. J. **909**, 143 (2021), arXiv:2012.14998 [astro-ph.GA].
- [72] S. Johnston and J. Wall, Science with ASKAP the Australian Square Kilometre Array Pathfinder, Exper. Astron. 22, 151 (2008), arXiv:0810.5187 [astro-ph].
- [73] J.-P. Macquart *et al.* (CRAFT), The Commensal Realtime ASKAP Fast Transients (CRAFT) survey, Publ. Astron. Soc. Austral. **27**, 272 (2010), arXiv:1001.2958 [astro-ph.HE].
- [74] K. Bannister *et al.*, The detection of an extremely bright fast radio burst in a phased array feed survey, Astrophys. J. Lett. **841**, L12 (2017), arXiv:1705.07581 [astroph.HE].
- [75] R. M. t. Shannon, The dispersion-brightness relation for fast radio bursts from a wide-field survey, Nature 562, 386 (2018).
- [76] J. P. Macquart, R. M. Shannon, K. W. Bannister, C. W. James, R. D. Ekers, and J. D. Bunton, The Spectral Properties of the Bright Fast Radio Burst Population, Astrophys. J. Lett. 872, L19 (2019), arXiv:1810.04353 [astro-ph.HE].
- [77] S. Bhandari, K. W. Bannister, C. W. James, R. M. Shannon, C. M. Flynn, M. Caleb, and J. D. Bunton, A Southern sky search for repeating Fast Radio Bursts using the Australian SKA Pathfinder, Mon. Not. Roy. Astron. Soc. 486, 70 (2019), arXiv:1903.06525 [astroph.HE].
- [78] J. M. Cordes and T. J. W. Lazio, NE2001. 1. A New model for the galactic distribution of free electrons and its fluctuations, (2002), arXiv:astro-ph/0207156.
- [79] K. Ioka, Cosmic dispersion measure from gamma-ray burst afterglows: probing the reionization history and the burst environment, Astrophys. J. Lett. 598, L79 (2003), arXiv:astro-ph/0309200.
- [80] S. Inoue, Probing the cosmic reionization history and local environment of gamma-ray bursts through radio dispersion, Mon. Not. Roy. Astron. Soc. **348**, 999 (2004), arXiv:astro-ph/0309364.
- [81] X.-H. Fan, C. L. Carilli, and B. G. Keating, Observational constraints on cosmic reionization, Ann. Rev. Astron. Astrophys. 44, 415 (2006), arXiv:astroph/0602375.
- [82] V. Mossa *et al.*, The baryon density of the Universe from an improved rate of deuterium burning, Nature 587, 210 (2020).
- [83] M. McQuinn, Locating the "missing" baryons with extragalactic dispersion measure estimates, Astrophys. J. Lett. 780, L33 (2014), arXiv:1309.4451 [astro-ph.CO].
- [84] J. Miralda-Escude, M. Haehnelt, and M. J. Rees, Reionization of the inhomogeneous universe, Astrophys. J. 530, 1 (2000), arXiv:astro-ph/9812306.

- [85] G. Q. Zhang, H. Yu, J. H. He, and F. Y. Wang, Dispersion measures of fast radio burst host galaxies derived from IllustrisTNG simulation, Astrophys. J. 900, 170 (2020), arXiv:2007.13935 [astro-ph.HE].
- [86] K. Dolag, B. M. Gaensler, A. M. Beck, and M. C. Beck, Constraints on the distribution and energetics of fast radio bursts using cosmological hydrodynamic simulations, Mon. Not. Roy. Astron. Soc. 451, 4277 (2015), arXiv:1412.4829 [astro-ph.CO].
- [87] J. X. Prochaska and Y. Zheng, Probing galactic haloes with fast radio bursts, Mon. Not. Roy. Astron. Soc. 485, 648 (2019).
- [88] C. W. James, J. X. Prochaska, J. P. Macquart, F. O. North-Hickey, K. W. Bannister, and A. Dunning, The z–DM distribution of fast radio bursts, Mon. Not. Roy. Astron. Soc. 509, 4775 (2021), arXiv:2101.08005 [astroph.HE].
- [89] M. Hollis, Closing in on the Hubble constant, Nat. Astron. 6, 770 (2022).
- [90] J.-P. Macquart and R. Ekers, FRB event rate counts II. Fluence, redshift, and dispersion measure distributions, Mon. Not. Roy. Astron. Soc. 480, 4211 (2018), arXiv:1808.00908 [astro-ph.HE].
- [91] L. Li, Y. Huang, Z. Zhang, D. Li, and B. Li, Intensity Distribution Function and Statistical Properties of Fast Radio Bursts, Res. Astron. Astrophys. 17, 6 (2017), arXiv:1602.06099 [astro-ph.HE].
- [92] D. Foreman-Mackey, D. W. Hogg, D. Lang, and J. Goodman, emcee: The MCMC Hammer, Publ. Astron. Soc. Pac. **125**, 306 (2013), arXiv:1202.3665 [astroph.IM].
- [93] R. Abbott *et al.* (LIGO Scientific, Virgo, KA-GRA), Constraints on the Cosmic Expansion History from GWTC-3, Astrophys. J. **949**, 76 (2023), arXiv:2111.03604 [astro-ph.CO].
- [94] Y. Li, J.-M. Zou, J.-G. Zhang, Z.-W. Zhao, J.-F. Zhang, and X. Zhang, Fast radio burst energy function in the presence of DM_{host} variation, (2023), arXiv:2303.16775 [astro-ph.HE].
- [95] B. F. Schutz, Determining the Hubble Constant from Gravitational Wave Observations, Nature **323**, 310 (1986).
- [96] D. E. Holz and S. A. Hughes, Using gravitationalwave standard sirens, Astrophys. J. 629, 15 (2005), arXiv:astro-ph/0504616.
- [97] B. P. Abbott *et al.* (LIGO Scientific, Virgo, 1M2H, Dark Energy Camera GW-E, DES, DLT40, Las Cumbres Observatory, VINROUGE, MASTER), A gravitationalwave standard siren measurement of the Hubble constant, Nature 551, 85 (2017), arXiv:1710.05835 [astroph.CO].
- [98] B. P. Abbott *et al.* (LIGO Scientific, Virgo), GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral, Phys. Rev. Lett. **119**, 161101 (2017), arXiv:1710.05832 [gr-qc].
- [99] B. P. Abbott et al. (LIGO Scientific, Virgo, Fermi GBM, INTEGRAL, IceCube, AstroSat Cadmium Zinc Telluride Imager Team, IPN, Insight-Hxmt, ANTARES, Swift, AGILE Team, 1M2H Team, Dark Energy Camera GW-EM, DES, DLT40, GRAWITA, Fermi-LAT, ATCA, ASKAP, Las Cumbres Observatory Group, OzGrav, DWF (Deeper Wider Faster Program), AST3, CAASTRO, VINROUGE, MAS-TER, J-GEM, GROWTH, JAGWAR, CaltechNRAO,

TTU-NRAO, NuSTAR, Pan-STARRS, MAXI Team, TZAC Consortium, KU, Nordic Optical Telescope, ePESSTO, GROND, Texas Tech University, SALT Group, TOROS, BOOTES, MWA, CALET, IKI-GW Follow-up, H.E.S.S., LOFAR, LWA, HAWC, Pierre Auger, ALMA, Euro VLBI Team, Pi of Sky, Chandra Team at McGill University, DFN, AT-LAS Telescopes, High Time Resolution Universe Survey, RIMAS, RATIR, SKA South Africa/MeerKAT), Multi-messenger Observations of a Binary Neutron Star Merger, Astrophys. J. Lett. 848, L12 (2017), arXiv:1710.05833 [astro-ph.HE].

- [100] W. Zhao, C. Van Den Broeck, D. Baskaran, and T. G. F. Li, Determination of Dark Energy by the Einstein Telescope: Comparing with CMB, BAO and SNIa Observations, Phys. Rev. D 83, 023005 (2011), arXiv:1009.0206 [astro-ph.CO].
- [101] S. Vitale and H.-Y. Chen, Measuring the Hubble constant with neutron star black hole mergers, Phys. Rev. Lett. **121**, 021303 (2018), arXiv:1804.07337 [astroph.CO].
- [102] L.-F. Wang, X.-N. Zhang, J.-F. Zhang, and X. Zhang, Impacts of gravitational-wave standard siren observation of the Einstein Telescope on weighing neutrinos in cosmology, Phys. Lett. B 782, 87 (2018), arXiv:1802.04720 [astro-ph.CO].
- [103] S.-J. Jin, D.-Z. He, Y. Xu, J.-F. Zhang, and X. Zhang, Forecast for cosmological parameter estimation with gravitational-wave standard siren observation from the Cosmic Explorer, JCAP 03, 051, arXiv:2001.05393 [astro-ph.CO].
- [104] E. Belgacem, Y. Dirian, S. Foffa, E. J. Howell, M. Maggiore, and T. Regimbau, Cosmology and dark energy from joint gravitational wave-GRB observations, JCAP 08, 015, arXiv:1907.01487 [astro-ph.CO].
- [105] C. Howlett and T. M. Davis, Standard siren speeds: improving velocities in gravitational-wave measurements of H₀, Mon. Not. Roy. Astron. Soc. **492**, 3803 (2020), arXiv:1909.00587 [astro-ph.CO].
- [106] J.-F. Zhang, M. Zhang, S.-J. Jin, J.-Z. Qi, and X. Zhang, Cosmological parameter estimation with future gravitational wave standard siren observation from the Einstein Telescope, JCAP 09, 068, arXiv:1907.03238 [astro-ph.CO].
- [107] J. M. Ezquiaga and D. E. Holz, Spectral Sirens: Cosmology from the Full Mass Distribution of Compact Binaries, Phys. Rev. Lett. **129**, 061102 (2022), arXiv:2202.08240 [astro-ph.CO].
- [108] L.-F. Wang, Z.-W. Zhao, J.-F. Zhang, and X. Zhang, A preliminary forecast for cosmological parameter estimation with gravitational-wave standard sirens from Tian-Qin, JCAP 11, 012, arXiv:1907.01838 [astro-ph.CO].
- [109] Z.-W. Zhao, L.-F. Wang, J.-F. Zhang, and X. Zhang, Prospects for improving cosmological parameter estimation with gravitational-wave standard sirens from Taiji, Sci. Bull. 65, 1340 (2020), arXiv:1912.11629 [astroph.CO].
- [110] L.-F. Wang, S.-J. Jin, J.-F. Zhang, and X. Zhang, Forecast for cosmological parameter estimation with gravitational-wave standard sirens from the LISA-Taiji network, Sci. China Phys. Mech. Astron. 65, 210411 (2022), arXiv:2101.11882 [gr-qc].
- [111] S.-J. Jin, L.-F. Wang, P.-J. Wu, J.-F. Zhang, and X. Zhang, How can gravitational-wave standard sirens

and 21-cm intensity mapping jointly provide a precise late-universe cosmological probe?, Phys. Rev. D **104**, 103507 (2021), arXiv:2106.01859 [astro-ph.CO].

- [112] J. Yu, H. Song, S. Ai, H. Gao, F. Wang, Y. Wang, Y. Lu, W. Fang, and W. Zhao, Multimessenger Detection Rates and Distributions of Binary Neutron Star Mergers and Their Cosmological Implications, Astrophys. J. 916, 54 (2021), arXiv:2104.12374 [astro-ph.HE].
- [113] S.-J. Jin, Y.-Z. Zhang, J.-Y. Song, J.-F. Zhang, and X. Zhang, Taiji-TianQin-LISA network: Precisely measuring the Hubble constant using both bright and dark sirens, Sci. China Phys. Mech. Astron. 67, 220412 (2024), arXiv:2305.19714 [astro-ph.CO].
- [114] L.-G. Zhu and X. Chen, The Dark Side of Using Dark Sirens to Constrain the Hubble–Lemaître Constant, Astrophys. J. 948, 26 (2023), arXiv:2302.10621 [astroph.CO].

- [115] T. Han, S.-J. Jin, J.-F. Zhang, and X. Zhang, A comprehensive forecast for cosmological parameter estimation using joint observations of gravitational-wave standard sirens and short γ-ray bursts, (2023), arXiv:2309.14965 [astro-ph.CO].
- [116] T.-N. Li, S.-J. Jin, H.-L. Li, J.-F. Zhang, and X. Zhang, Prospects for Probing the Interaction between Dark Energy and Dark Matter Using Gravitational-wave Dark Sirens with Neutron Star Tidal Deformation, Astrophys. J. 963, 52 (2024), arXiv:2310.15879 [astro-ph.CO].
- [117] H.-Y. Chen, Systematic Uncertainty of Standard Sirens from the Viewing Angle of Binary Neutron Star Inspirals, Phys. Rev. Lett. **125**, 201301 (2020), arXiv:2006.02779 [astro-ph.HE].
- [118] J. Seebeck, V. Ravi, L. Connor, C. Law, D. Simard, and B. Uzgil, The Effects of Selection Biases on the Analysis of Localised Fast Radio Bursts, (2021), arXiv:2112.07639 [astro-ph.HE].