A PDE APPROACH TO THE EXISTENCE AND REGULARITY OF SURFACES OF MINIMUM MEAN CURVATURE VARIATION

LUIS A. CAFFARELLI, PABLO RAÚL STINGA, AND HERNÁN VIVAS

ABSTRACT. We develop an analytic theory of existence and regularity of surfaces (given by graphs) arising from the geometric minimization problem

$$\min_{\mathcal{M}} \frac{1}{2} \int_{\mathcal{M}} \left| \nabla_{\mathcal{M}} H \right|^2 dA$$

where \mathcal{M} ranges over all *n*-dimensional manifolds in \mathbb{R}^{n+1} with prescribed boundary, $\nabla_{\mathcal{M}} H$ is the tangential gradient along \mathcal{M} of the mean curvature H of \mathcal{M} and dA is the differential of surface area. The minimizers, called surfaces of minimum mean curvature variation, are central in applications of computer-aided design, computer-aided manufacturing and mechanics. Our main results show the existence of both smooth surfaces and of variational solutions to the minimization problem together with geometric regularity results in the case of graphs. These are the first analytic results available on the literature for this problem.

1. INTRODUCTION

Our aim is to develop an analytic theory of existence and regularity of surfaces of minimum mean curvature variation, that is, surfaces arising from the geometric minimization problem

(1.1)
$$\min_{\mathcal{M}} \frac{1}{2} \int_{\mathcal{M}} |\nabla_{\mathcal{M}} H|^2 \, dA$$

in the case where the surface \mathcal{M} is the graph of a function u. Here \mathcal{M} ranges over all ndimensional manifolds in \mathbb{R}^{n+1} , $n \geq 1$, with prescribed boundary, H is the mean curvature of \mathcal{M} , $\nabla_{\mathcal{M}}$ represents the tangential gradient over \mathcal{M} and dA is the differential of surface area. Since (1.1) minimizes the quadratic variation of the mean curvature of \mathcal{M} , surfaces with constant mean curvature such as spheres, cylinders, planes and minimal surfaces have zero energy.

Our main results are the existence of smooth surfaces of minimum mean curvature variation, see Theorems 2.2 and 3.1, and the existence and regularity of variational solutions to (1.1), see Theorem 4.1.

There are several difficulties when considering the geometric minimization problem (1.1). One main obstacle is the highly nonlinear and degenerate nature of the problem. If a manifold \mathcal{M} is a smooth minimizer then, by performing normal variations, it can be seen that the Euler-Lagrange equation satisfied by the mean curvature H of \mathcal{M} is

$$\Delta_{\mathcal{M}}^2 H + 2(2H^2 - K)\Delta_{\mathcal{M}}H + 2\langle \nabla_{\mathcal{M}}H, \diamond H \rangle - 2H|H|^2 = 0$$

where $\Delta_{\mathcal{M}}$ is the Laplace–Beltrami operator on \mathcal{M} , K is the Gaussian curvature and $\diamond H$ denotes second tangent operator over \mathcal{M} , see [11]. This is a nonlinear sixth-order equation

²⁰¹⁰ Mathematics Subject Classification. Primary: 35B65, 49Q10, 65D17. Secondary: 35Q74, 53A10.

Key words and phrases. Mean curvature variation, prescribed mean curvature equation, computer-aided design, bending energy, existence and regularity.

Research partially supported by NSF grant 1500871 (USA), Simons Foundation grant 580911 (USA), and Agencia Nacional de Promoción Científica y Tecnológica grant PICT 2019-3530 (Argentina).

for u (the function that represents the graph of \mathcal{M}) which, at this moment, seems to be analytically quite intractable. In fact, comparison principles and uniqueness of solutions for this equation are not known. If one writes H in terms of a parametrization of \mathcal{M} and looks at (1.1) as an energy given by third order derivatives of the parametrization then the nonlinear degenerate structure prevents from applying minimization techniques in typical Hilbert spaces like the Sobolev space $W^{3,2}(\Omega)$. In addition, the solvability of the prescribed mean curvature equation depends in a nontrivial way on geometric properties of the boundary, see [5].

One of the motivations for studying (1.1) comes from computer-aided design (CAD) and computer-aided manufacturing (CAM) problems in engineering, aerospace industry, computer animation and architectural design. A typical problem in CAD and CAM is the robust design of fair surfaces and the creation of shapes such that some aspects of the design process are optimized, see [9]. In many instances, the goal is the creation of complex, smoothly shaped models and surfaces with specified geometric constraints, see [10]. Typically, these problems are approached via a variational principle. In this context, in 1992, Moreton and Séquin proposed in [8] a numerical algorithm to create 2-dimensional fair surfaces \mathcal{M} as minimizers of the geometric energy functional in (1.1). In [11], a finite difference method was proposed to construct surfaces as steady states of a sixth order flow derived from the Euler-Lagrange equation associated with (1.1).

Perhaps the most important aspect in the applications mentioned above is to obtain surfaces that preserve several degrees of geometric continuity where two different surfaces meet. In particular, global continuity of the mean curvature is fundamental in concrete problems such as the design of streamlined surfaces of aircrafts, ships and cars, and numerical evidence of this regularity has been observed in [11]. Furthermore, minimization of geometric functionals given in terms of the mean curvature are important in continuum mechanics as they account for the bending energy of elastic materials [2, 12].

Up to the best of our knowledge, the analytical theory for the existence of surfaces of minimum mean curvature variation in general is missing. No proof of regularity of minimizers and their mean curvature has been available thus far. Our aim is to fill these gaps and establish an analytical foundation from the PDE and variational perspectives.

Before describing our results, we set up the notation. From now on, we assume that \mathcal{M} is the graph of a real-valued function u defined on a bounded domain $\Omega \subset \mathbb{R}^n$, $n \geq 1$, namely, $\mathcal{M} = \{(x, u(x)) : x \in \overline{\Omega}\}$. For $x \in \Omega$, the upward pointing unit normal at (x, u(x)) is

$$\nu(x) = \frac{(-\nabla u(x), 1)}{(1 + |\nabla u(x)|^2)^{1/2}}$$

and the mean curvature H = H(x) is

$$H = \frac{1}{n} \operatorname{div} \left(\frac{\nabla u}{(1 + |\nabla u|^2)^{1/2}} \right).$$

We let $D(u) := (1 + |\nabla u|^2)^{1/2}$ so that dA = D(u) dx. It is not difficult to see that

$$|\nabla_{\mathcal{M}}H|^{2} = |\nabla H|^{2} - \left|\frac{\nabla u \cdot \nabla H}{D(u)}\right|^{2}$$

where ∇ is the usual Euclidean gradient. Then the geometric energy in (1.1) becomes

(1.2)
$$E[\mathcal{M}] := \frac{1}{2} \int_{\Omega} \left[|\nabla H|^2 - \left| \frac{\nabla u \cdot \nabla H}{D(u)} \right|^2 \right] D(u) \, dx.$$

By the Cauchy–Schwartz inequality,

$$\frac{|\nabla H|^2}{D(u)^2} \le |\nabla_{\mathcal{M}} H|^2 \le |\nabla H|^2.$$

Therefore, we will also consider the (larger) simplified energy functional

(1.3)
$$E[H, u] := \frac{1}{2} \int_{\Omega} |\nabla H|^2 D(u) \, dx.$$

In Section 2 we prove the existence of a smooth function u that satisfies the prescribed mean curvature equation and whose mean curvature H is regular up to the boundary $\partial\Omega$ and, in addition, minimizes the simplified energy functional (1.3), see Theorem 2.2. Then, in Section 3, we show how to modify our procedure to construct regular solutions for the case of the geometric energy functional (1.2), see Theorem 3.1. We believe that our method of proof, which is based on a fixed-point argument, will help establish numerical and computational schemes for the construction of minimizers whose mean curvature is guaranteed to be globally continuous. Indeed, our iterative procedure begins by linearizing the energy at some v, so that the numerical iteration may be initiated at, say, a minimal surface v, and continued by solving a linear Euler–Lagrange equation for H. Moreover, our regularity results are sharp and global, not requiring any extra additional assumptions on the boundary of the domain, so they will be useful for proving precise rates of convergence.

Next, in Section 4, we establish a variational formulation for constructing minimizers of (1.3) and show global regularity results for u and H, see Theorem 4.1. We highlight that setting up an appropriate functional setting to look for minimizers is a nontrivial task for reasons like the ones we mentioned before. For instance, the Euler-Lagrange equation seems quite intractable with the available tools and the nonlinear degenerate structure prevents us from using typical Hilbert space methods. We overcome these difficulties by looking for a minimizer pair (u, H), where u is the solution to the variational or weak formulation of the prescribed mean curvature equation with right hand side H. Our functional setting requires mild conditions on u and H on which we can use geometric measure theory tools to recover enough compactness. We also show that our minimizer pairs (u, H) satisfy the required mean curvature global continuity required in applications [8, 11].

2. EXISTENCE AND REGULARITY FOR THE SIMPLIFIED ENERGY

In this section we work with the simplified energy functional (1.3). Consider a bounded domain Ω such that $\partial \Omega \in C^{2,\alpha}$, for some $0 < \alpha < 1$ fixed. We assume that we are given prescribed boundary values $g \in C^{2,\alpha}(\overline{\Omega})$ for u and $h \in C^{1,\alpha}(\overline{\Omega})$ for H on $\partial \Omega$.

We address the following problem: given Ω and the boundary datum g, find a surface \mathcal{M} given by the graph of a function u such that u = g on $\partial \Omega$ and its mean curvature H is a minimizer of (1.3) among all functions with prescribed boundary values h.

Consider the Banach space $\mathfrak{B} = C^{1,\alpha}(\overline{\Omega})$ and its subset $\mathfrak{G} := \{v \in C^{1,\alpha}(\overline{\Omega}) : v = g \text{ on } \partial\Omega\}.$ Clearly, \mathfrak{G} is nonempty, closed and convex. For any $v \in \mathfrak{G}$, we define the functional

(2.1)
$$E[H,v] := \frac{1}{2} \int_{\Omega} |\nabla H|^2 D(v) \, dx.$$

A map $T : \mathfrak{G} \to \mathfrak{G}$ is constructed in a two-step process. First, given any $v \in \mathfrak{G}$, let $H \in W^{1,2}(\Omega)$ be the unique minimizer to (2.1) such that $H - h \in W^{1,2}_0(\Omega)$, which exists

because $1 \le D(v) \le (1 + \|\nabla v\|_{L^{\infty}(\Omega)}^2)^{1/2}$. Then *H* is the unique weak solution to

$$\begin{cases} \operatorname{div}(D(v)\nabla H) = 0 & \text{in } \Omega\\ H = h & \text{on } \partial\Omega. \end{cases}$$

Since $v \in C^{1,\alpha}(\overline{\Omega})$, by global Schauder estimates (see [5]),

(2.2)
$$\|H\|_{C^{1,\alpha}(\overline{\Omega})} \le C_n[\partial\Omega]_{C^{1,\alpha}} \|D(v)\|_{C^{0,\alpha}(\overline{\Omega})} \|h\|_{C^{1,\alpha}(\partial\Omega)}$$

where $C_n > 0$ is a constant that depends only on n.

Second, having constructed this H, we find the solution u to the prescribed mean curvature equation

(2.3)
$$\begin{cases} \operatorname{div}\left(\frac{\nabla u}{D(u)}\right) = nH & \text{in } \Omega\\ u = g & \text{on } \partial\Omega \end{cases}$$

For this, we use the following sharp existence result, see [5].

Theorem 2.1. Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with $C^{2,\alpha}$ boundary, for some $0 < \alpha < 1$. Suppose that $H \in C^1(\overline{\Omega})$ satisfies

(2.4) $||H||_{L^n(\Omega)} < |B_1|^{1/n}$

and, for any $y \in \partial \Omega$,

(2.5)
$$|H(y)| \le \frac{n-1}{n} H_{\partial\Omega}(y)$$

where $|B_1|$ is the measure of the unit ball in \mathbb{R}^n and $H_{\partial\Omega}$ is the mean curvature of $\partial\Omega$ corresponding to the inner unit normal vector to $\partial\Omega$. Then for any $g \in C^{2,\alpha}(\overline{\Omega})$ there exists a unique solution $u \in C^{2,\alpha}(\overline{\Omega})$ to (2.3). In particular, there exists a constant $C_* > 0$ depending only on n, α , $||H||_{L^n(\Omega)}, ||H||_{C^1(\overline{\Omega})}, ||g||_{C^{2,\alpha}(\overline{\Omega})}$ and Ω such that

$$(2.6) ||u||_{C^{2,\alpha}(\overline{\Omega})} \le C_*.$$

Now (2.4) and (2.5) impose further restrictions on the boundary values h of H (see also Remark 2.3). By the maximum principle, if we assume that

(2.7)
$$\max_{\partial\Omega} |h| < \left(\frac{|B_1|}{|\Omega|}\right)^{1/n}$$

then (2.4) holds. Condition (2.5) is natural and cannot be avoided (see Remark 2.3). Therefore, we assume that h additionally satisfies

(2.8)
$$|h(y)| \le \frac{n-1}{n} H_{\partial\Omega}(y)$$
 for all $y \in \partial\Omega$

Therefore, under the additional assumptions (2.7) and (2.8), we can apply Theorem 2.1 and find the unique solution $u \in C^{2,\alpha}(\overline{\Omega})$ to (2.3). This completes the second step.

We then define $T: \mathfrak{G} \to \mathfrak{G}$ by T(v) = u.

Let us prove that T is continuous. Fix $v_1 \in \mathfrak{G}$ and $\varepsilon > 0$. We need to show that there exists $\delta = \delta(\varepsilon, v_1) > 0$ such that for any $v_2 \in \mathfrak{G}$ satisfying $||v_1 - v_2||_{C^{1,\alpha}(\overline{\Omega})} < \delta$ we have $||u_1 - u_2||_{C^{1,\alpha}(\overline{\Omega})} < \varepsilon$, where $u_j = Tv_j$, for j = 1, 2. Let H_j denote the minimizer of $E[\cdot, v_j]$, j = 1, 2, as constructed in the first step. Then $H = H_1 - H_2$ is the unique weak solution to

$$\begin{cases} \operatorname{div}(D(v_1)\nabla H) = \operatorname{div}\left((D(v_2) - D(v_1))\nabla H_2\right) & \text{in } \Omega\\ H = 0 & \text{on } \partial\Omega. \end{cases}$$

By global Schauder estimates,

(2.9)
$$\|H\|_{C^{1,\alpha}(\overline{\Omega})} \leq C_n[\partial\Omega]_{C^{1,\alpha}} \|D(v_1)\|_{C^{0,\alpha}(\overline{\Omega})} \|(D(v_2) - D(v_1))\nabla H_2\|_{C^{0,\alpha}(\overline{\Omega})} \\ \leq C(n,\alpha,\Omega,v_1,\nabla H_2) \|v_1 - v_2\|_{C^{1,\alpha}(\overline{\Omega})} =: C_1 \|v_1 - v_2\|_{C^{1,\alpha}(\overline{\Omega})}.$$

To estimate the difference $u = u_1 - u_2 \in C^{2,\alpha}(\overline{\Omega})$, observe that

$$\begin{cases} \operatorname{div}\left(\frac{\nabla u_1}{D(u_1)} - \frac{\nabla u_2}{D(u_2)}\right) = nH & \text{in } \Omega\\ u = 0 & \text{on } \partial\Omega \end{cases}$$

The vector field

$$F(p) := \frac{p}{\sqrt{1+|p|^2}} \qquad p \in \mathbb{R}^n$$

is smooth and bounded and its gradient $\nabla F(p)$ is a bounded, symmetric matrix. Moreover, $\nabla F(p)$ is locally strictly elliptic, namely, for any $\xi \in \mathbb{R}^n$,

$$\sum_{i,j=1}^n \partial_j F_i(p)\xi_i\xi_j \ge \frac{|\xi|^2}{D(p)^3} \ge \theta(R)|\xi|^2$$

for all |p| < R, where $\theta(R) \to 0$ as $R \to \infty$. Furthermore,

$$F(\nabla u_1) - F(\nabla u_2) = \int_0^1 \frac{d}{dt} F(t\nabla u_1 + (1-t)\nabla u_2) dt$$
$$= \int_0^1 \nabla F(t\nabla u_1 + (1-t)\nabla u_2) \nabla (u_1 - u_2) dt =: A(x)\nabla u_1$$

where A(x) is symmetric and bounded. Now, $u_1 \in C^{2,\alpha}(\overline{\Omega})$ is fixed. By (2.6), the $C^{2,\alpha}(\overline{\Omega})$ norm of u_2 is uniformly controlled by the $C^1(\overline{\Omega})$ norm of H_2 , which in turn is uniformly close to the $C^1(\overline{\Omega})$ norm of the initially fixed H_1 . Therefore, A(x) is strictly elliptic. Moreover, since $\nabla F(p)$ and $D^2F(p)$ are bounded and $\nabla u_1, \nabla u_2 \in C^{0,\alpha}(\overline{\Omega})$, it can be verified that

 $\|A\|_{C^{0,\alpha}(\overline{\Omega})} \le M$

with M > 0 a constant depending only on n, α , $||H_1||_{L^n(\Omega)}$, $||H_1||_{C^1(\overline{\Omega})}$, $||g||_{C^{2,\alpha}(\overline{\Omega})}$, and Ω , see (2.6). All of these quantities are independent of u_2 if v_2 is close to v_1 in $C^{1,\alpha}(\overline{\Omega})$. In summary, we have found that u is a solution to

$$\begin{cases} \operatorname{div}(A(x)\nabla u) = nH & \text{in } \Omega\\ u = 0 & \text{on } \partial\Omega \end{cases}$$

and so, by Schauder estimates,

(2.10)
$$\|u\|_{C^{1,\alpha}(\overline{\Omega})} \le C_n[\partial\Omega]_{C^{1,\alpha}}M\|H\|_{C^{1,\alpha}(\overline{\Omega})} =: C_2\|H\|_{C^{1,\alpha}(\overline{\Omega})}.$$

Collecting estimates (2.9) and (2.10) and recalling that $u = u_1 - u_2 = Tv_1 - Tv_2$, and $H = H_1 - H_2$, we obtain $||Tv_1 - Tv_2||_{C^{1,\alpha}(\overline{\Omega})} \leq C_1 C_2 ||v_1 - v_2||_{C^{1,\alpha}(\overline{\Omega})}$. If we choose $\delta = \varepsilon/(C_1 C_2)$ then we conclude that T is continuous, as desired.

Let us next prove that $T(\mathfrak{G})$ is precompact. Let $\{v_k\}_{k\geq 1}$ be a sequence in \mathfrak{G} such that

$$\sup_{k\geq 1} \|v_k\|_{C^{1,\alpha}(\overline{\Omega})} \leq N_1 < \infty.$$

Consider the corresponding solutions $H_k \in C^{1,\alpha}(\overline{\Omega})$ found in the first step. Set $u_k = Tv_k$. By (2.6),

$$\|u_k\|_{C^{2,\alpha}(\overline{\Omega})} \le C_k$$

where $C_k > 0$ is a constant depending only on n, α , $\|H_k\|_{L^n(\Omega)}$, $\|H_k\|_{C^1(\overline{\Omega})}$, $\|h\|_{C^{2,\alpha}(\overline{\Omega})}$, and Ω . Since all H_k have the same boundary values h, by the maximum principle,

$$\sup_{k\geq 1} \|H_k\|_{L^n(\Omega)} = N_2 < \infty.$$

Furthermore, from the $C^{1,\alpha}$ estimate in (2.2),

$$\sup_{k\geq 1} \|H_k\|_{C^1(\overline{\Omega})} \leq C_n[\partial\Omega]_{C^{1,\alpha}} \|h\|_{C^{1,\alpha}(\partial\Omega)} \sup_{k\geq 1} \|D(v_k)\|_{C^{0,\alpha}(\overline{\Omega})} = N_3 < \infty$$

Consequently,

$$\sup_{k\geq 1} \|u_k\|_{C^{2,\alpha}(\overline{\Omega})} \leq \sup_{k\geq 1} C_k = N_4 < \infty.$$

By the Arzelà–Ascoli compact embedding theorem, there exist a subsequence $\{u_{k_j}\}_{j\geq 1}$ of $\{u_k\}_{k\geq 1}$ and $u \in \mathfrak{G}$ such that $u_{k_j} \to u$ in $C^{1,\alpha}(\overline{\Omega})$, as desired.

Thus, by Schauder's fixed point theorem, there exists $u \in \mathfrak{G}$ such that Tu = u. We have proved the following:

Theorem 2.2 (Existence for the simplified energy and regularity). Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with $C^{2,\alpha}$ boundary $\partial\Omega$, for $0 < \alpha < 1$. Fix $g \in C^{2,\alpha}(\overline{\Omega})$. Let $h \in C^{1,\alpha}(\overline{\Omega})$ such that

(2.11)
$$\max_{\partial\Omega} |h| < \left(\frac{|B_1|}{|\Omega|}\right)^{1/n}$$

and

(2.12)
$$|h(y)| \le \frac{n-1}{n} H_{\partial\Omega}(y) \quad \text{for all } y \in \partial\Omega,$$

where $H_{\partial\Omega}$ is the mean curvature of $\partial\Omega$ corresponding to the inner unit normal vector to $\partial\Omega$. Then there exist $u \in C^{2,\alpha}(\overline{\Omega})$ and $H \in C^{1,\alpha}(\overline{\Omega})$ such that H minimizes the energy

$$\frac{1}{2} \int_{\Omega} |\nabla H|^2 D(u) \, dx$$

among all $H \in W^{1,2}(\Omega)$ such that $H - h \in W^{1,2}_0(\Omega)$, or, equivalently, H is the unique weak solution to

$$\begin{cases} \operatorname{div}(D(u)\nabla H) = 0 & \text{in } \Omega\\ H = h & \text{on } \partial\Omega, \end{cases}$$

and, in addition, H is the mean curvature of the graph of u with prescribed values on $\partial \Omega$:

$$\begin{cases} \frac{1}{n} \operatorname{div} \left(\frac{\nabla u}{D(u)} \right) = H & \text{in } \Omega\\ u = g & \text{on } \partial \Omega \end{cases}$$

Remark 2.3 (Nonexistence of solutions). The conditions imposed on the curvature datum h at the boundary in Theorem 2.2 come from restrictions already present when one seeks for solutions to the prescribed mean curvature equation. Indeed, the equation for H is uniformly elliptic when u is, say, Lipschitz continuous and, therefore, is always solvable. Next, if condition (2.12) is not satisfied, that is,

$$|h(y_0)| > \frac{n-1}{n} H_{\partial\Omega}(y_0)$$
 for some $y_0 \in \partial\Omega$

and $h \ge 0$ (or $h \le 0$) on $\partial\Omega$ then $H \ge 0$ (or $H \le 0$) in Ω and we have that for any $\varepsilon > 0$ there exists $g \in C^{\infty}(\overline{\Omega})$ with $|g| < \varepsilon$ such that the prescribed mean curvature equation with curvature H and boundary values g is not solvable. This is due to the fact that one cannot guarantee boundary gradient estimates, see [5, Section 14.4]. On the other hand, a necessary condition for existence of solutions of the prescribed mean curvature equation is that H satisfies

(2.13)
$$\left| \int_{\Omega} H\eta \, dx \right| \le \frac{(1-\varepsilon_0)}{n} \int_{\Omega} |\nabla \eta| \, dx$$

for some $\varepsilon_0 > 0$, for all $\eta \in C_c^1(\Omega)$, see [4,5] and Section 4. It turns out that (2.4) implies (2.13). This structural condition on H can be guaranteed by imposing (2.11).

3. EXISTENCE AND REGULARITY FOR THE GEOMETRIC ENERGY

In this section we discuss how the technique we developed in the previous section can be applied to the geometric energy functional

$$E[\mathcal{M}] = \frac{1}{2} \int_{\Omega} \left[|\nabla H|^2 - \left| \frac{\nabla u \cdot \nabla H}{D(u)} \right|^2 \right] D(u) \, dx.$$

Let Ω , α , h and g be as in Section 2. Fix $v \in C^{1,\alpha}(\overline{\Omega})$ such that v = g on $\partial \Omega$. Let

$$E_{v}[H] := \frac{1}{2} \int_{\Omega} \left[|\nabla H|^{2} - \left| \frac{\nabla v \cdot \nabla H}{D(v)} \right|^{2} \right] D(v) \, dx = \int_{\Omega} L(\nabla H) \, dx$$

where the smooth Lagrangian L is given by

$$L(p) = \frac{1}{2} \left[|p|^2 - \left| \frac{\nabla v \cdot p}{D(v)} \right|^2 \right] D(v) \quad \text{for } p \in \mathbb{R}^n.$$

It can be seen that L is coercive and uniformly convex, with

$$L_{p_i p_j}(p)\xi_i\xi_j = D(v)|\xi|^2 - \frac{(\nabla v \cdot \xi)^2}{D(v)} \ge \frac{1}{D(v)}|\xi|^2.$$

Thus, there exists a unique minimizer $H \in W^{1,2}(\Omega)$ of $E_v[H]$ such that $H - h \in W^{1,2}_0(\Omega)$. In particular, H is the unique weak solution to

$$\begin{cases} \operatorname{div}(a(x)\nabla H) = 0 & \text{in } \Omega \\ H = h & \text{on } \partial\Omega \end{cases}$$

where $a^{ij}(x) = \delta_{ij}D(v) - v_{x_i}v_{x_j}/D(v) \in C^{0,\alpha}(\overline{\Omega})$ is uniformly elliptic. Then $H \in C^{1,\alpha}(\overline{\Omega})$ and

$$\|H\|_{C^{1,\alpha}(\overline{\Omega})} \le C_n[\partial\Omega]_{C^{1,\alpha}} \|v\|_{C^{1,\alpha}(\overline{\Omega})} \|h\|_{C^{1,\alpha}(\partial\Omega)}.$$

If h satisfies (2.7) and (2.8) then Theorem 2.1 guarantees the existence and uniqueness of the solution $u \in C^{2,\alpha}(\overline{\Omega})$ to (2.3). From here on we can continue with the arguments we did in Section 2 and conclude the following result.

Theorem 3.1 (Existence for the geometric functional and regularity). Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with $C^{2,\alpha}$ boundary $\partial\Omega$, for some $0 < \alpha < 1$. Fix $g \in C^{2,\alpha}(\overline{\Omega})$. Let $h \in C^{1,\alpha}(\overline{\Omega})$ such that

$$\max_{\partial\Omega} |h| < \left(\frac{|B_1|}{|\Omega|}\right)^{1/n}$$

and

$$|h(y)| \le \frac{n-1}{n} H_{\partial\Omega}(y) \quad \text{for all } y \in \partial\Omega,$$

where $H_{\partial\Omega}$ is the mean curvature of $\partial\Omega$ corresponding to the inner unit normal vector to $\partial\Omega$. Then there exist $u \in C^{2,\alpha}(\overline{\Omega})$ and $H \in C^{1,\alpha}(\overline{\Omega})$ such that H minimizes the energy

$$\frac{1}{2} \int_{\Omega} \left[|\nabla H|^2 - \left| \frac{\nabla u \cdot \nabla H}{D(u)} \right|^2 \right] D(u) \, dx$$

among all $H \in W^{1,2}(\Omega)$ such that $H - h \in W^{1,2}_0(\Omega)$, or, equivalently, H is the unique weak solution to

$$\begin{cases} \operatorname{div}(a(x)\nabla H) = 0 & \text{in } \Omega\\ H = h & \text{on } \partial\Omega \end{cases}$$

where

$$a^{ij}(x) = \delta_{ij}D(u) - \frac{u_{x_i}u_{x_j}}{D(u)}$$

and, in addition, H is the mean curvature of the graph of u with prescribed values on $\partial \Omega$:

$$\begin{cases} \frac{1}{n} \operatorname{div} \left(\frac{\nabla u}{D(u)} \right) = H & \text{in } \Omega\\ u = g & \text{on } \partial \Omega \end{cases}$$

4. EXISTENCE AND REGULARITY OF VARIATIONAL SOLUTIONS

We next develop the variational formulation to solve (1.1). It is important to notice that the main result of this section, Theorem 4.1, is of a different nature than Theorems 2.2 and 3.1. Indeed, in Theorem 4.1 we construct a minimizing pair (u, H).

We start by recalling (see [4]) that $u \in BV(\Omega)$ (the space of functions of bounded variation in Ω) is a generalized solution to the prescribed mean curvature equation with (weak) mean curvature $H \in L^1(\Omega)$ and boundary value $g \in L^1(\partial\Omega)$ if

(WPMC)
$$\mathcal{J}[u] = \min_{v \in \mathrm{BV}(\Omega)} \mathcal{J}[v]$$

where

$$\mathcal{J}[v] := \int_{\Omega} D(v) + \int_{\Omega} nHv \, dx + \int_{\partial\Omega} |v - g| \, dS$$

and

$$\int_{\Omega} D(v) := \sup \left\{ \int_{\Omega} \left[v \sum_{i=1}^{n} \partial_{x_i} \phi_i + \phi_{n+1} \right] dx : \phi_i \in C_c^1(\Omega), \ \sum_{i=1}^{n+1} \phi_i^2 \le 1 \right\}.$$

It can be seen that (see [7]) for $v \in W^{1,1}(\Omega)$ we have

$$\int_{\Omega} D(v) = \int_{\Omega} (1 + |\nabla v|^2)^{1/2} \, dx.$$

In [4], Giaquinta proved that if H is a measurable function then (WPMC) is solvable in $BV(\Omega)$ if and only if there is $\varepsilon_0 > 0$ such that, for any set of finite perimeter $A \subset \Omega$,

(4.1)
$$\left| \int_{A} H \, dx \right| \le (1 - \varepsilon_0) \frac{1}{n} P(\partial A)$$

where $P(\partial A)$ denotes the perimeter of A.

The area measure is defined by

$$D(u)(U) = \sup\left\{\int_{\Omega} \left[u\sum_{i=1}^{n} \partial_{x_i}\phi_i + \phi_{n+1}\right] dx : \phi_i \in C_c^1(U), \ \sum_{i=1}^{n+1} \phi_i^2 \le 1\right\}$$

for any $U \subset \Omega$ open and $D(u)(V) = \inf \{D(u)(U) : V \subset U \text{ and } U \text{ is open}\}$, whenever $V \subset \Omega$ is arbitrary. It can be seen that if $u \in BV(\Omega)$ then D(u) is a Radon measure on \mathbb{R}^n .

From now on, we fix a bounded C^2 domain Ω and $g \in C^{1,\alpha}(\partial \Omega)$ for some $0 < \alpha < 1$. We consider the minimization problem

$$\min_{(u,H)\in\mathcal{A}}\mathcal{I}[u,H]$$

where

(4.2)
$$\mathcal{I}[u,H] := \int_{\Omega} |\nabla H|^2 \, dD(u)$$

and dD(u) stands for the area measure defined above. The admissible set \mathcal{A} is defined as follows. Let $h \in W^{2,2}(\Omega) \cap \operatorname{Lip}(\partial\Omega)$ satisfying

(4.3)
$$|h(y)| \le \frac{n-1}{n} H_{\partial\Omega}(y), \ y \in \partial\Omega, \quad \text{and} \quad \max_{\partial\Omega} |h| \le (1-\varepsilon_0) \left(\frac{|B_1|}{|\Omega|}\right)^{1/n},$$

where $H_{\partial\Omega}(y)$ is the mean curvature of $\partial\Omega$ at $y \in \partial\Omega$, for some $0 < \varepsilon_0 < 1$. Define, for some $C_0 > 0$,

(4.4)
$$\mathcal{A} := \left\{ \begin{array}{cc} (u,H) \in \mathrm{BV}(\Omega) \times (\mathrm{Lip}(\overline{\Omega}) \cap W^{2,2}(\Omega)) : u \text{ solves } (\mathrm{WPMC}) \\ \mathrm{and} \ \|H\|_{\mathrm{Lip}(\overline{\Omega})} + \|H\|_{W^{2,2}(\Omega)} \le C_0, \ H = h \text{ on } \partial\Omega \end{array} \right\}.$$

Theorem 4.1 (Existence and regularity of variational solutions). Let Ω be a bounded domain with C^2 boundary $\partial\Omega$. Let $g \in C^{1,\alpha}(\partial\Omega)$ for some $0 < \alpha < 1$, and $h \in W^{2,2}(\Omega) \cap \operatorname{Lip}(\partial\Omega)$ satisfying (4.3) for some $0 < \varepsilon_0 < 1$. Let \mathcal{I} be defined by (4.2). Then there is $C_0 > 0$, depending only on $\partial\Omega$ and $\|h\|_{L^{\infty}(\partial\Omega)}$, such that the admissible set \mathcal{A} in (4.4) is nonempty and there exists a minimizer (u_{∞}, H_{∞}) of \mathcal{I} within the class \mathcal{A} . Moreover, $u_{\infty} \in C^{1,\alpha}(\overline{\Omega}) \cap C^{2,\alpha}_{\operatorname{loc}}(\Omega)$.

To prove Theorem 4.1 we need to recall the notion and properties of Γ -convergence (see [1]) in our context. Let $\mathcal{J}_k, k \geq 1$, and \mathcal{J}_∞ be functionals defined on BV(Ω) and taking values in $[-\infty, \infty]$. Then $\{\mathcal{J}_k\}_{k\geq 1}$ is said to Γ -converge to \mathcal{J}_∞ if the following two conditions hold:

(a) For every $v \in BV(\Omega)$ and every sequence $\{v_k\}_{k\geq 1} \subset BV(\Omega)$ such that $v_k \to v$ in $BV(\Omega)$ it holds

$$\liminf_{k \to \infty} \mathcal{J}_k(v_k) \ge \mathcal{J}_\infty(v).$$

(b) For every $v \in BV(\Omega)$ there exists a sequence $\{v_k\}_{k\geq 1} \subset BV(\Omega)$ such that $v_k \to v$ in $BV(\Omega)$ for which

$$\limsup_{k \to \infty} \mathcal{J}_k(v_k) \le \mathcal{J}_\infty(v).$$

We will use the following fact (see [1, Theorem 1.21]). Suppose that $\{\mathcal{J}_k\}_{k\geq 1}$ is an equi-mildly coercive sequence of functionals on BV(Ω) that Γ -converges to \mathcal{J}_{∞} . Then there exits

$$\min_{\mathrm{BV}(\Omega)} \mathcal{J}_{\infty} = \lim_{k \to \infty} \inf_{\mathrm{BV}(\Omega)} \mathcal{J}_k$$

Moreover, if $\{u_k\}_{k\geq 1} \subset BV(\Omega)$ is a precompact sequence in $BV(\Omega)$ such that

$$\lim_{k \to \infty} \mathcal{J}_k(u_k) = \lim_{k \to \infty} \inf_{\mathrm{BV}(\Omega)} \mathcal{J}_k$$

then every limit of $\{u_k\}_{k\geq 1}$ is a minimum point for \mathcal{J}_{∞} . A functional \mathcal{J} is mildly coercive in $\mathrm{BV}(\Omega)$ if there exists a nonempty compact set $K \subset \mathrm{BV}(\Omega)$ such that $\inf_K \mathcal{J} = \inf_{\mathrm{BV}(\Omega)} \mathcal{J}$, and equi-mild coercivity means that the set K is the same for the whole sequence $\{\mathcal{J}_k\}_{k\geq 1}$.

Proof of Theorem 4.1. We begin by showing that \mathcal{A} is nonempty. Let H be the harmonic function in Ω such that H = h on $\partial\Omega$. By elliptic regularity, $H \in \operatorname{Lip}(\overline{\Omega}) \cap W^{2,2}(\Omega)$ and $\|H\|_{\operatorname{Lip}(\overline{\Omega})} + \|H\|_{W^{2,2}(\Omega)} \leq C_0$, where $C_0 = C_0(\partial\Omega, \|h\|_{L^{\infty}(\partial\Omega)}) > 0$. Moreover, by the Hölder and isoperimetric inequalities,

$$\int_{A} H \, dx \bigg| \le \|H\|_{L^{n}(\Omega)} |A|^{\frac{n-1}{n}} \le \|H\|_{L^{n}(\Omega)} \frac{P(\partial A)}{n|B_{1}|^{1/n}}$$

The maximum principle and (4.3) give

$$||H||_{L^n(\Omega)} \le |\Omega|^{1/n} \max_{\partial \Omega} |h| \le (1 - \varepsilon_0) |B_1|^{1/n}$$

Therefore,

$$\left| \int_{A} H \, dx \right| \le \frac{(1 - \varepsilon_0)}{n} P(\partial A)$$

and (WPMC) is solvable for this H. If we let $u \in BV(\Omega)$ be the corresponding minimizer of \mathcal{J} then the pair (u, H) is in \mathcal{A} .

We further point out that $\int_{\Omega} D(u) < \infty$ and $H \in \operatorname{Lip}(\overline{\Omega})$ so that

$$0 \le m := \inf_{(u,H) \in \mathcal{A}} \mathcal{I}[u,H] \le \|\nabla H\|_{L^{\infty}(\Omega)}^2 D(u)(\Omega) < \infty.$$

Consider next a minimizing sequence $\{(u_k, H_k)\}_{k \ge 1} \subset \mathcal{A}$, that is, $\lim_{k \to \infty} \mathcal{I}[u_k, H_k] = m$. Since every u_k is a minimizer of the functional \mathcal{J}_k defined by

(4.5)
$$\mathcal{J}_k[v] := \int_{\Omega} D(v) + \int_{\Omega} nH_k v \, dx + \int_{\partial\Omega} |v - g| \, dS$$

we have that, for any $u_0 \in BV(\Omega)$, $\mathcal{J}_k(u_k) \leq \mathcal{J}_k(u_0)$, from where

(4.6)
$$\int_{\Omega} D(u_k) + \int_{\Omega} nH_k u_k \, dx \le C + \int_{\Omega} nH_k u_0 \, dx$$

for C > 0 independent of k. Now we estimate by below the second integral in the left hand side above as in [4]. Indeed, we extend H_k and u_k as 0 outside of Ω and write

$$\int_{\Omega} nH_k u_k \, dx = \int_{\mathbb{R}^n} nH_k u_k^+ \, dx - \int_{\mathbb{R}^n} nH_k u_k^- \, dx$$

where $u^+, u^- \ge 0$ denote the positive and negative parts of a function u, respectively. By (4.1) and the coarea formula for functions of bounded variation (see [3]),

$$\left| \int_{\mathbb{R}^n} nH_k u_k^{\pm} dx \right| = \left| \int_0^{\infty} \int_{\{x: u_k^{\pm}(x) > t\}} nH_k dx dt \right|$$
$$\leq (1 - \varepsilon_0) \int_0^{\infty} P(\partial\{x: u_k^{\pm}(x) > t\}) dt$$
$$= (1 - \varepsilon_0) \int_{\mathbb{R}^n} |\nabla u_k^{\pm}|.$$

Hence,

$$\int_{\Omega} nH_k u_k \, dx \ge -(1-\varepsilon_0) \int_{\mathbb{R}^n} |\nabla u_k|$$
$$= -(1-\varepsilon_0) \int_{\Omega} |\nabla u_k| - (1-\varepsilon_0) \int_{\partial \Omega} |u_k| \, dS$$
$$= -(1-\varepsilon_0) \int_{\Omega} |\nabla u_k| - C$$

as $u_k = g$ on $\partial \Omega$ for all k (see [4]). Using this in (4.6),

$$\int_{\Omega} D(u_k) \le (1 - \varepsilon_0) \int_{\Omega} |\nabla u_k| + n \|H_k\|_{L^{\infty}(\Omega)} \|u_0\|_{L^1(\Omega)} + C$$

for a new constant C > 0 that is independent of k. Moreover, the uniform bound on the $L^{\infty}(\Omega)$ norm of $\{H_k\}_{k\geq 1}$ gives

$$\int_{\Omega} |\nabla u_k| \le (1 - \varepsilon_0) \int_{\Omega} |\nabla u_k| + nC_0 ||u_0||_{L^1(\Omega)} + C.$$

so that

$$\varepsilon_0 \int_{\Omega} |\nabla u_k| \le nC_0 ||u_0||_{L^1(\Omega)} + C.$$

Hence, by compactness in $BV(\Omega)$, there exist a subsequence of $\{u_k\}_{k\geq 1}$, still denoted by the same indexes, and $u_{\infty} \in BV(\Omega)$ such that $u_k \to u_{\infty}$ in $L^1(\Omega)$ as $k \to \infty$, and $|\nabla u_{\infty}|(U) \leq \lim \inf_{k\to\infty} |\nabla u_k|(U)$ for any Borel set $U \subset \Omega$. In addition,

$$(4.7) D(u_{\infty})(U) \le \liminf_{k \to \infty} D(u_k)(U).$$

By Poincaré's inequality and the Rellich–Kondrachov compactness theorem, there exist a subsequence of $\{H_k\}_{k\geq 1}$, still denoted by the same indexes, and $H_{\infty} \in W^{2,2}(\Omega)$ such that

(4.8)
$$\nabla H_k \to \nabla H_\infty \quad \text{in } L^2(\Omega), \text{ as } k \to \infty.$$

Further, due to the uniform bound on $||H_k||_{\operatorname{Lip}(\overline{\Omega})}$, we may assume that H_k and ∇H_k converge weak-* in $L^{\infty}(\Omega)$ to $H_{\infty} \in \operatorname{Lip}(\overline{\Omega})$. Finally, this and the weak convergence of H_k to H_{∞} in $W^{2,2}(\Omega)$ ensure that

$$\|H_{\infty}\|_{\operatorname{Lip}(\overline{\Omega})} + \|H_{\infty}\|_{W^{2,2}(\Omega)} \le C_0.$$

Let us now prove that $(u_{\infty}, H_{\infty}) \in \mathcal{A}$. Recall the functionals \mathcal{J}_k defined in (4.5) for the subsequence H_k we just found, and define \mathcal{J}_{∞} analogously. We claim that $\{\mathcal{J}_k\}_{k\geq 1}$ Γ -converges to \mathcal{J}_{∞} . Indeed, it is sufficient to prove the Γ -convergence of

$$\widetilde{\mathcal{J}}_k(v) := \int_{\Omega} v H_k \, dx \quad \text{to} \quad \widetilde{\mathcal{J}}_{\infty}(v) := \int_{\Omega} v H_{\infty} \, dx$$

since the other two terms are continuous perturbations of \mathcal{J}_k (see [1]). To prove the limit inequality (a), let $\{v_k\}_{k\geq 1} \subset BV(\Omega)$ and $v \in BV(\Omega)$ such that $v_k \to v$ in $BV(\Omega)$. We write

$$\int_{\Omega} v_k H_k \, dx - \int_{\Omega} v H_\infty \, dx = I_k + II_k + III_k$$

with

$$I_k = \int_{\Omega} (v_k - v) H_{\infty} dx$$
$$II_k = \int_{\Omega} (v_k - v) (H_k - H_{\infty}) dx$$
$$III_k = \int_{\Omega} v (H_k - H_{\infty}) dx.$$

By lower semicontinuity (as in [6, Proposition 2.1]), $\liminf_{k\to\infty} I_k \ge 0$. Next, we bound

$$|II_k| \le ||v_k - v||_{L^1(\Omega)} \left(||H_k||_{L^{\infty}(\Omega)} + ||H_{\infty}||_{L^{\infty}(\Omega)} \right)$$

The convergence of v_k to v in $L^1(\Omega)$ and the uniform bound of H_k in $L^{\infty}(\Omega)$ give $\lim_{k\to\infty} II_k = 0$. Finally, $\lim_{k\to\infty} III_k = 0$ by the weak-* convergence of H_k to H_{∞} in $L^{\infty}(\Omega)$. As for the limsup inequality (b), given any $v \in BV(\Omega)$, consider the constant sequence $v_k = v$ for all $k \geq 1$ and notice that, using the weak-* convergence of H_k to H_{∞} in $L^{\infty}(\Omega)$,

$$\lim_{k \to \infty} \widetilde{\mathcal{J}}_k(v_k) = \widetilde{\mathcal{J}}_\infty(v).$$

Hence, $\{\mathcal{J}_k\}_{k\geq 1}$ converges to \mathcal{J}_{∞} in the Γ sense. Furthermore, the sequence $\{\mathcal{J}_k\}_{k\geq 1}$ is equimildly coercive. Indeed, by regularity estimates for the prescribed mean curvature equation, all the minimizers of each $\mathcal{J}_k, k \geq 1$, are in a ball in $C^{1,\alpha}(\overline{\Omega})$ of radius depending only on C_0, Ω, g and h, and this ball is compact in $BV(\Omega)$.

Consequently, u_{∞} is a minimizer of \mathcal{J}_{∞} and $(u_{\infty}, H_{\infty}) \in \mathcal{A}$.

Finally, by convexity,

$$\int_{\Omega} |\nabla H_k|^2 \, dD(u_k) \ge \int_{\Omega} |\nabla H_{\infty}|^2 \, dD(u_k) + 2 \int_{\Omega} \nabla H_{\infty} \cdot (\nabla H_k - \nabla H_{\infty}) \, dD(u_k).$$

As $k \to \infty$, the left hand side converges to m. As for the right hand side, (4.7) implies that

$$\liminf_{k \to \infty} \int_{\Omega} |\nabla H_{\infty}|^2 \, dD(u_k) \ge \int_{\Omega} |\nabla H_{\infty}|^2 \, dD(u_{\infty}).$$

For the second term on the right hand side above, the regularity of the prescribed mean curvature equation gives that $u_k \in \operatorname{Lip}(\Omega)$, with Lipschitz constant bounded by some constant c > 0 independent of k (see [4]). Hence,

$$\left| \int_{\Omega} \nabla H_{\infty} \cdot (\nabla H_k - \nabla H_{\infty}) \, dD(u_k) \right| \le (1 + c^2)^{1/2} \|\nabla H_{\infty}\|_{L^2(\Omega)} \|\nabla H_k - \nabla H_{\infty}\|_{L^2(\Omega)}.$$

In view of (4.8), this term goes to 0 as $k \to \infty$. We have shown that (u_{∞}, H_{∞}) is a minimizer.

Since the first relation in (4.3) holds, by the results in [4] (for which $\partial \Omega \in C^2$ is in fact enough, see also [5, Theorem 13.2] and the comment following it), we conclude that $u \in C^{1,\alpha}(\overline{\Omega}) \cap C^{2,\alpha}_{\text{loc}}(\Omega)$.

Remark 4.2. We point out that the first condition in (4.3) is only used to deduce the boundary regularity of u_{∞} , and is not actually needed in the construction of the pair (u_{∞}, H_{∞}) .

Acknowledgements. We would like to thank Irene Martínez Gamba and the referee for useful comments that helped improve the presentation of the paper.

STATEMENTS AND DECLARATIONS

Data Availability Statement. Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

Conflict of Interest Statement. The authors declare no conflict of interest.

References

- [1] A. Braides, Γ -convergence for Beginners, Oxford Lecture Series in Mathematics and its Applications 22, Oxford University Press, Oxford, 2002.
- [2] R. Capovilla, Elastic bending energy: a variational approach, Journal of Geometry and Symmetry in *Physics* **45** (2017), 1–45.
- [3] L. Evans and R. F. Gariepy, Measure Theory and Fine Properties of Functions, Revised Edition, Textbooks in Mathematics, CRC Press, Boca Raton, FL, 2015.
- [4] M. Giaquinta, On the Dirichlet problem for surfaces of prescribed mean curvature, Manuscripta Math. 12 (1974), 73-86.

- [5] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001.
- [6] E. Giusti, Boundary value problems for non-parametric surfaces of prescribed mean curvature, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 3 (1976), 501–548.
- [7] E. Giusti, *Minimal Surfaces and Functions of Bounded Variation*, Monographs in Mathematics **80**, Birkhäuser Verlag, Basel, 1984.
- [8] H. P. Moreton and C. H. Séquin Functional optimization for fair surface design, ACM SIGGRAPH Computer Graphics 2 (1992), 167-176.
- [9] K. L. Narayan, K. M. Rao and M. M. M. Sacar, Computer Aided Design and Manufacturing, PHI Learning Pvt. Ltd., 2008.
- [10] W. Welch and A. Witkin, Variational surface modeling, ACM SIGGRAPH computer graphics 2 (1992), 157-166.
- [11] G. Xu and Q. Zhang, Minimal mean-curvature-variation surfaces and their applications in surface modeling, in: International Conference on Geometric Modeling and Processing, 357–370, Springer, Berlin, Heidelberg, 2006.
- [12] D. Zorin, Curvature-based energy for simulation and variational modeling, in: International Conference on Shape Modeling and Applications 2005, 196–204, Cambridge, MA, 2005.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN, 2515 SPEEDWAY, AUSTIN, TX 78712, UNITED STATES OF AMERICA

Email address: caffarel@math.utexas.edu

DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY, 396 CARVER HALL, AMES, IA 50011, UNITED STATES OF AMERICA

Email address: stinga@iastate.edu

Centro Marplatense de Investigaciones Matemáticas/CONICET, Dean Funes 3350, 7600 Mar del Plata, Argentina

Email address: havivas@mdp.edu.ar