

Graphs with girth $2\ell + 1$ and without longer odd holes are 3-colorable

Rong Chen

Center for Discrete Mathematics, Fuzhou University
Fuzhou, P. R. China

January 3, 2023

1

Abstract

For a number $\ell \geq 2$, let \mathcal{G}_ℓ denote the family of graphs which have girth $2\ell + 1$ and have no odd hole with length greater than $2\ell + 1$. Plummer and Zha conjectured that every 3-connected and internally 4-connected graph in \mathcal{G}_2 is 3-colorable. Wu, Xu, and Xu conjectured that every graph in $\bigcup_{\ell \geq 2} \mathcal{G}_\ell$ is 3-colorable. Chudnovsky et al. and Wu et al., respectively, proved that every graph in \mathcal{G}_2 and \mathcal{G}_3 is 3-colorable. In this paper, we prove that every graph in $\bigcup_{\ell \geq 5} \mathcal{G}_\ell$ is 3-colorable.

Key Words: chromatic number; odd holes

1 Introduction

All graphs considered in this paper are finite, simple, and undirected. A *proper coloring* of a graph G is an assignment of colors to the vertices of G such that no two adjacent vertices receive the same color. A graph is *k-colorable* if it has a proper coloring using at most k colors. The *chromatic number* of G , denoted by $\chi(G)$, is the minimum number k such that G is k -colorable.

The *girth* of a graph G , denoted by $g(G)$, is the minimum length of a cycle in G . A *hole* in a graph is an induced cycle of length at least four. An *odd hole* means a hole of odd length. For any integer $\ell \geq 2$, let \mathcal{G}_ℓ be the family of graphs that have girth $2\ell + 1$ and have no odd holes of length at least $2\ell + 3$. Robertson conjectured in [3] that the Petersen graph is the only graph in \mathcal{G}_2 that is 3-connected and internally 4-connected. Plummer and Zha [4] disproved Robertson's conjecture

¹Mathematics Subject Classification: 05C15, 05C17, 05C69
Email: rongchen@fzu.edu.cn (R. Chen).

and proposed the conjecture that all 3-connected and internally 4-connected graphs in \mathcal{G}_2 have bounded chromatic number, and the strong conjecture that such graphs are 3-colorable. The first was proved by Xu, Yu, and Zha [7], who proved that all graphs in \mathcal{G}_2 is 4-colorable. Chudnovsky and Seymour [2] proved that every graph in \mathcal{G}_2 is 3-colorable. In the same paper, Chudnovsky and Seymour also gave a short and pretty proof of the result in [7]. Wu, Xu, and Xu [5] showed that graphs in $\bigcup_{\ell \geq 2} \mathcal{G}_\ell$ are 4-colorable and proposed the following conjecture.

Conjecture 1.1. ([5], Conjecture 6.1.) *Every graph in $\bigcup_{\ell \geq 2} \mathcal{G}_\ell$ is 3-colorable.*

We say that a graph G has an *odd K_4 -subdivision* if some subgraph H of G is isomorphic to a K_4 -subdivision and whose face cycles are all odd holes of G . The subgraph H is also induced by ([1], Lemma 2.7). In the same paper, the author and Zhou proved

Theorem 1.2. ([1], Theorem 1.1.) *For each graph G in $\bigcup_{\ell \geq 5} \mathcal{G}_\ell$, if G has an odd K_4 -subdivision, then $\chi(G) = 3$.*

Based on Theorem 1.2, in this paper we prove that Conjecture 1.1 holds for $\ell \geq 5$. That is,

Theorem 1.3. *All graphs in $\bigcup_{\ell \geq 5} \mathcal{G}_\ell$ are 3-colorable.*

We conjecture

Conjecture 1.4. *For a graph G without triangles, if all odd holes of G have the same length, then G is 3-colorable.*

Conjecture 1.5. *For an integer $\ell \geq 2$ and a graph G with $g(G) = 2\ell + 1$, if the set of lengths of odd holes of G have k members, then G is $(k + 2)$ -colorable.*

2 Preliminary

A *cycle* is a connected 2-regular graph. Let G be a graph. For any subset $U \subseteq V(G)$, let $G[U]$ be the induced subgraph of G defined on U . For subgraphs H and H' of G , set $|H| := |E(H)|$ and $H\Delta H' := E(H)\Delta E(H')$. Let $H \cup H'$ denote the subgraph of G with vertex set $V(H) \cup V(H')$ and edge set $E(H) \cup E(H')$. Let $H \cap H'$ denote the subgraph of G with edge set $E(H) \cap E(H')$ and without isolated vertex. Let $N(H)$ be the set of vertices in $V(G) - V(H)$ that have a neighbour in $V(H)$. Set $N[H] := N(H) \cup V(H)$.

Let P be an (x, y) -path and Q a (y, z) -path. When P and Q are internally disjoint paths, let PQ denote the (x, z) -path $P \cup Q$. Evidently, PQ is a path when $x \neq z$, and PQ is a cycle when $x = z$. Let P^* denote the set of internal vertices

of P . When $u, v \in V(P)$, let $P(u, v)$ denote the subpath of P with ends u, v . For simplicity, we will let $P^*(u, v)$ denote $(P(u, v))^*$.

For a number $k \geq 2$, a graph G is k -vertex-critical if $\chi(G) = k$ and $\chi(G - v) \leq k - 1$ for each vertex v of G .

Lemma 2.1. ([1], Lemma 2.2.) *For any number $k \geq 4$, each k -vertex-critical graph has no 2-edge-cut.*

For an i -vertex path P of a connected graph G , if $G - V(P)$ is disconnected, then we say that P is a P_i -cut. Usually, a P_2 -cut is also called a K_2 -cut. Evidently, every k -vertex-critical graph has no K_2 -cut. Chudnovsky and Seymour in [2] proved that every 4-vertex-critical graph G in \mathcal{G}_2 has no P_3 -cut. In fact, their methods can also be used to prove the following result.

Lemma 2.2. ([2]) *For any number $\ell \geq 2$, every 4-vertex-critical graph in \mathcal{G}_ℓ has neither K_2 -cut nor P_3 -cut.*

A *theta graph* is a graph that consists of a pair of distinct vertices joined by three internally disjoint paths. Let C be a hole of a graph G . A path P of G is a *chordal path* of C if $C \cup P$ is an induced theta-subgraph of G . Since $g(G) = 2\ell + 1$ and each odd hole of G has length $2\ell + 1$, by simple computation we have the following result, which will be frequently used.

Lemma 2.3. ([1], Lemma 2.3.) *Let C be an odd hole of a graph $G \in \mathcal{G}_\ell$. Let P be a chordal path of C , and P_1, P_2 be the internally disjoint paths of C that have the same ends as P . Assume that $|P|$ and $|P_1|$ have the same parity. Then*

$$|P_1| = 1 \text{ or } \ell \geq |P_2| < |P_1| = |P| \geq \ell + 1.$$

In particular, when $|P_1| \geq 2$, all chordal paths of C with the same ends as P_1 have length $|P_1|$.

Let C_1, C_2 be odd holes of a graph $G \in \mathcal{G}_\ell$ such that $C_1 \cup C_2$ is a theta subgraph G . When $C_1 \cup C_2$ is not induced, since $|C_1 \Delta C_2| \leq 4\ell$ and $g(G) = 2\ell + 1$, we have that $|C_1 \Delta C_2| = 4\ell$ and $G[V(C_1 \cup C_2)]$ is an odd K_4 -subdivision. For the similar reason, if C is an even hole of length 4ℓ of G , then C has at most two chords, and when C has two chords, $G[V(C)]$ is an odd K_4 -subdivision.

3 Graphs has a subgraph isomorphic to a K_4 -subdivision

Let H be a graph that is isomorphic to a K_4 -subdivision. For a path P of H whose ends are degree-3 vertices of H , if P contains exactly two degree-3 vertices of H , then we say it is an *arris* of H . Evidently, H has exactly six arrises. We say that a graph




Figure 1: u_1, u_2, u_3, u_4 are the degree-3 vertices of H . The face cycles C_1, C_2 are odd holes, and C_3, C_4 are even holes. $\{P_1, P_2\}, \{Q_1, Q_2\}, \{L_1, L_2\}$ are the pairs of vertex disjoint arrises of H .

G contains a *balanced K_4 -subdivision* if G has an induced subgraph H isomorphic to a K_4 -subdivision and exactly two face cycles of H are odd holes of G .

Lemma 3.1. *If a graph G in \mathcal{G}_ℓ contains a balanced K_4 -subdivision H that is pictured as the graph in Figure 1, then the following hold.*

- (1) $|P_1| \leq \ell$ and $1 \in \{|Q_1|, |Q_2|, |L_1|, |L_2|\}$.
- (2) $|P_2| \geq \ell$.
- (3) *Assume that G has no odd K_4 -subdivision. If $|Q_1| = 1$ and $|L_2| \geq 2$, then no vertex $v \in V(G) - V(H)$ has two neighbours in $V(H)$.*

Proof. Since C_1, C_2 are odd holes and $g(G) = 2\ell + 1$, we have $|P_1| \leq \ell$. Assume that $1 \notin \{|Q_1|, |Q_2|, |L_1|, |L_2|\}$. Since P_2Q_2 is a chordal path of C_1 and C_4 is an even hole, we have $|P_2| + |Q_2| = |L_1|$ by Lemma 2.3. Similarly, we have $|P_2| + |L_1| = |Q_2|$, which is a contradiction. So (1) holds.

Now we consider (2). By (1) and symmetry we may assume that $|Q_1| = 1$. Since $|P_1| \leq \ell$ by (1), we have $|L_1| \geq \ell$, so Q_1P_2 is a chordal path of C_2 . Then $|Q_1P_2| \geq \ell + 1$ by Lemma 2.3. So $|P_2| \geq \ell$. This proves (2).

Next, we prove that (3) is true. Assume not. Let $v \in V(G) - V(H)$ have at least two neighbours in $V(H)$. Since $|L_2| > 1$ and C_3 is an even hole, $C_2\Delta C_3$ is an odd hole and $|L_2| \geq \ell + 1$ by Lemma 2.3. Moreover, since a vertex not in an odd hole has at most one neighbour in the odd hole, each arris of H has at most one vertex adjacent to v .

3.1.1. *The vertex v has exactly one neighbour in $V(C_1 \cup C_2)$ and exactly one neighbour in $V(P_2^*)$.*

Subproof. Since v has at most one neighbour in $V(P_2)$, it suffices to show that v has at most one neighbour in $V(C_1 \cup C_2)$. Assume not. Then v has exactly two neighbours in $V(C_1 \cup C_2)$ with one in $V(C_1) - V(P_1)$ and the other in $V(C_2) - V(P_2)$. Then $G[V(C_1 \cup C_2) \cup \{v\}]$ is a balanced K_4 -subdivision as G has no odd K_4 -subdivision, which is not possible by (2). \square

Note that $C_2\Delta C_3$ is an odd hole. Since v can not have two neighbours in an odd hole of H , the vertex v has no neighbour in $V(P_1)$. For the same reason, if v has a neighbour in $V(Q_1 \cup Q_2)$, the neighbours of v in $V(H)$ must be in $V(C_1 \cup C_2)$, which is not possible by 3.1.1. Hence, $N(v) \cap V(C_1 \cup C_2) \subseteq V(L_1^* \cup L_2^*)$. Assume that v has a neighbour in $V(L_1^*)$. Since $|C_4| \leq 4\ell - 2$ and $g(G) = 2\ell + 1$, by 3.1.1, the two cycles in $G[V(C_4) \cup \{v\}]$ containing v are odd holes. Moreover, since $|L_2| \geq \ell + 1$, the subgraph $G[V(C_4) \cup \{v\}] \cup P_1 \cup Q_1$ is an odd K_4 -subdivision, which is not possible. Similarly we can show that v can not have a neighbour in $V(L_2^*)$. This proves (3). \square

Let H be an induced subgraph of G pictured as the graph in Figure 1. When $|Q_1| = 1$, we say that H is a *balanced K_4 -subdivision of type (1, 2)* if $|L_2| > 1$. Since $|Q_1| = 1$ implies that $|L_1| > 1$, this definition is well defined.

Let H_1, H_2 be vertex disjoint induced subgraphs of a graph G . An induced (v_1, v_2) -path P is a *direct connection* linking H_1 and H_2 if v_1 is the only vertex in $V(P)$ having a neighbour in H_1 and v_2 is the only vertex in $V(P)$ having a neighbour in H_2 . Evidently, the set of internal vertices of each shortest induced path joining H_1 and H_2 induces a direct connection linking H_1 and H_2 .

Theorem 3.2. *Let $\ell \geq 5$ be an integer and G be a graph in \mathcal{G}_ℓ . If G is 4-vertex critical, then G does not contain a balanced K_4 -subdivision of type (1, 2).*

Proof. By Theorem 1.2, G does not have an odd K_4 -subdivision. Assume to the contrary that G has a balanced K_4 -subdivision H of type (1, 2) that is pictured as the graph in Figure 1. Without loss of generality we may assume that H is chosen with $|H|$ as small as possible. By Lemma 3.1 (1) and symmetry we may assume that $|Q_1| = 1$. By Lemmas 3.1 (2) and 2.3, we have

$$|Q_1| = 1, |P_2| \geq \ell, |L_1|, |L_2| \geq \ell + 1, \text{ and } |P_1| < \ell. \quad (4.1)$$

Let e, f be the edges in P_2 incident with u_3, u_4 , respectively. Let P be a direct connection in $G \setminus \{e, f\}$ linking P_2^* and $H - V(P_2^*)$. Lemma 2.1 implies that such P exists. Let v_1, v_2 be the ends of P with v_2 having a neighbour in $V(P_2^*)$ and v_1 having a neighbour in $V(H) - V(P_2^*)$. By Lemma 3.1 (3), both v_1 and v_2 have a unique neighbour in $V(H)$. Let x be the neighbour of v_1 in $V(H)$ and y the

neighbour of v_2 in $V(H)$. Set $P' := xv_1Pv_2y$. Then $H \cup P'$ is an induced subgraph of G . By Lemma 3.1 (3) again, we have $|P'| \geq 3$.

3.2.1. $x \neq u_2$.

Subproof. Assume that $x = u_2$. Set $C'_3 := L_2P_2(u_4, y)P'$. Since $|L_2| \geq \ell + 1$, we have that C'_3 is an even hole by Lemma 2.3. Since $|P'| \geq 3$ and $C_2\Delta C'_3$ is an odd hole, $(H \cup P') - V(L_2^*)$ is a balanced K_4 -subdivision of type (1, 2) whose edge number is less than $|H|$, which is a contradiction to the choice of H . \square

3.2.2. $x \notin P_1$.

Subproof. Assume not. By 3.2.1, we have $x \in V(P_1) - \{u_2\}$. Set $C'_1 := P_1(x, u_2)Q_1P_2(u_3, y)P'$. Since $|P_1| < \ell$ by (4.1), it follows from Lemma 2.3 that C'_1 is an odd hole. Then $P_1(x, u_1)Q_2P_2(u_4, y)P'$ is an even hole. Moreover, since $|L_2| \geq \ell + 1$, we have $x = u_1$ and $|Q_2| = 1$ by Lemma 2.3. Since C_1 and C'_1 are odd holes, we have $|L_1| > |P'|$. Moreover, since $|P'| \geq 3$, the subgraph $C'_1 \cup C_2 \cup P_2$ is a balanced K_4 -subdivision of type (1, 2) whose edge number is less than $|H|$, which is a contradiction. \square

3.2.3. When $x = u_3$, we have $|P_2(u_3, y)| = 1$ or $|P_2(u_3, y)| \geq \ell + 1$.

Subproof. Assume that $|P_2(u_3, y)| \neq 1$. Since Q_1P_2 and $Q_1P'P_2(y, u_4)$ are chordal paths of C_2 that have the same ends as L_2 , they have length $|L_2|$ by Lemma 2.3. So $|P_2(u_3, y)P'| = 2|P_2(u_3, y)|$, implying that $|P_2(u_3, y)| \geq \ell + 1$. \square

3.2.4. If $x \in V(L_1^*)$, then $xu_3, yu_3 \in E(G)$, and $|Q_2| = 1$.

Subproof. Set $C'_3 := L_1(x, u_3)P_2(u_3, y)P'$. When C'_3 is an odd hole, since $C'_3\Delta C_4$ is an odd hole, $C_1 \cup C'_3 \cup P_2 \cup Q_2$ is an odd K_4 -subdivision, which is not possible. So C'_3 is an even hole. Assume that $xu_3 \notin E(G)$. Then $|C'_3| = 2|L_1(x, u_3)|$ by Lemma 2.3. Moreover, since $C_1\Delta C'_3$ is an odd hole, $(H \cup P') - V(L_1^*(x, u_3))$ is a balanced K_4 -subdivision of type (1, 2) whose edge number is less than $|H|$, which is a contradiction to the choice of H . So $xu_3 \in E(G)$.

Assume that $yu_3 \notin E(P_2)$. Since Q_1P_2 and $Q_1L_1(u_3, x)P'P_2(y, u_4)$ are chordal paths of C_2 that have the same ends as L_2 , they have length $|L_2|$ by Lemma 2.3. Moreover, since $xu_3 \in E(G)$ and $|L_1| \geq \ell + 1$ imply $|L_1(x, u_1)| \geq 2$, the subgraph $(H \cup P') - V(P_2^*(y, u_3))$ is a balanced K_4 -subdivision of type (1, 2) whose edge number is less than $|H|$, which is a contradiction. So $yu_3 \in E(P_2)$, implying that $|P'| \geq 2\ell$. Since $C'_3\Delta C_3\Delta C_1$ is an odd cycle of length at least $2\ell + 3$, we have $|Q_2| = 1$. This proves 3.2.4. \square

3.2.5. $x \notin V(Q_2^*)$.

Subproof. Assume not. Set $C'_3 := Q_2(x, u_4)P_2(u_4, y)P'$. Assume that C'_3 is an odd hole. Since $C_1 \cup C_2 \cup (C_3 \Delta C'_3)$ is an odd K_4 -subdivision when $yu_4 \notin E(P_2)$, we have $yu_4 \in E(P_2)$. Since $|Q_2| < \ell$ as $|L_2| \geq \ell + 1$, the graph $C_4 \Delta C'_3$ is an odd hole with length larger than 3ℓ by (4.1), which is not possible. So C'_3 is an even hole. Since $|Q_2| < \ell$, it follows from Lemma 2.3 that $xu_4 \in E(Q_2)$. Moreover, since P' is a chordal path of the odd hole $C_2 \Delta C_3$ and C'_3 is an even hole, $\ell + 1 \leq |P'| = |P_2(y, u_4)| + 1 \leq |P_2| < |L_1|$ by Lemma 2.3. Hence, $C_2 \cup C_3 \cup P'$ is a balanced K_4 -subdivision of type (1, 2) whose edge number is less than $|H|$, which is a contradiction. \square

Let u'_4 be the neighbour of u_4 in L_2 .

3.2.6. *If $x \in V(L_2)$, then $x \in \{u_4, u'_4\}$. In particular, when $x = u'_4$, we have $yu_4 \in E(P_2)$; and when $x = u_4$, we have $yu_4 \in E(P_2)$ or $|P_2(y, u_4)| \geq \ell + 1$.*

Subproof. Set $C'_3 := L_2(x, u_4)P_2(u_4, y)P'$. Assume that $x = u_4$ and $yu_4 \notin E(P_2)$. Since Q_1P_2 and $Q_1P_2(u_3, y)P'$ are chordal paths of C_2 with the same ends, $|C'_3| = 2|P_2(y, u_4)|$ by Lemma 2.3, so $|P_2(y, u_4)| \geq \ell + 1$. That is, when $x = u_4$, 3.2.6 holds. So we may assume that $x \neq u_4$.

By 3.2.2, we may assume that $x \in V(L_2^*)$. When C'_3 is an odd hole, since $x \notin \{u_2, u_4\}$, the graph $C_2 \cup C_3 \cup P'$ is an odd K_4 -subdivision. So C'_3 is an even hole. When $x \neq u'_4$, by Lemma 2.3, we have that $xu_2 \in E(H)$, $|C'_3| = 2|L_2(x, u_4)|$ and $C_3 \Delta C'_3$ is an odd hole, so $(H \cup P') - V(L_2^*(x, u_4))$ is a balanced K_4 -subdivision of type (1, 2) whose edge number is less than $|H|$, which is not possible. Hence, $x = u'_4$. When $yu_4 \notin E(P_2)$, since $Q_1P_2(u_3, y)P'$ is a chordal path of C_2 , we have $|C'_3| = 2|P_2(y, u_4)|$ by Lemma 2.3. So $(H \cup P') - V(P_2^*(y, u_4))$ is a balanced K_4 -subdivision of type (1, 2) whose edge number is less than $|H|$, which is not possible. So $yu_4 \in E(G)$. \square

Let v be a vertex in $V(P_2^*)$. When $|P_2| \geq \ell + 1$, let $|P_2(u_3, v)| = \ell - 1$, otherwise let $|P_2(u_3, v)| = \ell - 2 \geq 3$. The vertex v is well defined as $|P_2| \geq \ell$. Since $|P_2| \leq 2\ell - 2$, we have $2 \leq |P_2(u_4, v)| \leq \ell - 1$. Let $u, w \in N(v) \cap V(P_2^*)$ with u closer to u_4 and w closer to u_3 along P_2 . Since $\{uv, vw\}$ is not an edge cut of G by Lemma 2.1, there is a shortest induced path Q in $G \setminus \{uv, vw\}$ linking v and $H - v$. By Lemma 3.1 (3), $H \cup Q$ is an induced subgraph of G . Let v' be the other end of Q that is not equal to v . When $v' \in V(C_1 \cup C_2)$, since Q^* is a direct connection in $G \setminus \{e, f\}$ linking P_2^* and $H - V(P_2^*)$, we get a contradiction to 3.2.2-3.2.6. So $v' \in V(P_2^*) - \{v\}$. Assume that $v' \in V(P_2^*(u_3, v))$. When $v' \neq w$, either $Q_2P_1Q_1P_2(u_3, v')QP_2(v, u_4)$ is an odd hole of length at least $2\ell + 3$ or $QP_2(v, v')$ is a cycle of length $2|P_2(v, v')|$ that is at most $2\ell - 2$ by the choice of v , which is not possible. So $v' = w$. By symmetry, we have $v' \in \{u, w\}$.

Now we show that $\{v, v'\}$ is a K_2 -cut of G , implying that Theorem 3.2 holds from Lemma 2.2. Assume not. Let R' be a direct connection in $G - \{v, v'\}$ linking Q^* and $H - \{v, v'\}$. Let s_1, s_2 be the ends of R' with s_1 having a neighbour in $V(H) - \{v, v'\}$ and s_2 having a neighbour in $V(Q^*)$. By Lemma 3.1 (3), s_1 has a unique neighbour, say t_1 , in $V(H) - \{v, v'\}$. Since Q is chosen with $|Q|$ as small as possible, s_2 has a unique neighbour, say t_2 , in $V(Q^*)$. Set $R := t_1 s_1 R' s_2 t_2$. Then $H \cup Q \cup R$ is an induced subgraph of G or some vertex in $\{v, v'\}$ has a neighbour in $V(R')$. When $t_1 \in V(P_2)$, let P'_2 be an induced (u_3, u_4) -path in $P_2 \cup Q \cup R$ that is not equal to P_2 . Since $Q_1 P'_2$ is a chordal path of C_2 with length $|L_2|$ by Lemma 2.3, there is a cycle in $P_2 \cup P'_2$ with length at most 2ℓ by the choice of v , which is not possible. So $t_1 \in V(H) - V(P_2)$. Then $R \cup Q^*$ contains a direct connection in $G \setminus \{e, f\}$ linking P_2^* and $H - V(P_2^*)$. Since $|P_2(u_3, v)| \geq \ell - 2 \geq 3$ and $2 \leq |P_2(u_4, v)| \leq \ell - 1$, by 3.2.2-3.2.6, we have that $v' = u$, $t_1 = u'_4$, and $uu_4 \in E(P_2)$. Let P'_2 be an induced (u_3, u'_4) -path in $(P_2 - \{u_4\}) \cup Q \cup R$. Since $Q_1 P'_2$ and $Q_1 P_2$ are chordal paths of C_2 , by Lemma 2.3, we have $|P'_2(u'_4, v)| = |P_2(u_4, v)| - 1$, so $u'_4 v \in E(G)$, which is not possible. \square

4 Jumps over an odd hole

Let C be an odd hole of a graph G and $s, t \in V(C)$ nonadjacent. Let P be an induced (s, t) -path. If $V(C) \cap V(P^*) = \emptyset$, we call P a *jump* or an (s, t) -*jump* over C . Let Q_1, Q_2 be the internally disjoint (s, t) -paths of C . If some vertex in $V(Q_1^*)$ has a neighbour in $V(P^*)$ and no vertex in $V(Q_2^*)$ has a neighbour in $V(P^*)$, we say that P is a *local jump over C across Q_1^** . When there is no need to strengthen Q_1^* , we will also say that P is a *local jump over C* . In particular, when $|V(Q_1^*)| = 1$, we say that P is a *local jump over C across one vertex*. When no vertex in $V(Q_1^* \cup Q_2^*)$ has a neighbour in $V(P^*)$, we say that P is a *short jump over C* . Hence, short jumps over C are chordal paths of C . But chordal paths over C maybe not short jumps over C as the ends of a chordal path maybe adjacent. When P is a short jump over C , if PQ_1 is an odd hole, we say that PQ_1 is a *jump hole over C* and P is a *short jump over C across Q_1^** . Note that our definition of local jumps over C is a little different from the definition given in [2]. By our definitions, no short jump over C is local.

Lemma 4.1. *Let C be an odd hole of a graph $G \in \mathcal{G}_\ell$. Let P, Q_1, Q_2 be defined as the last paragraph. If P is a local or short jump over C across Q_1^* , then $|P|, |Q_2|$ have the same parity, that is, PQ_2 is an even hole and PQ_1 is an odd cycle.*

Proof. When P is short, the lemma holds from its definition. So we may assume that P is local. Since some vertex in $V(Q_1^*)$ has a neighbour in $V(P^*)$ and $g(G) = 2\ell + 1$, we have $|P| \geq 2\ell$. So PQ_2 is an even hole. This proves Lemma 4.1. \square

Let C be a cycle of a graph G and e, f be chords of C . If no cycle in $C \cup e$ containing e contains the two ends of f , we say e, f are *crossing*; otherwise they are *uncrossing*. Note that by our definition, e, f are uncrossing when they share an end. When C is an odd hole and P_i is an (u_i, v_i) -jump over C for each integer $1 \leq i \leq 2$, if u_1v_1 and u_2v_2 are uncrossing chords of C after we replace P_1 by the edge u_1v_1 and P_2 by the edge u_2v_2 , we say that P_1, P_2 are *uncrossing*; otherwise, they are *crossing*.

Lemma 4.2. *Let C be an odd hole of a graph $G \in \mathcal{G}_\ell$. If a jump P over C is not local, then $G[V(C \cup P)]$ contains a short jump over C .*

Proof. Without loss of generality we may assume that P is not a short jump over C . Let s, t be the ends of P and Q_1, Q_2 be the internally disjoint (s, t) -paths of C . Since P is neither local nor short, there are $u_1, u_2 \in V(P^*)$ such that u_1 has a neighbour in $V(Q_1^*)$ and u_2 has a neighbour in $V(Q_2^*)$, and such that no vertex in $V(P^*(u_1, u_2))$ has a neighbour in $V(C)$. Since no vertex outside an odd hole has two neighbours in the odd hole, $u_1 \neq u_2$ and u_i has a unique neighbour, say w_i , in $V(C)$ for each integer $1 \leq i \leq 2$. Then $w_1u_1P(u_1, u_2)u_2w_2$ is a short jump over C . This proves Lemma 4.2. \square

Lemma 4.3. *Let C be an odd hole of a graph $G \in \mathcal{G}_\ell$. If P is a local (v_1, v_2) -jump over C , then $G[V(C \cup P)]$ contains a local jump over C across one vertex or a short jump over C .*

Proof. Assume not. Among all local jumps over C not satisfying Lemma 4.3, let P be chosen with $|P|$ as small as possible. Let $u_1, u_2 \in V(P^*)$ have a neighbour in $V(C) - \{v_1, v_2\}$ that are closest to v_1, v_2 , respectively. Let w_1, w_2 be the unique neighbour of u_1, u_2 in $V(C)$, respectively. Since $w_1u_1P(u_1, v_1)$ is a short jump over C when $w_1v_1 \notin E(G)$, we have $w_1v_1 \in E(G)$. Similarly, we have $w_2v_2 \in E(G)$. Hence, when $u_1 = u_2$, $w_1 = w_2$ and w_1 is adjacent to v_1, v_2 , which is a contradiction. So $u_1 \neq u_2$. Since P is not a local jump over C across one vertex, we have $w_1 \neq w_2$. Let u_3 be the neighbour of w_1 in $V(P^*)$ closest to v_2 . Note that u_3 maybe equal to u_1 . By the choice of u_2 , we have $u_2 \in V(P(u_3, v_2))$. Set $P' := w_1u_3P(u_3, v_2)$. Since P' is a local jump over C with $|P'| < |P|$, by the choice of P , the subgraph $G[V(C \cup P')]$ contains a local jump over C across one vertex or a short jump over C , so is $G[V(C \cup P)]$, which is a contradiction. This proves Lemma 4.3. \square

Let C be a cycle and suppose that $x_1, x_2, \dots, x_n \in V(C)$ occur on C in this cyclic order with $n \geq 3$. For any two distinct x_i and x_j , the cycle C contains two (x_i, x_j) -paths. Let $C(x_i, x_{i+1}, \dots, x_j)$ denote the (x_i, x_j) -path in C containing x_i, x_{i+1}, \dots, x_j (and not containing x_{j+1} if $i \neq j + 1$), where subscripts are modulo n . Such path is uniquely determined as $n \geq 3$.

Lemma 4.4. *Let C be an odd hole of a graph $G \in \mathcal{G}_\ell$, and P_i be a short (u_i, v_i) -jump over C for each integer $1 \leq i \leq 2$. If P_1, P_2 are crossing, then G contains an odd K_4 -subdivision or a balanced K_4 -subdivision of type $(1, 2)$.*

Proof. Since P_1, P_2 are short jumps over C , either $P_1C(v_1, u_2)P_2C(v_2, u_1)$ or $P_1C(u_1, u_2)P_2C(v_2, v_1)$ has length at most 4ℓ by Lemma 2.3. By symmetry we may assume that the first cycle is shorter than the last one. Set $C' := P_1C(v_1, u_2)P_2C(v_2, u_1)$. Since $g(G) = 2\ell + 1$, either C' is a hole or $|C'| = 4\ell$ and C' has exactly one or two chords. When C' is a hole, implying that $|C(u_1, u_2)|, |C(v_2, v_1)| \geq 2$, the subgraph $C \cup P_1 \cup P_2$ is induced and isomorphic to a balanced K_4 -subdivision of type $(1, 2)$. So we may assume that $|C'| = 4\ell$ and C' has exactly one or two chords. Since $|C'| = 4\ell$ and $g(G) = 2\ell + 1$, all cycles in $G[V(C')]$ using exactly one chord of C' are odd holes. Hence, when C' has exactly two chords, the two chords of C' are crossing and $G[V(C')]$ is isomorphic to an odd K_4 -subdivision; and when C' has exactly one chord, G contains a balanced K_4 -subdivision of type $(1, 2)$. \square

The proofs of Lemma 4.5 and Theorem 4.6 use some ideas in [2].

Lemma 4.5. *Let $\ell \geq 4$ be an integer and C be an odd hole of a graph $G \in \mathcal{G}_\ell$. For any integer $1 \leq i \leq 2$, let P_i be a short or local (u_i, v_i) -jump over C such that $\{u_1, v_1\} \neq \{u_2, v_2\}$ and u_1, u_2, v_2, v_1 appear on C in this order. Assume that P_i is across $C^*(u_i, v_i)$ and if P_i is local, we have $|V(C^*(u_i, v_i))| = 1$ for any integer $1 \leq i \leq 2$. Then the following hold.*

- (1) *When P_1, P_2 are short, G has an odd K_4 -subdivision.*
- (2) *At most two vertices in $V(C(u_1, v_1) \cup C(u_2, v_2))$ are not in a jump hole over C .*

Proof. Without loss of generality we may assume that P_1, P_2 are chosen with $P_1^* \cup P_2^*$ minimal. Set $H := P_1 \cup P_2 \cup C(u_1, u_2) \cup C(v_1, v_2)$. By symmetry we may assume that $|C(u_1, u_2)| \leq |C(v_1, v_2)| \geq 1$. Note that u_1 maybe equal to u_2 .

4.5.1. *(1) is true.*

Subproof. Since P_1, P_2 are short jumps over C , by Lemmas 2.3 and 4.1, $|P_1 \cup P_2| \leq 4\ell - 2$ and $|H|$ is odd and at most $6\ell - 5$. Then $P_1 \cup P_2$ contains at most one cycle. By the choice of P_1 and P_2 , the subgraph $P_1 \cup P_2$ contains no cycle. So H has at most two cycles. When H has exactly two cycles, since $|H| \leq 6\ell - 5$, one cycle in H is a hole, so we can find a new short jump over C to replace one of P_1, P_2 and get a contradiction to the choice of P_1, P_2 . Hence, H contains a unique cycle C' . Since $g(G) = 2\ell + 1$, the cycle C' contains at most two chords, and the two chords are uncrossing when C' has two chords. No matter which case happens, $G[V(P_1 \cup P_2)]$

contains two short jumps Q_1, Q_2 over C such that $C \cup Q_1 \cup Q_2$ is isomorphic to an odd K_4 -subdivision. So (1) holds. \square

By 4.5.1, we may assume that some P_i is local. For any integer $1 \leq i \leq 2$, let a_i, b_i be the vertices in $V(P_i)$ adjacent to u_i, v_i , respectively, and set $D_i := P_i^* - \{a_i, b_i\}$. Since $\ell \geq 3$, both D_1 and D_2 are nonempty by Lemma 2.3. Since $v_1 \neq v_2$ and P_1, P_2 are jumps over C across $C^*(u_1, v_1)$ and $C^*(u_2, v_2)$ respectively, we have that $b_1 \neq b_2$ and they are nonadjacent.

We say two vertex disjoint subgraphs of a graph are *anticomplete* if there are no edges between them.

4.5.2. *Either (2) is true or $D_1 \cup \{b_1\}$ is disjoint and anticomplete to $D_2 \cup \{b_2\}$.*

Subproof. Assume not. Since $b_1 \neq b_2$ and they are nonadjacent, by symmetry we may assume that $G[V(D_1 \cup D_2) \cup \{b_1\}]$ is connected. We claim that P_2 is short. Assume not. Let t be the vertex in $V(C)$ adjacent to u_2, v_2 . Since $D_2 \neq \emptyset$, there is a minimal path Q linking $V(C(u_1, v_1)) - \{u_1\}$ and t with interior in $V(D_1 \cup D_2) \cup \{b_1\}$. Let s be the other end of Q . Since (2) holds when $s \neq v_1$, we have $s = v_1$. When Q is across $V(C^*(v_1, u_1, u_2, t))$, (2) holds; when Q is across $V(C^*(v_1, v_2, t))$, replacing P_2 by Q , we get a contradiction to the choice of P_1, P_2 . Hence, P_2 is short, implying that P_1 is local and $u_1 \neq u_2$, for otherwise (2) holds.

Let Q be a minimal (s, t) -path linking $V(C(u_1, v_1)) - \{u_1\}$ and $\{u_2, v_2\}$ with interior in $V(D_1 \cup P_2^*) \cup \{b_1\}$, with $s \in V(C(u_1, v_1)) - \{u_1\}$ and $t \in \{u_2, v_2\}$. Since no vertex in $V(C) - \{s, t\}$ has a neighbour in $V(Q^*)$, either Q is a short (s, t) -jump over C or $st \in E(C)$. Assume that $st \in E(C)$. Since $1 \leq |C(u_1, u_2)| \leq |C(v_1, v_2)|$, we have $s = v_1, t = v_2$, and $|C(u_1, u_2)| = 1$. Since P_2 is short and P_1 is across one vertex, by Lemmas 2.3 and 4.1, we have $|P_2| = |C(u_2, u_1, v_1, v_2)| = 4 \geq \ell + 1$, so $\ell \leq 3$, which contradicts to the fact that $\ell \geq 4$. Hence, Q is a short (s, t) -jump over C .

When $t = u_2$, since P_1 is local and $1 \leq |C(u_1, u_2)| \leq |C(v_1, v_2)|$, the short jump Q is across $V(C^*(s, u_1, u_2))$, so (2) holds as P_1 is short. When $t = v_2$, either (2) holds or we can replace P_1 by Q to get a contradiction to the choice of P_1, P_2 . This proves 4.5.2 as $t \in \{u_2, v_2\}$. \square

By 4.5.2, we may assume that $D_1 \cup \{b_1\}$ is disjoint and anticomplete to $D_2 \cup \{b_2\}$. By Lemma 4.1, $|H|$ is odd and at least $2\ell + 3$. By symmetry and 4.5.2, we may assume that a_1 has a neighbour in $V(D_2) \cup \{b_2\}$. Let w be the vertex in $N(a_1) \cap (V(D_2) \cup \{b_2\})$ closest to v_2 along P_2 . Set $Q := u_1 a_1 w P_2(w, v_2)$. Then Q is a jump of C . Since (2) holds when Q is short, we may assume that Q is not short, implying that P_2 is local and the vertex adjacent to v_2, u_2 has a neighbour in $V(P_2(w, v_2))$. Then $|Q| \geq 3\ell - 1$, implying that $P_1(a_1, v_1)C(v_1, v_2)Q(v_2, a_1)$ is an even hole by 4.5.2.

Hence, by Lemma 4.1, $C(u_1, v_1, v_2)Q$ is an odd hole of length at least $3\ell + 2$, which is not possible. \square

Theorem 4.6. *Let $\ell \geq 5$ be an integer and C be an odd hole of a graph $G \in \mathcal{G}_\ell$. Then one of the following holds.*

- (1) G has an odd K_4 -subdivision.
- (2) G contains a balanced K_4 -subdivision of type (1, 2).
- (3) G has a P_3 -cut.
- (4) G has a degree-2 vertex.

Proof. Assume that neither (1) nor (2) is true. Set $C := v_1v_2 \dots v_{2\ell+1}v_1$. Let P be a short jump over C with $|P|$ as small as possible. By symmetry we may assume that the ends of P are v_2, v_k with $k \leq \ell + 2$. By Lemmas 4.5 (1), 4.4 and 2.3, we may assume that all short jumps over C have ends in $\{v_2, v_3, \dots, v_k\}$. That is, (i) no jump hole over C contains a vertex in $V(C) - \{v_2, v_3, \dots, v_k\}$.

For each integer $1 \leq i \leq 2$, let P_i be a local (s_i, t_i) -jump over C across one vertex and Q_i be the (s_i, t_i) -path on C of length three. When P_1 and P_2 are uncrossing, by (i) and Lemma 4.5 (2), $|V(Q_1 \cup Q_2) - \{v_2, v_3, \dots, v_k\}| \leq 2$. When P_1 and P_2 are crossing, $|Q_1 \cup Q_2| = 4$. Since there is no short jump over C or at least one vertex in $\{s_i, t_i\}$ is in $\{v_2, v_3, \dots, v_k\}$ by Lemma 4.5 (2), by possibly reordering we may assume that all local jumps over C across one vertex have ends in $\{v_1, v_2, \dots, v_k, v_{k+1}\}$ with $4 \leq k \leq \ell + 2$. For any integer $1 \leq i \leq k + 1$, let X_i be the set of vertices adjacent to v_i that are in a local jump over C across one vertex with one end v_i or a short jump over C with one end v_i . Set $X := X_1 \cup X_2 \cup \dots \cup X_{k+1}$. Then X intersects all short jumps over C and all local jumps over C across one vertex in at least two vertices.

Since $\ell \geq 5$ and $k \leq \ell + 2$, we have $k + 3 \leq 2\ell$. Since no vertex in $V(G) - V(C)$ has two neighbours in $V(C)$, the vertex v_{k+3} has no neighbour in X . Assume that v_{k+3} has degree at least three, for otherwise (4) holds. There is a connected induced subgraph D such that v_{k+3} has a neighbour in $V(D)$ and $V(D) \cap (V(C) \cup X) = \emptyset$, and D is maximal with these properties. Let N be the set of vertices in $V(C) \cup X$ that have a neighbour in $V(D)$. Evidently, $v_{k+3} \in N$.

4.6.1. $N \cap (X \cup V(C)) \subseteq \{v_{k+2}, v_{k+3}, v_{k+4}\}$.

Subproof. Assume not. When $1 \leq i \leq k + 1$, set $W_i := X_i \cup \{v_i\}$; and when $k + 5 \leq i \leq 2\ell + 1$, set $W_i := \{v_i\}$. Assume that $N \cap W_i \neq \emptyset$ for some integer $i \notin \{k + 2, k + 3, k + 4\}$. Let Q be a minimal (v_i, v_{k+3}) -path with interior in $V(D) \cup (W_i - \{v_i\})$. Then Q is a jump over C and $|Q \cap X| \leq 1$. Since X intersects each

local jumps over C across one vertex and each short jump over C in at least two vertices, $G[V(C \cup Q)]$ does not contain a local jump over C across one vertex or a short jump over C , which is not possible by Lemmas 4.2 and 4.3 \square

By 4.6.1, the set $\{v_{k+2}, v_{k+3}, v_{k+4}\}$ is a P_3 -cut of G . So (3) holds. \square

Now, we can prove Theorem 1.3, which is restated here for convenience.

Theorem 4.7. *For any integer $\ell \geq 5$, each graph in \mathcal{G}_ℓ is 3-colorable.*

Proof. Assume not. Let $G \in \mathcal{G}_\ell$ be a minimal counterexample to Theorem 1.3. Then G is 4-vertex-critical. So G has no degree-2 vertex. By Theorems 1.2 and 3.2, G contains neither odd K_4 -subdivision nor balanced K_4 -subdivision of type (1, 2). Hence, G has a P_3 -cut by Theorem 4.6, which is a contradiction to Lemma 2.2. \square

5 Acknowledgments

This research was partially supported by grants from the National Natural Sciences Foundation of China (No. 11971111). The author thanks Yidong Zhou for carefully reading the paper and giving some suggestions.

References

- [1] R. Chen, Y. Zhou, On coloring of graphs with girth $2\ell + 1$ and without longer odd holes. odd K_4 -subdivisions, 2022, in arXiv:2210.12376v1.
- [2] M. Chudnovsky, P. Seymour, Proof of a conjecture of Plummer and Zha, to appear in J. Graph Theory, 2022. in arXiv:2201.11505v2.
- [3] D. Nelson, M. Plummer, N. Robertson, X. Zha, On a conjecture concerning the Petersen graph. Electron. J. Combin., 2011, 18: #P20.
- [4] M. Plummer, X. Zha, On a conjecture concerning the Petersen Graph: Part II. Electron. J. Combin., 2014, 21: #P1.34.
- [5] D. Wu, B. Xu, Y. Xu, On coloring of graphs of girth $2\ell + 1$ without longer odd holes (in Chinese), Sci Sin Math., 2022, 52: 1-18. in arXiv: 2204.06284v1.
- [6] D. Wu, B. Xu, Y. Xu, The chromatic number of heptagraphs, 2022, in arXiv: 2206.01400v1.
- [7] B. Xu, G. Yu, X. Zha, A note on chromatic number and induced odd cycles. Electron. J. Combin., 2017, 24(4): #P4.32.