

HIGH-PLIABILITY FANO HYPERSURFACES

LIVIA CAMPO

ABSTRACT. We show that five of Reid's Fano 3-fold hyperurfaces containing at least one compound Du Val singularity of type cA_n have pliability at least two. The two elements of the pliability set are the singular hypersurface itself, and another non-isomorphic Fano hypersurface of the same degree, embedded in the same weighted projective space, but with different compound Du Val singularities. The birational map between them is the composition of two birational links initiated by blowing up two Type I centres on a codimension 4 Fano 3-fold of $\mathbb{P}^2 \times \mathbb{P}^2$ -type having Picard rank 2.

1. INTRODUCTION

We work over the complex numbers. In recent years, algebraic geometers have gained a new perspective on the birational classification of Fano varieties by looking at them through the lenses of the Minimal Model Program (MMP)[BCHM10]. The strategy is to describe the birational classes of Fano varieties by studying "minimal" members of such families, called *Mori fibre spaces*, that are the outcomes of the MMP. For 3-dimensional Fano varieties, these outcomes are not unique, albeit birational, so it makes sense to understand how they relate to one another using techniques coming from the Sarkisov Program [Cor95, HM13] and variation of Geometric Invariant Theory (vGIT) [Cor00, BZ10, Cam20b]. The set of all the Mori fibre spaces that are birational to a given one has been formalised by Corti in the definition of *pliability* (also in [CR00, Foreword 4.6]).

Definition 1.1 (Definition 1.2 of [CM04]). Given a Mori fibre space $X \rightarrow S$, the *pliability* of X is the set

$$\mathcal{P}(X) := \{\text{Mfs } Y \rightarrow T \mid X \text{ is birational to } Y\} / \text{square equivalence}$$

where two Mori fibre spaces are *square equivalent* if there are two birational maps $X \dashrightarrow Y$ and $S \dashrightarrow T$ such that the maps induced on the generic fibres $X_f \dashrightarrow Y_f$ are biregular.

In this paper we focus on \mathbb{Q} -factorial terminal 3-dimensional Fano varieties (3-folds) having Fano index 1. They can be embedded via Graded Rings methods into suitable weighted projective spaces (see [ABR02, Rei00], and [BK⁺] for the list of Hilbert series associated to terminal Fano 3-folds). When we talk about *codimension* of a Fano 3-fold, we refer to its codimension with respect to its embedding into a weighted projective space $w\mathbb{P}^n$, for some $n > 0$.

Many efforts have been devoted to studying the pliability of Fano 3-folds. For instance, Reid's "famous 95" Fano hypersurfaces [Rei80, IF00] have been proven to have pliability $|\mathcal{P}| = 1$ both in the general quasi-smooth [CPR00] case and for all quasi-smooth members [CP17]. Of the 85 Fano 3-fold codimension 2 complete intersections, 19 have pliability $|\mathcal{P}| = 1$ [IP96, Oka14, AZ16]. In codimension 3, only 3 families have $|\mathcal{P}| = 1$ [AO18]. These are the Fano 3-folds

2020 *Mathematics Subject Classification.* 14E05, 14E30, 14J30, 14J45.

Key words and phrases. Fano 3-fold, Compound Du Val singularities, Pliability.

The author would like to thank Tiago Guerreiro for conversations and comments during the development of this work. The author is kindly supported by Korea Institute for Advanced Study, grant No. MG087901.

for which there is a structure theorem describing their equations (see also [BE74]). Starting from Fano 3-folds in codimension 4, a more sophisticated machinery is needed to retrieve their equations, called *unprojection* [KM83, Pap04, BKR12a], that pinpoints between two and four distinguished deformation families (evocatively called Tom and Jerry) for each Hilbert series (see 2.1 below for details). These have different Euler characteristics; we call *second Tom type* the Tom family with smaller Euler characteristic.

Here we focus on those codimension 4 Fano 3-folds having Picard rank 2 as in [BKQ18], whose equations resemble the equations of the Segre embedding of $\mathbb{P}^2 \times \mathbb{P}^2$: these are realised as 2×2 minors of graded 3×3 matrices, and second Tom type Fano 3-folds always present a $\mathbb{P}^2 \times \mathbb{P}^2$ -type structure. Such $\mathbb{P}^2 \times \mathbb{P}^2$ -type Fano 3-folds are not Mori fibre spaces, but we can run birational links in the same spirit of the Sarkisov Program by performing a toric vGIT of a blow up of $w\mathbb{P}^n$ similarly to [Ahm17, BZ10, Cam20b]. We are most interested in the endpoints of these birational links: in fact, we restrict to those terminating with a Fano 3-fold hypersurface in a weighted \mathbb{P}^4 . The possible hypersurfaces we obtain are listed in Table 1 below. We consider only codimension 4 Fano 3-folds of $\mathbb{P}^2 \times \mathbb{P}^2$ -type and of second Tom type that allow for two distinct birational links both terminating with a Fano hypersurface. We compare the two hypersurfaces thus obtained; they are factorial (see Theorem 4.3 below) and have terminal singularities, both cyclic quotient and compound Du Val [Rei87].

We use this approach to prove the following.

Theorem 1.1. *The 3-dimensional Fano hypersurfaces $\mathcal{N}1, 2, 3, 4, 5$ of Reid's 95 hypersurfaces having compound Du Val singularities as in Table 1 have pliability $|\mathcal{P}| \geq 2$.*

The pliability of quartic Fano 3-folds having one compound Du Val singularity has been studied in [CM04], where the authors prove that the pliability of such singular quartic 3-fold is exactly 2: it contains only another Mori fibre space, the quasi-smooth complete intersection of a cubic and a quartic in a weighted \mathbb{P}^5 .

In this paper we consider more singular Fano hypersurfaces whose singularities arise in the process of running a birational link starting from X of $\mathbb{P}^2 \times \mathbb{P}^2$ -type. In particular, we encounter a singular quartic Fano 3-fold too (i.e. Reid's $\mathcal{N}1$ and GRDB ID #20521, see entry #11125 of Table 1 below): however, our has three compound Du Val singularities instead of one as in [CM04] (one of which is the cA_2 singularity of [CM04]). We still retrieve the complete intersection $Y_{3,4}$ of [CM04] (see Table [Cam20a, Entry #11125]), which is now not quasi-smooth but has compound Du Val singularities. In addition, we find that the pliability set of our singular quartic Fano also contains another, non-isomorphic, quartic Fano 3-fold having different compound Du Val singularities (cf also [Oka18, Oka20]).

This is a recurrent phenomenon in the picture we study in this paper. In fact, the elements in the pliability sets of the singular Fano hypersurfaces we obtain are the hypersurface itself, and the same hypersurface with different compound Du Val singularities.

The main ingredient to prove Theorem 1.1 is understanding the behaviour of the birational links starting from a codimension 4 Fano 3-fold X of $\mathbb{P}^2 \times \mathbb{P}^2$ -type, and how they are affected by X having $\rho_X = 2$. The following theorem summarises Theorems 3.4 and 3.3 below.

Theorem 1.2. *Let X be a codimension 4 Fano 3-fold in second Tom format, and let $p \in X$ be a Type I centre. Suppose that the birational link centred at p terminates with a Fano 3-fold hypersurface X' . Then, the link terminates with two divisorial contractions.*

The fact that $\rho_X = 2$ is reflected on the birational link of (X, p) ; in other words, the birational link initiated by the Kawamata blow-up of a Type I centre p on X a codimension 4 Fano 3-fold having Picard rank $\rho_X = 2$ detects ρ_X by presenting two divisorial contractions. Hence, a birational link constructed in this way and run on a Fano 3-fold of unknown Picard rank ρ can give a lower bound to ρ itself. This is a remark that is used in [Cam20b, Section 6] to deduce the Picard rank of other deformation families of codimension 4 Fano 3-folds.

In Section 2 we set the notation and we introduce the $\mathbb{P}^2 \times \mathbb{P}^2$ formats of [BKQ18]. We then study the birational links starting from a codimension 4 Fano 3-fold of $\mathbb{P}^2 \times \mathbb{P}^2$ -type in Section 3. Finally, in Section 4 we analyse the hypersurfaces that are the endpoints of the birational links, and we discuss their generality under the unprojection assumptions. We conclude with a complete explicit example in Section 5. Below is the table containing the detailed results for each family we considered.

TABLE 1. Hypersurface endpoints

ID of X	Format	Link	ID of X'	$X' = X_d \subset w\mathbb{P}^4$	Singularities
#4839	$T_2 \bullet_{13,45}$	\ggg	#10980	$X_8 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_{y_3}, 2_{y_2}, 4_{x_3})$	$cA_5 @ P_{y_3}$
	$T_5 \bullet_{14}$	\ggg	#10980	$X_8 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_s, 2_{y_4}, 4_t)$	$cA_6 @ P_s$
#4915	$T_2 \bullet_{13,45}$	\ggg	#10981	$X_7 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_{y_3}, 2_{y_2}, 3_{x_3})$	$cA_4 @ P_{y_3}$
	$T_5 \bullet_{14}$	\ggg	#10981	$X_7 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_s, 2_{y_4}, 4_t)$	$cA_6 @ P_s$
#5002	$T_2 \bullet_{13,45}$	\Rightarrow	#16202	$X_6 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_{y_2}, 1_{y_3}, 3_{x_3})$	$cA_5 @ P_{y_2}$ $cA_3 @ P_{y_3}$ $cA_5 @ P := (0, 0, 0, -\frac{1}{2}, 1)$
	$T_4 \bullet_{13}$	\Rightarrow	#16202	$X_6 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_s, 1_{y_4}, 3_t)$	$cA_4 @ P_s$ $cA_5 @ P_{y_4}$ $cA_4 @ P := (0, 0, 0, 2, 1)$
#5163	$T_2 \bullet_{13,45}$	\ggg	#11001	$X_6 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_{y_3}, 2_{y_2}, 2_{x_3})$	$cA_3 @ P_{y_3}$
	$T_5 \bullet_{14}$	\ggg	#11001	$X_6 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_s, 2_{y_4}, 2_t)$	$cA_4 @ P_s$
#5306	$T_2 \bullet_{13,45}$	\Rightarrow	#16203	$X_5 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_{y_2}, 1_{y_3}, 2_{x_3})$	$cA_3 @ P_{y_2}$ $cA_2 @ P_{y_3}$ $cA_3 @ P := (0, 0, 0, 1, -1)$
	$T_4 \bullet_{13}$	\Rightarrow	#16203	$X_5 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_s, 1_{y_4}, 2_t)$	$cA_4 @ P_s$ $cA_3 @ P_{y_4}$ $cA_3 @ P := (0, 0, 0, 1, 1)$
#10985	$T_2 \bullet_{13,45}$	\ggg	#16203	$X_5 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_{y_3}, 1_{x_3}, 2_{y_2})$	$cA_2 @ P_{y_3}$
	$T_5 \bullet_{14}$	\ggg	#16203	$X_5 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_s, 1_{y_4}, 2_t)$	$cA_3 @ P_s$
#11125	$T_2 \bullet_{13,45}$	\Rightarrow	#20521	$X_4 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_{x_3}, 1_{y_2}, 1_{y_3})$	$cA_3 @ P_{y_2}$ $cA_2 @ P_{y_3}$ $cA_3 @ P := (0, 0, 0, 1, -1)$
	$T_4 \bullet_{13}$	\Rightarrow	#20521	$X_4 \subset \mathbb{P}(1_{x_1}, 1_{x_2}, 1_s, 1_{y_4}, 1_t)$	$cA_3 @ P_s$ $cA_2 @ P_{y_4}$ $cA_2 @ P := (0, 0, 0, 1, 1)$

Notation of Table 1. The first and fourth columns, e.g. numbers preceded by #, show the Graded Ring Database IDs of X and the links' endpoints respectively. The second column

records the format of X , e.g. $T_2 \bullet_{13,45}$ means that X is in second Tom_2 format and the entries $m_{13}, m_{45} = 0$ (see Definition 2.1 below). The third column contains information about the shape of the link, that is, the weights d_i of $w\mathbb{P}^7$ are either $d_1 > d_2 > d_3 > d_4$ for $>>>$ or $d_1 > d_2 = d_3 > d_4$ for $>=>$. The fifth column shows the Fano hypersurface endpoint obtained via a link from X in the given Tom format. Lastly, the sixth column reports the type of compound Du Val singularities of the endpoint, and at which point they are, e.g. the coordinate point $P_{y_3} \in X_8 \subset \mathbb{P}^4(1^3, 2, 4)$ is a cA_5 singularity.

2. FANO VARIETIES IN $\mathbb{P}^2 \times \mathbb{P}^2$ TOM FORMAT

2.1. Setting and notation. Codimension 4 Fano 3-folds obtained via Type I unprojections present between two and four formats (at most two Tom and Jerry formats respectively), identifying just as many distinguished deformation families (see [BKR12a, Section 3.4] and [PR04, KM83]). To build a codimension 4 Fano 3-fold X we need the following data.

- A fixed projective plane $D := \mathbb{P}^2(a_{x_1}, b_{x_2}, c_{x_3}) \subset \mathbb{P}^6(a_{x_1}, b_{x_2}, c_{x_3}, d_{1y_1}, d_{2y_2}, d_{3y_3}, d_{4y_4})$ with $d_1 \geq d_2 \geq d_3 \geq d_4$. So D is defined by the ideal $I_D := \langle y_1, y_2, y_3, y_4 \rangle$.
- A family \mathcal{Z}_1 of codimension 3 Fano 3-folds $Z \subset w\mathbb{P}^6$, each defined by maximal pfaffians of a graded skew-symmetric 5×5 syzygy matrix M with weighted entries

$$\begin{pmatrix} m_{1,2} & m_{1,3} & m_{1,4} & m_{1,5} \\ & m_{2,3} & m_{2,4} & m_{2,5} \\ & & m_{3,4} & m_{3,5} \\ & & & m_{4,5} \end{pmatrix}$$

where we omit the principal diagonal, and write only the upper-right triangle.

Each entry of M is occupied by a polynomial in the degree dictated by the grading (all possible gradings are listed in [BKR12b]). The plane D is a divisor inside $Z_1 \in \mathcal{Z}_1$ if the equations of Z_1 are the maximal pfaffians of M in either Tom or Jerry format, and Z_1 is nodal on D . In this paper we will only focus on the deformation families arising as Tom formats.

Definition 2.1 ([BKR12a], Definition 2.2). A 5×5 skew-symmetric matrix M is in Tom_k format if and only if each entry $a_{i,j}$ for $i, j \neq k$ is in the ideal I_D .

If M is in Tom format, we say that

Definition 2.2. A codimension 4 index 1 Fano 3-fold X is of *Tom Type* if it is obtained as Type I unprojection of the codimension 3 pair $Z := (\text{Pf}_i(M))_{i=1}^5 \supset D$ in a Tom family (cf [BKR12a, PR04]). It is said to be *general* if $Z \supset D$ is general in its Tom family. The image of $D \subset Z$ in X is a cyclic quotient singularity $X \ni p \sim \frac{1}{r}(a, b, c)$ called *Type I centre*.

Type I unprojections introduce a new coordinate s of weight r , called *unprojection variable*, and X is now embedded into a 7-dimensional weighted projective space $\mathbb{P}^7(a, b, c, d_1, \dots, d_4, r)$. For $\text{Pf}_i(M)$, $g_j \in \mathbb{C}[x_1, x_2, x_3, y_1, \dots, y_4]$, the equations of X are of the form

$$(\text{Pf}_i(M) = sy_j - g_j = 0, 1 \leq i \leq 5, 1 \leq j \leq 4)$$

When there are two possible Tom formats for M , one of the two (the one with smaller Euler characteristic, which we will often call *second Tom format* for simplicity) is such that some of the Tom constraints cannot be satisfied with the coordinates of $w\mathbb{P}^7$, e.g. a certain entry must be in I_D , but its degree is unattainable. As a consequence, some entries can only be filled with the

zero polynomial. The bullet notation in Table 1 records the zero entries in M , e.g. for M in $T_5 \bullet_{14}$ format, $m_{14} = 0$.

If X is of second Tom type, it has Picard rank $\rho_X = 2$ (see [BKQ18, Proposition 2.1]), and its equations can be written in the $\mathbb{P}^2 \times \mathbb{P}^2$ format of [BKQ18], that is they are the 2×2 minors of a 3×3 graded matrix

$$N := \begin{pmatrix} n_{11} & n_{12} & n_{13} \\ n_{21} & n_{22} & n_{23} \\ n_{31} & n_{32} & n_{33} \end{pmatrix}.$$

Building the matrix M from N consists in performing a Gorenstein projection with respect to $p \in X$ (see [BKQ18, Subsection 3.2] and the example 5 below).

3. BIRATIONAL LINKS

We consider those Fano 3-folds X in second Tom format that have at least two Type I centres whose blow up initiates a birational link terminating in a Fano hypersurface; we will then compare the Fano endpoints. Thus, this restricts the landscape to those Fano 3-folds X sitting inside weighted $w\mathbb{P}^7$ having either $d_1 > d_2 > d_3 > d_4$ or $d_1 > d_2 = d_3 > d_4$ (see [Cam20b, Lemma 4.8]). Each single birational link that we look at will therefore fall in one of these two cases: when $d_1 > d_2 > d_3 > d_4$ we have

$$(3.1) \quad \begin{array}{ccccccc} & & Y_1 & \overset{\Psi_1}{\dashrightarrow} & Y_2 & \overset{\Psi_2}{\dashrightarrow} & Y_3 \\ & \searrow \Phi & & \swarrow \alpha_1 & \swarrow \beta_1 & \swarrow \alpha_2 & \swarrow \beta_2 \\ w\mathbb{P} \supset X & & & Z = Z_1 & & Z_2 & & Z_3 = Y_4 \\ & & & & & & & \searrow \Phi'_1 := \Psi_3 = \alpha_3 \\ & & & & & & & X' \subset w\mathbb{P}' \\ & & & & & & & \searrow \Phi'_2 \end{array}$$

and when $d_1 > d_2 = d_3 > d_4$ we have

$$(3.2) \quad \begin{array}{ccccccc} & & Y_1 & \overset{\Psi_1}{\dashrightarrow} & Y_2 & \overset{\Psi_2}{\dashrightarrow} & Y_3 \\ & \searrow \Phi & & \swarrow \alpha_1 & \swarrow \beta_1 & \swarrow \alpha_2 & \swarrow \beta_2 \\ w\mathbb{P} \supset X & & & Z = Z_1 & & Z_2 & & X' \subset w\mathbb{P}' \\ & & & & & & & \searrow \Phi' \end{array}$$

where Φ' consists of two simultaneous divisorial contractions (see Theorem 3.4). Call \mathbb{F}_i the rank two toric variety that is the ambient space of Y_i .

We now list some theorems regarding the behaviour of the birational links starting from (X, p) . These ultimately justify the diagrams (3.1) and (3.2). Some of these results are proven in [Cam20b], and they hold for any Tom format. Each link is initiated by a Kawamata blow-up of X at its Type I centre p , and continues performing a variation of GIT quotient of the toric ambient space of Y_1 : this is described explicitly in [Cam20b, Section 3].

Theorem 3.1 (Theorem 4.3 of [Cam20b]). *Let n be the number of nodes on $D \subset Z_1$. Then, the birational map $\Psi_1: Y_1 \dashrightarrow Y_2$ is a flop of n smooth rational curves.*

Theorem 3.2 (Theorem 4.5 of [Cam20b]). *Since $d_1 > d_2$, the birational map $\Psi_2: Y_2 \dashrightarrow Y_3$ is a hypersurface flip.*

Proof. Refer to [Cam20b, Theorem 4.5] for the proof of the fact that Ψ_2 is a flip. Here we show that, in the considered cases, Ψ_2 is in particular a hypersurface flip. Note that the coordinate y_1 always occupies the entry $m_{35}, m_{34}, m_{25} \in I_D$ in the formats Tom₂, Tom₅, Tom₄ respectively, and also entry $m_{23} \notin I_D$ in format Tom₅, and entries $m_{35} \in I_D, m_{24}, m_{34} \notin I_D$ in format Tom₄. In the entries having degree d_1 occupied by y_1 , we have ty_1 in the Φ -pull-back. For this reason, t is locally eliminated in a neighbourhood of $P_{y_1} \in Z_2$ when M is either in Tom₄ or Tom₅ format. Instead, since x_3 occupies the entry m_{12} in Tom₂ format (see also (3.3) below), x_3 is locally eliminated when M is in Tom₂ format. Observe that y_4 occupies entry m_{14} in Tom₂, m_{12} in Tom₅, and m_{12} in Tom₄; hence, y_4 is locally eliminated too. Lastly, the unprojection variable s is globally eliminated because the unprojection equations are of the form $sy_i = g_i(t, x_1, x_2, x_3, y_1, y_2, y_3, y_4)$.

This shows that the locus flipped by Ψ_2 is: the weighted plane $\mathbb{P}^2(d_1, a, b)$ cut by the remaining equations of Y_2 (the ones not used to eliminate variables); the weighted plane $\mathbb{P}^2(a, b, c)$ cut by the remaining equations of Y_2 . Thus, these are hypersurface flips. \square

Theorem 3.3. *Suppose that $d_1 > d_2 > d_3 > d_4$ and X is in second Tom format. Then, both $\Phi'_1 := \Psi_3: Y_3 \dashrightarrow Y_4$ and $\Phi'_2: Y_4 \dashrightarrow X'$ are divisorial contractions to a point.*

Proof. Suppose that $d_1 > d_2 > d_3 > d_4$. In this instance and for the families we consider, the matrix M is either in second Tom₂ format (3.3) or is second Tom₅ format (3.4), having grading of the form

$$(3.3) \quad \begin{pmatrix} c & c + d_2 - d_3 & d_4 & d_3 \\ & d_4 & d_3 & d_2 \\ & & d_2 & d_1 \\ & & & d_1 + d_3 - d_4 \end{pmatrix} \quad \text{or} \quad (3.4) \quad \begin{pmatrix} d_4 & d_3 & d_3 & d_2 \\ & d_2 & d_2 & d_1 \\ & & d_1 & c \\ & & & c \end{pmatrix}$$

where the entries with degree written in purple are polynomials not in the ideal I_D . Then, the second Pfaffian $\text{Pf}_2(M)$ for M as in (3.3), or $\text{Pf}_5(M)$ for M as in (3.4), contains the pure monomial y_2y_3 . Thus, the localisation at $P_{y_2} \in Z_3$ is such that the coordinate y_3 can be locally eliminated in a neighbourhood of P_{y_2} . The unprojection variable s is always globally eliminated thanks to the unprojection equations. Moreover, the entry m_{25} in (3.3) and the entry m_{15} in (3.4) are polynomials in which y_2 appears as a pure monomial of degree 1: in the pull-back by Φ it will become ty_2 (see also [Cam20b, Subsection 3.3] for more details). Thus, the localisation $P_{y_2} \in Z_3$ locally eliminates the variable t .

The m_{12} entry of (3.3), and the m_{35} entry of (3.3) must feature x_3 . Therefore, the localisation at P_{y_2} also locally eliminates x_3 via $\text{Pf}_5(M)$ in (3.3) and via $\text{Pf}_1(M)$ in (3.4) (note that $m_{14} = 0$).

Therefore, the contracted locus of the toric flip $\Psi_3: \mathbb{F}_3 \dashrightarrow \mathbb{F}_4$ restricted to Y_3 is the weighted plane $\mathbb{P}^2(a, b, d_1 - d_2)$; this is contracted to $P_{y_2} \in Z_3$, and we call Φ'_1 the restriction $\Psi_3|_{Z_3}$.

The singularities of Y_3 are terminal: they are either isolated cyclic quotient singularities, or isolated compound Du Val singularities. Also, Y_3 is \mathbb{Q} -factorial and its anticanonical divisor is Φ'_1 -ample. The contracted locus is irreducible, and Φ'_1 has relative Picard rank 1. Thus, Φ'_1 is a divisorial contraction to $P_{y_2} \in Z_3$. Proving that Φ'_2 is a divisorial contraction is analogous. \square

Theorem 3.4. *Suppose that $d_1 > d_2 = d_3 > d_4$ and X is in second Tom format. Then, $\Phi': Y_3 \dashrightarrow X'$ consists of two simultaneous divisorial contractions to points.*

Proof. When $d_1 > d_2 = d_3 > d_4$ and for the families we consider, M is either in second Tom₂ format or in second Tom₄ format, whose gradings are as follows

$$(3.5) \quad \begin{pmatrix} c & c & d_4 & d_2 \\ & d_4 & d_2 & d_1 \\ & & d_2 & d_1 \\ & & & d_1 + d_4 - c \end{pmatrix} \quad \text{or} \quad (3.6) \quad \begin{pmatrix} d_4 & d_4 & d_2 & d_2 \\ & d_2 & d_1 & d_1 \\ & & d_1 & d_1 \\ & & & c \end{pmatrix}$$

where the entries with purple degree are not in the ideal I_D .

Torically, Φ' is a divisorial contraction to the curve $\mathbb{P}_{y_2:y_3}^1 \subset w\mathbb{P}'$. However, the generic point $p := (\sigma_1, \sigma_2) \in \mathbb{P}_{y_2:y_3}^1$ for $\sigma_1, \sigma_2 \neq 0$ is not contained in X' . Indeed, both $\text{Pf}_2(M)$ and $\text{Pf}_4(M)$ in (3.5), (3.6) respectively are equal to $y_2y_3 - y_1y_4$, which does not vanish at p . Therefore, the toric contraction restrict to two simultaneous divisorial contractions, one at $P_{y_2} \in X'$ and one at $P_{y_3} \in X'$. \square

Note that Theorems 3.3, 3.4 together with the fact that $\rho_X = 2$ imply that the Picard rank of the endpoint hypersurfaces is 1.

3.1. Birational links through $\mathbb{P}^2 \times \mathbb{P}^2$ formats. Suppose a codimension 4 Fano 3-fold X has two different Type I centres p, q , and that the birational links initiated by blowing up p, q end with two different hypersurfaces X' and X'' . For instance, if the weights of $w\mathbb{P}^7$ are $d_1 > d_2 > d_3 > d_4$ for both centres, we have diagrams like

$$(3.7) \quad \begin{array}{ccccc} & & W_3 \xleftarrow{\Xi_2} W_2 \xleftarrow{\Xi_1} W_1 & & Y_1 \xrightarrow{\Psi_1} Y_2 \xrightarrow{\Psi_2} Y_3 \\ & \Theta'_1 \searrow & & \Theta = \text{Bl}_q \searrow & \Phi = \text{Bl}_p \searrow & & \Phi'_1 \searrow \\ & W_4 & & Z' \dashrightarrow X \dashleftarrow Z & & & Y_4 \\ & \Theta'_2 \searrow & & & & & \Phi'_2 \searrow \\ & X'' & & & & & X' \end{array}$$

where $Z \dashrightarrow X$ is the unprojection from (Z, D) producing X with Type I centre p , and, similarly, $Z' \dashrightarrow X$ unprojects (Z', D') with Type I centre q . Note that Z and Z' are the bases for the flops of Ψ_1 and Ξ_1 respectively.

Our goal is to compare the hypersurface endpoints X', X'' . We prove the following theorem.

Theorem 3.5. *Let X be a codimension 4 Fano 3-fold of second Tom type with at least two Type I centres $p, q \in X$ as in Table 1. The endpoints X', X'' of the two birational links centred at p and q are non-isomorphic 3-dimensional Fano hypersurfaces having the same Hilbert series.*

The proof of Theorem 3.5 is contained in Section 4, as it relies on the analysis of the compound Du Val singularities of X' and X'' carried out in the proof of Theorem 4.1 below.

The heart of the construction is to realise (X, p) via Type I unprojection of (Z, D) where $Z = (\text{Pf}_i(M) = 0)_{i=1}^5$, and to then retrieve the defining matrix M' of (Z', D') by passing through the $\mathbb{P}^2 \times \mathbb{P}^2$ construction of N . Then, using the new pair (Z', D') we perform the Type I unprojection to obtain (X, q) . Note that the equations of (X, p) and (X, q) are different, but they are two general members of the same Tom deformation family.

Example 3.1. For instance, looking at Table 1 and in comparison with [BKR12b, Cam20a], consider #4839; the two Tom formats $\text{Tom}_2 \bullet_{13,45}$ and $\text{Tom}_5 \bullet_{14}$ represent the same deformation family with Hilbert series #4839, but with respect to the two centres $p \sim \frac{1}{5}(1, 1, 4)$ and $q \sim \frac{1}{9}(1, 1, 8)$. Thus, we have M in $\text{Tom}_2 \bullet_{13,45}$, M' in $\text{Tom}_5 \bullet_{14}$, and N as in [BKQ18, Table 2].

Consider the Type I centre $p \in X \subset \mathbb{P}^7(1_{x_1}, 1_{x_2}, 4_{x_3}, 5_s, 6_{y_4}, 7_{y_3}, 8_{y_2}, 9_{y_1})$ at the coordinate point P_s having local coordinates x_1, x_2, x_3 . The unprojection that produced such Type I centre started from (Z, D) for $D = \mathbb{P}^2(1_{x_1}, 1_{x_2}, 4_{x_3})$ inside $Z = (\text{Pf}_i(M) = 0)_{i=1}^5$ where $I_D := \langle y_1, y_2, y_3, y_4 \rangle$ and the grading of M is below (3.8). The circled weights in (3.8) mark the zero entries, and in purple are the entries that must be filled with polynomials not in I_D as before. The matrix N is built by ignoring the zero entries of M and by placing the unprojection variable s in the entry n_{31} of N (in a square box below).

Call N' the 3×3 matrix where the second and third row of N have been swapped. The $n'_{33} = n_{23}$ entry of N' (boxed) is therefore occupied by y_1 , which is the unprojection variable when the Type I centre is $q \sim \frac{1}{9}(1, 1, 8)$ having local coordinates x_1, x_2, y_2 . Projecting away from q we retrieve $D' = \mathbb{P}^2(1_{x_1}, 1_{x_2}, 8_{y_2})$ sitting inside $Z' = (\text{Pf}_i(M') = 0)_{i=1}^5$ where $I_{D'} := \langle x_3, s, y_3, y_4 \rangle$ and M' is in $\text{Tom}_{5 \bullet 14}$ format (entries not in $I_{D'}$ marked in blue).

$$(3.8) \quad N: \begin{pmatrix} 4 & 6 & 7 \\ 6 & 8 & 9 \\ \boxed{5} & 7 & 8 \end{pmatrix} \rightsquigarrow N': \begin{pmatrix} 4 & 6 & 7 \\ 5 & 7 & 8 \\ 6 & 8 & \boxed{9} \end{pmatrix}$$

Gorenstein p -projection ↙

$p \sim \frac{1}{5}(1, 1, 4)$

$$M: \begin{pmatrix} 4 & \textcircled{5} & 6 & 7 \\ & 6 & 7 & 8 \\ & & 8 & 9 \\ & & & \textcircled{10} \end{pmatrix}$$

↘ Gorenstein q -projection

$q \sim \frac{1}{9}(1, 1, 8)$

$$M': \begin{pmatrix} 4 & 5 & \textcircled{5} & 6 \\ & 6 & 6 & 7 \\ & & 7 & 7 \\ & & & 8 \end{pmatrix}$$

The codimension 4 Fano 3-fold X realised as Type I unprojection of (Z, D) lies in the same deformation family of the one realised as Type I unprojection of (Z', D') . We run the two birational links in (3.7) by blowing up p and q , and using the two sets of equations for X that we have just built using $\mathbb{P}^2 \times \mathbb{P}^2$ formats. We will give a detailed account on how to do so in the continuation of this explicit example in Section 5.

4. VERY SINGULAR FANO HYPERSURFACES

The endpoints of the birational links we have constructed are Fano hypersurfaces in weighted $w\mathbb{P}^4$. Here we want to study their singularities, and discuss the generality of these hypersurfaces. Recall that, by Theorem 3.2, the birational links in this paper all present a hypersurface flip. This introduces a compound Du Val singularity on Y_3 induced by the local equation in the flipped locus of Ψ_2 . Consequently, X' has at least one compound Du Val singularity (possibly more, depending on whether the contracted locus of the divisorial contractions has a local equation too).

Theorem 4.1. *Let X be a codimension 4 Fano 3-fold of second Tom type, and let X be general in its Tom format. Assume that the birational link initiated by the Kawamata blowup of one of X 's Type I centres terminates with a Fano hypersurface X' . Then, the compound Du Val singularities of X' are constant when X varies in its general Tom family.*

Proof. Suppose the Type I centres are $p, q \in X$ as before. For simplicity, we focus on the link initiated by blowing up p ; the proof for the link corresponding to blowing up q is analogous. Since we consider only weights $d_1 > d_2 > d_3 > d_4$ and $d_1 > d_2 = d_3 > d_4$, the second Tom formats we have are $\text{Tom}_2, \text{Tom}_5$, or Tom_4 . Firstly, consider the case in which $d_1 > d_2 > d_3 > d_4$.

To fix ideas, suppose that $p \in X$ is the cyclic quotient singularity with lowest index in the basket of X . So, by [BKR12b], M is in $\text{Tom}_2 \bullet_{13,45}$ format as follows

$$\begin{pmatrix} x_3 & 0 & y_4 & y_3 \\ & p_1 & p_2 & p_3 \\ & & y_2 & y_1 \\ & & & 0 \end{pmatrix}$$

for $p_1, p_2, p_3 \notin I_D$ homogeneous polynomials of prescribed degree as in (3.3).

The equations of X' can be found by imposing $y_4 = 1$ (e.g. blowing down) in the equations of Y_4 (or Y_3 when $Y_4 = Y_3$); some coordinates (like s via the unprojection equations) can be globally eliminated, and can be substituted in the rest of the equations of $Y_4|_{y_4=1}$. After the elimination of s , we are left with only the five pfaffian equations. In the cases we consider, the coordinates t and y_1 can always be globally eliminated by $\text{Pf}_5(M)$ and $\text{Pf}_2(M)$ respectively for M is in Tom_2 format. The fifth pfaffian equation is $\text{Pf}_5(M) = x_3y_2 - 0 \cdot p_2 + y_4p_1$. Here $p_1 \notin I_D$ is a homogeneous polynomial of degree d_4 of the form $p_1 = y_4 + x_1^{d_4} + h_1$, where h_1 is a homogeneous polynomial of degree d_4 , possibly in I_D (also recall that all these second Tom type Fano 3-folds have Fano index 1, so we can assume that the weight of x_1 is 1). The only monomials in h_1 that contribute to achieving quasi-smoothness of X and reducedness of $D \subset Z$ are the pure monomials in x_1, x_2, x_3 , and quasi-linear terms in x_1, x_2, x_3 (see [Cam20b]); indeed, monomials that are in terms of three or more variables vanish when in the restriction to D when checking reducedness and quasi-smoothness. Thus, we can assume without loss of generality that h_1 does not contain other monomials in I_D . As a consequence, in the pull-back of the pfaffian equations via Φ , the only term in $\text{Pf}_5(M)$ picking up a t factor is y_4 appearing in p_1 . Hence, the elimination of t is global. If we look at the second pfaffian instead, we have that $\text{Pf}_2(M) = y_2y_3 - y_1y_4$; here the global elimination is evident.

This leaves us with three pfaffian equations: $\text{Pf}_1(M)$, $\text{Pf}_3(M)$, and $\text{Pf}_4(M)$. After replacing the global expressions of t, y_1 , and since the equations are explicit, we can see that $\text{Pf}_4(M)$ is always identically 0, and that $\text{Pf}_1(M)$ is a y_2 -multiple of $\text{Pf}_3(M)$. Thus, the ideal of $\text{Pf}_1(M), \text{Pf}_3(M), \text{Pf}_4(M)$ is actually only generated by $\text{Pf}_3(M)$; this is the equation of the Fano (cf [Cam20b, Lemma 4.8]) hypersurface X' . Since $\text{Pf}_3(M) = 0$ is an explicit equation, we can run a simple Magma routine to find out where the compound Du Val singularities are on X' : it is enough to find the singular locus and its prime components. For $d_1 > d_2 > d_3 > d_4$ and M in Tom_2 format, then there is one compound Du Val singularity on X' at the coordinate point $P_{y_3} \in X' \subset w\mathbb{P}^4$. In order to understand what type of compound Du Val singularity there is at P_{y_3} , we look at $\text{Pf}_3(M) = 0$ after replacing t and y_1 with their local expressions. So, to $\text{Pf}_3(M) = y_3p_2(t, y_3, x_1, x_2, x_3) - y_4p_3(t, y_2, x_1, x_2, x_3)$ we need to impose: $y_4 = 1$ to blow down; $y_3 = 1$ to localise at the singular point; $t = x_1^{d_4} + h_1$ for the global elimination of t . Since $p_2 = y_3 + x_1^{d_3} + h_2$, the pullback via Φ and the localisation at P_{y_3} imply that $\text{Pf}_3(M) = t + x_1^{d_3} + h_2(t, y_3, x_1, x_2, x_3) - p_3(t, y_2, x_1, x_2, x_3)$. Thus, recalling the form of $\text{Pf}_5(M), h_1$ the equation of X' contains the monomials: $x_3y_2, x_1^{d_4}$ (taking the smallest pure power of x_1), and x_3^2 (that appears in either p_2 or p_3 if $d_3, d_4 = 2 \cdot \text{wts}(x_3)$). Note that, when $d_3, d_4 = 2 \cdot \text{wts}(x_3)$ does not happen, we can still perform a local change of variables in a neighbourhood of P_{y_3} to get two quadratic terms starting from x_3y_2 . Locally at P_{y_3} the equation of X' is of the form $x^2 + xy + z^n$ plus higher order terms, which, by a change of variables, becomes the standard $x^2 + y^2 + z^n$ that expresses a compound Du Val singularity of type cA_{n-1} .

The case of $d_1 > d_2 = d_3 > d_4$ is similar: the local expressions of t, y_1 are analogous to the case above, and $\text{Pf}_3(M)$ is still the equation of X' . What changes is that the compound Du Val singularities are three: two at the coordinate points P_{y_2}, P_{y_3} , and one at a point P indicated explicitly for each format in the right-most column of Table 1. The singularity analysis is conducted separately at each point in a similar fashion as above (with a simple change of variables to make P a coordinate point).

In conclusion, the only constraints that we imposed to the generality of these cA_n singularities are the following: the Tom constraints, the reducedness of $D \subset Z$, the quasi-smoothness of X . Thus, we conclude that the cA_n singularities on X' persist when X varies in its Tom family. \square

Remark 4.2. Note that there is possibly another compound Du Val singularity at orbifold points when $d_1 > d_2 > d_3 > d_4$ (more precisely, at the coordinate point P_{y_2}). Even though we do not study their singularity type here, their existence does not affect the statement of Theorem 3.5. However, this could lead to yet another Mori fibre space birational to X' . Thus, the inequality in Theorem 1.1 about the size of the pliability set of X' could be strict.

Proof of Theorem 3.5. The link initiated by $q \in X$ starts from X in either Tom_5 or Tom_4 type. For simplicity we assume that M' is in Tom_5 format; the proof for M' is in Tom_4 format is analogous.

Recall from the proof of Theorem 4.1 that the coordinate point $P_{y_3} \in X'$ has local equation $x^2 + y^2 + z^n$ where $n = d_4$; so, P_{y_3} is a cA_{d_4-1} singularity. Moreover, since the coordinates t, s, y_1 can be globally eliminated and y_4 has been set to 1, X' sits inside the weighed projective space $\mathbb{P}^4(1_{x_1}, b_{x_2}, c_{x_3}, (d_2 - d_4)_{y_2}, (d_3 - d_4)_{y_3})$ (with a little abuse of notation we still use the x_i, y_j coordinate notation here). The proof of Theorem 4.1 can be conducted similarly when considering the birational link initiated by blowing up the other Type I centre $q \in X$. In this case, the globally eliminated coordinates are y_1, y_2, y_3 (where y_1 is the new unprojection variable coming from the Type I centre q), so the Fano hypersurface X'' is embedded in the weighed projective space $\mathbb{P}^4(1_{x_1}, b_{x_2}, c_t, (r - c)_s, (d_4 - c)_{y_4})$. As it was for $P_{y_3} \in X'$, the coordinate point $P_s \in X''$ is always a compound Du Val singularity with local equation of the form $x^2 + y^2 + z^{n'}$. The equation of X'' is given by $\text{Pf}_4(M')|_{x_3=1} = p_3 - sp_2 + y_4p_1$, which contains the monomial $x_1^{d_2}$ coming from p_2 . Note that $x_1^{d_2}$ is the smallest pure power of x_1 appearing in $\text{Pf}_4(M')|_{x_3=1}$, because p_3 features $x_1^{d_1}$, and $d_1 > d_2$. Thus, $n' = d_2$ and P_s is a compound Du Val singularity of Type cA_{d_2-1} .

Note that the weighted projective spaces where X' and X'' are embedded have the same weights: indeed, in the cases we consider, $d_3 - d_4 = r - c = 1$, and $d_2 - d_4 = d_4 - c$.

The fourth pfaffian of M' is $\text{Pf}_4(M') = x_3p_3 - sp_2 + y_4p_1$, which is a homogeneous polynomial of degree $c + d_2$. The equation of X'' is given by $\text{Pf}_4(M')|_{x_3=1}$, which has therefore degree d_2 and is homogeneous in $\mathbb{P}^4(1_{x_1}, b_{x_2}, c_t, (r - c)_s, (d_4 - c)_{y_4})$.

On the other hand, the third pfaffian of M is $\text{Pf}_3(M) = y_3p_2 - y_4p_3$, which is homogeneous of degree $d_4 + d_2$. The equation of X' is $\text{Pf}_3(M)|_{y_4=1}$, whose degree is then d_2 , and is homogeneous in $\mathbb{P}^4(1_{x_1}, b_{x_2}, c_{x_3}, (d_2 - d_4)_{y_2}, (d_3 - d_4)_{y_3})$.

Thus, X' and X'' are two Fano hypersurfaces having the same degree and sitting inside the same weighted \mathbb{P}^4 . As a consequence, they have the same Hilbert series as in [BK⁺]. However, they are not isomorphic because they have different compound Du Val singularities. \square

Now we have all the elements to prove our Main Theorem 1.1.

Proof of Theorem 1.1. In the cases we consider, the square-equivalence classes in the pliability definition reduce to isomorphism classes because $S = T = \{\text{point}\}$. Theorem 3.5 shows that the two birational links starting from (X, p) and (X, q) respectively terminate with two birational non-isomorphic Fano hypersurfaces X', X'' having the same Hilbert series but different compound Du Val singularities. Hence, the pliability of X' contains at least X'' , so $|\mathcal{P}(X')| \geq 2$. \square

Lastly, we discuss the factoriality of the hypersurfaces we obtained. Factoriality is crucial for understanding how many compound Du Val singularities (including ordinary double points) a hypersurface can have in order to still have pliability $|\mathcal{P}| = 1$. This is in relation to Conjecture 1.4 of [CM04, extended version] and [Che05, Che10, CP06, AK16]. Factoriality is necessary, but not sufficient, to prove the birational rigidity (i.e. $|\mathcal{P}| = 1$) of a hypersurface; in fact, the singular quartic 3-fold in [CM04] is factorial but has $|\mathcal{P}| = 2$. The trend is that factoriality combined with a bounded number of ordinary double points gives birational rigidity. What we show is that all our very singular Fano hypersurfaces are factorial and yet are non-rigid. In our case, the factoriality property still derives from the second Tom formats of M, M' , and therefore from the shape of the equations of Z, Z' . We check this in the next theorem.

Theorem 4.3. *The Fano 3-fold hypersurfaces X' and X'' obtained as above are factorial.*

Proof. We refer to the proofs of Theorems 3.5 and 4.1 for the notation and the equations of X' and X'' . For simplicity, we focus on X' ; the proof for the hypersurface endpoint X'' is analogous. To prove factoriality of X' it is enough to show that they do not contain reducible curves such as $(x = y = 0) \cap X'$ for some coordinates x, y of $w\mathbb{P}^4$. We look at the equation of X' , which is given by $\text{Pf}_3(M) = 0$ after evaluating at $y_4 = 1$ and replacing the expressions of the global elimination of t, y_1 . Recall that the polynomial p_3 contains $x_1^{d_2}$ without loss of generality. Therefore, whenever we cut X' by $(x = y = 0)$ with both $x, y \neq x_1$, X' does not contain $(x = y = 0)$. Suppose instead that $x = x_1$; we have two possibilities. If $y = x_2$ or x_3 , since the global expression of t comes from $\text{Pf}_5(M)$, we have that at least one of the monomials $y_3^2 x_2^m, y_3^2 x_2^{m'}$ appear in $\text{Pf}_3(M)$ for some m, m' . Thus, X' does not contain $(x = y = 0)$. On the other hand, if $y = y_i$ for $i \in \{1, \dots, 4\}$, then p_3 contains at least one of $x_2^m, x_3^{m'}$ for $m := d_2 - \text{wts}(x_2)$ and $m' := d_2 - \text{wts}(x_3)$; so, X' does not contain $(x = y = 0)$. \square

5. EXAMPLE

We run the explicit computation of the matrices M, M', N, N' for the family X with Hilbert series #4839 expanding from Example 3.1 above. Moreover, we study the two links obtained by blowing up X at its two Type I centres $p \sim \frac{1}{5}(1, 1, 4)$ and $q \sim \frac{1}{9}(1, 1, 8)$; this produces a diagram as in (3.7). Then, we find the explicit equations of the endpoints X, X'' and we look at their compound Du Val singularities, following the proof of Theorem 4.1.

Continuing Example 3.1 above, the 5×5 skew-symmetric matrix M is

$$M := \begin{pmatrix} x_3 & 0 & y_4 & y_3 & \\ & p_1 & p_2 & p_3 & \\ & & y_2 & y_1 & \\ & & & & 0 \end{pmatrix} \quad \text{with grading} \quad \begin{pmatrix} 4 & \textcircled{5} & 6 & 7 & \\ & 6 & 7 & 8 & \\ & & 8 & 9 & \\ & & & & \textcircled{10} \end{pmatrix}$$

where $p_{5-i} = x_1^{d_i} + x_2^{d_i} + y_i + h_{5-i}(x_1, x_2, x_3)$ and h_{5-i} is a homogeneous polynomial of degree d_{5-i} , $i = 2, 3, 4$. In fact, we only need at least one of the x_i to appear as a pure power in the

polynomials p_j . The $\mathbb{P}^2 \times \mathbb{P}^2$ format N for (X, p) is therefore

$$N := \begin{pmatrix} x_3 & y_4 & y_3 \\ p_1 & y_2 & y_1 \\ s & p_2 & p_3 \end{pmatrix} \quad \text{with grading} \quad \begin{pmatrix} 4 & 6 & 7 \\ 6 & 8 & 9 \\ \boxed{5} & 7 & 8 \end{pmatrix}.$$

The other $\mathbb{P}^2 \times \mathbb{P}^2$ format N' , corresponding to (X, q) , is the matrix obtained by swapping the second and third row of N . With a Gorenstein projection from $q \in X$ we get

$$M' := \begin{pmatrix} x_3 & s & 0 & p_1 \\ & y_4 & y_4 & p_2 \\ & & y_3 & p_3 \\ & & & y_2 \end{pmatrix} \quad \text{with grading} \quad \begin{pmatrix} 4 & 5 & \textcircled{5} & 6 \\ & 6 & 6 & 7 \\ & & 7 & 7 \\ & & & 8 \end{pmatrix}.$$

To perform the blow-ups of X at p and q , we follow [Cam20b, Section 3.4]; thus, the proper transforms (after saturation by a t factor) Y_1 and W_1 of X via such blow-ups are respectively embedded into the rank 2 toric varieties

$$\mathbb{F}_1 := \left(t \ s \left| \begin{array}{cccccccc} x_1 & x_2 & x_3 & y_1 & y_2 & y_3 & y_4 \\ 0 & 5 & 1 & 1 & 4 & 9 & 8 & 7 & 6 \\ 1 & 1 & 0 & 0 & 0 & -1 & -1 & -1 & -1 \end{array} \right. \right), \quad \mathbb{F}'_1 := \left(t \ y_1 \left| \begin{array}{cccccccc} x_1 & x_2 & y_2 & y_3 & y_4 & s & x_3 \\ 0 & 9 & 1 & 1 & 8 & 7 & 6 & 5 & 4 \\ 1 & 1 & 0 & 0 & 0 & -1 & -1 & -1 & -1 \end{array} \right. \right).$$

For instance, for $p_1 = -x_1^6 + x_1^5 x_2 + x_2^6 + y_4$, $p_2 = x_1^6 x_2 - x_2^7 - y_3$, $p_3 = -x_1^4 x_3 - x_3^2 - y_2$ the equations of Y_1 and W_1 are

$$\left\{ \begin{array}{l} -tx_3y_3 - sy_4 + x_1^6x_2x_3 - x_2^7x_3 = 0 \\ ty_4^2 + x_1^6y_4 - x_1^5x_2y_4 - x_2^6y_4 - x_3y_2 = 0 \\ t^2y_4^2 + tx_1^6y_4 - tx_1^5x_2y_4 - tx_2^6y_4 + sy_3 \\ + x_1^4x_3^2 + x_3^3 = 0 \\ ty_3y_4 + x_1^6y_3 - x_1^5x_2y_3 - x_2^6y_3 - x_3y_1 = 0 \\ -t^2y_3y_4 + tx_1^6x_2y_4 - tx_1^6y_3 + tx_1^5x_2y_3 \\ -tx_2^7y_4 + tx_2^6y_3 - sy_2 + x_1^{12}x_2 - x_1^{11}x_2^2 \\ -2x_1^6x_2^7 + x_1^5x_2^8 + x_2^{13} = 0 \\ -tx_2y_3y_4 + ty_2y_4 - ty_3^2 + x_1^5x_2^2y_3 + x_1^4x_3y_4 \\ + x_2x_3y_1 + x_3^2y_4 = 0 \\ t^2y_2y_4 + tx_1^6y_2 - tx_1^5x_2y_2 + tx_1^4x_3y_4 - tx_2^6y_2 \\ + tx_2^2y_4 + sy_1 + x_1^{10}x_3 - x_1^9x_2x_3 + x_1^6x_3^2 \\ -x_1^5x_2x_3^2 - x_1^4x_2^6x_3 - x_2^6x_3^2 = 0 \\ -y_1y_4 + y_2y_3 = 0 \\ -ty_1y_3 + ty_2^2 + x_1^6x_2y_1 + x_1^4x_3y_2 \\ -x_2^7y_1 + x_3^2y_2 = 0 \end{array} \right. \quad \left\{ \begin{array}{l} -y_3x_3 + y_4s = 0 \\ ty_4^2 + x_1^6y_4 - x_1^5x_2y_4 - x_2^6y_4 - y_2x_3 = 0 \\ t^2x_3^3 + tx_1^4x_3^2 - ty_3s + x_1^6x_2s - x_2^7s + 2y_2x_3 = 0 \\ ty_3y_4 + x_1^6y_3 - x_1^5x_2y_3 - x_2^6y_3 - y_2s = 0 \\ t^2y_3y_4 - tx_1^6x_2y_4 + tx_1^6y_3 - tx_1^5x_2y_3 + tx_2^7y_4 \\ -tx_2^6y_3 + y_1x_3 - x_1^{12}x_2 + x_1^{11}x_2^2 \\ + 2x_1^6x_2^7 - x_1^5x_2^8 - x_2^{13} = 0 \\ t^2y_4x_3^2 + tx_1^4y_4x_3 - tx_2y_3y_4 \\ -ty_3^2 + x_1^5x_2^2y_3 + x_2y_2s + 2y_2y_4 = 0 \\ -t^2x_1^6x_3^2 + t^2x_1^5x_2x_3^2 + t^2x_2^6x_3^2 - t^2y_3^2 - tx_1^{10}x_3 \\ + tx_1^9x_2x_3 + tx_1^6x_2y_3 + tx_1^4x_2^6x_3 - tx_2^7y_3 \\ -y_1s - 2x_1^6y_2 + 2x_1^5x_2y_2 + 2x_2^6y_2 = 0 \\ ty_2y_3 + y_1y_4 - x_1^6x_2y_2 + x_2^7y_2 = 0 \\ -t^3y_4^2x_3 - t^2x_1^6y_4x_3 + t^2x_1^5x_2y_4x_3 \\ -t^2x_1^4y_4^2 + t^2x_2^6y_4x_3 - tx_1^{10}y_4 \\ + tx_1^9x_2y_4 + tx_1^4x_2^6y_4 - y_1y_3 - 2y_2^2 = 0 \end{array} \right.$$

respectively. The birational links in (3.7) are obtained by variation of GIT quotient on $\mathbb{F}_1, \mathbb{F}'_1$ respectively, and by intersecting the contracted loci of the maps Ψ_i, Ξ_i with Y_1, W_1 and their proper transforms (cf [Cam20b, Cam20a] for a more detailed description). Then, Ψ_1 is a flop of five rational curves and Ψ_2 is a hypersurface flip with weights $(9, 1, 1, -1, -2; 5)$; on the other

hand, Ξ_1 flops nine rational curves and Ξ_2 is a hypersurface flip with weights $(1, 1, 8, -1, -2; 6)$. Both Ψ_2 and Ξ_2 are followed by two consecutive divisorial contractions to points.

We carry out the explicit calculations to find the equation of X' ; the one for X'' is analogous.

The last divisorial contraction Φ'_2 is a blow-down to a point in $\mathbb{P}^7(6_t, 11_s, 1_{x_1}, 1_{x_2}, 4_{x_3}, 3_{y_1}, 2_{y_2}, 1_{y_3})$ after setting $y_4 = 1$. Looking at the equations of Y_4 (that are the same as Y_1), we see that the coordinates s, t, y_1 can be globally eliminated respectively by the unprojection equations, $\text{Pf}_5(M), \text{Pf}_2(M)$. In particular, the (global) expression of t, y_1 is $t = -x_1^6 + x_1^5 x_2 + x_2^6 + x_3 y_2$, $y_1 = y_2 y_3$. This leaves us with three remaining equations: $\text{Pf}_1(M), \text{Pf}_3(M), \text{Pf}_4(M) = 0$ evaluated at the global expressions of t, y_1 . It is easy to see that $\text{Pf}_4(M)$ is now identically zero, and that $\text{Pf}_1(M) = y_2 \text{Pf}_3(M)$. Thus, the ideal $\langle \text{Pf}_1(M), \text{Pf}_3(M), \text{Pf}_4(M) \rangle$ is actually generated only by $\text{Pf}_3(M)$. The equation of X' is then $\text{Pf}_3(M) = 0$ after setting $y_4 = 1$ and replacing the global expressions of t, y_1 . Hence, the degree 8 equation of $X' \subset \mathbb{P}^4(1_{x_1}, 1_{x_2}, 1_{y_3}, 2_{y_2}, 4_{x_3})$ is

$$X' = \{x_1^6 x_2 y_3 - x_2^7 y_3 - x_1^6 y_2 + x_1^5 x_2 y_2 + x_2^6 y_2 + x_3 y_2^2 + x_1^6 y_3^2 - x_1^5 x_2 y_3^2 - x_2^6 y_3^2 - x_3 y_2 y_3^2 + x_1^4 x_3 + x_3^2 = 0\}.$$

Analogously, the equation of $X'' \subset \mathbb{P}^4(1_{x_1}, 1_{x_2}, 1_s, 2_{y_4}, 4_t)$ is

$$X'' = \{t^2 + t x_1^4 - t y_4 s^2 + x_1^6 x_2 s - x_2^7 s + 2 t y_4^2 + 2 x_1^6 y_4 - 2 x_1^5 x_2 y_4 - 2 x_2^6 y_4 = 0\}.$$

The compound Du Val singularities of X' lie at the coordinate points P_{y_3} and P_{y_2} ; the latter is also a cyclic quotient singularity, and we will only focus on the former. To see which kind of compound Du Val singularity P_{y_3} is, we need to localise, i.e. set $y_3 = 1$. We see that the lowest-power monomials are $x_3^2 + x_3 y_2 + x_1^6$, so P_{y_3} is a cA_5 singularity.

Instead, the compound Du Val singularities of X'' lie at P_s and P_{y_4} (which is an orbifold point). Localising at $s = 1$ we obtain the local equation $t^2 + t y_4 + x_2^7$, which is a cA_6 singularity.

In conclusion we have just built two non-isomorphic X_8 hypersurfaces lying in the same birational class.

Declarations. This work was supported by Korea Institute for Advanced Study, grant No. MG087901. The author has no relevant financial or non-financial interests to disclose.

REFERENCES

- [ABR02] S. Altınok, G. Brown, and M. Reid. Fano 3-folds, $K3$ surfaces and graded rings. In *Topology and geometry: commemorating SISTAG*, volume 314 of *Contemp. Math.*, pages 25–53. Amer. Math. Soc., Providence, RI, 2002.
- [Ahm17] H. Ahmadinezhad. On pliability of del Pezzo fibrations and Cox rings. *J. Reine Angew. Math.*, 723:101–125, 2017.
- [AK16] H. Ahmadinezhad and A.-S. Kaloghiros. Non-rigid quartic 3-folds. *Compos. Math.*, 152(5):955–983, 2016.
- [AO18] H. Ahmadinezhad and T. Okada. Birationally rigid Pfaffian Fano 3-folds. *Algebr. Geom.*, 5(2):160–199, 2018.
- [AZ16] H. Ahmadinezhad and F. Zucconi. Mori dream spaces and birational rigidity of Fano 3-folds. *Adv. Math.*, 292:410–445, 2016.
- [BCHM10] C. Birkar, P. Cascini, C. D. Hacon, and J. McKernan. Existence of minimal models for varieties of log general type. *J. Amer. Math. Soc.*, 23(2):405–468, 2010.
- [BE74] D. A. Buchsbaum and D. Eisenbud. Some structure theorems for finite free resolutions. *Advances in Math.*, 12:84–139, 1974.
- [BK⁺] G. Brown, A. M. Kasprzyk, et al. Graded Ring Database. *Online. Access via <http://www.grdb.co.uk>.*
- [BKQ18] G. Brown, A. M. Kasprzyk, and M. I. Qureshi. Fano 3-folds in $\mathbb{P}^2 \times \mathbb{P}^2$ format, Tom and Jerry. *Eur. J. Math.*, 4(1):51–72, 2018.
- [BKR12a] G. Brown, M. Kerber, and M. Reid. Fano 3-folds in codimension 4, Tom and Jerry. Part I. *Compos. Math.*, 148(4):1171–1194, 2012.

- [BKR12b] G. Brown, M. Kerber, and M. Reid. Tom and Jerry table, part of "Fano 3-folds in codimension 4, Tom and Terry. Part I". *Compositio Mathematica*, 148(4):1171–1194, 2012.
- [BZ10] G. Brown and F. Zucconi. Graded rings of rank 2 Sarkisov links. *Nagoya Math. J.*, 197:1–44, 2010.
- [Cam20a] L. Campo. Big Table Links. *Online. Access via <http://www.grdb.co.uk/files/fanolinks/BigTableLinks.pdf>*, 2020.
- [Cam20b] L. Campo. Sarkisov links for index 1 fano 3-folds in codimension 4. *arXiv preprint arXiv:2011.12209, to appear in Math. Nachr.*, 2020.
- [Che05] I. Cheltsov. On factoriality of nodal threefolds. *J. Algebraic Geom.*, 14(4):663–690, 2005.
- [Che10] I. Cheltsov. Factorial threefold hypersurfaces. *J. Algebraic Geom.*, 19(4):781–791, 2010.
- [CM04] A. Corti and M. Mella. Birational geometry of terminal quartic 3-folds. I. *Amer. J. Math.*, 126(4):739–761, 2004.
- [Cor95] A. Corti. Factoring birational maps of threefolds after Sarkisov. *J. Algebraic Geom.*, 4(2):223–254, 1995.
- [Cor00] A. Corti. Singularities of linear systems and 3-fold birational geometry. In *Explicit birational geometry of 3-folds*, volume 281 of *London Math. Soc. Lecture Note Ser.*, pages 259–312. Cambridge Univ. Press, Cambridge, 2000.
- [CP06] I. Cheltsov and J. Park. Factorial hypersurfaces in \mathbb{P}^4 with nodes. *Geom. Dedicata*, 121:205–219, 2006.
- [CP17] I. Cheltsov and J. Park. Birationally rigid Fano threefold hypersurfaces. *Mem. Amer. Math. Soc.*, 246(1167):v+117, 2017.
- [CPR00] A. Corti, A. Pukhlikov, and M. Reid. Fano 3-fold hypersurfaces. In *Explicit birational geometry of 3-folds*, volume 281 of *London Math. Soc. Lecture Note Ser.*, pages 175–258. Cambridge Univ. Press, Cambridge, 2000.
- [CR00] Alessio Corti and Miles Reid, editors. *Explicit birational geometry of 3-folds*, volume 281 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2000.
- [HM13] C. D. Hacon and J. McKernan. The Sarkisov program. *J. Algebraic Geom.*, 22(2):389–405, 2013.
- [IF00] A. R. Iano-Fletcher. Working with weighted complete intersections. In *Explicit birational geometry of 3-folds*, volume 281 of *London Math. Soc. Lecture Note Ser.*, pages 101–173. Cambridge Univ. Press, Cambridge, 2000.
- [IP96] V. A. Iskovskikh and A. V. Pukhlikov. Birational automorphisms of multidimensional algebraic manifolds. volume 82, pages 3528–3613. 1996. *Algebraic geometry*, 1.
- [KM83] A. R. Kustin and M. Miller. Constructing big Gorenstein ideals from small ones. *J. Algebra*, 85(2):303–322, 1983.
- [Oka14] T. Okada. Birational Mori fiber structures of \mathbb{Q} -Fano 3-fold weighted complete intersections. *Proc. Lond. Math. Soc. (3)*, 109(6):1549–1600, 2014.
- [Oka18] Takuzo Okada. Birational Mori fiber structures of \mathbb{Q} -Fano 3-fold weighted complete intersections, II. *J. Reine Angew. Math.*, 738:73–129, 2018.
- [Oka20] Takuzo Okada. Birational Mori fiber structures of \mathbb{Q} -Fano 3-fold weighted complete intersections, III. *Kyoto J. Math.*, 60(3):825–893, 2020.
- [Pap04] S. A. Papadakis. Kustin-Miller unprojection with complexes. *J. Algebraic Geom.*, 13(2):249–268, 2004.
- [PR04] S. A. Papadakis and M. Reid. Kustin-Miller unprojection without complexes. *J. Algebraic Geom.*, 13(3):563–577, 2004.
- [Rei80] M. Reid. Canonical 3-folds. In *Journées de Géométrie Algébrique d'Angers, Juillet 1979/Algebraic Geometry, Angers, 1979*, pages 273–310. Sijthoff & Noordhoff, Alphen aan den Rijn—Germantown, Md., 1980.
- [Rei87] M. Reid. Young person's guide to canonical singularities. In *Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985)*, volume 46 of *Proc. Sympos. Pure Math.*, pages 345–414. Amer. Math. Soc., Providence, RI, 1987.
- [Rei00] M. Reid. Graded rings and birational geometry. pages pp. 1–72, 2000. In: Ohno, K., ed. *Proc. of algebraic symposium (Kinosaki, Oct 2000)*.

SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY (KIAS), 85 HOEGIRO, DONGDAEMUN-GU, SEOUL, 02455, REPUBLIC OF KOREA

Email address: liviacampo@kias.re.kr