Discrete Mathematics and Theoretical	Computer Science	
	vol. 2	26:2 #6 (2024)

On harmonious coloring of hypergraphs

Sebastian Czerwiński

Institute of Mathematics, University of Zielona Góra, Poland

revisions 21st Mar. 2023, 8th Nov. 2023; accepted 5th Apr. 2024.

A harmonious coloring of a k-uniform hypergraph H is a vertex coloring such that no two vertices in the same edge have the same color, and each k-element subset of colors appears on at most one edge. The harmonious number h(H) is the least number of colors needed for such a coloring.

The paper contains a new proof of the upper bound $h(H) = O(\sqrt[k]{k!m})$ on the harmonious number of hypergraphs of maximum degree Δ with m edges. We use the local cut lemma of A. Bernshteyn.

Keywords: Harmonious coloring, Hypergraphs

1 Introducion

Let $H=(V,\mathcal{E})$ be a k-uniform hypergraph with the set of vertices V and the set of edges \mathcal{E} . The set of edges is a family of k-element sets of V, where $k\geq 2$.

A rainbow coloring c of a hypergraph H is a map $c:V\mapsto\{1,\ldots,r\}$ in which no two vertices in the same edge have the same color. If two vertices are in the same edge e with the same color, we say that the edge e is bad.

A coloring c is called *harmonious* if $c(e) \neq c(f)$ for every pair of distinct edges $e, f \in \mathcal{E}$ and c is a rainbow coloring.

We say that distinct edges e and f have the *same pattern of colors* if $c(e \setminus f) = c(f \setminus e)$ and there is no uncolored vertex in the set $e \setminus f$.

Let h(H) be the least number of colors needed for a harmonious coloring of H. In Bosek et al. (2016), the following result is proved

Theorem 1 (Bosek et al. (2016)). For every $\varepsilon > 0$ and every $\Delta > 0$ there exist integers k_0 and m_0 such that every k-uniform hypergraph H with m edges (where $m \ge m_0$ and $k \ge k_0$) and maximum degree Δ satisfies

$$h(H) \le (1+\varepsilon) \frac{k}{k-1} \sqrt[k]{\Delta(k-1)k!m}.$$

Remark 1. The paper Bosek et al. (2016) contains the following upper bound on the harmonious number

$$h(H) \leq \frac{k}{k-1} \sqrt[k]{\Delta(k-1)k!m} + 1 + \Delta^2 + (k-1)\Delta + \sum_{i=2}^{k-1} \frac{i}{i-1} \sqrt[i]{(i-1)i\frac{(k-1)\Delta^2}{k-i}}.$$

ISSN 1365-8050

© 2024 by the author(s)

Distributed under a Creative Commons Attribution 4.0 International License

2 Sebastian Czerwiński

The proof of this theorem is based on the entropy compression method, see Grytczuk et al. (2013); Esperet and Parreau (2013).

Because a number r of used colors must satisfy the inequality $\binom{r}{k} \leq m$, we get the lower bound $\Omega(\sqrt[k]{k!m})$. By these observations, it is conjectured by Bosek et al. (2016) that

Conjecture 1. For each $k, \Delta \geq 2$ there exist a constant $c = c(k, \Delta)$ such that every k-uniform hypergraph H with m edges and maximum degree Δ satisfies

$$h(H) \le \sqrt[k]{k!m} + c.$$

A similar conjecture was posed by Edwards (1997b) for simple graphs G. He prove there that

$$h(G) \le (1 + o(1))\sqrt{2m}.$$

There are many results about the harmonious number of particular classes of graphs, see Aflaki et al. (2012); Akbari et al. (2012); Edwards (1997a); Edwards and McDiarmid (1994); Edwards (1996); Krasikov and Roditty (1994) or Aigner et al. (1992); Balister et al. (2002, 2003); Bazgan et al. (1999); Burris and Schelp (1997).

The present paper a new contains proof of the theorem of Bosek et al. We use a different method, the local cut lemma of Bernshteyn (2017, 2016). The proof is simpler and shorter than the original proof of Bosek et al.

2 A special version of the local cut lemma

Let A be a family of subsets of a powerset $\operatorname{Pow}(I)$, where I is a finite set. We say that it is downwards-closed if for each $S \in A$, $\operatorname{Pow}(S) \subseteq (A)$. A subset ∂A of I is called boundary of a downwards-closed family A if

$$\partial A := \{i \in I : S \in A \text{ and } S \cup \{i\} \notin A \text{ for some } S \subseteq I \setminus \{i\}\}.$$

Let $\tau \colon T \mapsto [1; +\infty)$ be a function, then for every $X \subseteq I$ we denote by $\tau(X)$ a number

$$\tau(X) := \prod_{x \in X} \tau(x).$$

Let B be a random event, $X \subseteq I$ and $i \in I$. We introduce two quantities:

$$\sigma_{\tau}^A(B,X) := \max_{Z \subseteq I \backslash X} \Pr(B \text{ and } Z \cup X \not \in A | Z \in A) \cdot \tau(X)$$

and

$$\sigma_{\tau}^{A}(B,i) := \min_{i \in X \subset I} \sigma_{\tau}^{A}(B,X).$$

If $\Pr(Z \in A) = 0$, then $\Pr(P|Z \in A) = 0$, for all events P.

Theorem 2 (Bernshteyn (2017)). Let I be a finite set. Let Ω be a probability space and let $A \colon \Omega \mapsto \operatorname{Pow}(\operatorname{Pow}(I))$ be a random variable such that with probability I, A is a nonempty downwards-closed family of subsets of I. For each $i \in I$, Let $\mathcal{B}(i)$ be a finite collection of random events such that whenever

 $i \in \partial A$, at least one of the events in $\mathcal{B}(i)$ holds. Suppose that there is a function $\tau \colon I \mapsto [1, +\infty)$ such that for all $i \in I$ we have

$$\tau(i) \ge 1 + \sum_{B \in \mathcal{B}(i)} \sigma_{\tau}^{A}(B, i).$$

Then $Pr(I \in A) \ge 1/\tau(I) > 0$.

3 Proof of Theorem 1

We choose a coloring $f: V \mapsto \{1, \dots, t\}$ uniformly at random. Let A be a subset of the power set of V given by

$$A := \{ S \subseteq V : c \text{ is a harmonious coloring of } H(V) \}.$$

It is a nonempty downwards-closed family with probability 1 (the empty set is an element of A)

By a set ∂A , we denote the set of all vertices v such that there is an element X of A such that the coloring c is not a harmonious coloring of $X \cup \{v\}$. If the coloring c is not a harmonious coloring, then there is a bad edge or there are two different edges with the same pattern of colors. So, we define for every $v \in V$, a collection $\mathcal{B}(v)$ as union of sets:

$$\mathcal{B}^1(v) := \{B_e : v \in e \in E(H) \text{ and } e \text{ is not proper colored}\},$$

and for every $i \in \{0, 1, ..., k-1\}$,

$$\mathcal{B}_{i}^{2}(v) := \{B_{e,f} : v \in e, f \in E(H) \text{ and } c(e) = c(f), |e \setminus f| = i\}.$$

That is $\mathcal{B}(v) = \mathcal{B}^1(v) \cup \bigcup_{i=1}^{k-1} \mathcal{B}_i^2(v)$.

We assume that the event B_e happens if and only if the edge e is the bad edge and the event $B_{e,f}$ happens if and only if the edges e and f have the same pattern of colors.

We also assume that a function τ is a constant function, that is, $\tau(v) = \tau \in [1, +\infty)$. This implies that for any subset S of V, we have $\tau(S) = \tau^{|S|}$.

Now, we must find an upper bound on

$$\sigma_{\tau}^{A}(B,v) = \min_{X \subseteq V: v \in X} \max_{Z \subseteq V \setminus X} \Pr(B \land Z \cup X \not\in A | Z \in A) \tau(X),$$

where $v \in V$ and $B \in \mathcal{B}(v)$. We will use an estimation

$$\sigma_{\tau}^{A}(B, v) \le \max_{Z \subseteq V \setminus X} \Pr(B|Z \in A)\tau(X).$$

Now, we consider two cases.

Case 1: $B \in \mathcal{B}^1$, i.e. $B = B_e$.

We choose as X the set $\{e\}$. Because the colors of distinct vertices are independent, we get an upper bound $\sigma_{\tau}^{A}(B_{e},v) \leq \Pr(B_{e})\tau^{k}$ (events B_{e} and " $Z \in A$ " are independent). The probability $\Pr(\overline{B_{e}})$, opposite to $\Pr(B_{e})$, fulfills

$$\Pr(\overline{B_e}) = 1 - \frac{t}{t} \cdot \frac{t-1}{t} \cdot \dots \cdot \frac{t-k+1}{t} \ge 1 - (1 - \frac{k-1}{t})^{k-1}.$$

4 Sebastian Czerwiński

By Bernoulli's inequality, we get

$$\Pr(\overline{B_e}) \ge 1 - (1 - \frac{k-1}{t} \cdot (k-1)) = \frac{(k-1)^2}{t}.$$

So, $\Pr(B_e) \leq \frac{k^2}{t}$.

Case 2: $B \in \mathcal{B}_i^2$, i.e. $B = B_{e,f}$ and $|e \setminus f| = i$. Now, we set $X = e \setminus f$. The probability $\Pr(B_{e,f})$ is bounded from above by $\frac{i!}{t^i}$. So, we get

$$\sigma_{\tau}^{A}(B_{e,f},v) \leq \Pr(B_{e,f})\tau^{i} \leq \frac{i!}{t^{i}}\tau^{i}.$$

To end the proof we must find an upper bound on sizes of sets $\mathcal{B}^1(v)$, $\mathcal{B}^2_0(v)$ and $\mathcal{B}^2_i(v)$, where i > 0. Because the degree of a vertex is bounded by Δ and the number of edges is m we get that

$$|\mathcal{B}^1(v)| \leq \Delta$$
 and $|\mathcal{B}_0^2(v)| \leq \Delta m$.

The hardest part is an upper bound on $\mathcal{B}_i^2(v)$, i > 0. The number of edges f such that $e \setminus f = i$ is bounded by $\frac{k\Delta}{k-i}$. There are at most $k\Delta$ edges with a nonempty intersection with the edge e, and the edge f has exactly k-i common elements with e. So, we have $|\mathcal{B}_i^2(v)| \leq \Delta \frac{k\Delta}{k-i}$. To apply Theorem 2 we must find $\tau \in [1, +\infty)$ and $c \in \mathbb{N}$ such that, for all $v \in V$, the inequality below holds

$$\tau \ge 1 + \Delta \frac{k^2}{t} \tau^k + \Delta m \frac{k!}{t^k} \tau^k + \sum_{i=1}^{k-1} \Delta \frac{k\Delta}{k-i} \frac{i!}{t^i} \tau^i.$$

If we choose $\tau = \frac{k}{k-1}$ and $t = \frac{k}{k-1} \sqrt[k]{\Delta(k-1)k!m}(1+\varepsilon)$, it is easy to see that the inequality holds for sufficiently large m.

Acknowledgments

I would like to thank the anonymous reviewer for his/her suggestions and comments.

References

- A. Aflaki, S. Akbari, K. J. Edwards, D. S. Eskandani, M. Jamaali, and H. Ravanbod. On harmonious colouring of trees. *Electron. J. Combin.*, 19(1):Paper 3, 9, 2012. URL https://doi.org/10. 37236/9.
- M. Aigner, E. Triesch, and Z. Tuza. Irregular assignments and vertex-distinguishing edge-colorings of graphs. In Combinatorics '90 (Gaeta, 1990), volume 52 of Ann. Discrete Math., pages 1-9. North-Holland, Amsterdam, 1992. URL https://doi.org/10.1016/S0167-5060(08)70896-3.
- S. Akbari, J. Kim, and A. Kostochka. Harmonious coloring of trees with large maximum degree. Discrete Math., 312(10):1633-1637, 2012. URL https://doi.org/10.1016/j.disc.2012. 02.009.

- P. N. Balister, B. Bollobás, and R. H. Schelp. Vertex distinguishing colorings of graphs with $\Delta(G)=2$. Discrete Math., 252(1-3):17–29, 2002. URL https://doi.org/10.1016/S0012-365X(01) 00287-4.
- P. N. Balister, O. M. Riordan, and R. H. Schelp. Vertex-distinguishing edge colorings of graphs. *J. Graph Theory*, 42(2):95–109, 2003. URL https://doi.org/10.1002/jgt.10076.
- C. Bazgan, A. Harkat-Benhamdine, H. Li, and M. Woźniak. On the vertex-distinguishing proper edge-colorings of graphs. *J. Combin. Theory Ser. B*, 75(2):288–301, 1999. URL https://doi.org/10.1006/jctb.1998.1884.
- A. Bernshteyn. New bounds for the acyclic chromatic index. *Discrete Math.*, 339(10):2543–2552, 2016. URL https://doi.org/10.1016/j.disc.2016.05.002.
- A. Bernshteyn. The local cut lemma. *European J. Combin.*, 63:95–114, 2017. URL https://doi.org/10.1016/j.ejc.2017.03.005.
- B. Bosek, S. Czerwiński, J. Grytczuk, and P. Rzążewski. Harmonious coloring of uniform hypergraphs. *Appl. Anal. Discrete Math.*, 10(1):73-87, 2016. URL https://doi.org/10.2298/AADM160411008B.
- A. C. Burris and R. H. Schelp. Vertex-distinguishing proper edge-colorings. *J. Graph Theory*, 26(2):73–82, 1997. URL https://doi.org/10.1002/(SICI)1097-0118(199710)26:2<73:: AID-JGT2>3.0.CO; 2-C.
- K. Edwards. The harmonious chromatic number of bounded degree trees. *Combin. Probab. Comput.*, 5 (1):15–28, 1996. URL https://doi.org/10.1017/S0963548300001802.
- K. Edwards. The harmonious chromatic number and the achromatic number, volume 241 of London Math. Soc. Lecture Note Ser., pages 13–47. Cambridge Univ. Press, Cambridge, 1997a. URL https: //doi.org/10.1017/CB09780511662119.003.
- K. Edwards. The harmonious chromatic number of bounded degree graphs. *J. London Math. Soc.* (2), 55 (3):435–447, 1997b. URL https://doi.org/10.1112/S0024610797004857.
- K. Edwards and C. McDiarmid. New upper bounds on harmonious colorings. *J. Graph Theory*, 18(3): 257–267, 1994. URL https://doi.org/10.1002/jgt.3190180305.
- L. Esperet and A. Parreau. Acyclic edge-coloring using entropy compression. *European J. Combin.*, 34 (6):1019–1027, 2013. URL https://doi.org/10.1016/j.ejc.2013.02.007.
- J. Grytczuk, J. Kozik, and P. Micek. New approach to nonrepetitive sequences. *Random Structures Algorithms*, 42(2):214–225, 2013. URL https://doi.org/10.1002/rsa.20411.
- I. Krasikov and Y. Roditty. Bounds for the harmonious chromatic number of a graph. *J. Graph Theory*, 18(2):205–209, 1994. URL https://doi.org/10.1002/jgt.3190180212.