Correspondences in log Hodge cohomology

Charles Godfrey Pacific Northwest National Laboratory

charles.godfrey@pnnl.gov

January 3, 2023

Abstract

We construct correspondences in logarithmic Hodge theory over a perfect field of arbitrary characteristic. These are represented by classes in the cohomology of sheaves of differential forms with log poles and, notably, log *zeroes* on cartesian products of varieties. From one perspective this generalizes work of Chatzistamatiou and Rülling, who developed (non-logarithmic) Hodge correspondences over perfect fields of arbitrary characteristic; from another we provide partial generalizations of more recent work of Binda, Park and Østvær on logarithmic Hodge correspondences by relaxing finiteness and strictness conditions on the correspondences considered.

1 Introduction

Generally speaking, a *correspondence* between two algebraic varieties X and Y over a field k is a cycle or cohomology class on the product $X \times Y$. The study of such objects dates back (at least) to Lefschetz [Lef53], and features prominently in famous conjectures on algebraic cycles (see e.g. [Voi14]) and Voevodsky's theory of motives (see e.g. [MVW06]).

In a number of algebro-geometric research areas it has become commonplace to work with pairs (X, Δ_X) consisting of a variety X together with a divisor Δ_X on X. Such areas include moduli of varieties (where pairs generalize the curves with marked points of [DM69]), birational geometry (where pairs appear naturally, for example as the output of strong resolution of singularities [KM98]) and logarithmic geometry (in this case vast generalizations of divisors Δ_X are allowed [Ogu18]). It is natural to wonder about analogues of correspondences in this category of pairs, and there have been efforts in this direction, for example development of categories of logarithmic motives [BPØ20].

In this paper, we focus on correspondences for logarithmic Hodge cohomology of pairs (X, Δ_X) , where X is a smooth (but not necessarily proper) variety over a perfect field k and Δ_X is a simple normal crossing divisor on X. These cohomology groups can be described as

$$H^*(X, \Delta_X) = \bigoplus H^q(X, \Omega_X^p(\log \Delta_X)), \tag{1.1}$$

where $\Omega_X(\log \Delta_X)$ is the sheaf of differential 1-forms on X with log poles along Δ_X and $\Omega_X^p(\log \Delta_X)$ the p-th exterior power thereof. In addition we consider a generalization where X comes with a family of supports Φ_X , and the ordinary cohomology groups on the right hand side of eq. (1.1) are replaced with cohomology with supports in Φ_X , namely $H_{\Phi_X}^q(X,\Omega_X^p(\log \Delta_X))$. Allowing for supports greatly expands the applicability of our results: for example, it permits us to construct a correspondence associated to a cycle $Z \subset X \times Y$ in a situation where neither X nor Y is proper over k, but Z is proper over both X and Y.

There are multiple motivations for investigating correspondences for this particular cohomology of pairs:

¹One way that such a cycle Z might naturally arise is as the closure of the graph of a birational equivalence $X \rightarrow Y$ of non-proper varieties.

This work was completed while the author was a PhD student in the University of Washington Department of Mathematics. The author was partially supported by the University of Washington Department of Mathematics Graduate Research Fellowship, and by the NSF grant DMS-1440140, administered by the Mathematical Sciences Research Institute, while in residence at MSRI during the program Birational Geometry and Moduli Spaces.

- By analogy with the case of varieties (that is, without auxiliary divisors/log structures), we suspect that correspondences at the level of Chow cycles are more fundamental, and that (many) correspondences in logarithmic Hodge cohomology are obtained from Chow correspondences via a cycle morphism. However, as of this writing there is no full-fledged theory of Chow cohomology of pairs or log schemes (though there has been considerable progress, for instance in [Bar18; BBG22]). Logarithmic Hodge cohomology is in contrast quite mature, appearing as early as [Del71].
- Correspondences in (non-logarithmic) Hodge cohomology have found remarkable applications. For example, [CR11] used them to prove birational invariance of the cohomology groups of the structure sheaf $H^i(X, \mathcal{O}_X)$ for smooth varieties X over perfect fields of positive characteristic. In fact, attempting to implement a similar strategy with logarithmic Hodge cohomology to obtain results on invariance of the cohomology groups $H^i(X, \mathcal{O}_X(-\Delta_X))$ with respect to (a restricted class of) birational equivalences was the initial inspiration for this work. Ultimately that attempt was unsuccessful, as we describe in Appendix A.
- There has been recent interest in logarithmic Hodge cohomology as a representable functor on a category of motives of log schemes over a perfect field [BPØ20, §9]. While that work does also construct some correspondences, they are restricted to those associated with logarithmic Hodge cohomology classes of cycles $Z \subset X \times Y$ which are *finite* over X and obey additional strictness (in the sense of logarithmic geometry) conditions; we remove these restrictions.

The correspondences we construct are obtained from certain Hodge classes with both log poles *and* log zeroes. Our main result is:

Theorem 1.2 (= Theorem 4.1). A class $\gamma \in H^j_{P(\Phi_X,\Phi_Y)}(X \times Y, \Omega^i_{X \times Y}(\log \Delta_{X \times Y})(-\operatorname{pr}^*_X \Delta_X))$ defines homomorphisms

$$\operatorname{cor}(\gamma): H^q_{\Phi_Y}(X, \Omega_X^p(\log \Delta_X)) \to H^{q+j-d_X}_{\Phi_Y}(Y, \Omega_Y^{p+i-d_X}(\log \Delta_Y))$$

by the formula $\operatorname{cor}(\gamma)(\alpha) := \operatorname{pr}_{Y*}(\operatorname{pr}_X^*(\alpha) \smile \gamma)$. Moreover if (Z, Δ_Z, Φ_Z) is another snc pair with supports and $\delta \in H^{j'}_{P(\Phi_V, \Phi_Z)}(Y \times Z, \Omega^{i'}_{Y \times Z}(\log \Delta_{Y \times Z})(-\operatorname{pr}_Y^* \Delta_Y))$, then

$$\mathrm{pr}_{X\times Z*}(\mathrm{pr}_{X\times Y}^*(\gamma)\smile\mathrm{pr}_{Y\times Z}^*(\delta))\in H^{j+j'-d_Y}_{P(\Phi_X,\Phi_Z)}(X\times Z,\Omega_{X\times Z}^{i+i'-d_Y}(\log\Delta_{X\times Z})(-\mathrm{pr}_X^*\Delta_X))\ and$$

$$\mathrm{cor}(\mathrm{pr}_{X\times Z*}(\mathrm{pr}_{X\times Y}^*(\gamma)\smile\mathrm{pr}_{Y\times Z}^*(\delta)))=\mathrm{cor}(\delta)\circ\mathrm{cor}(\gamma)$$

$$as\ homomorphisms\ H^q_{\Phi_X}(X,\Omega_X^p(\log\Delta_X))\to H^{q+j+j'-d_X-d_Y}_{\Phi_Z}(Z,\Omega_Z^{p+i+i'-d_X-d_Y}(\log\Delta_Z)).$$

In the above, $\Delta_{XY} := \operatorname{pr}_X^* \Delta_X + \operatorname{pr}_Y^* \Delta_Y$, a simple normal crossing divisor on $X \times Y$. There is a simple heuristic explanation for the appearance of differential forms in $\Omega^i_{X\times Y}(\log \Delta_{X\times Y})(-\operatorname{pr}_X^* \Delta_X)$: working over the complex numbers, in the case where X and Y are both proper the class $\operatorname{cor}(\gamma)(\alpha) := \operatorname{pr}_{Y*}(\operatorname{pr}_X^*(\alpha) \smile \gamma)$ can be computed explicitly as an integral of the form

$$\int_{X} \alpha(x) \wedge \gamma(x, y), \tag{1.3}$$

and this integral will only be finite when the log poles of α along Δ_X are cancelled by complementary zeroes of the form $\gamma(x, y)$ along the preimage $\operatorname{pr}_X^* \Delta_X$.

Our proof of Theorem 1.2 relies heavily on prior work on both Hodge cohomology with supports [CR11, §2] and its logarithmic variant [BPØ20, §9]. Section 2 is a rapid summary of those results. The key new technical ingredient is a base change formula on the interaction of pushforward and pullback operations in cartesian squares, proved in Section 3. Section 4 includes the proof of our main theorem.

1.1 Acknowledgements

Thanks to Daniel Bragg, Yun Hao, Sarah Scherotzke, Nicolò Sibilla and Mattia Talpo for helpful conversations, to Lawrence Jack Barrott for illuminating email correspondence regarding logarithmic aspects of Chow and Hodge, and to my advisor Sándor Kovács for many insightful discussions. Thanks also to the participants of the Spring 2019 MSRI graduate student seminar, in particular Giovanni Inchiostro and organizer Fatemeh Rezaee, for feedback on early work on this paper.

2 Functoriality properties of log Hodge cohomology with supports

2.1 Supports

In order to obtain results that apply to correspondences between varieties X and Y where neither X nor Y is proper, it is necessary to work with cohomology with supports, also known as local cohomology. A primary source for the material of this subsection is [R&D, §IV]. Let X be a noetherian scheme.

Definition 2.1 ([R&D, §IV], [CR11, §1.1]). A **family of supports** Φ **on** X is a non-empty collection Φ of closed subsets of X such that

- If $C \in \Phi$ and $D \subset C$ is a closed subset, then $D \in \Phi$.
- If $C, D \in \Phi$ then $C \cup D \in \Phi$.

Example 2.2. $\Phi = \{ \text{ all closed subsets of } X \}$ is a family of supports. More generally if $\mathcal C$ is any collection of closed subsets $C \subset X$, there is a *smallest* family of supports $\Phi(\mathcal C)$ containing $\mathcal C$ (explicitly, $\Phi(\mathcal C)$ consists of finite unions $\bigcup_i Z_i$ of closed subsets $Z_i \subset C_i$ of elements $C_i \in \mathcal C$). Taking $\Phi = \Phi(\{X\})$ recovers the previous example. A more interesting example is the case where for some fixed $p \in \mathbb{N}$, $\Phi = \{\text{closed sets } Z \subseteq X \mid \dim Z \le p\}$.

There is a close relationship between families of supports on X and certain collections of specialization-closed subsets of points on X, and we can also consider sheaves of families of supports — for further details we refer to [R&D, §IV.1].

If $f: X \to Y$ is a morphism of noetherian schemes and Ψ is a family of supports on Y, then $\{f^{-1}(Z) \mid Z \in \Psi\}$ is a family of closed subsets of X, and is closed under unions, but is *not* in general closed under taking closed subsets.

Definition 2.3. $f^{-1}(\Psi)$ is the smallest family of supports on X containing $\{f^{-1}(Z) \mid Z \in \Psi\}$.

Let Φ be a family of supports on X. The notation/terminology $f|_{\Phi}$ is **proper** will mean $f|_{C}$ is proper for every $C \in \Phi$. If $f|_{\Phi}$ is proper then $f(C) \subset Y$ is closed for every $C \in \Phi$ and in fact

$$f(\Phi) = \{ f(C) \subset Y \mid C \in \Phi \}$$
 (2.4)

is a family of supports on Y. The key point here is that if $D \subset f(C)$ is closed, then $f^{-1}(D) \cap C \in \Phi$ and $D = f(f^{-1}(D) \cap C)$.

Definition 2.5. A **scheme with supports** (X, Φ_X) is a scheme X together with a family of supports Φ_X on X.

Definition 2.6. A **pushing morphism** $f:(X,\Phi_X)\to (Y,\Phi_Y)$ of schemes with supports is a morphism $f:X\to Y$ of underlying schemes such that $f|_{\Phi_X}$ is proper and $f(\Phi_X)\subset \Phi_Y$. A **pulling morphism** $f:X\to Y$ is a morphism $f:X\to Y$ such that $f^{-1}(\Phi_Y)\subset \Phi_X$.

These morphisms provide two different categories with underlying set of objects schemes with supports (X, Φ_X) , and pushing/pulling morphisms respectively (the verification is elementary; for instance a composition of pushing morphisms is again a pushing morphism since compositions of proper morphisms are proper). Schemes with supports provide a natural setting for describing

functoriality properties of local cohomology. Let \mathcal{F} be a sheaf of abelian groups on a scheme with supports (X, Φ_X) .

Definition 2.7. The **sheaf of sections with supports** of \mathcal{F} , denoted $\underline{\Gamma}_{\Phi}(\mathcal{F})$, is obtained by setting

$$\underline{\Gamma}_{\Phi}(\mathcal{F})(U) = \{ \sigma \in \mathcal{F}(U) \mid \operatorname{supp} \sigma \in \Phi_X|_U \}$$
(2.8)

for each open $U \subset X$ (here $\Phi_X|_U$ is short for $\iota^{-1}\Phi_X$ where $\iota: U \to X$ is the inclusion). More explicitly: for a local section $\sigma \in \mathcal{F}(U)$, $\sigma \in \underline{\Gamma}_{\Phi}(\mathcal{F})(U)$ means $\operatorname{supp} \sigma = C \cap U$ for a closed set $C \subset \Phi_X$.

The functor $\underline{\Gamma}_{\Phi}$ is right adjoint to an exact functor, for instance the inclusion of the subcategory $\mathbf{Ab}_{\Phi}(X) \subset \mathbf{Ab}(X)$ of abelian sheaves on X with supports in Φ ; so, $\underline{\Gamma}_{\Phi}$ is left exact and preserves injectives. In the case $\Phi = \Phi(Z)$ for some closed $Z \subset X$, this is proved in [Stacks, Tag 0A39, Tag 0G6Y, Tag 0G7F] — the general case can then be obtained by writing $\underline{\Gamma}_{\Phi}$ as a filtered colimit:

$$\underline{\Gamma}_{\Phi} = \operatorname{colim}_{Z \in \Phi} \underline{\Gamma}_{Z}.$$

The right derived functor of $\underline{\Gamma}_{\Phi}$ will be denoted $R\underline{\Gamma}_{\Phi}$. Taking global sections on X gives the **sections** with supports of \mathcal{F} : $\Gamma_{\Phi}(\mathcal{F}) := \Gamma_{X}(\underline{\Gamma}_{\Phi}(\mathcal{F}))$ This is also left exact, and (the cohomologies of) its derived functor give the **cohomology with supports in** Φ : $H^{i}_{\Phi}(X,\mathcal{F}) := R^{i}\Gamma_{\Phi}(\mathcal{F})$.

Proposition 2.9. Cohomology with supports enjoys the following functoriality properties:

(i) If $f:(X,\Phi_X)\to (Y,\Phi_Y)$ is a pulling morphism of schemes with supports, \mathcal{F},\mathcal{G} are sheaves of abelian groups on X,Y respectively, and if

$$\varphi: \mathcal{G} \to f_* \mathcal{F}$$
 is a morphism of sheaves, (2.10)

then there is a natural morphism $R\underline{\Gamma}_{\Phi}\mathscr{G}\to Rf_*R\underline{\Gamma}_{\Phi}\mathscr{F}$. Similarly if \mathscr{F} and \mathscr{G} are quasicoherent then there are natural morphisms $R\underline{\Gamma}_{\Phi}\mathscr{G}\to Rf_*R\underline{\Gamma}_{\Phi}\mathscr{F}$.

(ii) If $f:(X,\Phi_X)\to (Y,\Phi_Y)$ is a pushing morphism, \mathcal{F},\mathcal{G} are sheaves of abelian groups on X,Y respectively, and

$$\psi: Rf_*\mathcal{F} \to \mathcal{G}$$
 is a morphism in the derived category of X, (2.11)

then there is a natural morphism $Rf_*R\underline{\Gamma}_{\Phi}(\mathcal{F}) \to R\underline{\Gamma}_{\Phi}\mathcal{G}$.

Both parts of the proposition follow from [Stacks, Tag 0G78]; (i) is discussed in detail in [CR11, $\S2.1$] and (ii) can be extracted from [CR11, $\S2.2$] (although it doesn't appear to be stated explicitly). See also [BPØ20, Constructions 9.4.2, 9.5.3]

2.2 Differential forms with log poles

Let *k* be a perfect field.

Definition 2.12. A **snc pair with supports** (X, Δ_X, Φ_X) over k is a smooth scheme X separated and of finite type over k with a family of supports Φ_X together with a reduced, effective divisor Δ_X on X such that supp Δ_X has simple normal crossings, in the sense that for any point $x \in X$ there are regular parameters $z_1, \ldots, z_c \in \mathcal{O}_{X,x}$ such that supp $\Delta_X = V(z_1 \cdot z_2 \cdots z_r)$ on a Zariski neighborhood of x.³ The **interior** U_X of a snc pair with supports (X, Δ_X, Φ_X) is

$$U_X := X \setminus \operatorname{supp} \Delta_X \tag{2.13}$$

The inclusion of U_X in X is denoted by ι_X : $U_X \to X$.

²Simply put \mathcal{F} is a sheaf of abelian groups on X.

³This is equivalent to the more general definition [BPØ20, Def. 7.2.1] in the case where the base scheme is Spec k, which is all we need.

Here supp Δ_X denotes the **support** of Δ_X (if $\Delta_X = \sum_i a_i D_i$ where the D_i are prime divisors, then supp $\Delta_X = \cup_i D_i$). Similarly let j_X : supp $\Delta_X \to X$ denote the evident inclusion.

Definition 2.14 (compare with [CR11, Def. 1.1.4]). A **pulling morphism** $f:(X,\Delta_X,\Phi_X)\to (Y,\Delta_Y,\Phi_Y)$ **of snc pairs with supports** is a pulling morphism $f:X\to Y$ of underlying schemes with support such that $f^{-1}(\sup \Delta_Y)\subset \sup \Delta_X$; equivalently, f restricts to a morphism $f|_{U_X}:U_X\to U_Y$. A **pushing morphism** $f:(X,\Delta_X,\Phi_X)\to (Y,\Delta_Y,\Phi_Y)$ **of snc pairs with supports** is a pushing morphism of underlying schemes with support such that $f^*\Delta_Y=\Delta_X$.

Note that if $f:(X,\Delta_X,\Phi_X)\to (Y,\Delta_Y,\Phi_Y)$ is a pushing morphism then $U_X=f^{-1}(U_Y)$, so for example if $f:X\to Y$ is proper then so is the induced map $U_X\to U_Y$.

Convention 2.15 (compare with [CR11, p. 1.1.5]). A morphism of snc pairs with supports $f:(X,\Delta_X,\Phi_X)\to (Y,\Delta_Y,\Phi_Y)$ is flat, proper, an immersion, etc. if and only if the same is true of the underlying morphism of schemes $f:X\to Y$. A diagram of snc pairs with supports

$$(X', \Delta_{X'}, \Phi_{X'}) \xrightarrow{g'} (X, \Delta_X, \Phi_X)$$

$$f' \downarrow \qquad \qquad \downarrow f$$

$$(Y', \Delta_{Y'}, \Phi_{Y'}) \xrightarrow{g} (Y, \Delta_Y, \Phi_Y)$$

$$(2.16)$$

is cartesian if and only if the induced diagram of underlying schemes

$$X' \xrightarrow{g'} X$$

$$f' \downarrow \qquad \qquad \downarrow f$$

$$Y' \xrightarrow{g} Y$$

$$(2.17)$$

is cartesian.4

The terminology is meant to suggest that pushing (resp. pulling) morphisms induce pushforward (resp. pullback) maps on log Hodge cohomology, as we now describe.

If (X, Δ_X) is an snc pair, or more generally a normal separated scheme of finite type X over k together with a sequence of effective Cartier divisors $D_1, \dots, D_N \subseteq X$ with sum $\Delta_X = \sum_i D_i$, then it comes with a sheaf of differential forms with log poles $\Omega_X(\log \Delta_X)$. In the case where (X, Δ_X, Φ_X) is snc, this sheaf and its properties are described in [EV92, §2]. For a definition and treatment of $\Omega_X(\log \Delta_X)$ in the much greater generality of logarithmic schemes we refer to [Ogu18, §IV].

In some of the calculations below the following concrete local description will be very useful. Let $z_1, z_2, ..., z_n$ be local coordinates at a point $x \in X$ such that supp $\Delta_X = V(z_1 z_2 \cdots z_r)$ in a neighborhood of x. Recall that as X is smooth the differentials $d z_1, d z_2, ..., d z_n$ freely generate Ω_X on a neighborhood of x.

Lemma 2.18 (see e.g. [EV92, §2]). The sections $\frac{dz_1}{z_1}, \dots, \frac{dz_r}{z_r}, dz_{r+1}, \dots, dz_n$ freely generate $\Omega_X(\log \Delta_X)$ on a neighborhood of x.

Given $\Omega_X(\log \Delta_X)$, we can form the exterior powers

$$\Omega_X^p(\log \Delta_X) := \bigwedge^p \Omega_X(\log \Delta_X), \tag{2.19}$$

and combining Lemma 2.18 with (2.19) gives concrete local descriptions of the $\Omega_X^p(\log \Delta_X)$; in particular, we see that $\Omega_X^{\dim X}(\log \Delta_X) = \omega_X(\Delta_X)$.

⁴If we take the red pill of logarithmic geometry, it starts to seem almost more reasonable to only require flatness, properness, cartesianness and so on of the induced maps of *interiors* $U_X \to U_Y$. However we do use the stronger restrictions of the given definition in some of the proofs below.

Definition 2.20. The **log-Hodge cohomology with supports** of a log-smooth pair with supports (X, Δ_X, Φ_X) is defined by

$$H^{d}(X, \Delta_{X}, \Phi_{X}) = \bigoplus_{p+q=d} H^{q}_{\Phi}(X, \Omega_{X}^{p}(\log \Delta_{X}))$$
(2.21)

Here H^q_{Φ} denotes local cohomology with respect to the family of supports Φ_X . For connected X, we $define H_d(X, \Delta_X, \Phi_X) := H^{2\dim X - d}(X, \Delta_X, \Phi_X)$, and in general we set $H_d(X, \Delta_X, \Phi_X) = \bigoplus_i H_d(X_i, \Delta_{X_i}, \Phi_{X_i})$ where X_i are the connected components of X.

Let $f:(X,\Delta_X,\Phi_X)\to (Y,\Delta_Y,\Phi_Y)$ be pulling morphism of snc pairs with supports.

Lemma 2.22 ([Ogu18, Prop. 2.3.1] + (2.19)). The map f induces a morphism of sheaves

$$f^*\Omega_Y^p(\log \Delta_Y) \xrightarrow{d \ f^{\vee}} \Omega_X^p(\log \Delta_X) \ adjoint \ to \ a \ morphism$$

$$f^*\Omega_Y^p(\log \Delta_Y) \xrightarrow{d \ f^{\vee}} \Omega_X^p(\log \Delta_X) \ for \ all \ p.$$
(2.23)

The essential content of this lemma is that when we pull back a log differential form σ on (Y, Δ_Y) , it doesn't *develop* poles of order ≥ 1 along Δ_X . Combining the previous lemma with proposition 2.9 gives:

Proposition 2.24 ([BPØ20, §9.1-2], see also [CR11, §2.1]). For every pulling morphism $f:(X, \Delta_X, \Phi_X) \to (Y, \Delta_Y, \Phi_Y)$ there are functorial morphisms

$$R\Gamma_{\Phi}\Omega_{V}^{p}(\log \Delta_{Y}) \to Rf_{*}R\Gamma_{\Phi}\Omega_{V}^{p}(\log \Delta_{Y})$$
 for all p (2.25)

In particular, for each p, q there are functorial homomorphisms

$$f^*: H^q_{\Phi}(Y, \Omega^p_Y(\log \Delta_Y)) \to H^q_{\Phi}(X, \Omega^p_X(\log \Delta_X))$$
 (2.26)

and hence (summing over p + q = d) functorial homomorphisms

$$f^*: H^d(X, \Delta_X, \Phi_X) \to H^d(Y, \Delta_Y, \Phi_Y)$$
 (2.27)

The maps $f_*: H_d(X, \Delta_X, \Phi_X) \to H_d(Y, \Delta_Y, \Phi_Y)$ induced by a pushing morphism $f: (X, \Delta_X, \Phi_X) \to (Y, \Delta_Y, \Phi_Y)$ can be obtained from a combination of Nagata compactification and Grothendieck duality.

Lemma 2.28 ([BPØ20, §9.5], see also [CR11, §2.3]). Let $f: (X, \Delta_X, \Phi_X) \to (Y, \Delta_Y, \Phi_Y)$ be a pushing morphism of equidimensional log-smooth pairs with support such that. Then letting $c = \dim Y - \dim X$, for each p there are functorial morphisms of complexes of coherent sheaves

$$Rf_*R\underline{\Gamma}_{\Phi_Y}(\Omega_X^p(\log \Delta_X)) \to R\underline{\Gamma}_{\Phi_Y}\Omega_Y^{p+c}(\log \Delta_Y)[c]$$
 (2.29)

inducing maps on cohomology

$$f_*: H^q_{\Phi_X}(X, \Omega_X^p(\log \Delta_X)) \to H^{q+c}_{\Phi_Y}(Y, \Omega_Y^{p+c}(\log \Delta_Y))$$
(2.30)

for all q.

Since they enter into the calculations below, we give a description of these pushforward morphisms. Before beginning, a word on duality in our current setup: since we are working exclusively over Spec k, we can make use of compatible normalized dualizing complexes — namely, if $\pi:Z\to \operatorname{Spec} k$ is a separated finite type k-scheme then $\pi^!\mathfrak{G}_{\operatorname{Spec} k}$ is a dualizing complex [Stacks, Tag 0E2S, Tag 0FVU]. We will make repeated use of the behavior of dualizing with respect to differentials: as a consequence of Lemma 2.18, wedge product gives a perfect pairing

$$\Omega_X^p(\log \Delta_X)(-\Delta_X) \otimes \Omega_X^{\dim X - p}(\log \Delta_X) \to \omega_X$$
 (2.31)

(see also [Har77, Cor. III.7.13]) and so $\Omega_X^{\dim X-p}(\log \Delta_X) \simeq R\mathcal{H}om_X(\Omega_X^p(\log \Delta_X)(-\Delta_X),\omega_X)$. Here the derived sheaf Hom $R\mathcal{H}om_X$ agrees with the regular sheaf Hom as $\Omega_X^p(\log \Delta_X)(-\Delta_X)$ is locally free. On the other hand, the *dualizing functor* of X is $R\mathcal{H}om_X(\Omega_X^p(\log \Delta_X)(-\Delta_X),\omega_X[\dim X])$ where $\omega_X = \Omega_X^{\dim X}$. An upshot is that Grothendieck duality calculations involving the sheaves of differential forms become more symmetric and predictable if we work with the shifted versions $\Omega_X^p(\log \Delta_X)(-\Delta_X)[p]$; for example then we have the identity

$$\Omega_X^{\dim X - p}(\log \Delta_X)[\dim X - p] \simeq R\mathcal{H}om_X(\Omega_X^p(\log \Delta_X)(-\Delta_X)[p], \omega_X[\dim X])$$

Now, we need to compactify $f: X \to Y$.

Theorem 2.32 ([Nag63, §4 Thm. 2], [Con07, Thm. 4.1]). Let S be a quasi-compact quasi-separated scheme and let $X \to S$ be a separated morphism of finite type. Then there is a dense open immersion of S-schemes $X \hookrightarrow \overline{X}$ such that \overline{X} is proper.

Using Theorem 2.32 we obtain morphisms of schemes

$$X \xrightarrow{\iota} \bar{X}$$

$$\downarrow^{\bar{f}}$$

$$Y$$

$$(2.33)$$

where $\iota: X \to \bar{X}$ is a dense open immersion and $\bar{f}: \bar{X} \to Y$ is *proper*. Note that \bar{X} need not be smooth over k, and in the absence of resolutions of singularities⁵ there is not even a way to make \bar{X} smooth. This means we cannot hope to upgrade \bar{X} to a simple normal crossing pair $(\bar{X}, \Delta_{\bar{X}})$. However, we do still have a divisor $\Delta_{\bar{X}} := \bar{f}^* \Delta_y$ on \bar{X} . One way to overcome these difficulties is to equip the possibly singular \bar{X} with a *logarithmic structure*, in some sense associated to $\Delta_{\bar{X}}$, whose restriction to X coincides with a logarithmic structure naturally defined by the simple normal crossing divisor Δ_X .

Formally, we use the log structure on \bar{X} pulled back from the log structure on (Y, Δ_Y) [Ogu18, §III.1.6-7] along the morphism $\bar{f}: \bar{X} \to Y$. Since $(Y, \Delta_Y = \sum_{i=1}^N D_i^Y)$ is a simple normal crossing pair, its associated log structure is Deligne-Faltings [Ogu18, §III.1.7] and can be encoded in the sequence of inclusions of ideal sheaves $\mathcal{O}_Y(-D_i^Y) \hookrightarrow \mathcal{O}_Y$. The pullback log structure on \bar{X} can then be encoded in the sequence of inclusions of ideal sheaves

$$\bar{f}^{-1} \mathcal{O}_Y (-D_i^Y) \cdot \mathcal{O}_{\bar{X}} = \mathcal{O}_{\bar{X}} (-\bar{f}^* D_i^Y) \hookrightarrow \mathcal{O}_{\bar{X}}.$$

The pushforward morphisms of Lemma 2.28 are defined using the sheaves of log differential p-forms on \bar{X} over k as described in [Ogu18, §IV.1, V.2] — these will be denoted⁶ by $\Omega_{\overline{X}}^p(\log \Delta_{\overline{X}})$. The essential properties that we need are:

• $\Omega_{\overline{X}}^{p}(\log \Delta_{\overline{X}})$ is a coherent sheaf on \overline{X} together with a functorial morphism

$$\Omega_Y^p(\log \Delta_Y) \to \overline{f}_*\Omega_{\overline{X}}^p(\log \Delta_{\overline{X}}).$$

Coherence can be obtained as follows: first, the log structure on (Y, Δ_Y) is coherent ([Ogu18, §III.1.9]), and hence so is its pullback to \bar{X} (see for example [Ogu18, Def. III.1.1.5, Rmk III.1.1.6]). Then [Ogu18, Cor. IV.1.2.8] implies $\Omega_{\bar{X}}^1(\log \Delta_{\bar{X}})$ is a coherent sheaf, and it follows that its p-th exterior powers are coherent sheaves as well. The desired functorial morphism can be obtained from [Ogu18, Prop. IV.1.2.15].

⁵At the time of this writing, this applies to the cases char k = p > 0 and dim X > 3.

⁶This is an abuse of notation since the construction of this sheaf is (as far as we know) not the same as the one for simple normal crossing pairs described above Lemma 2.18, however the notation of [Ogu18] seems unsatisfactory for our purposes as we wish to stress that these are not the ordinary differential forms Ω^p_{∇} ,

• There is a natural isomorphism $\Omega^{\underline{p}}_{\overline{X}}(\log \Delta_{\overline{X}})|_{X} \simeq \Omega^{\underline{p}}_{X}(\Delta_{X})$. This can be seen by observing that the log structures on (X, Δ_{X}) and \bar{X} are obtained as pullbacks of the log structure on (Y, Δ_{Y}) with respect to f and \bar{f} respectively (in the case of (X, Δ_{X}) this follows from Definition 2.14, and in the latter case it is how we defined the log structure on \bar{X}). Hence considering eq. (2.33) we find that the log structure on \bar{X} restricts to that on (X, Δ_{X}) .

Hence in particular $\Omega_{\overline{X}}^p(\log \Delta_{\overline{X}})$ is a *functorial coherent extension* of $\Omega_X^p(\Delta_X)$ to the possibly non-snc log scheme \overline{X} . Starting with the log differential

$$d \operatorname{pr}_{Y}^{\vee} : \Omega_{Y}^{p}(\log \Delta_{Y})[p] \to R\overline{f}_{*}\Omega_{\overline{X}}^{p}(\log \Delta_{\overline{X}})[p],$$

twisting by $-\Delta_Y$ and using the projection formula gives a morphism (*note*: this is where we use the assumptions that $f^*\Delta_Y = \Delta_X$ and $\bar{f}^*\Delta_Y = \Delta_{\bar{X}}$)

$$\Omega_{Y}^{p}(\log \Delta_{Y})(-\Delta_{Y})[p] \to R\overline{f}_{*}\Omega_{\overline{X}}^{p}(\log \Delta_{\overline{X}})(-\Delta_{\overline{X}})[p]$$
(2.34)

to which we apply Grothendieck duality:

Theorem 2.35 (Grothendieck duality, [R&D, Cor. VII.3.4], [Con00, Thm. 3.4.4]). Let $f: X \to Y$ be a proper morphism of finite-dimensional noetherian schemes and assume Y admits a dualizing complex (for example X and Y could be schemes of finite type over K). Then for any pair of objects $\mathcal{F}^{\bullet} \in D_{qc}^{-}(X)$ and $\mathcal{G}^{\bullet} \in D_{c}^{+}(Y)$ there is a natural isomorphism

$$Rf_*R\underline{Hom}_X(\mathcal{F}^{\bullet}, f^!\mathcal{G}^{\bullet}) \simeq R\underline{Hom}_Y(Rf_*\mathcal{F}^{\bullet}, \mathcal{G}^{\bullet}) \text{ in } D^b_c(Y)$$

Combining Theorem 2.35 with eq. (2.34) gives a morphism

where the equality is Theorem 2.35 and the vertical map is induced by (2.34). Adding supports gives a morphism

$$Rf_*R\underline{\Gamma}_{\Phi_X}R\mathcal{H}om_X(\Omega_X^p(\log\Delta_X)(-\Delta_X)[p],\omega_X[\dim X]) = R\overline{f}_*R\underline{\Gamma}_{\Phi_X}R\mathcal{H}om_{\overline{X}}(\Omega_{\overline{X}}^p(\log\Delta_{\overline{X}})(-\Delta_{\overline{X}})[p],\omega_{\overline{X}})$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad (2.37)$$

$$R\underline{\Gamma}_{\Phi_Y}R\mathcal{H}om_Y(\Omega_Y^p(\log\Delta_Y)(-\Delta_Y)[p],\omega_Y[\dim Y])$$

where the equality is obtained from the *excision* property of local cohomology, compatibility of the dualizing functor with restriction *and* the natural isomorphism $\Omega_{\overline{X}}^p(\log \Delta_{\overline{X}})|_X \simeq \Omega_X^p(\Delta_X)$. Using (2.31) we obtain

$$\Omega_X^{\dim X-p}(\log \Delta_X) \simeq \mathcal{H}om_X(\Omega_X^p(\log \Delta_X)(-\Delta_X),\omega_X) = R\mathcal{H}om_X(\Omega_X^p(\log \Delta_X)(-\Delta_X),\omega_X)$$

where the last equality uses the fact that $\Omega_X^p(\log \Delta_X)(-\Delta_X)$ is locally free. A similar calculation on Y transforms (2.37) into:

$$Rf_*R\underline{\Gamma}_{\Phi_Y}\Omega_X^{\dim X-p}(\log\Delta_X)[\dim X-p]\to R\underline{\Gamma}_{\Phi_Y}\Omega_Y^{\dim Y-p}(\log\Delta_Y)[\dim Y-p]$$

and reindexing like $p \leftrightarrow \dim X - p$ recovers Lemma 2.28.

3 A base change formula

Lemma 3.1 (compare with [CR11, Prop. 2.3.7]). *Let*

$$(X', \Delta_{X'}, \Phi_{X'}) \xrightarrow{g'} (X, \Delta_X, \Phi_X)$$

$$\downarrow f' \qquad \qquad \downarrow f$$

$$(Y', \Delta_{Y'}, \Phi_{Y'}) \xrightarrow{g} (Y, \Delta_Y, \Phi_Y)$$

$$(3.2)$$

be a cartesian diagram of equidimensional snc pairs with supports, where f, f' (resp. g, g') are pushing (resp. pulling) morphisms and g is either flat or a closed immersion transverse to f. Then

$$g^*f_* = f'_*g'^* : H^*(X, \Delta_X, \Phi_X) \to H^*(Y', \Delta_{Y'}, \Phi_{Y'}).$$

We will prove this following Chatzistamation and Rülling's argument [CR11, Prop. 2.3.7] quite closely, at various points reducing to statements proved therein. In the proofs we will make use of a slight variant of Definition 2.3.

Definition 3.3. If $f: X \to Y$ is a morphism of noetherian schemes and let Φ_Y is a family of supports on Y, then

$$f_*^{-1}(\Phi_V) := \{Z \subseteq X \mid f|_Z \text{ is proper and } f(Z) \in \Phi_V\}$$

Lemma 3.4. It suffices to prove Lemma 3.1 in the cases where f is either

- (i) a projection morphism of the form $\operatorname{pr}_Y: (X \times Y, \operatorname{pr}_Y^* \Delta_Y, \operatorname{pr}_{Y*}^{-1}(\Phi_Y)) \to (Y, \Delta_Y, \Phi_Y)$, or
- (ii) a closed immersion.

Remark 3.5. This lemma makes essential use of the functoriality part of Lemma 2.28.

Proof. We can decompose (3.2) as a concatenation of cartisian diagrams

$$(X', \Delta_{X'}, \Phi_{X'}) \xrightarrow{g'} (X, \Delta_{X}, \Phi_{X})$$

$$\downarrow^{h'} (2) \qquad \downarrow^{h}$$

$$(X \times Y', \operatorname{pr}_{Y'}^{*} \Delta_{Y}, \operatorname{pr}_{Y'*}^{-1}(\Phi_{Y}')) \xrightarrow{\operatorname{id} \times g} (X \times Y, \operatorname{pr}_{Y}^{*} \Delta_{Y}, \operatorname{pr}_{Y*}^{-1}(\Phi_{Y}))$$

$$\downarrow^{\operatorname{pr}_{Y'}} (1) \qquad \downarrow^{\operatorname{pr}_{Y}}$$

$$(Y', \Delta_{Y'}, \Phi_{Y'}) \xrightarrow{g} (Y, \Delta_{Y}, \Phi_{Y})$$

$$(3.6)$$

where $h = id \times f$ is the graph morphism of f and $h' = g' \times f'$. If g is flat or a closed immersion transverse to f then $id \times g$ is flat or a closed immersion transverse to h (by base change).

Here the only new feature not covered in [CR11, Prop. 2.3.7] is the presence of divisors, and we simply note that $\Delta_X = f^*\Delta_X = h^* \operatorname{pr}_Y^*\Delta_Y$ and similarly for $\Delta_{X'}$, so that both pr_Y and h are pushing morphisms in the sense of Definition 2.14, and similarly for the left vertical maps. In other words, the supports and divisors in the middle row have been chosen precisely so that the vertical morphisms are all "pushing."

We proceed to consider case (i), and wish to point out that for this case g can be arbitrary (we will need the flatness/transversality restrictions in case (ii)). In what follows we set $d_X = \dim X$, $d_Y = \dim Y$ and similarly for X', Y'. Using Theorem 2.32 we obtain a compactification $\iota: X \hookrightarrow \overline{X}$ over k of the smooth, separated and finite type k-scheme X in the upper right corner of (3.2) and (3.6).

This results in a compactification of the square (1) in (3.6) which we write as

$$(X \times Y', \operatorname{pr}_{Y'}^* \Delta_Y, \operatorname{pr}_{Y'*}^{-1}(\Phi_Y')) \xrightarrow{\operatorname{id} \times g} (X \times Y, \operatorname{pr}_Y^* \Delta_Y, \operatorname{pr}_{Y}^{-1}(\Phi_Y))$$

$$\downarrow_{\iota \times \operatorname{id}} \qquad \qquad \downarrow_{\iota \times \operatorname{id}}$$

$$(\overline{X} \times Y', \overline{\operatorname{pr}}_{Y'}^* \Delta_Y, \overline{\operatorname{pr}}_{Y'*}^{-1}(\Phi_Y')) \xrightarrow{\operatorname{id} \times g} (\overline{X} \times Y, \overline{\operatorname{pr}}_Y^* \Delta_Y, \overline{\operatorname{pr}}_{Y}^{-1}(\Phi_Y))$$

$$\downarrow_{\overline{\operatorname{pr}}_{Y'}} \qquad \qquad \downarrow_{\overline{\operatorname{pr}}_Y}$$

$$(Y', \Delta_{Y'}, \Phi_{Y'}) \xrightarrow{g} (Y, \Delta_Y, \Phi_Y)$$

$$(3.7)$$

By the description following Lemma 2.28, we know that

$$\operatorname{pr}_{Y*}: H^*(X \times Y, \operatorname{pr}_Y^* \Delta_Y, \operatorname{pr}_{Y*}^{-1}(\Phi_Y)) \to H^*(Y, \Delta_Y, \Phi_Y)$$

stems from a morphism

$$R\overline{\mathrm{pr}}_{Y*}R\mathcal{H}om_{\overline{X}\times Y}(\Omega^{p}_{\overline{X}\times Y}(\log\mathrm{pr}_{Y}^{*}\Delta_{Y})(-\mathrm{pr}_{Y}^{*}\Delta_{Y})[p],\omega_{\overline{X}\times Y}^{\bullet})\to\Omega^{d_{Y}-p}_{Y}(\log\Delta_{Y})[d_{Y}-p] \tag{3.8}$$

obtained as the Grothendieck dual of a log differential of \overline{pr}_Y (here and throughout what follows, a similar statement holds for $\overline{pr}_{Y'}$). By an observation of Chatzistamatiou-Rülling , this map factors as

$$R\overline{\operatorname{pr}}_{Y*}R\mathcal{H}om_{\overline{X}\times Y}(\Omega_{\overline{X}\times Y}^{p}(\log \overline{\operatorname{pr}}_{Y}^{*}\Delta_{Y})(-\overline{\operatorname{pr}}_{Y}^{*}\Delta_{Y})[p], \omega_{\overline{X}\times Y}^{\bullet})$$

$$\to R\overline{\operatorname{pr}}_{Y*}R\mathcal{H}om_{\overline{X}\times Y}(L\overline{\operatorname{pr}}_{Y}^{*}\Omega_{Y}^{p}(\log \Delta_{Y})(-\Delta_{Y})[p], \omega_{\overline{X}\times Y}^{\bullet})$$

$$\xrightarrow{\simeq} R\mathcal{H}om_{Y}(\Omega_{Y}^{p}(\log \Delta_{Y})(-\Delta_{Y})[p], R\overline{\operatorname{pr}}_{Y*}\omega_{\overline{X}\times Y}^{\bullet})$$

$$\xrightarrow{\operatorname{trace}} R\mathcal{H}om_{Y}(\Omega_{Y}^{p}(\log \Delta_{Y})(-\Delta_{Y})[p], \omega_{Y}[d_{Y}])$$

$$\xrightarrow{\simeq} \Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})[d_{Y}-p]$$

$$(3.9)$$

where the adjunction isomorphism is [R&D, Prop. II.5.10], and the map labeled trace is induced by the Grothendieck trace $R\overline{\mathrm{pr}}_{Y*}\omega_{\overline{X}\times Y}^{\bullet}\to \omega_{Y}[d_{Y}]$. If it were the case that \overline{X} were smooth, then the usual "box product" decomposition

$$\omega_{\overline{X}\times Y}^{\bullet} \simeq \omega_{\overline{X}}[d_X] \boxtimes \omega_Y[d_Y] := \operatorname{pr}_{\overline{X}}^{*} \omega_{\overline{X}}[d_X] \otimes \overline{\operatorname{pr}}_{Y*} \omega_Y[d_Y]$$

together with the perect pairings (2.31) and the local freeness of $\Omega_Y^p(\log \Delta_Y)(-\Delta_Y)[p]$ would give an identification

$$R\mathcal{H}om_{\overline{X}\times Y}(L\overline{\operatorname{pr}}_{Y}^{*}\Omega_{Y}^{p}(\log \Delta_{Y})(-\Delta_{Y})[p], \omega_{\overline{X}\times Y}^{\bullet}) \simeq \operatorname{pr}_{\overline{X}}^{*}\omega_{\overline{X}}[d_{X}] \otimes \overline{\operatorname{pr}}_{Y}^{*}\Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})[d_{Y}-p] \quad (3.10)$$

In fact a more careful version of this argument, carrying out the above calculation on the smooth locus $X \times Y$ and using excision, shows that $H^*(X \times Y, \operatorname{pr}_Y^* \Delta_Y, \operatorname{pr}_{Y^*}^{-1}(\Phi_Y)) \to H^*(Y, \Delta_Y, \Phi_Y)$ always factors through the summand $H^*_{\Phi_X}(X \times Y, \operatorname{pr}_{\overline{X}}^* \omega_{\overline{X}} \otimes \overline{\operatorname{pr}}_Y^* \Omega_Y^{d_Y - p}(\log \Delta_Y))$.

Our next lemma implies that even when \overline{X} is not known to be smooth, (3.8) still factors through something like $R\overline{\operatorname{pr}}_{Y*}(\operatorname{pr}_{\overline{X}}^*\omega_{\overline{X}}[d_X]\otimes \overline{\operatorname{pr}}_Y^*\Omega_Y^{d_Y-p}(\log \Delta_Y)[d_Y-p])$, provided we replace $\operatorname{pr}_{\overline{X}}^*\omega_{\overline{X}}[d_X]$ with $\overline{\operatorname{pr}}_Y^!$ \mathfrak{O}_Y .

Lemma 3.11 (compare with [CR11, Lem. 2.2.16]). For each p there is a natural map

$$\gamma: \overline{\mathrm{pr}}_{Y}^{!} \odot_{Y} \otimes \overline{\mathrm{pr}}_{Y}^{*} \Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p] \to R\mathcal{H}om_{\overline{X} \times Y}(\overline{\mathrm{pr}}_{Y}^{*} \Omega_{Y}^{p}(\log \Delta_{Y})(-\Delta_{Y})[p], \omega_{\overline{Y} \times Y}^{\bullet})$$

such that the restriction of γ to $X \times Y$ agrees with the isomorphism

 $\operatorname{pr}_X^* \omega_X[d_X] \otimes \operatorname{pr}_Y^* \Omega_Y^{d_Y-p}(\log \Delta_Y)(-\Delta_Y)[d_Y-p] \xrightarrow{\simeq} R\mathcal{H}om_{X\times Y}(L\operatorname{pr}_Y^* \Omega_Y^p(\log \Delta_Y)(-\Delta_Y)[p], \omega_{X\times Y}^{\bullet})$ and such that the composition

$$R\overline{\operatorname{pr}}_{Y*}(\operatorname{pr}_{X}^{*} \omega_{X}[d_{X}] \otimes \operatorname{pr}_{Y}^{*} \Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p])$$

$$\xrightarrow{R\overline{\operatorname{pr}}_{Y*}(Y)} R\overline{\operatorname{pr}}_{Y*}R\mathcal{H}om_{X\times Y}(\operatorname{pr}_{Y}^{*} \Omega_{Y}^{p}(\log \Delta_{Y})(-\Delta_{Y})[p], \omega_{X\times Y}^{\bullet})$$

$$\xrightarrow{\cong} R\mathcal{H}om_{X\times Y}(\Omega_{Y}^{p}(\log \Delta_{Y})(-\Delta_{Y})[p], R\overline{\operatorname{pr}}_{Y*}\omega_{X\times Y}^{\bullet})$$

$$\xrightarrow{trace} R\mathcal{H}om_{X\times Y}(\Omega_{Y}^{p}(\log \Delta_{Y})(-\Delta_{Y})[p], \omega_{Y}[d_{Y}]) \simeq \Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p]$$

$$(3.12)$$

coincides with the composition

$$R\overline{\operatorname{pr}}_{Y*}(\overline{\operatorname{pr}}_{Y}^{!} \mathfrak{G}_{Y} \otimes \overline{\operatorname{pr}}_{Y}^{*} \Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p])$$

$$\xrightarrow{\operatorname{proj.}} R\overline{\operatorname{pr}}_{Y*}(\overline{\operatorname{pr}}_{Y}^{!} \mathfrak{G}_{Y}) \otimes \overline{\operatorname{pr}}_{Y}^{*} \Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p]$$

$$\xrightarrow{\operatorname{tr} \otimes \operatorname{id}} \Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p]$$

$$(3.13)$$

By base change for dualizing complexes ([Stacks, Tag 0BZX, Tag 0E2S]) applied to the cartesian diagram

$$\overline{X} \times Y \longrightarrow \overline{X}$$

$$\downarrow \qquad \qquad \downarrow$$

$$Y \longrightarrow \operatorname{Spec} k$$

(note that this is a very mild situation: $\overline{X} \to \operatorname{Spec} k$ is flat and proper and $Y \to \operatorname{Spec} k$ is smooth) we see that $\overline{\operatorname{pr}}_Y^! \mathcal{O}_Y \simeq \operatorname{pr}_{\overline{X}}^* \omega_{\overline{X}}^{\bullet}$. This makes the map γ look even more like (3.10).

Proof. Following [CR11, Lem. 2.2.16] we begin with the morphism

$$e: \overline{\operatorname{pr}}_{Y}^{!} \mathfrak{G}_{Y} \otimes^{L} L \overline{\operatorname{pr}}_{Y}^{*} \omega_{Y}^{\bullet} \to \overline{\operatorname{pr}}_{Y}^{!} \omega_{Y}^{\bullet} =: \omega_{\overline{X} \times Y}^{\bullet}$$

of [Con00, p. 4.3.12], which as explained therein agrees with

$$\operatorname{pr}_X^* \omega_X[d_X] \otimes \operatorname{pr}_Y^* \omega_Y[d_Y] \xrightarrow{\simeq} \omega_{X \times Y}[d_X + d_Y]$$

on locus $X \times Y$, and has the property that

$$R\overline{pr}_{Y*}(\overline{\operatorname{pr}}_{Y}^{!} \mathbb{O}_{Y} \otimes^{L} L\overline{\operatorname{pr}}_{Y}^{*} \omega_{Y}^{\bullet}) \xrightarrow{R\overline{pr}_{Y*}e} R\overline{pr}_{Y*} \omega_{\overline{X} \times Y}^{\bullet}$$

$$\downarrow \operatorname{proj. form} \qquad \qquad \downarrow \operatorname{tr}$$

$$R\overline{pr}_{Y*}\overline{\operatorname{pr}}_{Y}^{!} \mathbb{O}_{Y} \otimes^{L} \omega_{Y}^{\bullet} \xrightarrow{\operatorname{tr} \otimes \operatorname{id}} \omega_{Y}^{\bullet}$$

⁷See Conrad's comment "It is easy to check that e_f coincides with (3.3.21) in the smooth case and is compatible with composites in f (using (4.3.6)."

commutes [Con00, Thm. 4.4.1]. We then define our version of γ as the composition

$$\overline{\operatorname{pr}}_{Y}^{l} \circ_{Y} \otimes^{L} L \overline{\operatorname{pr}}_{Y}^{*} \Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p]$$

$$\underline{\operatorname{id} \otimes^{L}(2.31)} \quad \overline{\operatorname{pr}}_{Y}^{l} \circ_{Y} \otimes^{L} L \overline{\operatorname{pr}}_{Y}^{*} R \mathcal{H} o m_{Y}(\Omega_{Y}^{p}(\log \Delta_{Y})[p], \omega_{Y}^{\bullet})$$

$$\underline{\operatorname{functoriality}} \quad R \mathcal{H} o m_{\overline{X} \times Y}(L \overline{\operatorname{pr}}_{Y}^{*} \Omega_{Y}^{p}(\log \Delta_{Y})[p], \overline{\operatorname{pr}}_{Y}^{l} \circ_{Y} \otimes^{L} \omega_{Y}^{\bullet})$$

$$\underline{\operatorname{induced by}} \quad R \mathcal{H} o m_{\overline{X} \times Y}(L \overline{\operatorname{pr}}_{Y}^{*} \Omega_{Y}^{p}(\log \Delta_{Y})[p], \omega_{\overline{X} \times Y}^{\bullet})$$

$$\underline{\operatorname{induced by}} \quad R \mathcal{H} o m_{\overline{X} \times Y}(L \overline{\operatorname{pr}}_{Y}^{*} \Omega_{Y}^{p}(\log \Delta_{Y})[p], \omega_{\overline{X} \times Y}^{\bullet})$$
(3.14)

Note that we may drop the "L"s as $\Omega_Y^{d_Y-p}(\log \Delta_Y)(-\Delta_Y)$ and $\Omega_Y^p(\log \Delta_Y)$ are locally free. Verification of the stated compatibilities is as in [CR11, Lem. 2.2.16].

Remark 3.15. It seems like we could have also used the more general version of [Con00, p. 4.3.12]

$$e': \overline{\operatorname{pr}}_Y^! \mathfrak{G}_Y \otimes^L L \overline{\operatorname{pr}}_Y^* \Omega_Y^{d_Y-p} (\log \Delta_Y) (-\Delta_Y) [d_Y-p] \to \overline{\operatorname{pr}}_Y^! \Omega_Y^{d_Y-p} (\log \Delta_Y) (-\Delta_Y) [d_Y-p]$$

together with the description

$$\overline{\operatorname{pr}}_Y^! \Omega_Y^{d_Y - p} (\log \Delta_Y) (-\Delta_Y) [d_Y - p] = D_{\overline{X} \times Y} (L \overline{\operatorname{pr}}_Y^* D_Y (\Omega_Y^{d_Y - p} (\log \Delta_Y) (-\Delta_Y) [d_Y - p]))$$

where $D_Y(-) = R\mathcal{H}om(-, \omega_Y^{\bullet})$ and similarly for $D_{\overline{X} \times Y}$.

Using this modified γ , we obtain a modified version of the diagram [CR11, p. 732 during Lem. 2.3.4], namely (3.16) in Figure 1). To make this diagram legible, we use a few abbreviations: all functors are derived, we use the dualizing functors of the form $D_Y(-) = R\mathcal{H}om_Y(-, \omega_Y^{\bullet})$ and we let $d = d_X + d_Y$. Lemma 3.11 shows that triangles involving γ commute, and (3.9) gives commutativity of the rest of the diagram. The usefulness of this diagram is that by *definition* beginning in the top left corner and following the path $\rightarrow \downarrow$ we obtain the pushforward on Hodge cohomology

$$\operatorname{pr}_{Y*} \underline{\Gamma}_{\operatorname{pr}_{V*}^{-1} \Phi_{Y}} \Omega_{X \times Y}^{d-p} (\log \operatorname{pr}_{Y}^{*} \Delta_{Y}) [d-p] \to \underline{\Gamma}_{\Phi_{Y}} \Omega_{\times Y}^{d_{Y}-p} (\log \Delta_{Y}) (-\Delta_{Y}) [d_{Y}-p]$$

but following $\downarrow \rightarrow$ gives a composition whose behavior with respect to (3.7) is easier to analyze. Namely, we have a diagram like (3.16) on Y', and in fact a map from (3.16) to g_* of the analogous diagram on Y', and hence from the preceding discussion it will suffice to prove commutativity of (3.17) of Figure 1.

Applying excision together with Lemma 3.11 we may rewrite the top row of (3.17) as

$$R \operatorname{pr}_{Y*} R\underline{\Gamma}_{\operatorname{pr}_{Y*}^{-1} \Phi_{Y}} \Omega_{X \times Y}^{d-p}(\log \operatorname{pr}_{Y}^{*} \Delta_{Y})[d-p]$$

$$\xrightarrow{\operatorname{project}} R \operatorname{pr}_{Y*} R\underline{\Gamma}_{\operatorname{pr}_{Y*}^{-1} \Phi_{Y}}(\operatorname{pr}_{X}^{*} \omega_{X}[d_{X}] \otimes \operatorname{pr}_{Y}^{*} \Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p])$$

$$\xrightarrow{\operatorname{proj.}} R \operatorname{pr}_{Y*} R\underline{\Gamma}_{\operatorname{pr}_{Y*}^{-1} \Phi_{Y}}(\operatorname{pr}_{X}^{*} \omega_{X}[d_{X}]) \otimes \Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p]$$

$$\xrightarrow{\operatorname{tr} \otimes \operatorname{id}} R\underline{\Gamma}_{\Phi_{Y}} \Omega_{Y}^{d_{Y}-p}(\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p]$$

$$(3.18)$$

where the first map is induced by a projection


$$\Omega^{d-p}_{V \vee V}(\log \operatorname{pr}_V^* \Delta_Y)[d-p] \to \operatorname{pr}_V^* \omega_X[d_X] \otimes \operatorname{pr}_V^* \Omega_V^{d_Y-p}(\log \Delta_Y)(-\Delta_Y)[d_Y-p]$$

coming from a Künneth-type decomposition of $\Omega_{X\times Y}^{d-p}(\log \operatorname{pr}_Y^*\Delta_Y)$, the second is the projection formula, and the last map is induced by a trace map with supports defined as the composition

$$R \operatorname{pr}_{Y*} R \underline{\Gamma}_{\operatorname{pr}_{Y*}^{-1} \Phi_{Y}} (\operatorname{pr}_{X}^{*} \omega_{X}[d_{X}]) \xrightarrow{\operatorname{excision}} R \overline{\operatorname{pr}}_{Y*} R \underline{\Gamma}_{\overline{\operatorname{pr}}_{Y*}^{-1} \Phi_{Y}} (\overline{\operatorname{pr}}_{Y}^{!} \mathfrak{G}_{Y})$$

$$\xrightarrow{\operatorname{Proposition 2.9}} R \underline{\Gamma}_{\Phi_{Y}} R \overline{\operatorname{pr}}_{Y*} (\overline{\operatorname{pr}}_{Y}^{!} \mathfrak{G}_{Y}) \xrightarrow{\operatorname{tr}} R \underline{\Gamma}_{\Phi_{Y}} \mathfrak{G}_{Y}$$

$$(3.19)$$



$$\operatorname{pr}_{Y*} \underbrace{\Gamma_{\overline{\operatorname{pr}}_{Y*}^{-1} \Phi_{Y}}^{-1} \Omega_{X \times Y}^{d-p} (\log \operatorname{pr}_{Y}^{*} \Delta_{Y})[d-p]} \xrightarrow{} \overline{\operatorname{pr}}_{Y*} \underbrace{\Gamma_{\overline{\operatorname{pr}}_{Y*}^{-1} \Phi_{Y}}^{-1} (\overline{\operatorname{pr}}_{Y}^{-1} \Theta_{Y} \otimes \overline{\operatorname{pr}}_{Y}^{*} \Omega_{Y}^{d_{Y}-p} (\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p])} \xrightarrow{} \underbrace{\Gamma_{\Phi_{Y}} \Omega_{Y}^{d_{Y}-p} (\log \Delta_{Y})(-\Delta_{Y})[d_{Y}-p]} \xrightarrow{} \underbrace{\Gamma_$$

Figure 1: Modified versions of diagrams appearing in the proof of [CR11, Lem. 2.3.4] (all functors derived)

Here the second map comes from the functoriality properties of Proposition 2.9, since there is an inclusion $\operatorname{pr}_{Y*}^{-1}\Phi_Y\subseteq\operatorname{pr}_Y^{-1}\Phi_Y$. The decomposition (3.18) maps to a similar decomposition of the bottom row of (3.17), and the only commutativity not guaranteed by standard functoriality properties (e.g. functoriality of the projection formula appearing in the second map of (3.18)) is that of

$$R \operatorname{pr}_{Y*} R \underline{\Gamma}_{\operatorname{pr}_{Y*}^{-1} \Phi_{Y}} (\operatorname{pr}_{X}^{*} \omega_{X}[d_{X}]) \otimes \Omega_{Y}^{d_{Y}-p} (\log \Delta_{Y}) (-\Delta_{Y}) [d_{Y}-p] \xrightarrow{\operatorname{tr} \otimes \operatorname{id}} R \underline{\Gamma}_{\Phi_{Y}} \Omega_{Y}^{d_{Y}-p} (\log \Delta_{Y}) (-\Delta_{Y}) [d_{Y}-p]$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$R g_{*}(R \operatorname{pr}_{Y'*} R \underline{\Gamma}_{\operatorname{pr}_{Y'*}^{-1} \Phi_{Y'}} (\operatorname{pr}_{X}^{*} \omega_{X}[d_{X}]) \otimes \Omega_{Y'}^{d_{Y}-p} (\log \Delta_{Y'}) (-\Delta_{Y'}) [d_{Y}-p]) \xrightarrow{\operatorname{tr} \otimes \operatorname{id}} R g_{*}(R \underline{\Gamma}_{\Phi_{Y'}} \Omega_{Y'}^{d_{Y}-p} (\log \Delta_{Y'}) (-\Delta_{Y'}) [d_{Y}-p])$$

$$(3.20)$$

But applying one more projection formula to the bottom row of (3.20), we see (3.20) is obtained by tensoring the differential

$$\Omega_Y^{d_Y-p}(\log \Delta_Y)(-\Delta_Y)[d_Y-p] \to Rg_*\Omega_{Y'}^{d_Y-p}(\log \Delta_{Y'})(-\Delta_{Y'})[d_Y-p]$$

with

$$R \operatorname{pr}_{Y_{*}} R\underline{\Gamma}_{\operatorname{pr}_{Y_{*}}^{-1} \Phi_{Y}}(\operatorname{pr}_{X}^{*} \omega_{X}[d_{X}]) \xrightarrow{\operatorname{tr} \otimes \operatorname{id}} R\underline{\Gamma}_{\Phi_{Y}} \mathfrak{G}_{Y}$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$Rg_{*}(R \operatorname{pr}_{Y'_{*}} R\underline{\Gamma}_{\operatorname{pr}_{Y'_{*}}^{-1} \Phi_{Y'}}(\operatorname{pr}_{X}^{*} \omega_{X}[d_{X}]))^{\operatorname{tr}' \otimes \operatorname{id}} Rg_{*}(R\underline{\Gamma}_{\Phi_{Y'}} \mathfrak{G}_{Y'})$$

$$(3.21)$$

and the commutativity of (3.21) is proved in [CR11, Lem. 2.3.4]. So far we have proved:

Lemma 3.22. Lemma 3.1 holds in case (i) of Lemma 3.4.

It remains to deal with case (ii) of Lemma 3.4, and for this we use the following lemma.

Lemma 3.23 (compare with [CR11, Cor. 2.2.22]). Consider a diagram of pure-dimensional snc pairs

$$(X', \Delta_{X'}) \xrightarrow{g'} (X, \Delta_X)$$

$$\downarrow^{l'} \qquad \downarrow^{l}$$

$$(Y', \Delta_{Y'}) \xrightarrow{g} (Y, \Delta_Y)$$

$$(3.24)$$

where ι, ι' are pushing closed immersions and $\dim Y - \dim X = \dim Y' - \dim X' = : c$. Then, for all q the diagram

$$\iota_*\Omega_X^q(\log \Delta_X)[q] \xrightarrow{dg^{\vee}} Rg_*\iota_*'\Omega_{X'}^q(\log \Delta_{X'})$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$\Omega_Y^{q+c}(\log \Delta_Y)[q+c] \xrightarrow{dg^{\vee}} Rg_*\Omega_{Y'}^{q+c}(\log \Delta_{Y'})[q+c]$$

$$(3.25)$$

commutes, where the horizontal maps are induced by log differentials and the left vertical map is the composition

$$\iota_* \Omega_X^q (\log \Delta_X)[q] \xrightarrow{\simeq} \iota_* R \mathcal{H}om(\Omega_X^{d_X - q}(\log \Delta_X)(-\Delta_X)[d_X - q], \omega_X^{\bullet}) \\
\xrightarrow{duality} R \mathcal{H}om(\iota_* \Omega_X^{d_X - q}(\log \Delta_X)(-\Delta_X)[d_X - q], \omega_Y^{\bullet}) \\
\xrightarrow{d\iota^{\vee}} R \mathcal{H}om(\Omega_Y^{d_X - q}(\log \Delta_Y)(-\Delta_Y)[d_X - q], \omega_Y^{\bullet}) \\
\xrightarrow{\simeq} \Omega_Y^{q+c}(\log \Delta_Y)[q+c]$$
(3.26)

and the right vertical arrow is Rg_* of a similar composition on Y'.

Note that the codimension hypotheses hold if g is flat or a closed immersion transverse to ι .

Proof. While it seems a proof following [CR11, Cor. 2.2.22] step-by-step is possible, we instead *reduce* to the case proved there as follows: first, observe that there is an evident map from the cartesian diagram

$$U_{X'} \longrightarrow U_X$$

$$\downarrow \qquad \qquad \downarrow$$

$$U_{Y'} \longrightarrow U_Y$$

$$(3.27)$$

of interiors to (3.24). Noting that (3.25) will map to a similar diagram obtained from (3.27), that the compositions (3.26) are at least compatible with Zariski localization, *and* that the situation of (3.27) is covered by [CR11, Cor. 2.2.22], it will suffice to show that the natural map

$$h^{0}R\mathcal{H}om_{Y}(\iota_{*}\Omega_{X}^{q}(\log \Delta_{X})[q], Rg_{*}\Omega_{Y'}^{q+c}(\log \Delta_{Y'})[q+c]) \rightarrow h^{0}R\mathcal{H}om_{U_{Y}}(\iota_{*}\Omega_{U_{X}}^{q}[q], Rg_{*}\Omega_{U_{Y'}}^{q+c}[q+c])$$
(3.28)

is *injective*. This can be checked Zariski-locally at a point $x \in X \subseteq Y$, so we may assume $X \subseteq Y$ is a global complete intersection, say of $t_1, \ldots, t_c \in \mathcal{O}_Y$. In that case the t_i define a Koszul resolution $\mathcal{K}^{\bullet}(t_i) \to \mathcal{O}_X$, and *because* $X' = Y' \times_Y X = V(t_1 \circ g, \cdots t_c \circ g)$ is smooth of codimension c by hypotheses, it must be that the $t_i \circ g$ are also a regular sequence, hence

$$L^{i}g^{*}\mathcal{O}_{X} = h^{-i}g^{*}\mathcal{K}^{\bullet}(t_{i}) = \begin{cases} \mathcal{O}_{X'}, & i = 0\\ 0 & \text{otherwise} \end{cases}$$

in other words $Lg^* \mathfrak{O}_X = \mathfrak{O}_{X'}$. Now using the fact that $\Omega_X^q(\log \Delta_X)$ is locally free on X' we conclude

$$Lg^*\iota_*\Omega_Y^q(\log \Delta_X)[q] = g^*\iota_*\Omega_X^q(\log \Delta_X)[q] = \iota_*'g'^*\Omega_X^q(\log \Delta_X)[q]$$

Next, applying derived adjunction to both sides of (3.28) gives a commutative diagram

$$R\mathcal{H}om_{Y}(\iota_{*}\Omega_{X}^{q}(\log \Delta_{X})[q], Rg_{*}\Omega_{Y'}^{q+c}(\log \Delta_{Y'})[q+c]) \longrightarrow R\mathcal{H}om_{U_{Y}}(\iota_{*}\Omega_{U_{X}}^{q}[q], Rg_{*}\Omega_{U_{Y'}}^{q+c}[q+c])$$

$$\parallel \qquad \qquad \parallel \qquad \qquad \parallel$$

$$Rg_{*}R\mathcal{H}om_{Y'}(Lg^{*}\iota_{*}\Omega_{X}^{q}(\log \Delta_{X})[q], \Omega_{Y'}^{q+c}(\log \Delta_{Y'})[q+c]) \longrightarrow Rg_{*}R\mathcal{H}om_{U_{Y'}}(Lg^{*}\iota_{*}\Omega_{U_{X}}^{q}[q], \Omega_{U_{Y'}}^{q+c}[q+c])$$

$$\parallel \qquad \qquad \parallel \qquad \qquad \parallel$$

$$Rg_{*}R\mathcal{H}om_{Y'}(\iota'_{*}g'^{*}\Omega_{X}^{q}(\log \Delta_{X})[q], \Omega_{Y'}^{q+c}(\log \Delta_{Y'})[q+c]) \longrightarrow Rg_{*}R\mathcal{H}om_{U_{Y'}}(\iota'_{*}g'^{*}\Omega_{U_{X}}^{q}[q], \Omega_{U_{Y'}}^{q+c}[q+c])$$

$$(3.2)$$

Getting even more Zariski-local we may assume $\Omega_X^q(\log \Delta_X)$ is *free*, say generated by dx_1, \dots, dx_n and in that case

$$R\mathcal{H}om_{Y'}(\iota_{*}'g'^{*}\Omega_{X}^{q}(\log \Delta_{X})[q], \Omega_{Y'}^{q+c}(\log \Delta_{Y'})[q+c])$$

$$= (\prod_{i} R\mathcal{H}om_{Y'}(\mathcal{O}_{X'}dx_{i}[q], \mathcal{O}_{Y'}[q+c])) \otimes \Omega_{Y'}^{q+c}(\log \Delta_{Y'})$$
(3.30)

and by Grothendieck's fundamental local isomorphism [Con00, §2.5]

$$R\mathcal{H}om_{Y'}(\mathcal{O}_{X'}[q], \mathcal{O}_{Y'}[q+c])) \simeq \mathcal{E}xt_{Y'}^{c}(\mathcal{O}_{X'}, \mathcal{O}_{Y'}) \simeq \det(\mathcal{I}_{X'}/\mathcal{I}_{X'})^{\vee}$$
(3.31)

(the last 2 as sheaves supported in degree 0). In particular, this is an *invertible sheaf on* X', and it follows that the left hand side of (3.30) is a locally free sheaf (supported in degree 0) on X'. Recalling X' is smooth and so in particular reduced, and since $U_{Y'} \cap X'$ is a dense open (this is part of the

hypothesis that $X' \to Y'$ is a pulling map) the natural map

$$h^{0}R\mathcal{H}om_{Y'}(\iota'_{*}g'^{*}\Omega_{X}^{q}(\log \Delta_{X})[q], \Omega_{Y'}^{q+c}(\log \Delta_{Y'})[q+c])$$

$$\to h^{0}R\mathcal{H}om_{Y'}(\iota'_{*}g'^{*}\Omega_{X}^{q}(\log \Delta_{X})[q], \Omega_{Y'}^{q+c}(\log \Delta_{Y'})[q+c])|_{U_{Y'}}$$

$$\simeq h^{0}R\mathcal{H}om_{U_{Y'}}(\iota'_{*}g'^{*}\Omega_{X}^{q}(\log \Delta_{X})|_{U_{Y'}}[q], \Omega_{Y'}^{q+c}(\log \Delta_{Y'})|_{U_{Y'}}[q+c])$$
(3.32)

is injective, where on the third line we have applied localization for $\mathcal{E}xt$. Now left-exactness of g_* gives an injection

$$h^{0}Rg_{*}R\mathcal{H}om_{Y'}(\iota'_{*}g'^{*}\Omega_{X}^{q}(\log \Delta_{X})[q], \Omega_{Y'}^{q+c}(\log \Delta_{Y'})[q+c])$$

$$\to h^{0}Rg_{*}R\mathcal{H}om_{U_{Y'}}(\iota'_{*}g'^{*}\Omega_{X}^{q}(\log \Delta_{X})|_{U_{Y'}}[q], \Omega_{Y'}^{q+c}(\log \Delta_{Y'})|_{U_{Y'}}[q+c])$$
(3.33)

To complete the proof, we use (3.29) to identify the map (3.33) with (3.28).

Corollary 3.34. Lemma 3.1 holds in case (ii) of Lemma 3.4.

Proof. This follows by applying cohomology with supports to (3.25).

This completes our proof of Lemma 3.1.

Corollary 3.35 (projection formula, compare with [CR11, Prop. 1.1.16]). Let $f: X \to Y$ be a map of smooth schemes admitting two different enhancements to maps of smooth schemes with supports,

$$(X, \Delta_X, \Phi_X) \rightarrow (Y, \Delta_Y, f(\Phi_X))$$
 pushing and $(X, f^*(\Delta_Y'), f^{-1}(\Phi_Y)) \rightarrow (Y, \Delta_Y', \Phi_Y)$ pulling

Assume in addition that $\Delta_X + f^*(\Delta_Y')$ and $\Delta_Y + \Delta_Y'$ are (reduced) snc divisors. Then

$$(X, \Delta_X + f^*(\Delta_Y'), \Phi_X \cap f^{-1}(\Phi_Y)) \to (Y, \Delta_Y + \Delta_Y', f(\Phi_X) \cap \Phi_Y)$$

is also a pushing map, and

$$f_*(\beta \smile f^*\alpha) = f_*\beta \smile \alpha \in H^*(Y, \Delta_Y + \Delta_Y', f(\Phi_X) \cap \Phi_Y)$$

for any $\alpha \in H^*(Y, \Delta_Y', \Phi_Y)$ and $\beta \in (X, \Delta_X, \Phi_X)$, where \smile is the cup product on log Hodge cohomology defined along the lines of [CR11, §1.1.4, 2.4]

Proof. This is a formal consequence of Lemma 3.1 and can be derived following the proof of [CR11, Prop. 1.1.16]. Again we use a factorization through the graph like

$$(X, \Delta_{X} + f^{*}(\Delta'_{Y}), \Phi_{X} \cap f^{-1}(\Phi_{Y})) \xrightarrow{f} (Y, \Delta_{Y} + \Delta'_{Y}, f(\Phi_{X}) \cap \Phi_{Y})$$

$$\downarrow^{\mathrm{id}_{X} \times \mathrm{id}_{X}}$$

$$(X \times X, \mathrm{pr}_{1}^{*} \Delta_{X} + \mathrm{pr}_{2}^{*} f^{*}(\Delta'_{Y}), \Phi_{X} \times f^{-1}(\Phi_{Y}))$$

$$\downarrow^{\mathrm{id}_{X} \times f}$$

$$(X \times Y, \mathrm{pr}_{1}^{*} \Delta_{X} + \mathrm{pr}_{2}^{*} \Delta'_{Y}, \Phi_{X} \times \Phi_{Y}) \xrightarrow{f \times \mathrm{id}_{Y}} (Y \times Y, \mathrm{pr}_{1}^{*} \Delta_{Y} + \mathrm{pr}_{2}^{*} \Delta'_{Y}, f(\Phi_{X}) \times \Phi_{Y})$$

$$(3.36)$$

Here $f \times \operatorname{id}_Y$ on the bottom is a pushing morphism (since $f|_{\Phi_X}$ is proper and $f^*\Delta_Y = \Delta_X$) and the right vertical map $\operatorname{id}_Y \times \operatorname{id}_Y$ is a closed immersion transverse to $f \times \operatorname{id}_Y$ since the outer rectangle is cartesian and X is smooth of the correct codimension. This means we are in a situation to apply Lemma 3.1, and that lemma plus the definition of cup products in terms of pullbacks along diagonals gives the desired identity.

4 Correspondences

Given snc pairs with familes of supports (X, Δ_X, Φ_X) and (Y, Δ_Y, Φ_Y) with dimensions d_X and d_Y , as in [CR11, §1.3] we may define a family of supports $P(\Phi_X, \Phi_Y)$ on $X \times Y$ by

$$\begin{split} P(\Phi_X, \Phi_Y) := & \{ \text{closed subsets } Z \subseteq X \times Y \mid \operatorname{pr}_Y|_Z \text{ is proper and for all } W \in \Phi_X, \\ & \operatorname{pr}_Y(\operatorname{pr}_X^{-1}(W) \cap Z) \in \Phi_Y \} \end{split}$$

(the conditions of Definition 2.1 are straightforward to verify). For convenience we will let $\Delta_{X\times Y}:=\operatorname{pr}_X^*\Delta_X+\operatorname{pr}_Y^*\Delta_Y$.

Theorem 4.1. A class $\gamma \in H^j_{P(\Phi_Y,\Phi_Y)}(X \times Y, \Omega^i_{X \times Y}(\log \Delta_{X \times Y})(-\operatorname{pr}^*_X \Delta_X))$ defines homomorphisms

$$\operatorname{cor}(\gamma): H^q_{\Phi_X}(X,\Omega_X^p(\log \Delta_X)) \to H^{q+j-d_X}_{\Phi_Y}(Y,\Omega_Y^{p+i-d_X}(\log \Delta_Y))$$

by the formula $\operatorname{cor}(\gamma)(\alpha) := \operatorname{pr}_{Y*}(\operatorname{pr}_X^*(\alpha) \smile \gamma)$. Moreover if (Z, Δ_Z, Φ_Z) is another snc pair with supports and $\delta \in H^{j'}_{P(\Phi_V, \Phi_Z)}(Y \times Z, \Omega^{i'}_{Y \times Z}(\log \Delta_{Y \times Z})(-\operatorname{pr}_Y^* \Delta_Y))$, then

$$\operatorname{pr}_{X\times Z*}(\operatorname{pr}_{X\times Y}^*(\gamma) \smile \operatorname{pr}_{Y\times Z}^*(\delta)) \in H^{j+j'-d_Y}_{P(\Phi_X,\Phi_Z)}(X\times Z, \Omega_{X\times Z}^{i+i'-d_Y}(\log \Delta_{X\times Z})(-\operatorname{pr}_X^*\Delta_X)) \ and$$

$$\operatorname{cor}(\operatorname{pr}_{X\times Z*}(\operatorname{pr}_{Y\times Z}^*(\operatorname{pr}_{Y\times Y}^*(\gamma) \smile \operatorname{pr}_{Y\times Z}^*(\delta))) = \operatorname{cor}(\delta) \circ \operatorname{cor}(\gamma)$$

as homomorphisms
$$H^q_{\Phi_X}(X,\Omega_X^p(\log \Delta_X)) \to H^{q+j+j'-d_X-d_Y}_{\Phi_Z}(Z,\Omega_Z^{p+i+i'-d_X-d_Y}(\log \Delta_Z)).$$

Remark 4.2. The sheaves $\Omega^i_{X\times Y}(\log \Delta_{X\times Y})(-\operatorname{pr}_X^*\Delta_X)$ are particular instances of the sheaves $\Omega^i_X(A,B)$ appearing in [DI87, §4.2].

Such correspondences involving both log poles and "log zeroes" appear to have been considered before at least in crystalline cohomology, for example in work of Mieda [Mie09a; Mie09b]. However, I was unable to find any published proof of Theorem 4.1 in the literature.

Proof. We make two observations: first, using Lemma 2.18 there are natural wedge product pairings

$$\Omega^p_{X\times Y}(\log \Delta_{X\times Y}) \otimes \Omega^i_{X\times Y}(\log \Delta_{X\times Y})(-\operatorname{pr}_X^*\Delta_X) \xrightarrow{\wedge} \Omega^{p+i}_{X\times Y}(\log \Delta_Y)$$

Second, essentially by the definition of $P(\Phi_X, \Phi_Y)$ the Künneth morphism on cohomology for the tensor product $\Omega^p_{X\times Y}(\log \Delta_{X\times Y})\otimes \Omega^i_{X\times Y}(\log \Delta_{X\times Y})(-\operatorname{pr}^*_X\Delta_X)$ can be enhanced with supports as

$$\begin{split} H^q_{\mathrm{pr}_X^{-1}(\Phi_X)}(X \times Y, \Omega^p_{X \times Y}(\log \Delta_{X \times Y})) \otimes H^j_{P(\Phi_X, \Phi_Y)}(X \times Y, \Omega^i_{X \times Y}(\log \Delta_{X \times Y})(-\mathrm{pr}_X^* \Delta_X)) \\ & \to H^{p+j}_{\Psi}(X \times Y, \Omega^p_{X \times Y}(\log \Delta_{X \times Y}) \otimes \Omega^i_{X \times Y}(\log \Delta_{X \times Y})(-\mathrm{pr}_X^* \Delta_X)) \end{split}$$

where $\Psi:=\mathrm{pr}_{Y*}^{-1}(\Phi_Z)$ (see [CR11, §1.3.7, Prop. 1.3.10]). Combining these 2 observations gives a pairing

$$H^{q}_{\mathrm{pr}_{X}^{-1}(\Phi_{X})}(X \times Y, \Omega_{X \times Y}^{p}(\log \Delta_{X \times Y})) \otimes H^{j}_{P(\Phi_{X}, \Phi_{Y})}(X \times Y, \Omega_{X \times Y}^{i}(\log \Delta_{X \times Y})(-\mathrm{pr}_{X}^{*}\Delta_{X}))$$

$$\stackrel{\smile}{\longrightarrow} H^{p+j}_{W}(X \times Y, \Omega_{X \times Y}^{p+i}(\log \Delta_{Y}))$$

Now note that $\operatorname{pr}_X: (X\times Y, \Delta_{X\times Y}, \operatorname{pr}_X^{-1}(\Phi_X)) \to (X, \Delta_X, \Phi_X)$ is a pulling morphism, so by Proposition 2.24 there is an induced map $\operatorname{pr}_X^*: H^q_{\Phi_X}(X, \Omega_X^p(\log \Delta_X)) \to H^q_{\operatorname{pr}_X^{-1}(\Phi_X)}(X\times Y, \Omega_{X\times Y}^p(\log \Delta_{X\times Y}))$. On the other hand since $\operatorname{pr}_Y: (X\times Y, \Delta_Y, \Psi) \to (Y, \Delta_Y, \Phi_Y)$ is a pushing morphism, Lemma 2.28 provides

a morphism $\operatorname{pr}_{Y*}: H^{p+j}_{\Psi}(X\times Y, \Omega^{p+i}_{X\times Y}(\log \Delta_Y)) \to H^{q+j-d_X}_{\Phi_Y}(Y, \Omega^{p+i-d_X}_Y(\log \Delta_Y))$. Composing, we obtain the desired homomorphism

$$\begin{split} H^q_{\Phi_X}(X, \Omega_X^p(\log \Delta_X)) &\xrightarrow{\operatorname{pr}_X^*} H^q_{\operatorname{pr}_X^{-1}(\Phi_X)}(X \times Y, \Omega_{X \times Y}^p(\log \Delta_{X \times Y})) \\ &\xrightarrow{\sim \gamma} H^{p+j}_{\Psi}(X \times Y, \Omega_{X \times Y}^{p+i}(\log \Delta_Y)) \\ &\xrightarrow{\operatorname{pr}_{Y*}} H^{q+j-d_X}_{\Phi_Y}(Y, \Omega_Y^{p+i-d_X}(\log \Delta_Y)) \end{split}$$

For the "moreover" half of the lemma, we again begin with a certain wedge product pairing, this time on $X \times Y \times Z$:

$$\Omega_{X\times Y\times Z}^{i}(\log \operatorname{pr}_{X\times Y}^{*}\Delta_{X\times Y})(-\operatorname{pr}_{X}^{*}\Delta_{X}) \otimes \Omega_{X\times Y\times Z}^{i'}(\log \operatorname{pr}_{Y\times Z}^{*}\Delta_{Y\times Z})(-\operatorname{pr}_{Y}^{*}\Delta_{Y})
\stackrel{\wedge}{\to} \Omega_{X\times Y\times Z}^{i+i'}(\log \operatorname{pr}_{X\times Z}^{*}\Delta_{X\times Z})(-\operatorname{pr}_{X}^{*}\Delta_{X})$$
(4.3)

If $V \in P(\Phi_X, \Phi_Y)$, $W \in P(\Phi_Y, \Phi_Z)$ then unravelling definitions (again we refer to [CR11, §1.3.7, Prop. 1.3.10] for a similar claim) we find:

• $\operatorname{pr}_{X\times Z}|_{\operatorname{pr}_{X\times Y}^{-1}(V)\cap\operatorname{pr}_{Y\times Z}^{-1}(W)}$ is proper and • $\operatorname{pr}_{X\times Z}(\operatorname{pr}_{X\times Y}^{-1}(V)\cap\operatorname{pr}_{Y\times Z}^{-1}(W))\in P(\Phi_X,\Phi_Z)$ so that the Künneth morphism on cohomology associated to the left hand side of (4.3) can be enhanced with supports like

$$\begin{split} &H^{j}_{\mathrm{pr}_{X\times Y}^{-1}(P(\Phi_{X},\Phi_{Y}))}(X\times Y\times Z,\Omega_{X\times Y\times Z}^{i}(\log\mathrm{pr}_{X\times Y}^{*}\Delta_{X\times Y})(-\mathrm{pr}_{X}^{*}\Delta_{X}))\\ &\otimes H^{j'}_{\mathrm{pr}_{Y\times Z}^{-1}(P(\Phi_{Y},\Phi_{Z}))}(X\times Y\times Z,\Omega_{X\times Y\times Z}^{i'}(\log\mathrm{pr}_{Y\times Z}^{*}\Delta_{Y\times Z})(-\mathrm{pr}_{Y}^{*}\Delta_{Y}))\\ &\to H^{j+j'}_{\Sigma}(X\times Y\times Z,\Omega_{X\times Y\times Z}^{i}(\log\mathrm{pr}_{X\times Y}^{*}\Delta_{X\times Y})(-\mathrm{pr}_{X}^{*}\Delta_{X})\otimes\Omega_{X\times Y\times Z}^{i'}(\log\mathrm{pr}_{Y\times Z}^{*}\Delta_{Y\times Z})(-\mathrm{pr}_{Y}^{*}\Delta_{Y})) \end{split}$$

where $\Sigma := \operatorname{pr}_{X \times Z *}^{-1}(P(\Phi_X, \Phi_Z)).$

Since $\operatorname{pr}_{X\times Y}: (X\times Y\times Z,\operatorname{pr}_{X\times Y}^*\Delta_{X\times Y},\operatorname{pr}_{X\times Y}^{-1}(P(\Phi_X,\Phi_Y)))\to (X\times Y,\Delta_{X\times Y},P(\Phi_X,\Phi_Y))$ is a pulling morphism, Proposition 2.24 gives an induced morphism

$$\Omega^{i}_{X\times Y}(\log \Delta_{X\times Y}) \to Rf_*\Omega^{i}_{X\times Y\times Z}(\log \operatorname{pr}^*_{X\times Y}\Delta_{X\times Y});$$

twisting by $-\Delta_{X\times Y}$ and applying the projection formula gives a morphism

$$\Omega^i_{X\times Y}(\log \Delta_{X\times Y})(-\Delta_{X\times Y})\to Rf_*\big(\Omega^i_{X\times Y\times Z}(\log \mathrm{pr}^*_{X\times Y}\Delta_{X\times Y})(-\mathrm{pr}^*_{X\times Y}\Delta_{X\times Y})\big)$$

and then taking cohomology with supports along $P(\Phi_X, \Phi_Y)$ and using Proposition 2.9 gives a modified pullback map

$$H^{j}_{P(\Phi_{X},\Phi_{Y})}(X \times Y, \Omega^{i}_{X \times Y}(\log \Delta_{X \times Y})(-\Delta_{X \times Y}))$$

$$\to H^{j}_{\operatorname{pr}_{X \times Y}^{-1}(P(\Phi_{X},\Phi_{Y}))}(X \times Y \times Z, \Omega^{i}_{X \times Y \times Z}(\log \operatorname{pr}_{X \times Y}^{*}\Delta_{X \times Y})(-\operatorname{pr}_{X}^{*}\Delta_{X}))$$

$$(4.4)$$

and a similar argument gives a modified pullback

$$H_{P(\Phi_{Y},\Phi_{Z})}^{j'}(Y \times Z, \Omega_{Y \times Z}^{i'}(\log \Delta_{Y \times Z})(-\Delta_{Y \times Z}))$$

$$\to H_{\text{pr}_{Y \times Z}^{-1}(P(\Phi_{Y},\Phi_{Z}))}^{j'}(X \times Y \times Z, \Omega_{X \times Y \times Z}^{i'}(\log \text{pr}_{Y \times Z}^{*}\Delta_{Y \times Z})(-\text{pr}_{X}^{*}\Delta_{Y}))$$

$$(4.5)$$

On the other hand, $\operatorname{pr}_{X\times Z}:(X\times Y\times Z,\operatorname{pr}_{X\times Z}^*\Delta_{X\times Y},\Sigma)\to (X\times Z,\Delta_{X\times Z},P(\Phi_X,\Phi_Z))$ is a pushing morphism and hence by Lemma 2.28 induces morphisms

$$R\mathrm{pr}_{X\times Z*}R\underline{\Gamma}_{\Sigma}(\Omega^{\dim X\times Y\times Z-k}_{X\times Y\times Z}(\log \mathrm{pr}^*_{X\times Z}\Delta_{X\times Y}))\to R\underline{\Gamma}_{P(\Phi_{X},\Phi_{Z})}\Omega^{\dim X\times Z-k}_{X\times Z}(\log \Delta_{X\times Z})[-\dim Z]$$

for all k; twisting by $-pr_X^*\Delta_X$ and applying the projection formula this becomes

$$R \operatorname{pr}_{X \times Z *} R \underline{\Gamma}_{\Sigma} (\Omega_{X \times Y \times Z}^{\dim X \times Y \times Z - k} (\log \operatorname{pr}_{X \times Z}^{*} \Delta_{X \times Y}) (-\operatorname{pr}_{X}^{*} \Delta_{X}))$$

$$\to R \underline{\Gamma}_{P(\Phi_{X}, \Phi_{Z})} \Omega_{X \times Z}^{\dim X \times Z - k} (\log \Delta_{X \times Z}) (-\operatorname{pr}_{X}^{*} \Delta_{X}) [-\dim Z]$$

$$(4.6)$$

Now letting $k = \dim X \times Y \times Z - i - i'$, the induced morphisms of cohomology with supports are

$$H_{\Sigma}^{j+j'}(X \times Y \times Z, \Omega_{X \times Y \times Z}^{i+i'}(\log \operatorname{pr}_{X \times Z}^* \Delta_{X \times Y})(-\operatorname{pr}_X^* \Delta_X))$$

$$\to H_{P(\Phi_X, \Phi_Z)}^{j+j'-\dim Z}(X \times Z, \Omega_{X \times Z}^{i+i'-\dim Z}(\log \Delta_{X \times Z})(-\operatorname{pr}_X^* \Delta_X))$$
(4.7)

Combining the above ingredients, we obtain a bilinear pairing


$$H^{j}_{P(\Phi_{X},\Phi_{Y})}(X \times Y, \Omega^{i}_{X \times Y}(\log \Delta_{X \times Y})(-\Delta_{X \times Y})) \otimes H^{j'}_{P(\Phi_{Y},\Phi_{Z})}(Y \times Z, \Omega^{i'}_{Y \times Z}(\log \Delta_{Y \times Z})(-\Delta_{Y \times Z}))$$

$$\to H^{j+j'-\dim Z}_{P(\Phi_{X},\Phi_{Z})}(X \times Z, \Omega^{i+i'-\dim Z}_{X \times Z}(\log \Delta_{X \times Z})(-\operatorname{pr}^{*}_{X}\Delta_{X}))$$

sending $\gamma \otimes \delta \longmapsto \operatorname{pr}_{X \times Z*}(\operatorname{pr}^*_{X \times Y}(\gamma) \smile \operatorname{pr}^*_{Y \times Z}(\delta))$. It remains to be seen that

$$\operatorname{cor}(\operatorname{pr}_{X\times Z*}(\operatorname{pr}_{X\times Y}^*(\gamma)\smile\operatorname{pr}_{Y\times Z}^*(\delta)))=\operatorname{cor}(\delta)\circ\operatorname{cor}(\gamma)$$

and for this we will make repeated use of Lemma 3.1. Consider the diagram of smooth schemes



where all morphisms are projections. There are various ways to enhance this to include supports; here we add the family of supports Ψ on $X \times Y$ defined above. Then in the cartesian diagram (*), $\operatorname{pr}_Y: (X \times Y, \Psi) \to (Y, \Phi_Y)$ and $\operatorname{pr}_{Y \times Z}: (X \times Y \times Z, \operatorname{pr}_{X \times Y}^{-1} \Psi) \to (Y \times Z, \operatorname{pr}_Y^{-1} \Phi_Y)$ are pushing morphisms, whereas $\operatorname{pr}_{X \times Y}$ and pr_Y are pulling morphisms. At the same time, we have a pulling morphism $\operatorname{pr}_{X \times Z}: (X \times Y \times Z, \operatorname{pr}_{X \times Z}^{-1}(P(\Phi_Y, \Phi_Z))) \to (Y \times Z, P(\Phi_Y, \Phi_Z))$. To be precise in what follows, whenever ambiguity is possible we will use notation like $\operatorname{pr}_X^{X \times Y}$ to denote the projection $X \times Y \to X$, $\operatorname{pr}_X^{X \times Y \times Z}$ to denote the projection $X \times Y \times Z \to X$ and so on.

Applying Corollary 3.35 first to $pr_{X\times Z}$ we see that

$$\mathrm{pr}_{Y\times Z*}(\mathrm{pr}_{X\times Y}^*(\mathrm{pr}_X^{X\times Y*}\alpha\smile\gamma)\smile\mathrm{pr}_{Y\times Z}^*\delta)=\mathrm{pr}_{Y\times Z*}(\mathrm{pr}_{X\times Y}^*(\mathrm{pr}_X^{X\times Y*}\alpha\smile\gamma))\smile\delta$$

and then applying Lemma 3.1 to (*) shows

$$\mathrm{pr}_{Y\times Z*}(\mathrm{pr}_{X\times Y}^*(\mathrm{pr}_X^{X\times Y*}\alpha\smile\gamma))=\mathrm{pr}_Y^{Y\times Z*}(\mathrm{pr}_{Y*}^{X\times Y}(\mathrm{pr}_X^{X\times Y*}\alpha\smile\gamma))=\mathrm{pr}_Y^{Y\times Z*}\mathrm{cor}(\gamma)(\alpha)$$

so that

$$\mathrm{pr}_{Y\times Z*}(\mathrm{pr}_{X\times Y}^*(\mathrm{pr}_X^{X\times Y*}\alpha\smile\gamma)\smile\mathrm{pr}_{Y\times Z}^*\delta)=\mathrm{pr}_Y^{Y\times Z*}\operatorname{cor}(\gamma)(\alpha)\smile\delta$$

Applying $\operatorname{pr}_{Z*}^{Y \times Z}$ we conclude that

$$\operatorname{cor} \delta(\operatorname{cor} \gamma)(\alpha)) = \operatorname{pr}_{Z*}^{X \times Y \times Z}(\operatorname{pr}_{X}^{X \times Y \times Z*} \alpha \smile \operatorname{pr}_{X \times Y}^{*} \gamma \smile \operatorname{pr}_{Y \times Z}^{*} \delta)$$
(4.8)

Finally, we rewrite the right hand side as

$$\operatorname{pr}_{Z*}^{X\times Z}\operatorname{pr}_{X\times Z*}(\operatorname{pr}_{X\times Z}^*\operatorname{pr}_X^{X\times Z*}\alpha\smile\operatorname{pr}_{X\times Y}^*\gamma\smile\operatorname{pr}_{Y\times Z}^*\delta)$$

and apply Corollary 3.35 to $\operatorname{pr}_{X\times Z}$ (with the pushing morphism $(X\times Y\times Z,\Sigma)\to (X\times Z,P(\Phi_X,\Phi_Z))$ and pulling morphism $(X\times Y\times Z,\operatorname{pr}_X^{X\times Y\times Z-1}(\Phi_X))\to (X\times Z,\operatorname{pr}_X^{X\times Z-1}(\Phi_X)))$ to arrive at

$$\mathrm{pr}_{X\times Z*}(\mathrm{pr}_{X\times Z}^*\mathrm{pr}_X^{X\times Z*}\alpha\smile\mathrm{pr}_{X\times Y}^*\gamma\smile\mathrm{pr}_{Y\times Z}^*\delta)=\mathrm{pr}_X^{X\times Z*}\alpha\smile\mathrm{pr}_{X\times Z*}(\mathrm{pr}_{X\times Y}^*\gamma\smile\mathrm{pr}_{Y\times Z}^*\delta)$$

Applying $\operatorname{pr}_{Z*}^{X\times Z}$ on both sides shows that the right hand side of (4.8) is $\operatorname{cor}(\operatorname{pr}_{X\times Z*}(\operatorname{pr}_{X\times Y}^*\gamma \smile \operatorname{pr}_{Y\times Z}^*\delta)(\alpha)$, as desired.

Remark 4.9. There is a Grothendieck-Serre dual approach to such correspondences, where classes $\gamma \in H^j_{P(\Phi_Y,\Phi_Y)}(X \times Y, \Omega^i_{X \times Y}(\log \Delta_{X \times Y})(-\mathrm{pr}^*_Y \Delta_Y))$ define homomorphisms

$$H^q(X,\Omega_X^p(\log \Delta_X)(-\Delta_X)) \to H^{q+j-d_X}(Y,\Omega_Y^{p+i-d_X}(\log \Delta_Y)(-\Delta_Y)).$$

The construction is formally similar.

References

- [Bar18] Lawrence Jack Barrott. "Logarithmic Chow theory". In: *arXiv preprint arXiv:1810.03746* (2018).
- [BBG22] Christian Böhning, Hans-Christian Graf von Bothmer, and Michel van Garrel. "Prelog Chow rings and degenerations". In: *Rendiconti del Circolo Matematico di Palermo Series 2* (2022), pp. 1–34.
- [BPØ20] Federico Binda, Doosung Park, and Paul Arne Østvær. "Triangulated Categories of Logarithmic Motives over a Field". In: *arXiv:2004.12298* [math] (Apr. 2020). arXiv: 2004.12298 [math].
- [Con00] Brian Conrad. *Grothendieck duality and base change*. Vol. 1750. Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2000, pp. vi+296. ISBN: 3-540-41134-8. DOI: 10.1007/b75857. URL: https://doi.org/10.1007/b75857.
- [Con07] Brian Conrad. "Deligne's notes on Nagata compactifications". In: *J. Ramanu-jan Math. Soc.* 22.3 (2007), pp. 205–257. ISSN: 0970-1249.
- [CR11] Andre Chatzistamatiou and Kay Rülling. "Higher direct images of the structure sheaf in positive characteristic". In: *Algebra Number Theory* 5.6 (2011), pp. 693–775. ISSN: 1937-0652. DOI: 10.2140/ant.2011.5.693. URL: https://doi.org/10.2140/ant.2011.
- [CR15] Andre Chatzistamatiou and Kay Rülling. "Vanishing of the higher direct images of the structure sheaf". In: *Compos. Math.* 151.11 (2015), pp. 2131–2144.

 ISSN: 0010-437X. DOI: 10.1112/S0010437X15007435. URL: https://doi.org/10.1112/S0010437X
- [Del71] Pierre Deligne. "Théorie de Hodge. II". In: *Inst. Hautes Études Sci. Publ. Math.*40 (1971), pp. 5–57. ISSN: 0073-8301. URL: http://www.numdam.org/item?id=PMIHES_1971__40__5
- [DI87] Pierre Deligne and Luc Illusie. "Relèvements modulo p^2 et décomposition du complexe de de Rham". In: *Invent. Math.* 89.2 (1987), pp. 247–270. ISSN: 0020-9910. DOI: 10.1007/BF01389078. URL: https://doi.org/10.1007/BF01389078.
- [DM69] P. Deligne and D. Mumford. "The Irreducibility of the Space of Curves of given Genus". In: *Inst. Hautes Études Sci. Publ. Math.* 36 (1969), pp. 75–109. ISSN: 0073-8301. URL: http://www.numdam.org/item?id=PMIHES_1969__36__75_0.
- [EV92] Hélène Esnault and Eckart Viehweg. *Lectures on vanishing theorems*. Vol. 20. DMV Seminar. Birkhäuser Verlag, Basel, 1992, pp. vi+164. ISBN: 3-7643-2822-3. DOI: 10.1007/978-3-0348-8600-0. URL: https://doi.org/10.1007/978-3-0348-8600-0.

- [God22] Charles Godfrey. Higher direct images of snc ideal sheaves. 2022. DOI: 10.48550/ARXIV.2207.01142. URL: https://arxiv.org/abs/2207.01142.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977, pp. xvi+496. ISBN: 0-387-90244-9.
- [Hir64] Heisuke Hironaka. "Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II". In: *Ann. of Math. (2)* **79** (1964), 109–203; ibid. (2) 79 (1964), pp. 205–326. ISSN: 0003-486X. DOI: 10.2307/1970547. URL: https://doi.org/10.2307/1970547.
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*. Vol. 134. Cambridge Tracts in Mathematics. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original. Cambridge University Press, Cambridge, 1998, pp. viii+254. ISBN: 0-521-63277-3. DOI: 10.1017/CB09780511662560. URL: https://doi.org/10.1017/CB09780511662560.
- [Kol13] János Kollár. Singularities of the minimal model program. Vol. 200. Cambridge Tracts in Mathematics. With a collaboration of Sándor Kovács. Cambridge University Press, Cambridge, 2013, pp. x+370. ISBN: 978-1-107-03534-8. DOI: 10.1017/CB0978113954789 URL: https://doi.org/10.1017/CB09781139547895.
- [Kov20] Sándor J. Kovács. "Rational Singularities". In: *arXiv:1703.02269* [math] (July 2020). arXiv: 1703.02269 [math].
- [KX16] János Kollár and Chenyang Xu. "The dual complex of Calabi-Yau pairs". In: *Invent. Math.* 205.3 (2016), pp. 527–557. ISSN: 0020-9910. DOI: 10.1007/s00222-015-0640-6. URL: https://doi.org/10.1007/s00222-015-0640-6.
- [Lef53] Solomon Lefschetz. *Algebraic Geometry*. Princeton University Press, Princeton, NJ, 1953, pp. ix+233.
- [Mie09a] Yoichi Mieda. "Cycle classes, Lefschetz trace formula and integrality for p-adic cohomology". en. In: Algebraic Number Theory and Related Topics 2007, RIMS

 Kôkyûroku Bessatsu B12 (2009). URL: https://www.kurims.kyoto-u.ac.jp/~kenkyubu/bessatsu
- [Mie09b] Yoichi Mieda. "Integral Log Crystalline Cohomology and Algebraic Correspondences". en. In: *Proceedings of Kinosaki Algebraic Geometry Symposium* (2009). URL: https://www.ms.u-tokyo.ac.jp/~mieda/pdf/kinosaki2009.pdf.
- [MVW06] Carlo Mazza, Vladimir Voevodsky, and Charles Weibel. Lecture Notes on Motivic Cohomology. Vol. 2. Clay Mathematics Monographs. American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2006, pp. xiv+216. ISBN: 978-0-8218-3847-1 0-8218-3847-4.
- [Nag63] Masayoshi Nagata. "A generalization of the imbedding problem of an abstract variety in a complete variety". In: *J. Math. Kyoto Univ.* 3 (1963), pp. 89–102. ISSN: 0023-608X. DOI: 10.1215/kjm/1250524859. URL: https://doi.org/10.1215/kjm/1250524859.
- [Ogu18] Arthur Ogus. Lectures on logarithmic algebraic geometry. Vol. 178. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2018, pp. xviii+539. ISBN: 978-1-107-18773-3. DOI: 10.1017/9781316941614. URL: https://doi.org/10.1017/9781316941614.


- [R&D] Robin Hartshorne. *Residues and duality*. Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, No. 20. Springer-Verlag, Berlin-New York, 1966, pp. vii+423.
- [Stacks] The Stacks project authors. The Stacks project. 2021. URL: https://stacks.math.columbia.edu.
- [Voi14] Claire Voisin. Chow Rings, Decomposition of the Diagonal, and the Topology of Families. Princeton University Press, 2014. ISBN: 9780691160511. URL: http://www.jstor.org/stal (visited on 12/29/2022).

A Attempts to construct a fundamental class of a thrifty birational equivalence

As mentioned in Section 1 inspiration for this work was the following remarkable theorem of Chatzistamatiou-Rülling:

Theorem A.1 ([CR11, Thm. 3.2.8] (see also [CR15, Thm. 1.1], [Kov20, Thm. 1.6])). Let k be a perfect field and let S be a scheme. Suppose X and Y are two separated, finite type k-schemes which are

- (i) smooth over k and
- (ii) **properly birational** over S in the sense that there is a commutative diagram



with r and s proper birational morphisms.

Let $n = \dim X = \dim Y = \dim Z$. Then, there are isomorphisms of sheaves

$$R^{i}f_{*}\mathcal{O}_{X} \xrightarrow{\sim} R^{i}g_{*}\mathcal{O}_{Y} \text{ and } R^{i}f_{*}\omega_{X} \xrightarrow{\sim} R^{i}g_{*}\omega_{Y} \text{ for all } i,$$
 (A.3)

This result implies, for example, that if S is a variety over a perfect field k with a **rational resolution**, that is, a resolution of singularities $f: X \to S$ such that $Rf_* \mathcal{O}_X = \mathcal{O}_S$, then every other resolution $g: Y \to S$ satisfies $Rg_* \mathcal{O}_Y = \mathcal{O}_S$ and is hence also rational. In characteristic 0 this was a corollary of Hironaka's resolution of singularities [Hir64]; in positive characteristic it remained open until 2011.

The original proof in [CR11, Thm. 3.2.8] makes use of a cycle morphism cl: $CH^*(X) \to H^*(X, \Omega_X^*)$ from Chow cohomology to Hodge cohomology, which is ultimately applied to a cycle $Z \subset X \times Y$ obtained from a proper birational equivalence. That cycle morphism satisfies 2 essential properties: the first is that it is compatible with *correspondences*: here Chow correspondences are homomorphisms

$$CH^*(X) \to CH^*(Y)$$
 of the form $\alpha \longmapsto \operatorname{pr}_{Y*}(\operatorname{pr}_X^* \alpha \smile \gamma)$ for some $\gamma \in CH^*(X \times Y)$

where \smile is the cup product induced by intersecting cycles; Hodge correspondences are defined in a similar way. The second key property is a compatibility with the filtrations

$$CH^n(X \times Y) = F^0CH^n(X \times Y) \supseteq F^1CH^n(X \times Y) \supseteq \cdots \supseteq F^{\dim Y}CH^n(X \times Y) \supseteq 0$$

where $F^cCH^n(X\times Y)$ is the subgroup generated by cycles $Z\subseteq X\times Y$ such that $\operatorname{codim}(\operatorname{pr}_YZ\subseteq Y)\geq c$, and

$$H^n(X\times Y,\Omega^m_{X\times Y})=F^0H^n(X\times Y,\Omega^m_{X\times Y})\supseteq F^1CH^*(X\times Y)\supseteq\cdots\supseteq F^{\dim Y}H^n(X\times Y,\Omega^m_{X\times Y})\supseteq 0$$

where $F^cH^n(X\times Y,\Omega^m_{X\times Y})$ is the image of the map $H^n(X\times Y,\oplus_{j=c}^m\Omega^{m-j}_X\boxtimes\Omega^j_Y)\to H^n(X\times Y,\Omega^m_{X\times Y})$ coming from the Künneth decomposition.

It is natural to ask if a similar method can be applied to prove an analogue of Theorem A.1 for pairs, which might read something like Conjecture A.7 below. In order to state this analogue, we require a few additional definitions. For the remainder of this appendix we work over a fixed perfect field k.

Definition A.4 (slightly simplified version of [Kol13, Def. 1.5]). A **pair** (X, Δ_X) over k will mean

- a reduced, equidimensional and S_2 scheme X of finite type over k admitting a dualizing complex, together with
- a Q-Weil divisor $\Delta_X = \sum_i a_i D_i$ on X such that no irreducible component D_i of Δ_X is contained

Definition A.5. A *stratum* of a simple normal crossing pair $(X, \Delta_X = \sum_i D_i)$ is a connected (equivalently, irreducible) component of an intersection $D_J = \bigcap_{j \in J} D_j$.

Given any pair (X, Δ_X) , there is a largest open set $U \subseteq X$ such that $(U, \Delta_X|_U)$ is a simple normal crossing pair, and we will refer to the resulting simple normal crossing pair as $\operatorname{snc}(X, \Delta_X) :=$ $(U,\Delta_X|_U).$


Definition A.6 (compare with [Kol13, Def. 2.79-2.80], [KX16, §1, discussion before Def. 10]). Let $(S, \Delta_S = \sum_i D_i)$ be a pair, and assume Δ_S is reduced and effective. A separated, finite type birational

- morphism $f: X \to S$ is thrifty with respect to Δ_S if and only if

 (i) f is an isomorphism over the generic point of every stratum of $\operatorname{snc}(S, \Delta_S)$ and

 (ii) letting $\tilde{D}_i = f_*^{-1}D_i$ for i = 1, ..., N be the strict transforms of the divisors D_i , and setting $\Delta_X := \sum_i \tilde{D}_i$, the map f is an isomorphism at the generic point of every stratum of $\operatorname{snc}(X, \Delta_X)$.

Conjecture A.7. Let k be a perfect field, let S be a scheme and let (X, Δ_X) and (Y, Δ_Y) be simple normal crossing pairs over k. Suppose (X, Δ_X) and (Y, Δ_Y) are properly birational over S in the sense that there is a commutative diagram



where r, s are proper and birational morphisms, and assume $\Delta_Z = r_*^{-1} \Delta_X = s_*^{-1} \Delta_Y$. If r and s are thrifty, then there are quasi-isomorphisms

$$Rf_* \mathcal{O}_Y(-\Delta_Y) \simeq Rg_* \mathcal{O}_Y(-\Delta_Y) \text{ and } Rf_* \omega_Y(\Delta_Y) \simeq Rg_* \omega_Y(\Delta_Y).$$
 (A.9)

Following [CR11] closely, one might begin by replacing the ordinary sheaves of differentials Ω_X appearing in Hodge cohomology with sheaves of differentials with log poles $\Omega_X(\log \Delta_X)$ and attempt to implement a similar strategy, i.e. starting a cycle $Z \subset X \times Y$ representing a thrifty proper birational equivalince, producing a correspondence in logarithmic Hodge cohomology and analyzing its properties.

Ultimately even the correspondences of Section 4 seem to be insufficient to deal with thrifty proper birational equivalences, as we illustrate in Appendix A.1 below. The problem we encounter is elementary: looking at the recipe for the Hodge class cl(Z) of a subvariety $Z \subseteq X$, where Z and X are smooth an projective (outlined in [Har77, Ex. III.7.4]), we see that cl(Z) ultimately comes from the trace linear functional tr : $H^{\dim Z}(Z, \omega_Z) \to k$, or Serre-dually the element $1 \in H^0(Z, \mathcal{O}_Z)$. Due to the introduction of log poles and zeroes in Section 4, trying to follow that recipe we pass through cohomology groups of the form $H^{\dim Z}(Z,\omega_Z(D))$, or dually $H^0(Z,\mathcal{O}_Z(-D))$ where D is an (often non-0 in cases of interest) effective Cartier divisor on Z, and so there simply is no "1" to be had.

Beyond the difficulties described in the previous paragraph, when attempting to formulate a logarithmic variant of Chatzistamatiou-Rülling's cycle morphism argument one is hampered by the fact that we are still in the early days of logarithmic Chow theory. It is not clear to the author which logarithmic variant of Fulton's CH^* , if any, could be used to construct a logarithmic cycle morphism with all of the desired properties. Further investigation of this question could be an interesting topic

Despite the aforementioned challenges, it is possible to prove a result almost identical to Conjecture A.7 by entirely different methods [God22].8

A.1 Obstructions to obtaining log Hodge correspondences from thrifty birational equivalences

Let (X, Δ_X) , (Y, Δ_Y) be simple normal crossing pairs, and assume in addition that X, Y are connected and proper. Let $Z \subseteq X \times Y$ be a smooth closed subvariety with codimension c. In this situation the fundamental class of $\mathrm{cl}(Z) \in H^c(X \times Y, \Omega^c_{X \times Y})$ (no log poles yet) can be described using only Serre duality, as follows (we refer to [Har77, Ex. III.7.4]). the composition

$$H^{\dim Z}(X \times Y, \Omega_{X \times Y}^{\dim Z}) \to H^{\dim Z}(Z, \Omega_Z^{\dim Z}) \xrightarrow{\operatorname{tr}} k$$
 (A.10)

(where tr is the trace map of Serre duality) is an element of

$$H^{\dim Z}(X \times Y, \Omega_{X \times Y}^{\dim Z})^{\vee} \simeq H^{c}(X \times Y, \Omega_{X \times Y}^{c})$$
(A.11)

which we may define to be cl(Z). In light of Theorem 4.1 we might hope to modify eqs. (A.10) and (A.11) to obtain a class in $H^c(X\times Y,\Omega^c_{X\times Y}(\log\Delta_{X\times Y})(-\operatorname{pr}_X^*\Delta_X))$. Let us focus on the case where $\operatorname{pr}_X|_Z:Z\to X,\operatorname{pr}_Y|_Z:Z\to Y$ are both thrifty and birational, so in particular $c=\dim X=$

- $\dim Y = : d$ and

• $(\operatorname{pr}_X|_Z)_*^{-1}\Delta_X = (\operatorname{pr}_Y|_Z)_*^{-1}\Delta_Y =: \Delta_Z$ To keep the notation under control, set $\pi_X := \operatorname{pr}_X|_Z$ and $\pi_Y := \operatorname{pr}_Y|_Z$. In this situation letting $\iota: Z \to X \times Y$ be the inclusion there is a natural map

$$d\iota^\vee: \Omega^d_{X\times Y}(\log \Delta_{X\times Y}) \to \iota_*\Omega^d_Z(\log \Delta_{X\times Y}|_Z) \text{ and twisting by } -\mathrm{pr}_Y^*\Delta_Y \text{ gives a map}$$

$$\Omega^d_{X\times Y}(\log \Delta_{X\times Y})(-\mathrm{pr}_Y^*\Delta_Y) \to \iota_*\Omega^d_Z(\log \Delta_{X\times Y}|_Z)(-\mathrm{pr}_Y^*\Delta_Y|_Z) = \iota_*\Omega^d_Z(\log \Delta_{X\times Y}|_Z)(-\pi_Y^*\Delta_Y)$$

To identify $\Omega_Z^d(\log \Delta_{X\times Y}|_Z)(-\operatorname{pr}_X^*\Delta_X|_Z)$, write

$$(\pi_X)^* \Delta_X = (\pi_X)_*^{-1} \Delta_X + E_X = \Delta_Z + E_X$$
 and $(\pi_Y)^* \Delta_Y = (\pi_Y)_*^{-1} \Delta_Y + E_Y = \Delta_Z + E_Y$

so that $\Delta_{X\times Y}|_Z=(\pi_X)^*\Delta_X+(\pi_Y)^*\Delta_Y=2\Delta_Z+E_X+E_Y$. While the hypotheses guarantee Δ_Z is reduced it may be that E_X , E_Y are non-reduced — however something can be said about their multiplicities. If $E_X = \sum_i a_X^i E_X^i$, $E_Y = \sum_i a_Y^i E_Y^i$ where the E_X^i , E_Y^i are irreducible, then by a generalization of [Har77, Prop. 3.6] (see also [Kol13, §2.10]),

$$a_X^i = \mathrm{mlt}(\pi_X(E_X^i) \subseteq \Delta_X)$$

and since Δ_X is a reduced effective simple normal crossing divisor, if in addition we write $\Delta_X = \sum_i D_X^i$, then $\mathrm{mlt}(\pi_X(E_X^i) \subseteq \Delta_X) = |\{i \mid \pi_X(E_X^i) \subseteq D_X^i\}|$. The thriftiness hypothesis that $\pi_X(E_X^i)$ is not

⁸The reason the result is only "almost identical" is that in [God22] we require ostensibly stronger hypotheses on the base scheme S (namely that it is excellent and noetherian), but it is possible that even in the situation of Theorem A.1 and Conjecture A.7 one can reduce to this case, for example using noetherian approximation.

⁹It may then be non-trivial to verify this agrees with other definitions, especially if we worry about signs, but we will not need that level of detail for what follows.

a stratum then implies $a_X^i = \mathrm{mlt}(\pi_X(E_X^i) \subseteq \Delta_X) < \mathrm{codim}(\pi_X(E_X^i) \subset X)$. Since differentials with log poles are insensitive to multiplicities, we have

$$\Omega_Z^d(\log \Delta_{X \times Y}|_Z) = \omega_Z(\Delta_Z + E_X^{\text{red}} + E_Y^{\text{red}})$$

where $-^{\text{red}}$ denotes the associated reduced effective divisor. Then

$$\begin{split} \Omega^d_Z(\log \Delta_{X\times Y}|_Z)(-\pi_Y^*\Delta_Y) &= \omega_Z(\Delta_Z + E_X^{\mathrm{red}} + E_Y^{\mathrm{red}} - \Delta_Z - E_Y) \\ \omega_Z(E_X^{\mathrm{red}} + (E_Y^{\mathrm{red}} - E_Y)) &= \omega_Z(\sum_i E_X^i + \sum_i (1 - a_Y^i) E_Y^i) \end{split}$$

The upshot is that we have an induced map

$$H^{d}(X \times Y, \Omega^{d}_{X \times Y}(\log \Delta_{X \times Y})(-\operatorname{pr}_{Y}^{*} \Delta_{Y})) \to H^{d}(Z, \omega_{Z}(E_{X}^{\operatorname{red}} + (E_{Y}^{\operatorname{red}} - E_{Y})))$$
(A.12)

Here the left hand side is Serre dual to $H^d(X \times Y, \Omega^d_{X \times Y}(\log \Delta_{X \times Y})(-\operatorname{pr}_X^* \Delta_X))$, so the k-linear dual of (A.12) is a morphism

$$H^d(Z,\omega_Z(E_X^{\mathrm{red}} + (E_Y^{\mathrm{red}} - E_Y)))^\vee \to H^d(X \times Y,\Omega^d_{X \times Y}(\log \Delta_{X \times Y})(-\mathrm{pr}_X^*\Delta_X))$$

Unfortunately $^{10}H^d(Z, \omega_Z(E_X^{\mathrm{red}} + (E_Y^{\mathrm{red}} - E_Y)))$ is often 0. If E_X and E_Y are both reduced (an explicit example where this holds will be given below), then $H^d(Z, \omega_Z(E_X^{\mathrm{red}} + (E_Y^{\mathrm{red}} - E_Y))) = H^d(Z, \omega_Z(E_X))$. If in addition $E_X \neq 0$, we obtain $H^d(Z, \omega_Z(E_X)) = 0$ by an extremely weak (but characteristic independent) sort of Kodaira vanishing:

Lemma A.13. Let Z be a proper variety over a field k with dimension d, and assume Z is normal and Cohen-Macaulay. If $D \subset Z$ is a non-0 effective Cartier divisor on Z then $H^d(Z, \omega_Z(D)) = 0$.

Proof. By Serre duality $H^d(Z, \omega_Z(D)) = H^0(Z, \mathcal{O}_Z(-D))$, which vanishes by the classic fact that "a nontrivial line bundle and its inverse can't both have non-0 global sections." Since I am not aware of a specific reference, here is a proof:

Suppose towards contraditction that there is a non-0 global section $\sigma \in H^0(Z, \mathbb{O}_Z(-D))$ — then the composition

$$\mathcal{O}_Z \xrightarrow{\sigma} \mathcal{O}_Z(-D) \xrightarrow{\tau} \mathcal{O}_Z$$

is non-0. By [Stacks, Tag 0358] $H^0(Z, \mathcal{O}_Z)$ is a (normal) domain, and since it's also a finite dimensional k-vector space it must be an extension field of k. But then $\tau \in H^0(Z, \mathcal{O}_Z)$ is invertible hence surjective, so $\mathcal{O}_Z(-D) \hookrightarrow \mathcal{O}_Z$ is surjective, which is a contradiction since by hypothesis the cokernel $\mathcal{O}_D \neq 0$.

Example A.14. Let $X=\mathbb{P}^2$ and let $\Delta_X\subset X$ be a line. Let $p\in L$ be a k-point, let $Y=\operatorname{Bl}_pX$ and let $\Delta_Y=\tilde{L}=\mathbb{P}$ the strict transform of L. Finally let $f:Y\to X$ be the blowup map and let $Z=(f\times\operatorname{id})(Y)\subset X\times Y$. In this case (with all notation as above) $\pi_X\circ(f\times\operatorname{id})=f$ and $\pi_Y\circ(f\times\operatorname{id})=\operatorname{id}_Y$, so under the isomorphism $f\times\operatorname{id}:Y\simeq Z$, E_X is the exceptional divisor of f (with multiplicity 1). On the other hand $E_Y=0$. In particular E_X and E_Y are reduced and $E_X\neq 0$ so from the above discussion $H^2(Z,\omega_Z(E_X))=0$.

¹⁰at least for the purposes of constructing log Hodge cohomology classes of subvarieties ...