

# ON THE LIFTING PROBLEM OF REPRESENTATIONS OF A METACYCLIC GROUP.

ARISTIDES KONTOGEORGIS AND ALEXIOS TEREZAKIS

ABSTRACT. We give a necessary and sufficient condition for a modular representation of a group  $G = C_{p^h} \rtimes C_m$  in a field of characteristic zero to be lifted to a representation over local principal ideal domain of characteristic zero containing the  $p^h$  roots of unity.

## 1. INTRODUCTION

The lifting problem for a representation

$$\rho : G \rightarrow \mathrm{GL}_n(k),$$

where  $k$  is a field of characteristic  $p > 0$ , is about finding a local ring  $R$  of characteristic 0, with maximal ideal  $\mathfrak{m}_R$  such that  $R/\mathfrak{m}_R = k$ , so that the following diagram is commutative:

$$\begin{array}{ccc} & & \mathrm{GL}_n(R) \\ & \nearrow & \downarrow \\ G & \longrightarrow & \mathrm{GL}_n(k) \end{array}$$

Equivalently one asks if there is a free  $R$ -module  $V$ , which is also an  $R[G]$ -module such that  $V \otimes_R R/\mathfrak{m}_R$  is the  $k[G]$ -module corresponding to our initial representation. We know that projective  $k[G]$ -modules lift in characteristic zero, [16, chap. 15], but for a general  $k[G]$ -module such a lifting is not always possible, for example, see [10, prop. 15]. This article aims to study the lifting problem for the group  $G = C_q \rtimes C_m$ , where  $C_q$  is a cyclic group of order  $p^h$  and  $C_m$  is a cyclic group of order  $m$ ,  $(p, m) = 1$  and give necessary and sufficient condition in order to lift. We assume that the local ring  $R$  contains the  $q$ -roots of unity and  $k$  is algebraically closed, and we might need to consider a ramified extension of  $R$ , in order to ensure that certain  $q$ -roots of unit are distant in the  $\mathfrak{m}_R$ -topology, see remark 35. An example of such a ring  $R$  is the ring of Witt vectors  $W(k)[\zeta_q]$  with the  $q$ -roots of unity adjoined to it.

We notice that a decomposable  $R[G]$ -module  $V$  gives rise to a decomposable  $R$ -module modulo  $\mathfrak{m}_R$  and also an indecomposable  $R[G]$ -module can break in the reduction modulo  $\mathfrak{m}_R$  into a direct sum of indecomposable  $k[G]$ -summands. We also give a classification of  $k[C_q \rtimes C_m]$ -modules in terms of Jordan decomposition and give the relation with the more usual uniserial description in terms of their socle [1].

---

*Date:* January 4, 2023.

*Key words and phrases.* Lifting of representations, modular representation theory, integral representation theory, Generalized Oort conjecture, metacyclic groups.

Our interest to this problem comes from the problem of lifting local actions. The local lifting problem considers the following question: Does there exist an extension  $\Lambda/W(k)$ , and a representation

$$\tilde{\rho} : G \hookrightarrow \text{Aut}(\Lambda[[T]]),$$

such that if  $t$  is the reduction of  $T$ , then the action of  $G$  on  $\Lambda[[T]]$  reduces to the action of  $G$  on  $k[[t]]$ ?

If the answer to the above question is positive, then we say that the  $G$ -action lifts to characteristic zero. A group  $G$  for which every local  $G$ -action on  $k[[t]]$  lifts to characteristic zero is called a *local Oort group for  $k$* . Notice that cyclic groups are always local Oort groups. This result was known as the ‘‘Oort conjecture’’, which was recently proved by F. Pop [15] using the work of A. Obus and S. Wewers [14].

There are a lot of obstructions that prevent a local action to lift in characteristic zero. Probably the most important of these obstructions in the KGB-obstruction [4]. It is believed that this is the only obstruction for the local lifting problem, see [11], [12]. In [10, Thm. 3] the authors have given a criterion for the local lifting which involves the lifting of a linear representation of the same group. The case  $G = C_q \rtimes C_m$  and especially the case of dihedral groups  $D_q = C_q \rtimes C_2$ , is a problem of current interest in the theory of local liftings, see [12], [6], [18]. For more details on the local lifting problem we refer to [3], [4], [5], [11].

Keep also in mind that the  $C_q \rtimes C_m$  groups were important to the study of group actions in holomorphic differentials of curves defined over fields of positive characteristic  $p$ , where the group involved has cyclic  $p$ -Sylow subgroup, see [2].

Let us now describe the method of proof. For understanding the splitting of indecomposable  $R[G]$ -modules modulo  $\mathfrak{m}_R$ , we develop a version of Jordan normal form in lemma 16 for endomorphisms  $T : V \rightarrow V$  of order  $p^h$ , where  $V$  is a free module of rank  $d$ . We give a way to select this basis, by selecting an initial suitable element  $E \in V$ , see lemma 15. The normal form (as given in eq. (9)) of the element  $T$  of order  $q$ , determines the decomposition of the reduction. We show that for every indecomposable summand  $V_i$  of  $V$ , we can select  $E$  as an eigenvalue of the generator  $\sigma$  of  $C_m$  and then by forcing the relation  $\Gamma T = T^\alpha \Gamma$  to hold, we see how the action of  $\sigma$  can be extended recursively to an action of  $\sigma$  on  $V_i$ , this is done in lemma 24. Proving that this gives indeed a well defined action is a technical computation and is done in lemmata 26, 27, 28, 32, 33.

The important thing here is that the definition of the action of  $\sigma$  on  $E$  is the ‘‘initial condition’’ of a dynamical system that determines the action of  $C_m$  on the indecomposable summand  $V_i$ . The  $R[C_q \rtimes C_m]$  indecomposable module  $V_i$  can break into a direct sum  $V_\alpha(\epsilon_\nu, \kappa_\nu)$ -modules  $1 \leq \nu \leq s$  (for a precise definition of them see definition 9, notice that  $\kappa_i$  denotes the dimension). The action of  $\sigma$  on each  $V_\alpha(\epsilon_\nu, \kappa_\nu)$  can be uniquely determined by the action of  $\sigma$  on an initial basis element as shown in section 3, again by a ‘‘dynamical system’’ approach, where we need  $s$  initial conditions, one for each  $V_\alpha(\epsilon_\nu, \kappa_\nu)$ . The lifting condition essentially means that the indecomposable summands  $V_\alpha(\epsilon, \kappa)$  of the special fibre, should be able to be rearranged in a suitable way, so that they can be obtained as reductions of indecomposable  $R[C_q \rtimes C_q]$ -modules. The precise expression of our lifting criterion is given in the following proposition:

**Proposition 1.** *Consider a  $k[G]$ -module  $M$  which is decomposed as a direct sum*

$$M = V_\alpha(\epsilon_1, \kappa_1) \oplus \cdots \oplus V_\alpha(\epsilon_s, \kappa_s).$$


The module lifts to an  $R[G]$ -module if and only if the set  $\{1, \dots, s\}$  can be written as a disjoint union of sets  $I_\nu$ ,  $1 \leq \nu \leq t$  so that

- a  $\sum_{\mu \in I_\nu} \kappa_\mu \leq q$ , for all  $1 \leq \nu \leq t$ .
- b  $\sum_{\mu \in I_\nu} \kappa_\mu \equiv a \pmod{m}$  for all  $1 \leq \nu \leq t$ , where  $a \in \{0, 1\}$ .
- c For each  $\nu$ ,  $1 \leq \nu \leq t$  there is an enumeration  $\sigma : \{1, \dots, \#I_\nu\} \rightarrow I_\nu \subset \{1, \dots, s\}$ , such that

$$\epsilon_{\sigma(2)} = \epsilon_{\sigma(1)} \alpha^{\kappa_{\sigma(1)}}, \epsilon_{\sigma(3)} = \epsilon_{\sigma(2)} \alpha^{\kappa_{\sigma(2)}}, \dots, \epsilon_{\sigma(s)} = \epsilon_{\sigma(s-1)} \alpha^{\kappa_{\sigma(s-1)}}$$

In the above proposition, each set  $I_\nu$  corresponds to a collection of modules  $V_\alpha(\epsilon_\mu, \kappa_\mu)$ ,  $\mu \in I_\nu$  which come as the reduction of an indecomposable  $R[C_q \rtimes C_m]$ -module  $V_\nu$  of  $V$ .

**Acknowledgements** A. Terezakis is a recipient of financial support in the context of a doctoral thesis (grant number MIS-5113934). The implementation of the doctoral thesis was co-financed by Greece and the European Union (European Social Fund-ESF) through the Operational Programme—Human Resources Development, Education and Lifelong Learning—in the context of the Act—Enhancing Human Resources Research Potential by undertaking a Doctoral Research—Sub-action 2: IKY Scholarship Programme for Ph.D. candidates in the Greek Universities.



## 2. NOTATION

Let  $\tau$  be a generator of the cyclic group  $C_q$  and  $\sigma$  be a generator of the cyclic group  $C_m$ . The group  $G$  is given in terms of generators and relations as follows:

$$G = \langle \sigma, \tau \mid \tau^q = 1, \sigma^m = 1, \sigma\tau\sigma^{-1} = \tau^\alpha \text{ for some } \alpha \in \mathbb{N}, 1 \leq \alpha \leq p^h - 1, (\alpha, p) = 1 \rangle.$$

The integer  $\alpha$  satisfies the following congruence:

$$(1) \quad \alpha^m \equiv 1 \pmod{q}$$

as one sees by computing  $\tau = \sigma^m \tau \sigma^{-m} = \tau^{\alpha^m}$ . Also the integer  $\alpha$  can be seen as an element in the finite field  $\mathbb{F}_p$ , and it is a  $(p-1)$ -th root of unity, not necessarily primitive. In particular the following holds:

**Lemma 2.** *Let  $\zeta_m \in k$  be a fixed primitive  $m$ -th root of unity. There is a natural number  $a_0$ ,  $0 \leq a_0 < m-1$  such that  $\alpha = \zeta_m^{a_0}$ .*

*Proof.* The integer  $\alpha$  if we see it as an element in  $k$  is an element in the finite field  $\mathbb{F}_p \subset k$ , therefore  $\alpha^{p-1} = 1$  as an element in  $\mathbb{F}_p$ . Let  $\text{ord}_p(\alpha)$  be the order of  $\alpha$  in  $\mathbb{F}_p^*$ . By eq. (1) we have that  $\text{ord}_p(\alpha) \mid p-1$  and  $\text{ord}_p(\alpha) \mid m$ , that is  $\text{ord}_p(\alpha) \mid (p-1, m)$ .

The primitive  $m$ -th root of unity  $\zeta_m$  generates a finite field  $\mathbb{F}_p(\zeta_m) = \mathbb{F}_{p^\nu}$  for some integer  $\nu$ , which has cyclic multiplicative group  $\mathbb{F}_{p^\nu} \setminus \{0\}$  containing both the cyclic groups  $\langle \zeta_m \rangle$  and  $\langle \alpha \rangle$ . Since for every divisor  $\delta$  of the order of a cyclic group  $C$  there is a unique subgroup  $C' < C$  of order  $\delta$  we have that  $\alpha \in \langle \zeta_m \rangle$ , and the result follows.  $\square$

**Definition 3.** For each  $p^i \mid q$  we define  $\text{ord}_{p^i} \alpha$  to be the smallest natural number  $o$  such that  $\alpha^o \equiv 1 \pmod{p^i}$ .

It is clear that for  $\nu \in \mathbb{N}$

$$\alpha^\nu \equiv 1 \pmod{p^i} \Rightarrow \alpha^\nu \equiv 1 \pmod{p^j} \text{ for all } j \leq i.$$

Therefore

$$\text{ord}_{p^j} \alpha \mid \text{ord}_{p^i} \alpha \text{ for } j \leq i.$$

On the other hand  $\alpha \in \mathbb{N}$  and  $\alpha^{p-1} \equiv 1 \pmod{p}$  so  $\text{ord}_p \alpha \mid p-1$ . Also since  $\sigma^t \tau \sigma^{-t} = \tau^{\alpha^t}$  we have that  $\alpha^m \equiv 1 \pmod{p^h}$ , therefore  $\text{ord}_p \alpha \mid \text{ord}_{p^i} \alpha \mid \text{ord}_{p^h} \alpha \mid m$ , for  $1 \leq i \leq h$ .

**Lemma 4.** *The center  $\text{Cent}_G(\tau) = \langle \tau, \sigma^{\text{ord}_{p^h} \alpha} \rangle$ . Moreover*

$$\frac{|\text{Cent}_G(\tau)|}{p^h} = \frac{m}{\text{ord}_{p^h}(\alpha)} =: m'$$

*Proof.* The result follows by observing  $(\tau^\nu \sigma^t) \tau (\tau^\nu \sigma^t)^{-1} = \tau^{\alpha^t}$ , for all  $1 \leq \nu \leq q$ ,  $1 \leq t \leq m$ .  $\square$

**Remark 5.** If  $\text{ord}_p \alpha = m$  then  $\text{ord}_{p^i} \alpha = m$  for all  $1 \leq i \leq h$ .

**Lemma 6.** *If the group  $G = C_q \rtimes C_m$  is a subgroup of  $\text{Aut}(k[[t]])$ , then all orders  $\text{ord}_{p^i} \alpha = m/m'$ , for all  $1 \leq i \leq h$ .*

*Proof.* We will use the notation of the book of J.P.Serre on local fields [17]. By [13, Th.1.1b] we have that the first gap  $i_0$  in the lower ramification filtration of the cyclic group  $C_q$  satisfies  $(m, i_0) = m'$ .

The ramification relation [17, prop. 9 p. 69]

$$\alpha \theta_{i_0}(\tau) = \theta_{i_0}(\tau^\alpha) = \theta_{i_0}(\sigma \tau \sigma^{-1}) = \theta_0(\sigma)^{i_0} \theta_{i_0}(\tau),$$

implies that  $\theta_0(\sigma)^{i_0} = \alpha \in \mathbb{N}$ . From  $(m, i_0) = m'$  and the fact that  $\text{ord} \theta_0(\sigma) = m$  we obtain

$$\frac{m}{m'} = \text{ord} \theta_0(\sigma)^{i_0} = \text{ord}_p(\alpha).$$

Thus

$$\frac{m}{m'} = \text{ord}_p \alpha \mid \text{ord}_{p^i} \alpha \mid \text{ord}_{p^h} \alpha = \frac{m}{m'}.$$

Hence all orders  $\text{ord}_{p^i} \alpha = m/m'$ .  $\square$

**Remark 7.** If the KGB-obstruction vanishes and  $\alpha \neq 1$ , then by [11][prop. 5.9]  $i_0 \equiv -1 \pmod{m}$  and  $\text{ord}_{p^i} \alpha = m$  for all  $1 \leq i \leq h$ .

### 3. INDECOMPOSABLE $C_q \rtimes C_m$ MODULES, MODULAR REPRESENTATION THEORY

In this section we will describe the indecomposable  $C_q \rtimes C_m$ -modules. We will give two methods in studying them. The first one is needed since it is in accordance to the method we will give in order to describe indecomposable  $R[C_q \rtimes C_m]$ -modules. The second one, using the structure of the socle, is the standard method of describing  $k[C_q \rtimes C_m]$ -modules in modular representation theory.

**3.1. Linear algebra method.** The indecomposable modules of the  $C_q$  are determined by the Jordan normal forms of the generator  $\tau$  of the cyclic group  $C_q$ . So for each  $1 \leq \kappa \leq p^h$  there is exactly one  $C_q$  indecomposable module denoted by  $J_\kappa$ . Therefore we have the following decomposition of an indecomposable  $C_q \rtimes C_m$ -module  $M$  considered as a  $C_q$ -module.

$$(2) \quad M = J_{\kappa_1} \oplus \cdots \oplus J_{\kappa_r}.$$

**Lemma 8.** *In the indecomposable module  $J_\kappa$  for every element  $E$  such that*

$$(\tau - \text{Id}_{\kappa_i})^{\kappa_i-1} E \neq 0$$

*the elements  $B = \{E, (\tau - \text{Id}_\kappa)E, \dots, (\tau - \text{Id}_\kappa)^{\kappa-1}E\}$  form a basis of  $J_\kappa$  such that the matrix of  $\tau$  with respect to this basis is given by*

$$(3) \quad \tau = \text{Id}_\kappa + \begin{pmatrix} 0 & \cdots & \cdots & \cdots & 0 \\ 1 & \ddots & & & \vdots \\ 0 & \ddots & \ddots & & \vdots \\ \vdots & \ddots & 1 & 0 & \vdots \\ 0 & \cdots & 0 & 1 & 0 \end{pmatrix}.$$

*Proof.* Since the set  $B$  has  $k$ -elements it is enough to prove that it consists of linear independent elements. Indeed, consider a linear relation

$$\lambda_0 E + \lambda_1 (\tau - \text{Id}_\kappa)E + \cdots + \lambda_{\kappa-1} (\tau - \text{Id}_\kappa)^{\kappa-1} E = 0.$$

By applying  $(\tau - \text{Id}_\kappa)^{\kappa-1}$  we obtain  $\lambda_0 (\tau - \text{Id}_\kappa)^{\kappa-1} = 0$ , which gives us  $\lambda_0 = 0$ . We then apply  $(\tau - \text{Id}_\kappa)^{\kappa-2}$  to the linear relation and by the same argument we obtain  $\lambda_1 = 0$  and we continue this way proving that  $\lambda_0 = \cdots = \lambda_{\kappa-1} = 0$ . The matrix form of  $\tau$  in this basis is immediate.  $\square$

We will now prove that  $\sigma$  acts on each  $J_\kappa$  of eq. (2) proving that  $r = 1$ . Since the field  $k$  is algebraically closed and  $(m, p) = 1$  we know that there is a basis of  $M$  consisting of eigenvectors of  $\sigma$ . There is an eigenvector  $E$  of  $\sigma$ , which is not in the kernel of  $(\tau - \text{Id}_\kappa)^{\kappa-1}$ . Then the elements of the set  $B = \{E, (\tau - \text{Id}_\kappa)E, \dots, (\tau - \text{Id}_\kappa)^{\kappa-1}E\}$  are linearly independent and form a direct  $C_q$  summand of  $M$  isomorphic to  $J_{\kappa_1}$ .

We will now show that this module is an  $k[C_q \rtimes C_m]$ -module. For this, we have to show that the generator  $\sigma$  of  $C_m$  acts on the basis  $B$ . Observe that for every  $0 \leq i \leq \kappa_1 - 1 < p^h$

$$\sigma(\tau - 1)^{i-1} = (\tau^\alpha - 1)^{i-1} \sigma.$$

Set  $e = E_1$  and  $\kappa = \kappa_1$ . This means that the action of  $\sigma$  on  $e$  determines the action of  $\sigma$  on all other basis elements  $e_\nu := (\tau - 1)^{\nu-1} e$ ,  $1 \leq \nu \leq \kappa_1$ .

Let us compute:

$$\sigma e_{i+1} = \sigma(\tau - 1)^i e = (\tau^\alpha - 1)^i \zeta_m^\lambda e$$

On the basis  $\{e_1, \dots, e_{\kappa_1}\}$  the matrix  $\tau$  is given by eq. (3) hence using the binomial formula we compute

$$(4) \quad \tau^\alpha = \begin{pmatrix} 1 & 0 & \cdots & \cdots & \cdots & 0 \\ \binom{\alpha}{1} & 1 & \ddots & & & \vdots \\ \binom{\alpha}{2} & \binom{\alpha}{1} & \ddots & \ddots & & \vdots \\ \binom{\alpha}{3} & \binom{\alpha}{2} & \ddots & 1 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \binom{\alpha}{1} & 1 & 0 \\ \binom{\alpha}{k} & \binom{\alpha}{k-1} & \cdots & \binom{\alpha}{2} & \binom{\alpha}{1} & 1 \end{pmatrix}.$$

Thus  $\tau^\alpha - 1$  is a nilpotent matrix  $A = (a_{ij})$  of the form:

$$a_{ij} = \begin{cases} \binom{\alpha}{\mu} & \text{if } j = i - \mu \text{ for some } \mu, 1 \leq \mu \leq \kappa \\ 0 & \text{if } j \geq i \end{cases}$$

The  $\ell$ -th power  $A^\ell = (a_{ij}^{(\ell)})$  of  $A$  is then computed by (keep in mind that  $a_{ij} = 0$  for  $i \leq j$ )

$$a_{ij}^{(\ell)} = \sum_{i < \nu_1 < \cdots < \nu_{\ell-1} < j} a_{i, \nu_1} a_{\nu_1, \nu_2} a_{\nu_2, \nu_3} \cdots a_{\nu_{\ell-1}, j}$$

This means that  $i - j > \ell$  in order to have  $a_{ij} \neq 0$ . Moreover for  $i = j + \ell$  (which is the the first non zero diagonal below the main diagonal) we have

$$a_{i, i+\ell} = a_{i, i+1} a_{i+1, i+2} \cdots a_{i+\ell-1, i+\ell} = \binom{\alpha}{1}^\ell = \alpha^\ell.$$

Therefore, the matrix of  $A^\ell$  is of the following form:

$$(5) \quad \begin{pmatrix} \overbrace{0 \cdots \cdots}^{k-\ell} 0 & \overbrace{0 \cdots 0}^{\ell} \\ \vdots & \vdots \\ 0 \cdots \cdots 0 & 0 \cdots 0 \\ \alpha^\ell \ddots & 0 \vdots \\ * \alpha^\ell \ddots & \vdots \vdots \\ \vdots \ddots \ddots & 0 \vdots \\ * \cdots * \alpha^\ell & 0 \cdots 0 \end{pmatrix}$$

**Definition 9.** We will denote by  $V_\alpha(\lambda, \kappa)$  the indecomposable  $\kappa$ -dimensional  $G$ -module given by the basis elements  $\{(\tau - 1)^\nu e, \nu = 0, \dots, \kappa - 1\}$ , where  $\sigma e = \zeta_m^\lambda e$ .

This definition is close to the notation used in [9].

**Lemma 10.** *The action of  $\sigma$  on the basis element  $e_i$  of  $V_\alpha(\lambda, \kappa)$  is given by:*

$$(6) \quad \sigma e_i = \alpha^{i-1} \zeta_m^\lambda e_i + \sum_{\nu=i+1}^{\kappa} a_\nu e_\nu,$$

for some coefficients  $a_i \in k$ . In particular the matrix of  $\sigma$  with respect to the basis  $e_1, \dots, e_\kappa$  is lower triangular.

*Proof.* Recall that  $e_i = (\tau - 1)^{i-1}e_1$ . Therefore

$$\sigma e_i = \sigma(\tau - 1)^{i-1}e_1 = (\tau^\alpha - 1)^{i-1}\sigma e_1 = \zeta_m^\lambda (\tau^\alpha - 1)^{i-1}e_1.$$

The result follows by eq. (5)  $\square$

We have constructed a set of indecomposable modules  $V_\alpha(\lambda, \kappa)$ . Apparently  $V_\alpha(\lambda, \kappa)$  can not be isomorphic to  $V_\alpha(\lambda', \kappa')$  if  $\kappa \neq \kappa'$ , since they have different dimensions.

Assume now that  $\kappa = \kappa'$ . Can the modules  $V_\alpha(\lambda, \kappa)$  and  $V_\alpha(\lambda', \kappa)$  be isomorphic for  $\lambda \neq \lambda'$ ?

The eigenvalues of the prime to  $p$  generator  $\sigma$  on  $V_\alpha(\lambda, \kappa)$  are

$$\zeta_m^\lambda, \alpha\zeta_m^\lambda, \dots, \alpha^{\kappa-1}\zeta_m^\lambda.$$

Similarly the eigenvalues for  $\sigma$  when acting on  $V_\alpha(\lambda', \kappa)$  are

$$\zeta_m^{\lambda'}, \alpha\zeta_m^{\lambda'}, \dots, \alpha^{\kappa-1}\zeta_m^{\lambda'}.$$

If the two sets of eigenvalues are different then the modules can not be isomorphic. But even if  $\lambda \neq \lambda' \pmod{n}$  the two sets of eigenvalues can still be equal. Even in this case the modules can not be isomorphic.

**Lemma 11.** *The modules  $V_\alpha(\lambda_1, \kappa)$  and  $V_\alpha(\lambda_2, \kappa)$  are isomorphic if and only if  $\lambda_1 \equiv \lambda_2 \pmod{m}$ .*

*Proof.* Indeed, the module  $V_\alpha(\lambda_1, \kappa)$  has an eigenvector for the action of  $\sigma$  which generates the  $V_\alpha(\lambda_1, \kappa)$  by powers of  $(\tau - 1)$ , i.e. the vectors

$$(7) \quad e, (\tau - 1)e, (\tau - 1)^2e, \dots, (\tau - 1)^{\kappa-1}e$$

form a basis of  $V_\alpha(\lambda_1, \kappa)$ .

The elements  $E$  which can generate  $V_\alpha(\lambda_1, \kappa)$  by powers of  $(\tau - 1)$  are linear combinations

$$E = \sum_{\nu=0}^{\kappa-1} \lambda_\nu (\tau - 1)^\nu e,$$

for  $\lambda_i \in k$  and  $\lambda_0 \neq 0$ .

On the other hand using eq. (6) we see that  $\sigma$  with respect to the basis given in eq. (7) admits the matrix form:

$$\begin{pmatrix} \zeta_m^\lambda & 0 & \cdots & \cdots & 0 \\ 0 & \alpha\zeta_m^\lambda & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & \vdots \\ 0 & \cdots & \cdots & 0 & \alpha^{\kappa-1}\zeta_m^\lambda \end{pmatrix}.$$

It is now easy to see from the above matrix that every eigenvector of the eigenvalue  $\alpha^\nu \lambda_1$ ,  $\nu > 1$  is expressed as a linear combination of the basis given in eq. (7), where the coefficient of  $e$  is zero.

Therefore, the eigenvector of the eigenvalue  $\alpha^\nu \zeta_m$  can not generate the module  $V_\alpha(\lambda, \kappa)$  by powers of  $(\sigma - 1)^\nu$ .  $\square$

**3.2. The uniserial description.** We will now give an alternative description of the indecomposable  $C_q \rtimes C_m$ -modules, which is used in [2].

It is known that  $\text{Aut}(C_q) \cong \mathbb{F}_p^* \times Q$ , for some abelian  $p$ -group  $Q$ . The representation  $\psi : C_m \rightarrow \text{Aut}(C_q)$  given by the action of  $C_m$  on  $C_q$  is known to factor through a character  $\chi : C_m \rightarrow \mathbb{F}_p^*$ . The order of  $\chi$  divides  $p-1$  and  $\chi^{p-1} = \chi^{-(p-1)}$  is the trivial one dimensional character.

For all  $i \in \mathbb{Z}$ ,  $\chi^i$  defines a simple  $k[C_m]$ -module of  $k$  dimension one, which we will denote by  $S_{\chi^i}$ . For  $0 \leq \ell \leq m-1$  denote by  $S_\ell$  the simple module where on which  $\sigma$  acts as  $\zeta_m^\ell$ . Both  $S_{\chi^i}$ ,  $S_\ell$  can be seen as  $k[C_q \rtimes C_m]$ -modules using inflation. Finally for  $0 \leq \ell \leq m-1$  we define  $\chi^i(\ell) \in \{0, 1, \dots, m-1\}$  such that  $S_{\chi^i(\ell)} \cong S_\ell \otimes_k S_{\chi^i}$ .

There are  $q \cdot m$  isomorphism classes of indecomposable  $k[C_q \rtimes C_m]$ -modules and are all uniserial. An indecomposable  $k[C_q \rtimes C_m]$ -module  $U$  is unique determined by its socle, which is the kernel of the action of  $\tau - 1$  on  $U$ , and its  $k$ -dimension. For  $0 \leq \ell \leq m-1$  and  $1 \leq \mu \leq q$ , let  $U_{\ell, \mu}$  be the indecomposable  $k[C_q \rtimes C_m]$  module with socle  $S_a$  and  $k$ -dimension  $\mu$ . Then  $U_{\ell, \mu}$  is uniserial and its  $\mu$  ascending composition factors are the first  $\mu$  composition factors of the sequence

$$S_\ell, S_{\chi^{-1}(\ell)}, S_{\chi^{-2}(\ell)}, \dots, S_{\chi^{-(p-2)}(\ell)}, S_\ell, S_{\chi^{-1}(\ell)}, S_{\chi^{-2}(\ell)}, \dots, S_{\chi^{-(p-2)}(\ell)}.$$

Notice that in our notation  $V_\alpha(\lambda, \kappa) = U_{\lambda+\kappa, \kappa}$ .

**Remark 12.** The condition  $\text{ord}_{p^i} = m$  for all  $1 \leq i \leq h$ , is equivalent to requiring that  $\psi_i : C_m \rightarrow \text{Aut}(C_{p^i})$  is faithful for all  $i$ .

#### 4. LIFTING OF REPRESENTATIONS

**Proposition 13.** *Let  $G = C_q \rtimes C_m$ . Assume that for all  $1 \leq i \leq h$ ,  $\text{ord}_{p^i} a = m$ . If the  $G$ -module  $V$  lifts to an  $R[G]$ -module  $\tilde{V}$ , where  $K = \text{Quot}(R)$  is a field of characteristic zero, then*

$$m \mid \left( \dim(\tilde{V} \otimes_R K) - \dim(\tilde{V} \otimes_R K)^{C_q} \right).$$

Moreover, if  $\tilde{V}(\zeta_q^{\alpha^i \kappa})$  is the eigenspace of the eigenvalue  $\zeta_q^{\alpha^i \kappa}$  of  $T$  acting on  $\tilde{V}$ , then

$$\dim \tilde{V}(\zeta_q^\kappa) = \dim \tilde{V}(\zeta_q^{\alpha \kappa}) = \dim \tilde{V}(\zeta_q^{\alpha^2 \kappa}) = \dots = \dim \tilde{V}(\zeta_q^{\alpha^{m-1} \kappa}).$$

*Proof.* Consider a lifting  $\tilde{V}$  of  $V$ . The generator  $\tau$  of the cyclic part  $C_q$  has eigenvalues  $\lambda_1, \dots, \lambda_s$  which are  $p^n$ -roots of unity. Let  $\zeta_q$  be a primitive  $q$ -root of unity. Consider any eigenvalue  $\lambda \neq 1$ . It is of the form  $\lambda = \zeta_q^\kappa$  for some  $\kappa \in \mathbb{N}$ ,  $q \nmid \kappa$ . If  $E$  is an eigenvector of  $T$  corresponding to  $\lambda$ , that is  $\tau E = \zeta_q^\kappa E$  then

$$\tau \sigma^{-1} E = \sigma^{-1} \tau^\alpha E = \zeta_q^{\kappa \alpha^{m-1}} \sigma^{-1} E$$

and we have a series of eigenvectors  $E, \sigma^{-1} E, \sigma^{-2} E, \dots$  with corresponding eigenvalues  $\zeta_q^\kappa, \zeta_q^{\kappa \alpha}, \zeta_q^{\kappa \alpha^2}, \dots, \zeta_q^{\kappa \alpha^o}$ , where  $o = \text{ord}_{q/(q, \kappa)}$ . Indeed, the integer  $o$  satisfies the

$$\kappa \alpha^o \equiv \kappa \pmod{q} \Rightarrow \alpha^m \equiv 1 \pmod{\frac{q}{(q, \kappa)}}.$$

Therefore the eigenvalues  $\lambda \neq 1$  form orbits of size  $m$ , while the eigenspace of the eigenvalue 1 is just the invariant space  $V^G$  and the result follows.  $\square$



5. INDECOMPOSABLE  $C_q \rtimes C_m$  MODULES, INTEGRAL REPRESENTATION THEORY

From now on  $V$  be a free  $R$ -module, where  $R$  is an integral local principal ideal domain with maximal ideal  $\mathfrak{m}_R$ ,  $R$  has characteristic zero and that  $R$  contains all  $q$ -th roots of unity and has characteristic zero. Let  $K = \text{Quot}(R)$ .

The indecomposable modules for a cyclic group both in the ordinary and in the modular case are described by writing down the Jordan normal form of a generator of the cyclic group. Since in integral representation theory there are infinitely many non-isomorphic indecomposable  $C_q$ -modules for  $q = p^h$ ,  $h \geq 3$ , one is not expecting to have a theory of Jordan normal forms even if one works over complete local principal ideal domains [7], [8].

**Lemma 14.** *Let  $T$  be an element of order  $q = p^h$  in  $\text{End}(V)$ , then the minimal polynomial of  $T$  has simple eigenvalues and  $T$  is diagonalizable when seen as an element in  $\text{End}(V \otimes K)$ .*

*Proof.* Since  $T^q = \text{Id}_V$ , the minimal polynomial of  $T$  divides  $x^q - 1$ , which has simple roots over a field of characteristic zero. This ensures that  $T \in \text{End}(V \otimes K)$  is diagonalizable.  $\square$

**Lemma 15.** *Let  $f(x) = (x - \lambda_1)(x - \lambda_2) \cdots (x - \lambda_d)$  be the minimal polynomial of  $T$  on  $V$ . There is an element  $E \in V$ , such that*

*$E, (T - \lambda_1 \text{Id}_V)E, (T - \lambda_2 \text{Id}_V)(T - \lambda_1 \text{Id}_V)E, \dots, (T - \lambda_{d-1} \text{Id}_V) \cdots (T - \lambda_1 \text{Id}_V)E$*   
*are linear independent elements in  $V \otimes K$ .*

*Proof.* Consider the endomorphisms for  $i = 1, \dots, d$

$$\Pi_i = \prod_{\substack{\nu=1 \\ \nu \neq i}}^d (T - \lambda_\nu \text{Id}_V).$$

In the above product notice that  $T - \lambda_i \text{Id}_V$ ,  $T - \lambda_j \text{Id}_V$  are commuting endomorphisms. Since the minimal polynomial of  $T$  has degree  $d$  all  $R$ -modules  $\text{Ker} \Pi_i$  are strictly less than  $V$ . Moreover there is an element  $E$  such that  $E \notin \text{Ker}(\Pi_i)$  for all  $1 \leq i \leq d$ . Consider a relation

$$(8) \quad \sum_{\mu=0}^d \gamma_\mu \prod_{\nu=0}^{\mu} (T - \lambda_\nu \text{Id}_V)E,$$

where  $\prod_{\nu=0}^0 (T - \lambda_\nu \text{Id}_V)E = E$ . We first apply the operator  $\prod_{\nu=2}^d (T - \lambda_\nu \text{Id}_V)$  to eq. (8) and we obtain

$$0 = \gamma_0 \Pi_1 E,$$

and by the selection of  $E$  we have that  $\gamma_0 = 0$ . We now apply  $\prod_{\nu=3}^d (T - \lambda_\nu \text{Id}_V)$  to eq. (8). We obtain that

$$0 = \gamma_1 \prod_{\nu=3}^d (T - \lambda_\nu \text{Id}_V)(T - \lambda_1 \text{Id}_V)E = \gamma_1 \Pi_2 E,$$

and by the selection of  $E$  we have that  $\gamma_1 = 0$ . We now apply  $\prod_{\nu=4}^d (T - \lambda_\nu \text{Id}_V)$  to eq. (8) and we obtain

$$0 = \gamma_2 \prod_{\nu=4}^d (T - \lambda_\nu \text{Id}_V)(T - \lambda_2 \text{Id}_V)(T - \lambda_1 \text{Id}_V)E = \gamma_2 \Pi_3 E$$

and by the selection of  $E$  we obtain  $\gamma_3 = 0$ . Continuing this way we finally arrive at  $\gamma_0 = \gamma_1 = \cdots = \gamma_{d-1} = 0$ .  $\square$

**Lemma 16.** *Let  $V$  be a free  $R$ -module of rank  $R$  acted on by an automorphism  $T : V \rightarrow V$  of order  $p^h$ . Assume that the minimal polynomial of  $T$  is of degree  $d$  and has roots  $\lambda_1, \dots, \lambda_d$ . Then  $T$  can be written as a matrix with respect to the basis as follows:*

$$(9) \quad \begin{pmatrix} \lambda_1 & 0 & \cdots & \cdots & 0 \\ a_1 & \lambda_2 & \ddots & & \vdots \\ 0 & a_2 & \lambda_3 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & a_{d-1} & \lambda_d \end{pmatrix}$$

*Proof.* By lemma 15 the elements

$$E, (T - \lambda_1 \text{Id}_V)E, (T - \lambda_2 \text{Id}_V)(T - \lambda_1 \text{Id}_V)E, \dots, (T - \lambda_{d-1} \text{Id}_V) \cdots (T - \lambda_1 \text{Id}_V)E$$

form a free submodule of  $V$  of rank  $d$ . The theory of submodules of principal ideal domains, there is a basis  $E_1, E_2, \dots, E_d$  of the free module  $V$  such that

$$(10) \quad \begin{aligned} E_1 &= E, \\ a_1 E_2 &= (T - \lambda_1 \text{Id}_V)E_1, \\ a_2 E_3 &= (T - \lambda_2 \text{Id}_V)E_2, \\ &\dots \\ a_{s-1} E_d &= (T - \lambda_{d-1} \text{Id}_V)E_{d-1}. \end{aligned}$$

Let us consider the module  $V_1 = \langle E_1, \dots, E_d \rangle \subset V$ . By construction, the map  $T$  restricts to an automorphism  $V_1 \rightarrow V_1$  with respect to the basis  $E_1, \dots, E_d$  has the desired form. We then consider the free module  $V/V_1$  and we repeat the procedure for the minimal polynomial of  $T$ , which again acts on  $V/V_1$ . The desired result follows.  $\square$

**Remark 17.** The element  $T$  as defined in eq. (9) has order equal to the higher order of the eigenvalues  $\lambda_1, \dots, \lambda_d$  involved. Indeed, since we have assumed that the eigenvalues are different the matrix is diagonalizable in  $\text{Quot}(R)$  and has order equal to the maximal order of the eigenvalues involved. In particular it has order  $q$  if there is at least one  $\lambda_i$  that is a primitive  $q$ -root of unity. The statement about the order of  $T$  is not necessarily true if some of the eigenvalues are the same. For instance the matrix  $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$  has infinite order over a field of characteristic zero.

**Remark 18.** The number of indecomposable  $R[T]$ -summands of  $V$  is given by  $\#\{i : a_i = 0\} + 1$ .

A lift of a sum of indecomposable  $kC_q$ -modules  $J_{\kappa_1} \oplus \cdots \oplus J_{\kappa_n}$  can form an indecomposable  $RC_q$ -module. For example the indecomposable module where the

generator  $T$  of  $C_q$  has the form

$$T = \begin{pmatrix} \lambda_1 & 0 & \cdots & \cdots & 0 \\ a_1 & \lambda_2 & \ddots & & \vdots \\ 0 & a_2 & \lambda_3 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & a_{s-1} & \lambda_d \end{pmatrix}$$

where  $a_1 = \cdots = a_{\kappa_1-1} = 1$ ,  $a_{\kappa_1} \in \mathfrak{m}_R$ ,  $a_{\kappa_1+1}, \dots, a_{\kappa_2+\kappa_1-1} = 1$ ,  $a_{\kappa_2+\kappa_1} \in \mathfrak{m}_R$ , etc reduces to a decomposable direct sum of Jordan normal forms of sizes  $J_{\kappa_1}, J_{\kappa_2-\kappa_1}, \dots$ .

**Remark 19.** It is an interesting question to classify these matrices up to conjugation with a matrix in  $\mathrm{GL}_d(R)$ . It seems that the valuation of elements  $a_i$  should also play a role.

**Definition 20.** Let  $h_i(x_1, \dots, x_j)$  be the complete symmetric polynomial of degree  $i$  in the variables  $x_1, \dots, x_j$ . For instance

$$h_3(x_1, x_2, x_3) = x_1^3 + x_1^2x_2 + x_1^2x_3 + x_1x_2^2 + x_1x_2x_3 + x_1x_3^2 + x_2^3 + x_2^2x_3 + x_2x_3^2 + x_3^3.$$

Set

$$L(\kappa, j, \nu) = h_\kappa(\lambda_j, \lambda_{j+1}, \dots, \lambda_{j+\nu})$$

$$A(i, j) = \begin{cases} a_i a_{i+1} \cdots a_{i+j} & \text{if } j \geq 0 \\ 0 & \text{if } j < 0 \end{cases}$$

**Lemma 21.** The matrix  $T^\alpha = (t_{ij}^{(\alpha)})$  is given by the following formula:

$$t_{ij}^{(\alpha)} = \begin{cases} \lambda_i^\alpha & \text{if } i = j \\ A(j, i-j-1) \cdot L(\alpha - (i-j), j, i-j) & \text{if } j < i \\ 0 & \text{if } j > i \end{cases}$$

*Proof.* For  $j \geq i$  the proof is trivial. When  $j < i$  and  $\alpha = 1$  it is immediate, since  $L(x, \cdot, \cdot) \equiv 0$ , for every  $x \leq 0$ . Assume this holds for  $\alpha = n$ . If  $\alpha = n+1$ ,

$$\begin{aligned} t_{ij}^{(n+1)} &= t_{ij}^{(n)} t_{ij} = \sum_{k=1}^r t_{ik}^{(\alpha)} t_{kj} = \lambda_j t_{ij}^{(\alpha)} + a_j t_{ij+1}^{(\alpha)} = \lambda_j A(j, i-j-1) L(\alpha - (i-j), j, i-j) + \\ &+ a_j A(j+1, i-j-2) L(\alpha - (i-j-1), j+1, i-j-1) = \\ &= A(j, i-j-1) (\lambda_j h_{\alpha-(i-j)}(\lambda_j, \dots, \lambda_j) + h_{\alpha-(i-j)+1}(\lambda_{j+1}, \dots, \lambda_i)) = \\ &= A(j, i-j-1) h_{\alpha-(i-j)+1}(\lambda_j, \dots, \lambda_i) = \\ &= A(j, i-j-1) L(\alpha - (i-j) + 1, i, i-j). \end{aligned}$$

□

**Remark 22.** The space of homogeneous polynomials of degree  $k$  in  $n$ -variables has dimension  $\binom{n-1+c}{n-1}$ . Since all  $q$ -roots of unity are reduced to 1 modulo  $\mathfrak{m}_R$  the quantity  $L(\alpha - (i-j), j, i-j)$  is reduced to  $n = (i-j) + 1$ ,  $c = \alpha - (i-j)$

$$\binom{n-1+c}{n-1} = \binom{\alpha}{i-j}.$$

This equation is compatible with the computation of  $\tau^\alpha$  given in eq. (4).

**Lemma 23.** *There is an eigenvector  $E$  of the generator  $\sigma$  of the cyclic group  $C_m$  which is not an element in  $\bigcup_{i=1}^s \text{Ker}(\Pi_i \otimes K)$ .*

*Proof.* The eigenvectors  $E_1, \dots, E_d$  of  $\sigma$  form a basis of the space  $V \otimes K$ . By multiplying by certain elements in  $R$ , if necessary, we can assume that all  $E_i$  are in  $V$  and their reductions  $E_i \otimes R/\mathfrak{m}_R$ ,  $1 \leq i \leq d$  give rise to a basis of eigenvectors of a generator of the cyclic group  $C_m$  acting on  $V \otimes R/\mathfrak{m}_R$ . If every eigenvector  $E_i$  is an element of some  $\text{Ker}(\Pi_\nu)$  for  $1 \leq i \leq d$ , then their reductions will be elements in  $\text{Ker}(T - 1)^{d-1}$ , a contradiction since the later kernel has dimension  $< d$ .  $\square$

**Lemma 24.** *Let  $V$  be a free  $C_q \rtimes C_m$ -module, which is indecomposable as a  $C_q$ -module. Consider the basis given in lemma 16. Then the value of  $\sigma(E_1)$  determines  $\sigma(E_i)$  for  $2 \leq i \leq d$ .*

*Proof.* Let  $\sigma$  be a generator of the cyclic group  $C_m$ . We will use the notation of lemma 15. We use lemma 23 in order to select a suitable eigenvector of  $E_1$  of  $\sigma$  and then form the basis  $E_1, E_2, \dots, E_d$  as given in eq. (10). We can compute the action of  $\sigma$  on all basis elements  $E_i$  by

$$(11) \quad \sigma(a_{i-1}E_i) = \sigma(T - \lambda_{i-1}\text{Id}_V)E_{i-1} = (T^a - \lambda_{i-1}\text{Id}_V)\sigma(E_{i-1}).$$

This means that one can define recursively the action of  $\sigma$  on all elements  $E_i$ . Indeed, assume that

$$\sigma(E_{i-1}) = \sum_{\nu=1}^d \gamma_{\nu, i-1} E_\nu.$$

We now have

$$\begin{aligned} (T^a - \lambda_{i-1}\text{Id}_V)E_\nu &= \sum_{\mu=1}^d t_{\mu, \nu}^{(\alpha)} E_\mu - \lambda_{i-1} E_\nu \\ &= (\lambda_\nu^\alpha - \lambda_{i-1})E_\nu + \sum_{\mu=\nu+1}^d t_{\mu, \nu}^{(\alpha)} E_\mu \end{aligned}$$

We combine all the above to

$$\begin{aligned} a_{i-1}\sigma(E_i) &= \sum_{\nu=1}^d \gamma_{\nu, i-1} (\lambda_\nu^\alpha - \lambda_{i-1})E_\nu + \sum_{\nu=1}^d \gamma_{\nu, i-1} \sum_{\mu=\nu+1}^d t_{\mu, \nu}^{(\alpha)} E_\mu \\ (12) \quad &= \sum_{\nu=1}^d \tilde{\gamma}_{\nu, i} E_\nu, \end{aligned}$$

for a selection of elements  $\gamma_{\nu, i} \in R$ , which can be explicitly computed by collecting the coefficients of the basis elements  $E_1, \dots, E_d$ .

Observe that the quantity on the right hand side of eq. (12) must be divisible by  $a_{i-1}$ . Indeed, let  $v$  be the valuation of the local principal ideal domain  $R$ . Set

$$e_0 = \min_{1 \leq \nu \leq d} \{v(\tilde{\gamma}_{\nu, i})\}.$$

If  $e_0 < v(a_{i-1})$  then we divide eq. (12) by  $\pi^{e_0}$  where  $\pi$  is the local uniformizer of  $R$ , that is  $\mathfrak{m}_R = \pi R$ . We then consider the divided equation modulo  $\mathfrak{m}_R$  to obtain

a linear dependence relation among the elements  $E_i \otimes k$ , which is a contradiction. Therefore  $e_0 \geq v(a_{i-1})$  and we obtain an equation

$$\sigma(E_i) = \sum_{\nu=1}^d \frac{\tilde{\gamma}_{\nu,i}}{a_{i-1}} E_\nu = \sum_{\nu=1}^d \gamma_{\nu,i} E_\nu.$$

□

For example  $\sigma(E_1) = \zeta_m^\epsilon E_1$ . We compute that

$$a_1 \sigma(E_2) = (T^\alpha - \lambda_1 \text{Id}) \sigma(E_1)$$

and

$$\begin{aligned} \sigma(E_2) &= \frac{(\lambda_1^\alpha - \lambda_1)}{a_1} \zeta_m^\epsilon E_1 + \zeta_m^\epsilon \sum_{\mu=2}^d \frac{t_{\mu,1}^{(\alpha)}}{a_1} E_\mu \\ &= \frac{(\lambda_1^\alpha - \lambda_1)}{a_1} \zeta_m^\epsilon E_1 + \zeta_m^\epsilon \sum_{\mu=2}^d \frac{A(1, \mu-2) L(\alpha - (\mu-1), 1, \mu-1)}{a_1} E_\mu \\ &= \frac{(\lambda_1^\alpha - \lambda_1)}{a_1} \zeta_m^\epsilon E_1 + \zeta_m^\epsilon \sum_{\mu=2}^d \frac{a_1 a_2 \cdots a_{\mu-1} h_{\alpha - (\mu-1)}(\lambda_1, \lambda_2, \dots, \lambda_\mu)}{a_1} E_\mu. \end{aligned}$$

**Proposition 25.** *Assume that no element  $a_1, \dots, a_{d-1}$  given in eq. (9) is zero. Given  $\alpha \in \mathbb{N}, \alpha \geq 1$  and an element  $E_1$ , which is not an element in  $\bigcup_{i=1}^d \text{Ker}(\Pi_i \otimes K)$ , if there is a matrix  $\Gamma = (\gamma_{ij})$ , such that  $\Gamma T \Gamma^{-1} = T^\alpha$  and  $\Gamma E_1 = \zeta_m^\epsilon E_1$ , then this matrix  $\Gamma$  is unique.*

*Proof.* We will use the idea leading to equation (11) replacing  $\sigma$  with  $\Gamma$ . We will compute recursively and uniquely the entries  $\gamma_{\mu,i}$ , arriving at the explicit formula of eq. (18).

Observe that trivially  $\gamma_{\nu,1} = 0$  for all  $\nu < 1$  since we only allow  $1 \leq \nu \leq d$ . We compute

(13)

$$\begin{aligned} \tilde{\gamma}_{\mu,i} &= \gamma_{\mu,i-1} (\lambda_\mu^\alpha - \lambda_{i-1}) + \sum_{\nu=1}^{\mu-1} \gamma_{\nu,i-1} t_{\mu,\nu}^{(\alpha)} \\ &= \gamma_{\mu,i-1} (\lambda_\mu^\alpha - \lambda_{i-1}) + \sum_{\nu=1}^{\mu-1} \gamma_{\nu,i-1} A(\nu, \mu - \nu - 1) L(\alpha - (\mu - \nu), \nu, \mu - \nu) \\ &= \gamma_{\mu,i-1} (\lambda_\mu^\alpha - \lambda_{i-1}) + \sum_{\nu=1}^{\mu-1} \gamma_{\nu,i-1} a_\nu a_{\nu+1} \cdots a_{\mu-1} h_{\alpha - \mu + \nu}(\lambda_\nu, \lambda_{\nu+1}, \dots, \lambda_\mu) \end{aligned}$$

Define

$$\begin{aligned} [\lambda_m^\alpha - \lambda_x]_i^j &= \prod_{x=i}^j (\lambda_\mu^\alpha - \lambda_x) \\ [a]_i^j &= \prod_{x=i}^j a_x \end{aligned}$$

for  $i \leq j$ . If  $i > j$  then both of the above quantities are defined to be equal to 1.

Observe that for  $\mu = 1$  eq. (13) becomes

$$(14) \quad \gamma_{1,i} = \frac{1}{a_{i-1}} \gamma_{1,i-1} (\lambda_1^\alpha - \lambda_{i-1})$$

and we arrive at (assuming that  $\Gamma(E_1) = \zeta_m^\epsilon E_1$ )

$$(15) \quad \gamma_{1,i} = \frac{\zeta_m^\epsilon}{a_1 a_2 \cdots a_{i-1}} \prod_{x=1}^{i-1} (\lambda_1^\alpha - \lambda_x) = \frac{\zeta_m^\epsilon}{a_1 a_2 \cdots a_{i-1}} [\lambda_1^\alpha - \lambda_x]_1^{i-1}.$$

For  $\mu \geq 2$  we have  $\gamma_{\mu,1} = 0$ , since by assumption  $TE_1 = \zeta_m^\epsilon E_1$ . Therefore eq. (13) gives us

$$(16) \quad \begin{aligned} \gamma_{\mu,i} &= \sum_{\kappa_1=0}^{i-2} \frac{[\lambda_\mu^\alpha - \lambda_x]_{i-\kappa_1}^{i-1}}{[a]_{i-1-\kappa_1}^{i-1}} \sum_{\mu_2=1}^{\mu-1} \gamma_{\mu_2,i-1-\kappa_1} [a]_{\mu_2}^{\mu-1} h_{\alpha-\mu+\mu_2}(\lambda_{\mu_2}, \dots, \lambda_\mu) \\ &= \sum_{\mu_2=1}^{\mu-1} [a]_{\mu_2}^{\mu-1} h_{\alpha-\mu+\mu_2}(\lambda_{\mu_2}, \dots, \lambda_\mu) \sum_{\kappa_1=0}^{i-2} \frac{[\lambda_\mu^\alpha - \lambda_x]_{i-\kappa_1}^{i-1}}{[a]_{i-1-\kappa_1}^{i-1}} \gamma_{\mu_2,i-1-\kappa_1}. \end{aligned}$$

We will now prove eq. (16) by induction on  $i$ . For  $i = 2, \mu \geq 2$  we have

$$\begin{aligned} \gamma_{\mu,2} &= \frac{1}{a_1} \gamma_{\mu,1} (\lambda_\mu^\alpha - \lambda_1) + \frac{1}{a_1} \sum_{\mu_2=1}^{\mu-1} \gamma_{\mu_2,1} [a]_{\mu_2}^{\mu-1} h_{\alpha-\mu+\mu_2}(\lambda_{\mu_2}, \dots, \lambda_\mu) \\ &= \frac{1}{a_1} [a]_1^{\mu-1} h_{\alpha-\mu+1}(\lambda_1, \dots, \lambda_\mu) \gamma_{1,1}. \end{aligned}$$

Assume now that eq. (16) holds for computing  $\gamma_{\mu, i-1}$ . We will treat the  $\gamma_{\mu, i}$  case. We have

$$\begin{aligned}
 \gamma_{\mu, i} &= \frac{(\lambda_{\mu}^{\alpha} - \lambda_{i-1})}{a_{i-1}} \gamma_{\mu, i-1} + \frac{1}{a_{i-1}} \sum_{\mu_2=1}^{\mu-1} \gamma_{\mu_2, i-1} [a]_{\mu_2}^{\mu-1} h_{\alpha-\mu+\mu_2}(\lambda_{\mu_2}, \dots, \lambda_{\mu}) \\
 &= \frac{(\lambda_{\mu}^{\alpha} - \lambda_{i-1})}{a_{i-1}} \sum_{\mu_2=1}^{\mu-1} [a]_{\mu_2}^{\mu-1} h_{\alpha-\mu+\mu_2}(\lambda_{\mu_2}, \dots, \lambda_{\mu}) \sum_{\kappa_1=0}^{i-3} \frac{[\lambda_{\mu}^{\alpha} - \lambda_x]_{i-1-\kappa_1}^{i-2}}{[a]_{i-2-\kappa_1}^{i-2}} \gamma_{\mu_2, i-2-\kappa_1} \\
 &+ \frac{1}{a_{i-1}} \sum_{\mu_2=1}^{\mu-1} \gamma_{\mu_2, i-1} [a]_{\mu_2}^{\mu-1} h_{\alpha-\mu+\mu_2}(\lambda_{\mu_2}, \dots, \lambda_{\mu}) \\
 &= \sum_{\mu_2=1}^{\mu-1} [a]_{\mu_2}^{\mu-1} h_{\alpha-\mu+\mu_2}(\lambda_{\mu_2}, \dots, \lambda_{\mu}) \sum_{\kappa_1=0}^{i-3} \frac{[\lambda_{\mu}^{\alpha} - \lambda_x]_{i-1-\kappa_1}^{i-1}}{[a]_{i-2-\kappa_1}^{i-1}} \gamma_{\mu_2, i-2-\kappa_1} \\
 &+ \frac{1}{a_{i-1}} \sum_{\mu_2=1}^{\mu-1} \gamma_{\mu_2, i-1} [a]_{\mu_2}^{\mu-1} h_{\alpha-\mu+\mu_2}(\lambda_{\mu_2}, \dots, \lambda_{\mu}) \\
 &= \sum_{\mu_2=1}^{\mu-1} [a]_{\mu_2}^{\mu-1} h_{\alpha-\mu+\mu_2}(\lambda_{\mu_2}, \dots, \lambda_{\mu}) \sum_{\kappa_1=1}^{i-2} \frac{[\lambda_{\mu}^{\alpha} - \lambda_x]_{i-\kappa_1}^{i-1}}{[a]_{i-1-\kappa_1}^{i-1}} \gamma_{\mu_2, i-1-\kappa_1} \\
 &+ \sum_{\mu_2=1}^{\mu-1} [a]_{\mu_2}^{\mu-1} h_{\alpha-\mu+\mu_2}(\lambda_{\mu_2}, \dots, \lambda_{\mu}) \frac{1}{a_{i-1}} \gamma_{\mu_2, i-1} \\
 &= \sum_{\mu_2=1}^{\mu-1} [a]_{\mu_2}^{\mu-1} h_{\alpha-\mu+\mu_2}(\lambda_{\mu_2}, \dots, \lambda_{\mu}) \sum_{\kappa_1=0}^{i-2} \frac{[\lambda_{\mu}^{\alpha} - \lambda_x]_{i-\kappa_1}^{i-1}}{[a]_{i-1-\kappa_1}^{i-1}} \gamma_{\mu_2, i-1-\kappa_1}
 \end{aligned}$$

and equation (16) is now proved.

We proceed recursively applying eq. (16) to each of the summands  $\gamma_{\mu_2, i-1-\kappa_1}$  if  $\mu_2 > 1$  and  $i-1-\kappa_1 > 1$ . If  $\mu_2 = 1$ , then  $\gamma_{\mu_2, i-1-\kappa_1}$  is computed by eq. (14) and if  $\mu_2 > 1$  and  $i-1-\kappa_1 \leq 1$  then  $\gamma_{\mu_2, i-1-\kappa_1} = 0$ . We can classify all iterations needed by the set  $\Sigma_{\mu}$  of sequences  $(\mu_s, \mu_{s-1}, \dots, \mu_3, \mu_2)$  such that

$$(17) \quad 1 = \mu_s < \mu_{s-1} < \dots < \mu_3 < \mu_2 < \mu = \mu_1.$$


For example for  $\mu = 5$  the set of such sequences is given by

$$\Sigma_{\mu} = \{(1), (1, 2), (1, 3), (1, 2, 3), (1, 4), (1, 2, 4), (1, 3, 4), (1, 2, 3, 4)\}$$

corresponding to the tree of iterations given in figure 1. The length of the sequence  $(\mu_s, \mu_{s-1}, \dots, \mu_2)$  is given in eq. (17) is  $s-1$ . In each iteration the  $i$  changes to  $i-1-k$  thus we have the following sequence of indices

$$i_1 = i \rightarrow i_2 = i-1-\kappa_1 \rightarrow i_3 = i-2-(\kappa_1+\kappa_2) \rightarrow \dots \rightarrow i_s = i-(s-1)-(\kappa_1+\dots+\kappa_{s-1})$$

For the sequence  $i_1, i_2, \dots$ , we might have  $i_t = 1$  for  $t < s-1$ . But in this case, we will arrive at the element  $\gamma_{\mu_{t+1}, i_t} = \gamma_{\mu_t, 1} = 0$  since  $\mu_t > 1$ . This means that we will have to consider only selections  $\kappa_1, \dots, \kappa_{s-1}$  such that  $i_{s-1} \geq 1$ . Therefore we

FIGURE 1. Iteration tree for  $\mu = 5$ 

arrive at the following expression for  $\mu \geq 2$

$$\begin{aligned}
\gamma_{\mu,i} &= \sum_{(\mu_s, \dots, \mu_2) \in \Sigma_\mu} [a]_{\mu_2}^{\mu-1} [a]_{\mu_3}^{\mu_2-1} \dots [a]_{\mu_s}^{\mu_{s-1}-1} \prod_{\nu=2}^s h_{\alpha-\mu_{\nu-1}+\mu_\nu}(\lambda_{\mu_\nu}, \dots, \lambda_{\mu_{\nu-1}}) \\
&\cdot \sum_{i=i_1 > i_2 > \dots > i_s \geq 1} \prod_{\nu=1}^{s-1} \frac{[\lambda_{\mu_\nu}^\alpha - \lambda_x]_{i_{\nu+1}+1}^{i_\nu-1}}{[a]_{i_{\nu+1}}^{i_\nu-1}} \cdot \gamma_{1, i_s}. \\
&= \sum_{(\mu_s, \dots, \mu_2) \in \Sigma_\mu} \prod_{\nu=2}^s h_{\alpha-\mu_{\nu-1}+\mu_\nu}(\lambda_{\mu_\nu}, \dots, \lambda_{\mu_{\nu-1}}) \\
&\cdot \sum_{i=i_1 > i_2 > \dots > i_s \geq 1} \frac{[a]_1^{\mu-1}}{[a]_{i_s}^{i-1}} \prod_{\nu=1}^{s-1} [\lambda_{\mu_\nu}^\alpha - \lambda_x]_{i_{\nu+1}+1}^{i_\nu-1} \frac{\zeta_m^\epsilon [\lambda_1^\alpha - \lambda_x]_1^{i_s-1}}{[a]_1^{i_s-1}} \\
(18) \quad &= \sum_{(\mu_s, \dots, \mu_2) \in \Sigma_\mu} \prod_{\nu=2}^s h_{\alpha-\mu_{\nu-1}+\mu_\nu}(\lambda_{\mu_\nu}, \dots, \lambda_{\mu_{\nu-1}}) \frac{[a]_1^{\mu-1}}{[a]_1^{i-1}} \zeta_m^\epsilon \sum_{i=i_1 > i_2 > \dots > i_s \geq 1} \prod_{\nu=1}^s [\lambda_{\mu_\nu}^\alpha - \lambda_x]_{i_{\nu+1}+1}^{i_\nu-1}
\end{aligned}$$

where  $i_{s+1} + 1 = 1$  that is  $i_{s+1} = 0$ .  $\square$

We will now prove that the matrix  $\Gamma$  of lemma 25 exists by cheking that  $\Gamma T = T^\alpha \Gamma$ . Set  $(a_{\mu,i}) = \Gamma T$ ,  $(b_{\mu,i}) = T^\alpha \Gamma$ . For  $i < d$  we have

$$\begin{aligned}
a_{\mu,i} &= \sum_{\nu=1}^d \gamma_{\mu,\nu} t_{\nu,i} = \gamma_{\mu,i} t_{ii} + \gamma_{\mu,i+1} t_{i+1,i} \\
&= \gamma_{\mu,i} \lambda_i + \gamma_{\mu,i} (\lambda_\mu^\alpha - \lambda_i) + \sum_{\nu=1}^{\mu-1} \gamma_{\nu,i} t_{\mu,\nu}^{(\alpha)} \\
&= \gamma_{\mu,i} \lambda_\mu^\alpha + \sum_{\nu=1}^{\mu-1} \gamma_{\nu,i} t_{\mu,\nu}^{(\alpha)} = \sum_{\nu=1}^{\mu} t_{\mu,\nu}^{(\alpha)} \gamma_{\nu,i} = b_{\mu,i}.
\end{aligned}$$

For  $i = d$  we have:

$$a_{\mu,d} = \sum_{\nu=1}^d \gamma_{\mu,\nu} t_{\nu,d} = \gamma_{\mu,d} t_{d,d} = \gamma_{\mu,d} \lambda_d$$



while

$$b_{\mu,d} = \sum_{\nu=1}^d t_{\mu,\nu}^{(\alpha)} \gamma_{\nu,d} = \sum_{\nu=1}^{\mu-1} t_{\mu,\nu}^{(\alpha)} \gamma_{\nu,d} + \lambda_{\mu}^{\alpha} \gamma_{\mu,d}$$

This gives us the relation

$$(19) \quad (\lambda_d - \lambda_{\mu}^{\alpha}) \gamma_{\mu,d} = \sum_{\nu=1}^{\mu-1} t_{\mu,\nu}^{(\alpha)} \gamma_{\nu,d}$$

For  $\mu = 1$  using eq. (15) we have

$$\gamma_{1,d} \lambda_d = \gamma_{1,d} \lambda_1^{\alpha} \Rightarrow [\lambda_1^{\alpha} - \lambda_x]_1^d = 0.$$

This relation is satisfied if  $\lambda_1^{\alpha}$  is one of  $\{\lambda_1, \dots, \lambda_d\}$ . Without loss of generality we assume that

$$(20) \quad \lambda_i^{(a)} = \begin{cases} \lambda_{i+1} & \text{if } m \nmid i \\ \lambda_{i-m+1} & \text{if } m \mid i \end{cases}$$

We have the following conditions:

$$\begin{aligned} \mu = 2 & \quad (\lambda_d - \lambda_2^{\alpha}) \gamma_{2,d} = t_{2,1}^{(\alpha)} \gamma_{1,d} \\ \mu = 3 & \quad (\lambda_d - \lambda_3^{\alpha}) \gamma_{3,d} = t_{3,1}^{(\alpha)} \gamma_{1,d} + t_{3,2}^{(\alpha)} \gamma_{2,d} \\ \mu = 4 & \quad (\lambda_d - \lambda_4^{\alpha}) \gamma_{4,d} = t_{4,1}^{(\alpha)} \gamma_{1,d} + t_{4,2}^{(\alpha)} \gamma_{2,d} + t_{4,3}^{(\alpha)} \gamma_{3,d} \\ & \quad \vdots \\ \mu = d-1 & \quad (\lambda_d - \lambda_{d-1}^{\alpha}) \gamma_{d-1,d} = t_{d-1,1}^{(\alpha)} \gamma_{1,d} + t_{d-1,2}^{(\alpha)} \gamma_{2,d} + \dots + t_{d-1,d-2}^{(\alpha)} \gamma_{d-2,d} \end{aligned}$$

All these equations are true provided that  $\gamma_{1,d}, \dots, \gamma_{d-2,d} = 0$ . Finally, for  $\mu = d$ , we have

$$(21) \quad (\lambda_d - \lambda_d^{\alpha}) \gamma_{d,d} = \sum_{\nu=1}^{d-1} t_{d,\nu}^{(\alpha)} \gamma_{\nu,d}$$

which is true provided that  $(\lambda_d - \lambda_d^{\alpha}) \gamma_{d,d} = t_{d,d-1}^{(\alpha)} \gamma_{d-1,d}$ .

**Lemma 26.** For  $n \geq 2$  the vertical sum  $S_n$  of the products of every line of the following array

$y$						
1	1	$(x_1 - x_2)$	$(x_1 - x_3)$	$\dots$	$\dots$	$(x_1 - x_n)$
2	$(z - x_1)$	1	$(x_1 - x_3)$	$\dots$	$\dots$	$(x_1 - x_n)$
3	$(z - x_1)$	$(z - x_2)$	1	$\ddots$	$\ddots$	$\vdots$
$\vdots$	$\vdots$	$\ddots$	$\ddots$	$\ddots$	$\ddots$	$\vdots$
$\vdots$	$\vdots$	$\ddots$	$\ddots$	$\ddots$	$\ddots$	$\vdots$
$n-1$	$(z - x_1)$	$(z - x_2)$	$\dots$	$(z - x_{n-2})$	1	$(x_1 - x_n)$
$n$	$(z - x_1)$	$(z - x_2)$	$\dots$	$(z - x_{n-2})$	$(z - x_{n-1})$	1

is given by

$$S_n = \sum_{y=1}^n \prod_{\nu=y+1}^n (x_1 - x_{\nu}) \prod_{\mu=1}^{y-1} (z - x_{\mu}) = (z - x_2) \dots (z - x_n).$$

In particular when  $z = x_n$  the sum is zero.

*Proof.* We will prove the lemma by induction. For  $n = 2$  we have  $S_2 = (x_1 - x_2) + (z - x_1) = z - x_2$ . Assume that the equality holds for  $n$ . The sum  $S_{n+1}$  corresponds to the array:

$y$						
1	1	$(x_1 - x_2)$	$(x_1 - x_3)$	$\cdots$	$(x_1 - x_n)$	$(x_1 - x_{n+1})$
2	$(z - x_1)$	1	$(x_1 - x_3)$	$\cdots$	$(x_1 - x_n)$	$(x_1 - x_{n+1})$
3	$(z - x_1)$	$(z - x_2)$	1	$\ddots$	$\vdots$	$\vdots$
$\vdots$	$\vdots$		$\ddots$	$\ddots$	$\vdots$	$\vdots$
$n - 1$	$(z - x_1)$	$\cdots$	$(z - x_{n-2})$	1	$(x_1 - x_n)$	$(x_1 - x_{n+1})$
$n$	$(z - x_1)$	$(z - x_2)$	$\cdots$	$(z - x_{n-1})$	1	$(x_1 - x_{n+1})$
$n + 1$	$(z - x_1)$	$(z - x_2)$	$\cdots$	$(z - x_{n-1})$	$(z - x_n)$	1

We have by definition  $S_{n+1} = S_n(x_1 - x_{n+1}) + (z - x_1)(z - x_2) \cdots (z - x_n)$ , which by induction gives

$$\begin{aligned} S_{n+1} &= (z - x_2) \cdots (z - x_n)(x_1 - x_{n+1}) + (z - x_1)(z - x_2) \cdots (z - x_n) \\ &= (z - x_2) \cdots (z - x_n)(x_1 - x_{n+1} + z - x_1) \end{aligned}$$

and gives the desired result.  $\square$

**Lemma 27.** Consider  $A < l < L < B$ . The quantity

$$\sum_{l \leq y \leq L} [\lambda_a - \lambda_x]_A^{y-1} \cdot [\lambda_b - \lambda_x]_{y+1}^B$$

equals to

$$[\lambda_a - \lambda_x]_A^{l-1} \cdot [\lambda_b - \lambda_x]_{L+1}^B \cdot \frac{[\lambda_a - \lambda_x]_l^L - [\lambda_b - \lambda_x]_l^L}{(\lambda_a - \lambda_b)}$$

*Proof.* We write

$$\begin{aligned} &\sum_{l \leq y \leq L} [\lambda_a - \lambda_x]_A^{y-1} \cdot [\lambda_b - \lambda_x]_{y+1}^B \\ &= [\lambda_a - \lambda_x]_A^{l-1} \cdot [\lambda_b - \lambda_x]_{L+1}^B \cdot \sum_{l \leq y \leq L} [\lambda_a - \lambda_x]_l^{y-1} \cdot [\lambda_b - \lambda_x]_{y+1}^L \end{aligned}$$

The last sum can be read as the vertical sum  $S$  of the products of every line in the following array:

$y$					
$l$	1	$(\lambda_b - \lambda_{l+1})(\lambda_b - \lambda_{l+2})$	$\cdots$	$(\lambda_b - \lambda_{L-1})(\lambda_b - \lambda_L)$	
$l + 1$	$(\lambda_a - \lambda_l)$	1	$(\lambda_b - \lambda_{l+2})$	$\cdots$	$(\lambda_b - \lambda_{L-1})(\lambda_b - \lambda_L)$
$l + 2$	$(\lambda_a - \lambda_l)(\lambda_a - \lambda_{l+1})$	1		$\vdots$	$\vdots$
$\vdots$	$\vdots$	$\vdots$	$\ddots$	$\ddots$	$\vdots$
$L - 2$	$(\lambda_a - \lambda_l)(\lambda_a - \lambda_{l+1})$	$\cdots$	1	$(\lambda_b - \lambda_{L-1})(\lambda_b - \lambda_L)$	
$L - 1$	$(\lambda_a - \lambda_l)(\lambda_a - \lambda_{l+1})$	$\cdots$	$(\lambda_a - \lambda_{L-2})$	1	$(\lambda_b - \lambda_L)$
$L$	$(\lambda_a - \lambda_l)(\lambda_a - \lambda_{l+1})$	$\cdots$	$(\lambda_a - \lambda_{L-2})(\lambda_a - \lambda_{L-1})$		1

If  $l = b$ , then lemma 26 implies that  $S = [\lambda_a - \lambda_x]_{b+1}^L$ . Furthermore, if  $L = a$  then  $S = 0$ .

The quantity  $S$  cannot be directly computed using lemma 26, if  $l \neq b$ . We proceed by forming the array:

$\frac{y}{b}$	1	$(\lambda_b - \lambda_{b+1})$	$\cdots$	$(\lambda_b - \lambda_l)$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$(\lambda_b - \lambda_L)$
$\vdots$				$\vdots$					$\vdots$
$l-1$	$(\lambda_a - \lambda_b)$	$\cdots$	1	$(\lambda_b - \lambda_l)$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$(\lambda_b - \lambda_L)$
$l$	$(\lambda_a - \lambda_b)$	$\cdots$	$(\lambda_a - \lambda_{l-1})$	1	$(\lambda_b - \lambda_{l+1})$	$(\lambda_b - \lambda_{l+2})$	$\cdots$	$(\lambda_b - \lambda_{L-1})$	$(\lambda_b - \lambda_L)$
$l+1$	$(\lambda_a - \lambda_b)$	$\cdots$	$(\lambda_a - \lambda_{l-1})$	$(\lambda_a - \lambda_l)$	1	$(\lambda_b - \lambda_{l+2})$	$\cdots$	$(\lambda_b - \lambda_{L-1})$	$(\lambda_b - \lambda_L)$
$l+2$	$(\lambda_a - \lambda_b)$	$\cdots$	$(\lambda_a - \lambda_{l-1})$	$(\lambda_a - \lambda_l)(\lambda_a - \lambda_{l+1})$	1				
$\vdots$				$\vdots$	$\vdots$	$\ddots$	$\ddots$	$\vdots$	$\vdots$
$L-2$	$(\lambda_a - \lambda_b)$	$\cdots$	$(\lambda_a - \lambda_{l-1})$	$(\lambda_a - \lambda_l)(\lambda_a - \lambda_{l+1})$	$\cdots$	1	$(\lambda_b - \lambda_{L-1})$	$(\lambda_b - \lambda_L)$	
$L-1$	$(\lambda_a - \lambda_b)$	$\cdots$	$(\lambda_a - \lambda_{l-1})$	$(\lambda_a - \lambda_l)(\lambda_a - \lambda_{l+1})$	$\cdots$	$(\lambda_a - \lambda_{L-2})$	1	$(\lambda_b - \lambda_L)$	
$L$	$(\lambda_a - \lambda_b)$	$\cdots$	$(\lambda_a - \lambda_{l-1})$	$(\lambda_a - \lambda_l)(\lambda_a - \lambda_{l+1})$	$\cdots$	$(\lambda_a - \lambda_{L-2})(\lambda_a - \lambda_{L-1})$	1		

The value of this array is computed using lemma 26 to be equal to  $[\lambda_a - \lambda_x]_{b+1}^L$ . We observe that the sum of the products of the top left array can be computed using lemma 26, while the sum of the products of the lower right array is  $S$ .

$$[\lambda_a - \lambda_x]_b^{l-1} \cdot S + [\lambda_a - \lambda_x]_{b+1}^{l-1} \cdot [\lambda_b - \lambda_x]_l^L = [\lambda_a - \lambda_x]_{b+1}^L$$

we arrive at

$$[\lambda_a - \lambda_x]_b^{l-1} S = [\lambda_a - \lambda_x]_{b+1}^{l-1} ([\lambda_a - \lambda_x]_l^L - [\lambda_b - \lambda_x]_l^L)$$

or equivalently

$$(\lambda_a - \lambda_b) \cdot S = [\lambda_a - \lambda_x]_l^L - [\lambda_b - \lambda_x]_l^L$$

□

**Lemma 28.** For all  $1 \leq \mu \leq d-2$  we have  $\gamma_{\mu,d} = 0$ .

*Proof.* Let  $\mu_1 = \mu > \mu_2 > \cdots > \mu_s = 1 \in \Sigma_\mu$  be a selection of iterations and  $d = i_1 > i_2 > \cdots > i_s \geq 1 > i_{s+1} = 0$  be the sequence of  $i$ 's. Using eq. (20) we see that the quantity  $[\lambda_{\mu_\nu}^\alpha - \lambda_x]_{i_{\nu+1}+1}^{i_\nu-1} \neq 0$  if and only if one of the following two inequalities hold:

$$(22) \quad \text{either} \quad i_{\nu+1} > \mu_\nu - mf(\mu_\nu)$$

$$(23) \quad \text{or} \quad i_\nu < \mu_\nu + 2 - mf(\mu_\nu),$$

where

$$f(x) = \begin{cases} 1 & \text{if } m \mid x \\ 0 & \text{if } m \nmid x \end{cases}$$

We will denote the above two inequalities by  $(22)_\nu, (23)_\nu$  when applied for the integer  $\nu$ . Assume, that for all  $1 \leq \nu \leq s$  one of the two inequalities  $(22)_\nu, (23)_\nu$  hold, that is  $[\lambda_{\mu_\nu}^\alpha - \lambda_x]_{i_{\nu+1}+1}^{i_\nu-1} \neq 0$ . Inequality  $(22)_s$  can not hold for  $\nu = s$  since it gives us  $0 = i_{s+1} > 1 = \mu_s$ , we have  $m \nmid 1 = \mu_s$ .

We will keep the sequence  $\bar{\mu} : \mu_1 > \mu_2 > \cdots > \mu_s$  fixed and we will sum over all possible selections of sequences of  $i_1 > \cdots > i_s > i_{s+1} = 0$ , that is we will show that the sum

$$(24) \quad \Gamma_{\bar{\mu}, i} := \sum_{i=i_1 > i_2 > \cdots > i_s \geq 1} \prod_{\nu=1}^s [\lambda_{\mu_\nu}^\alpha - \lambda_x]_{i_{\nu+1}+1}^{i_\nu-1}$$

is zero, which will show that  $\gamma_{\mu,d} = 0$  using eq. (18).

Observe now that if  $(23)_\nu$  holds and  $m \nmid \nu, \nu - 1$ , then  $(23)_{\nu-1}$  also holds. Indeed the combination of  $(23)_\nu$  and  $(22)_{\nu-1}$  gives the impossible inequality

$$\mu_\nu + 2 \stackrel{(23)_\nu}{>} i_\nu \stackrel{(22)_{\nu-1}}{>} \mu_{\nu-1}.$$

Assume now that  $m \mid \nu$  and  $(23)_\nu$  holds, then  $(23)_{\nu-1}$  also holds. Indeed the combination of  $(23)_\nu$  and  $(22)_{\nu-1}$  gives us

$$\mu_\nu + 2 - m \stackrel{(23)_\nu}{>} i_\nu \stackrel{(22)_{\nu-1}}{>} \mu_{\nu-1} - mf(\mu_{\nu-1}).$$

If  $m \nmid \mu_{\nu-1}$ , then the above inequality is impossible since it implies that

$$\mu_\nu + 2 - m > \mu_{\nu-1} > \mu_\nu.$$

If  $m \mid \mu_{\nu-1}$ , then the inequality is also impossible since it implies that  $\mu_\nu + 2 > \mu_{\nu-1}$  so if we write  $\mu_{\nu-1} = k'm$  and  $\mu_\nu = km$ ,  $k, k' \in \mathbb{N}$ ,  $k' > k$ , we arrive at  $2 > (k' - k)m \geq m$ . This proves the following

**Lemma 29.** *The inequality  $(22)_{\nu-1}$  might be correct only in cases where  $m \mid \mu_{\nu-1}$ ,  $m \nmid \mu_\nu$ .*

Assume that for all  $\nu$  inequality (23) holds. Then for  $\nu = 1$  it gives us (recall that  $\mu \leq d - 2$ )

$$(25) \quad \mu + 2 \leq d = i_1 < \mu_1 + 2 - mf(\mu_1) = \mu + 2 - mf(\mu),$$

which is impossible. Therefore either there are  $\nu$  such that none of the two inequalities  $(22)_\nu, (23)_\nu$  hold (in this case the contribution to the sum is zero) or there are cases where (22) holds.

The summands appearing in eq. (24) can be zero, for example the sequence  $\mu_1 = m > \mu_2 = 1$  with  $i_2 = 2 < i_1 = d$ ,  $s = 2$  give the contribution

$$[\lambda_{\mu_2}^\alpha - \lambda_x]_1^{i_2-1} [\lambda_{\mu_1}^\alpha - \lambda_x]_{i_2}^{d-1} = [\lambda_1^\alpha - \lambda_x]_1^1 [\lambda_m^\alpha - \lambda_x]_{i_2+1}^{d-1} = (\lambda_2 - \lambda_1) [\lambda_1 - \lambda_x]_3^{d-1}$$

while for  $i_2 = 1 < i_1 = d$  it gives the contribution

$$[\lambda_{\mu_2}^\alpha - \lambda_x]_1^{i_2-1} [\lambda_{\mu_1}^\alpha - \lambda_x]_{i_2+1}^{d-1} = [\lambda_1^\alpha - \lambda_x]_1^0 [\lambda_m^\alpha - \lambda_x]_2^{d-1} = [\lambda_1 - \lambda_x]_2^{d-1}$$

It is clear that these non-zero contributions cancel out when added.

**Lemma 30.** *Assume that  $m \mid \mu_{\nu_0-1}$  and  $m \nmid \mu_{\nu_0}$ , where  $(23)_{\nu_0}$  and  $(22)_{\nu_0-1}$  hold. Then, we can eliminate  $\mu_{\nu_0-1}$  and  $i_{\nu_0}$  from both selections of the sequence of  $\mu$ 's and  $i$ 's, i.e. we can form the sequence of length  $s - 1$*

$$\bar{\mu}_{s-1} = \mu_s < \bar{\mu}_{s-2} = \mu_{s-1} < \cdots < \bar{\mu}_{\nu_0-1} = \mu_{\nu_0} < \bar{\mu}_{\nu_0-2} = \mu_{\nu_0-2} < \cdots < \bar{\mu}_1 = \mu_1.$$

and the corresponding sequence of equal length

$$\bar{i}_{s-1} = i_s < \bar{i}_{s-2} = i_{s-1} < \cdots < \bar{i}_{\nu_0-1} = i_{\nu_0-1} < \bar{i}_{\nu_0} = i_{\nu_0+1} < \cdots < \bar{i}_1 = i_1 = d,$$

so that

$$\Gamma_{\bar{\mu}, \bar{i}} = \sum_{i_1 > \cdots > i_s} \prod_{\nu=1}^s [\lambda_{\mu_\nu}^\alpha - \lambda_x]_{i_{\nu+1}+1}^{i_\nu-1} = (\star) \sum_{\bar{i}_1 > \cdots > \bar{i}_{s-1}} \prod_{\substack{\nu=1 \\ \nu \neq \nu_0-1}}^s [\lambda_{\mu_\nu}^\alpha - \lambda_x]_{i_{\nu+1}+1}^{i_\nu-1},$$

where  $(\star)$  is a non zero element.

*Proof.* (of lemma 30) We are in the case  $m \mid \mu_{\nu_0-1}$  and  $m \nmid \mu_{\nu_0}$ , where  $(23)_{\nu_0}$  and  $(22)_{\nu_0-1}$  hold,

$$(26) \quad \mu_{\nu_0-1} - m \stackrel{(22)_{\nu_0-1}}{<} i_{\nu_0} \stackrel{(23)_{\nu_0}}{<} \mu_{\nu_0} + 2,$$

or equivalently

$$\mu_0 := \mu_{\nu_0-1} - m + 1 \leq i_{\nu_0} \leq \mu_{\nu_0} + 1$$

For  $i_{\nu_0+1}$  the inequality  $(22)_{\nu_0} i_{\nu_0+1} > \mu_{\nu_0} - mf(\mu_{\nu_0})$  can not hold, since it implies

$$i_{\nu_0+1} < i_{\nu_0} \stackrel{(23)_{\nu_0}}{<} \mu_{\nu_0} + 2 < i_{\nu_0+1} + 2.$$

Observe that also

$$i_{\nu_0+1} + 1 \leq i_{\nu_0} \leq i_{\nu_0-1} - 1.$$

Set  $l = \max\{\mu_0, i_{\nu_0+1} + 1\}$  and  $L = \min\{\mu_{\nu_0} + 1, i_{\nu_0-1} - 1\}$ . Then  $y = i_{\nu_0}$  satisfies

$$l \leq y \leq L.$$

By lemma 27 the quantity

$$\sum_{l \leq y \leq L} [\lambda_{\mu_{\nu_0+1}} - \lambda_x]_{i_{\nu_0+1}+1}^{y-1} \cdot [\lambda_{\mu_0} - \lambda_x]_{y+1}^{i_{\nu_0-1}-1}$$

equals to

$$(27) \quad \frac{[\lambda_{\mu_{\nu_0+1}} - \lambda_x]_{i_{\nu_0+1}+1}^{l-1} \cdot [\lambda_{\mu_0} - \lambda_x]_{L+1}^{i_{\nu_0-1}-1} \cdot \frac{[\lambda_{\mu_{\nu_0+1}} - \lambda_x]_l^L - [\lambda_{\mu_0} - \lambda_x]_l^L}{(\lambda_{\mu_{\nu_0+1}} - \lambda_{\mu_0})}}{[\lambda_{\mu_{\nu_0+1}} - \lambda_x]_{i_{\nu_0+1}+1}^L \cdot [\lambda_{\mu_0} - \lambda_x]_{L+1}^{i_{\nu_0-1}-1} - [\lambda_{\mu_{\nu_0+1}} - \lambda_x]_{i_{\nu_0+1}+1}^{l-1} \cdot [\lambda_{\mu_0} - \lambda_x]_l^{i_{\nu_0-1}-1}} \cdot (\lambda_{\mu_{\nu_0+1}} - \lambda_{\mu_0})$$

**Case A1**  $l = \mu_0 \geq i_{\nu_0+1} + 1$ . Then  $[\lambda_{\mu_0} - \lambda_x]_l^L = 0$ .

**Case A2**  $l = i_{\nu_0+1} + 1 > \mu_0$ . We set  $z := i_{\nu_0+1}$ , which is bounded by eq.  $(23)_{\nu_0+1}$  that is

$$\text{Case A2} \quad \mu_0 \leq z \stackrel{(23)_{\nu_0+1}}{\leq} \mu_{\nu_0+1} + 1.$$

Notice that in this case  $m \nmid \mu_{\nu_0+1}$ . Indeed, we have assumed that inequality  $(23)_{\nu_0+1}$  holds which gives us

$$\mu_{\nu_0-1} - m = \mu_0 - 1 \stackrel{\text{Case A2}}{<} i_{\nu_0+1} \stackrel{(23)_{\nu_0+1}}{<} \mu_{\nu_0+1} + 2 - m,$$

which implies that  $\mu_{\nu_0-1} < \mu_{\nu_0+1} + 2$ , a contradiction. Thus for  $l = z + 1$  we compute

$$\begin{aligned} & \sum_{\mu_0 \leq z \leq \mu_{\nu_0+1}+1} [\lambda_{\mu_{\nu_0+1}}^\alpha - \lambda_x]_{i_{\nu_0+2}+1}^{i_{\nu_0+1}-1} \cdot [\lambda_{\mu_0} - \lambda_x]_l^L = \\ & = \sum_{\mu_0 \leq z \leq \mu_{\nu_0+1}+1} [\lambda_{\mu_{\nu_0+1}+1} - \lambda_x]_{i_{\nu_0+2}+1}^{z-1} \cdot [\lambda_{\mu_0} - \lambda_x]_{z+1}^L = \\ & = (\star) \cdot \frac{[\lambda_{\mu_{\nu_0+1}+1} - \lambda_x]_{\mu_0}^{\mu_{\nu_0+1}+1} - [\lambda_{\mu_0} - \lambda_x]_{\mu_0}^{\mu_{\nu_0+1}+1}}{\lambda_{\mu_{\nu_0+1}+1} - \lambda_{\mu_0+1}} = 0. \end{aligned}$$

**Case B1**  $L = \mu_{\nu_0} + 1 \leq i_{\nu_0-1} - 1$ . In this case  $[\lambda_{\mu_{\nu_0+1}} - \lambda_x]_l^L = 0$ .

**Case B2**  $L = i_{\nu_0-1} - 1 < \mu_{\nu_0} + 1$ . In this case eq. (27) is reduced to

$$\frac{[\lambda_{\mu_{\nu_0}+1} - \lambda_x]_{i_{\nu_0+1}+1}^{i_{\nu_0-1}-1}}{(\lambda_{\mu_{\nu_0}+1} - \lambda_{\mu_0})}$$

This means that we have erased the  $\mu_{\nu_0-1}$  from the product and we have

$$\sum_{i_1 > \dots > i_s} \prod_{\nu=1}^s [\lambda_{\mu_\nu}^\alpha - \lambda_x]_{i_{\nu+1}+1}^{i_\nu-1} = (\star) \sum_{i_1 > \dots > i_s} \prod_{\substack{\nu=1 \\ \nu \neq \nu_0-1}}^s [\lambda_{\mu_\nu}^\alpha - \lambda_x]_{i_{\nu+1}+1}^{i_\nu-1},$$

where  $(\star)$  is a non zero element. This procedure gives us that the original quantity

$$[\lambda_{\mu_{\nu_0}}^\alpha - \lambda_x]_{i_{\nu_0+1}+1}^{i_{\nu_0}-1} \cdot [\lambda_{\mu_{\nu_0-1}}^\alpha - \lambda_x]_{i_{\nu_0+1}}^{i_{\nu_0-1}-1}$$

after summing over  $i_{\nu_0}$  becomes the quantity

$$[\lambda_{\mu_{\nu_0}}^\alpha - \lambda_x]_{i_{\nu_0+1}+1}^{i_{\nu_0-1}-1} = [\lambda_{\bar{\mu}_{\nu_0-1}}^\alpha - \lambda_x]_{\bar{i}_{\nu_0+1}}^{\bar{i}_{\nu_0-1}-1},$$

that is we have eliminated the  $\mu_{\nu_0-1}$  and  $i_{\nu_0}$  from both selections of the sequence of  $\mu$ 's and  $i$ 's, i.e. we have the sequence of length  $s-1$

$$\bar{\mu}_{s-1} = \mu_s < \bar{\mu}_{s-2} = \mu_{s-1} < \dots < \bar{\mu}_{\nu_0-1} = \mu_{\nu_0} < \bar{\mu}_{\nu_0-2} = \mu_{\nu_0-2} < \dots < \bar{\mu}_1 = \mu_1.$$

and the corresponding sequence of equal length

$$\bar{i}_{s-1} = i_s < \bar{i}_{s-2} = i_{s-1} < \dots < \bar{i}_{\nu_0-1} = i_{\nu_0-1} < \bar{i}_{\nu_0} = i_{\nu_0+1} < \dots < \bar{i}_1 = i_1 = d.$$

□

**Remark 31.** One should be careful here since  $\bar{i}_{\nu_0-1} = i_{\nu_0-1} > i_{\nu_0} > \bar{i}_{\nu_0} = i_{\nu_0+1}$ , so  $\bar{i}_{\nu_0-1} > \bar{i}_{\nu_0} + 1$ . This means that the new sequence of  $\bar{i}_{s-1} > \dots > \bar{i}_1$  satisfies a stronger inequality in the  $\nu_0$  position, unless  $\nu_0 - 1 = d$  in the computation of  $\gamma_{d,d}$ .

Consider the set  $s, s-1, \dots, \nu_0$  such that  $m \nmid \mu_\nu$  for  $s \geq \nu \geq \nu_0$  and assume that  $m \mid \mu_{\nu_0-1}$  and  $(23)_{\nu_0}$  and  $(22)_{\nu_0-1}$  hold. We apply lemma 30 and we obtain a new sequence of  $\mu$ 's with  $\mu_{\nu_0-1}$  removed, provided that  $\nu_0 - 1 > 1$ . We continue this way and in the sequence of  $\mu$ 's we eliminate all possible inequalities like (26) obtaining a series of  $\mu$  which involves only inequalities of type (23). But this is not possible if  $\mu \leq d-2$ , according to equation (25). This proves that all  $\gamma_{\mu,d} = 0$  for  $1 \leq \mu \leq d-2$ , this completes the proof of lemma 28. □

**Lemma 32.** *If  $\mu_2 \neq d-1$ , then the contribution of the corresponding summand  $\Gamma_{\bar{\mu},i}$  to  $\gamma_{d,d}$  is zero.*

*Proof.* We are in the case  $\mu = d = i$ . We begin the procedure of eliminating all sequences of inequalities of the form  $(23)_{\nu_0}, (22)_{\nu_0-1}$ , where  $m \mid \nu_0 - 1, m \nmid \nu_0$ , using lemma 30. For  $\nu = 1$  inequality  $(23)_1$  can not hold since it implies the impossible inequality  $d = i_1 < d + 2 - m$ . Therefore,  $(22)_1$  holds, that is  $i_2 > d - m$ . On the other hand we can assume that  $(23)_2$  holds by the elimination process, so we have

$$d - m \stackrel{(22)_1}{<} i_2 \stackrel{(23)_2}{<} \mu_2 + 2.$$

Following the analysis of the proof of lemma 28 we see that the contribution to  $\gamma_{d,d}$  is non zero if case B2 holds, that is ( $\nu_0 = 2$  in this case)  $d-1 = i_{\nu_0-1} - 1 < \mu_2 + 1$ , obtaining that  $\mu_2 = d-1$ . □

**Lemma 33.** *Equation (21) holds, that is*

$$(\lambda_d - \lambda_d^\alpha)\gamma_{d,d} = \sum_{\nu=1}^{d-1} t_{d,\nu}^{(\alpha)} \gamma_{\nu,d} = t_{d,d-1}^{(\alpha)} \gamma_{d-1,d}.$$

*Proof.* We will use the procedure of the proof of lemma 30. We recall that for each fixed sequence of  $\mu_s > \dots > \mu_1$  we summed over all possible sequences  $i_1 > \dots > i_{s+1} = 0$ . In the final step the inequality (26) appears, for  $\nu_0 = 2$ , and  $\mu_{\nu_0} = \mu_2 = d - 1$  and  $\nu_0 - 1 = 1$  and  $\mu_{\nu_0-1} = \mu = d$ , that is:

$$0 = \mu_{\nu_0-1} - m \stackrel{(22)_2}{<} i_{\nu_0} \stackrel{(23)_1}{<} \mu_{\nu_0} + 2 = d + 1.$$

As in the proof of lemma 30 we sum over  $y = i_\nu$  and the result is either zero in case B1 or in the B2 case, where  $\mu_{\nu_0} = \mu_2 = d - 1$  and  $\mu_0 = \mu_{\nu_0-1} - m + 1 = d - m + 1$ , the contribution is computed to be equal to

$$\frac{[\lambda_{\mu_{\nu_0}+1}^\alpha - \lambda_x]_{i_{\nu_0+1}+1}^{i_{\nu_0-1}-1}}{(\lambda_{\mu_{\nu_0}+1} - \lambda_{m_0})} = \frac{[\lambda_d^\alpha - \lambda_x]_{i_3+1}^{d-1}}{\lambda_d - \lambda_d^\alpha}.$$

The last  $\mu_{\nu_0-1} = \mu_1 = d$  is eliminated in the above expression. This means that for a fixed sequence  $\mu_1 > \dots > \mu_s$  the contribution of the inner sum in eq. (24) is given by

$$\frac{1}{\lambda_d - \lambda_d^\alpha} \cdot \sum_{d-1=i_2 > i_3 > \dots > i_s \geq 1} \prod_{\nu=2}^s [\lambda_{\mu_\nu}^\alpha - \lambda_x]_{i_{\nu+1}+1}^{i_\nu-1}.$$

Observe that  $\mu_1 = d$  does not appear in this expression and this expression corresponds to the sequence  $\bar{\mu}_1 = \mu_2 = d - 1 > \bar{\mu}_2 = \mu_3 > \dots > \bar{\mu}_{s-1} = \bar{\mu}_s = 1$ . Notice, also that the problem described in remark 31 does not appear here, since we erased  $i_1$  which is not between some  $i$ 's but the first one. Therefore, we can relate it to a similar expression that contributes to  $\gamma_{d-1,d}$ . Conversely every contribution of  $\gamma_{d-1,d}$  gives rise to a contribution in  $\gamma_{d,d}$ , by multiplying by  $\lambda_d - \lambda_d^\alpha$ . The desired result follows by the expression of  $\gamma_{\mu,d}$  given in eq. (18).  $\square$

We have shown so far how to construct matrices  $\Gamma, T$  so that

$$(28) \quad T^q = 1, \Gamma T \Gamma^{-1} = T^\alpha.$$

We will now prove that  $\Gamma$  has order  $m$ . By equation (28)  $\Gamma^k$  should satisfy equation

$$\Gamma^k T \Gamma^{-k} = T^{\alpha^k}.$$

Using proposition 25 asserting the uniqueness of such  $\Gamma^k$  with  $\alpha$  replaced by  $\alpha^k$  we have that the matrix multiplication of the entries of  $\Gamma$  giving rise to  $(\gamma_{\mu,i}^{(k)}) = \Gamma^k$  coincide to the values by the recursive method of proposition (28) applied for  $\Gamma' = \Gamma^k$ ,  $\alpha' = \alpha^k$  and  $\Gamma' E_1 = \zeta_m^{\epsilon k} E_1$ . In particular for  $k = m$ , we have  $\alpha^m \equiv 1 \pmod{p^\nu}$  for all  $1 \leq \nu \leq h$ , that is the matrix  $\Gamma^k$  should be recursively constructed using proposition (28) for the relation  $\Gamma^m T \Gamma^m = T$ ,  $\Gamma^m E_1 = E_1$ , leading to the conclusion  $\Gamma^m = \text{Id}$ . Notice that the first eigenvalue of  $\Gamma$  is a primitive root of unity, therefore  $\Gamma$  has order exactly  $m$ .

By lemma 10 the action of  $\sigma$  in the special fibre is given by a lower triangular matrix. Therefore, we must have

$$(29) \quad \gamma_{\nu,i} \in \mathfrak{m}_r \text{ for } \nu < i.$$

**Proposition 34.** *If*

$$(30) \quad v(\lambda_i - \lambda_j) > v(a_\nu) \text{ for all } 1 \leq i, j \leq d \text{ and } 1 \leq \nu \leq d - 1$$

*then the matrix  $(\gamma_{\mu,i})$  has entries in the ring  $R$  and is lower triangular modulo  $\mathfrak{m}_R$ .*

*Proof.* Assume that the condition of eq. (30) holds. In equation (18) we compute the fraction

$$(31) \quad \frac{[a]_1^{\mu-1}}{[a]_1^{i-1}} = \begin{cases} \frac{1}{[a]_\mu^{i-\mu}} & \text{if } i > \mu \\ 1 & \text{if } i = \mu \\ [a]_i^{\mu-1} & \text{if } i < \mu \end{cases}$$

The number of  $(\lambda_\mu^\alpha - \lambda_x)$  factors in the numerator is equal to (recall that  $i_{s+1} = 0$ )

$$\sum_{\nu=1}^s (i_\nu - 1 - i_{\nu+1} - 1 + 1) = i - s,$$

and  $i > \mu \geq s$ , so  $i - s > 0$ . Therefore, for the upper part of the matrix  $i > \mu$  we have  $i - s$  factors of the form  $(\lambda_i^\alpha - \lambda_j)$  in the numerator and  $i - \mu$  factors  $a_x$  in the denominator. Their difference is equal to  $(i - s) - (i - \mu) = \mu - s \geq 0$ . By assumption the matrix reduces to an upper triangular matrix modulo  $\mathfrak{m}_R$ .  $\square$

**Remark 35.** The condition given in equation (30) can be satisfied in the following way: It is clear that  $\lambda_i - \lambda_j \in \mathfrak{m}_R$ . Even in the case  $v_{\mathfrak{m}_R}(\lambda_i - \lambda_j) = 1$  we can consider a ramified extension  $R'$  of the ring  $R$  with ramification index  $e$ , in order to make the valuation  $v_{\mathfrak{m}_{R'}}(\lambda_i - \lambda_j) = e$  and then there is space to select  $v_{\mathfrak{m}_{R'}}(a_i) < v_{\mathfrak{m}_{R'}}(\lambda_i - \lambda_j)$ .

**Proposition 36.** *We have that*

$$(32) \quad \gamma_{i,i} \equiv \zeta_m^\epsilon \alpha^{i-1} \pmod{\mathfrak{m}_R}$$

*Let  $A = \{a_1, \dots, a_{d-1}\} \in R$  be the set of elements below the diagonal in eq. (9). If  $a_i \in \mathfrak{m}_R$ , then*

$$\gamma_{\mu,i} \in \mathfrak{m}_R \text{ for } \mu \neq i,$$

*that is  $E_i$  is an eigenvector for the reduced action of  $\Gamma$  modulo  $\mathfrak{m}_R$ . If  $a_{\kappa_1}, \dots, a_{\kappa_r}$  the elements of the set  $A$  which are in  $\mathfrak{m}_R$ , then the reduced matrix of  $\Gamma$  has the form:*

$$\begin{pmatrix} \Gamma_1 & 0 & \cdots & 0 \\ 0 & \Gamma_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & \Gamma_r \end{pmatrix}$$

*where  $\Gamma_1, \Gamma_2, \dots, \Gamma_{r+1}$  for  $1 \leq \nu \leq r + 1$  are  $(\kappa_\nu - \kappa_{\nu-1}) \times (\kappa_\nu - \kappa_{\nu-1})$  lower triangular matrices (we set  $\kappa_0 = 0, \kappa_{r+1} = d$ ).*



*Proof.* Consider the matrix  $\Gamma$ :

$$\left( \begin{array}{ccc|ccc} \boxed{\begin{array}{ccc} \gamma_{11} \\ \vdots \\ \gamma_{\kappa_1,1} \end{array}} & & & & & 0 \\ & & & & & \\ & & & & & \\ \hline & \boxed{\begin{array}{ccc} \gamma_{\mu,i} \\ 1 \leq i \leq \kappa_1 < m \leq d \end{array}} & \boxed{\begin{array}{ccc} \gamma_{\kappa_1+1,\kappa_1+1} \\ \vdots \\ \gamma_{\kappa_2,\kappa_1+1} \end{array}} & \cdots & & \\ & & \boxed{\begin{array}{ccc} \gamma_{\mu,i} \\ \kappa_1 < i \leq \kappa_2 < \mu \leq d \end{array}} & \cdots & & \\ & & & & \boxed{\begin{array}{ccc} \gamma_{\kappa_r+1,\kappa_r+1} \\ \vdots \\ \gamma_{d,\kappa_r+1} \end{array}} & \end{array} \right)$$

We have that  $\mu = i$  and the only element in  $\Sigma_\mu$  which does not have any factor of the form  $(\lambda_y^\alpha - \lambda_x)$  is the sequence

$$1 = \mu_s = \mu_{s-1} - 1 < \mu_{s-1} < \cdots < \mu_2 = \mu_1 - 1 < \mu_1 = \mu$$

For this sequence eq. (18) becomes

$$\gamma_{i,i} = \prod_{\nu=2}^s h_{\alpha-1}(\lambda_{\mu_\nu}, \lambda_{\mu_{\nu-1}}) \zeta_m^\epsilon \text{ mod } \mathfrak{m}_R,$$

which gives the desired result since  $h_{\alpha-1}(\lambda_{\mu_\nu}, \lambda_{\mu_{\nu-1}}) \equiv \binom{\alpha}{1} = \alpha \text{ mod } \mathfrak{m}_R$ .

For proving that all entries  $\gamma_{\mu,i} \in \mathfrak{m}_R$  for  $\kappa_\nu < i \leq \kappa_{\nu+1} < \mu \leq d$ , that is for all entries below the central blocks, we observe that from equation (18) combined with eq. (31) that  $\gamma_{\mu,i}$  is divisible by  $[a]_i^{\mu-1} = a_i a_{i+1} \cdots a_{\kappa_\nu+1} \cdots a_{\mu-1} \in \mathfrak{m}_R$ .  $\square$

Recall that by lemma 2 there is an  $1 \leq a_0 \leq m$  such that  $\alpha = \zeta_m^{a_0}$ .

**Proposition 37.** *The indecomposable module  $V$  modulo  $\mathfrak{m}_R$  breaks into a direct sum of  $r+1$  indecomposable  $k[C_q \rtimes C_m]$  modules  $V_\nu$ ,  $1 \leq \nu \leq r+1$ . Each  $V_\nu$  is isomorphic to  $V_\alpha(\epsilon + a_0 \kappa_{\nu-1}, \kappa_\nu - \kappa_{\nu-1})$ .*

*Proof.* By eq. (32) the first eigenvalue of the reduced matrix block  $\Gamma_\nu$  is

$$\zeta_m^\epsilon \alpha^{\kappa_{\nu-1}} = \zeta_m^{\epsilon + (\kappa_{\nu-1})a_0}.$$

Since that first eigenvalue together with the size of the block determine the last eigenvalue, that is the action of  $C_m$  on the socle the reduced block is uniquely determined up to isomorphism.  $\square$

This way we arrive at a new obstruction. Assume that the indecomposable representation given by the matrix  $T$  as in lemma 16 reduces modulo  $\mathfrak{m}_R$  to a sum of Jordan blocks. Then the  $\sigma$  action on the leading elements of each Jordan block in the special fibre should be described by the corresponding action of  $\sigma$  on the leading eigenvector  $E$  of  $V$ . The corresponding actions on the special fibre should be compatible.

This observation is formally given in proposition 1, which we now prove: Each set  $I_\nu$ ,  $1 \leq \nu \leq t$  corresponds to an indecomposable  $R[G]$ -module, which decomposes to the indecomposables  $V_\alpha(\epsilon_\mu, \kappa_\mu)$ ,  $\nu \in I_\nu$  of the special fiber. Indecomposable

summands have different roots of unity in  $R$ , therefore  $\sum_{\mu \in I_\nu} k_\nu \leq q$ , this is condition (1.a). The second condition (1.b) comes from proposition 13. If 1 is one of the possible eigenvalues of the lift  $T$ , then  $\sum_{\mu \in I_\nu} \kappa_\mu \equiv 1 \pmod{m}$ . If all eigenvalues of the lift  $T$  are different than one, then  $\sum_{\mu \in I_\nu} \kappa_\mu \equiv 0 \pmod{m}$ . If  $\#I_\nu = q$ , then there is one zero eigenvalue and the sum equals 1 mod  $m$ .

It is clear by eq. (32) that condition (1.c) is a necessary condition. On the other hand if (1.c) is satisfied we can write (after a permutation if necessary) the set  $\{1, \dots, S\}$ ,  $S = \sum_{\nu=1}^t \#I_\nu$  as

$$J_1 = \{1, 2, \dots, \kappa_1^{(1)}, \kappa_1^{(1)} + 1, \dots, \kappa_1^{(1)} + \kappa_2^{(1)}, \dots, \sum_{j=1}^{r_1} \kappa_j^{(1)} = b_1\}, I_1 = \{\kappa_1^{(1)}, \dots, \kappa_{r_1}^{(1)}\}$$

$$J_2 = \{b_1 + 1, b_1 + 2, \dots, b_2 = b_1 + \sum_{j=1}^{r_2} \kappa_j^{(2)}\}, I_2 = \{\kappa_1^{(2)}, \dots, \kappa_{r_2}^{(2)}\}$$

.....

$$J_s = \{b_{s-1} + 1, b_{s-1} + 2, \dots, b_t = S\}, I_s = \{\kappa_1^{(s)}, \dots, \kappa_{r_s}^{(s)}\}$$

The matrix given in eq. (9), where

$$a_i = \begin{cases} 0 & \text{if } i \in \{b_1, \dots, b_{s-1}\} \\ \pi & \text{if } i \in \{\kappa_1^{(\nu)}, \kappa_1^{(\nu)} + \kappa_2^{(\nu)}, \kappa_1^{(\nu)} + \kappa_2^{(\nu)} + \kappa_3^{(\nu)}, \dots, \kappa_1^{(\nu)} + \kappa_2^{(\nu)} + \dots + \kappa_{r_\nu-1}^{(\nu)}\} \\ 1 & \text{otherwise} \end{cases}$$

lifts the  $\tau$  generator, and by (12) there is a well defined extended action of the  $\sigma$  as well.

**Example:** Consider the group  $q = 5^2, m = 4, \alpha = 7$ ,

$$G = C_{5^2} \rtimes C_4 = \langle \sigma, \tau \mid \sigma^4 = \tau^{25} = 1, \sigma\tau\sigma^{-1} = \tau^7 \rangle.$$

Observe that  $\text{ord}_5 7 = \text{ord}_{5^2} 7 = 4$ .

- The module  $V(\epsilon, 25)$  is projective and is known to lift in characteristic zero. This fits well with proposition 1, since  $4 \mid 25 - 1 = 4 \cdot 6$ .
- The modules  $V(\epsilon, \kappa)$  do not lift in characteristic zero if  $4 \nmid \kappa$  or  $4 \nmid \kappa - 1$ . Therefore only  $V(\epsilon, 1), V(\epsilon, 4), V(\epsilon, 5), V(\epsilon, 8), V(\epsilon, 9), V(\epsilon, 12), V(\epsilon, 13), V(\epsilon, 16), V(\epsilon, 17), V(\epsilon, 20), V(\epsilon, 21), V(\epsilon, 24), V(\epsilon, 25)$  lift.
- The module  $V(1, 2) \oplus V(3, 2)$  lift to characteristic zero, where the matrix of  $T$  with respect to a basis  $E_1, E_2, E_3, E_4$  is given by

$$T = \begin{pmatrix} \zeta_q & 0 & 0 & 0 \\ 1 & \zeta_q^2 & 0 & 0 \\ 0 & \pi & \zeta_q^3 & 0 \\ 0 & 0 & 1 & \zeta_q^4 \end{pmatrix}$$

and  $\sigma(E_1) = \zeta_q E_1$ .

- The module  $V(1, 2) \oplus V(1, 2)$  does not lift in characteristic zero. There is no way to permute the direct summands so that the eigenvalues of  $\sigma$  are given by  $\zeta_m^\epsilon, \alpha \zeta_m^\epsilon, \alpha^2 \zeta_m^\epsilon, \alpha^3 \zeta_m^\epsilon$ . Notice that  $\alpha = 2 = \zeta_m$ .
- The module  $V(\epsilon_1, 21) \oplus V(2^{21} \cdot \epsilon_1, 23)$  does not lift in characteristic zero. The sum  $21 + 24$  is divisible by 4,  $\epsilon_2 = 2^{21} \epsilon_1$  is compatible, but  $21 + 23 = 44 > 25$  so the representation of  $T$  in the supposed indecomposable module formed

by their sum can not have different eigenvalues which should be 25-th roots of unity.

## REFERENCES

- [1] J. L. Alperin. *Local representation theory*, volume 11 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1986. Modular representations as an introduction to the local representation theory of finite groups.
- [2] Frauke M. Bleher, Ted Chinburg, and Aristides Kontogeorgis. Galois structure of the holomorphic differentials of curves. *J. Number Theory*, 216:1–68, 2020.
- [3] T. Chinburg, R. Guralnick, and D. Harbater. Oort groups and lifting problems. *Compos. Math.*, 144(4):849–866, 2008.
- [4] Ted Chinburg, Robert Guralnick, and David Harbater. The local lifting problem for actions of finite groups on curves. *Ann. Sci. Éc. Norm. Supér. (4)*, 44(4):537–605, 2011.
- [5] Ted Chinburg, Robert Guralnick, and David Harbater. Global Oort groups. *J. Algebra*, 473:374–396, 2017.
- [6] Huy Dang, Soumyadip Das, Kostas Karagiannis, Andrew Obus, and Vaidehee Thatte. Local oort groups and the isolated differential data criterion, 2019.
- [7] A. Heller and I. Reiner. Representations of cyclic groups in rings of integers. I. *Ann. of Math. (2)*, 76:73–92, 1962.
- [8] A. Heller and I. Reiner. Representations of cyclic groups in rings of integers. II. *Ann. of Math. (2)*, 77:318–328, 1963.
- [9] Sotiris Karanikolopoulos and Aristides Kontogeorgis. Representation of cyclic groups in positive characteristic and Weierstrass semigroups. *J. Number Theory*, 133(1):158–175, 2013.
- [10] Aristides Kontogeorgis and Alexios Terezakis. The canonical ideal and the deformation theory of curves with automorphisms, 2021.
- [11] Andrew Obus. The (local) lifting problem for curves. In *Galois-Teichmüller theory and arithmetic geometry*, volume 63 of *Adv. Stud. Pure Math.*, pages 359–412. Math. Soc. Japan, Tokyo, 2012.
- [12] Andrew Obus. A generalization of the Oort conjecture. *Comment. Math. Helv.*, 92(3):551–620, 2017.
- [13] Andrew Obus and Rachel Pries. Wild tame-by-cyclic extensions. *J. Pure Appl. Algebra*, 214(5):565–573, 2010.
- [14] Andrew Obus and Stefan Wewers. Cyclic extensions and the local lifting problem. *Ann. of Math. (2)*, 180(1):233–284, 2014.
- [15] Florian Pop. The Oort conjecture on lifting covers of curves. *Ann. of Math. (2)*, 180(1):285–322, 2014.
- [16] Jean-Pierre Serre. *Linear representations of finite groups*. Springer-Verlag, New York, 1977. Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42.
- [17] Jean-Pierre Serre. *Local fields*. Springer-Verlag, New York, 1979. Translated from the French by Marvin Jay Greenberg.
- [18] Bradley Weaver. The local lifting problem for  $D_4$ . *Israel J. Math.*, 228(2):587–626, 2018.

DEPARTMENT OF MATHEMATICS, NATIONAL AND KAPODISTRIAN UNIVERSITY OF ATHENS PANEPISTIMIOUPOLIS, 15784 ATHENS, GREECE

*Email address:* `kontogar@math.uoa.gr`

DEPARTMENT OF MATHEMATICS, NATIONAL AND KAPODISTRIAN UNIVERSITY OF ATHENS, PANEPIS-TIMIOUPOLIS, 15784 ATHENS, GREECE

*Email address:* `aleksistere@math.uoa.gr`