

Holomorphic tensors on Vaisman manifolds

Liviu Ornea¹, Misha Verbitsky²

Dedicated to Professor Valentin Poenaru at his ninetieth birthday

Abstract

An LCK (locally conformally Kähler) manifold is a complex manifold admitting a Hermitian form ω which satisfies $d\omega = \omega \wedge \theta$, where θ is a closed 1-form, called **the Lee form**. An LCK manifold is called **Vaisman** if the Lee form is parallel with respect to the Levi-Civita connection. The dual vector field, called **the Lee field**, is holomorphic and Killing. We prove that any holomorphic tensor on a Vaisman manifold is invariant with respect to the Lee field. This is used to compute the Kodaira dimension of Vaisman manifolds. We prove that the Kodaira dimension of a Vaisman manifold obtained as a \mathbb{Z} -quotient of an algebraic cone over a projective manifold X is equal to the Kodaira dimension of X . This can be applied to prove the deformational stability of the Kodaira dimension of Vaisman manifolds.

Contents

| | | |
|----------|--|----------|
| 1 | Introduction | 2 |
| 2 | Preliminaries | 3 |
| 2.1 | Locally conformally Kähler manifolds | 3 |
| 2.2 | LCK manifolds with potential | 4 |
| 2.2.1 | Algebraic cones | 5 |
| 2.2.2 | Vaisman manifolds | 6 |
| 3 | Holomorphic tensors on LCK manifolds with potential | 7 |
| 3.1 | Extension of reflexive coherent sheaves | 7 |
| 3.2 | Invariance of holomorphic tensor fields | 8 |

¹Liviu Ornea is partially supported by Romanian Ministry of Education and Research, Program PN-III, Project number PN-III-P4-ID-PCE-2020-0025, Contract 30/04.02.2021

²Misha Verbitsky is partially supported by the HSE University Basic Research Program, FAPERJ E-26/202.912/2018 and CNPq - Process 313608/2017-2.

Keywords: Locally conformally Kähler, LCK potential, algebraic group, Zariski closure
2010 Mathematics Subject Classification: 53C55, 32G05.

| | | |
|----------|---|-----------|
| 4 | Zariski closures and the Chevalley theorem | 9 |
| 5 | Holomorphic tensors on Vaisman manifolds | 11 |
| 6 | An application: the Kodaira dimension of Vaisman manifolds | 12 |

1 Introduction

Locally conformally Kähler geometry is probably the simplest complex geometry beyond the Kähler which admits many examples, explicit and non-explicit. A manifold is called **LCK (locally conformally Kähler)** if it admits a Kähler covering, with the deck group acting by holomorphic homotheties ([Definition 2.1](#)). Every Hopf manifold is LCK, because its universal cover admits a Kähler potential, with the deck transform multiplying the potential by a number. A manifold with this property is called **an LCK manifold with potential**; an LCK manifold admits a potential if and only if it can be realized as a complex submanifold in a Hopf manifold ([\[OV3\]](#)).

Recall that a **diagonal Hopf manifold** is a manifold of form $\frac{\mathbb{C}^n \setminus 0}{\langle A \rangle}$, where A is a linear contraction which can be diagonalized in appropriate basis. A more special class of LCK manifolds with potential are **Vaisman manifolds**, [Subsection 2.2.2](#), which can be characterized as complex manifolds admitting a holomorphic embedding to a diagonal Hopf manifold ([\[OV9, Theorem 4.9\]](#)).

Every Vaisman manifold M is equipped with an action of the Lee and anti-Lee fields, which are Killing and holomorphic ([Proposition 2.17](#)). In [\[Ve2\]](#), it was shown that any stable holomorphic bundle on M is equivariant with respect to this action. From [Theorem 2.18](#), it follows that any complex subvariety is invariant under this action as well.

Let G be the group of holomorphic isometries of a Vaisman manifold M obtained as the closure of the group generated by the Lee and the anti-Lee flow. In this paper, we prove that any holomorphic tensor field on M is G -invariant.

For holomorphic vector fields and differential forms this result was obtained by K. Tsukada in [\[T1, Theorem 3.3, Theorem 4.2\]](#).

We prove a similar result for LCK manifolds with potential. Let M be an LCK manifold with potential, and \tilde{M} its Kähler \mathbb{Z} -cover. We assume that \tilde{M} is an open algebraic cone ([Definition 2.10](#)); this is true automatically when $\dim_{\mathbb{C}} M \geq 3$ ([\[OV8\]](#)). When $\dim_{\mathbb{C}} M = 2$, this would follow if we assume the GSS conjecture ([\[OV7, Section 25.6\]](#)). Since \tilde{M} is algebraic, it makes

sense to speak of the Zariski closure of the \mathbb{Z} -action on \tilde{M} . We consider the Zariski closure \mathcal{G} of the \mathbb{Z} -action as the smallest algebraic subgroup of $\text{Aut}(\tilde{M})$ containing this \mathbb{Z} -action; this group is clearly commutative and positive-dimensional. Since the action of \mathcal{G} commutes with the action of the deck transform group $\mathbb{Z} \subset \mathcal{G}$ on \tilde{M} , the quotient \mathcal{G}/\mathbb{Z} naturally acts on M .

We prove that any holomorphic tensor field on an LCK manifold with potential is \mathcal{G} -invariant ([Proposition 3.3](#)).

Let $H = \frac{\mathbb{C}^n \setminus 0}{\langle A \rangle}$ be a diagonal Hopf manifold, and \mathcal{G} the Zariski closure of $\langle A \rangle$. Let α_i be the eigenvalues of A , and A_1 an operator which has the eigenvalues $|\alpha_i|$ in the same basis. In [Theorem 4.3](#), we show that the linear field $\log A_1$ belongs to the Zariski closure of $\langle A \rangle$, and in [Theorem 5.1](#), step 3, we prove that $\log A_1$ can be obtained as the Lee field of a Vaisman structure on H .

This implies that the Lee and anti-Lee field of a Vaisman manifold belong to the Lie algebra of \mathcal{G} (the Zariski closure of the \mathbb{Z} -action). Therefore, any holomorphic tensor field is Lee and anti-Lee invariant.

2 Preliminaries

For the material in this section, we refer the reader to [\[OV7\]](#).

2.1 Locally conformally Kähler manifolds

Definition 2.1: A complex manifold (M, I) is called **locally conformally Kähler** (LCK, for short) if it admits a covering (\tilde{M}, I) equipped with a Kähler metric $\tilde{\omega}$ such that the deck group of the cover acts on $(\tilde{M}, \tilde{\omega})$ by holomorphic homotheties. An **LCK metric** on an LCK manifold is an Hermitian metric on (M, I) such that its pullback to \tilde{M} is conformal with $\tilde{\omega}$.

Proposition 2.2: A Hermitian manifold (M, I, g) is LCK if and only if there exists a closed 1-form θ (called **the Lee form**) such that the fundamental form $\omega(x, y) := g(Ix, y)$ satisfies $d\omega = \theta \wedge \omega$.

Definition 2.3: An LCK structure (ω, θ) on a complex manifold (M, I) is called **strict** if the Lee form is not exact. When the Lee form is exact, the structure (ω, θ) is called **globally conformally Kähler** (GCK).

Example 2.4: Let $H := (\mathbb{C}^n \setminus 0)/\langle A \rangle$, $A = \lambda \text{Id}$, $|\lambda| > 1$, be a **classical**

Hopf manifold. It is LCK because the flat Kähler metric $\tilde{g}_0 = \sum dz_i \otimes d\bar{z}_i$ is multiplied by 2 by the deck group \mathbb{Z} . The Lee form on the covering $\mathbb{C}^n \setminus 0$ is $\theta = -d \log |z|^2$. Since the Hopf manifold is diffeomorphic with $S^1 \times S^{2n-1}$, the LCK structure is strict.

Remark 2.5: All complex submanifolds of an LCK manifold are still LCK (but not necessarily strict).

2.2 LCK manifolds with potential

Definition 2.6: An LCK manifold has **LCK potential** if it admits a Kähler covering on which the Kähler metric has a global and positive potential function ψ such that the deck group multiplies ψ by a constant.¹ In this case, M is called **LCK manifold with potential**.

Example 2.7: The diagonal Hopf manifold in [Example 2.4](#) is LCK with potential because \tilde{g}_0 has the global automorphic potential $\psi := \sum |z_i|^2$.

By [\[OV3\]](#), compact LCK manifolds with potential are stable to small deformations of the complex structure. This implies that all **linear Hopf manifolds** $(\mathbb{C}^n \setminus 0)/\langle A \rangle$, $A \in \text{GL}(n, \mathbb{C})$, with eigenvalues of absolute value > 1 , are LCK with potential.

Remark 2.8: One can easily prove that all complex submanifolds of an LCK manifold with potential are still LCK with potential.

One of the most important properties of LCK manifolds with potential is the following Kodaira-type embedding result:

Theorem 2.9: ([\[OV3\]](#)) Let M be a compact LCK manifold with potential of complex dimension at least 3. Then M admits a holomorphic embedding into a linear Hopf manifold. ■

The idea of the proof is that the Kähler cover of an LCK manifold with potential is not a cone in the Riemannian sense (it lacks the homothety), but is very close to the idea of cone, because it can be completed with just one (singular) point. This completion can be realized using a theorem by Rossi and Andreotti-Siu which only works in dimension greater than 3. In dimension 2, the result is true, if one assumes the GSS conjecture ([\[OV6\]](#),

¹In the sequel, a differential form which is multiplied by a constant factor by the action of the deck group is called **automorphic**.

Theorem 5.6]), which is a long-standing conjecture, often assumed to be true.

Below, we give a precise description of the Kähler cover of an LCK manifold with potential.

2.2.1 Algebraic cones

Definition 2.10: ([OV4]) A **closed algebraic cone** is an affine variety \mathcal{C} admitting a \mathbb{C}^* -action ρ with a unique fixed point x_0 , called **the origin** or **the apex**, and satisfying the following:

- (i) \mathcal{C} is smooth outside of x_0 , and \mathbb{C}^* acts freely on $\mathcal{C} \setminus x_0$.
- (ii) ρ acts on the Zariski tangent space $T_{x_0}\mathcal{C}$ with all eigenvalues $|\alpha_i| < 1$.

An **open algebraic cone** is a complex manifold which is biholomorphic to a closed algebraic cone without the origin.

Theorem 2.11: ([OV8]) Let M be an LCK manifold with proper potential, and \tilde{M} its Kähler \mathbb{Z} -cover. Then \tilde{M} is an open algebraic cone. ■

Remark 2.12: The restriction to proper potentials is not significant, since any LCK metric with potential can be approximated in the C^∞ -topology (while keeping the complex structure fixed) by an LCK metric with proper potential ([OV2, OV5]).

As a topological space, the closed algebraic cone \tilde{M}_c is obtained as a one-point completion of an open algebraic cone \tilde{M} . However, the complex structure on a closed algebraic cone is not uniquely determined by \tilde{M} . We refer the reader to [OV8] for this intricate and fascinating subject.

The manifold \tilde{M} does not define the LCK manifold M uniquely: $M = \tilde{M}/\mathbb{Z}$ is determined by the choice of the \mathbb{Z} -action which may vary.

In [OV8], we proved that an algebraic cone of an LCK manifold with potential is isomorphic to an affine cone over a projective orbifold. This nearly transforms the geometry of LCK manifolds with potential into a chapter of algebraic geometry.

We end these preliminaries by citing a result which gives a method of constructing LCK manifolds with potential once the algebraic cone which will be the covering is known. We first need the following notion.

Definition 2.13: Let \tilde{M} be an open algebraic cone, \tilde{M}_c the corresponding closed cone, and $\tilde{r} \in T\tilde{M}_c$ a holomorphic vector field such that for all $t > 0$ the diffeomorphism $e^{t\tilde{r}}$ is a holomorphic contraction of \tilde{M}_c to the origin. In this situation, the 1-parameter family $e^{t\tilde{r}}$ is called **the flow of contractions** (or **contraction flow**). A strictly pseudoconvex hypersurface $S \subset \tilde{M}$ is called a **pseudoconvex shell** if S intersects each orbit of $e^{t\tilde{r}}$, $t \in \mathbb{R}$, exactly once.

Pseudoconvex shells are used to obtain automorphic plurisubharmonic functions, as follows.

Theorem 2.14: ([OV4]) Let \tilde{M} be an algebraic cone, $e^{t\tilde{r}}$ a contraction flow, and $S \subset \tilde{M}$ a pseudoconvex shell. Then for each $\lambda \in \mathbb{R}$ there exists a unique function φ_λ such that $\text{Lie}_{\tilde{r}}\varphi = \lambda\varphi$ and $\varphi_\lambda|_S = 1$. Moreover, such φ_λ is strictly plurisubharmonic when $\lambda \gg 0$. ■

2.2.2 Vaisman manifolds

Among the LCK manifolds with potential, the best understood subclass is formed by the **Vaisman manifolds**. These are LCK manifolds with the Lee form parallel with respect to the Levi-Civita connection of the LCK metric.

Let (M, I, ω, θ) be a Vaisman manifold and $\pi : \tilde{M} \rightarrow M$ a Kähler cover on which $\pi^*\theta$ is exact. Then it can be seen that the $\tilde{\omega}$ -squared norm of $\pi^*\theta$ is a positive, automorphic potential for the Kähler metric $\tilde{\omega}$ ([OV3]).

Example 2.15: Some examples of Vaisman manifolds:

- (i) Diagonal Hopf manifolds $(\mathbb{C}^n \setminus 0)/\langle A \rangle$ where A is semi-simple and with eigenvalues α_i of absolute value > 1 , [GO, OV4].
- (ii) Elliptic complex surfaces (see [B] for the complete classification of Vaisman compact surfaces; see also [VVO]).
- (iii) All compact submanifolds of a Vaisman manifold ([Ve1]).

Remark 2.16: The class of Vaisman manifolds is strict: neither the LCK Inoue surfaces, nor the non-diagonal Hopf manifolds can bear Vaisman metrics ([B], [OV4]).

The metric dual θ^\sharp of the Lee form is called **the Lee field**.

Proposition 2.17: ([Vai1, Vai2]) Let (M, g, I, θ) be a Vaisman manifold. Then the Lee field θ^\sharp is Killing and holomorphic; moreover, it commutes with $I\theta^\sharp$. ■

Denote by Σ the holomorphic 1-dimensional foliation generated by θ^\sharp and $I\theta^\sharp$. It is called **the canonical foliation** (the motivation is given in the next theorem).

Theorem 2.18: Let M be a compact Vaisman manifold, and $\Sigma \subset TM$ its canonical foliation. Then:

- (i) Σ is independent from the choice of the Vaisman metric ([T3]).
- (ii) $d^c\theta = \omega - \theta \wedge I\theta$ ([Vai2]) and the exact (1,1)-form $\omega_0 := d^c\theta$ is semi-positive ([Ve1]). Therefore, $\Sigma = \ker \omega_0$, and ω_0 is transversally Kähler with respect to Σ .

The following result is almost obvious. We give its proof for consistency.

Lemma 2.19: Let M be a compact Vaisman manifold, and G the closure of the Lie group generated by the action of the Lee and the anti-Lee fields on M . Then G is a compact torus.

Proof: Since the Lee and anti-Lee fields are acting on M by isometries (Proposition 2.17), the group G is a closed subgroup of the isometry group of M , and the latter is compact. Then G is compact; since it is the closure of an abelian group, it is abelian. ■

3 Holomorphic tensors on LCK manifolds with potential

The purpose of this section is to prove Proposition 3.3. For its proof, we shall recall some facts related to coherent sheaves.

3.1 Extension of reflexive coherent sheaves

Given a coherent sheaf \mathcal{F} over X , let $\mathcal{F}^* := \text{Hom}(\mathcal{F}, \mathcal{O}_X)$ be **the dual sheaf**, that is, the sheaf of module homomorphisms to the ring of regular

functions. The natural morphism of sheaves $\mathcal{F} \rightarrow \mathcal{F}^{**}$ does not need to be an isomorphism: for example, its kernel contains the torsion of \mathcal{F} . A coherent sheaf is called **reflexive** if the natural map $\mathcal{F} \rightarrow \mathcal{F}^{**}$ is an isomorphism. For an introduction to the reflexive sheaves, see [OSS]. Here we state some results which are relevant to our work.

First of all, notice that the sheaf \mathcal{F}^* is already reflexive ([OSS], Ch. II, Lemma 1.1.8). Moreover, the singular set of a reflexive sheaf over a normal variety has codimension ≥ 3 ; in particular, a reflexive sheaf over a complex surface is locally free ([OSS], Ch. II, 1.1.10). Also, a reflexive sheaf of rank one over a smooth manifold is locally free ([OSS], Ch. II, Lemma 1.1.15).

The most important property of reflexive sheaves is **normality**. Let $Z \subset X$ be a subvariety of a normal complex variety, $\text{codim}_X Z \geq 2$. Consider the open embedding map $j : X \setminus Z \hookrightarrow X$, and let j^* and j_* be the sheaf pullback and pushforward, For any sheaf \mathcal{F} , there exists a natural sheaf morphism $\mathcal{F} \rightarrow j_* j^* \mathcal{F}$ taking a section of \mathcal{F} to its restriction to $X \setminus Z$. A coherent sheaf \mathcal{F} is **normal** if the natural map $\mathcal{F} \rightarrow j_* j^* \mathcal{F}$ is an isomorphism, for any subvariety $Z \subset X$ of codimension ≥ 2 .

The following theorem can be understood as a sheaf version of the Hartogs extension theorem.

Theorem 3.1: A coherent sheaf \mathcal{F} over a normal complex variety is normal if and only if it is reflexive.

Proof: [Se, Proposition 7]. ■

Theorem 3.2: ([OV7, Theorem 27.6]) Let M be a normal complex variety with isolated singularities, $\dim_{\mathbb{C}} M \geq 3$, $x \in M$ a point, and $M_0 := M \setminus \{x\} \xrightarrow{j} M$. Consider a reflexive sheaf \mathcal{F} on M_0 . Then $j_* \mathcal{F}$ is a reflexive coherent sheaf. Moreover, $j_* \mathcal{F}$ is the unique coherent reflexive sheaf on M which is isomorphic to \mathcal{F} on M_0 .

Proof: Uniqueness of a reflexive extension follows from the fact that a coherent sheaf \mathcal{F} over a normal variety is normal if and only if it is reflexive (Theorem 3.1). The existence of coherent extension of \mathcal{F} follows from [AS, Proposition 6.1]. ■

3.2 Invariance of holomorphic tensor fields

We are now ready to prove the main result of this section.

Proposition 3.3: Let M be a compact LCK manifold with potential, \tilde{M} its Kähler \mathbb{Z} -cover, considered as an open algebraic cone, and $\Phi \in H^0(M, B)$ be a holomorphic tensor field on M , where $B = (\Omega^1 M)^{\otimes k} \otimes TM^{\otimes l}$. Denote by $\tilde{\Phi}$ its lift to \tilde{M} , and let \mathcal{G} be the Zariski closure of the \mathbb{Z} -action on \tilde{M} . Then $\tilde{\Phi}$ is \mathcal{G} -invariant.

Proof. Step 1: Consider a \mathbb{Z} -action on a finite-dimensional space W . Then any \mathbb{Z} -invariant vector $w \in W$ is invariant under the Zariski closure of \mathbb{Z} . This is why the statement of [Proposition 3.3](#) is not surprising. However, the space of tensor fields on \tilde{M} is not finite-dimensional, which makes [Proposition 3.3](#) non-trivial. Denote by \mathfrak{m} the maximal ideal of the origin in the closed algebraic cone \tilde{M}_c . We think of the quotients $\mathcal{O}_{\tilde{M}_c}/\mathfrak{m}^k$ as of the spaces of jets of holomorphic (or algebraic) functions. As in [\[OV8\]](#), we choose \tilde{M}_c normal, so that the \mathbb{Z} -action is extended to \tilde{M}_c . Then any \mathbb{Z} -invariant jet $u \in \mathcal{O}_{\tilde{M}_c}/\mathfrak{m}^k$ is also \mathcal{G} -invariant.

Step 2: From now on, we use the same letter Φ to denote the lift of Φ to \tilde{M} . Consider Φ as a \mathbb{Z} -invariant section of the appropriate tensor bundle $B = (\Omega^1 M)^{\otimes k} \otimes TM^{\otimes l}$, which is by construction \mathbb{Z} -equivariant. Using [Theorem 3.2](#), we extend B to a reflexive coherent sheaf B_c on \tilde{M}_c . Since coherent reflexive sheaves are normal ([\[OSS, Ch. II, Lemma 1.1.12\]](#), [\[Se, Proposition 7\]](#)), the section Φ admits a holomorphic extension to \tilde{M}_c , denoted as $\Phi_c \in H^0(\tilde{M}_c, B_c)$. Consider the space $J_B^k := \frac{B_c}{\mathfrak{m}^{k+1} B_c}$ of k -jets of B_c . The \mathbb{Z} -action on $H^0(\tilde{M}_c, B_c)$ preserves $H^0(\tilde{M}_c, \mathfrak{m}^k B_c) \subset H^0(\tilde{M}_c, B_c)$. Let $\Phi_c^k \in J_B^k$ be the k -jet of Φ_c . Since Φ_c is \mathbb{Z} -invariant, and the space of k -jets is finite-dimensional, the k -jet Φ_c^k is \mathcal{G} -invariant. Let $\hat{B}_c := \varprojlim J_B^k$ be the \mathfrak{m} -adic completion of B_c , and $\hat{\Phi}_c$ the image of Φ_c in this completion. Since all k -jets Φ_c^k are \mathcal{G} -invariant, the completion $\hat{\Phi}_c$ is also \mathcal{G} -invariant. Since the ring of germs of holomorphic functions on \tilde{M}_c in c is Noetherian ([\[GN, Chapter II, Theorem B.9\]](#)), and B_c , being reflexive, is torsion-free, the completion map $H^0(\tilde{M}_c, B_c) \rightarrow \hat{B}_c$ is injective ([\[AM, Theorem 10.17\]](#)). Therefore, $\Phi \in H^0(\tilde{M}_c, B_c)$ is also \mathcal{G} -invariant. ■

4 Zariski closures and the Chevalley theorem

We recall the following famous theorem due to Chevalley.

Let $G \subset \mathrm{GL}(V)$ be an algebraic group, and $W = V^{\otimes k} \otimes (V^*)^{\otimes l}$ a tensor representation of G . A G -invariant vector $v \in W$ is called a **tensor invariant of G** . A point $x \in \mathbb{P}W$ is called a **projective tensor invariant of G** if it is G -invariant.

Theorem 4.1: (Chevalley theorem, [M]).

An algebraic group is uniquely determined by the set of its projective tensor invariants. ■

When an algebraic group is *reductive* (over \mathbb{C} , a group is reductive if and only if it has a compact real form), a stronger version of Chevalley theorem is available.

Theorem 4.2: A reductive algebraic group is uniquely determined by the set of its tensor invariants.

Proof: [D, Proposition 3.1 (c)]. ■

We are going to prove the following result, used further on to determine the Lee field action on a Vaisman manifold.

Theorem 4.3: Let $A \in \mathrm{GL}(n, \mathbb{C})$ be a diagonal linear operator with the eigenvalues $\alpha_1, \dots, \alpha_n$, and A_1 an operator which is diagonal in the same basis, with the eigenvalues $|\alpha_1|, \dots, |\alpha_n|$. Denote by \mathcal{G} the Zariski closure of the group $\langle A \rangle$ in $\mathrm{GL}(n, \mathbb{C})$. Then \mathcal{G} contains A_1 .

Proof: Since the operator A is diagonalizable, the algebraic closure of $\langle A \rangle$ is a commutative group which contains only semisimple elements. By [BT, Proposition 1.5], the connected component of \mathcal{G} is isomorphic to $(\mathbb{C}^*)^d$, hence \mathcal{G} is reductive. Then Theorem 4.2 implies that \mathcal{G} is the set of all $g \in \mathrm{GL}(n, \mathbb{C})$ which preserve all A -invariant vectors $w \in W = V^{\otimes k} \otimes (V^*)^{\otimes l}$. The eigenvalues of A on W are products of k instances of α_i and l instances of α_i^{-1} . The eigenvectors $w \in W$ are $w = \bigotimes_{i=1}^k z_{m_i} \otimes \bigotimes_{i=1}^l \zeta_{n_i}$, where z_{n_i} are the eigenvectors in V with the eigenvalues α_{n_i} and ζ_{m_i} are the eigenvectors in V^* with the eigenvalues $\alpha_{m_i}^{-1}$. Then $A(w) = \prod_{i=1}^k \alpha_{n_i} \prod_{i=1}^l \alpha_{m_i}^{-1} w$. The eigenvector w is A -invariant if and only if $\prod_{i=1}^k \alpha_{n_i} \prod_{i=1}^l \alpha_{m_i}^{-1} = 1$. This implies $\prod_{i=1}^k |\alpha_{n_i}| \prod_{i=1}^l |\alpha_{m_i}^{-1}| = 1$, hence w is A_1 -invariant as well. ■

5 Holomorphic tensors on Vaisman manifolds

Let M be a Vaisman manifold, and θ^\sharp its Lee field. Then θ^\sharp and $I(\theta^\sharp)$ are holomorphic Killing fields ([Proposition 2.17](#)). The closure G of the group $e^{t_1\theta^\sharp+t_2I(\theta^\sharp)}$ in the group of isometries of M is compact and commutative, hence it is a compact torus ([Lemma 2.19](#)).

Theorem 5.1: Let M be a compact Vaisman manifold, $\Phi \in H^0(M, B)$ a holomorphic tensor, where $B = (\Omega^1 M)^{\otimes k} \otimes TM^{\otimes l}$, and G the smallest closed Lie group containing the flows generated by the Lee and the anti-Lee fields. Then Φ is G -invariant.

Proof. Step 1: Choose a Vaisman metric on M with LCK rank 1, and let \tilde{M} be the corresponding Kähler \mathbb{Z} -cover of M , which is considered as an open algebraic cone. Let $\mathcal{G} \subset \text{Aut}(\tilde{M})$ be the Zariski closure of the \mathbb{Z} -action. Then Φ is \mathcal{G} -invariant by [Proposition 3.3](#). To finish the proof of [Theorem 5.1](#), it would suffice to show that $G \subset \mathcal{G}$; since \mathcal{G} is closed, this would follow if we prove that the Lee field θ^\sharp is tangent to \mathcal{G} .

Step 2: Let $M \rightarrow H$ be a holomorphic embedding of M to a diagonal Hopf manifold. By [Theorem 2.18](#), this embedding commutes with the group generated by the Lee and the anti-Lee flow, hence it would suffice to show that θ^\sharp is tangent to \mathcal{G} when $M = H$. It remains to prove [Theorem 5.1](#) assuming that M is a Hopf manifold. For this purpose, we compute the groups G and \mathcal{G} for a diagonal Hopf manifold. This can be done explicitly using the expression for the Vaisman metric on a Hopf manifold obtained in [Theorem 2.14](#) via the pseudoconvex shells.

Step 3: The rank 2 algebra generated by the Lee and anti-Lee flows on a Vaisman manifold M is uniquely determined by the complex structure on M ([Theorem 2.18](#)). We express the Lee flow on a Hopf manifold using a Vaisman structure we shall construct explicitly, and prove that it belongs to $\text{Lie}(\mathcal{G})$. This would imply that the original Lee flow also belongs to $\text{Lie}(\mathcal{G})$.

Let $H = \frac{\mathbb{C}^n \setminus 0}{\langle A \rangle}$, where $A \in \text{GL}(n, \mathbb{C})$ is a diagonal linear contraction with eigenvalues $\alpha_1, \dots, \alpha_n$. Let A_1 be the matrix which is diagonal in the same basis as A , and has eigenvalues $|\alpha_i|$. Then $\vec{r} := \log A_1$ is a holomorphic contraction vector field, and $e^{t\vec{r}}$ is an A -invariant contraction flow. The sphere S^{2n-1} is a pseudoconvex shell compatible with this contraction flow. By [\[OV4\]](#), we obtain a $e^{t \log A_1}$ -automorphic plurisubharmonic function φ . We are going to prove that φ is A -automorphic.

Consider the diagonal $U(1)^n$ -action on \mathbb{C}^n . The plurisubharmonic function φ is $U(1)^n$ -invariant because \vec{r} and S^{2n-1} are $U(1)^n$ -invariant. Since $A \in U(1)^n \cdot A_1$, and φ is A_1 -automorphic, it is A -automorphic. Therefore, $\frac{dd^c \varphi}{\varphi}$ defines a Vaisman metric on H . By definition, the Lee flow maps the level sets of φ to the level sets of φ , with $\varphi(e^{t\theta^\sharp} z) = t\varphi(z)$. By construction of φ , the same is true for $e^{-t \log A_1}$, that is, $\varphi(e^{-t \log A_1} z) = t\varphi(z)$. Therefore, $\theta^\sharp = -\log A_1$, where $\log A_1$ is the diagonal matrix with the eigenvalues $\log |\alpha_i|$ in the same basis. This is the Lee flow for the Vaisman metric $\frac{dd^c \varphi}{\varphi}$.

The vector field $\log A_1$ belongs to $\text{Lie}(\mathcal{G})$, because any power of A_1 fixes all tensor invariants of \mathcal{G} by [Theorem 4.3](#), hence the corresponding 1-parametric subgroup also belongs to $\text{Lie}(\mathcal{G})$; this gives $\log A_1 \in \text{Lie}(\mathcal{G})$. ■

6 An application: the Kodaira dimension of Vaisman manifolds

Definition 6.1: Let M be a compact complex manifold. Its **pluricanonical bundle** is the tensor power $K_M^n = K_M^{\otimes n}$, $n \geq 0$. The **Kodaira dimension** $\kappa(M)$ is defined as $\kappa(M) := \limsup_n \frac{\log(\dim H^0(K_M^n))}{\log n}$.

Remark 6.2: The Kodaira dimension is equal to $-\infty$ if $H^0(K_M^n) = 0$ for almost all $n > 0$, and to 0 if $\dim H^0(K_M^n)$ is bounded, but non-zero for infinitely many n . If the function $n \mapsto \dim H^0(K_M^n)$ grows as a polynomial of degree d , the Kodaira dimension of M is d .

Let M be a Vaisman manifold, G the closure of the group generated by the Lee and the anti-Lee action, \tilde{M} a Kähler \mathbb{Z} -covering of M , and \tilde{G} the lift of G to \tilde{M} . Denote by \mathfrak{G} the Zariski closure of \tilde{G} in $\text{Aut}(\tilde{M})$, considered as an algebraic variety ([\[OV8\]](#)).

Since \tilde{G} acts on \tilde{M} by homotheties, and contains contractions, there is an open set $U \subset \mathfrak{G}$ in the connected component of \mathfrak{G} such that for all $\gamma' \in U$, the quotient $M' := \frac{\tilde{M}}{\langle \gamma' \rangle}$ is LCK ([\[OV8\]](#)). The automorphism γ commutes with the action of the group generated by the Lee and the anti-Lee field, hence by [\[KO\]](#) M' is Vaisman.¹ By [\[BT, Proposition 1.5\]](#), the connected component of \mathfrak{G} is isomorphic to $(\mathbb{C}^*)^k$. It is not hard to see that the union of all one-dimensional closed subgroups is dense in any algebraic group, hence U contains a dense subset of γ' which are contained in a subgroup

¹As shown in [\[KO\]](#), an LCK manifold which admits a holomorphic conformal \mathbb{C} -action is Vaisman, assuming that this action cannot be lifted to an isometry of the Kähler cover.

$\mathbb{C}^* \subset \mathfrak{G}$. Since $\mathbb{C}^*/\langle\gamma'\rangle$ is an elliptic curve, the Vaisman manifold $M' = \frac{\tilde{M}}{\langle\gamma'\rangle}$ is elliptically fibered.² As shown in [OV1], the leaf space X is a projective orbifold.

Theorem 6.3: Let M be a Vaisman manifold, M' its quasi-regular deformation obtained above, and X the corresponding leaf space, which is a projective orbifold. Then $\kappa(M) = \kappa(M') = \kappa(X)$, where κ denotes the Kodaira dimension.

Proof. Step 1: The equality $\kappa(M') = \kappa(X)$ is clear from the adjunction formula: since $\pi : M' \rightarrow X$ is an elliptic fibration, one has $H^0(K_{M'}^k) = H^0(K_X^k)$ for all k .

Step 2: By Theorem 5.1, any section h of the pluricanonical bundle K_M^k on M is \mathfrak{G} -invariant. Then, for any $\gamma' \in U$, the lift of h to \tilde{M} is γ' -invariant, hence can be obtained as a pullback of a section of $K_{M'}^k$ under the quotient map $\tilde{M} \rightarrow M'$. This defines an injective map $\Psi : H^0(K_M^k) \rightarrow H^0(K_{M'}^k)$.

Step 3: To finish the proof of Theorem 6.3 it remains to show that $\Psi : H^0(K_M^k) \rightarrow H^0(K_{M'}^k)$ is surjective. This would follow if we show that any section $h' \in H^0(K_{M'}^k)$ is \mathfrak{G} -invariant. Since $H^0(K_{M'}^k) = H^0(K_X^k)$ (Step 1), the \mathfrak{G} -invariance of h' would follow if we prove that the corresponding section of K_X^k is invariant under the natural action of \mathfrak{G} on X .

Recall that **the pluricanonical representation** is the natural action of $\text{Aut}(X)$ on $H^0(K_X^k)$. Since X is an orbifold, its singularities are klt ([P, Proposition 1.2.1]). By [KK, Theorem 10.61], the image of the pluricanonical representation is finite; since \mathfrak{G} is connected, it acts on $H^0(K_X^k)$ trivially. This gives $H^0(K_M^k) = H^0(K_{M'}^k) = H^0(K_X^k)$. ■

As an application, we outline the proof of the deformational stability of Kodaira dimension for Vaisman manifolds.

Conjecture 6.4: Let $M_t, t \in \mathbb{R}$, be a smooth family of compact Vaisman manifolds. Then the Kodaira dimension of M_t is constant.

Sketch of a proof: We sketch a proof of this conjecture, reducing it to Conjecture 6.5 below.

²Elliptically fibered Vaisman manifolds are called **quasi-regular**.

Let \tilde{M}_t be the family of open algebraic cones obtained from each M_t by taking a Kähler \mathbb{Z} -cover; this is possible to do by [OV1]. Each \tilde{M}_t is equipped with a natural algebraic structure by [OV8].

What follows is a conjecture, which most likely can be proven by the methods used in [OV8].

Let $\tilde{M}_t, t \in \mathbb{R}$, be a smooth family of open algebraic cones. As shown in [OV8], for an appropriate choice of the \mathbb{C}^* -action on \tilde{M}_t the quotient \tilde{M}_t/\mathbb{C}^* is isomorphic to a complex projective orbifold X_t ; this isomorphism is extended to an isomorphism between \tilde{M}_t and $\text{Tot}^\circ(L_t)$, where L_t is an ample line bundle over X_t .

Conjecture 6.5: Let $\tilde{M}_t, t \in \mathbb{R}$, be a smooth family of open algebraic cones. Then for any sufficiently small open subset $U \subset \mathbb{R}$, the choice of \mathbb{C}^* -action as above can be performed in such a way that the family (X_t, L_t) depends smoothly on $t \in U$.

Now, by Theorem 6.3, the Kodaira dimension of M_t is equal to the Kodaira dimension of X_t , which is constant, as shown by Siu in [Si]. ■

Acknowledgements: We are grateful to Nicolina Istrati for her interest and valuable advice.

References

- [AS] A. Andreotti, Y. T. Siu, *Projective embeddings of pseudoconcave spaces*, Ann. Scuola Norm. Sup. Pisa **24**, 231-278 (1970). (Cited on page 8.)
- [AM] M. F. Atiyah, I. G. MacDonald, *Introduction to commutative algebra*, Addison-Wesley, 1969. (Cited on page 9.)
- [B] F. A. Belgun, *On the metric structure of non-Kähler complex surfaces*, Math. Ann. **317** (2000), 1-40. (Cited on page 6.)
- [BT] A. Borel, J. Tits, *Groupes réductives*, Publications Mathématiques de l’IHÉS, **27** (1965), 55-151. (Cited on pages 10 and 12.)
- [D] P. Deligne, *Hodge cycles on abelian varieties* (Notes by J. Milne.) In: P. Deligne, et al., *Hodge cycles, motives, and Shimura varieties*. Lecture Notes in Mathematics **900**. Springer-Verlag, Berlin-New York, 1982. (Cited on page 10.)
- [GO] P. Gauduchon, L. Ornea, *Locally conformally Kähler metrics on Hopf surfaces*, Ann. Inst. Fourier (Grenoble) **48** (1998), 1107-1128. (Cited on page 6.)

- [GN] R. C. Gunning, H. Rossi, *Analytic functions of several complex variables*, Reprint of the 1965 original. AMS Chelsea Publishing, Providence, RI, 2009. (Cited on page 9.)
- [KO] Y. Kamishima, L. Ornea, *Geometric flow on compact locally conformally Kähler manifolds*, Tohoku Math. J., **57** (2) (2005), 201-221. (Cited on page 12.)
- [KK] Kollár, J., with collaboration of Kovács, S., *Singularities of the Minimal Model Program*, 2013, Cambridge University Press. (Cited on page 13.)
- [M] D. W. Morris, *Ratner's Theorems on Unipotent Flows*, Univ. of Chicago Press, Chicago, 2005. (Cited on page 10.)
- [OSS] C. Okonek, M. Schneider, H. Spindler, *Vector bundles on complex projective spaces*, Corrected reprint of the 1980 edition. With an appendix by S. I. Gelfand. Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel, 2011. (Cited on pages 8 and 9.)
- [OV1] L. Ornea, M. Verbitsky, *An immersion theorem for compact Vaisman manifolds*, Math. Ann. **332** (2005), 121-143. (Cited on pages 13 and 14.)
- [OV2] L. Ornea, M. Verbitsky, *Morse-Novikov cohomology of locally conformally Kähler manifolds*, J. Geom. Phys. **59**, No. 3 (2009), 295-305. (Cited on page 5.)
- [OV3] L. Ornea, M. Verbitsky, *Locally conformal Kähler manifolds with potential*, Math. Ann. **348** (2010), 25-33. (Cited on pages 2, 4, and 6.)
- [OV4] L. Ornea, M. Verbitsky, *Locally conformally Kähler metrics obtained from pseudoconvex shells*, Proc. Amer. Math. Soc. **144** (2016), 325-335. (Cited on pages 5, 6, and 11.)
- [OV5] L. Ornea, M. Verbitsky, *LCK rank of locally conformally Kähler manifolds with potential*, J. Geom. Phys. **107** (2016), 92-98. (Cited on page 5.)
- [OV6] L. Ornea, M. Verbitsky, *Embedding of LCK manifolds with potential into Hopf manifolds using Riesz-Schauder theorem*, "Complex and Symplectic Geometry", Springer INdAM series, 2017, 137-148. (Cited on page 5.)
- [OV7] L. Ornea, M. Verbitsky, *Principles of locally conformally Kähler geometry*. (Cited on pages 2, 3, and 8.)
- [OV8] L. Ornea, M. Verbitsky, *Algebraic cones on LCK manifolds with potential*, arXiv:2208.05833. (Cited on pages 2, 5, 9, 12, and 14.)
- [OV9] L. Ornea, M. Verbitsky, *Lee classes on LCK manifolds with potential*, arXiv:2112.03363. (Cited on page 2.)
- [P] Prokhorov, Yuri G., *Lectures on complements on log surfaces*, MSJ Memoirs, 10. Mathematical Society of Japan, Tokyo, 2001. (Cited on page 13.)
- [Se] J.-P. Serre, *Prolongement de faisceaux analytiques cohérents*, Ann. Inst. Fourier, **16** (1966) no. 1, 363-374. (Cited on pages 8 and 9.)
- [Si] Y.-T. Siu, *Extension of twisted pluricanonical sections with plurisubharmonic weight and invariance of semipositively twisted plurigenera for manifolds not necessarily of general type*, Complex geometry (Göttingen, 2000), 223-277, Springer, Berlin, 2002. (Cited on page 14.)
- [T1] K. Tsukada, *Holomorphic forms and holomorphic vector fields on compact generalized Hopf manifolds*, Compositio Math. **93** (1994), no. 1, 1-22. (Cited on page 2.)

- [T3] K. Tsukada, *The canonical foliation of a compact generalized Hopf manifold*, Differential Geom. Appl. **11** (1999), no. 1, 13-28. (Cited on page 7.)
- [Vai1] I. Vaisman, *Locally conformal Kähler manifolds with parallel Lee form*, Rend. Mat. (6) **12** (1979), no. 2, 263-284. (Cited on page 7.)
- [Vai2] I. Vaisman, *Generalized Hopf manifolds*, Geom. Dedicata, **13** (1982), 231-255. (Cited on page 7.)
- [Ve1] M. Verbitsky, *Theorems on the vanishing of cohomology for locally conformally hyper-Kähler manifolds*, Proc. Steklov Inst. Math. no. 3 **246** (2004), 54-78. (Cited on pages 6 and 7.)
- [Ve2] M. Verbitsky, *Stable bundles on positive principal elliptic fibrations*, Math. Res. Lett. **12** (2005), no. 2-3, 251-264. (Cited on page 2.)
- [VVO] M. Verbitsky, V. Vuletescu, L. Ornea *Classification of non-Kähler surfaces and locally conformally Kähler geometry*, Russian Math. Surv. **76** (2021), 261-290. arxiv:1810.05768. (Cited on page 6.)

LIVIU ORNEA

UNIVERSITY OF BUCHAREST, FACULTY OF MATHEMATICS AND INFORMATICS,
14 ACADEMIEI STR., 70109 BUCHAREST, ROMANIA, AND:
INSTITUTE OF MATHEMATICS "SIMION STOILOW" OF THE ROMANIAN ACADEMY,
21, CALEA GRIVITEI STR. 010702-BUCHAREST, ROMANIA
lornea@fmi.unibuc.ro, liviu.ornea@imar.ro

MISHA VERBITSKY

INSTITUTO NACIONAL DE MATEMÁTICA PURA E APLICADA (IMPA)
ESTRADA DONA CASTORINA, 110
JARDIM BOTÂNICO, CEP 22460-320
RIO DE JANEIRO, RJ - BRASIL

ALSO:

LABORATORY OF ALGEBRAIC GEOMETRY,
FACULTY OF MATHEMATICS, NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL
OF ECONOMICS, 6 USACHEVA STR. MOSCOW, RUSSIA
verbit@verbit.ru, verbit@impa.br