

ON CYCLES AND MERGE TREES

JULIAN BRÜGGEMANN AND NICHOLAS A. SCOVILLE

ABSTRACT. In this paper, we extend the notion of a merge tree to that of a generalized merge tree, a merge tree that includes 1-dimensional cycle birth information. Given a discrete Morse function on a 1-dimensional regular CW complex, we construct the induced generalized merge tree. We give several notions of equivalence of discrete Morse functions based on the induced generalized merge tree and how these notions relate to one another. As a consequence, we obtain a complete solution to the inverse problem between discrete Morse functions on 1-dimensional regular CW complexes and generalized merge trees. After characterizing which generalized merge trees can be induced by a discrete Morse function on a simple graph, we give an algorithm based on the induced generalized merge tree of a discrete Morse function $f: X \rightarrow \mathbb{R}$ that cancels the critical simplices of f and replaces it with an optimal discrete Morse function.

CONTENTS

1. Introduction	1
2. Preliminaries on dMfs and Merge Trees	2
3. Inverse Problem for Multigraphs	8
4. Realization problem with simple graphs	11
5. How to find cancellations with merge trees	16
References	21

1. INTRODUCTION

Let X be a simplicial complex along with a sequence of subcomplexes $\emptyset = X_0 \subseteq X_1 \subseteq \dots \subseteq X_n = X$ known as a filtration. In the burgeoning field of topological data analysis, a filtration is often given by a sampling of points based on some increasing parameter. Geometrical and topological features of X are then estimated by studying the persistence of certain topological features [PRSZ20]. When the topological feature in question is the number of connected components, the persistence over the lifetime of the filtration is given by birth and death information and is summarized in a barcode or persistence diagram [Oud15, CVJ22]. If one wishes to not only determine birth and death information from the filtration but also how the components are evolving, i.e., which components are merging with which, one associates a merge tree to the filtration. Because the merge tree carries with it this extra information, merge trees are a rich topic of study in both the theoretical and computational settings [CHM⁺22, Cur18, MBW13, GMO⁺, CCLL22].

Date: January 5, 2023.

2020 Mathematics Subject Classification. (Primary) 57Q70; (Secondary) 05C90, 55N31.

Key words and phrases. Discrete Morse Theory, merge trees.

One way to induce a filtration on X is with a discrete Morse function [For98, For02]. Such a function f induces a filtration by considering subcomplexes associated to each critical value of f . The induced merge tree of a discrete Morse function on a tree, or 1-dimensional acyclic complex, was introduced in [JS22]. There the authors showed that a certain class of merge trees could be realized as the induced merge tree of a star graph. The authors went on to conjecture that any merge tree could be the induced merge tree of a certain discrete Morse function on a path. This conjecture was recently proved in [Brü22].

The goal of this paper is to extend the theory of merge trees and discrete Morse theory to include cycles. More specifically, given any 1-dimensional regular CW complex (i.e. a graph with or without multiedges) equipped with a discrete Morse function, we define a generalized induced Morse labeled merge tree (Definition 2.6) associated to this discrete Morse function. The generalized induced Morse labeled merge tree keeps track of not only component birth, death, and merge information but also cycle birth information via a node with a single child. After defining some basic properties, we introduce an equivalence relation on regular connected graphs called component-merge equivalence (cm-equivalence, Definition 2.10) and show that there is a one-to-one correspondence between the set of cm-equivalence classes of discrete Morse functions with only critical cells and the set of isomorphism classes of generalized Morse labeled merge tree in Theorem 3.1. In addition, we determine when a given generalized merge tree can be realized by an induced Morse function on a graph without multiedges. Unlike the case of merge trees, not all generalized merge trees can be realized. Theorem 4.1 gives a simple counting condition for when a generalized merge tree can be realized. The proof is constructive and builds off of the merge tree construction in [Brü22, Theorem 5.9]. Finally in Section 5, we give an algorithm on merge tree induced by a discrete Morse function in order to cancel critical cells of the discrete Morse function. The algorithm allows for some options depending on whether one wishes to preserve homeomorphism type of the graph or find an optimal matching. We briefly compare the algorithm to similar algorithms from the literature [LLT03a, RS20].

2. PRELIMINARIES ON DMFS AND MERGE TREES

We recall and introduce the necessary notions for this work. In this article, we use the term graph for finite abstract multigraphs without degenerate loops. That is, graphs in this work may have multiple edges between two given vertices, but they cannot have any degenerate loops, i.e., edges of the form (x, x) . This notion of graph can be geometrically interpreted as one-dimensional regular CW complexes.

On the other hand, we will use the term simple graph when we mean a graph in which there is at most one edge between two given vertices (degenerate loops are still not allowed). Simple graphs correspond to one-dimensional simplicial complexes. Since we consider graphs as geometrical objects, we also use geometrical terms like cells, simplices, and faces to describe them. For any graph X , we use $v(X)$, $e(X)$, and $b_1(X)$ to denote the number of vertices, edges, and cycles of X , respectively.

We continue with one of the most central notions of the article, namely that of a discrete Morse function.

Definition 2.1. Let X be a graph, not necessarily connected. A function $f: X \rightarrow \mathbb{R}$ is called *weakly increasing* if $f(v) \leq f(e)$ whenever vertex v is a face of edge e . A *discrete Morse function* $f: X \rightarrow \mathbb{R}$ is a weakly increasing function which is at most 2–1 and satisfies

the property that if $f(v) = f(e)$, then v is incident with e . A cell s of X is *critical* if s is the unique preimage of $f(s)$. Otherwise, s is called *matched*.

For any $a \in \mathbb{R}$, the *sublevel subcomplex* of X at a is $X_a = \{s \in X : f(s) \leq a\}$. The connected component of $s \in X$ is denoted $X[s]$. We use the notation $X_{a-\varepsilon}$ to denote the sublevel subcomplex of X immediately preceding a , i.e., $X_{a-\varepsilon} := \{\sigma : f(\sigma) < a\}$.

Remark 2.1. This definition of discrete Morse functions, due to B. Benedetti, is not equivalent to the more general definition originally given by Forman. Nonetheless, the given definition is generic in the sense that any discrete Morse function in the sense of Forman can be modified to fulfill the definition above without changing the induced acyclic matching. The definition stated above has the advantage that critical cells are distinguished by their critical values and at each level, at most either one critical cell or one pair of matched cells is added to the sublevel complex.

Definition 2.2. A *generalized merge tree* T is a rooted chiral binary tree T such that each leaf has a sibling, and inner nodes without a sibling have the same chirality as their parent node. By convention, we say that the root always has chirality L. Furthermore, the root is never regarded as a leaf, even if it only has one child node.

For nodes c of generalized merge trees we use the notation c_l/c_r for the *left/right* child node of c .

Remark 2.2. Generalized merge trees may have nodes without siblings. The notion of chirality of only children does not really deserve the name chirality because there is only one total ordering on a set with one element. We impose the condition that an only child has the same chirality as its parent node for technical reasons. We need this convention so the constructions in [Definition 2.7](#) and [Definition 2.6](#) make part 2 of [Theorem 3.1](#) work.

Generalized merge trees generalize merge trees in the sense that they keep track of more information than merge trees do.

Definition 2.3. Let T be a generalized merge tree. We call a total order \leq on the nodes of T a *Morse order* if it fulfills the following two properties for all generalized merge subtrees T' of T :

- (1) The restriction $\leq|_{T'}$ attains its maximum on the root p of T' .
- (2) The minimum of $\leq|_{T'}$ has the same chirality as p .

We call a generalized merge tree together with a Morse order (T, \leq) a *generalized Morse ordered merge tree (gMo tree)*

Remark 2.3. Assuming property 2 of [Definition 2.3](#) for every subtree T' with root p of T is equivalent to either of the following:

- For any subtree T' with root p of T , the restriction $\leq|_{T'}$ attains its minimum on the subtree with root p_l/p_r if L/R is the chirality of the root p of T' .
- For any subtree T' with root p of T , all nodes on the shortest path between p and the minimum of $\leq|_{T'}$ have the same chirality as p .

The equivalence can be proved by an inductive argument over all nodes of the shortest path between p and the minimum.

Definition 2.4. We call a generalized merge tree (T, λ) with an injective map $\lambda: T \rightarrow \mathbb{R}$ such that λ induces a Morse order on T a *generalized Morse labeled merge tree (gMl tree)*. Any such map λ is called a *Morse labeling* on T .

Let (T, λ) and (T', λ') be gMl trees. An *order equivalence* $(\varphi, \psi): (T, \lambda) \rightarrow (T', \lambda')$ of gMl trees is a pair of maps consisting of an isomorphism of the underlying generalized merge trees $\varphi: T \rightarrow T'$ and a bijection $\psi: \mathbb{R} \rightarrow \mathbb{R}$ such that the restriction $\psi|_{\text{im}(\lambda)}: \text{im}(\lambda) \rightarrow \text{im}(\lambda')$ is order preserving.

Proposition 2.1. *Let $gMoT$ be the set of generalized Morse ordered merge trees and let $gMIT$ be the set of generalized Morse labeled merge trees. Then taking the Morse order induced by a Morse labeling and using a Morse order and the labels $\{0, 1, \dots, |V(T)| - 1\}$ to induce a Morse labeling define inverse bijections*

$$iMl: gMoT/\cong \longleftrightarrow gMIT/\sim: iMo$$

where \sim denotes order equivalence.

Proof. The proof is completely analogous to [Brü22, Proposition 3.26]. \square

Definition 2.5. Let (X, f) be a discrete Morse function on a graph. We call a critical edge $\sigma \in X$ a *closing edge* if σ is part of a regular subdivision of S^1 in X such that $f(\sigma)$ is the maximum on said subdivision.

We define $C(X, f) := \{c \in X | c \text{ is closing}\}$ to be the *set of closing edges* of (X, f) and $(\bar{X}, \bar{f}) := (X \setminus C(X, f), f|_{X \setminus C(X, f)})$ to be the *spanning tree induced by f* of X .

Remark 2.4. In the previous definition, the subdivision of S^1 that any closing edge σ must be part of does not need to be unique. Nonetheless, the removal of σ would lead to the reduction of the first Betti number by one. Moreover, the notion of closing edges is well-defined because the edge σ being closing implies that it is the unique maximal edge of all subdivisions of S^1 in $X_{f(\sigma)[\sigma]}$.

Furthermore, it is immediate that $(\bar{X}, \bar{f}) := (X \setminus C(X, f), f|_{X \setminus C(X, f)})$ is a discrete Morse function on a tree.

Definition 2.6 (Induced Morse Labeled Merge Tree). Let $f: X \rightarrow \mathbb{R}$ be a discrete Morse function on a connected graph X . Let $\sigma_0 < \sigma_1 < \dots < \sigma_n$ be the critical edges of (X, f) ordered by their values under f . The generalized Morse labeled merge tree induced by (X, f) , denoted $M(X, f)$ with labeling λ_f , is constructed inductively as follows:

For the base case we construct a node $M(\sigma_n)$ with label $f(\sigma_n)$ and left chirality as root node. For any other critical edge σ_i between two 0-simplices v and w there are two cases:

- (1) The critical edge σ_i is closing $\Leftrightarrow b_1(X_{f(\sigma_i)} \setminus \sigma_i) = b_1(X_{f(\sigma_i)}) - 1 \Leftrightarrow b_0(X_{f(\sigma_i)} \setminus \sigma_i) = b_0(X_{f(\sigma_i)})$,
- (2) The critical edge σ_i is not closing $\Leftrightarrow b_1(X_{f(\sigma_i)} \setminus \sigma_i) = b_1(X_{f(\sigma_i)}) \Leftrightarrow b_0(X_{f(\sigma_i)} \setminus \sigma_i) = b_0(X_{f(\sigma_i)}) + 1$.

Depending on the case at hand, we perform the following step for the construction of $M(X, f)$:

- (1) We construct a child node c of $M(\sigma_i)$ with label $\lambda := \max\{f(\sigma) | \sigma \in X_{f(\sigma_i)-\varepsilon}, \sigma \text{ critical}\}$ and the same chirality as $M(\sigma_i)$. The node c then corresponds to the edge of X labeled λ .
- (2) We construct two child nodes c_{λ_v} and c_{λ_w} of $M(\sigma_i)$. Define $\lambda_v := \max\{f(\sigma) | \sigma \in X_{\sigma_i-\varepsilon}[v], \sigma \text{ critical}\}$ and $\lambda_w := \max\{f(\sigma) | \sigma \in X_{\sigma_i-\varepsilon}[w], \sigma \text{ critical}\}$. Then label the new nodes $\lambda_f(c_{\lambda_v}) := \lambda_v$ and $\lambda_f(c_{\lambda_w}) := \lambda_w$. If $\min\{f(\sigma) | \sigma \in X_{\sigma_i-\varepsilon}[v]\} < \min\{f(\sigma) | \sigma \in X_{\sigma_i-\varepsilon}[w]\}$

$X_{\sigma_i - \varepsilon}[w]\}$, we assign c_{λ_w} the same chirality (L or R) as c_{σ_i} and give c_{λ_w} the opposite chirality.

Continue the induction over the rest of the critical edges of X .

Remark 2.5. There is a one-to-one correspondence between vertices of X and leaves of $M(X, f)$, non-closing edges of X and parents with two children of $M(X, f)$, and closing edges (cycles) of X and parents with one child in $M(X, f)$.

Furthermore, the proof that the construction indeed produces a gMl tree is completely analogous to [Brü22, Proposition 2.20], respectively [JS22, Theorem 9].

Lemma 2.1. *Let (X, f) be a discrete Morse function on a graph and let $M(X, f)$ be the induced generalized Morse labeled merge tree. For any simplex $s \in X$, the rooted subtree $T(M(s))$ of $M(X, f)$ is induced by the connected component $X_{f(s)}[s]$ of s in the sublevel complex of level $f(s)$. Moreover, the rooted subtree $T(M(s))$ is isomorphic to $M(X_{f(s)}[s], f|_{X_{f(s)}[s]})$ as merge trees if and only if $M(s)$ has chirality L. If $M(s)$ has chirality R, then $T(M(s))$ is isomorphic to $M(X_{f(s)}[s], f|_{X_{f(s)}[s]})$ as rooted binary trees but the chiralities of all nodes are opposite to the ones of their respective nodes in the other tree.*

Proof. We observe that by Definition 2.6 the label of $M(s)$ is $f(s)$ and the chirality of $M(s)$ is decided by the minimum of $f|_{X_{f(s)}[s]}$ in comparison to the minimum of the connected component that $X_{f(s)}[s]$ got divided from at level $f(s)$. It follows inductively by construction that all nodes of the subtree $T(M(s))$ are induced by critical cells of $X_{f(s)}[s]$ because they are constructed by removing critical edges of $X_{f(s)}[s]$.

The isomorphism as rooted binary trees is constructed by the same inductive argument. Since the chirality depends on the chirality of the respective parent node, said isomorphism is compatible with the chirality if and only if the root of the rooted subtree $T(M(s))$, namely $M(s)$, has chirality L. This is true because the root of $M(X_{f(s)}[s], f|_{X_{f(s)}[s]})$ by convention always has chirality L. \square

Definition 2.7. Let (T, λ) be a generalized Morse labeled merge tree. Let $C(T) \subset V(T)$ be the set of nodes that have exactly one child node. We refer to the elements of $C(T)$ as *cycle nodes*. We denote by (\bar{T}, λ) the Morse labeled merge tree that is obtained from (T, λ) by removing the cycle nodes by connecting their parent nodes directly to their child nodes. We call (\bar{T}, λ) the underlying Morse labeled merge tree of (T, λ) .

We obtain a discrete Morse function on a graph $f_\lambda: X \rightarrow \mathbb{R}$ from (T, λ) in two steps as follows: In a first step, we construct the induced discrete Morse function on a path (P, f_λ) as in [Brü22, Definition 3.21]. For the second step, for each node c of $C(T)$ we add an edge parallel to the edge corresponding to c 's oldest descendant which has two children to P . We denote the graph obtained this way by X and extend the function $f_\lambda: P \rightarrow \mathbb{R}$ to X using the values of λ on the corresponding nodes. We denote the pair (P, f_λ) by $\Phi(T, \lambda)$.

Lemma 2.2. *We have $M(\bar{X}, f|_{\bar{X}}) \cong \bar{M}(X, f)$ as Morse labeled merge trees.*

Proof. The construction of the induced generalized merge tree induces a bijection $M: X \rightarrow V(M(X, f))$. It follows immediately by construction that M bijectively maps closing edges to nodes of $C(M(X, f))$. Hence removing the closing edges from (X, f) , that is, passing on to (\bar{X}, f) , precisely removes the nodes of $C(M(X, f))$, which corresponds to passing on to $\bar{M}(X, f)$. Hence, the statement holds because the values of f on non-closing edges are not changed. \square

Definition 2.8. Let $f: X \rightarrow \mathbb{R}$ be a dMf on a graph. For each non-empty connected component $X_c^f[v]$ of a sublevel complex X_c^f we denote by $\text{Aut}(X_c^f[v])$ the group of simplicial automorphisms of said connected component of said sublevel complex. Each $a \in \text{Aut}(X_c^f[v])$ can be extended by the identity to a set function that is a self-bijection $X \rightarrow X$. The group $\widetilde{\text{Aut}}(X_c^f[v])$ is defined to be the group of said extensions of elements of $\text{Aut}(X_c^f[v])$ by the identity. We consider $\widetilde{\text{Aut}}(X_c^f[v])$ as a subgroup of the group of all self-bijections of X . The total order on $\text{Cr}(f)$ induced by f induces chains $\widetilde{\text{Aut}}(X_{c_0}^f[v]) \subset \widetilde{\text{Aut}}(X_{c_1}^f[v]) \subset \dots$ of inclusions of subgroups. Moreover, we have inclusions $\widetilde{\text{Aut}}(X_{c_i}^f[v]) \subset \widetilde{\text{Aut}}(X_{c_j}^f[v]) = \widetilde{\text{Aut}}(X_{c_j}^f[w]) \supset \widetilde{\text{Aut}}(X_{c_i}^f[w])$ if v and w are in different connected components of some sublevel complex $X_{c_i}^f$ that merge together in some other sublevel complex $X_{c_j}^f$ for $j > i$. We define the *sublevel automorphism group* of (X, f) , denoted by $\text{Aut}_{sl}(X, f)$, to be the subgroup generated by $\bigcup_{c \in \text{Cr}(f), v \in X} \widetilde{\text{Aut}}(X_c^f[v])$. We call the elements of $\text{Aut}_{sl}(X, f)$ *sublevel automorphisms*.

Note that an element $a \in \text{Aut}_{sl}(X, f)$ is not necessarily an automorphism of X , but only a self-bijection of X that restricts to an automorphism on some connected component of some sublevel complex corresponding to a critical simplex.

Definition 2.9. Let $f: X \rightarrow \mathbb{R}$ and $g: X \rightarrow \mathbb{R}$ be dMfs on a graph X . We call f and g *sublevel-equivalent* if they have the same critical values and isomorphic sublevel complexes. If additionally $g = f * a$ holds for a sublevel automorphism $a \in \text{Aut}_{sl}(X, f) = \text{Aut}_{sl}(X, g)$, then we call f and g *symmetry-equivalent*. We call the map a a *symmetry equivalence* from f to g .

We call two dMfs $f: X \rightarrow \mathbb{R}$ and $g: Y \rightarrow \mathbb{R}$ *symmetry-equivalent* if there is a simplicial isomorphism $\varphi: X \rightarrow Y$ such that f and $g \circ \varphi$ are symmetry-equivalent.

Definition 2.10. Let (X, f) and (X', f') be critical dMfs on connected graphs. A *component-merge equivalence (cm equivalence)* is a bijection $\varphi: X \rightarrow X'$ such that at least one of the following cases holds:


- (1) φ is a symmetry equivalence.
- (2) φ fulfills the following:
 - $f' \circ \varphi = f$,
 - φ induces a bijection between the sets of connected components of sublevel complexes such that the restriction $\varphi|_{X_{a-\varepsilon}[v]}: X_{a-\varepsilon}[v] \rightarrow X'_{a-\varepsilon}[\varphi(v)]$ to each connected component is a cm equivalence, and
 - the edge $\sigma \in X$ with $f(\sigma) = a$ merges two connected components $X_{a-\varepsilon}[v_1]$ and $X_{a-\varepsilon}[v_2]$ in $X_a[v_1] = X_a[v_2]$ if and only if the edge $\varphi(\sigma)$ merges the corresponding two connected components $X'_{a-\varepsilon}[\varphi(v_1)]$ and $X'_{a-\varepsilon}[\varphi(v_2)]$ in $X'_a[\varphi(v_1)] = X'_a[\varphi(v_2)]$. Otherwise, if the edge $\sigma \in X$ with $f(\sigma) = a$ does not merge two connected components but rather closes a circle within a connected component $X_{a-\varepsilon}[v]$, then and only then $\varphi(\sigma)$ closes a circle within $X'_{a-\varepsilon}[\varphi(v)]$.

If φ re-attaches the critical edge labeled a , we call φ *non-trivial*. Moreover, if φ re-attaches the critical edge of level a and acts as a symmetry equivalence everywhere else, we say that φ is *of level a* . If φ does not re-attach any critical edge, i.e., if φ is a symmetry equivalence, we call φ a *trivial cm equivalence*.

Remark 2.6. Extending the notion of cm equivalences to dMfs with matched cells is a bit tedious. We would like to suggest getting rid of matched cells by identifying arbitrary dMfs

on graphs with critical dMfs on the corresponding graph that arises by collapsing matched cells beforehand but degenerate loops might arise in this process. Nonetheless, the newly created degenerate loops are critical by construction and the definition above works in this context.

Example 2.1. Let $f: X \rightarrow \mathbb{R}$ be the complex with discrete Morse function on the left and $f': X' \rightarrow \mathbb{R}$ be the complex with discrete Morse function on the right.



Then a cm-equivalence of critical levels $a = 7$ is given by $\varphi: X \rightarrow X'$ where $\varphi(v) = v'$ whenever $f(v) = f'(v')$ on vertices and $\varphi(e) = e'$ whenever $f(e) = f'(e')$ on edges.

Remark 2.7. It is clear from the case distinction made in [Definition 2.10](#) that any cm equivalence $\varphi: (X, f) \rightarrow (X', f')$ restricts to a bijection $\varphi|_{C(X, f)}: C(X, f) \rightarrow C(X', f')$.

Lemma 2.3. *Let $f: X \rightarrow \mathbb{R}$ and $f': X' \rightarrow \mathbb{R}$ be cm-equivalent dMfs on multigraphs. Then $M(X, f) \cong M(X', f')$ holds as generalized Morse labeled merge trees.*

Proof. Let φ be a cm equivalence $\varphi: (X, f) \rightarrow (X', f')$. Since we work with a version of discrete Morse functions which are at most 2-1, at most one non-trivial cm equivalence of level a can occur for any level a because there is at most one critical edge labeled a in $(X, f), (X', f')$, respectively. Thus, we can decompose any cm equivalence into a sequence $(\varphi_a)_a$ of non-trivial cm equivalences of decreasing levels such that each φ_a only changes the attachment of the single edge σ with $f(\sigma) = a$ and acts as a symmetry equivalence on the rest of graph and dMf. It suffices to consider a single level a because the statement then follows by induction from highest to lowest over all levels a .

For such a non-trivial cm equivalence φ_a we consider the step of the construction of the induced Ml trees that considers the critical edge σ with $f(\sigma) = a$ and the critical edge $\varphi(\sigma)$. If σ is not closing, neither is $\varphi(\sigma)$ by [Remark 2.7](#) and the inductive step follows by [[Brü22](#), Proposition 2.52]. In the case that σ is closing, so is $\varphi(\sigma)$ and we inductively assume that φ induces an isomorphism of induced generalized Ml trees everywhere outside the subtree corresponding to the connected component of $X_{a-\varepsilon}^f$ that the edge σ with $f(\sigma) = a$ is attached to. That is, on the rest of $M(X, f)$ the map $M(\varphi)$ is a bijection compatible with the chiral child relation onto $M(X', f')$ except possibly for the subtree of $M(X', f')$ which corresponds to the connected component of $X_{a-\varepsilon}^{f'}$ that the edge $\varphi(\sigma)$ is attached to.

Since the map φ is compatible with the dMfs and because it restricts to a cm equivalence $X_{a-\varepsilon}^f \rightarrow X_{a-\varepsilon}^{f'}$, the dMf f attains the same minima and maxima on the two relevant connected component of $X_{a-\varepsilon}^f$ as f' does on its counterpart of $X_{a-\varepsilon}^{f'}$ via φ . Since [Definition 2.6](#) only considers which connected component the considered edge is attached to, it makes no difference for the isomorphism type of the induced Ml trees that in general σ is attached to

said connected component of $X_{a-\varepsilon}^f$ at vertices that do not correspond via φ to the ones adjacent to $\varphi(\sigma)$ in $X_{a-\varepsilon}^{f'}$. Thus, the construction of the induced generalized MI tree produces nodes with the same chirality and label for both induced MI trees in the steps that consider $\sigma, \varphi(\sigma)$, respectively. By assumption, the restriction $\varphi_{X_{a-\varepsilon}^f} : X_{a-\varepsilon}^f \rightarrow X_{a-\varepsilon}^{f'}$ is a symmetry equivalence, so the isomorphism of MI trees extends to the subtrees that correspond to the respective connected components. \square

3. INVERSE PROBLEM FOR MULTIGRAPHS

In this section we want to describe the relationship between dMfs on graphs, generalized MI trees, generalized Mo trees, and generalized merge trees:

$$\begin{array}{ccc}
 & \Phi \circ iMl \circ \leq_{sc} & \\
 gMer & \xleftarrow{\hspace{10em}} & DMF_{graphs}^{crit} \\
 \leq_{sc} \downarrow & \uparrow \text{forget} & \downarrow \Phi \\
 & M(-, -) & \\
 gMoT & \xleftarrow{\hspace{10em}} & gMIT \\
 & iMo &
 \end{array}$$

FIGURE 1. Relationships between dMfs and merge trees

Theorem 3.1. *Let DMF_{mult}^{crit} denote the set of cm-equivalence classes of discrete Morse functions with only critical cells on multigraphs. Let MIT denote the set of isomorphism classes of generalized Morse labeled merge trees. Then the induced discrete Morse function Φ , [Definition 2.7](#), and the induced Morse labeled merge tree $M(-, -)$, [Definition 2.6](#), define maps $M(-, -): DMF_{mult}^{crit} \leftrightarrow MIT: \Phi$ that are inverse of each other in the sense that:*

- (1) *for any discrete Morse function (X, f) with only critical cells, the discrete Morse function $\Phi(M(X, f), \lambda_f)$ is cm-equivalent to (X, f) , and*
- (2) *for any generalized Morse labeled merge tree (T, λ) , we have $M(\Phi T, f_\lambda) \cong (T, \lambda)$.*

Proof. (1) Let (X, f) be a discrete Morse function with only critical cells on a graph X . We construct a cm equivalence $\varphi: (X, f) \rightarrow (\Phi(M(X, f)), f_{\lambda_f})$ as follows: First we consider the spanning trees induced by (X, f) and $(\Phi(M(X, f)), f_{\lambda_f})$ and show that they are cm equivalent. Then we define φ on the closing edges and prove that φ is a cm equivalence.

By application of [[Brü22](#), Theorem 5.6] we have a cm equivalence $\tilde{\varphi}: (\bar{X}, \bar{f}) \rightarrow (\Phi(M(\bar{X}, \bar{f})), \bar{f}_{\lambda_{\bar{f}}})$. We extend $\tilde{\varphi}$ to a cm equivalence $\varphi: (X, f) \rightarrow (\Phi(M(X, f)), f_{\lambda_f})$ by mapping each closing edge $\sigma \in X$ such that $f(\sigma) = a$ to the unique edge $\sigma' \in \Phi(M(X, f))$ with $f_{\lambda_f}(\sigma') = a$. The edge $\sigma' \in \Phi(M(X, f))$ is closing because a does not appear as a label on $(M(\bar{X}, \bar{f}), \lambda_{\bar{f}}) \cong (\bar{M}(X, f), \bar{\lambda}_f)$ since a is the value of the closing critical edge $\sigma \in X$. Furthermore, the connected component of $X_{a-\varepsilon}$ that σ is attached to corresponds to the subtree of $M(X, f)$ that consists of all descendants of $M(\sigma)$. By [Definition 2.7](#), the edge σ' is attached to the connected component of $\Phi(M(X, f))_{a-\varepsilon}$ that corresponds to said subtree. It follows that φ is a cm equivalence.

- (2) Let (T, λ) be a generalized Morse labeled Merge tree. Let $c_0 < c_1 < \dots < c_n$ be the critical values of f_λ and let $\sigma_i \in \Phi T$ such that $f_\lambda(\sigma_i) = c_i$. We recall that the induced merge tree M defines in particular a bijection between the critical cells of ΦT and

the nodes of $M(\Phi T, f_\lambda)$. For any cell $\sigma \in \Phi T$, we recall that we denote the node of $M(\Phi T, f_\lambda)$ that corresponds to σ by $M(\sigma)$. An isomorphism $(\varphi, \text{id}_{\mathbb{R}}): (T, \lambda) \rightarrow M(\Phi T, f_\lambda)$ is given by $\varphi := M \circ \phi^{-1}$. It is immediate that φ is a bijection because M and ϕ are. Furthermore, φ is by construction compatible with the respective Morse labelings. It is only left to show that φ is compatible with the chiral child relation and the respective roots.

Consider $\sigma_n \in \Phi T$. For both trees, the cell σ_n corresponds to the root of the respective tree. In $M(\Phi T, f_\lambda)$ this is the case because f_λ attains its maximum on σ_n . In (T, λ) this holds because $\phi(\sigma_n)$ holds the maximal Morse label $\lambda(\phi(\sigma_n)) = c_n$. Thus, the map φ maps the root of (T, λ) to the root of $M(\Phi T, f_\lambda)$.

For each critical edge $\sigma_i \in \Phi T$ we have one of the two cases:

- a) σ_i is closing, or
- b) σ_i is not closing.

For case b), the proof is identical to the proof of case (2) of [Brü22, Theorem 5.4]. For case a), let σ_i be a closing critical edge. In this case, the compatibility with the chiral child relation follows directly by case 1. of Definition 2.6 and the property that only children of generalized merge trees need to have the same chirality as their parent node.

□

Corollary 3.1. Since the bijection from Theorem 3.1 is compatible with the Morse labels, it induces a bijection $M(-, -): DMF_{mult}^{crit}/\leq \leftrightarrow MIT/\leq: \Phi$ where $/\leq$ denotes dividing by order equivalence.

Definition 3.1. Let \leq and \leq' be two Morse orders on a generalized merge tree T . We call \leq and \leq' *merge equivalent* if

- (1) for each inner node a of T , the node a is the maximum of a subtree T' of T with respect to \leq if and only if a is the maximum of T' with respect to \leq' , and
- (2) for each leaf a of T , the node a is the minimum of a subtree T' of T with respect to \leq if and only if a is the minimum of T' with respect to \leq' .

A merge equivalence $(T, \leq) \rightarrow (T, \leq')$ of Mo trees is a self-bijection $\psi: V(T) \cong V(T)$ such that ψ preserves conditions 1 and 2. A merge equivalence $(T, \leq) \rightarrow (T', \leq')$ is a concatenation of an isomorphism $\varphi: T \rightarrow T'$ of underlying merge trees and a merge equivalence $(T, \leq) \xrightarrow{\psi} (T, \varphi^* \leq') \xrightarrow{\varphi} (T', \leq')$.

Proposition 3.1. *Any two Morse orders \leq and \leq' on a generalized merge tree T are merge equivalent.*

Proof. The statement is proved inductively. Let a be the minimal leaf of a subtree T' of T with respect to \leq . Then a needs to be the minimal leaf of T' with respect to \leq' because otherwise \leq' would fail to be a Morseorder due to Remark 2.3. The statement for inner nodes follows similarly. □

Corollary 3.2. Two generalized Mo trees have isomorphic underlying generalized merge trees if and only if they are merge equivalent. In particular, two (not generalized) Mo trees have isomorphic underlying (not generalized) merge trees if and only if they are merge equivalent.

For any generalized merge tree T , there are several ways to induce canonical Morse orders on T . We introduce the sublevel-connected Morse order (generalization of [Brü22, Definition 4.1]) on any given generalized merge tree in the following:

To define the sublevel-connected Morse order, we first observe that every node a of T is uniquely determined by the shortest path from the root to a . We recall that the depth of T is the maximal length of any path in T that appears as the shortest path from the root to a leaf. Because T is chiral, we can identify such shortest paths with certain words:

Definition 3.2. Let T be a generalized merge tree of depth n and let a be a node of T . The *path word* corresponding to a is a word $a_0a_1\dots a_n \in \{L, R, _ \}^{n+1}$ where $_$ denotes the empty letter. If a is of depth k , the letters $a_0\dots a_k$ are given by the chirality of the nodes belonging to the shortest path from the root to a . The letters $a_{k+1}\dots a_n$ are then empty.


Remark 3.1. Let a, b be nodes of a generalized merge tree T and let $a_0a_1\dots a_n$ be the path word corresponding to a and $b_0b_1\dots b_n$ be the path word corresponding to b . Then the equation $a_0 = b_0 = L$ always holds because we consider paths that begin at the root. Because $a_0 = b_0 = L$ and because we consider finite trees, there is always a maximal $k \in \mathbb{N}$ such that $a_i = b_i$ holds for all $i \leq k$. Furthermore, the last non-empty letter of a path word is always the chirality of the considered node.

Definition 3.3. Let T be a generalized merge tree. We define the *sublevel-connected Morse order* \leq_{sc} on the nodes of T as follows:

Let a, b be arbitrary nodes of T . Let $a_0a_1\dots a_n$ be the path word corresponding to a and $b_0b_1\dots b_n$ the path word corresponding to b (see Definition 3.2). Furthermore, let $k \in \mathbb{N}$ be maximal such that $a_i = b_i$ for all $i \leq k$. If $a_k = b_k = L/R$ we define $a \leq_{sc} b$ if and only if one of the following cases hold:

- a) $a_{k+1} = L$ and $b_{k+1} = R/a_{k+1} = R$ and $b_{k+1} = L$
- b) $b_{k+1} = _$
- c) $a = b$

Example 3.1. We depict the sublevel-connected Morse order in the following example:



Proposition 3.2. *The construction of the sublevel-connected Morse order and forgetting the Morse order defines a pair of inverse bijections*

$$\leq_{sc}: Mer/\cong \longleftarrow gMoT/\sim: \text{forget}$$

where \sim denotes merge equivalence.

Proof. The statement follows directly by [Corollary 3.2](#). \square

To summarize our results of this section, we take a look at how [Proposition 2.1](#), [Theorem 3.1](#), and [Proposition 3.2](#) turn the different maps from [Figure 1](#) into bijections by dividing out the needed notion of equivalence. If we do not divide out any equivalence relation, the map Φ is not even well-defined. The maps $M(-, -)$, iMo , and forget are surjective, but not injective. The maps \leq_{sc} and iMl are injective but not surjective.

Identifying cm-equivalent dMfs makes Φ a well-defined map and, moreover, a bijection which is inverse to $M(-, -): DMF_{graphs}^{crit} \rightarrow gMIT$ by [Theorem 3.1](#). Inverting order equivalences turns iMo and iMl into inverse bijections. Finally, inverting merge equivalences makes \leq_{sc} and forget inverse of each other. As a consequence, we have a complete description of the inverse problem for critical discrete Morse functions on multigraphs and their induced merge trees. The characterization for arbitrary discrete Morse functions on 1-dim regular CW complexes follows by collapsing matched cells and then applying a version of [Theorem 3.1](#) that incorporates [Remark 2.6](#). However, this procedure secretly makes use of a feature which might become problematic if one tries to generalize the result to higher dimensions: we are starting with regular CW complexes. Hence, the complex that arises by performing the simple collapses described by a Morse matching is not arbitrary but subject to being simple homotopy equivalent to a regular CW complex. It is a feature of dimension one that all 1-dimensional CW complexes are simple homotopy equivalent to a 1-dimensional regular CW complex. Hence, defining cm-equivalences becomes more difficult in a higher-dimensional setting, in particular, if one wants to work with non-critical discrete Morse functions. This would lead to the need to analyze which CW complexes are simple homotopy equivalent to regular CW complexes in order to know for which generality a notion of cm-equivalence is needed.

4. REALIZATION PROBLEM WITH SIMPLE GRAPHS

Let T be a generalized merge tree. Recall that $C(T) = C$ denotes the set of all cycle nodes of T . For any $c \in C$, let c_u denote the unique child of c . For any $v \in T$, let $T(v)$ denote the subtree of T with root v and let $\ell(v)$ denote the number of leafs of $T(v)$.

Theorem 4.1. *Let T be a generalized merge tree. Then there exists a simple graph X and discrete Morse function $f: X \rightarrow \mathbb{R}$ such that $M(X, f) = T$ if and only if for every $c \in C(T)$,*

$$|C(T(c_u))| < \frac{(\ell(c_u) - 2)(\ell(c_u) - 1)}{2}.$$

Furthermore, X can be made planar if and only if

$$|C(T(c_u))| < 2 \cdot \ell(c_u) - 5.$$

Proof. Suppose there exists a simple graph X and discrete Morse function $f: X \rightarrow \mathbb{R}$ such that $M(X, f) = T$, and suppose by contradiction that there is a $c \in C(T)$ with the property

that

$$|C(T(c_u))| \geq \frac{(\ell(c_u) - 2)(\ell(c_u) - 1)}{2}.$$

By [Lemma 2.1](#), the rooted subtree $T(c_u)$ is isomorphic as rooted binary trees to the induced Morse labeled merge tree of $X_{f(s)}[s]$ where s is the simplex of X such that $M(s) = c_u$. Letting v be the number of vertices in $X_{f(s)}[s]$, e the number of edges in $X_{f(s)}[s]$, and b_1 the number of cycles in $X_{f(s)}[s]$, we see that

$$\begin{aligned} e &= v - 1 + b_1 \\ &\geq v - 1 + \frac{(v - 1)(v - 2)}{2} \\ &= v - 1 + \frac{v(v - 1)}{2} + 1 - v \\ &= \frac{v(v - 1)}{2} \end{aligned}$$

which is the maximum number of edges any connected component can have. Hence it is impossible to add a cycle to this connected component so that

$$|C(T(c_u))| < \frac{(\ell(c_u) - 2)(\ell(c_u) - 1)}{2}.$$

for all $c \in C$. Now suppose further that X is planar, and suppose by contradiction that $|C(T(c_u))| \geq 2 \cdot \ell(c_u) - 5$. Using the same notation as above, we have

$$\begin{aligned} e &= v - 1 + b_1 \\ &\geq v - 1 + 2v - 5 \\ &= 3v - 6. \end{aligned}$$

But it is well known that a simple planar graph satisfies $e \leq 3v - 6$ [[Bic20](#), Theorem 5.9]. Hence either $X_{f(s)}[s]$ is not planar or maximal planar in the case of equality. In either case, another edge cannot be added to $X_{f(s)}[s]$ without breaking planarity, and thus the result.

For the other direction, given the generalized Merge tree T , construct the sublevel-connected Morse order \leq_{sc} ([Definition 3.3](#)) on the nodes of T . Associate to this Morse order a Morse labeling $\lambda: T \rightarrow \mathbb{R}$ such that $a \leq_{sc} b$ if and only if $\lambda(a) \leq \lambda(b)$. Apply the construction in [Definition 2.7](#) to (T, λ) to obtain the underlying merge tree $(\bar{T}, \bar{\lambda})$. By [[Brü22](#), Theorem 6.5], there is a path P and discrete Morse function $\bar{f}: P \rightarrow \mathbb{R}$ such that $M(P, \bar{f}) = (\bar{T}, \bar{\lambda})$. We will inductively attach edges to P in one-to-one correspondence with cycle nodes of T . Each edge will be labeled with the same label as its corresponding cycle node.

Induce on the cycle nodes of T with respect to the sublevel-connected Morse order $c_1 \leq_{sc} c_2 \leq_{sc} \dots$. For the base case $i = 1$, write $P = X^1$. We have by hypothesis that

$$|C(T(c_{1u}))| < \frac{(\ell(c_{1u}) - 2)(\ell(c_{1u}) - 1)}{2}.$$

In addition, $M(P, \bar{f}) = (\bar{T}, \bar{\lambda})$ so $c_{1u} = M(s_1)$ for some simplex $s_1 \in P = X^1$. Applying the correspondence noted in [Remark 2.5](#), this inequality means that


$$b_1(X^1[s_1]) < \frac{(v(X^1[s_1]) - 2)(v(X^1[s_1]) - 1)}{2}.$$

By the computation in the forward direction, this implies that $e(X^1[s_1]) < \frac{v(X^1[s_1])(v(X^1)-1)}{2}$. Hence there are at least two vertices in $X^1[s_1]$ not connected by an edge. A choice of vertex can be made by defining a lexicographic ordering on a subset of ordered pairs of the vertex set of P where an ordered pair (v, u) satisfies $\bar{f}(v) < \bar{f}(u)$ and $(v, u) < (v', u')$ if $\bar{f}(v) < \bar{f}(v')$ or $\bar{f}(u) < \bar{f}(u')$ when $\bar{f}(v) = \bar{f}(v')$. Since all the vertices of P are given distinct values, $<$ is a total order. Add an edge e_1 incident with the vertices in the minimum pair over all available pairs to create $X^2 = X^1 \cup \{e_1\}$ and extend \bar{f} to $f^1(e_1) := \lambda(c_1)$. Then $M(X^2, f^1) \simeq (T_{\leq \lambda(c_1)}, \lambda|_{T_{\leq \lambda(c_1)}})$. The inductive step is identical to the base case.


Now suppose that $|C(T(c_u))| < 2 \cdot \ell(c_u) - 5$ for all cycle nodes $c \in T$. By the forward direction, this is equivalent to $e < 3v - 6$ in the corresponding sublevel complex of X . The method of construction is analogous to the above construction and utilizes the fact that if a planar simple graph satisfies $e < 3v - 6$, then it is not maximal planar and hence an edge can be added while maintaining planarity [Bic20, Corollary 5.11]. \square

Remark 4.1. While the choices made in the construction of the simple graph X in Theorem 4.1 may be thought of as one canonical choice, the sublevel-connected Morse order is only one possible representative for the Morse order. Another just as natural (and shuffle equivalent) order would be the index Morse order [Brü22, Definition 3.3]. Furthermore, once a Morse order is picked, there are often several possible simple graphs with discrete Morse functions all related by cm equivalence that represent the given generalized merge tree.


Example 4.1. To illustrate the construction in the planar case, consider the generalized merge tree T pictured below:




We constructed the sublevel-connected Morse order and induced Morse labeling λ in Example 3.1.




We then pass to the underlying merge tree \bar{T} and restrict λ to \bar{T} in order to apply [Brü22, Theorem 6.5] to obtain the index-ordered discrete Morse function on the graph below with induced merge tree \bar{T} .




We induce on the cycle nodes ordered by their generalized Morse label. The first cycle to be introduced is cycle node with label 8. This will be a cycle added to the graph




to the component with the edge labeled 7.




We then add the cycle corresponding to the node labeled 9 to this same graph.




Skipping to the cycle node labeled 23, we see that we need to add a cycle to the component with edge labeled 22:




We add this edge



and must add another cycle corresponding to cycle node labeled 24 to this same connected component.



Notice that this component is now a complete graph and that no more cycles can be added. The final graph with discrete Morse function that induces the given generalized merge tree is



5. HOW TO FIND CANCELLATIONS WITH MERGE TREES

In this section, we present a way to find cancellations of critical cells of dMfs with the help of the induced merge tree. The idea is to start with an arbitrary dMf that only has critical cells and to perform cancellations along the merge tree. It will turn out that depending on the chosen critical dMf, one sometimes has to decide whether one wants to keep the homeomorphism type of X or to produce an optimal dMf.

Remark 5.1. In order to obtain an arbitrary dMf on a graph X that has only critical cells, one can simply choose any total order on the vertices and a total order on the edges. Then assign the values $0, \dots, |V(X)| - 1$ to the vertices according to the chosen order and the numbers $|V(X)|, \dots, |V(X)| + |E(X)|$ to the edges. This always produces an indexed dMf which is not necessary for the following algorithm. Perhaps more sophisticated approaches to finding a critical dMf might be useful, but for now we are satisfied with this simple one.

Given a critical dMf $f: X \rightarrow \mathbb{R}$, the algorithm proceeds as follows:

- (1) Calculate the induced generalized Morse labeled merge tree $M(X, f)$.
- (2) Consider the leaves of $M(X, f)$ in descending order with respect to their labels. Suppose we are considering the leaf c with the maximal label k such that c is critical. Let p be the youngest ancestor of c such that p is neither a cycle node nor matched. We have the following cases:
 - a) The vertex $M^{-1}(c)$ is adjacent to the edge $M^{-1}(p)$.
 - b) The vertex $M^{-1}(c)$ is not adjacent to the edge $M^{-1}(p)$ but there is a symmetry equivalence a of (X, f) such that $a(M^{-1}(c))$ is adjacent to $a(M^{-1}(p))$.
 - c) The vertex $M^{-1}(c)$ is not adjacent to the edge $M^{-1}(p)$ and there is no symmetry equivalence as in case b).

In case a) we match $M^{-1}(c)$ and $M^{-1}(p)$. This does not produce cycles because we explicitly exclude cycle nodes from the matching.

In case b) we apply the symmetry equivalence a and then proceed as in case a). Since symmetry equivalences are only automorphisms of connected components of sublevel complexes, we do not alter the homeomorphism type of X in the process.

In case c) we have to make a decision, we could

- i) simply skip c and leave $M^{-1}(c)$ critical,
- ii) apply a cm equivalence in order to make $M^{-1}(c)$ and $M^{-1}(p)$ adjacent, then proceed as in case a), or
- iii) observe that there is a unique gradient flow line from $M^{-1}(c)$ to $M^{-1}(p)$ and cancel the two cells along this flow line.

If we choose possibility i), we might not obtain an optimal matching but we preserve the homeomorphism type of X . In case ii), we produce an optimal matching but may change the homeomorphism type of X . In case iii), we preserve the homeomorphism type of X and obtain an optimal matching but we change the order of the vertices induced by f .

In the preceding algorithm, many claims are made, most of which are straightforward to prove. For example, the fact that the cases 2a), 2b), 2c)i), and 2c)ii) work as described follows immediately from the definition of the used equivalences. However in general it does not appear easy to decide whether case 2b) or 2c) holds. Nonetheless, case 2c)iii) is not so obvious, so we consider it in the following lemma:


Lemma 5.1. *Let X be a graph, $f: X \rightarrow \mathbb{R}$ a critical dMf , and $M(X, f)$ the induced generalized Morse labeled merge tree. At any point of the cancellation algorithm, there is always a unique gradient flow line from the vertex $M^{-1}(c)$ corresponding to the maximally labeled unmatched leaf c to the edge $M^{-1}(c)$ corresponding to its youngest unmatched ancestor p .*

Proof. If $M^{-1}(c)$ and $M^{-1}(p)$ are adjacent, there is nothing to prove. If $M^{-1}(c)$ and $M^{-1}(p)$ are not adjacent then there is no other non-closing critical edge in $X_{f(M^{-1}(p)-\varepsilon)}[M^{-1}(c)]$ because otherwise said other younger critical edge would induce a younger unmatched ancestor of c .


Since $M^{-1}(c)$ is a critical vertex with no adjacent critical edge, all adjacent edges of $M^{-1}(c)$ are matched with their respective other vertex. This means that on all adjacent edges, there is a gradient flow line pointing towards $M^{-1}(c)$. Following these gradient flow lines backwards either leads to matched vertices that are adjacent only to the edge they are matched with, or to the unique critical edge of $X_{f(M^{-1}(p)-\varepsilon)}[M^{-1}(c)]$. One of the flow lines eventually leads to $M^{-1}(p)$ because $X_{f(M^{-1}(p)-\varepsilon)}[M^{-1}(c)]$ is connected.¹ The flow line is unique because closing edges remain critical, that is, because we only match cells along a subtree of X . \square

We apply the cancellation algorithm in the following example:


Example 5.1. We consider the graph:




We put some critical discrete Morse function on it and calculate the induced generalized merge tree:



We apply step 2a) as long as possible:




Now is the first time we run into case 2b)




In this example, the cases 2a) and 2b) sufficed.

We consider the following example in order to see how quickly things can fail:


Example 5.2. We consider the following dMf and its induced merge tree:




After twofold application of step 2a), we have the following:



Now we have reached case 2c). We would need to have the vertex labeled 1 adjacent to the edge labeled 10. But this is not possible because all symmetry equivalences leave the vertex labeled 1 adjacent to the edge labeled 9 and no other edge. The three different solutions result in the following:




Example 5.3. A sublevel symmetry of the last sublevel complex before the “merge tree algorithm” fails may not always be sufficient. Consider the graph with discrete Morse function given below.




Proceeding as before, we obtain a matching on the graph until the algorithm specifies to match the vertex labeled 1 with the edge labeled 13. Since these simplices are not incident, we need to find a sublevel-symmetry of sublevel 12. However, the sublevel subcomplex X_{12}

is given by



which is well-known to have no non-trivial automorphisms. There is also no symmetry equivalence of a lower level than 12 that makes the vertex labeled 1 and the edge labeled 13 adjacent. However, the three different workarounds mentioned earlier result in the following:



At the end of this section, we compare our algorithm for finding cancellations of critical cells to similar algorithms from the literature. In [LLT03b] the authors introduce an algorithm to find optimal discrete Morse functions on 2-dimensional manifolds which they generalize to higher dimensions and more general complexes in [LLT03a], even though losing the guarantee for optimality in the process. The main similarity to our approach is the use of an auxiliary tree structure, in our case the generalized merge tree, in the case of [LLT03a] a spanning hyperforest of a hypergraph associated to the Hasse diagram of a discrete Morse function.

In [RS20], the authors provide an algorithm to find optimal discrete Morse functions on trees. Said algorithm, combined with any standard algorithm to find spanning trees, can easily be generalized to provide optimal discrete Morse functions on graphs with a prescribed critical vertex.

The main feature of our new approach, compared to the pre-existing ones, seems to be that our algorithm allows to preserve certain properties of a given discrete Morse function. In certain cases, such a discrete Morse function might be given by an application and, therefore, might be worth preserving. We conjecture that, given a suitable version of higher merge trees, our algorithm can be generalized to higher dimensions. Since finding optimal Morse matchings is MAX-SNP hard, such a generalization might either fail to be optimal or be inconvenient to work with in practice. Nonetheless, we hope to find interesting classes of examples in which such a generalized algorithm happens to be performative and informative.

REFERENCES

- [Bic20] Allan Bickle. *Fundamentals of graph theory*, volume 43 of *Pure and Applied Undergraduate Texts*. American Mathematical Society, Providence, RI, [2020] ©2020. [12](#), [13](#)
- [Brü22] Julian Brüggemann. On Merge Trees and Discrete Morse Functions on Paths and Trees. *J Appl. and Comput. Topology*, 11 2022. DOI: <https://doi.org/10.1007/s41468-022-00101-w>. [2](#), [4](#), [5](#), [7](#), [8](#), [9](#), [10](#), [12](#), [13](#), [14](#)
- [CCLL22] Robert Cardona, Justin Curry, Tung Lam, and Michael Lesnick. The universal ℓ^p -metric on merge trees. In *38th International Symposium on Computational Geometry*, volume 224 of *LIPICs. Leibniz Int. Proc. Inform.*, pages Art. No. 24, 20. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2022. [1](#)
- [CHM⁺22] Justin Curry, Haibin Hang, Washington Mio, Tom Needham, and Osman Berat Okutan. Decorated merge trees for persistent topology. *J. Appl. Comput. Topol.*, 6(3):371–428, 2022. [1](#)
- [Cur18] Justin Curry. The fiber of the persistence map for functions on the interval. *J. Appl. Comput. Topol.*, 2(3-4):301–321, 2018. [1](#)
- [CVJ22] Gunnar Carlsson and Mikael Vejdemo-Johansson. *Topological data analysis with applications*. Cambridge University Press, Cambridge, 2022. [1](#)
- [For98] Robin Forman. Morse theory for cell complexes. *Adv. Math.*, 134(1):90–145, 1998. [2](#)
- [For02] Robin Forman. A user’s guide to discrete Morse theory. *Sém. Lothar. Combin.*, 48:Art. B48c, 35, 2002. [2](#)
- [GMO⁺] Ellen Gasparovic, Elizabeth Munch, Steve Oudot, Katharine Turner, Bei Wang, and Yusu Wang. Intrinsic interleaving distance for merge trees. *trees*, 38(37):32. [1](#)
- [JS22] Benjamin Johnson and Nicholas A. Scoville. Merge trees in discrete Morse theory. *Res. Math. Sci.*, 9:Paper No. 49, 07 2022. [2](#), [5](#)
- [LLT03a] Thomas Lewiner, Hélio Lopes, and Geovan Tavares. Optimal discrete morse functions for 2-manifolds. *Computational Geometry*, 26(3):221–233, 2003. [2](#), [20](#)
- [LLT03b] Thomas Lewiner, Hélio Lopes, and Geovan Tavares. Toward optimality in discrete morse theory. *Experimental Mathematics*, 12(3):271–285, 2003. [20](#)
- [MBW13] Dmitriy Morozov, Kenes Beketayev, and Gunther Weber. Interleaving distance between merge trees. *Discrete and Computational Geometry*, 49(22-45):52, 2013. [1](#)
- [Oud15] Steve Y. Oudot. *Persistence theory: from quiver representations to data analysis*, volume 209 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015. [1](#)
- [PRSZ20] Leonid Polterovich, Daniel Rosen, Karina Samvelyan, and Jun Zhang. *Topological persistence in geometry and analysis*, volume 74 of *University Lecture Series*. American Mathematical Society, Providence, RI, [2020] ©2020. [1](#)
- [RS20] Ian Rand and Nicholas A. Scoville. Discrete Morse functions, vector fields, and homological sequences on trees. *Involve*, 13(2):219–229, 2020. [2](#), [20](#)

(Julian Brüggemann) MAX PLANCK INSTITUTE FOR MATHEMATICS, BONN, GERMANY
Email address: brueggemann@mpim-bonn.mpg.de

(Nicholas A. Scoville) DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, URSINUS COLLEGE,
 COLLEGEVILLE PA 19426
Email address: nscoville@ursinus.edu