Exact Controllability for a Schrödinger equation with dynamic boundary conditions^{*}

Alberto Mercado[†] Roberto Morales[‡]

January 9, 2023

Abstract

In this paper, we study the controllability of a Schrödinger equation with mixed boundary conditions on disjoint subsets of the boundary: dynamic boundary condition of Wentzell type, and Dirichlet boundary condition. The main result of this article is given by new Carleman estimates for the associated adjoint system, where the weight function is constructed specially adapted to the geometry of the domain. Using these estimates, we prove the exact controllability of the system with a boundary control acting only in the part of the boundary where the Dirichlet condition is imposed. Also, we obtain a distributed exact controllability result for the system.

Keywords Schrödinger equation, dynamic boundary conditions, exact controllability, Carleman estimates.

AMS 35Q41, 93B05, 93B07, 93C05, 35M13.

1 Introduction

In this article, controllability properties of a non-conservative Schrödinger equation with dynamic boundary conditions of Wentzell type are studied. Let $\Omega \subset \mathbb{R}^n$, $n \ge 2$, be a bounded domain with regular boundary $\partial\Omega$ such that $\partial\Omega = \Gamma_0 \cup \Gamma_1$ with Γ_0 and Γ_1 are two closed subsets and $\Gamma_0 \cap \Gamma_1 = \emptyset$; a typical example is given by the annulus $\Omega = \{x \in \mathbb{R}^n : R_1 < |x| < R_2\}$. Then we consider the system given by

$$\begin{cases} i\partial_t y + d\Delta y - \vec{q}_1 \cdot \nabla y + q_0 y = 0, & \text{in } \Omega \times (0, T), \\ i\partial_t y_{\Gamma} - d\partial_{\nu} y + \delta \Delta_{\Gamma} y_{\Gamma} - \vec{q}_{\Gamma,1} \cdot \nabla_{\Gamma} y_{\Gamma} + q_{\Gamma,0} y_{\Gamma} = 0, & \text{on } \Gamma_1 \times (0, T), \\ y = y_{\Gamma}, & \text{on } \Gamma_1 \times (0, T), \\ y = \mathbb{1}_{\Gamma_*} h, & \text{on } \Gamma_0 \times (0, T), \\ (y(0), y_{\Gamma}(0)) = (y_0, y_{\Gamma,0}), & \text{in } \Omega \times \Gamma_1, \end{cases}$$
(1.1)

where $h \in L^2(\Gamma_* \times (0,T);\mathbb{C})$ (with $\Gamma_* \subseteq \Gamma_0$) is a control acting on a subset of $\Gamma_0 \times (0,T)$. Besides, d > 0 and $\delta > 0$ are parameters representing the diffusion on the bulk and on the boundary, respectively. Moreover, $(\vec{q_1}, \vec{q_{\Gamma,1}}) \in [L^{\infty}(\Omega \times (0,T);\mathbb{C})]^n \times [L^{\infty}(\Gamma_1 \times (0,T);\mathbb{C})]^n$ and $(q_0, q_{\Gamma,0}) \in$

^{*}The first author ... The second author has been supported by FONDECYT 3200830

[†]Departamento de Matemática, Universidad Técnica Federico Santa María, Casilla 110-V, Valparaíso, Chile e-mail: alberto.mercado@usm.cl

[‡]Departamento de Matemática, Universidad Técnica Federico Santa María, Casilla 110-V, Valparaíso, Chile e-mail: roberto.moralesp@usm.cl

 $L^{\infty}(\Omega \times (0,T); \mathbb{C}) \times L^{\infty}(\Gamma_1 \times (0,T); \mathbb{C})$ are lower order potentials. We denote by Δ_{Γ} the Laplace-Beltrami operator, ∇_{Γ} is the tangential gradient and $\partial_{\nu} y$ the normal derivative associated to the outward normal ν of Ω . We wish to investigate controllability properties of the system (1.1). Roughly speaking, we are interested in finding conditions on the geometry of the domain and the parameters of the system such that the associated solution (y, y_{Γ}) can be driven to any given state at time T > 0. More precisely, we study the exact controllability of (1.1), which can be defined as follows:

Definition 1.1. System (1.1) is said to be **exactly controllable** at time T > 0 in space X if for every states $(y_0, y_{\Gamma,0}), (y_T, y_{\Gamma,T}) \in X$, there exists a (boundary) control $h \in L^2(\Gamma_* \times (0,T); \mathbb{C})$ such that the associated solution (y, y_{Γ}) satisfies

$$(y(T), y_{\Gamma}(T)) = (y_T, y_{\Gamma,T}), \text{ in } \Omega \times \Gamma_1.$$

We point out that the control h acts only on a portion of the boundary Γ_0 . This means that the equation in the bulk is controlled directly by h, while the equation on the boundary Γ_1 is being controlled indirectly through the side condition $y = y_{\Gamma}$ on $\Gamma_1 \times (0, T)$.

Linear and nonlinear Schrödinger equations have been intensely studied due to their applications to plasma physics and laser optics, see e.g. [2] and [12]. On the other hand, Schrödinger equation can be represented as two suitable diffusion equations which their solutions are in duality, see [26]. In our case, the system (1.1) can model the evolution of diffusion processes in an object Ω and its interaction with another one, having a thick border Γ_1 .

Controllability properties of the Schrödinger equation has been studied by several authors in the last 30 years. In [20], G. Lebeau proved that the Geometric Control Condition (GCC) for the exact controllability of the wave equation is sufficient for the exact controllability of the Schrödinger equation in any time T > 0. The proof of this result is based on the diadic decomposition of the Fourier representation of solutions of the Schrödinger equations with velocity of propagation tending to infinity. Besides, a particular case of this work is due to E. Machtyngier in [22]. In this case, the exact controllability in $H^{-1}(\Omega)$ with L^2 -boundary control and exact controllability in $L^2(\Omega)$ with L^2 -controls supported in a neighborhood of the boundary are achieved. These results were obtained using multiplier techniques, Hilbert Uniqueness Method (HUM) and Holmgren unique continuation principle, a property which relies on the analyticity of the coefficients.

On the other hand, contrary to the results of hyperbolic equations, there are relevant controllability results for the Schrödinger equation in some situations in which the GCC is not fulfilled in any time T. We refer to [15] and [7] where the main results are based on the decomposition of the Plate operator $\partial_t^2 + \Delta^2$ in two conjugate Schrödinger operators in the following form:

$$\partial_t^2 u + \Delta^2 u = (i\partial_t + \Delta)(-i\partial_t + \Delta)u.$$

A useful tool to obtain observability inequalities is given by the so-called Carleman estimates. In the case of Schrödinger equation with Dirichlet boundary conditions, several authors established controllability and stability results using these estimates combining another techniques. In [5], the authors derived a Carleman estimate under a strict pseudoconvexity condition, or equivalently, under a strong convexity of the weight function in the space variable. We also mention that the observation is taken in regions according to the classical geometric conditions typically used for the wave equation. As a consequence of this result, the authors proved a Lipschitz stability estimate for an inverse problem for the potential of the Schrödinger operator in $H^1(\Omega)$. On the other hand, in [24] the authors proved a more general Carleman estimate replacing the strong pseudoconvexity condition for a weaker one, where the Hessian of the weight function may be degenerate at some points, allowing to consider weights of the form $\psi(x) = x \cdot e$, where $e \in \mathbb{R}^n$, resulting in observation regions not satisfying the geometric control conditions. However, in these estimates only a part of the weighted- H^1 energy may be bounded by boundary/internal observations. For other results concerning controllability of the Schrödinger equation with Dirichlet boundary conditions we refer to [27], [1], [28], [29] and [33].

Controllability properties of PDEs with dynamic boundary conditions have also been intensively studied in the last years. We mention the works [23] and [17] where the authors studied controllability properties of parabolic equations with these kind of boundary conditions. Using the approach of Fursikov and Imanuvilov and using the fact that the tangential derivatives of the associated weight function vanish, the authors determined the null controllability of such systems with arbitrary small regions. Based on these results, in [32] it is studied the existence of insensitizing controls for systems of parabolic equations with Wentzell boundary conditions. We also refer the works [13] and [14] where some inverse problems for such models are considered.

Controllability properties in the case of the wave equation with dynamic boundary conditions were also recently studied. In [11], the authors studied a non-conservative wave system acting in a domain Ω with the same topology structure that the case studied here. They proved a global Carleman estimate with observations on both sides of the boundary, i.e., on the usual observation region satisfying the geometric control condition and on the part of the boundary where the dynamic conditions are given; as a consequence, it is obtained a controllability result with two controls. On the other hand, in [6], the authors determined that, in the particular situation when Ω is an *n*-dimensional interval, the control acting on the subset of the boundary where the Wentzell boundary conditions are imposed can be neglected. This result is based on Fourier expansions and a generalization of the Ingham inequalities due to Mehrenger for *n*-dimensional intervals.

Recently, some uniform stability results on the solutions for the Schrödinger and Ginzburg-Landau equations with dynamic boundary conditions were obtained in [8] and [9], respectively. We also mention the articles [18] and [19] where the authors studied Carleman estimates in H^1 and L^2 , respectively, for non-conservative Schrödinger equations with different types of boundary conditions. However, to the best of the authors' knowledge, this is the first time that controllability properties of the Schrödinger equation with Wentzell boundary conditions like (1.1) are studied.

1.1 General setting

In this section, we set up the notation and terminology used in this paper. We consider the set $\Gamma_1 \subset \partial\Omega$ as an (n-1)-dimensional compact Riemannian submanifold equipped by the Riemannian metric g, induced by the natural embedding $\Gamma_1 \subset \mathbb{R}^n$. It is possible to define the differential operators on Γ_1 in terms of the Riemannian metric g. However, for the purposes of this article, it will be enough to use the most important properties of the underlaying operators and spaces. The details can be found, for instance, in [16] and [30]. For the sake of completeness, we recall some of those properties.

The tangential gradient ∇_{Γ} of y_{Γ} at each point $x \in \Gamma_1$ can be seen as the projection of the standard Euclidean gradient ∇y onto the tangent space of Γ_1 at $x \in \Gamma_1$, where y_{Γ} is the trace of y on Γ_1 . That is to say,

$$\nabla_{\Gamma} y_{\Gamma} = \nabla y - \nu \partial_{\nu} y,$$

where $y = y_{\Gamma}$ on Γ_1 and $\partial_{\nu} y$ is the normal derivative associated to the outward normal ν . In this

way, the tangential divergence $\operatorname{div}_{\Gamma}$ in Γ_1 is defined by

$$\operatorname{div}_{\Gamma}(F_{\Gamma}): H^{1}(\Gamma_{1}; \mathbb{R}) \to \mathbb{R}, \quad y_{\Gamma} \mapsto -\int_{\Gamma_{1}} F_{\Gamma} \cdot \nabla_{\Gamma} y_{\Gamma} \, d\sigma.$$

The Laplace Beltrami operator is given by $\Delta_{\Gamma} y_{\Gamma} = \operatorname{div}_{\Gamma}(\nabla_{\Gamma} y_{\Gamma})$ for all $y_{\Gamma} \in H^2(\Gamma_1; \mathbb{R})$. In particular, the surface divergence theorem holds:

$$\int_{\Gamma_1} \Delta_{\Gamma} y z \, d\sigma = -\int_{\Gamma_1} \nabla_{\Gamma} y \cdot \nabla_{\Gamma} z \, d\sigma, \quad \forall y \in H^2(\Gamma_1; \mathbb{R}), \quad \forall z \in H^1(\Gamma_1; \mathbb{R})$$

In order to simplify the notation, here and subsequently, the function spaces refer to complexvalued functions unless otherwise stated.

We introduce the Hilbert space $\mathcal{H} = L^2(\Omega) \times L^2(\Gamma_1)$ in \mathbb{C} equipped with the scalar product

$$\langle (u, u_{\Gamma}), (v, v_{\Gamma}) \rangle_{\mathcal{H}} = \int_{\Omega} u \overline{v} \, dx + \int_{\Gamma_1} u_{\Gamma} \overline{v_{\Gamma}} \, d\sigma.$$

Moreover, for $m \in \mathbb{N}$, we consider the space

$$H^m_{\Gamma_0}(\Omega) = \{ u \in H^m(\Omega) : u = 0 \text{ on } \Gamma_0 \},\$$

which is a closed subspace of the Sobolev space $H^m(\Omega)$. In the same manner, we define the space

$$\mathcal{V}^m = \{(u, u_{\Gamma}) \in H^m_{\Gamma_0}(\Omega) \times H^m(\Gamma_1) : u = u_{\Gamma} \text{ on } \Gamma_1\}.$$

and by simplicity we write $\mathcal{V} = \mathcal{V}^1$. Due to the Poincaré inequality and a trace theorem, we have

$$\int_{\Omega} |u|^2 dx + \int_{\Gamma_1} |u_{\Gamma}|^2 d\sigma \leqslant C \int_{\Omega} |\nabla u|^2 dx$$

for all $u \in \mathcal{V}$, and then we deduce that \mathcal{V} is a Hilbert space in \mathbb{C} with the inner product given by

$$\langle (u, u_{\Gamma}), (v, v_{\Gamma}) \rangle_{\mathcal{V}} = \int_{\Omega} \nabla u \cdot \nabla \overline{v} \, dx + \int_{\Gamma_1} \nabla_{\Gamma} u_{\Gamma} \cdot \nabla_{\Gamma} \overline{v}_{\Gamma} \, d\sigma.$$

Now, we present a definition related with the geometric hypothesis we will assume for the interior boundary.

Definition 1.2. An open, bounded and convex set $U \subset \mathbb{R}^n$, is said to be strongly convex if ∂U is of class C^2 and all the principal curvatures are strictly positive functions on ∂U .

We point out that $U \subset \mathbb{R}^n$ is strongly convex if and only if for all plane $\Pi \subset \mathbb{R}^n$ intersecting U, the curve $\Pi \cap \partial U$ has strictly positive curvature at each point. In particular, a strongly convex set is geometrically strictly convex, in the sense that it has, at each of its boundary points, a supporting hyperplane with exactly one contact point.

We assume that $\Omega = \Omega_0 \setminus \overline{\Omega}_1$, where Ω_1 is strongly convex. Also, without loss of generality, we suppose that $0 \in \Omega_1$ (if it is not the case, we can take $x_0 \in \Omega_1$ and then perform a translation by $-x_0$). Then, we set $\Gamma_k = \partial \Omega_k$ for k = 0, 1.

Now we can define the Carleman weight function we will use in this work. For each $x \in \mathbb{R}^n$, we set

$$\mu(x) = \inf\{\lambda > 0 : x \in \lambda\Omega_1\}.$$
(1.2)

We define for each $x \in \Omega$, $\psi(x) = \mu^2(x)$, and for $\lambda > 0$ we set

$$\theta(x,t) = \frac{e^{\lambda\psi(x)}}{t(T-t)}, \quad \varphi(x,t) = \frac{\alpha - e^{\lambda\psi(x)}}{t(T-t)}, \quad \forall (x,t) \in \Omega \times (0,T),$$
(1.3)

where $\alpha > \|e^{\lambda \psi}\|_{L^{\infty}(\Omega)}$.

In order to guarantee enough regularity of the function μ , we will assume that $\partial \Omega_1$ has a regular parametrization. In order to be explicit, we will ask the following property.

The bijection
$$\Phi: x \in \partial \Omega_1 \mapsto \frac{x}{\|x\|} \in S^{n-1}$$
 is a C^4 diffeomorphism. (1.4)

We recall that Φ is well-defined and is a bijective function thanks to the fact that Ω_1 is convex and it contains the origin.

We point out that the weight functions defined in (1.3) have been previously used to deduce Carleman estimates for transmission problems for wave and Schrödinger equations with Dirichlet boundary conditions, see [4] and [3]. However, to the author's knowledge it is the first time that this function is used to deduce controllability results for Schrödinger equation with dynamic boundary conditions.

Remark 1.3. The function μ defined in (1.2) is called the Minkowski functional of the set Ω_1 . By definition, given $x \in \Omega$, if $\lambda = \lambda(x) > 0$ is such that $x \in \lambda \partial \Omega_1 = \lambda \Gamma_1$ (by convexity, there exists exactly only such λ), then $\mu(x) = \lambda$. This function has the well-known property of, under adequate hypothesis on the open set Ω_1 , defining a norm in \mathbb{R}^n such that their unit ball is Ω_1 .

1.2 Main results

In this section, we give the main results of this article. The first result is a Carleman estimate for a Schrödinger equation with dynamic boundary conditions, whose proof is given in Section 3.

Theorem 1.4. Suppose that $\Omega = \Omega_0 \setminus \overline{\Omega}_1$, where Ω_0 is an open bounded set with C^2 boundary and Ω_1 is an open strongly convex set with boundary $\partial\Omega_1$ satisfying regularity hypothesis (1.4). Let $(\vec{q}_1, \vec{q}_{\Gamma,1}) \in [L^{\infty}(\Omega \times (0,T))]^n \times [L^{\infty}(\Gamma_1 \times (0,T))]^n$ and $(q_0, q_{\Gamma,0}) \in L^{\infty}(\Omega \times (0,T)) \times L^{\infty}(\Gamma_1 \times (0,T))$. Also, assume that δ and d are positive constants satisfying

$$\delta > d. \tag{1.5}$$

Then, there exist constants C, s_0 and λ_0 such that

$$\int_{0}^{T} \int_{\Omega} e^{-2s\varphi} \left(s^{3}\lambda^{4}\theta^{3}|v|^{2} + s\lambda\theta|\nabla v|^{2} + s\lambda^{2}\theta|\nabla\psi\cdot\nabla v|^{2} \right) dxdt
+ \int_{0}^{T} \int_{\Gamma_{1}} e^{-2s\varphi} \left(s^{3}\lambda^{3}\theta^{3}|v_{\Gamma}|^{2} + s\lambda\theta|\partial_{\nu}v|^{2} + s\lambda\theta|\nabla_{\Gamma}v_{\Gamma}|^{2} \right) d\sigma dt
\leqslant C \int_{0}^{T} \int_{\Omega} e^{-2s\varphi}|L(v)|^{2} dxdt + C \int_{0}^{T} \int_{\Gamma_{1}} e^{-2s\varphi}|N(v,v_{\Gamma})|^{2} d\sigma dt
+ Cs\lambda \int_{0}^{T} \int_{\Gamma_{*}} e^{-2s\varphi}\theta|\partial_{\nu}v|^{2} d\sigma dt,$$
(1.6)

for all $\lambda \ge \lambda_0$, $s \ge s_0$ and $(v, v_{\Gamma}) \in L^2(0, T; \mathcal{V})$ where

$$L(v) := i\partial_t v + d\Delta v + \vec{q}_1 \cdot \nabla v + q_0 v \in L^2(\Omega \times (0,T)),$$

$$N(v,v_{\Gamma}) := i\partial_t v - d\partial_\nu v + \delta\Delta_{\Gamma} v_{\Gamma} + \vec{q}_{\Gamma,1} \cdot \nabla_{\Gamma} u_{\Gamma} + q_{\Gamma,0} v_{\Gamma} \in L^2(\Gamma_1 \times (0,T))$$

and $\partial_{\nu} v \in L^2(\partial \Omega \times (0,T))$, with

$$\Gamma_* := \{ x \in \partial\Omega : \partial_\nu \psi(x) \ge 0 \} \subseteq \Gamma_0.$$
(1.7)

Remark 1.5. The hypothesis about the convexity of Ω_1 allows us to define a weight function ψ adapted to the geometry of our problem (see (1.2) and (1.3)): it can be used as a Carleman weight function, and has the particularity that it is constant on Γ_1 , which implies that $\nabla_{\Gamma}\psi = 0$ and $\partial_{\nu}\psi < 0$ in Γ_1 .

Remark 1.6. Hypothesis (1.5) is used in order to estimate the term with $\nabla_{\Gamma} v_{\Gamma}$ on $\Gamma_1 \times (0,T)$ in the left-hand side of the Carleman estimate (1.6) (see inequality (3.20)).

Remark 1.7. We recall that $\partial \Omega = \Gamma_0 \cup \Gamma_1$, where Γ_0 and Γ_1 are connected and closed. In Figure 1, we illustrate an example of the shape of Ω under the geometrical assumption of Theorem 1.4.



Figure 1: Geometric assumptions of Theorem 1.4.

As a direct consequence of the Theorem 1.4, we can obtain a Carleman estimate where the observation is taken in a boundary neighborhood of Γ_* (see for example the region ω in gray in Figure 1).

Corollary 1.8. Let $\omega \subset \Omega$ be an open set such that there exists $\varepsilon > 0$ such that

$$\omega \supset \{x \in \Omega : dist(x, \Gamma_*) \leqslant \varepsilon\},\$$

where Γ_* is given by (1.7). Then, under the assumptions of Theorem 1.4, there exist positive constants C, s_0, λ_0 such that

$$\int_{0}^{T} \int_{\Omega} e^{-2s\varphi} (s^{3}\lambda^{4}\theta^{3}|v|^{2} + s\lambda\theta|\nabla v|^{2}) dxdt$$

$$+ \int_{0}^{T} \int_{\Gamma_{1}} e^{-2s\varphi} (s^{3}\lambda^{3}\theta^{3}|v|^{2} + s\lambda\theta|\nabla_{\Gamma}v_{\Gamma}|^{2} + s\lambda\theta|\partial_{\nu}v|^{2}) d\sigma dt$$

$$\leq C \int_{0}^{T} \int_{\Omega} e^{-2s\varphi} |L(v)|^{2} dxdt + C \int_{0}^{T} \int_{\Gamma_{1}} e^{-2s\varphi} |N(v,v_{\Gamma})|^{2} d\sigma dt$$

$$+ C \int_{0}^{T} \int_{\omega} e^{-2s\varphi} \left(s^{3}\lambda^{4}\theta^{3}|v|^{2} + s\lambda\theta|\nabla v|^{2}\right) dxdt$$
(1.8)

for all $\lambda \ge \lambda_0$ and $s \ge s_0$, and for all pair $(v, v_{\Gamma}) \in L^2(0, T; \mathcal{V})$ satisfying $L(v) \in L^2(\Omega \times (0, T))$, $N(v, v_{\Gamma}) \in L^2(\Gamma_1 \times (0, T))$ and $\partial_{\nu} v \in L^2(\partial \Omega \times (0, T))$.

With Theorem 1.4 and Corollary 1.8 at hand, we derive exact controllability results for Schrödinger equations with dynamic boundary conditions. To this end, we suppose that

$$\vec{q}_1 = d\nabla \pi - i\vec{r}, \quad \vec{q}_{\Gamma,1} = \delta \nabla_{\Gamma} \pi_{\Gamma} - i\vec{r}_{\Gamma}, \tag{1.9}$$

where

$$\begin{aligned} \pi \in L^{\infty}(0,T;W^{3,\infty}(\Omega;\mathbb{R})) \cap W^{1,\infty}(0,T;W^{1,\infty}(\Omega;\mathbb{R})), \\ \pi_{\Gamma} \in L^{\infty}(0,T;W^{3,\infty}(\Gamma_1;\mathbb{R})) \cap W^{1,\infty}(0,T;W^{1,\infty}(\Gamma_1;\mathbb{R})) \end{aligned}$$

with $\pi = \pi_{\Gamma}$ on $\Gamma_1 \times (0, T)$, and

$$\vec{r} \in [L^{\infty}(0,T;W^{2,\infty}(\Omega;\mathbb{R}))]^n, \quad \vec{r}_{\Gamma} \in [L^{\infty}(0,T;W^{2,\infty}(\Gamma_1;\mathbb{R}))]^n$$

with $\vec{r} = \vec{r}_{\Gamma}$ on $\Gamma_1 \times (0,T)$, $\vec{r} \cdot \nu \leq 0$ on $\partial \Omega \times (0,T)$. We also consider that

$$(q_0, q_{\Gamma,0}) \in L^{\infty}(0, T; W^{1,\infty}(\Omega) \times W^{1,\infty}(\Gamma_1)).$$
 (1.10)

We point out that the choice of $(\vec{q}_1, \vec{q}_{\Gamma,1})$, $(q_0, q_{\Gamma,0})$ and its assumptions comes from the wellposedness of system (1.1) and also of its adjoint, which requires more space regularity. For more details about this subject, see Section 2, problem (4.1) and (4.2).

Then, thanks to Theorem 1.4 we obtain

Theorem 1.9. Assume the hypotheses of Theorem 1.4. In addition, assume (1.9) and (1.10). Then, for all initial states $(y_0, y_{\Gamma,0}), (y_T, y_{\Gamma,T}) \in \mathcal{V}'$, there exists a control $h \in L^2(\Gamma_* \times (0,T))$ such that the associated solution (y, y_{Γ}) of (1.1) (defined in the sense of transposition) satisfies

$$(y(T), y_{\Gamma}(T)) = (y_T, y_{\Gamma,T})$$
 in $\Omega \times \Gamma_1$.

On the other hand, as a consequence of Corollary 1.8 we can obtain a controllability result where a (distributed) control is acting on a subdomain of Ω . In order to formulate this result, let (y, y_{Γ}) be a solution of

$$\begin{cases} i\partial_t y + d\Delta y - \vec{q}_1 \cdot \nabla y + q_0 y = \mathbb{1}_{\omega} h, & \text{in } \Omega \times (0, T), \\ i\partial_t y_{\Gamma} - d\partial_{\nu} y + \delta \Delta_{\Gamma} y_{\Gamma} - \vec{q}_{\Gamma,1} \cdot \nabla_{\Gamma} y_{\Gamma} + q_{\Gamma,0} y_{\Gamma} = 0, & \text{on } \Gamma_1 \times (0, T), \\ y = y_{\Gamma}, & \text{on } \Gamma_1 \times (0, T), \\ y = 0, & \text{on } \Gamma_0 \times (0, T), \\ (y, y_{\Gamma})(0) = (y_0, y_{\Gamma,0}), & \text{in } \Omega \times \Gamma_1, \end{cases}$$
(1.11)

with $h \in L^2(\omega \times (0,T))$, with $\omega \subset \Omega$. Then, we have the following result:

Theorem 1.10. Assume the hypotheses of Corollary 1.8. Besides, we consider (1.9) and (1.10). Then, for all T > 0 and $(y_0, y_{\Gamma,0}), (y_T, y_{\Gamma,T}) \in \mathcal{V}'$, there exists a control $h \in L^2(0, T; (H^1_{\Gamma_0}(\omega))')$ such that the associated solution (y, y_{Γ}) of (1.11) (defined in the sense of transposition) satisfies

$$(y(T), y_{\Gamma}(T)) = (y_T, y_{\Gamma,T}), \text{ in } \Omega \times \Gamma_1.$$

The proof of Theorem 1.10 follows the same spirit of the proof of Theorem 1.9. For this reason, in this paper we omit the proof of this result.

Remark 1.11. We point out that some controllability results for conservative Schrödinger equation are obtained from the exact controllability for the wave equation, see for instance [20], [10] and generalized in [25] for conservative systems by using a transmutation control technique. In case of (1.1) with $(\vec{q}_1, \vec{q}_{\Gamma,1}) = (0,0)$ and $(q_0, q_{\Gamma,0}) = (0,0)$, a result can be obtained by applying these arguments but with an additional control acting on a subset of Γ_1 .

The rest of the paper is organized as follows. In Section 2 we prove some results concerning existence and uniqueness of systems like (1.1) and (1.11). In Section 3 we prove the Carleman estimate obtained in Theorem 1.4 and in Section 3.2 we prove the Corollary 1.8 by using a suitable cut-off function. Finally, in Section 4 we prove the Theorem 1.9, which is equivalent to prove the so-called observability inequality associated to the adjoint system.

2 Existence and uniqueness of solutions

In this section, we provide existence and uniqueness results of solutions for Schrödinger equation with dynamic boundary conditions. We also give a hidden regularity property and define solutions in the sense of transposition for such systems.

Without loss of generality, it is sufficient to analyze the problem (1.1) considering (1.9) with $(\pi, \pi_{\Gamma}) = (0, 0)$. That is to say, from now on, we will consider

$$\begin{cases} i\partial_t y + d\Delta y + i\vec{r} \cdot \nabla y + q_0 y = 0, & \text{in } \Omega \times (0, T), \\ i\partial_t y_{\Gamma} - d\partial_{\nu} y + \delta \Delta_{\Gamma} y_{\Gamma} + i\vec{r}_{\Gamma} \cdot \nabla_{\Gamma} y_{\Gamma} + q_{\Gamma,0} y_{\Gamma} = 0, & \text{on } \Gamma_1 \times (0, T), \\ y = y_{\Gamma}, & \text{on } \Gamma_1 \times (0, T), \\ y = \mathbb{1}_{\Gamma_*} h, & \text{on } \Gamma_0 \times (0, T), \\ (y(0), y_{\Gamma}(0)) = (y_0, y_{\Gamma,0}), & \text{in } \Omega \times \Gamma_1, \end{cases}$$
(2.1)

with \vec{r} and \vec{r}_{Γ} real valued. Otherwise, we apply the change of variables $\tilde{y}(x,t) = e^{-\pi(x,t)/2}y(x,t)$ in $\Omega \times (0,T)$ and $\tilde{y}_{\Gamma}(x,t) = e^{-\pi(x,t)/2}y_{\Gamma}(x,t)$ on $\Gamma_1 \times (0,T)$, where (y,y_{Γ}) is a solution of (1.1). Then, arguing as [18, Appendix A], the new variable $(\tilde{y}, \tilde{y}_{\Gamma})$ satisfies a problem of the form (2.1).

2.1 Existence and regularity of solutions

Given $d, \delta > 0$, we consider the problem

$$\begin{cases} \partial_t u - di\Delta u + \vec{\rho}_1 \cdot \nabla u + \rho_0 u = g, & \text{in } \Omega \times (0, T), \\ \partial_t u + di\partial_\nu u - \delta i\Delta_\Gamma u_\Gamma + \vec{\rho}_{\Gamma,1} \cdot \nabla_\Gamma u_\Gamma + \rho_{\Gamma,0} u_\Gamma = g_\Gamma, & \text{on } \Gamma_1 \times (0, T), \\ u = u_\Gamma, & \text{on } \Gamma_1 \times (0, T), \\ u = 0, & \text{on } \Gamma_0 \times (0, t), \\ (u(0), u_\Gamma(0)) = (u_0, u_{\Gamma,0}), & \text{in } \Omega \times \Gamma_1, \end{cases}$$
(2.2)

where $(\vec{\rho}_1, \vec{\rho}_{\Gamma,1})$ is real valued and $(\rho_0, \rho_{\Gamma,0}), (g, g_{\Gamma})$ are complex-valued. The next results establish L^2 and H^1 estimates of the weak solutions of the problem (2.2), respectively. The proofs, as usual, are based on energy estimates and density arguments.

Proposition 2.1. Suppose that $(u_0, u_{\Gamma,0}) \in \mathcal{H}$, $\vec{\rho_1} \in [L^{\infty}(0, T; W^{1,\infty}(\Omega; \mathbb{R}))]^n$. Moreover, assume that $\vec{\rho}_{\Gamma,1} \in [L^{\infty}(0, T; W^{1,\infty}(\Gamma_1; \mathbb{R}))]^n$, with $\vec{\rho_1} = \vec{\rho}_{\Gamma,1}$ on $\Gamma_1 \times (0, T)$, $(\rho_0, \rho_{\Gamma,0}) \in L^{\infty}(0, T; L^{\infty}(\Omega \times \mathbb{R}))$

 (Γ_1) , and $(g, g_{\Gamma}) \in L^1(0, T; \mathcal{H})$. Then, there exists a positive constant C such that the weak solution $(u, u_{\Gamma}) \in C^0([0, T]; \mathcal{H})$ of (2.2) satisfies

$$\max_{t \in [0,T]} \|(u, u_{\Gamma})(t)\|_{\mathcal{H}}^2 \leqslant C\left(\|(g, g_{\Gamma})\|_{L^1(0,T;\mathcal{H})}^2 + \|(u_0, u_{\Gamma,0})\|_{\mathcal{H}}^2\right).$$
(2.3)

Proof. Firstly, we multiply the first equation of (2.2) by \overline{u} and integrate in Ω . Secondly, we multiply the second equation by \overline{u}_{Γ} and integrate on $\Gamma_1 \times (0, T)$. Next, we add these identities and take the real part on the obtained equation. This yields

$$\Re \int_{\Omega} \overline{u} \partial_t u dx + d\Im \int_{\Omega} \overline{u} \Delta u dx + \Re \int_{\Omega} \overline{u} (\vec{\rho}_1 \cdot \nabla u) dx + \Re \int_{\Omega} \rho_0 |u|^2 dx + \Re \int_{\Gamma_1} \overline{u}_{\Gamma} \partial_t u_{\Gamma} d\sigma - d\Im \int_{\Gamma_1} \overline{u}_{\Gamma} \partial_{\nu} u d\sigma + d\Im \int_{\Gamma_1} \overline{u}_{\Gamma} \Delta_{\Gamma} u_{\Gamma} d\sigma + \Re \int_{\Gamma_1} \overline{u}_{\Gamma} (\vec{\rho}_{\Gamma,1} \cdot \nabla_{\Gamma} u_{\Gamma}) d\sigma + \Re \int_{\Gamma_1} \rho_{\Gamma,0} |u_{\Gamma}|^2 d\sigma = \Re \int_{\Omega} \overline{u} g dx + \Re \int_{\Gamma_1} \overline{u}_{\Gamma} g_{\Gamma} d\sigma,$$

$$(2.4)$$

a.e. in (0, T). We notice that,

$$\Re \int_{\Omega} \overline{u} \partial_t u dx + \Re \int_{\Gamma_1} \overline{u}_{\Gamma} \partial_t u_{\Gamma} d\sigma = \frac{d}{dt} \left(\frac{1}{2} \int_{\Omega} |u|^2 dx + \frac{1}{2} \int_{\Gamma_1} |u_{\Gamma}|^2 d\sigma \right).$$
(2.5)

Moreover, integration by parts and surface divergence theorem implies that

$$d\Im \int_{\Omega} \overline{u} \Delta u dx - d\Im \int_{\Gamma_1} \overline{u}_{\Gamma} \partial_{\nu} u d\sigma + \delta\Im \int_{\Gamma_1} \overline{u}_{\Gamma} \Delta_{\Gamma} u_{\Gamma} d\sigma = 0, \qquad (2.6)$$

where we have used that $u = u_{\Gamma}$ on $\Gamma_1 \times (0, T)$. In addition,

$$\Re \int_{\Omega} \overline{u}(\vec{\rho}_{1} \cdot \nabla u) dx + \Re \int_{\Gamma_{1}} \overline{u}_{\Gamma}(\vec{\rho}_{\Gamma,1} \cdot \nabla_{\Gamma} u_{\Gamma}) d\sigma$$

$$= -\frac{1}{2} \int_{\Omega} \operatorname{div}(\vec{\rho}_{1}) |u|^{2} dx - \frac{1}{2} \int_{\Gamma_{1}} \operatorname{div}_{\Gamma}(\vec{\rho}_{\Gamma,1}) |u_{\Gamma}|^{2} d\sigma + \frac{1}{2} \int_{\Gamma_{1}} (\vec{\rho}_{1} \cdot \nu) |u|^{2} d\sigma.$$
(2.7)

Substituting (2.5), (2.6) and (2.7) into (2.4) and applying Cauchy-Scharwz inequality we deduce that

$$\frac{d}{dt}\|(u,u_{\Gamma})(t)\|_{\mathcal{H}}^2 \leqslant C\|(u,u_{\Gamma})(t)\|_{\mathcal{H}}^2 + C\|(u,u_{\Gamma})(t)\|_{\mathcal{H}}\|(g,g_{\Gamma})(t)\|_{\mathcal{H}}$$

Then, by Gronwall's and Holder's inequalities, we obtain (2.3). This ends the proof of the Proposition 2.1. $\hfill \Box$

Under additional assumptions on $(\vec{\rho}_1, \vec{\rho}_{\Gamma,1})$ and $(\rho_0, \rho_{\Gamma,0})$, we get the following result.

Proposition 2.2. Suppose that $(u_0, u_{\Gamma,0}) \in \mathcal{V}$ and $(g, g_{\Gamma}) \in L^1(0, T; \mathcal{V})$. Moreover, assume that

$$\vec{\rho}_1 \in [L^{\infty}(0,T;W^{1,\infty}(\Omega;\mathbb{R}))]^n, \quad \vec{\rho}_{\Gamma,1} \in [L^{\infty}(0,T;W^{1,\infty}(\Gamma_1;\mathbb{R}))]^n,$$
(2.8)

with $\vec{\rho_1} = \vec{\rho_{\Gamma,1}}$, on $\Gamma_1 \times (0,T)$ and $\vec{\rho_1} \cdot \nu \leq 0$ on $\partial \Omega \times (0,T)$. We also assume that

$$\rho_0 \in L^{\infty}(0,T; W^{1,\infty}(\Omega)), \quad \rho_{\Gamma,0} \in L^{\infty}(0,T; W^{1,\infty}(\Gamma_1)),$$

with $\rho_0 = \rho_{\Gamma,0}$ on $\Gamma_1 \times (0,T)$. Then, the weak solution of (2.2) belongs to $C^0([0,T]; \mathcal{V})$. Moreover, there exists C > 0 such that

$$\max_{t \in [0,T]} \|(u, u_{\Gamma})\|_{\mathcal{V}}^2 \leqslant C\left(\|(g, g_{\Gamma})\|_{L^1(0,T;\mathcal{V})}^2 + \|(u_0, u_{\Gamma,0})\|_{\mathcal{V}}^2\right).$$
(2.9)

Proof. We multiply the first equation of (2.2) by $-d\Delta \overline{u}$ and integrate in $\Omega \times (0, T)$. Next, we multiply the second equation of (2.2) by $(-\delta \Delta_{\Gamma} \overline{u}_{\Gamma} + d\partial_{\nu} \overline{u})$ and integrate on $\Gamma_1 \times (0, T)$. Then, adding these identities and taking the real part we have

$$-d\Re \int_{\Omega} \partial_{t} u \Delta \overline{u} dx - d\Re \int_{\Omega} (\vec{\rho}_{1} \cdot \nabla u) \Delta \overline{u} dx - d\Re \int_{\Omega} p u \Delta \overline{u} dx + \Re \int_{\Gamma_{1}} \partial_{t} u_{\Gamma} (-\delta \Delta_{\Gamma} \overline{u}_{\Gamma} + d \partial_{\nu} \overline{u}) d\sigma + \Re \int_{\Gamma_{1}} (\vec{\rho}_{1} \cdot \nabla_{\Gamma} u_{\Gamma}) (-\delta \Delta_{\Gamma} \overline{u}_{\Gamma} + d \partial_{\nu} \overline{u}) d\sigma + \Re \int_{\Gamma_{1}} p_{\Gamma} u_{\Gamma} (-\delta \Delta_{\Gamma} \overline{u}_{\Gamma} + d \partial_{\nu} \overline{u}) d\sigma = -d\Re \int_{\Omega} g \Delta \overline{u} dx + \Re \int_{\Gamma_{1}} g_{\Gamma} (-\delta \Delta_{\Gamma} \overline{u}_{\Gamma} + d \partial_{\nu} \overline{u}) d\sigma.$$

$$(2.10)$$

Integration by parts and surface divergence theorem imply that

$$-d\Re \int_{\Omega} \partial_t u \Delta \overline{u} dx + \Re \int_{\Gamma_1} \partial_t u_{\Gamma} (-\delta \Delta_{\Gamma} \overline{u}_{\Gamma} + d\partial_{\nu} \overline{u}) d\sigma$$

$$= \frac{d}{dt} \left(\frac{1}{2} d \int_{\Omega} |\nabla u|^2 dx + \frac{1}{2} \delta \int_{\Gamma_1} |\nabla_{\Gamma} u_{\Gamma}|^2 d\sigma \right).$$
(2.11)

On the other hand,

$$\begin{split} \Lambda &:= -d\Re \int_{\Omega} (\vec{\rho}_{1} \cdot \nabla u) \Delta \overline{u} dx + \Re \int_{\Gamma_{1}} (\vec{\rho}_{\Gamma,1} \cdot \nabla_{\Gamma} u_{\Gamma}) (-\delta \Delta_{\Gamma} \overline{u}_{\Gamma} + d\partial_{\nu} \overline{u}) d\sigma \\ &= d\Re \int_{\Omega} \nabla (\vec{\rho}_{1} \cdot \nabla u) \cdot \nabla \overline{u} dx + \frac{1}{2} d \int_{\partial \Omega} (\vec{\rho}_{1} \cdot \nabla u) \partial_{\nu} \overline{u} d\sigma \\ &+ \delta \Re \int_{\Gamma_{1}} \nabla_{\Gamma} (\vec{\rho}_{\Gamma,1} \cdot \nabla_{\Gamma} u_{\Gamma}) \cdot \nabla_{\Gamma} \overline{u}_{\Gamma} d\sigma + d\Re \int_{\Gamma_{1}} (\vec{\rho}_{\Gamma,1} \cdot \nabla_{\Gamma} u_{\Gamma}) \partial_{\nu} \overline{u} d\sigma. \end{split}$$

Since

$$\begin{split} \nabla(\vec{\rho_1}\cdot\nabla u)\cdot\nabla\overline{u} &= \vec{\nabla}\rho_1(\nabla u,\nabla\overline{u}) + \frac{1}{2}\vec{\rho_1}\cdot\nabla(|\nabla u|^2),\\ \nabla_{\Gamma}(\vec{\rho_{\Gamma,1}}\cdot\nabla_{\Gamma}u_{\Gamma})\cdot\nabla_{\Gamma}\overline{u}_{\Gamma} &= \vec{\nabla}_{\Gamma}\vec{\rho_{\Gamma,1}}(\nabla_{\Gamma}u_{\Gamma},\nabla_{\Gamma}\overline{u}_{\Gamma}) + \frac{1}{2}\vec{\rho_{\Gamma,1}}\cdot\nabla_{\Gamma}(|\nabla_{\Gamma}u_{\Gamma}|^2), \end{split}$$

and that $\nabla u = \nabla_{\Gamma} u + (\partial_{\nu} u) \nu$ on $\Gamma_1 \times (0, T)$, we obtain

$$\begin{split} \Lambda = d\Re \int_{\Omega} \vec{\nabla} \vec{\rho}_{1} (\nabla u, \nabla \overline{u}) dx + \delta \Re \int_{\Gamma_{1}} \vec{\nabla} \vec{\rho}_{\Gamma,1} (\nabla_{\Gamma} u_{\Gamma}, \nabla_{\Gamma} \overline{u}_{\Gamma}) d\sigma \\ &- \frac{1}{2} d \int_{\Omega} \operatorname{div}(\vec{\rho}_{1}) |\nabla u|^{2} - \frac{1}{2} \delta \int_{\Gamma_{1}} \operatorname{div}(\vec{\rho}_{\Gamma,1}) |\nabla_{\Gamma} u_{\Gamma}|^{2} d\sigma \\ &- \frac{1}{2} d \int_{\partial\Omega} (\vec{\rho}_{1} \cdot \nu) |\partial_{\nu} u|^{2} d\sigma + \frac{1}{2} d \int_{\Gamma_{1}} (\vec{\rho}_{1} \cdot \nu) |\nabla_{\Gamma} u_{\Gamma}|^{2} d\sigma, \end{split}$$
(2.12)

where we have integrated by parts and used the fact that $\vec{\rho}_1 = \vec{\rho}_{\Gamma,1}$ on $\Gamma_1 \times (0,T)$. In addition, since $\rho_0 = \rho_{\Gamma,0}$ on $\Gamma_1 \times (0,T)$, we have

$$-d\Re \int_{\Omega} pu\Delta \overline{u}dx + \Re \int_{\Gamma_{1}} p_{\Gamma}u_{\Gamma}(-\delta\Delta_{\Gamma}\overline{u}_{\Gamma} + d\partial_{\nu}\overline{u})d\sigma$$

$$=d\int_{\Omega} \Re(p)|\nabla u|^{2} + \delta \int_{\Gamma_{1}} \Re(p_{\Gamma})|\nabla_{\Gamma}u_{\Gamma}|^{2}d\sigma$$

$$+d\Re \int_{\Omega} u\nabla p \cdot \nabla \overline{u}dx + \delta\Re \int_{\Gamma_{1}} u_{\Gamma}\nabla_{\Gamma}p_{\Gamma} \cdot \nabla_{\Gamma}\overline{u}_{\Gamma}d\sigma.$$

(2.13)

Now, since g = 0 on $\Gamma_0 \times (0, T)$ and $g = g_{\Gamma}$ on $\Gamma_1 \times (0, T)$, we deduce that

$$-d\Re \int_{\Omega} g\Delta \overline{u} dx + \Re \int_{\Gamma_{1}} g_{\Gamma}(-\delta \Delta_{\Gamma} \overline{u}_{\Gamma} + d\partial_{\nu} \overline{u}) d\sigma$$

=
$$d\Re \int_{\Omega} \nabla g \cdot \nabla \overline{u} dx + \delta \Re \int_{\Gamma_{1}} \nabla_{\Gamma} g \cdot \nabla_{\Gamma} \overline{u}_{\Gamma} d\sigma.$$
 (2.14)

Then, substituting (2.11), (2.12), (2.13), (2.14) into (2.10) and using the fact that $\vec{\rho_1} \cdot \nu \leq 0$ on $\partial \Omega \times (0,T)$, we can assert that

$$\frac{d}{dt} \| (u, u_{\Gamma})(t) \|_{\mathcal{V}}^2 \leq C \| (u, u_{\Gamma})(t) \|_{\mathcal{V}}^2 + C \| (u, u_{\Gamma})(t) \|_{\mathcal{V}} \| (g, g_{\Gamma})(t) \|_{\mathcal{V}}$$

Then, applying Gronwall's and Hölder's inequalities we easily deduce (2.9) and the proof of the Proposition 2.2 is finished. $\hfill \Box$

Remark 2.3. It is also possible to obtain (2.9) considering regularity assumptions on $(\vec{\rho}_1, \vec{\rho}_{\Gamma,1})$ and $(\rho_0, \rho_{\Gamma,0})$ in the time variable. However, in order to simplify the presentation of the rest of the results in this paper, we do not consider these situations.

2.2 Hidden regularity

In this section, we devote to deduce a hidden regularity result for solutions of the Schrödinger equation with dynamic boundary conditions. For this purposes, we start giving the following identity.

Proposition 2.4. Suppose that $\vec{q} \in C^2(\overline{\Omega} \times [0,T]; \mathbb{R}^n)$. Moreover, consider $\vec{\rho_1} \in [L^{\infty}(0,T; W^{1,\infty}(\Omega; \mathbb{R}))]^n$, $\vec{\rho_{\Gamma,1}} \in [L^{\infty}(0,T; W^{1,\infty}(\Gamma_1; \mathbb{R}))]^n$ and $\rho_0 \in L^{\infty}(\Omega \times (0,T))$, $\rho_{\Gamma,0} \in L^{\infty}(\Gamma_1 \times (0,T))$. Then, for each $d, \delta > 0$, $(u_0, u_{\Gamma,0}) \in \mathcal{V}$ and $(g, g_{\Gamma}) \in L^1(0,T; \mathcal{V})$, the solution (u, u_{Γ}) of (2.2) satisfies

$$\frac{1}{2}d\int_0^T \int_{\partial\Omega} (\vec{q} \cdot \nu) |\partial_\nu u|^2 dx dt = X + Y + F, \qquad (2.15)$$

where X, Y and F are given by

$$\begin{split} X &= -\frac{1}{2}\delta\int_{0}^{T}\int_{\Gamma_{1}}(\vec{q}\cdot\nu)|\nabla_{\Gamma}u_{\Gamma}|^{2}\,d\sigma dt - \frac{1}{2}\Im\int_{0}^{T}\int_{\Gamma_{1}}(\vec{q}\cdot\nu)\rho_{\Gamma,0}|u_{\Gamma}|^{2}\,d\sigma dt \\ &+ \frac{1}{2}d\int_{0}^{T}\int_{\Omega}\vec{\nabla}\vec{q}(\nabla u,\nabla \overline{u})\,dx dt + \frac{1}{2}d\int_{0}^{T}\int_{\Omega}div(\vec{q})|\nabla u|^{2}\,dx dt \\ &+ \frac{1}{2}\delta\int_{0}^{T}\int_{\Gamma_{1}}\vec{\nabla}_{\Gamma}\vec{q}(\nabla_{\Gamma}u_{\Gamma},\nabla_{\Gamma}\overline{u}_{\Gamma})d\sigma dt + \frac{1}{2}\delta\int_{0}^{T}\int_{\Gamma_{1}}div(\vec{q})|\nabla_{\Gamma}u_{\Gamma}|^{2}\,d\sigma dt \\ &+ \Im\int_{0}^{T}\int_{\Omega}(\vec{\rho}_{1}\cdot\nabla u)(\vec{q}\cdot\nabla \vec{u})\,dx dt + \frac{1}{2}\Im\int_{0}^{T}\int_{\Omega}\rho_{0}div(\vec{q})|u|^{2}\,dx dt \\ &+ \Im\int_{0}^{T}\int_{\Gamma_{1}}(\vec{\rho}_{\Gamma,1}\cdot\nabla_{\Gamma}u_{\Gamma})(\vec{q}\cdot\nabla_{\Gamma}u_{\Gamma})\,d\sigma dt + \frac{1}{2}\Im\int_{0}^{T}\int_{\Gamma_{1}}\rho_{\Gamma,0}div(\vec{q})|u_{\Gamma}|^{2}\,d\sigma dt, \end{split}$$

$$\begin{split} Y = &\frac{1}{2} \Im \int_{0}^{T} \int_{\Omega} \overline{u} \nabla u \cdot \partial_{t} \vec{q} \, dx dt - \frac{1}{2} \Im \int_{\Omega} \overline{u} \nabla u \cdot \vec{q} \Big|_{0}^{T} dx \\ &- \frac{1}{2} d\Re \int_{0}^{T} \int_{\Gamma_{1}} (\vec{q} \cdot \nu) \overline{u}_{\Gamma} \partial_{\nu} u \, d\sigma dt - \frac{1}{2} \Im \int_{0}^{T} \int_{\Gamma_{1}} (\vec{q} \cdot \nabla_{\Gamma} \nu) (\vec{\rho}_{\Gamma,1} \cdot u_{\Gamma}) \overline{u}_{\Gamma} \, d\sigma dt \\ &+ \frac{1}{2} \Im \int_{0}^{T} \int_{\Gamma_{1}} (\nabla_{\Gamma} u_{\Gamma} \cdot \partial_{t} \vec{q}) \overline{u}_{\Gamma} \, d\sigma dt - \frac{1}{2} \Im \int_{\Gamma_{1}} (\vec{q} \cdot \nabla_{\Gamma} u_{\Gamma}) \overline{u}_{\Gamma} \Big|_{0}^{T} d\sigma \\ &+ \frac{1}{2} d\Re \int_{0}^{T} \int_{\Omega} \overline{u} \nabla (div(\vec{q})) \cdot \nabla u \, dx dt + \frac{1}{2} d\Re \int_{0}^{T} \int_{\Gamma_{1}} (\partial_{\nu} u) \vec{q} \cdot \nabla_{\Gamma} u_{\Gamma} \, d\sigma dt \\ &+ \frac{1}{2} \delta \Re \int_{0}^{T} \int_{\Gamma_{1}} \nabla_{\Gamma} (div(\vec{q}) \cdot \nabla_{\Gamma} u_{\Gamma}) \overline{u}_{\Gamma} \, d\sigma dt + \frac{1}{2} \Im \int_{0}^{T} \int_{\Omega} div(\vec{q}) (\vec{\rho} \cdot \nabla u) \overline{u} \, dx dt \\ &+ \Im \int_{0}^{T} \int_{\Omega} \rho_{0} (\vec{q} \cdot \nabla \overline{u}) u \, dx dt + \frac{1}{2} \int_{0}^{T} \int_{\Gamma_{1}} div(\vec{q}) (\vec{\rho}_{\Gamma,1} \cdot \nabla_{\Gamma} u_{\Gamma}) \overline{u}_{\Gamma} \, d\sigma dt \\ &+ \Im \int_{0}^{T} \int_{\Gamma_{1}} \rho_{\Gamma,0} (\vec{q} \cdot \nabla_{\Gamma} \overline{u}_{\Gamma}) u_{\Gamma} \, d\sigma dt, \end{split}$$

and

$$\begin{split} F = &\frac{1}{2}\Im\int_{0}^{T}\int_{\Gamma_{1}}(\vec{q}\cdot\nu)\overline{u}_{\Gamma}g_{\Gamma}\,d\sigma dt - \Im\int_{0}^{T}\int_{\Omega}g\left(\vec{q}\cdot\nabla\overline{u} + \frac{1}{2}div(\vec{q})\overline{u}\right)\,dxdt \\ &-\Im\int_{0}^{T}\int_{\Gamma_{1}}g_{\Gamma}\left(\vec{q}\cdot\nabla_{\Gamma}u_{\Gamma} + \frac{1}{2}div(\vec{q})\overline{u}_{\Gamma}\right)\,d\sigma dt. \end{split}$$

Proof. The proof of the Proposition 2.4 relies on the use of classical ideas concerning multiplier techniques. That is to say, we multiply the first equation of (2.2) by $-\left(\vec{q}\cdot\nabla\overline{u}+\frac{1}{2}\operatorname{div}(\vec{q})\overline{u}\right)$ and we integrate in $\Omega \times (0,T)$; similarly, we multiply the second one by $-\left(\vec{q}\cdot\nabla_{\Gamma}\overline{u}_{\Gamma}+\frac{1}{2}\operatorname{div}(\vec{q})\overline{u}_{\Gamma}\right)$ and the result is integrated on $\Gamma_1 \times (0,T)$. Then we add the above identities and take the imaginary part. Contrary to the case of the homogeneous Dirichlet boundary conditions (see for instance [22]), in our case several new boundary terms defined on $\Gamma_1 \times (0,T)$ appear. To deal with them, we use the surface divergence theorem and the second equation of (2.2) to estimate $\partial_t u_{\Gamma}$ on $\Gamma_1 \times (0,T)$. The rest of the proof runs as in [21] and [22] for wave and Schrödinger equation with Dirichlet boundary conditions, respectively. We omit the details.

With this estimate at hand, we have the next result of hidden regularity for the Schrödinger equation with dynamic boundary conditions:

Proposition 2.5. Consider the same assumptions of Proposition 2.2. Then, the derivative $\partial_{\nu} u$ of solution of (2.2) belongs to $L^2(\partial\Omega \times (0,T))$. Moreover, there exists a positive constant C such that

$$\int_{0}^{T} \int_{\partial \Omega} |\partial_{\nu} u|^{2} d\sigma dt \leq C \left(\|(g, g_{\Gamma})\|_{L^{1}(0,T;\mathcal{V})}^{2} + \|(u_{0}, u_{\Gamma,0})\|_{\mathcal{V}}^{2} \right).$$
(2.16)

Proof. We start choosing a smooth vector field $\vec{q} \in C^2(\overline{\Omega}, \mathbb{R}^n)$ such that $q = \nu$ on $\partial\Omega$. See, for instance, [21] for the proof of the existence of such a vector field. Then, by Proposition 2.4, equation (2.15) reads

$$\frac{1}{2}d\|\partial_{\nu}u\|_{L^{2}(0,T;L^{2}(\partial\Omega))}^{2} = X + Y + F.$$
(2.17)

Since (p, p_{Γ}) and \vec{q} and its derivatives are bounded in L^{∞} , we have that

$$X \| \leq C \| (u, u_{\Gamma}) \|_{C^{0}([0,T];\mathcal{V})} \leq C \left(\| (u_{0}, u_{\Gamma,0}) \|_{\mathcal{V}}^{2} + C \| (g, g_{\Gamma}) \|_{L^{1}(0,T;\mathcal{V})}^{2} \right),$$
(2.18)

where we have applied the Proposition 2.2 in the last inequality. In the same manner, by Young's inequality, Y can be estimated as

$$|Y| \leq C \|(u_0, u_{\Gamma,0})\|_{\mathcal{V}}^2 + \|(g, g_{\Gamma})\|_{L^1(0,T;\mathcal{V})}^2 + \frac{1}{4}d\|\partial_{\nu}u\|_{L^2(0,T;L^2(\partial\Omega))}^2.$$
(2.19)

Moreover, by Cauchy-Scharwz inequality we can assert that

|

$$|F| \leq C \|(u_0, u_{\Gamma,0})\|_{\mathcal{V}}^2 + C \|(g, g_{\Gamma})\|_{L^1(0,T;\mathcal{V})}^2.$$
(2.20)

Substituting (2.18), (2.19) and (2.20) into (2.17), we obtain (2.16). This finishes the proof of the Proposition 2.5. \Box

2.3 Solutions by transposition

To define solutions of the Schrödinger equation with dynamic boundary condition and boundary control (2.1), we consider the auxiliary system given by

$$\begin{cases} i\partial_t \phi + d\Delta \phi + i\vec{r} \cdot \nabla \phi + q\phi = f, & \text{in } \Omega \times (0, T), \\ i\partial_t \phi_{\Gamma} - d\partial_{\nu} \phi + \delta \Delta_{\Gamma} \phi_{\Gamma} + i\vec{r}_{\Gamma} \cdot \nabla_{\Gamma} \phi_{\Gamma} + q_{\Gamma} \phi_{\Gamma} = f_{\Gamma}, & \text{on } \Gamma_1 \times (0, T), \\ \phi = \phi_{\Gamma}, & \text{on } \Gamma_1 \times (0, T), \\ \phi = 0, & \text{on } \Gamma_0 \times (0, T), \\ (\phi, \phi_{\Gamma})(T, \cdot) = (\phi_T, \phi_{\Gamma, T}), & \text{in } \Omega \times \Gamma_1, \end{cases}$$
(2.21)

where the pair (q, q_{Γ}) is given by

$$q = i \operatorname{div}(\vec{r}) + \overline{q}_0, \quad q_{\Gamma} = i \operatorname{div}_{\Gamma}(\vec{r}_{\Gamma}) - i \vec{r} \cdot \nu + \overline{q}_{\Gamma,0}.$$

$$(2.22)$$

According to Proposition 2.2 we have that, for each $(\phi_T, \phi_{\Gamma,T}) \in \mathcal{V}$ and $(f, f_{\Gamma}) \in L^1(0, T; \mathcal{V})$, system (2.21) posses a unique solution

$$(\phi, \phi_{\Gamma}) \in C([0, T]; \mathcal{V}). \tag{2.23}$$

We can now define a solution by transposition.

Definition 2.6. Given $(y_0, y_{\Gamma 0}) \in \mathcal{V}'$ and a control $h \in L^2(\Gamma_* \times (0,T))$, we will say that the pair $(y, y_{\Gamma}) \in C([0,T]; \mathcal{V}')$ is a solution of system (1.1) in the sense of transposition if the following identity holds

$$\int_{0}^{T} \langle (y, y_{\Gamma}), (f, f_{\Gamma}) \rangle_{\mathcal{V}', \mathcal{V}} dt = d \int_{0}^{T} \int_{\Gamma_{0}} \mathbb{1}_{\Gamma_{*}} h \overline{\partial_{\nu} \phi} d\sigma dt + i \langle (y_{0}, y_{\Gamma_{0}}), (\phi, \phi_{\Gamma})(0, \cdot) \rangle_{\mathcal{V}', \mathcal{V}}$$
(2.24)

for each $(f, f_{\Gamma}) \in L^1(0, T; \mathcal{V})$, where (ϕ, ϕ_{Γ}) is the solution of (2.21) with $(\phi_T, \phi_{\Gamma,T}) = (0, 0)$.

Theorem 2.7. For each $(y_0, y_{\Gamma,0}) \in \mathcal{V}'$ and $h \in L^2(\Gamma_* \times (0,T))$, system (1.1) has a unique solution $(y, y_{\Gamma}) \in C([0,T]; \mathcal{V}')$. Moreover, there exists C > 0 such that

$$\|(y,y_{\Gamma})\|_{L^{\infty}(0,T;\mathcal{V}')} \leqslant C \left(\|(y_0,y_{\Gamma,0})\|_{\mathcal{V}'} + \|h\|_{L^2(\Gamma_* \times (0,T))}\right).$$

Proof. Given $(y_0, y_{\Gamma_0}) \in \mathcal{V}'$ and $h \in L^2(\Gamma_* \times (0, T))$, from Proposition 2.5 and (2.23), the linear functional defined by

is well defined and continuous. By duality, there exists $(y, y_{\Gamma}) \in L^{\infty}(0, T; \mathcal{V}')$ such that we have (2.24). By classical density arguments we get that $(y, y_{\Gamma}) \in C([0, T]; \mathcal{V}')$.

Now we will consider the problem of exact controllability. By linearity, it is enough to study the case of null initial conditions. It is direct to see that h is a control for system (1.1) driving $(y, y_{\Gamma}) = (0, 0)$ to $(y(T, \cdot), y_{\Gamma}(T, \cdot)) \in \mathcal{V}'$ if

$$d\int_0^T \int_{\Gamma_0} \mathbb{1}_{\Gamma_*} h \overline{\partial_\nu \phi} d\sigma dt = i \left\langle (y(T, \cdot), y_\Gamma(T, \cdot)), (\phi_T, \phi_{\Gamma, T}) \right\rangle_{\mathcal{V}', \mathcal{V}}$$
(2.25)

for each $(\phi_T, \phi_{\Gamma,T}) \in \mathcal{V}$, where (ϕ, ϕ_{Γ}) is the solution of (2.21) with $(f, f_{\Gamma}) = (0, 0)$. Therefore we get, by classical abstract results (see for instance [31]), the following characterization.

Proposition 2.8. System (1.1) is exactly controllable in time T > 0 if and only if there exists a constant C > 0 such that

$$\|(\phi_T, \phi_{\Gamma,T})\|_{\mathcal{V}}^2 \leqslant C \int_0^T \int_{\Gamma_*} |\partial_\nu \phi|^2 d\sigma dt, \qquad (2.26)$$

for all $(\phi_T, \phi_{\Gamma,T}) \in \mathcal{V}$.

3 Proofs of Theorem 1.4 and Corollary 1.8

In order to prove these results, we point out that it is sufficient to prove inequalities (1.6) and (1.8) for functions $(v, v_{\Gamma}) \in C^{\infty}((\overline{\Omega} \times \Gamma_1) \times [0, T])$ which satisfies v = 0 on $\Gamma_0 \times (0, T)$ and $v = v_{\Gamma}$ on $\Gamma_1 \times (0, T)$ and argue by density as usual. Then, due to the regularity of each function and the fact that $v = v_{\Gamma}$, from now on we shall write v instead of v_{Γ} on $\partial \Omega \times (0, T)$.

3.1 Proof of Theorem 1.4

Here we will prove the Carleman estimate of Theorem 1.4. In order to do that, we shall follow the classical conjugation for suitable operators involving the Schrödinger equation with dynamic boundary conditions. However, contrary to the case of Dirichlet boundary conditions, several additional boundary terms arise from the dynamic boundary conditions of Wentzell type. To treat them, we have to repeat some steps from [3] in a modified form.

In order to simplify the presentation, the proof of Theorem 1.4 is done in several steps.

• Step 1: Setting.

In this step, we define the operators to conjugate in $\Omega \times (0,T)$ and on $\Gamma_1 \times (0,T)$. To do this, we write $v = e^{s\varphi}w$ and compute $Pw = e^{-s\varphi}L(e^{s\varphi}w)$ in $\Omega \times (0,T)$ and $Qw = e^{-s\varphi}N(e^{s\varphi}w)$ on $\Gamma_1 \times (0,T)$.

Firstly, we write the operator P as follows

$$Pw = P_1w + P_2w + Rw, \quad \text{in } \Omega \times (0,T), \tag{3.1}$$

where the operators P_1 , P_2 and R are given by

$$P_1w = ds^2 |\nabla \varphi|^2 w + d\Delta w + i\partial_t w, \quad P_2w = ds\Delta \varphi w + 2ds\nabla \varphi \cdot \nabla w + is\partial_t \varphi w,$$

and

$$Rw = \vec{q_1} \cdot \nabla w + s \nabla \varphi \cdot \vec{q_1} w + q_0 w.$$

Secondly, we write

$$Qw = Q_1w + Q_2w + R_{\Gamma}w, \text{ on } \Gamma_1 \times (0,T), \qquad (3.2)$$

where Q_1, Q_2 and R_{Γ} are given by

$$Q_1w = \delta\Delta_{\Gamma}w + i\partial_t w, \quad Q_2w = -ds\partial_{\nu}\varphi w + is\partial_t\varphi w,$$

and

$$R_{\Gamma}w = -d\partial_{\nu}w + \vec{q}_{\Gamma,1} \cdot \nabla_{\Gamma}w + s\nabla_{\Gamma}\varphi \cdot \vec{q}_{\Gamma,1}w + q_{\Gamma,0}w$$

Now, taking the $L^2(\Omega \times (0,T))$ -norm in (3.1) and the $L^2(\Gamma_1 \times (0,T))$ -norm in (3.2) we get

$$\int_{0}^{T} \int_{\Omega} |Pw - Rw|^{2} dx dt + \int_{0}^{T} \int_{\Gamma_{1}} |Qw - R_{\Gamma}w|^{2} d\sigma dt$$

=
$$\int_{0}^{T} \int_{\Omega} \left(|P_{1}w|^{2} + |P_{2}w|^{2} \right) dx dt + \int_{0}^{T} \int_{\Gamma_{1}} \left(|Q_{1}w|^{2} + |Q_{2}w|^{2} \right) d\sigma dt$$

+
$$2\Re \int_{0}^{T} \int_{\Omega} P_{1}w \overline{P_{2}w} dx dt + 2\Re \int_{0}^{T} \int_{\Gamma_{1}} Q_{1}w \overline{Q_{2}w} d\sigma dt.$$
 (3.3)

The next two steps of the proof are devoted to compute the last two terms of the right-hand side of (3.3).

• Step 2: Estimates in $\Omega \times (0,T)$.

In this step, we compute the terms

$$\Re \int_0^T \int_\Omega P_1 w \overline{P_2 w} \, dx dt = \sum_{j=1}^3 \sum_{k=1}^3 I_{jk},$$

where I_{jk} denotes the L^2 real inner product of the j^{th} -term of P_1w with the k^{th} -term of P_2w . In order to do that, we shall take into account some properties of the function μ defined in (1.2),

which are stated in the following result.

Proposition 3.1. Under the hypotheses of Theorem 1.4, the function μ defined by (1.2) satisfies the following properties:

- 1. $\mu \in C^4(\overline{\Omega})$.
- 2. $\mu = 1$ on Γ_1 .
- 3. $\nabla \mu \neq 0$ in $\overline{\Omega}$.
- 4. There exists c > 0 such that $D^2 \mu(\xi, \xi) \ge c |\xi|^2$ in $\overline{\Omega}$ for all $\xi \in \mathbb{R}^n$.

Proof. We recall that the origin belongs to the open set Ω_1 , and then, there exists $\varepsilon > 0$ such that $\overline{B_{\varepsilon}(0)} \cap \overline{\Omega} = \emptyset$. Taking into account hypothesis (1.4), it is not difficult to see that, for each $x \in \Omega$, we have

$$\mu(x) = \frac{\|x\|}{\left\|\Phi^{-1}\left(\frac{x}{\|x\|}\right)\right\|},$$

and then, we obtain that μ is of class C^4 in $\overline{\Omega}$. Moreover, directly from the definition of μ we get that $\Gamma_1 = \{x \in \overline{\Omega} : \mu(x) = 1\}$. Finally, assertions 3 and 4 are proved in [3] (see also [4]).

In the following, we consider the following computations

$$\nabla \varphi = -\lambda \theta \nabla \psi, \quad \nabla \theta = \lambda \theta \nabla \psi, \quad \Delta \varphi = -\lambda^2 \theta |\nabla \psi|^2 - \lambda \theta \Delta \psi, \text{ in } \Omega \times (0, T).$$
(3.4)

Moreover,

$$\partial_{\nu}\varphi = -\lambda\theta\partial_{\nu}\psi$$
, on $\partial\Omega \times (0,T)$, and $\nabla_{\Gamma}\theta = \nabla_{\Gamma}\varphi = 0$, on $\Gamma_1 \times (0,T)$. (3.5)

Here and subsequently, C stands for a positive constant which is independent of the parameters $\lambda > 0$ and s > 0 and that might change from line to line in our computations.

According to (3.4), the term I_{11} can be written as

$$I_{11} = d^2 s^3 \int_0^T \int_\Omega \Delta \varphi |\nabla \varphi|^2 |w|^2 dx dt$$

= $-d^2 s^3 \lambda^3 \int_0^T \int_\Omega (\lambda |\nabla \psi|^2 + \Delta \psi) |\nabla \psi|^2 \theta^3 |w|^2 dx dt.$ (3.6)

Also, we have

$$\begin{split} I_{12} =& d^2 s^3 \int_0^T \int_\Omega |\nabla \varphi|^2 \nabla \varphi \cdot \nabla |w|^2 \, dx dt \\ =& -d^2 s^3 \int_0^T \int_\Omega \nabla (|\nabla \varphi|^2) \cdot \nabla \varphi |w|^2 \, dx dt - d^2 s^3 \int_0^T \int_\Omega \Delta \varphi |\nabla \varphi|^2 |w|^2 \, dx dt \\ &+ d^2 s^3 \int_0^T \int_{\Gamma_1} |\nabla \varphi|^2 \partial_\nu \varphi |w|^2 \, d\sigma dt \\ =& 3d^2 s^3 \lambda^4 \int_0^T \int_\Omega |\nabla \psi|^4 \theta^3 |w|^2 \, dx dt - d^2 s^3 \lambda^3 \int_0^T \int_{\Gamma_1} (\partial_\nu \psi)^3 \theta^3 |w|^2 \, d\sigma dt \\ &+ d^2 s^3 \lambda^3 \int_0^T \int_\Omega (2\nabla^2 \psi (\nabla \psi, \nabla \psi) + \Delta \psi |\nabla \psi|^2) \theta^3 |w|^2 \, dx dt, \end{split}$$
(3.7)

where we have used that w = 0 on $\Gamma_0 \times (0, T)$. Now, the term I_{13} is

$$I_{13} = -ds^3 \Re i \int_0^T \int_\Omega \partial_t \varphi |\nabla \varphi|^2 |w|^2 \, dx dt = ds^3 \Im \int_0^T \int_\Omega \partial_t \varphi |\nabla \varphi|^2 |w|^2 \, dx dt = 0. \tag{3.8}$$

Adding up the equations (3.6), (3.7) and (3.8) we get

$$\sum_{k=1}^{3} I_{1k} = 2d^2 s^3 \lambda^4 \int_0^T \int_\Omega |\nabla \psi|^4 \theta^3 |w|^2 dx dt + 2d^2 s^3 \lambda^3 \int_0^T \int_\Omega \nabla^2 \psi (\nabla \psi, \nabla \psi) \theta^3 |w|^2 dx dt - d^2 s^3 \lambda^3 \int_0^T \int_{\Gamma_1} (\partial_\nu \psi)^3 \theta^3 |w|^2 d\sigma dt \geqslant 2d^2 \gamma^4 s^3 \lambda^4 \int_0^T \int_\Omega \theta^3 |w|^2 dx dt + d^2 \gamma^3 s^3 \lambda^3 \int_0^T \int_{\Gamma_1} \theta^3 |w|^2 d\sigma dt.$$
(3.9)

On the other hand,

$$\begin{split} I_{21} = & d^2 s \Re \int_0^T \int_{\Omega} \Delta \varphi \overline{w} \Delta w \, dx dt \\ = & -d^2 s \int_0^T \int_{\Omega} \Delta \varphi |\nabla w|^2 \, dx dt - \frac{d^2 s}{2} \int_0^T \int_{\Omega} \nabla (\Delta \varphi) \cdot \nabla |w|^2 \, dx dt \\ & + d^2 s \Re \int_0^T \int_{\Gamma_1} \Delta \varphi \overline{w} \partial_{\nu} w \, d\sigma dt \\ = & -d^2 s \int_0^T \int_{\Omega} \Delta \varphi |\nabla w|^2 \, dx dt + \frac{d^2 s}{2} \int_0^T \int_{\Omega} \Delta^2 \varphi |w|^2 \, dx dt \\ & - \frac{d^2}{2} s \int_0^T \int_{\Gamma_1} \partial_{\nu} (\Delta \varphi) |w|^2 \, d\sigma dt + d^2 s \Re \int_0^T \int_{\Gamma_1} \Delta \varphi \overline{w} \partial_{\nu} w \, d\sigma dt. \end{split}$$

Now, explicit computations show that

$$|\Delta^2 \varphi| \ge C \lambda^4 \theta$$
, in $\Omega \times (0, T)$, and $|\partial_{\nu} (\Delta \varphi)| \ge C \lambda^3 \theta$, on $\Gamma_1 \times (0, T)$.

According to these estimates, I_{21} can be bounded in the following way

$$I_{21} \geq s \int_{0}^{T} \int_{\Omega} \left(\lambda^{2} \theta |\nabla \psi|^{2} + \lambda \theta \Delta \psi \right) |\nabla w|^{2} dx dt - Cs \lambda^{4} \int_{0}^{T} \int_{\Omega} \theta |w|^{2} dx dt - Cs \lambda^{2} \int_{0}^{T} \int_{\Gamma_{1}} \theta |w| \partial_{\nu} w \, d\sigma dt - Cs \lambda^{3} \int_{0}^{T} \int_{\Gamma_{1}} \theta |w|^{2} \, d\sigma dt.$$

$$(3.10)$$

On the other hand,

$$\begin{split} I_{22} =& 2d^2 s \Re \int_0^T \int_{\Omega} \Delta w \nabla \varphi \cdot \nabla \overline{w} \, dx dt \\ =& -2d^2 s \Re \int_0^T \int_{\Omega} \nabla w \cdot \nabla (\nabla \varphi \cdot \nabla \overline{w}) \, dx dt + 2d^2 s \Re \int_0^T \int_{\partial \Omega} (\nabla \varphi \cdot \nabla \overline{w}) \partial_{\nu} w d\sigma dt \\ =& K_1 + K_2. \end{split}$$

On the one hand, we have

$$\begin{split} K_1 &= -2d^2 s \Re \int_0^T \int_\Omega \left(\nabla^2 \varphi(\nabla w, \nabla \overline{w}) + \frac{1}{2} \nabla \varphi \cdot \nabla(|\nabla w|^2) \right) \, dx dt \\ &= -2d^2 s \Re \int_0^T \int_\Omega \nabla^2 \varphi(\nabla w, \nabla \overline{w}) \, dx dt + d^2 s \int_0^T \int_\Omega \Delta \varphi |\nabla w|^2 \, dx dt \\ &- d^2 s \int_0^T \int_{\partial \Omega} \partial_\nu \varphi |\nabla w|^2 d\sigma dt. \end{split}$$

Explicit computations show that

$$\nabla^2 \varphi(\nabla w, \nabla \overline{w}) = -\lambda^2 \theta |\nabla w \cdot \nabla \psi|^2 - \lambda \theta \nabla^2 \psi(\nabla w, \nabla w),$$

and

$$\nabla w = \partial_{\nu} w \nu, \text{ on } \Gamma_0 \times (0,T), \quad \partial_{\nu} \varphi = -\lambda \theta \partial_{\nu} \psi \text{ on } \partial \Omega \times (0,T).$$

Then,

$$\begin{split} K_{1} =& 2d^{2}s \Re \int_{0}^{T} \int_{\Omega} (\lambda^{2}\theta |\nabla\psi \cdot \nabla w|^{2} + \lambda\theta \nabla^{2}\psi(\nabla w, \nabla \overline{w})) \, dx dt \\ &- d^{2}s \lambda \int_{0}^{T} \int_{\Omega} (\lambda |\nabla\psi|^{2} + \Delta\psi)\theta |\nabla w|^{2} \, dx dt \\ &+ d^{2}s \lambda \int_{0}^{T} \int_{\Gamma_{0}} \partial_{\nu}\psi\theta |\partial_{\nu}w|^{2} \, d\sigma dt + d^{2}s \lambda \int_{0}^{T} \int_{\Gamma_{1}} \partial_{\nu}\psi\theta(|\partial_{\nu}w|^{2} + |\nabla_{\Gamma}w|^{2}) \, d\sigma dt. \end{split}$$

On the other hand,

$$K_{2} = 2d^{2}s \int_{0}^{T} \int_{\Gamma_{0}} \partial_{\nu}\varphi |\partial_{\nu}w|^{2} d\sigma dt + 2d^{2}s \int_{0}^{T} \int_{\Gamma_{1}} \partial_{\nu}\varphi |\partial_{\nu}w|^{2} d\sigma dt$$
$$= -2d^{2}s\lambda \int_{0}^{T} \int_{\Gamma_{0}} \partial_{\nu}\psi \theta |\partial_{\nu}w|^{2} d\sigma dt - 2d^{2}s\lambda \int_{0}^{T} \int_{\Gamma_{1}} \partial_{\nu}\psi \theta |\partial_{\nu}w|^{2} d\sigma dt.$$

Thus,

$$I_{22} = 2d^{2}s\lambda \int_{0}^{T} \int_{\Omega} \left(\lambda\theta |\nabla\psi \cdot \nabla w|^{2} + \theta\nabla^{2}\psi(\nabla w, \nabla \overline{w})\right) dxdt$$

$$- d^{2}s\lambda \int_{0}^{T} \int_{\Omega} (\lambda |\nabla\psi|^{2} + \Delta\psi)\theta |\nabla w|^{2} dxdt$$

$$- d^{2}s\lambda \int_{0}^{T} \int_{\Gamma_{0}} \partial_{\nu}\psi\theta |\partial_{\nu}w|^{2} d\sigma dt - d^{2}s\lambda \int_{0}^{T} \int_{\Gamma_{1}} \partial_{\nu}\psi\theta |\partial_{\nu}w|^{2} d\sigma dt$$

$$+ d^{2}s\lambda \int_{0}^{T} \int_{\Gamma_{1}} \partial_{\nu}\psi\theta |\nabla_{\Gamma}w|^{2} d\sigma dt.$$

(3.11)

Now, I_{23} is given by

$$I_{23} = -ds \Re i \int_0^T \int_\Omega \partial_t \varphi \overline{w} \Delta w \, dx dt$$

= $-ds \Im \int_0^T \int_\Omega \overline{w} \nabla (\partial_t \varphi) \cdot \nabla w \, dx dt + ds \Im \int_0^T \int_{\Gamma_1} \partial_t \varphi \overline{w} \partial_\nu w \, d\sigma dt.$

By Young's inequality, for all $\epsilon > 0$ there exists a constant $C = C(\epsilon)$ such that

$$I_{23} \ge -\int_0^T \int_\Omega \epsilon s\lambda \theta |\nabla w|^2 + C(\epsilon) s\lambda \theta^3 |w|^2 \, dx dt -\int_0^T \int_{\Gamma_1} (\epsilon s\lambda \theta |\partial_\nu w|^2 + C(\epsilon) s\lambda \theta^3 |w|^2) \, d\sigma dt.$$
(3.12)

Therefore, adding inequalities (3.10), (3.11) and (3.12), choosing $\epsilon > 0$ small enough and taking λ_0 and s_0 sufficiently large, we get

$$\sum_{k=1}^{3} I_{2k} \geq Cs\lambda \int_{0}^{T} \int_{\Omega} \theta |\nabla w|^{2} dx dt + Cs\lambda \int_{0}^{T} \int_{\Gamma_{0}} \theta |\partial_{\nu}w|^{2} d\sigma dt + Cs\lambda \int_{0}^{T} \int_{\Gamma_{1}} \theta |\partial_{\nu}w|^{2} d\sigma dt + d^{2}s\lambda \int_{0}^{T} \int_{\Gamma_{1}} \partial_{\nu}\psi \theta |\nabla_{\Gamma}w|^{2} d\sigma dt - Cs\lambda \int_{0}^{T} \int_{\Omega} \theta^{3} |w|^{2} dx dt - Cs\lambda \int_{0}^{T} \int_{\Gamma_{1}} \theta^{3} |w|^{2} d\sigma dt,$$

$$(3.13)$$

for all $\lambda \ge \lambda_0$ and for all $s \ge s_0$.

Moreover,

$$I_{31} = ds \Re i \int_0^T \int_\Omega \Delta \varphi \overline{w} \partial_t w \, dx dt = -ds \Im \int_0^T \int_\Omega \Delta \varphi \overline{w} \partial_t w \, dx dt \tag{3.14}$$

Integrating by parts, first in time and then in space, we have

$$\begin{split} I_{32} = & 2ds \Im \int_0^T \int_\Omega \partial_t \overline{w} \nabla \varphi \cdot \nabla w \, dx dt \\ = & -2ds \Im \int_0^T \int_\Omega \overline{w} \nabla \varphi \cdot \nabla \partial_t w \, dx dt - 2ds \Im \int_0^T \int_\Omega \overline{w} \nabla \partial_t \varphi \cdot \nabla w \, dx dt \\ = & -ds \Im \int_0^T \int_\Omega \overline{w} \nabla \partial_t \varphi \cdot \nabla w \, dx dt + ds \Im \int_0^T \int_\Omega \Delta \varphi \overline{w} \partial_t w \, dx dt \\ & - ds \Im \int_0^T \int_{\Gamma_1} \partial_\nu \varphi \overline{w} \partial_t w \, d\sigma dt. \end{split}$$

By Young's inequality, for all $\epsilon > 0$ there exists a constant $C = C(\epsilon)$ such that

$$I_{32} \ge -\epsilon s\lambda \int_0^T \int_\Omega \theta |\nabla w|^2 \, dx dt - C(\epsilon) s\lambda \int_0^T \int_\Omega \theta^3 |w|^2 \, dx dt + ds \Im \int_0^T \int_\Omega \Delta \varphi \overline{w} \partial_t w \, dx dt - ds \Im \int_0^T \int_{\Gamma_1} \partial_\nu \varphi \overline{w} \partial_t w \, d\sigma dt.$$
(3.15)

For the term I_{33} we have:

$$I_{33} = \Re s \int_0^T \int_\Omega \partial_t \varphi \overline{w} \partial_t w \, dx dt = +\frac{1}{2} s \int_0^T \int_\Omega \partial_t \varphi \partial_t (|w|^2) \, dx dt$$

$$\geq -Cs \int_0^T \int_\Omega \theta^3 |w|^2 \, dx dt.$$
(3.16)

Now, adding (3.14), (3.15) and (3.16) we conclude that

$$\sum_{k=1}^{3} I_{3k} \ge -\epsilon s\lambda \int_{0}^{T} \int_{\Omega} \theta |\nabla w|^{2} dx dt - C(\epsilon) s\lambda^{3} \int_{0}^{T} \int_{\Omega} \theta^{3} |w|^{2} dx dt + ds\lambda \Im \int_{0}^{T} \int_{\Gamma_{1}} \partial_{\nu} \psi \theta \overline{w} \partial_{t} w d\sigma dt.$$
(3.17)

Now, sum up inequalities (3.9), (3.13) and (3.17), using Young's inequality and taking λ_0 and s_0 large enough, we obtain

$$\begin{split} \Re \int_{0}^{T} \int_{\Omega} P_{1} w \overline{P_{2}w} \, dx dt \\ \geqslant C \int_{0}^{T} \int_{\Omega} \left(s^{3} \lambda^{4} \theta^{3} |w|^{2} + s \lambda \theta |\nabla w|^{2} + s \lambda^{2} \theta |\nabla \psi \cdot \nabla w|^{2} \right) \, dx dt \\ &+ C \int_{0}^{T} \int_{\Gamma_{1}} \left(s^{3} \lambda^{3} \theta^{3} |w|^{2} + s \lambda \theta |\partial_{\nu} w|^{2} \right) \, d\sigma dt - d^{2} s \lambda \int_{0}^{T} \int_{\Gamma_{0}} \partial_{\nu} \psi \theta |\partial_{\nu} w|^{2} \, d\sigma dt \\ &+ d^{2} s \lambda \int_{0}^{T} \int_{\Gamma_{1}} \partial_{\nu} \psi \theta |\nabla_{\Gamma} w|^{2} \, d\sigma dt + ds \lambda \Im \int_{0}^{T} \int_{\Gamma_{1}} \partial_{\nu} \psi \theta \overline{w} \partial_{t} w \, d\sigma dt. \end{split}$$

$$(3.18)$$

Notice that the global term of $\nabla_{\Gamma} w$ in the last inequality is not strictly positive due to $\partial_{\nu} \psi < 0$ on Γ_1 . As we shall see in the next step, this difficulty can be avoided choosing d and δ according to assumption (1.5).

• Step 3: Estimates on $\Gamma_1 \times (0,T)$.

In this step, we devote to compute the terms defined on $\Gamma_1 \times (0, T)$. By definition of Q_1 and Q_2 , we have

$$\Re \int_0^T \int_{\Gamma_1} Q_1 w \overline{Q_2 w} \, d\sigma dt = \sum_{j=1}^2 \sum_{k=1}^2 J_{jk},$$

where J_{jk} stands for the $L^2(\Gamma_1 \times (0, T))$ -inner product between the j^{th} -term of $Q_1 w$ and the k^{th} -term of $Q_2 w$.

Firstly, notice that the term J_{11} can be written as follows:

$$J_{11} = -\delta ds \Re \int_0^T \int_{\Gamma_1} \partial_\nu \varphi \overline{w} \Delta_\Gamma w \, d\sigma dt$$

= $-\delta ds \lambda \int_0^T \int_{\Gamma_1} \partial_\nu \psi \theta |\nabla_\Gamma w|^2 \, d\sigma dt + \delta ds \Re \int_0^T \int_{\Gamma_1} \overline{w} \nabla_\Gamma (\partial_\nu \varphi) \cdot \nabla_\Gamma w \, d\sigma dt.$

Using the formula $\operatorname{div}_{\Gamma}(w\nabla_{\Gamma}(\partial_{\nu}\psi)) = \nabla_{\Gamma}(\partial_{\nu}\psi) \cdot \nabla_{\Gamma}w + w\Delta_{\Gamma}(\partial_{\nu}\psi)$ and integrating by parts we easily deduce that

$$\begin{split} &-\delta ds\lambda \Re \int_0^T \int_{\Gamma_1} \overline{w} \theta \nabla_{\Gamma}(\partial_{\nu}\psi) \cdot \nabla_{\Gamma} w \, d\sigma dt \\ =& \delta ds\lambda \Re \int_0^T \int_{\Gamma_1} w \nabla_{\Gamma}(\overline{w}\theta) \nabla_{\Gamma}(\partial_{\nu}\psi) \, d\sigma dt + \delta ds\lambda \int_0^T \int_{\Gamma_1} \Delta_{\Gamma}(\partial_{\nu}\psi) \theta |w|^2 \, d\sigma dt \\ =& \frac{1}{2} \delta ds\lambda \int_0^T \int_{\Gamma_1} \Delta_{\Gamma}(\partial_{\nu}\psi) \theta |w|^2 \, d\sigma dt, \end{split}$$

where we have used that $\nabla_{\Gamma} \theta = 0$ on $\Gamma_1 \times (0, T)$.

In the same manner, J_{12} can be estimate as follows:

$$J_{12} = \delta s \Im \int_0^T \int_{\Gamma_1} \partial_t \varphi \overline{w} \Delta_{\Gamma} w \, d\sigma dt$$

= $-\delta s \Im \int_0^T \int_{\Gamma_1} \overline{w} (\nabla_{\Gamma} \partial_t \varphi) \cdot \nabla_{\Gamma} w \, d\sigma dt - \delta s \Im \int_0^T \int_{\Gamma_1} \partial_t \varphi |\nabla_{\Gamma} w|^2 \, d\sigma dt$
= $0,$

since $\nabla_{\Gamma}\psi = 0$ on Γ_1 .

On the other hand, the term J_{21} is given by

$$J_{21} = -ds \Re i \int_0^T \int_{\Gamma_1} \partial_\nu \varphi \overline{w} \partial_t w \, d\sigma dt = ds \Im \int_0^T \int_{\Gamma_1} \partial_\nu \varphi \overline{w} \partial_t w \, d\sigma dt$$
$$= -ds \lambda \Im \int_0^T \int_{\Gamma_1} \partial_\nu \psi \theta \overline{w} \partial_t w \, d\sigma dt.$$

Moreover,

$$J_{22} = s\Re \int_0^T \int_{\Gamma_1} \partial_t \varphi \overline{w} \partial_t w \, d\sigma dt \ge -Cs \int_0^T \int_{\Gamma_1} \theta^2 |w|^2 \, d\sigma dt.$$

According to the above computations, we easily deduce that

$$\Re \int_{0}^{T} \int_{\Gamma_{1}} Q_{1} w \overline{Q_{2}w} \, d\sigma dt$$

$$\geq -\delta ds \lambda \int_{0}^{T} \int_{\Gamma_{1}} \partial_{\nu} \psi |\nabla_{\Gamma}w|^{2} \, d\sigma dt - Cs \lambda \int_{0}^{T} \int_{\Gamma_{1}} \theta |w|^{2} \, d\sigma dt \qquad (3.19)$$

$$-Cs \int_{0}^{T} \int_{\Gamma_{1}} \theta^{2} |w|^{2} \, d\sigma dt - ds \lambda \Im \int_{0}^{T} \int_{\Gamma_{1}} \partial_{\nu} \psi \theta \overline{w} \partial_{t} w \, d\sigma dt.$$

In the next step, we gather the terms obtained in the steps 2 and 3 to conclude the desired Carleman estimate.

• Step 4: Last arrangements and conclusion

Adding the inequalities (3.18) and (3.19), using Young's inequality, and taking s_0 and λ_0 large enough if it is necessary, we obtain

$$\int_{0}^{T} \int_{\Omega} (s^{3}\lambda^{4}\theta^{3}|w|^{2} + s\lambda|\nabla w|^{2} + s\lambda^{2}\theta|\nabla\psi\cdot\nabla w|^{2}) dxdt$$

$$+ \int_{0}^{T} \int_{\Gamma_{1}} (s^{3}\lambda^{3}|w|^{2} + s\lambda|\partial_{\nu}w|^{2}) d\sigma dt$$

$$+ d(\delta - d)s\lambda \int_{0}^{T} \int_{\Gamma_{1}} |\partial_{\nu}\psi||\nabla_{\Gamma}w|^{2} d\sigma dt$$

$$\leq C \int_{0}^{T} \int_{\Omega} |Pw|^{2} dxdt + C \int_{0}^{T} \int_{\Gamma_{1}} |Qw|^{2} d\sigma dt + Cs\lambda \int_{0}^{T} \int_{\Gamma_{0}} |\partial_{\nu}w|^{2} d\sigma dt$$

$$+ C \int_{0}^{T} \int_{\Omega} |Rw|^{2} dxdt + C \int_{0}^{T} \int_{\Gamma_{1}} |R_{\Gamma}w|^{2} d\sigma dt,$$
(3.20)

for all $\lambda \ge \lambda_0$ and $s \ge s_0$. We notice that the global term of $\nabla_{\Gamma} w$ on $\Gamma_1 \times (0, T)$ is positive since we assumed the condition (1.5). Moreover, it is easy to see that

$$\begin{split} &\int_0^T \int_\Omega |Rw|^2 \, dx dt + \int_0^T \int_{\Gamma_1} |R_{\Gamma}w|^2 \, d\sigma dt \\ \leqslant C \int_0^T \int_\Omega \left(s^2 \lambda^2 \theta^2 |w|^2 + |\nabla w|^2 \right) \, dx dt \\ &+ C \int_0^T \int_{\Gamma_1} \left(s^2 \lambda^2 \theta^2 |w|^2 + |\partial_\nu w|^2 + |\nabla_\Gamma w|^2 \right) \, d\sigma dt. \end{split}$$

Therefore, the last two terms of the right-hand side of (3.20) can be absorbed by the left hand side taking s_0 and λ_0 sufficiently large.

Finally, we come back to the original variable $v = e^{s\varphi}w$. Taking into account that

$$e^{-2s\varphi}|\nabla v|^2 \leqslant C(s^2\lambda^2\theta^2|w|^2 + |\nabla w|^2)$$

and that, for each $v \in \mathcal{V}$,

$$\partial_{\nu}w = e^{s\varphi}\partial_{\nu}v \quad \text{on } \Gamma_0,$$

we obtain (1.6). This concludes the proof of Theorem 1.4.

3.2 Proof of the Corollary 1.8

We start considering a real function $\eta \in C^{\infty}(\overline{\Omega})$ such that $\eta = 1$ in $\overline{\Omega} \setminus \omega$ and vanishing close to Γ_* . if $w = \eta v$ in $\Omega \times (0, T)$ and $w_{\Gamma} = \eta v_{\Gamma}$ on $\Gamma_1 \times (0, T)$, then it is easy to see that (w, w_{Γ}) satisfies the equations

$$L(w) = \eta L(v) + 2d\nabla \eta \cdot \nabla v + \Delta \eta v, \text{ in } \Omega \times (0,T), \quad N(w,w_{\Gamma}) = 0, \text{ on } \Gamma_1 \times (0,T),$$

and

$$w = \partial_{\nu} w = 0$$
, on $\Gamma_* \times (0, T)$, $w = w_{\Gamma}$, on $\Gamma_1 \times (0, T)$.

Besides, since v = w in $\Omega \setminus \omega$, we point out that

$$\int_{0}^{T} \int_{\Omega} e^{-2s\varphi} \theta |\nabla w|^{2} dx dt \ge \int_{0}^{T} \int_{\Omega \setminus \omega} e^{-2s\varphi} \theta |\nabla w|^{2} dx dt$$

$$= \int_{0}^{T} \int_{\Omega \setminus \omega} e^{-2s\varphi} \theta |\nabla v|^{2} dx dt.$$
(3.21)

Then, applying the Carleman inequality (1.6) to $w = \eta v$ and taking into account the estimate (3.21) we easily get

$$\int_{0}^{T} \int_{\Omega} e^{-2s\varphi} (s^{3}\lambda^{4}\theta^{3}|v|^{2} + s\lambda\theta|\nabla v|^{2} dxdt + \int_{0}^{T} \int_{\Gamma_{1}} e^{-2s\varphi} (s^{3}\lambda^{3}\theta^{3}|v|^{2} + s\lambda\theta|\nabla_{\Gamma}v_{\Gamma}|^{2} + s\lambda\theta|\partial_{\nu}v|^{2}) d\sigma dt \leqslant C \int_{0}^{T} \int_{\Omega} e^{-2s\varphi} |L(v)|^{2} dxdt + C \int_{0}^{T} \int_{\Gamma_{1}} e^{-2s\varphi} |N(v,v_{\Gamma})|^{2} d\sigma dt$$
(3.22)
$$+ C \int_{0}^{T} \int_{\omega} e^{-2s\varphi} (s^{3}\lambda^{4}\theta^{3}|v|^{2} + s\lambda\theta|\nabla v|^{2}) dxdt + C \int_{0}^{T} \int_{\Omega} e^{-2s\varphi} (|v|^{2} + |\nabla v|^{2}) dxdt,$$

for all $\lambda \ge \lambda_0$ and for all $s \ge s_0$. We notice that, the last two terms of the right-hand side of (3.22) can be absorbed by taking λ_0 and s_0 sufficiently large if it is necessary. This completes the proof of the Corollary 1.8.

4 Proof of the Theorem 1.9

We introduce the adjoint system of (1.1)

$$\begin{cases} i\partial_t z + d\Delta z + i\vec{r} \cdot \nabla z + qz = 0, & \text{in } \Omega \times (0,T), \\ i\partial_t z_{\Gamma} - d\partial_{\nu} z + \delta \Delta_{\Gamma} z_{\Gamma} + i\vec{r}_{\Gamma} \cdot \nabla_{\Gamma} z_{\Gamma} + q_{\Gamma} z_{\Gamma} = 0, & \text{on } \Gamma_1 \times (0,T), \\ z = z_{\Gamma}, & \text{on } \Gamma_1 \times (0,T), \\ z = 0, & \text{on } \Gamma_0 \times (0,T), \\ (z, z_{\Gamma})(T) = (z_T, z_{\Gamma,T}), & \text{in } \Omega \times \Gamma_1, \end{cases}$$
(4.1)

where (q, q_{Γ}) are given by

$$q := i \operatorname{div}(\vec{r}) + \overline{q}_0, \quad q_{\Gamma} := i \operatorname{div}_{\Gamma}(\vec{r}_{\Gamma}) - i \vec{r} \cdot \nu + \overline{q}_{\Gamma,0}.$$

$$(4.2)$$

Since $\vec{r} \in [L^{\infty}(0,T;W^{2,\infty}(\Omega;\mathbb{R}))]^n$ and $\vec{r}_{\Gamma} \in [L^{\infty}(0,T;W^{2,\infty}(\Gamma;\mathbb{R}))]^n$, q and q_{Γ} defined in (4.2) belongs to $L^{\infty}(0,T;W^{1,\infty}(\Omega))$ and $L^{\infty}(0,T;W^{1,\infty}(\Gamma_1))$, respectively, and therefore (using the change of variables $t \mapsto T - t$) problem (4.1) has a unique solution $(z,z_{\Gamma}) \in C^0([0,T];\mathcal{V})$.

On the other hand, according to Proposition 2.8, the exact controllability of (1.1) is equivalent to prove the observability inequality

$$\|(z_T, z_{\Gamma,T})\|_{\mathcal{V}}^2 \leqslant C \int_0^T \int_{\Gamma_*} |\partial_\nu z|^2 d\sigma dt,$$
(4.3)

for all $(z_T, z_{\Gamma,T}) \in \mathcal{V}$ and (z, z_{Γ}) being the associated solution of (4.1) and Γ_* is given by (1.7). Firstly, we shall introduce a cut-off function $\zeta \in C^1([0, T]; \mathbb{R})$ such that

$$0\leqslant \zeta\leqslant 1,\quad \zeta=0,\quad \forall t\in [0,T/4],\quad \zeta=1,\quad \forall t\in [3T/4,T].$$

Then, the variables $(w, w_{\Gamma}) = (\zeta z, \zeta z_{\Gamma})$ solves the problem

$$\begin{cases} i\partial_t w + d\Delta w + i\vec{r} \cdot \nabla w + qw = i\zeta' z, & \text{in } \Omega \times (0,T), \\ i\partial_t w_{\Gamma} - d\partial_{\nu} w + \delta\Delta_{\Gamma} w_{\Gamma} + i\vec{r}_{\Gamma} \cdot \nabla_{\Gamma} w_{\Gamma} + q_{\Gamma} w = i\zeta' z_{\Gamma}, & \text{on } \Gamma_1 \times (0,T), \\ w = w_{\Gamma}, & \text{on } \Gamma_1 \times (0,T), \\ w = 0, & \text{on } \Gamma_0 \times (0,T), \\ (w, w_{\Gamma})(0) = (0,0), & \text{in } \Omega \times \Gamma_1, \end{cases}$$

$$(4.4)$$

By Proposition 2.2, $(w, w_{\Gamma}) \in C^0([0, T]; \mathcal{V})$ and since $\operatorname{supp}(\zeta') \subseteq (T/4, 3T/4)$ the following inequality holds

$$\|(w, w_{\Gamma})\|_{C^{0}([T/4, T]; \mathcal{V})}^{2} \leq C \|(z, z_{\Gamma})\|_{L^{1}(T/4, 3T/4; \mathcal{V})}^{2}.$$

In particular, we can assert that

$$\|(z_T, z_{\Gamma,T})\|_{\mathcal{V}}^2 \leqslant C \|(z, z_{\Gamma})\|_{L^2(T/4, 3T/4; \mathcal{V})}^2.$$
(4.5)

Moreover, by Carleman estimate (see Theorem 1.4) applied to (z, z_{Γ}) we obtain

$$\int_{T/4}^{3T/4} \int_{\Omega} e^{-2s\varphi} (s^{3}\lambda^{4}\theta^{3}|z|^{2} + s\lambda^{2}\theta|\nabla z|^{2}) dx dt + \int_{T/4}^{3T/4} \int_{\Gamma_{1}} e^{-2s\varphi} (s^{3}\lambda^{3}\theta^{3}|z_{\Gamma}|^{2} + s\lambda\theta|\nabla_{\Gamma}z_{\Gamma}|^{2}) d\sigma dt$$

$$\leq Cs\lambda \int_{0}^{T} \int_{\Gamma_{*}} e^{-2s\varphi}\theta|\partial_{\nu}z|^{2} d\sigma dt,$$

$$(4.6)$$

for all $s \ge s_0$ and $\lambda \ge \lambda_0$. Now, we fix s and λ . Since φ is bounded, there exists a constant C > 0 such that

$$1 \leqslant Ce^{-2s\varphi}, \quad \forall x \in \overline{\Omega} \times (T/4, 3T/4).$$
 (4.7)

Now, combining (4.5), (4.6) and (4.7) we easily deduce (4.3) and therefore the proof of the Theorem 1.9 is done.

Acknowledgments

The authors are indebted to the anonymous referees for their comments and suggestions, which in particular allowed us to improve the scope of our problem.

References

- M. Aassila. Exact controllability of the Schrödinger equation. Appl. Math. Comput., 144(1):89– 106, 2003.
- [2] A. D. Bandrauk. *Molecules in laser fields*. CRC Press, 1993.
- [3] L. Baudouin and A. Mercado. An inverse problem for Schrödinger equations with discontinuous main coefficient. Appl. Anal., 87(10-11):1145–1165, 2008.
- [4] L. Baudouin, A. Mercado, and A. Osses. A global Carleman estimate in a transmission wave equation and application to a one-measurement inverse problem. *Inverse Problems*, 23(1):257– 278, 2007.
- [5] L. Baudouin and J.-P. Puel. Uniqueness and stability in an inverse problem for the Schrödinger equation. *Inverse Problems*, 23(3):1327–1328, 2007.
- [6] I. Benabbas and D. E. Teniou. Observability of wave equation with Ventcel dynamic condition. Evol. Equ. Control Theory, 7(4):545–570, 2018.
- [7] N. Burq. Contrôle de l'équation des plaques en présence d'obstacles strictement convexes. Number 55. 1993.
- [8] M. M. Cavalcanti, W. J. Corrêa, I. Lasiecka, and C. Lefler. Well-posedness and uniform stability for nonlinear Schrödinger equations with dynamic/Wentzell boundary conditions. *Indiana* Univ. Math. J., 65(5):1445–1502, 2016.
- [9] W. J. Corrêa and T. Özsarı. Complex Ginzburg-Landau equations with dynamic boundary conditions. Nonlinear Anal. Real World Appl., 41:607–641, 2018.
- [10] S. Ervedoza, C. Zheng, and E. Zuazua. On the observability of time-discrete conservative linear systems. J. Funct. Anal., 254(12):3037–3078, 2008.
- [11] C. G. Gal and L. Tebou. Carleman inequalities for wave equations with oscillatory boundary conditions and application. SIAM J. Control Optim., 55(1):324–364, 2017.
- [12] A. Giusti-Suzor and F. H. Mies. Vibrational trapping and suppression of dissociation in intense laser fields. *Physical review letters*, 68(26):3869, 1992.
- [13] M. I. Ismailov. Inverse source problem for heat equation with nonlocal Wentzell boundary condition. *Results Math.*, 73(2):Paper No. 68, 11, 2018.
- [14] M. I. Ismailov, I. Tekin, and S. Erkovan. An inverse problem for finding the lowest term of a heat equation with Wentzell-Neumann boundary condition. *Inverse Probl. Sci. Eng.*, 27(11):1608–1634, 2019.
- [15] S. Jaffard. Contrôle interne exact des vibrations d'une plaque rectangulaire. Portugal. Math., 47(4):423–429, 1990.

- [16] J. Jost. *Riemannian geometry and geometric analysis*. Universitext. Springer, Cham, seventh edition, 2017.
- [17] A. Khoutaibi and L. Maniar. Null controllability for a heat equation with dynamic boundary conditions and drift terms. Evol. Equ. Control Theory, 9(2):535–559, 2020.
- [18] I. Lasiecka, R. Triggiani, and X. Zhang. Global uniqueness, observability and stabilization of nonconservative Schrödinger equations via pointwise Carleman estimates. I. H¹(Ω)-estimates. J. Inverse Ill-Posed Probl., 12(1):43–123, 2004.
- [19] I. Lasiecka, R. Triggiani, and X. Zhang. Global uniqueness, observability and stabilization of nonconservative Schrödinger equations via pointwise Carleman estimates. II. L₂(Ω)-estimates. J. Inverse Ill-Posed Probl., 12(2):183–231, 2004.
- [20] G. Lebeau. Contrôle de l'équation de Schrödinger. J. Math. Pures Appl. (9), 71(3):267–291, 1992.
- [21] J.-L. Lions. Contrôlabilité exacte, perturbations et stabilisation de systèmes distribués. Tome 1, volume 8 of Recherches en Mathématiques Appliquées. Masson, Paris, 1988. Contrôlabilité exacte., With appendices by E. Zuazua, C. Bardos, G. Lebeau and J. Rauch.
- [22] E. Machtyngier. Exact controllability for the Schrödinger equation. SIAM J. Control Optim., 32(1):24–34, 1994.
- [23] L. Maniar, M. Meyries, and R. Schnaubelt. Null controllability for parabolic equations with dynamic boundary conditions. *Evol. Equ. Control Theory*, 6(3):381–407, 2017.
- [24] A. Mercado, A. Osses, and L. Rosier. Inverse problems for the Schrödinger equation via Carleman inequalities with degenerate weights. *Inverse Problems*, 24(1):015017, 18, 2008.
- [25] L. Miller. Controllability cost of conservative systems: resolvent condition and transmutation. J. Funct. Anal., 218(2):425–444, 2005.
- [26] M. Nagasawa. Schrödinger equations and diffusion theory. Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel, 1993.
- [27] K.-D. Phung. Observability and control of Schrödinger equations. SIAM J. Control Optim., 40(1):211–230, 2001.
- [28] L. Rosier and B.-Y. Zhang. Exact boundary controllability of the nonlinear Schrödinger equation. J. Differential Equations, 246(10):4129–4153, 2009.
- [29] L. Rosier and B.-Y. Zhang. Null controllability of the complex Ginzburg-Landau equation. volume 26, pages 649–673, 2009.
- [30] M. E. Taylor. Partial differential equations I. Basic theory, volume 115 of Applied Mathematical Sciences. Springer, New York, second edition, 2011.
- [31] M. Tucsnak and G. Weiss. Observation and control for operator semigroups. Birkhäuser Advanced Texts: Basler Lehrbücher. Birkhäuser Verlag, Basel, 2009.
- [32] M. Zhang, J. Yin, and H. Gao. Insensitizing controls for the parabolic equations with dynamic boundary conditions. J. Math. Anal. Appl., 475(1):861–873, 2019.
- [33] E. Zuazua. Remarks on the controllability of the Schrödinger equation. 33:193–211, 2003.