

HADAMARD MATRICES: SKEW OF ORDER 276 AND SYMMETRIC OF ORDER 372

DRAGOMIR Ž. ĐOKOVIĆ

ABSTRACT. The smallest integer $v > 0$ for which no skew-Hadamard matrix of order $4v$ is known is $v = 69$. We show how to construct several such matrices. We also construct the first examples of symmetric Hadamard matrices of order 372.

In memory of my son Dejan Djokovic (1962-2022).

1. INTRODUCTION

According to Table 7.1 of the survey paper of Seberry and Yamada [10], published in 1992, there were only six odd integers $v < 100$ for which no skew-Hadamard matrix of order $4v$ was known at that time, namely the integers

47, 59, 69, 81, 89, 97.

Subsequently, the skew-Hadamard matrices of order $4v$ were constructed in 1994 for $v = 81$ [4], in 2004 for $v = 59$ [9], and in 2008 for $v = 47, 97$ [5]. In this note we construct 19 skew-Hadamard matrices of order 276 ($= 4 \cdot 69$) and 4 symmetric Hadamard matrices of order 372 ($= 4 \cdot 93$).

Let us make two remarks about the skew-Hadamard matrices. First, the case $v = 63$ is listed as unknown in the handbook [2, Table 1.51, p. 277] published in 2007. However the existence of a skew-Hadamard matrix of order $4 \cdot 63$ was known since 1969, as it belongs to an infinite series of such matrices constructed by Szekeres [12]. As this handbook does not list $v = 69$ as unknown, it is probable that this was just a misprint: 63 should be replaced by 69? Second, in the more recent book [11, Table 9.2, pp. 198-200], the cases $v = 39, 49, 65$ are listed as unknown. However the corresponding skew-Hadamard matrices have been constructed long ago in [3].

2. THE FIRST SKEW-HADAMARD MATRICES OF ORDER 276

As far as we know, the smallest integer $v > 0$ for which no skew-Hadamard matrix of order $4v$ is known is $v = 69$ [1, p. 1436]). In this section we construct several such matrices.

Our construction uses the Goethals-Seidel array (GS-array) shown below

$$\begin{bmatrix} A_0 & A_1R & A_2R & A_3R \\ -A_1R & A_0 & -RA_3 & RA_2 \\ -A_2R & RA_3 & A_0 & -RA_1 \\ -A_3R & -RA_2 & RA_1 & A_0 \end{bmatrix}.$$

We shall assume that the A_i are circulants and R is the back-diagonal identity matrix (i.e. the matrix obtained from the identity matrix by reversing the order of

rows). The circulants are obtained from a cyclic difference family $\{X_0, X_1, X_2, X_3\}$ with parameters

$$(v = 69; k_0 = 34, k_1 = 34, k_2 = 31, k_3 = 27; \lambda = 57).$$

For instance, for the first row $(a_0, a_1, \dots, a_{v-1})$ of A_0 we have $a_i = -1$ if $i \in X_0$ and $a_i = 1$ otherwise. Moreover it is required that the block X_0 is skew, i.e. $a_0 = 1$ and $a_i + a_{v-i} = 0$ for $i = 1, 2, \dots, 34$.

A special feature of our difference families is that they break up into two pieces $\{X_0, X_1\}$ and $\{X_2, X_3\}$ which are also difference families.

First, we need a difference family $\{X_0, X_1\}$ with parameters $(69; 34, 34; 33)$ with X_0 skew. This is provided by the well known family of Szekeres difference sets [12, 11]:

$$\begin{aligned} X_0 &= \{1, 2, 6, 7, 9, 13, 14, 16, 17, 18, 21, 27, 31, 34, 36, 37, 39, 40, \\ &\quad 41, 43, 44, 45, 46, 47, 49, 50, 54, 57, 58, 59, 61, 64, 65, 66\}; \\ X_1 &= \{1, 4, 5, 7, 9, 10, 11, 12, 15, 17, 18, 19, 24, 26, 27, 28, 30, 39, \\ &\quad 41, 42, 43, 45, 50, 51, 52, 54, 57, 58, 59, 60, 62, 64, 65, 68\}. \end{aligned}$$

Note that X_0 is skew and X_1 is symmetric. Further, all 19 difference families share the same first two blocks, X_0 and X_1 .

Second, we need a difference family $\{X_2, X_3\}$ with parameters $(69; 31, 27; 24)$. This is exactly the parameter set for a D-optimal design of order $2 \cdot 69 = 138$. In a joint paper with I. Kotsireas [8, Section 4.2], we have constructed 19 nonequivalent such difference families. Anyone of them can be used in our construction. As an example, let us choose the first one:

$$\begin{aligned} X_2 &= \{0, 1, 3, 4, 6, 9, 10, 11, 13, 14, 17, 18, 20, 22, 26, 28, 29, \\ &\quad 32, 33, 34, 39, 41, 43, 45, 46, 48, 51, 59, 60, 62, 63\}, \\ X_3 &= \{0, 2, 3, 4, 8, 9, 10, 11, 12, 15, 16, 17, 21, 25, 26, \\ &\quad 32, 33, 35, 36, 37, 39, 41, 46, 51, 54, 57, 59\}. \end{aligned}$$

By constructing the circulants A_i from the blocks X_i and by plugging the A_i into the GS-array we obtain a skew-Hadamard matrix of order 276.

Consequently, the smallest positive integer v for which the existence of a skew-Hadamard matrix of order $4v$ is still undecided is now 89.

For the readers convenience, we provide (for the difference family chosen above) the first rows of the blocks A_i :

$$\begin{aligned} &+ - - + + + - - + - + + + - - + - - - + + - + + + + - + + + - + + - \\ &+ - - + - - - + - - - - + - - + + + - + + - - - + - + + - - - + +; \\ &+ - + + - - + - + - - - - + + - + - - - + + + + - + - - - + - + + + + \\ &+ + + + - + - - - + - + + + + - - - + - + + - - - - + - + - - + + -; \\ &- - + - - + - + + - - - + - - + + - - + - + - + + + - + - - + + - - - \\ &+ + + + - + - + - + - - + - + + - + + + + + + - - + - - + + + + +; \\ &- + - - - + + + - - - - + + - - - + + + - + + + - - + + + + + - - + \\ &- - - + - + - - + + + + - + + + + - + + - + + - + - + + + + + + + + + . \end{aligned}$$

(The + and - signs stand for +1 and -1, respectively.)

As far as we know, the odd integers $v > 0$ less than 200 for which the existence of skew-Hadamard matrices of order $4v$ is still undecided are the following:

89, 101, 107, 119, 149, 153, 167, 177, 179, 191, 193.

After taking into account the papers [5, 6] (and correcting the hypothetical misprint mentioned earlier), this list agrees with [2, Table 1.51, p. 277].

3. THE FIRST EXAMPLES OF SYMMETRIC HADAMARD MATRICES OF ORDER 372

There are only three odd integers $v < 100$ for which no symmetric Hadamard matrix of order $4v$ is known [7]. These integers are 65, 89 and 93. We have constructed four non-equivalent such matrices for $v = 93$. Our construction uses the so called *propus array* shown below

$$\begin{bmatrix} -A_0 & A_1R & A_2R & A_3R \\ A_2R & RA_3 & A_0 & -RA_1 \\ A_1R & A_0 & -RA_3 & RA_2 \\ A_3R & -RA_2 & RA_1 & A_0 \end{bmatrix}.$$

Note that this array is obtained from the GS-array by multiplying the first column by -1 and switching the second and third rows. If the blocks A_i are circulant matrices and $A_1 = A_2$ then the propus array gives a symmetric matrix.

In our four examples the blocks A_i are circulants of size $v = 93$. Since $A_1 = A_2$ we need only to provide the first rows of the blocks A_0, A_1 and A_3 . We label the 93 positions of the first row by integers $0, 1, 2, \dots, 92$ in that order. It suffices to list the positions of the entries -1 in the first row. These positions are provided by a difference family (X_0, X_1, X_2, X_3) in \mathbf{Z}_{93} with parameters $(93; 45, 41, 41, 41; 75)$ and with $X_1 = X_2$. In the families that we use each block X_i is a union of orbits of the subgroup $H = \{1, 25, 67\}$ of \mathbf{Z}_{93}^* . Instead of listing the elements of X_i we list below just a set of representatives of the H -orbits contained in X_i for $i = 0, 1, 3$. Our four non-equivalent difference families are given by the following four listings:

$$\begin{aligned} &[[0, 1, 3, 4, 9, 11, 16, 17, 18, 20, 26, 31, 40, 44, 48, 55, 62], \\ &[2, 3, 9, 13, 16, 20, 22, 29, 31, 36, 43, 44, 48, 51, 62], \\ &[1, 2, 3, 8, 10, 13, 17, 26, 31, 36, 37, 40, 43, 51, 62]]; \end{aligned}$$

$$\begin{aligned} &[[0, 6, 8, 12, 13, 16, 17, 22, 24, 31, 33, 36, 40, 44, 47, 51, 62], \\ &[1, 4, 5, 6, 9, 11, 13, 20, 31, 33, 44, 48, 51, 55, 62], \\ &[1, 3, 5, 6, 8, 11, 17, 18, 22, 26, 31, 37, 40, 55, 62]]; \end{aligned}$$

$$\begin{aligned} &[[0, 2, 6, 8, 12, 17, 20, 22, 24, 31, 33, 36, 40, 43, 51, 55, 62], \\ &[1, 2, 3, 5, 8, 9, 10, 13, 16, 24, 29, 31, 51, 55, 62], \\ &[1, 4, 5, 6, 9, 12, 13, 20, 22, 31, 36, 43, 47, 51, 62]]; \end{aligned}$$

$$\begin{aligned} &[[0, 1, 2, 6, 9, 10, 12, 16, 26, 29, 31, 33, 36, 43, 44, 48, 62], \\ &[2, 4, 5, 6, 13, 16, 29, 31, 33, 36, 40, 43, 44, 51, 62], \\ &[1, 3, 6, 8, 9, 12, 13, 17, 20, 22, 31, 43, 51, 55, 62]]. \end{aligned}$$

For instance the difference family of the first example is explicitly:

$$\begin{aligned} X_0 &= \{0, 1, 3, 4, 7, 9, 11, 15, 16, 17, 18, 20, 23, 25, 26, 28, 31, 35, 38, 39, \\ &\quad 40, 44, 45, 48, 49, 53, 54, 55, 58, 62, 65, 67, 68, 70, 73, 75, 76, 77, 78, \\ &\quad 82, 84, 86, 89, 90, 92\}, \\ X_1 &= \{2, 3, 9, 13, 15, 16, 20, 22, 28, 29, 31, 34, 35, 36, 38, 39, 41, 43, 44, \\ &\quad 45, 46, 48, 49, 50, 51, 52, 54, 62, 63, 65, 66, 69, 74, 75, 77, 79, 83, 84, \\ &\quad 85, 87, 91\}, \\ X_2 &= X_1, \\ X_3 &= \{1, 2, 3, 8, 10, 13, 14, 15, 17, 19, 23, 25, 26, 31, 34, 36, 37, 40, 41, \\ &\quad 43, 46, 50, 51, 52, 53, 61, 62, 63, 64, 66, 67, 68, 69, 70, 71, 75, 76, 87, \\ &\quad 88, 91, 92\}. \end{aligned}$$

In all four difference families the block X_0 is symmetric i.e. $-X_0 = X_0$ in \mathbf{Z}_{93} and the representative 0 represents the trivial H -orbit, namely $\{0\}$.

4. ACKNOWLEDGEMENTS

This research was enabled in part by support provided by SHARCNET (<http://www.sharcnet.ca>) and the Digital Research Alliance of Canada (alliancecan.ca).

REFERENCES

- [1] C. Bright, D. Ž. Đoković, I. Kotsireas, V. Ganesh, A SAT+CAS Approach to Finding Good Matrices: New Examples and Counterexamples, The Thirty-Third AAAI Conference on Artificial Intelligence (AAAI-19)
- [2] R. Craigen and H. Kharaghani, Hadamard matrices and Hadamard designs, in Handbook of Combinatorial Designs, 2nd ed. C. J. Colbourn, J. H. Dinitz (eds) pp. 273–280. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, 2007.
- [3] D. Ž. Đoković, Ten new orders for Hadamard matrices of skew type, Univ. Beograd, Publ. Elektrotehn. Fak. Ser. Mat. 3 (1992), 47–59.
- [4] D. Ž. Đoković, Five new orders for Hadamard matrices of skew type. Australasian J. Combin. 10 (1994), 259–264.
- [5] D. Ž. Đoković, Skew-Hadamard matrices of orders 188 and 388 exist. International Mathematical Forum, 3, 22 (2008), 1063–1068.
- [6] D. Ž. Đoković, Skew-Hadamard matrices of orders 436, 580, and 988 exist, J. Combin. Designs, 16 (2008), 493–498.
- [7] D. Ž. Đoković, Some new symmetric Hadamard matrices. Informatsionno-upravliaiushchie sistemy [Information and Control Systems]. 2022 (2): 1–10. Saint-Petersburg State University of Aerospace Instrumentation, Russian Federation.
- [8] D. Ž. Đoković and I. S. Kotsireas, D-optimal matrices of orders 118, 138, 150, 154 and 174. In: C. J. Colbourn (ed.) Algebraic Design Theory and Hadamard Matrices, pp. 71–82, ADTHM, Lethbridge, Alberta, Canada, July 2014. Springer Proceedings in Mathematics & Statistics, vol. 133. Springer 2015.
- [9] R. J. Fletcher, C. Koukouvinos and J. Seberry, New skew-Hadamard matrices of order $4 \cdot 59$ and new D-optimal designs of order $2 \cdot 59$, Discrete Math. 286 (2004), 251–253.
- [10] J. Seberry, M. Yamada, Hadamard matrices, sequences, and block designs. In Contemporary design theory, 431–560, Wiley-Intersci. Ser. Discrete Math. Optim., Wiley, New York, 1992.
- [11] J. Seberry, M. Yamada, Hadamard Matrices, Constructions using Number Theory and Algebra, 2022 John Wiley & Sons, Inc.
- [12] . G. Szekeres, Tournaments and Hadamard matrices, Enseignement Math. 15 (1969), 269–278.

UNIVERSITY OF WATERLOO, DEPARTMENT OF PURE MATHEMATICS, WATERLOO, ONTARIO, N2L
3G1, CANADA
Email address: dragomir@rogers.com