Criticality in Sperner's Lemma

Tomáš Kaiser 1

Matěj Stehlík 2

Riste Škrekovski³

To the memory of Frank H. Lutz (1968–2023)

Abstract

We answer a question posed by T. Gallai in 1969 concerning criticality in Sperner's lemma, listed as Problem 9.14 in the collection of Jensen and Toft [Graph coloring problems, John Wiley & Sons, Inc., New York, 1995].

Sperner's lemma states that if a labelling of the vertices of a triangulation of the d-simplex Δ^d with labels $1, 2, \ldots, d+1$ has the property that (i) each vertex of Δ^d receives a distinct label, and (ii) any vertex lying in a face of Δ^d has the same label as one of the vertices of that face, then there exists a rainbow facet (a facet whose vertices have pairwise distinct labels). For $d \leq 2$, it is not difficult to show that for every facet σ , there exists a labelling with the above properties where σ is the unique rainbow facet. For every $d \geq 3$, however, we construct an infinite family of examples where this is not the case, which implies the answer to Gallai's question as a corollary. The construction is based on the properties of a 4-polytope which had been used earlier to disprove a claim of T. S. Motzkin on neighbourly polytopes.

¹Department of Mathematics and European Centre of Excellence NTIS (New Technologies for the Information Society), University of West Bohemia, Pilsen, Czech Republic. E-mail: kaisert@kma.zcu.cz. Supported by project GA20-09525S of the Czech Science Foundation.

²Université Paris Cité, CNRS, IRIF, F-75006, Paris, France. E-mail: matej@irif.fr. Partially supported by ANR project DISTANCIA (ANR-17-CE40-0015).

³Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana & Faculty of Information Studies, 8000 Novo Mesto, Slovenia. E-mail: skrekovski@gmail.com. Partially supported by the Slovenian Research Agency Program P1-0383, Project J1-3002, and BI-FR/22-23-Proteus-011.

1 Introduction

A central result of combinatorial topology, Sperner's lemma [21] has found wide-ranging applications in diverse areas of mathematics and beyond; for example, it can be used to give an elementary proof of Brouwer's fixed point theorem [18, p. 117] from algebraic topology, and to solve various fair-division problems, such as rental harmony [23, 24].

Let Δ^d be a *d*-simplex with vertices v_1, \ldots, v_{d+1} , and let K be a finite triangulation of Δ^d (see Section 2 for a precise definition). A Sperner labelling of K is a map $\lambda : V(K) \to \{1, \ldots, d+1\}$ such that $\lambda(v_i) = i$, and every vertex lying in a face of Δ^d has the same label as one of the vertices of that face. A simplex is said to be rainbow if its vertices receive pairwise distinct labels. The following classical result, proved by Sperner [21], is usually referred to as Sperner's lemma (see e.g. [5, p. 423], [6, p. 3]).

Theorem 1.1 (Sperner's lemma). If K is a finite triangulation of the dsimplex Δ^d and $\lambda : V(K) \rightarrow \{1, \ldots, d+1\}$ is any Sperner labelling of K, then there is a rainbow facet.

A natural question asked by Gallai in 1969 (see Problem 9.14 in Jensen and Toft [12] and Problem 3.2 below) concerns 'criticality' in Sperner's lemma. His question contains the following subproblem.

Problem 1.2. Given any triangulation K of Δ^d , and any facet $\sigma \in K$, is there a Sperner labelling of K where σ is the unique rainbow facet?

We will see later that if one could answer Problem 3.2 in the affirmative, then the same would follow for Problem 1.2. In dimension 1, the answer to Problem 1.2 is is easily seen to be positive. The same can be shown in dimension 2, as follows. (Gallai in fact obtained a slightly stronger result, settling Problem 3.2 in the affirmative in dimension 2; see [12, Problem 9.14].)

Theorem 1.3. For every triangulation K of Δ^2 and every facet $\sigma \in K$, there exists a Sperner labelling of K where σ is the unique rainbow facet.

Proof. Let K be a triangulation of the 2-simplex Δ^2 ; add edges and faces as shown in Figure 1 to make it into a triangulation K' of S^2 . Let τ be the 2-simplex corresponding to the outer face, and assume that $\sigma \in K$ is a given facet. Every planar triangulation is 3-connected, so by Steinitz's theorem [16, p. 63], it is the graph of a convex polyhedron, and the dual graph is also 3-connected. Hence, by Menger's theorem [16, p. 14], there exist three internally vertex-disjoint paths between σ and τ in the dual graph. The union of these paths (in the given embedding) has three faces; let us assign



Figure 1: A triangulation K of Δ^2 (in gray) extended to a triangulation K' of S^2 (thick edges and white faces; the outer face is labelled τ). Three internally vertex-disjoint paths (dashed) in the dual graph partition the vertices into three subsets (red, green and blue).

the vertices of K' embedded in the *i*-th face $(1 \le i \le 3)$ label *i*. It is not hard to see that this determines a Sperner labelling of K. Furthermore, if a 3-cycle of K' contains vertices of all three labels and differs from the face boundaries of σ and τ , then it separates a vertex of σ from a vertex of τ , so it is not a face boundary. Thus, σ is the unique rainbow facet in the Sperner labelling of K, as claimed.

In this paper, we will show that the answer to Problem 1.2 (and consequently, to Problem 3.2) is negative in dimension 3 and higher. Namely, we will prove the following theorem in Section 5.

Theorem 1.4. For every integer $d \geq 3$, there exist infinitely many triangulations K of Δ^d such that there exists a facet $\sigma \in K$ with the property that every Sperner labelling of K has a rainbow facet distinct from σ .

2 Terminology and preliminary results

For topological and graph-theoretic concepts not defined here, the reader is referred to Munkres [18] and to Bondy and Murty [3], respectively. An introduction to Sperner's lemma—and combinatorial topology in general can be found in de Longueville [6]. The book by Mohar and Thomassen [16] provides an introduction to topological graph theory. For polytope theory, we refer the reader to Grünbaum [10] and to Ziegler [26]. A k-colouring of a graph G is a function $c: V(G) \to \{1, \ldots, k\}$ such that $c(u) \neq c(v)$ whenever $uv \in E(G)$. The minimum k for which a k-colouring of G exists is known as the chromatic number of G, denoted by $\chi(G)$. If $\chi(G) \leq k$ or $\chi(G) = k$, then G is said to be k-colourable or k-chromatic, respectively. Furthermore, a graph G is said to be k-critical (resp. k-vertex-critical) if G is k-chromatic and deleting any edge (resp. vertex) results in a (k-1)-colourable graph. A graph is said to be k-vertex-connected if it has at least k + 1 vertices, and removing any set of k - 1 vertices does not disconnect the graph.

A (geometric) simplicial complex K is a collection of geometric simplices of various dimensions satisfying two conditions: firstly, if $\sigma \in K$ and τ is a face of σ , then $\tau \in K$; secondly, if $\sigma, \tau \in K$ and $\sigma \cap \tau \neq \emptyset$, then $\sigma \cap \tau$ is a face of both σ and τ . We denote the regular d-dimensional simplex, unit d-ball and unit d-sphere by Δ^d , B^d and S^d , respectively. Simplices of K that are not strictly contained in other simplices are the facets of K. The union of all simplices of K, known as the polyhedron of K, is denoted by |K|. The kskeleton of a simplicial complex K, denoted $K^{(k)}$, consists of all the simplices of K of dimension at most k; note that $K^{(1)}$ consists of vertices and edges, i.e., it is a graph. If two topological spaces X and Y are homeomorphic, we write $X \cong Y$. A simplicial complex K such that $X \cong |K|$, if one exists, is called a triangulation of X. The join of two simplicial complexes K and L is denoted by $K \star L$.

A quadrangulation of a surface (i.e., a 2-dimensional manifold) X is a graph embedded in X so that every face of the embedding is bounded by a 4cycle. In [13], the definition of quadrangulations was extended to triangulable topological spaces in the following way. A graph G is a quadrangulation of a topological space X if G is a spanning subgraph of the 1-skeleton $K^{(1)}$ of some triangulation K of X, with the property that the vertices of every facet of K induce a complete bipartite subgraph of G with at least one edge. This definition coincides with the usual definition in the case when X is a 2dimensional manifold. In [13, Lemma 3.2], the following characterisation was given of non-bipartite quadrangulations of the d-dimensional real projective space \mathbb{RP}^d :

Lemma 2.1. A graph G is a non-bipartite quadrangulation of \mathbb{RP}^d if and only if there exists a centrally symmetric triangulation K of S^d and a 2-colouring of the vertices with colours black and white, such that no pair of antipodal vertices is adjacent, antipodal vertices receive different colours, every facet of K contains a black and a white vertex, and G is obtained from the 1-skeleton $K^{(1)}$ by first deleting all monochromatic edges, and then identifying antipodal vertices and edges.



Figure 2: A triangulation K (dashed edges) of a 2-simplex (in gray), and the graph G_K (solid edges).

If P is a (convex) d-polytope, its (d-1)-dimensional faces are its facets. If every facet is a simplex, then the boundary of P, denoted by ∂P , is a simplicial complex such that $|\partial P| \cong S^{d-1}$, i.e., ∂P is a triangulation of S^{d-1} . The convex hull of the vectors of the standard orthonormal basis and their negatives in \mathbb{R}^d is known as the the d-dimensional cross polytope.

3 Relation to chromatic number

Gallai used Sperner's lemma to construct a family of 4-chromatic planar graphs; see [12, Problem 9.14] or [19]. Let K be a triangulation of Δ^d , and define the graph G_K as follows (see Figure 2 for an illustration). There are three types of vertices of G_K :

- (V1) The vertices u_1, \ldots, u_n of K;
- (V2) A vertex v_{ρ} for every d-simplex $\rho \in K$;
- (V3) A vertex w_{σ} for every (d-1)-simplex $\sigma \in \Delta^d$.

The edge set of G_K is defined as $E(G_K) = E_1 \cup E_2 \cup E_3$, where:

- (E1) $u_i v_\rho \in E_1$ if $u_i \in \rho$, where u_i is a vertex of K and ρ is a d-simplex of K;
- (E2) $u_i w_\sigma \in E_2$ if u_i belongs to the facet σ of Δ^d ;

(E3) $w_{\sigma}w_{\tau} \in E_3$ for any pair of distinct (d-1)-simplices $\sigma, \tau \in \Delta^d$.

Note that the vertices of type (V1) and (V2) form independent sets in G_K , the vertices of type (V3) induce a (d + 1)-clique in G_K , and there are no edges between the vertices of type (V2) and of type (V3). It follows that G_K is (d + 2)-colourable—for example, by colouring the clique induced by (V3) using colours $1, \ldots, d+1$, the vertices of type (V2) by colour 1, and the vertices of type (V1) by colour d + 2.

What is more interesting is that G_K is not (d+1)-colourable (and therefore G_K is (d+2)-chromatic). Indeed, Gallai observed that the statement that G_K is not (d+1)-colourable is equivalent to Sperner's lemma; we formalise it as the following proposition.

Proposition 3.1. G_K is not (d + 1)-colourable if and only if every Sperner labelling of K has a rainbow facet.

Proof. Suppose for a contradiction that there exists a Sperner labelling λ of K with no rainbow facet. Then we can define a (d + 1)-colouring c of G_K as follows. For each vertex u_i of type (V1), let $c(u_i) = \lambda(u_i)$. For each vertex w_{σ} of type (V3), let u_j be the unique vertex of Δ^d not belonging to σ , and set $c(w_{\sigma}) = \lambda(u_j)$. Finally, each vertex v_{ρ} of type (V2) is adjacent to the d + 1 vertices of a facet $\rho \in K$, which is not rainbow by assumption. Hence, the colouring c can be extended to a (d + 1)-colouring of G_K .

Conversely, suppose that there is a (d+1)-colouring of G_K . The vertices of G_K of type (V3) ensure that the labelling of K induced by the colouring is a Sperner labelling, and there is no rainbow facet in K since the vertices of every facet in K are adjacent to a vertex of type (V2) in G_K .

Observe that if K is a triangulation of the 2-dimensional simplex, then G_K is a 4-chromatic planar graph with exactly 4 triangles. This is best possible in view of the sharpening of Grötzsch's theorem due to Grünbaum [9] and Aksenov [1], which states that every planar graph with at most 3 triangles is 3-colourable. Indeed, Borodin et al. [4] characterised the 4-chromatic planar graphs with exactly 4 triangles; in this characterisation, the graphs G_K belong among those graphs that can be obtained from the complete graph on 4 vertices by replacing a single vertex of degree 3 by a 'critical' patch (in a sense made precise in [4]).

It is easy to see that for any triangulation K of the 1-simplex, the graph G_K is an odd cycle and thus 3-critical. Gallai [12, 19] proved that G_K is 4-critical whenever K is a triangulation of the 2-simplex; we note that this also follows from Theorem 4.1 in conjunction with [8, Theorem 5.4] (see the discussion at the end of Section 4).

In 1969 Gallai asked the following question, presented as Problem 9.14 by Jensen and Toft [12]. The question is also mentioned in [19, 20].

Problem 3.2. If $d \ge 3$, and K is a triangulation of Δ^d , is the graph G_K (d+2)-critical?

The following observation shows that a positive answer to Problem 3.2 would imply a positive answer to Problem 1.2.

Observation 3.3. Let K be a triangulation of Δ^d . If G_K is (d+2)-critical, then for any facet $\sigma \in K$, there exists a Sperner labelling of K where σ is the unique rainbow facet.

Proof. Suppose G_K is (d+2)-critical, for some triangulation K of Δ^d . Then for any facet $\sigma \in K$, the graph $G_K - v_\sigma$ is (d+1)-colourable. Hence, the labelling of K induced by the (d+1)-colouring of $G - v_\sigma$ is a Sperner labelling with no rainbow facet distinct from σ .

It is easy to see that Observation 3.3, in conjunction with Theorem 1.4, immediately settles Gallai's question in the negative.

Theorem 3.4. For every $d \ge 3$, there exist infinitely many triangulations K of Δ^d such that G_K is not (d+2)-critical.

4 G_K as projective quadrangulations

Jensen and Toft [12, Problem 9.14] also raised the question of whether the graphs G_K belong to a 'larger class of (d + 2)-chromatic graphs defined in purely graph-theoretic terms'. While we cannot do away with the topology, we can show that the graphs G_K belong to a larger class of graphs whose chromatic number is large for 'topological reasons', namely the class of quadrangulations of projective spaces. Indeed, a result by the first two authors [13, Theorem 1.1], generalising an earlier result of Youngs [25], shows that all non-bipartite quadrangulations of the *d*-dimensional projective space \mathbb{RP}^d are at least (d + 2)-chromatic.

Theorem 4.1. If K is a triangulation of the d-simplex, then G_K is a nonbipartite quadrangulation of the d-dimensional projective space \mathbb{RP}^d .

Proof. Let P be the 2-coloured complex obtained from the boundary complex ∂Q^{d+1} of the (d+1)-dimensional cross polytope Q^{d+1} by colouring all the vertices of one facet (say α_w) white, and all the vertices of the complementary



Figure 3: The complexes $P - \alpha_w$ (left), $P - \alpha_w - \alpha_b$ (centre) and $P - \alpha_b$ (right) in the proof of Theorem 4.1 (case d = 2).

facet (say α_b) black. The complex $P - \alpha_w - \alpha_b$ obtained by deleting the facets α_w and α_b from P contains $\partial \alpha_w$ and $\partial \alpha_b$ as subcomplexes. Similarly, $P - \alpha_w$ and $P - \alpha_b$ contain $\partial \alpha_w$ and $\partial \alpha_b$, respectively (see Figure 3 for an illustration). Glue $P - \alpha_w - \alpha_b$ to $P - \alpha_w$ along $\partial \alpha_w$ (by identifying the corresponding vertices of $\partial \alpha_w$), and then glue the result to $P - \alpha_b$ along $\partial \alpha_b$. The resulting complex C is a centrally symmetric triangulation of S^d . Note that C has a unique facet α'_w with all vertices white (coming from the facet α_w of $P - \alpha_b$) and a unique facet α'_b with all vertices black.

Since K is a subdivision of a d-simplex Δ^d , there is a sequence of stellar moves (see e.g. [14, Theorem 4.5]) that transforms Δ^d into K. Assign to each vertex v of the all-white facet α'_w of C a vertex of Δ^d using an arbitrary bijection, and assign the same vertex of Δ^d to the antipode of v. Perform the corresponding sequence of stellar moves on α'_w , colouring all new vertices white. Proceed analogously for α'_b with colours reversed. Finally, perform a stellar subdivision of each all-white facet with a black vertex, and of each all-black facet with a white vertex. Call the resulting 2-coloured complex \tilde{K} .

Since K is a centrally symmetric subdivision of the centrally symmetric triangulation C of S^d , it follows that \tilde{K} is a centrally symmetric triangulation of S^d . Moreover, \tilde{K} is equipped with an antisymmetric 2-colouring such that no facet is monochromatic. Therefore, by Lemma 2.1, the graph G obtained from the 1-skeleton of \tilde{K} by deleting monochromatic edges, and by identifying antipodal vertices, is a non-bipartite quadrangulation of \mathbb{RP}^d . We claim that $G \cong G_K$. To see this, note that \tilde{K} contains three types of vertices, each appearing in antipodal pairs consisting of a white vertex and a black vertex:

(W1) vertices in the black and white copies of K not contained in $P - \alpha_w - \alpha_b$;

(W2) vertices subdividing each facet of the black or white copy of K;

(W3) vertices in $P - \alpha_w - \alpha_b$.

There is an obvious bijection between the vertices of G and those of G_K : the vertices of G of type (W1), (W2) and (W3) are in bijection with the vertices

of G_K of type (V1), (V2) and (V3), respectively. It is not hard to check that the edges of G are in bijection with the edges of G_K .

Non-bipartite quadrangulations of \mathbb{RP}^1 are odd cycles, so they are 3critical. Youngs [25] showed that non-bipartite quadrangulations of \mathbb{RP}^2 are 4-chromatic, and conjectured that they are 4-critical as long as they contain no non-bipartite quadrangulation of \mathbb{RP}^2 as a proper subgraph; this was subsequently proved by Gimbel and Thomassen [8]. For $d \geq 3$, non-bipartite quadrangulations of \mathbb{RP}^d are not (d + 2)-critical in general; indeed, their chromatic number is unbounded [13].

5 Proof of the main result

Our construction uses a specific convex 4-polytope known as H_8 and described in [10, Section 7.2]. This polytope is 2-neighbourly (recall that a polytope is *t*-neighbourly if every set of *t* vertices constitutes a face). Motzkin [17] claimed that for even *d*, every d/2-neighbourly *d*-polytope is combinatorially equivalent to a cyclic polytope (see [10] for the necessary definitions). This assertion is true for *d*-polytopes with at most d + 3 vertices [7] but fails in general; H_8 is one of the two counterexamples for d = 4 constructed in [10]. Our example is in line with Sturmfels' remark [22] that d + 4 vertices seem to be a 'threshold of counterexamples'.

The convex 4-polytope H_8 has 8 vertices A, B, C, D, E, F, G, Z and the following are the 20 facets of its boundary complex.

ABCD	ABCG	<u>ABDE</u>	ABEF	ABFZ
ABGZ	ACDZ	ACGZ	ADEZ	AEFZ
BCDE	BCEF	BCFG	BFGZ	CDEF
CDFG	CDGZ	DEFG	DEGZ	EFGZ

We underlined the facets relevant for the proof of Lemma 5.1 below. The two facets in bold play a key role in the proof of Theorem 1.4, and it will be convenient to name them. Throughout the remainder of this paper, we let $\sigma_0 := ABGZ$ and $\sigma_1 := CDEF$.

Lemma 5.1. Let the vertices of H_8 be labelled by 1, 2, 3, 4 in such a way that the facets σ_0 and σ_1 of ∂H_8 are rainbow. Then there exists another rainbow facet of ∂H_8 .

Proof. We proceed by contradiction. Let φ be the given labelling. Suppose that σ_0 and σ_1 are the only rainbow facets of ∂H_8 , and assume (without loss of generality) that the vertices A, B, G, Z are labelled 1, 2, 3, 4 in order. Observe the following direct consequence of the assumptions.

Claim 1. If a facet τ of ∂H_8 shares a 2-dimensional face with σ_i (for some $i \in \{0, 1\}$), then the two vertices in the symmetric difference of τ and σ_i have different labels.

Applying Claim 1 to the facet ABCG (and σ_0) we find $\varphi(C) \neq \varphi(Z) = 4$. Similarly, $\varphi(C) \neq 2$ (considering ACGZ and σ_0) and $\varphi(C) \neq 3$ (considering DEFG and σ_1). Consequently, $\varphi(C) = 1$.

Neither *E* nor *F* have label 3, as seen by applying Claim 1 to the facets *CDFG* and *ABFZ*, respectively. Since σ_1 is rainbow, these vertices do not have label 1 either, since it is already taken by *C*. Thus, $\{\varphi(E), \varphi(F)\} = \{2, 4\}$. This implies that $\varphi(D) = 3$.

The facet ABDE shares two vertices A, B with σ_0 . Since ABDE is not rainbow, we deduce that $\{\varphi(D), \varphi(E)\} \neq \{\varphi(G), \varphi(Z)\} = \{3, 4\}$. Hence, $\varphi(E) \neq 4$ as we know that $\varphi(D) = 3$. This leaves only the option $\varphi(E) = 2$ for the label of E. But then the facet ADEZ is rainbow, a contradiction. \Box

We remark that H_8 is far from the only polytope satisfying an analogue of Lemma 5.1. While it is the only such polytope among 4-polytopes on 8 vertices which are *simplicial* (all their faces are simplices; these polytopes are listed in [11]), the situation is much different when the number of vertices is increased. For instance, a complete search of the 23 2-neighbourly 4polytopes on 9 vertices listed in [2] showed that 18 of them have the same property as H_8 .

Proof of Theorem 1.4. Let K be the simplicial complex obtained from the boundary complex of H_8 by deleting the simplex σ_0 . Since H_8 is a 4-polytope, we have $H_8 \cong B^4$, so $\partial H_8 \cong S^3$, and therefore $K \cong B^3$, or, equivalently, K a triangulation of the 3-simplex with vertices A, B, G, Z.

For an integer $d \geq 3$, let ρ be a (d-4)-simplex, and define $K_d = K \star \rho$ (we take the convention that the empty simplex has dimension -1; in particular, $K_3 = K$). We can view K_d as a triangulation of the *d*-simplex with vertices $\{A, B, G, Z\} \cup V(\rho)$. We will show that in any Sperner labelling of K_d , there is a rainbow facet distinct from $\sigma := \sigma_1 \star \rho$. Fix any Sperner labelling $\lambda : V(K_d) \to \{1, \ldots, d+1\}$ of K_d (observe that ρ is rainbow with respect to λ). If σ is not rainbow, then by Sperner's lemma, there must be a rainbow facet distinct from σ_1 receive the same set of labels. In particular, the labelling λ' of V(K) induced by λ uses 4 labels. By Lemma 5.1, some 3-simplex $\tau \in K$ distinct from σ_1 is rainbow with respect to λ' . Hence, $\tau \star \rho$ is a rainbow *d*-simplex with respect to λ . We have thus shown that any Sperner labelling of K_d has a rainbow *d*-simplex distinct from σ . This provides a negative answer to Gallai's question for all $d \geq 3$.

It remains to show that, for every $d \geq 3$, there are infinitely many triangulations of Δ^d with this property. Consider any facet $\pi \in K_d$ distinct from σ , and let L be a triangulation of π such that $\partial L = \partial \pi$. Let K'_d be the complex obtained from K_d by replacing π by L. Fix any Sperner labelling $\lambda : V(K'_d) \to \{1, \ldots, d+1\}$, and suppose that no facet of K'_d distinct from σ is rainbow. The d+1 vertices in ∂L must receive labels $1, \ldots, d+1$. Therefore, the labelling induced by L is a Sperner labelling of L, so by Sperner's lemma, there is at least one rainbow facet in L. This shows that any Sperner labelling of K'_d has a rainbow d-simplex distinct from σ .

By iterating this construction, we can obtain infinitely many triangulations K of Δ^d with the property that every Sperner labelling of K has a rainbow d-simplex distinct from σ .

It may well be that even in higher dimensions, the property in Problem 1.2 holds for triangulations of a special type. For example, one could take cyclic polytopes (a simple and well-understood class) and consider the triangulations associated with them in the following sense. Given a polytope P, let us say that a triangulation is *associated with* P if it is obtained from the boundary of P by removing a facet τ and projecting the rest of the boundary into the convex hull of the vertex set of τ .

Conjecture 5.2. Problem 1.2 has an affirmative answer if K is a triangulation of Δ^d associated with a d-dimensional cyclic polytope.

Acknowledgements

We are indebted to Frank Lutz for the valuable information on small triangulations of manifolds collected in [15]. We used Sage to test these triangulations (as well as those obtained from [2, 11]) for specific properties, and we thank the developers of this fine piece of software. Finally, we would like to thank the referees for making some useful suggestions to improve the readability of the paper.

References

- V. A. Aksenov. On continuation of 3-colouring of planar graphs (in Russian). Diskret. Anal. Novosibirsk, 26:3–19, 1974.
- [2] A. Altshuler and L. Steinberg. Neighborly 4-polytopes with 9 vertices. J. Combin. Theory Ser. A, 15:270–287, 1973.

- [3] J. A. Bondy and U. S. R. Murty. *Graph Theory*, volume 244 of *Graduate Texts in Mathematics*. Springer, New York, 2008.
- [4] O. V. Borodin, Z. Dvořák, A. V. Kostochka, B. Lidický, and M. Yancey. Planar 4-critical graphs with four triangles. *European J. Combin.*, 41:138–151, 2014.
- [5] J. A. De Loera, X. Goaoc, F. Meunier, and N. H. Mustafa. The discrete yet ubiquitous theorems of Carathéodory, Helly, Sperner, Tucker, and Tverberg. Bull. Amer. Math. Soc. (N.S.), 56(3):415–511, 2019.
- [6] M. de Longueville. A course in Topological Combinatorics. Universitext. Springer, New York, 2013.
- [7] D. Gale. Neighborly and cyclic polytopes. *Convexity* (2nd ed.), Proc. Symp. Pure Math., 7:225–232, 1963.
- [8] J. Gimbel and C. Thomassen. Coloring graphs with fixed genus and girth. Trans. Amer. Math. Soc., 349(11):4555–4564, 1997.
- [9] B. Grünbaum. Grötzsch's theorem on 3-colorings. Michigan Math. J., 10:303–310, 1963.
- [10] B. Grünbaum. *Convex Polytopes*, volume 221 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 2003. Prepared and with a preface by Volker Kaibel, Victor Klee and Günter M. Ziegler.
- [11] B. Grünbaum and V. P. Sreedharan. An enumeration of simplicial 4polytopes with 8 vertices. J. Combin. Theory, 2:437–465, 1967.
- [12] T. R. Jensen and B. Toft. Graph Coloring Problems. Wiley-Interscience Series in Discrete Mathematics and Optimization. John Wiley & Sons, Inc., New York, 1995. A Wiley-Interscience Publication.
- [13] T. Kaiser and M. Stehlík. Colouring quadrangulations of projective spaces. J. Combin. Theory Ser. B, 113:1–17, 2015.
- [14] W. B. R. Lickorish. Simplicial moves on complexes and manifolds. In Proceedings of the Kirbyfest (Berkeley, CA, 1998), volume 2 of Geom. Topol. Monogr., pages 299–320. Geom. Topol. Publ., Coventry, 1999.
- [15] F. Lutz. The Manifold Page. Available at https://page.math.tu-berlin.de/~lutz/stellar, retrieved on December 21, 2022.

- [16] B. Mohar and C. Thomassen. Graphs on Surfaces. Johns Hopkins Studies in the Mathematical Sciences. Johns Hopkins University Press, Baltimore, MD, 2001.
- [17] T. S. Motzkin. Comonotone curves and polyhedra. Abstract 111. Bull. Amer. Math. Soc., 63:35, 1957.
- [18] J. R. Munkres. Elements of Algebraic Topology. Addison-Wesley Publishing Company, Menlo Park, CA, 1984.
- [19] F. Nielsen and B. Toft. On a class of planar 4-chromatic graphs due to T. Gallai. In *Recent Advances in Graph Theory (Proc. Second Czechoslo*vak Sympos., Prague, 1974), pages 425–430. Academia, Prague, 1975.
- [20] V. Rödl and Zs. Tuza. On color critical graphs. J. Combin. Theory Ser. B, 38:204–213, 1985.
- [21] E. Sperner. Neuer Beweis für die Invarianz der Dimensionszahl und des Gebietes. Abh. Math. Sem. Univ. Hamburg, 6(1):265–272, 1928.
- [22] B. Sturmfels. Some applications of affine Gale diagrams to polytopes with few vertices. SIAM J. Discrete Math., 1(1):121–133, 1988.
- [23] F. E. Su. Rental harmony: Sperner's lemma in fair division. Amer. Math. Monthly, 106(10):930–942, 1999.
- [24] A. Sun. To divide the rent, start with a triangle. *New York Times*, April 29, 2014.
- [25] D. A. Youngs. 4-chromatic projective graphs. J. Graph Theory, 21(2):219–227, 1996.
- [26] G. M. Ziegler. Lectures on Polytopes, volume 152 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1995.