

A NOTE ON THE LIMIT LAW OF ONE-SIDED TREE DESTRUCTION

MARKUS KUBA AND ALOIS PANHOLZER

ABSTRACT. This short note serves an addendum to the article "Destruction of very simple trees" by Fill, Kapur and Panholzer (2004). Therein, the limit law of one-sided tree destruction was determined by its moment sequence. We add an identification of the limit law, using recent results of Bertoin (2022), in terms of the local time of a noise reinforced Bessel process.

1. ADDENDUM: IDENTIFICATION OF THE LIMIT LAW

In Theorem 5.1 of [4] the limit law of the cost of root isolation Y_n in the class of very simple trees, a subclass of simply generated trees, was obtained. The random variable Y_n itself satisfies the distributional equation

$$Y_n \stackrel{\mathcal{L}}{=} Y_{K_n} + t_n \quad n \geq 2,$$

for some random variable K_n [4], with initial value $Y_1 = t_1$ and toll function $t_n = n^a$, with $a \geq 0$. It was shown that for a parameter $\sigma > 0$ and $a' = a + \frac{1}{2}$ it holds

$$\frac{Y_n}{\sigma n^{a'}} \xrightarrow{\mathcal{L}} Y_{a'},$$

where $Y_{a'}$ is uniquely determined by its moment sequence $m_s = \mathbb{E}(Y_{a'}^s)$, for positive integers $s \geq 1$:

$$m_s = \frac{s!}{2^{s/2}} \prod_{k=1}^s \frac{\Gamma(ka')}{\Gamma(ka' + \frac{1}{2})}.$$

Recent results of Bertoin [2] on the noise reinforced Bessel process and its local time allows us to observe the following Corollary to Theorem 5.1 of [4].

Corollary 1. *The limit law $Y_{a'}$ has the same distribution as a moment-shifted (also called size biased or moment-tilted) \hat{L}_t of a local time of a noise reinforced Bessel process, scaled by $\frac{1}{\sqrt{2}}$:*

$$Y_{a'} \stackrel{\mathcal{L}}{=} \frac{1}{\sqrt{2}} \cdot \text{tilt}_1(\hat{L}_t / \kappa(p, t)),$$

with factor $\kappa(p, t)$ given in (1), dimension $d = 1$ and reinforcement parameter $p = \frac{1}{2} - \frac{1}{4a'}$.

Additionally, $Y_{a'} \stackrel{\mathcal{L}}{=} \frac{1}{\kappa(p, 1)\sqrt{2}} \cdot \hat{I}$, where the scaled exponential functional \hat{I} is given by

$$\hat{I} = \int_0^\infty \exp(-\frac{1}{2}\hat{\xi}_t) dt,$$

with $\hat{\xi}_t$ denoting the subordinator with Laplace-Exponent $\hat{\Phi}$, determined by

$$\hat{\Phi}(r) = 2^{-\frac{1}{2}} \left(\frac{1}{4a'}\right)^{\frac{1}{2}} \frac{\Gamma(\frac{1}{2})}{\frac{1}{2}B(\frac{1}{2}, 4a'r)}.$$

2000 *Mathematics Subject Classification.* 05C05, 05A15, 05A19.

Key words and phrases. Tree destruction, cutting down, local time, limit law.

Remark 1 (Moments of Gamma type). For $a' = \frac{1}{2}$ one obtains a Rayleigh distribution, as noted before [4]. For $m \in \mathbb{N}$ and $m \cdot a' = \frac{1}{2}$, i.e. $a' = \frac{1}{4}$, we observe cancellations and obtain interesting moments of Gamma type [5]. For example, for $a' = \frac{1}{4}$ we get

$$m_s = \frac{1}{2^{s/2}} \frac{\Gamma(\frac{1}{4})\Gamma(\frac{1}{2})\Gamma(s+1)}{\Gamma(\frac{s+1}{4})\Gamma(\frac{s+2}{4})}, \quad s \geq 1.$$

Density functions for such moment sequences can be obtained using inverse Mellin computations; compare with [1, Proof of Theorem 4.1] or see [5, Theorem 5.4].

Proof. The identification is directly done using moment sequences appearing in [2]. There, a noise reinforced Bessel process of dimension $d > 0$ and with reinforcement parameter $p \in (-\infty, 1/2)$ was considered. Set

$$\alpha = 1 - d/2 \in (0, 1) \text{ and } \beta = \frac{\alpha}{1 - 2p},$$

or equivalently

$$p = \frac{1}{2} - \frac{\alpha}{2\beta}, \text{ with } \beta > 0.$$

The local time \hat{L}_t of the noise reinforced Bessel process has power moments (see [2, Theorem 1.2]), which can be written as

$$\mathbb{E}(\hat{L}_t^s) = \kappa(p, t)^s \cdot \frac{1 - 2p}{\Gamma(1 + \alpha)} \cdot \Gamma(s) \cdot \prod_{j=1}^{s-1} \frac{\Gamma(j\beta)}{\Gamma(\alpha + j\beta)}.$$

with scale factor given by

$$\kappa(p, t) = \frac{(2t)^\alpha \cdot \Gamma(1 + \alpha)}{(1 - 2p)^\alpha \cdot \Gamma(1 - \alpha)}. \quad (1)$$

This implies that the scaled random variable $L = \hat{L}_t / \kappa(p, t)$ has moment sequence $\mu_s = \mathbb{E}(L^s)$, given by

$$\mu_s = \frac{1 - 2p}{\Gamma(1 + \alpha)} \cdot \Gamma(s) \cdot \prod_{j=1}^{s-1} \frac{\Gamma(j\beta)}{\Gamma(\alpha + j\beta)}, \quad s \geq 1.$$

Using standard results on moment shifts (see for example [1, Lemma 3.4]), also called size biased distributions or tilted distributions, we observe that the random variable $T = \text{tilt}(L)$ has moment sequence $\mathbb{E}(T^s) = \frac{\mu_{s+1}}{\mu_1}$, or more explicitly

$$\mathbb{E}(T^s) = \Gamma(s + 1) \cdot \prod_{j=1}^s \frac{\Gamma(j\beta)}{\Gamma(\alpha + j\beta)}.$$

Since $\Gamma(s + 1) = s!$, we finally set $\beta = a'$ and $\alpha = \frac{1}{2}$. This leads to the desired moment sequence. Finally, we note that the choices lead to a dimension $d = 1$ and $p = \frac{1}{2} - \frac{1}{4a'}$, such that $1 - 2p = \frac{1}{4a'}$.

Concerning the second statement we use the definition of \hat{I} and its Laplace exponent [2, Equations 4.3 and 4.4]. We use Corollary 4.3 of [2], which relates \hat{I} and \hat{L}_1 . Note that there is a small misprint [3] in the stated equation, as the prefactor

$$c = (1/2 - p)^{-\alpha} \frac{1 - 2p}{\Gamma(1 - \alpha)},$$

which can be rewritten into

$$\left(\frac{1 - 2p}{2}\right)^{-\alpha} \frac{1 - 2p}{\Gamma(1 - \alpha)} = \frac{2^\alpha (1 - 2p)^{1-\alpha}}{\Gamma(1 - \alpha)} = \mathbb{E}(\hat{L}_1),$$

should be replaced by its reciprocal. We state here the corrected version:

$$\mathbb{E}(f(\hat{I})) = \frac{1}{\mathbb{E}(\hat{L}_1)} \cdot \mathbb{E}(\hat{L}_1 f(\hat{L}_1)).$$

Choosing f as the power functions, we observe a tilt of the moments [1, Lemma 3.4], leading to the moment sequence of \hat{I} as the shifted moments of \hat{L}_1 . \square

Remark 2 (More moments of Gamma type). As already pointed out [2], for $p = 0$, such that $\alpha = \beta$, the random variable \hat{L}_t simplifies to a Mittag-Leffler distribution:

$$\mathbb{E}(\hat{L}_t^s) = \kappa(0, t)^s \cdot \frac{1}{\Gamma(1 + \alpha)} \cdot \Gamma(s) \cdot \frac{\Gamma(\alpha)}{\Gamma(s\alpha)} = \kappa(0, t)^s \frac{\Gamma(s + 1)}{\Gamma(s\alpha + 1)}.$$

The random variable T also leads to moments of Gamma types for small β , proportional to α : $\beta = \alpha/m$, $m \in \mathbb{N}$, leading to moments

$$\mathbb{E}(T^s) = \Gamma(s + 1) \cdot \prod_{j=1}^m \frac{\Gamma(\frac{j\alpha}{m})}{\Gamma(\frac{(s+j)\alpha}{m})}, \quad s \geq 1.$$

Again, density functions can be obtained for such moments.

2. OUTLOOK AND ACKNOWLEDGMENTS

The authors are currently investigating into other occurrences of such limit laws in combinatorial probability.

The authors warmly thank Jean Bertoin for feedback on his work and explanations.

REFERENCES

- [1] Cyril Banderier, Markus Kuba and Michael Wallner. [Phase transitions of composition schemes: Mittag-Leffler and Mixed Poisson distributions](#). *Submitted*, 2021.
- [2] Jean Bertoin. [On the local times of noise reinforced Bessel processes](#). *Annales Henri Lebesgue*, volume 5, pages 1277–1294, 2022.
- [3] Jean Bertoin. Personal communication. 2022.
- [4] James Allen Fill, Nevin Kapur and Alois Panholzer. [Destruction of very simple trees](#). *Algorithmica*, volume 46, pages 345–366, 2006.
- [5] Svante Janson. [Moments of Gamma type and the Brownian supremum process area](#). *Probability Surveys*, 7, 1–52, 2010.

MARKUS KUBA, DEPARTMENT APPLIED MATHEMATICS AND PHYSICS, UNIVERSITY OF APPLIED SCIENCES - TECHNIKUM WIEN, HÖCHSTÄDTPLATZ 5, 1200 WIEN

Email address: kuba@technikum-wien.at

ALOIS PANHOLZER, INSTITUT FÜR DISKRETE MATHEMATIK UND GEOMETRIE, TECHNISCHE UNIVERSITÄT WIEN, WIEDNER HAUPTSTR. 8-10/104, 1040 WIEN, AUSTRIA

Email address: Alois.Panholzer@tuwien.ac.at