

# Efficient Risk Estimation for the Credit Valuation Adjustment

Michael B. Giles\*    Abdul-Lateef Haji-Ali†    Jonathan Spence‡

## Abstract

**AMS Subject Classification:** 65C05, 62P05

**Keywords:** Multilevel Monte Carlo, Nested simulation, Risk estimation

The valuation of over-the-counter derivatives is subject to a series of valuation adjustments known as xVA, which pose additional risks for financial institutions. Associated risk measures, such as the value-at-risk of an underlying valuation adjustment, play an important role in managing these risks. Monte Carlo methods are often regarded as inefficient for computing such measures. As an example, we consider the value-at-risk of the Credit Valuation Adjustment (CVA-VaR), which can be expressed using a triple nested expectation. Traditional Monte Carlo methods are often inefficient at handling several nested expectations. Utilising recent developments in multilevel nested simulation for probabilities, we construct a hierarchical estimator of the CVA-VaR which reduces the computational complexity by 3 orders of magnitude compared to standard Monte Carlo.

## 1 Introduction

Abstractly, this paper constructs an efficient multilevel estimator [12] for triple nested expectations, of the kind which we later show arise in risk estimation involving the Credit Valuation Adjustment (CVA). We consider quantities of the form

$$\eta = \mathbb{E} \left[ \mathbb{H} \left( \mathbb{E} \left[ f \left( \mathbb{E} [g(X_T) | X_{t_1}], t_1 \right) \middle| X_{t_0} \right] \right) \right]. \quad (1)$$

Here,  $X_t$  is a stochastic process driven by an underlying  $d$ -dimensional SDE which is evaluated at (possibly random) times  $t_0 < t_1 < T$ ,  $f(x, y)$  and  $g(x)$  are Lipschitz or twice differentiable functionals of  $x$  and  $\mathbb{H}(x)$  is the Heaviside function taking value 1 if  $x \geq 0$  and value 0 otherwise. In many applications, including for CVA,  $f$  takes the form

$$f \left( \mathbb{E} [g(X_T) | X_{t_1}], t_1 \right) = \Lambda \left( \mathbb{E} [g(X_T) | X_{t_1}], t_1 \right) - L_\eta, \quad (2)$$

---

\*University of Oxford ([mike.giles@maths.ox.ac.uk](mailto:mike.giles@maths.ox.ac.uk)).

†Heriot-Watt University ([a.hajiali@hw.ac.uk](mailto:a.hajiali@hw.ac.uk)).

‡Corresponding author. Heriot-Watt University ([jws5@hw.ac.uk](mailto:jws5@hw.ac.uk)).

where  $\Lambda(\mathbb{E}[g(X_T) | X_{t_1}], t_1)$  represents random future losses conditional on a risk scenario  $X_{t_0}$  and  $L_\eta$  is a constant representing the value-at-risk (VaR). Often it is of interest to fix  $\eta$  and solve (1) for the corresponding VaR,  $L_\eta$ . Such problems arise frequently in risk analysis involving x-Valuation Adjustments (xVA) [22]. As motivation for this work, we consider an application computing the probability of large loss due to fluctuations in the Credit Valuation Adjustment (CVA) [2, 22]. Within this context,  $X_t$  denotes relevant market and credit factors, whereas  $g(X_T)$  represents future payoffs given the market state at time  $T$  and  $f(\mathbb{E}[g(X_T) | X_{t_1}], t_1)$  represents losses above a threshold  $L_\eta$  due to fluctuations in market and credit risk factors up to the short risk horizon  $t_0$ .

In practical settings, all three expectations in (1) must be approximated by, for example, a Monte Carlo average. Moreover, the underlying SDE of the process  $X_t$  will not typically admit an analytical solution and we must resort to Euler-Maruyama or Milstein approximations instead.

For traditional Monte Carlo approximation of (1), we use  $M^o$  Monte Carlo samples to compute the outermost expectation,  $M^m$  samples for the middle expectation and  $M^i$  samples for the innermost expectation. Additionally,  $\{X_t\}_{0 \leq t \leq T}$  is approximated using Euler-Maruyama or Milstein discretisation with step-size  $h$  given by

$$h = \begin{cases} h^o & t \in (0, t_0) \\ h^m & t \in (t_0, t_1) \\ h^i & t \in (t_1, T) \end{cases}.$$

By taking  $h^o \propto (M^m)^{-1}$  and  $h^m \propto (M^i)^{-1}$ , it follows from the weak convergence properties of the Euler-Maruyama and Milstein schemes [24] that the bias induced by approximating the underlying SDE is of the same order as that induced by using  $M^m$  and  $M^i$  samples to approximate the expectations determined by  $X_{t_0}$  and  $X_{t_1}$ , respectively.

Due to the nested structure of the problem, it follows that the total cost of the resulting estimator is of order  $M^o M^m M^i (h^i)^{-1}$ . To achieve a root mean square error of TOL, it follows from [21, Proposition 1] that we must take  $M^o = \mathcal{O}(\text{TOL}^{-2})$ ,  $M^m, M^i = \mathcal{O}(\text{TOL}^{-1})$  and  $h^i = \mathcal{O}(\text{TOL})$ . Hence, we obtain a total computational cost of order  $\text{TOL}^{-5}$ . The probability of large loss due to fluctuations in the CVA is typically small,  $\eta \ll 1$  in (1). Therefore, a small value of TOL is required to achieve reasonable results. The computational cost of order  $\text{TOL}^{-5}$  is therefore extremely prohibitive.

To reduce this cost, in the current work we consider an alternative estimator utilising the multilevel Monte Carlo setup [12]. A hierarchy of unbiased multilevel Monte Carlo corrections [28] are used to replace  $f(\mathbb{E}[g(X_T) | X_{t_1}], t_1)$  with an alternative random variable with mean  $\mathbb{E}[f(\mathbb{E}[g(X_T) | X_{t_1}], t_1) | X_{t_0}]$  and an  $\mathcal{O}(1)$  sampling cost and variance. Multilevel methods are less effective at estimating discontinuous observables such as the Heaviside function in (1).

Adaptive refinement around the discontinuity [14, 23] is used to construct an efficient multilevel estimator in this instance.

The paper is structured as follows:

- Section 2 describes the construction of a novel and efficient estimator for (1) using unbiased and adaptive multilevel Monte Carlo techniques.
- Section 2.3 discusses how the methods used can be incorporated in value-at-risk computations using stochastic root finding algorithms.
- The task of computing the probability of large loss due to fluctuations in CVA using the method of Section 2 is discussed in Section 3. The theoretical properties of the estimator are supported by numerical studies in Section 4.
- Appendix A contains an analysis of the unbiased multilevel Monte Carlo estimator proposed in Section 2.1.2.

## 2 Hierarchical Multilevel Monte Carlo Estimation

In this section, we use a hierarchy of multilevel Monte Carlo (MLMC) estimators to significantly improve upon the complexity of standard Monte Carlo approximation of (1). The discussion focuses on combining two methods:

1. Unbiased MLMC which allows us to express

$$\mathbb{E}[f(\mathbb{E}[g(X_T) | X_{t_1}], t_1) | X_{t_0}] = \mathbb{E}[Y | X_{t_0}], \quad (3)$$

where  $Y$  can be sampled with  $\mathcal{O}(1)$  cost and variance.

2. Adaptive MLMC to approximate

$$\mathbb{E}[\mathbb{H}(\mathbb{E}[Y | X_{t_0}])],$$

with Monte Carlo approximation of the inner expectation  $\mathbb{E}[Y | X_{t_0}]$  using exact samples of  $Y$ , and refinement around the discontinuity in  $\mathbb{H}(\cdot)$ .

### 2.1 Unbiased MLMC

We first recall the basic idea of unbiased MLMC [28]. Consider the general problem of approximating  $\mathbb{E}[Z]$ , where  $Z$  is a random variable which cannot be sampled directly. Assume we can sample from a hierarchy of increasingly accurate approximations  $Z_\ell \approx Z$  for all integers  $\ell \geq 0$ , with  $\lim_{\ell \rightarrow \infty} Z_\ell = Z$ , admitting a sequence of correction terms  $\Delta_\ell Z$  with the property

$$\mathbb{E}[\Delta_\ell Z] = \begin{cases} \mathbb{E}[Z_0] & \ell = 0 \\ \mathbb{E}[Z_\ell - Z_{\ell-1}] & \ell > 0 \end{cases}.$$

The following condition is required of the correction terms  $\Delta_\ell Z$ .

**Assumption 2.1.** *There exists constants  $\beta > \gamma > 0$  and  $a_0, a_1 > 0$ , independent of  $\ell$ , such that*

- $\text{cost}(\Delta_\ell Z) \leq a_0 2^{\gamma\ell}$ ,
- $\mathbb{E}[\Delta_\ell Z^2] \leq a_1 2^{-\beta\ell}$ .

Let  $\tilde{\ell}$  denote a random, non-negative integer with mass function

$$p_\ell := \mathbb{P}[\tilde{\ell} = \ell] \propto 2^{-\zeta\ell},$$

for  $\gamma < \zeta < \beta$ . Then, it follows from Assumption 2.1 that [28]

$$\mathbb{E}[Z] = \sum_{\ell=0}^{\infty} \mathbb{E}[\Delta_\ell Z] = \mathbb{E}\left[\Delta_{\tilde{\ell}} Z p_{\tilde{\ell}}^{-1}\right], \quad (4)$$

and  $\Delta_{\tilde{\ell}} Z p_{\tilde{\ell}}^{-1}$  has  $\mathcal{O}(1)$  sampling cost and variance. It is shown in [28] that to optimally balance the cost and variance of the unbiased Monte Carlo, one should take  $\zeta = (\beta + \gamma)/2$ . Alternatively, it may be of interest to ensure  $\Delta_{\tilde{\ell}} Z p_{\tilde{\ell}}^{-1}$  has finite  $q$ 'th moment for some  $q > 2$ . Under general assumptions, this can be shown to hold if  $\zeta < q\beta/2(q-1)$  [15, Remark 3.1].

We now consider the application of this method to (3).

### 2.1.1 Unbiased Estimation of $\mathbb{E}[g(X_T) | X_{t_1}]$

Given (an approximation of)  $X_{t_1}$ , we cannot exactly sample  $X_T$  and must instead discretize the underlying SDE on the interval  $[t_1, T]$ . In the context described above, let  $X_{T,\ell}$  denote the Milstein approximation to  $X_T$  given  $X_{t_1}$  and using time-step  $h_\ell^i \propto 2^{-\gamma\ell}$  on  $[t_1, T]$ , for  $\gamma$  as in Assumption 2.1. Define the correction terms  $\Delta_\ell g := g(X_{T,\ell}) - g(X_{T,\ell-1})$  for  $\ell > 0$  and  $\Delta_0 g := (X_{T,0})$ . It follows under regularity conditions on the coefficients of the underlying SDE [24, Chapter 10] and for Lipschitz  $g$  that Assumption 2.1 holds for  $\Delta_\ell g$  with  $\beta = 2\gamma$ . Thus,

$$\mathbb{E}[g(X_T) | X_{t_1}] = \mathbb{E}\left[\Delta_{\tilde{\ell}} g p_{\tilde{\ell}}^{-1} \mid X_{t_1}\right],$$

where  $\tilde{\ell}$  has probability mass function  $p_\ell \propto 2^{-\zeta\ell}$  for  $\gamma < \zeta < 2\gamma$  and  $\Delta_{\tilde{\ell}} g p_{\tilde{\ell}}^{-1}$  has finite sampling cost and variance. For later analysis (see Appendix A), we require the unbiased Monte Carlo term to have finite 3'rd moment, which instead requires  $\zeta < 3\gamma/2$ .

For high dimensional SDEs the Milstein scheme often requires the numerical simulation of Lévy areas, which can be computationally prohibitive. In this case, the antithetic Milstein scheme proposed in [18] may be useful to retain the required property on  $\beta$  while avoiding the simulation of Lévy areas.

### 2.1.2 Unbiased Estimation of (3)

Using the discussion above, we can express the left hand side of (3) as

$$\mathbb{E} \left[ f \left( \mathbb{E} \left[ \Delta_{\tilde{\ell}} g p_{\tilde{\ell}}^{-1} \mid X_{t_1} \right], t_1 \right) \mid X_{t_0} \right].$$

There are two remaining aspects of approximation which add bias to an estimator of the above expectation:

- Monte Carlo approximation of  $\mathbb{E}[\Delta_{\tilde{\ell}} g p_{\tilde{\ell}}^{-1} \mid X_{t_1}]$ .
- Approximation of  $X_{t_1}$  given  $X_{t_0}$ .

For the following discussion, let  $\Delta_{\ell} g = \Delta_{\ell} g(X_{t_1}, W_{t_1, T})$  be determined by  $X_{t_1}$  along with the Brownian path  $W_{t_1, T} = \{W_t\}_{t_1 \leq t \leq T}$  for an underlying Brownian Motion  $W_t$ . We can approximate  $\mathbb{E}[\Delta_{\tilde{\ell}} g p_{\tilde{\ell}}^{-1} \mid X_{t_1}]$  by

$$E_{\tilde{\ell}}^i(X_{t_1}) := \frac{1}{M_{\tilde{\ell}}^i} \sum_{m=1}^{M_{\tilde{\ell}}^i} \Delta_{\tilde{\ell}^{(m)}} g(X_{t_1}, W_{t_1, T}^{(m)}) p_{\tilde{\ell}^{(m)}}^{-1},$$

where  $\tilde{\ell}^{(m)}$  are i.i.d. realisations of  $\tilde{\ell}$ ,  $W_{t_1, T}^{(m)}$  are independent Brownian paths and  $M_{\tilde{\ell}}^i \propto 2^{\gamma \ell}$  for  $\gamma$  as in Assumption 2.1. The path of  $X_{t_1}$ , given  $X_{t_0}$  and a discretisation level  $\ell$ , can be computed using Milstein discretisation with time-step  $h_{\tilde{\ell}}^m \propto 2^{-\gamma \ell}$  on  $[t_0, t_1]$ .

A natural choice for the correction terms is then

$$\Delta_{\ell} f = f(E_{\tilde{\ell}}^i(X_{t_1, \ell}), t_1) - f(E_{\tilde{\ell}-1}^i(X_{t_1, \ell-1}), t_1),$$

where  $X_{t_1}$  is approximated using Milstein approximation at level  $\ell$  for the fine term and a correlated approximation at level  $\ell - 1$  for the coarse term. Independent samples of  $\tilde{\ell}$  and the Brownian path on  $W_{t_1, T}$  are used for the Monte Carlo estimators at  $E_{\tilde{\ell}}^i$  and  $E_{\tilde{\ell}-1}^i$ . However, the resulting error between the fine and coarse Monte Carlo estimators in  $\Delta_{\ell} f$  leads to  $\beta = \gamma$  in the context of Assumption 2.1, thus failing the conditions for unbiased MLMC. To improve upon this result, let  $\gamma = 1$ , giving  $M_{\tilde{\ell}}^i \propto 2^{\ell}$ . Then, for  $j = 0, 1$  define

$$\begin{aligned} E_{\tilde{\ell}-1, c, j}^i(X_{t_1}) &:= \frac{1}{M_{\tilde{\ell}-1}^i} \sum_{m=1}^{M_{\tilde{\ell}-1}^i} \Delta_{\tilde{\ell}^{(m+jM_{\tilde{\ell}-1}^i)}} g \left( X_{t_1}, W_{t_1, T}^{(m+jM_{\tilde{\ell}-1}^i)} \right) p_{\tilde{\ell}^{(m+jM_{\tilde{\ell}-1}^i)}}^{-1} \\ E_{\tilde{\ell}, f}^i(X_{t_1}) &:= \frac{1}{2} \left( E_{\tilde{\ell}-1, c, 0}^i(X_{t_1}) + E_{\tilde{\ell}-1, c, 1}^i(X_{t_1}) \right), \end{aligned}$$

so that the fine Monte Carlo average  $E_{\tilde{\ell}, f}^i(X_{t_1})$  is composed of the same Brownian paths  $W_{t_1, T}^{(m)}$  and randomised levels  $\tilde{\ell}^{(m)}$  used for two coarse averages

$E_{\ell-1,c,0}^i(X_{t_1}), E_{\ell-1,c,1}^i(X_{t_1})$ . Now, consider the antithetic estimator

$$\begin{aligned} \Delta_\ell^{\text{ant}} f &= f(E_{\ell,f}^i(X_{t_1,\ell}), t_1) \\ &\quad - \frac{1}{2} \left( f(E_{\ell-1,c,0}^i(X_{t_1,\ell-1}), t_1) + f(E_{\ell-1,c,1}^i(X_{t_1,\ell-1}), t_1) \right). \end{aligned} \quad (5)$$

For Lipschitz and piecewise twice differentiable functions  $f$ , it can be shown using the strong rate of convergence of the Milstein scheme combined with [4, Theorem 2.3] and [14, Theorem 4.1] that  $\Delta_\ell^{\text{ant}} f$  satisfies Assumption 2.1 with  $\beta = 3\gamma/2 = 3/2$ . A more detailed discussion of this result is given in Appendix A. Thus, we have

$$\mathbb{E} \left[ f \left( \mathbb{E} \left[ \Delta_{\hat{\ell}} g p_{\hat{\ell}}^{-1} \mid X_{t_1} \right], t_1 \right) \mid X_{t_0} \right] = \mathbb{E} \left[ \Delta_{\hat{\ell}}^{\text{ant}} f q_{\hat{\ell}}^{-1} \mid X_{t_0} \right],$$

where  $\hat{\ell}$  is a random, non-negative integer with probability mass function (using  $\zeta = (\beta + \gamma)/2$ )  $q_\ell \propto 2^{-5\ell/4}$  and  $\Delta_{\hat{\ell}}^{\text{ant}} f q_{\hat{\ell}}^{-1}$  has  $\mathcal{O}(1)$  sampling cost and variance. The process of generating a single sample of  $\Delta_{\hat{\ell}}^{\text{ant}} f$ , given  $X_{t_0}$ , is detailed in Algorithm 1.

## 2.2 Adaptive MLMC

The problem of computing (1) has now been reduced to estimating

$$\mathbb{E}[\mathbb{H}(\mathbb{E}[Y \mid X_{t_0}])], \quad (6)$$

where independent samples of  $Y$  are readily available. Given  $X_{t_0}$ , we thus approximate  $\mathbb{E}[Y \mid X_{t_0}]$  by

$$E_\ell^m(X_{t_0}) := \frac{1}{M_\ell^m} \sum_{m=1}^{M_\ell^m} Y^{(m)}(X_{t_0}),$$

where  $Y^{(m)}$  are i.i.d. samples of  $Y$  given  $X_{t_0}$ . In many financial applications,  $t_0 \ll t_1$  represents a short risk horizon. It is therefore possible to approximate  $X_{t_0}$  to a much higher degree of accuracy than  $X_{t_1}$  with negligible cost. Therefore, to simplify the following discussion, we assume  $X_{t_0}$  (or at least a suitably accurate approximation) is available with  $\mathcal{O}(1)$  cost. As before, we define the multilevel correction terms

$$\Delta_\ell \mathbb{H} = \Delta_\ell \mathbb{H}(X_{t_0}) := \mathbb{H}(E_\ell^m(X_{t_0})) - \mathbb{H}(E_{\ell-1}^m(X_{t_0})).$$

However, due to the possibility of making an  $\mathcal{O}(1)$  error in the Heaviside function when  $E_\ell^m(X_{t_0})$  and  $\mathbb{E}[Y \mid X_{t_0}]$  fall on opposite sides of zero, which occurs for  $\mathcal{O}\left((M_\ell^m)^{-1/2}\right)$  samples of  $E_\ell^m(X_{t_0})$ , it follows that Assumption 2.1 is satisfied only allowing for  $\beta = \gamma/2$  [13, 16]. Moreover, this property is not improved

---

**Algorithm 1:** Sampling  $\Delta_\ell^{\text{ant}} f$ 

---

**Input:**  $\ell, X_{t_0}, \gamma, M_\ell^i, h_\ell^m, \{h_\ell^i\}_{i \geq 0}$   
**Result:** Sample of  $\Delta_\ell^{\text{ant}} f$ , given  $X_{t_0}$   
**Function** Milstein( $T_0, T_1, h, x_0, \omega$ ):  
    **return** Milstein approximation of  $X_{T_1}$  with step-size  $h$  given  
        Brownian path  $\omega$  on the interval  $[T_0, T_1]$  with  $X_{T_0} = x_0$   
Sample Brownian path  $\omega_{t_1} = \{W_{t_0+nh_\ell} \mid n = 0, \dots, (t_1 - t_0)/h_\ell^m\}$ ;  
Set  $X_{t_1, \ell} = \text{Milstein}(t_0, t_1, h_\ell^m, X_{t_0}, \omega_{t_1})$ ;  
**if**  $\ell > 0$  **then**  
    **Set**  $X_{t_1, \ell-1} = \text{Milstein}(t_0, t_1, h_{\ell-1}^{(1)}, X_{t_0}, \omega_{t_1})$ ;  
    /\* Define arrays  $\Delta_j$  to store samples of  $\Delta_\ell g p_\ell^{-1}$  using  
         $X_{t_1} \approx X_{t_1, j}$  for  $j = \ell, \ell - 1$ . \*/  
    Set  $\Delta_\ell = \text{array}[M_\ell^i]$ ;  
    Set  $\Delta_{\ell-1} = \text{array}[M_\ell^i]$ ;  
    **for**  $m = 1, \dots, M_\ell^i$  **do**  
        Sample random integer  $k$  with probability mass function  
             $p_\ell \propto 2^{-3\ell/2}$ ;  
        Sample Brownian path  $\omega_T = \{W_{t_1+nh_k^i} \mid n = 0, \dots, (T - t_1)/h_k^i\}$ ;  
        Set  $X_{T, \ell, k} = \text{Milstein}(t_1, T, h_k^i, X_{t_1, \ell}, \omega_T)$ ;  
        **if**  $k > 0$  **then**  
            **Set**  $X_{T, \ell, k-1} = \text{Milstein}(t_1, T, h_{k-1}^i, X_{t_1, \ell}, \omega_T)$ ;  
        /\* Convention: When  $k = 0$  set  $g(X_{T, \cdot, -1}) \equiv 0$  \*/  
        Set  $\Delta_\ell^{(m)} = (g(X_{T, \ell, k}) - g(X_{T, \ell, k-1})) p_k^{-1}$ ;  
        **if**  $\ell > 0$  **then**  
            **Set**  $X_{T, \ell-1, k} = \text{Milstein}(t_1, T, h_k^i, X_{t_1, \ell-1}, \omega_T)$ ;  
            **if**  $k > 0$  **then**  
                **Set**  $X_{T, \ell-1, k-1} = \text{Milstein}(t_1, T, h_{k-1}^i, X_{t_1, \ell-1}, \omega_T)$ ;  
            **Set**  $\Delta_{\ell-1}^{(m)} = (g(X_{T, \ell-1, k}) - g(X_{T, \ell-1, k-1})) p_k^{-1}$ ;  
    **return**  
     $f(\text{Mean}(\Delta_\ell)) - \frac{1}{2} \sum_{i=0}^1 f(\text{Mean}(\Delta_{\ell-1}[iM_{\ell-1}^i + 1 : (i+1)M_{\ell-1}^i]))$

---

by taking an antithetic difference  $\Delta^{\text{ant}}$  as above [14]. Thus, we are unable to use unbiased MLMC for this case.

Instead, we truncate the telescoping sum appearing in (4) at some maximum level  $L$  and compute  $\{\mathbb{E}[\Delta_\ell \mathbb{H}(X_{t_0})]_{\ell=0}^L\}$  using a Monte Carlo average, reverting to standard MLMC [12]:

$$\begin{aligned} \mathbb{E}[\mathbb{H}(\mathbb{E}[Y | X_{t_0}])] &= \sum_{\ell=0}^{\infty} \mathbb{E}[\Delta_\ell \mathbb{H}(X_{t_0})] \approx \sum_{\ell=0}^L \mathbb{E}[\Delta_\ell \mathbb{H}(X_{t_0})] \\ &\approx \sum_{\ell=0}^L \frac{1}{M_\ell^o} \sum_{m=1}^{M_\ell^o} \Delta_\ell \mathbb{H}^{(m)}(X_{t_0}^{(m)}), \end{aligned} \quad (7)$$

with  $\Delta_\ell \mathbb{H}^{(m)}(X_{t_0}^{(m)}) \stackrel{\text{i.i.d.}}{\sim} \Delta_\ell \mathbb{H}(X_{t_0})$ . The key idea is that by using many samples at the cheaper levels of approximation and since the variance of  $\Delta_\ell \mathbb{H}$  decreases as  $\ell$  increases, we require fewer samples at the expensive level of simulation  $L$ . Defining  $V_\ell = \mathbb{E}[\Delta_\ell \mathbb{H}^2] \propto 2^{-\beta\ell}$ ,  $C_\ell = \text{cost}(\Delta_\ell \mathbb{H}) \propto 2^{\gamma\ell}$  by Assumption 2.1, it can be shown that by optimizing the number of samples  $M_\ell$  per level, the work required for the estimator (7) to achieve root mean square error TOL is proportional to [5, 13]

$$\text{TOL}^{-2} \left( \sum_{\ell=0}^L \sqrt{V_\ell C_\ell} \right)^2,$$

with  $L \propto \log(\text{TOL}^{-1})$ . In the best case,  $\beta > \gamma$  and the above term is  $\mathcal{O}(\text{TOL}^{-2})$ , which is frequently referred to as the canonical complexity. Instead, for  $\beta = \gamma/2$ , the estimator (7) can be shown to have  $\mathcal{O}(\text{TOL}^{-5/2})$  cost [5, 14]. For comparison, a standard Monte Carlo average used to compute (6), with  $\mathbb{E}[Y | X_{t_0}] \approx E_L^m(X_{t_0})$ , has  $\mathcal{O}(\text{TOL}^{-3})$  cost [21]. Thus, MLMC is able to improve the cost by a factor of order  $\text{TOL}^{-1/2}$  over standard Monte Carlo. However, the cost remains higher than the canonical  $\mathcal{O}(\text{TOL}^{-2})$  complexity.

Several methods have been proposed which remedy the effect of the Heaviside function on the performance of MLMC for this problem. In [17], the authors approximate  $\mathbb{H}(\cdot)$  by a Lipschitz function with better convergence properties as seen above for unbiased MLMC. However, this approximation adds an additional level of bias which is inversely proportional to the Lipschitz constant of the approximation to  $\mathbb{H}(\cdot)$ , which in turn affects the cost of MLMC. In [1] an implicit smoothing technique is applied which relies on a change of variables and a pre-integration step which smooths over the discontinuity. An alternative approach, and the one adopted for the remainder of this paper, relies on adaptively refining the number of Monte Carlo samples  $Y$  required for a given  $X_{t_0}$  [3, 9, 10, 14, 15, 23]. In the adaptive framework, we define the correction term

$$\Delta_\ell^{\text{ad}} \mathbb{H}(X_{t_0}) = \mathbb{H}(E_{\ell+\eta_\ell}^m(X_{t_0})) - \mathbb{H}(E_{\ell-1+\eta_{\ell-1}}^m(X_{t_0})),$$



with adapted Monte Carlo averages  $E_{\ell+\eta_\ell}^m(X_{t_0})$  obtained using Algorithm 2. Here,  $0 \leq \eta_\ell \leq \ell$  is a random integer, depending on  $X_{t_0}$ , which reflects uncertainty in the sign of  $E_{\ell+\eta_\ell}^m(X_{t_0})$ . Since  $\mathbb{H}(E_{\ell+\eta_\ell}^m(X_{t_0}))$  is incorrect only when  $E_{\ell+\eta_\ell}^m(X_{t_0})$  is estimated to be on the opposite side of 0 to  $\mathbb{E}[Y | X_{t_0}]$ , it is sensible to use a more accurate estimator, with larger  $\eta_\ell$ , when  $\mathbb{E}[Y | X_{t_0}]$  is thought to be close to the origin. The expected cost of sampling  $\Delta_\ell^{\text{ad}}\mathbb{H}$ ,  $\mathbb{E}[C_{\ell+\eta_\ell}]$ , is shown to remain of order  $M_\ell^m$ , whereas the variance reduces to  $V_\ell \propto (M_\ell^m)^{-1}$  under general assumptions, giving  $\beta = \gamma$  and reducing the cost of MLMC to  $\mathcal{O}(\text{TOL}^{-2}(\log \text{TOL})^2)$ . The lack of regularity in  $\mathbb{H}(\cdot)$  leads to a high kurtosis for the multilevel correction terms with and without adaptive sampling, reducing the reliability of error estimates of the MLMC estimator. One approach to this problem involves biased estimates of the first and second moments of  $\Delta_\ell\mathbb{H}$  and is discussed in Section 4.2.

---

**Algorithm 2:** Adaptive sampling at level  $\ell$  [23, Algorithm 1]

---

**Input:**  $X_{t_0}, \ell, r > 1, c > 0$   
**Result:** Adaptively refined sample  $\tilde{E}_{\ell+\eta_\ell}$   
Set  $\eta_\ell = 0$ ;  
Sample  $M_\ell^m$  independent samples of  $Y$  given  $X_{t_0}$ ;  
/\*  $\sigma_\ell(X_{t_0})$  is the sample variance given the  $M_\ell^m$  samples \*/  
Compute  $(E_\ell^m(X_{t_0}), \sigma_\ell(X_{t_0}))$  from the existing samples;  
**while**  $\sigma_{\ell+\eta_\ell}(X_{t_0})^{-1}|E_{\ell+\eta_\ell}^m(X_{t_0})| < c2^{(\ell(1-r)-\eta_\ell)/r}$  **and**  $\eta_\ell < \ell$  **do**  
    Sample  $M_{\ell+\eta_\ell+1}^m - M_{\ell+\eta_\ell}^m$  new, independent samples from  $Y$  given  
     $X_{t_0}$ ;  
    Refine  $(E_{\ell+\eta_\ell}^m(X_{t_0}), \sigma_{\ell+\eta_\ell}(X_{t_0}))$  to  $(E_{\ell+\eta_\ell+1}^m(X_{t_0}), \sigma_{\ell+\eta_\ell+1}(X_{t_0}))$   
    using all existing samples;  
    Set  $\eta_\ell = \eta_\ell + 1$ ;  
**Output:**  $E_{\ell+\eta_\ell}^m(X_{t_0})$

---

### 2.3 Quantile Estimation and Value-at-Risk

As mentioned in the introduction, it is often of interest to solve (1), (2) for  $L_\eta$  with a pre-defined value of  $\eta$ . This equates to finding the unique root of

$$F(\lambda) = \mathbb{E} \left[ \mathbb{H} \left( \mathbb{E} \left[ \Lambda(\mathbb{E}[g(X_T) | X_{t_1}], t_1) \mid X_{t_0} \right] - \lambda \right) \right] - \eta.$$

This problem has been studied using stochastic approximation methods in [25, 26, 29] and recently applied to the MLMC framework in [7, 8, 11]. An alternative algorithm is proposed in [14, Algorithm 2], although a complete analysis of this method was left to future work. The estimator of  $F(\lambda)$  constructed in previous sections can be included in such algorithms to efficiently approximate  $L_\eta$ .

### 3 CVA Capital Charge

This section describes the model CVA problem [2, 22, 27] used for the numerical experiments in Section 4. For ease of notation, the main discussion is based around a single counterparty. However, the method is intended for use in the more practical case of many counterparties. In Section 3.4, we discuss the extension from a single counterparty to an arbitrary number of counterparties.

#### 3.1 Market Setup

Consider a market based on a stock process  $S_t \in \mathbb{R}^d$  given by

$$dS_t = a(t, S_t)dt + b(t, S_t)dW_t; \quad S_0 \geq 0,$$

up to the contract maturity  $T > 0$ , where  $W_t$  is a Wiener process under the physical measure  $\mathbb{P}$  and  $a, b$  represent the drift and volatility of the market, respectively. We denote the (possibly stochastic) interest rate by  $r_t$  so that the value of an initial investment  $B_0$  in a risk-free bond at time  $t$  is  $B_t = B_0 e^{\int_0^t r_t dt}$ . We consider a contract between a bank and a counterparty with discounted future cash-flow to the bank given by  $\pi_t(S_T) = B_t B_T^{-1} \Pi(S_T)$ , where  $\Pi(S_T)$  denotes the payoff at maturity  $T$  given stock value  $S_T$ . We assume the existence of a risk-neutral measure  $\mathbb{Q}$  under which the discounted stock process  $B_t^{-1} S_t$  is a  $\mathbb{Q}$ -martingale. The arbitrage-free value of the contract at time  $t$  is then given by

$$V_t(S_t) := \mathbb{E}^{\mathbb{Q}}[\pi_t(S_T) | S_t]. \quad (8)$$

#### 3.2 Credit Valuation Adjustment

Suppose that the counterparty will default at a random time  $\tau > 0$ . In practice, the default time  $\tau$  is modelled according to the counterparties credit spread. In the event of default, the bank stands to lose a proportion LGD (Loss Given Default) of any positive cash flow. The impact this has on the valuation of the banks position at time  $0 \leq t \leq T$  is the CVA. After discounting to time 0, the CVA is given by [2]

$$\text{CVA}_t := \mathbb{E}^{\mathbb{Q}} \left[ \mathbb{I}_{\tau_t \leq T} \frac{1}{B_{\tau_t}} \text{LGD} (\mathbb{E}^{\mathbb{Q}}[\pi_{\tau_t}(S_T) | S_{\tau_t}, \tau_t])^+ \middle| S_t, c_t \right],$$

where  $c_t$  represents the counterparties credit spread at time  $t$  and  $\tau_t = \tau_t(c_t)$  defines the random default time based on the market conditions at time  $t$ . The factor  $B_t/B_{\tau_t}$  discounts the CVA to time  $t$ , and the positive part of the value process,  $(\mathbb{E}^{\mathbb{Q}}[\pi_{\tau_t}(S_T) | S_{\tau_t}, \tau_t])^+$ , is known as the exposure at default. For convenience, we express  $\text{CVA}_t = \mathbb{E}^{\mathbb{Q}}[U_{\tau_t}(S_{\tau_t}) | S_t, c_t]$ , where

$$U_{\tau_t}(x) := \mathbb{I}_{t \leq T} \frac{\text{LGD}}{B_{\tau_t}} (\mathbb{E}^{\mathbb{Q}}[\pi_{\tau_t}(S_T) | S_{\tau_t} = x, \tau_t])^+. \quad (9)$$

By an abuse of notation, we will consider  $U_{\tau_t}$  both as a function of  $x$  and of  $\mathbb{E}^{\mathbb{Q}}[\pi_{\tau_t}(S_T)|S_{\tau_t} = x, \tau_t]$  depending on the context.

We introduce the following terms, which are used frequently in what follows:

- $S_H^{\mathbb{P}}$  denotes the risky value of the stock price simulated under the physical measure  $\mathbb{P}$  at a short risk horizon  $H$ .
- $S_t^{t_0, s}$  denotes the stock process with  $S_{t_0}^{t_0, s} = s$ .

The CVA-VaR is defined as the threshold  $L_\eta$  above which losses due to changes in CVA between time 0 and a short risk horizon  $H \ll T$  occur with small probability  $\eta$  (typically  $\eta = 1\%$ ) under  $\mathbb{P}$ . Specifically,

$$\begin{aligned} \eta &= \mathbb{P}[\text{CVA}_H - \text{CVA}_0 > L_\eta] \\ &= \mathbb{E}^{\mathbb{P}}[\mathbb{H}(\mathbb{E}^{\mathbb{Q}}[\Lambda - L_\eta | S_H^{\mathbb{P}}, c_H])], \end{aligned} \tag{10}$$

where the (random) loss due to a shift in the CVA is

$$\Lambda := U_{\tau_H}(S_{\tau_H}^{H, S_H^{\mathbb{P}}}) - U_{\tau_0}(S_{\tau_0}),$$

where  $U_{\tau_t}(x)$  given by (9) is a Lipschitz function of  $\mathbb{E}^{\mathbb{Q}}[\pi_{\tau_t}(S_T)|S_{\tau_t} = x, \tau_t]$  and hence (10) may be written in the form (1). In the context of (1),  $X_t$  represents both the stock process  $S_t$  and credit spread  $c_t$ ,  $f$  is equivalent to  $\Lambda - L_\eta$  whereas  $g$  represents the discounted payoff  $\pi_\tau(S_T)$  and  $t_1$  represents both  $\tau_0$  and  $\tau_H$ . In Section 3.5, we construct an equivalent estimator which requires sampling only  $t_1 = \tau_0$ . Letting  $t_1$  be a random variable and  $g(x) = \pi_\tau(x)$  depend on  $\tau$  does not change the results of Section 2, and is discussed in Remark A.3. The risk horizon  $H$  usually represents a short interval, for example 1 week or 10 days, whereas the contract maturity  $T$  could represent several years.

### 3.3 Modelling the Default Time

As in [27], we assume a flat credit term structure so that  $c_t$  is constant under  $\mathbb{Q}$ . Following the model for default probabilities in [22, Chapter 12], it follows that

$$\tau_t = t + \hat{\tau}_t,$$

where  $\hat{\tau}_t$  are exponentially distributed random variables with rate  $c_t/\text{LGD}$ . The choice of a flat credit term structure simplifies the analysis in the following sections, however we emphasize that the multilevel Monte Carlo framework used is general and would also work with a variable credit structure.

In practice, the initial credit spread can be inferred from market data. The risky dynamics of  $c_t$  up to the risk horizon under  $\mathbb{P}$  are assumed to follow [27]

$$dc_t = \sigma^{\text{cs}} c_t dW_t^{\text{cs}},$$

where  $\sigma^{\text{cs}} > 0$  is a fixed volatility constant, and  $W_t^{\text{cs}}$  is a Wiener process under  $\mathbb{P}$ , independent of  $W_t$  which drives the underlying stock process in Section 3.1. The product  $\sigma^{\text{cs}}c_0$  is linked with the external credit rating of the counterparty in [27, Table A].

### 3.4 General Formulation for Multiple Counterparties

Typically, the CVA capital charge considers large portfolios containing many options with  $K \geq 1$  counterparties, opposed to the one-dimensional problem considered above. In this case, the adjustment made to the valuation of the banks portfolio is

$$\text{CVA}_t := \sum_{k=1}^K \text{CVA}_t^{(k)},$$

where  $\text{CVA}^{(k)}$  denotes the CVA owing to counterparty  $k$ . In this approach, the credit spread and loss given default will differ for each counterparty. The VaR becomes the value  $L_\eta$  solving

$$\eta = \mathbb{E}^{\mathbb{P}} \left[ \mathbb{H} \left( \mathbb{E}^{\mathbb{Q}} \left[ \sum_{k=1}^K \Lambda^{(k)} - L_\eta \mid S_H^{\mathbb{P}}, \{c_H^{(i)}\}_{i=1}^K \right] \right) \right], \quad (11)$$

where  $\Lambda^{(k)}$  represents the loss due to CVA for the counterparty  $k$ . While we focus on the case  $K = 1$  in this paper, the discussion applies in the extended case where  $K > 1$ . Some additional points to consider when  $K \gg 1$  are discussed in Section 4.4.

### 3.5 Variance Reduction

We now turn our attention to the approximation of (10) using the MLMC methods in Section 2. As mentioned in the introduction this approach reduces the  $\mathcal{O}(\text{TOL}^{-5})$  cost for standard Monte Carlo approximation to achieve root mean square error TOL to  $\mathcal{O}(\text{TOL}^{-2}(\log \text{TOL})^2)$ . The performance of (multilevel) Monte Carlo approximation depends strongly on the variance of  $\Lambda$  as defined in Section 3. As such we first consider variance reduction techniques, which are able to significantly improve the performance of MLMC.

There are two key factors which decrease the correlation between  $U_{\tau_H} \left( S_{\tau_H}^{H, S_H^{\mathbb{P}}} \right)$  and  $U_{\tau_0} \left( S_{\tau_0} \right)$  significantly, giving  $\Lambda$  a large variance:

- $\tau_H, \tau_0$  have different rates and thus are difficult to correlate. It is unlikely that both of these terms will default before  $T$ .
- Due to simulation of the market risk factor under  $\mathbb{P}$ , the stocks  $S_t^{H, S_H^{\mathbb{P}}}$  and  $S_t$  follow different stochastic processes up to the risk horizon  $H$ .

Note that for the exponentially distributed default times in Section 3.3 and some function  $F$  we have

$$\begin{aligned}\mathbb{E}^{\mathbb{Q}}[F(\tau_H)|c_H] &= \int_H^\infty F(t) \frac{c_H}{\text{LGD}} e^{-c_H(t-H)/\text{LGD}} dt \\ &= \frac{c_H}{c_0} e^{c_H H/\text{LGD}} \int_0^\infty \mathbb{I}_{t \geq H} F(t) \frac{c_0}{\text{LGD}} e^{-(c_H - c_0)t/\text{LGD}} e^{-c_0 t/\text{LGD}} dt \\ &= \mathbb{E}^{\mathbb{Q}} \left[ \mathbb{I}_{\tau_0 \geq H} F(\tau_0) g_{\tau_0}(c_H) e^{c_H H/\text{LGD}} | c_H \right],\end{aligned}$$

where  $g_t(x) := c_0^{-1} x e^{-(x-c_0)t/\text{LGD}}$ . For ease of notation, we will express  $\tau_0 \equiv \tau$  for the remainder of this paper. The above computation then motivates replacing  $\Lambda$  with

$$\begin{aligned}\Lambda^{\text{I}} = \mathbb{I}_{\tau \geq H} & \left( g_\tau(c_H) e^{c_H H/\text{LGD}} \underbrace{\left( U_\tau(S_\tau^{H, S_H^{\mathbb{P}}}) - U_\tau(S_\tau) \right)}_{\text{Market risk}} \right. \\ & \left. + \underbrace{\left( g_\tau(c_H) e^{c_H H/\text{LGD}} - 1 \right)}_{\text{Credit risk}} U_\tau(S_\tau) \right),\end{aligned}\tag{12}$$

whenever  $\tau \geq H$ . The component of  $\Lambda$  reflecting default before  $H$  is  $-\mathbb{I}_{\tau < H} U_\tau(S_\tau)$ . Since this is independent of the risk factors at the horizon  $H$ , we can cancel the bias induced by considering only  $\tau \geq H$  in  $\Lambda^{\text{I}}$  by setting

$$L_\eta^{\text{I}} := L_\eta + \mathbb{E}^{\mathbb{Q}}[\mathbb{I}_{\tau < H} U_\tau(S_\tau)],$$

where  $\mathbb{E}^{\mathbb{Q}}[\mathbb{I}_{\tau < H} U_\tau(S_\tau)]$  can be pre-computed offline at insignificant cost compared with the cost of computing  $\eta$ . By the change of measure from  $\tau_H$  to  $\tau$  above, it follows that  $\mathbb{E}^{\mathbb{Q}}[\Lambda - L_\eta | S_H^{\mathbb{P}}, c_H] = \mathbb{E}^{\mathbb{Q}}[\Lambda^{\text{I}} - L_\eta^{\text{I}} | S_H^{\mathbb{P}}, c_H]$ . Note that we have decomposed  $\Lambda^{\text{I}}$  into two terms with variance driven primarily by the market or credit risk factors as discussed above. We consider each term in turn.

### 3.5.1 Credit Risk

In [27], it is shown that the standardised CVA capital charge can be derived by linearising

$$g_\tau(c_H) e^{c_H H/\text{LGD}} - 1 = \underbrace{(c_H - c_0) \left( \frac{1}{c_0} - \frac{\tau}{\text{LGD}} \right)}_{\mathcal{O}(\sqrt{H})} + \mathcal{O}(H).\tag{13}$$

To remove order  $\sqrt{H}$  terms, we should take

$$\Lambda^{\text{II}} := \Lambda^{\text{I}} - \mathbb{I}_{\tau \geq H} (c_H - c_0) \chi_\tau(S_\tau),$$

where we introduce the control variate

$$\chi_\tau(S_\tau) := \left( \frac{1}{c_0} - \frac{\tau}{\text{LGD}} \right) U_\tau(S_\tau).$$

To ensure this does not bias the expected value of  $\Lambda - L_\eta$ , we set

$$L_\eta^{\text{II}} := L_\eta^{\text{I}} - (c_H - c_0) \mathbb{E}^{\mathbb{Q}}[\chi_\tau(S_\tau)],$$

so that again  $\mathbb{E}^{\mathbb{Q}}[\Lambda^{\text{II}} - L_\eta^{\text{II}} | S_H^{\mathbb{P}}, c_H] = \mathbb{E}^{\mathbb{Q}}[\Lambda - L_\eta | S_H^{\mathbb{P}}, c_H]$ .

### 3.5.2 Market Risk

The use of antithetic sampling and delta control variates to reduce the variance owed to market risk in VaR calculations is well documented [15, 20]. Following this approach leads to the term

$$\begin{aligned} \Lambda^{\text{III}} := & \mathbb{I}_{\tau \geq H} \left( g_\tau(c_H) e^{c_H H / \text{LGD}} U_\tau(S_\tau^{H, S_H^{\mathbb{P}}}) \right. \\ & - \frac{1}{2} (U_\tau(S_\tau^+) + U_\tau(S_\tau^-)) \\ & - \frac{1}{2} (c_H - c_0) (\chi_\tau(S_\tau^+) + \chi_\tau(S_\tau^-)) \\ & \left. - \frac{1}{2} (S_H^{H, S_H^{\mathbb{P}}} - S_0) (\nabla_{S_0} \cdot U_\tau(S_\tau^+) + \nabla_{S_0} \cdot U_\tau(S_\tau^-)) \right), \end{aligned}$$

where the antithetic pair  $S_t^\pm$  follow the Brownian paths  $\{\pm W_t\}_{0 \leq t \leq H}$  respectively. After  $H$ , all of the processes  $S_t^\pm, S_t^{H, S_H^{\mathbb{P}}}$  follow the same Brownian path. The delta control variates  $\nabla_{S_0} \cdot U_\tau(S_\tau^\pm)$  arise from removing the leading order term in a Taylor expansion of  $U_\tau(S_\tau^{H, S_H^{\mathbb{P}}}) - U_\tau(S_\tau)$ . The factor  $g_\tau(c_H) e^{c_H H / \text{LGD}}$  of the market risk in (12) can be ignored in this expansion since we see in (13) that this term is of the form  $1 + \mathcal{O}(\sqrt{H})$ . As before, to compensate for the bias induced by the delta control variate, we replace  $L_\eta^{\text{II}}$  with

$$L_\eta^{\text{III}} := L_\eta^{\text{II}} - (S_H^{H, S_H^{\mathbb{P}}} - S_0) \mathbb{E}^{\mathbb{Q}}[\nabla_{S_0} \cdot U_\tau(S_\tau)]$$

so that  $\mathbb{E}^{\mathbb{Q}}[\Lambda^{\text{III}} - L_\eta^{\text{III}} | S_H^{\mathbb{P}}, c_H] = \mathbb{E}^{\mathbb{Q}}[\Lambda - L_\eta | S_H^{\mathbb{P}}, c_H]$ . Since all of the expectations appearing in the definition of  $L_\eta^{\text{III}}$  are independent of  $S_H^{\mathbb{P}}$  and  $c_H$ , they may be pre-computed offline at insignificant cost compared to  $\mathbb{E}^{\mathbb{Q}}[\Lambda^{\text{III}}]$ .

## 3.6 Sampling Rare Default Times

Observe that the CVA loss,  $\Lambda^{\text{III}}$ , is non-zero only when  $H \leq \tau \leq T$ . For credible counterparties this is a rare event, making it costly to simulate a single instance of default. To address this issue note that, defining  $\tilde{p} := \mathbb{P}[H \leq \tau \leq T]$ , we have

$$\mathbb{E}^{\mathbb{Q}} \left[ \Lambda^{\text{III}} - L_\eta^{\text{III}} \middle| S_H^{\mathbb{P}}, c_H \right] = \mathbb{E}^{\mathbb{Q}} \left[ \Lambda^{\text{III}} \tilde{p} - L_\eta^{\text{III}} \middle| S_H^{\mathbb{P}}, c_H, H \leq \tau \leq T \right]. \quad (14)$$

For  $H \leq t \leq T$  and the model for  $\tau$  in Section 3.3, we have

$$\mathbb{P}[\tau \leq t | H \leq \tau \leq T] = \frac{\mathbb{P}[H \leq \tau \leq t]}{\mathbb{P}[H \leq \tau \leq T]} = \frac{e^{-c_0 H / \text{LGD}} - e^{-c_0 t / \text{LGD}}}{e^{-c_0 H / \text{LGD}} - e^{-c_0 T / \text{LGD}}}.$$

By inverting this equation for  $t$ , we can sample directly from the conditional distribution using inverse transform sampling.

This approach relies explicitly on the model for default times described in Section 3.3. For more complex models of the default time, more advanced importance sampling techniques are required to achieve a similar result.

### 3.7 Unbiased Estimation of the Loss

As previously remarked, it is not typically possible to sample the CVA loss  $\Lambda$  directly due to complicated market dynamics. The approach used in Section 2.1 provides an unbiased estimator of the form  $\mathbb{E}[\Lambda | S_H^{\mathbb{P}}, c_H] = \mathbb{E}[\Delta_{\hat{\ell}}^{(\text{ant})} \Lambda q_{\hat{\ell}}^{-1}]$  where the multilevel corrections  $\Delta_{\ell} \Lambda$  rely on Milstein simulation of the market and Monte Carlo valuation of the underlying contract. It is also possible to implement this approach for the estimator  $\Lambda^{\text{III}}$  above with the exception that the delta control variate  $\nabla_{S_0} \cdot U$  is discontinuous, and the multilevel correction term involving this control variate does not satisfy Assumption 2.1, precluding the application of unbiased MLMC. However, to retain the benefits of the delta control variate in variance reduction, one may include this term only for a fixed, deterministic range of levels as in [15].

## 4 Numerical Experiments

To illustrate the preceding methods, we consider the estimation of the probability of large loss,  $\eta$ , for a toy CVA problem (10) consisting of a single stock process and counterparty outlined in Section 4.1. In Section 4.4, we discuss how the methods can be readily extended to larger scale problems.

### 4.1 Simple Synthetic Portfolio

The underlying contract between the bank and counterparty is given by a linear combination of two European call options with discounted future cash-flow

$$\pi_t(S_T) = \frac{B_t}{B_T} (c_0(S_T - K_0)^+ + c_1(S_T - K_1)^+),$$

where constants  $c_0, c_1$  are determined by  $V_0$  and the delta-neutral condition  $\partial_{S_0} V_0 = 0$ . We assume the stock process follows a Geometric Brownian Motion under the Black-Scholes market assumptions with fixed interest rate  $r = 1\%$  so that  $B_t = e^{t/100}$ . Then, under  $\mathbb{P}$  and  $\mathbb{Q}$ , respectively, we have

$$\begin{aligned} dS_t &= \mu S_t dt + \sigma S_t dW_t, \\ dS_t &= r S_t dt + \sigma S_t dW_t^{\mathbb{Q}}. \end{aligned}$$

We take  $T = 1$  year, as well as  $S_0 = 1$  and  $\mu = \sigma = 0.1$ . To test the methods in greater generality, we ignore the existing analytic formulae for the value

and stock processes. Instead, we resort to Milstein simulation of  $S_t$  and Monte Carlo estimation of the exposure at default. Thus, we rely upon the methods of Section 3.7 to construct an unbiased estimator of the CVA loss.

Following the model for default times in Section 3.3 we take  $c_0 = 5\%$  and  $\sigma^{\text{cs}} = 0.8\%/c_0$ , corresponding to an A-rated counterparty according to [27, Table A]. The probability of large loss (10) is computed with risk horizon  $H = 10$  days and a fixed loss threshold  $L_\eta = 5 \times 10^{-4}$ , giving  $\eta \approx 2\%$ .

## 4.2 MLMC Setup

Simulation of the outer expectation in (10) over  $\mathbb{P}$  is performed using the methods in Section 2.2. We consider both adaptive sampling with  $M_\ell^{\text{m}} = M_0^{\text{m}}2^\ell$  middle simulation samples per level and non-adaptive sampling with  $M_\ell^{\text{m}} = M_0^{\text{m}}2^{\gamma\ell}$  for  $\gamma = 1, 2$ . At level  $\ell = 0$  we use  $M_0^{\text{m}} = 8$  samples per level. The expected value of the loss  $\Lambda$  over  $\mathbb{Q}$  is approximated using a hierarchy of unbiased MLMC methods as in Section 3.7, and using the control variates discussed therein.

### Optimal Starting Level

For an effective comparison, it is important all methods are tested with ideal hyper parameters. For nested simulation, it is often optimal to modify the MLMC estimator to start not from  $\ell = 0$  but at level  $\ell = \ell_0 > 0$  [14, 15]. The optimal starting level may be approximated by iteratively estimating the ratio  $R_{\ell_0}$  between the cost of a two level estimator starting at level  $\ell_0$  and a single level estimator at level  $\ell_0 + 1$ . If  $R_\ell > 1$ , it is more efficient to start at level  $\ell_0 + 1$ .

### Error Estimation

Robust computation of the MLMC estimator (7) requires accurate estimates of the error terms  $V_\ell := \text{Var}[\Delta_\ell \mathbb{H}]$  and  $E_\ell := |\mathbb{E}[\Delta_\ell \mathbb{H}]|$  in order to approximate the optimal maximum level  $L$  and number of samples  $\{M_\ell\}_{\ell=\ell_0}^L$  to use per level [13]. Reliable estimates for  $V_\ell$  need  $\mathcal{O}(\kappa_\ell)$  outer samples, where  $\kappa_\ell$  is the Kurtosis of  $\Delta_\ell \mathbb{H}$ . For the Heaviside function, it follows that  $\kappa_\ell \approx V_\ell^{-1}$ . This contradicts the methodology of MLMC to reduce the number of samples required at fine levels, and can severely impact the robustness of the estimator.

In [30], the authors approximate  $\mathbb{H}(\cdot)$  using a sequence of smooth logistic sigmoid functions which depend on  $\ell$  and significantly reduce  $V_\ell, \kappa_\ell$  by a constant factor. Alternatively, in [6] the authors assume  $V_\ell \approx a_0 2^{-\beta\ell}, E_\ell \approx a_1 2^{-\alpha\ell}$  and use Bayesian estimates for the parameters  $a_0, a_1, \alpha, \beta$ .

Here, we describe an alternative, heuristic method based on a similar ap-



proach in [10]. The multilevel correction terms  $\Delta_\ell \mathbb{H}$  take the form

$$\Delta_\ell \mathbb{H} = \begin{cases} 1 & \text{probability } p_\ell \\ -1 & \text{probability } q_\ell \\ 0 & \text{probability } 1 - p_\ell - q_\ell \end{cases},$$

with

$$V_\ell \leq p_\ell + q_\ell, \quad E_\ell = |p_\ell - q_\ell|.$$

When  $\ell$  is large, both  $p_\ell, q_\ell \ll 1$ . Heuristically, when the number of samples  $M_\ell^\circ$  of  $\Delta_\ell \mathbb{H}$  is small, it is extremely unlikely we will observe a non-zero sample, creating unstable approximations to  $V_\ell, E_\ell$ . The idea is to impose a Bayesian prior on  $p_\ell, q_\ell$  which incorporates knowledge of the theoretical convergence rates of  $V_\ell$  and  $E_\ell$  such that when the number of samples  $M_\ell^\circ$  is low, we rely more heavily on the theoretical rates than on the observations, and vice-versa as  $M_\ell^\circ$  becomes large.

Assume that we know a relationship of the form

$$V_\ell \approx a_0 2^{-\beta \ell}, \quad (15)$$

where the constant  $a_0 < 1$  may be reasonably estimated using information from coarse levels where samples of  $\Delta_\ell \mathbb{H}$  are inexpensive. Assume a Dirichlet prior density on  $p_\ell, q_\ell$

$$\rho_\ell(p, q) = (1 - p - q)^{c_\ell - 1} p^k q^k,$$

for constants  $c_\ell, k > 0$ . For the robustness of the MLMC estimator, it is preferable to over-estimate opposed to under-estimate  $V_\ell$ . Hence, it is reasonable to impose

$$\mathbb{E}_{\rho_\ell}[p_\ell + q_\ell] \geq V_\ell,$$

which is true provided

$$c_\ell \leq 2(k + 1)(V_\ell^{-1} - 1).$$

Hence, using (15), a reasonable conservative estimate is

$$c_\ell = 2k(a_0^{-1} 2^{\beta \ell} - 1). \quad (16)$$

Given observations  $\{\Delta_\ell \mathbb{H}^{(m)}\}_{m=1}^{M_\ell^\circ}$ , define

$$\nu_\ell^\pm := \sum_{m=1}^{M_\ell^\circ} \mathbb{I}_{\Delta_\ell \mathbb{H}^{(m)} = \pm 1}.$$

Then, the maximum-a-posteriori estimate of  $p_\ell + q_\ell$  is

$$\hat{V}_\ell = \frac{\nu_\ell^+ + \nu_\ell^- + 2k}{M_\ell^\circ + 2k + c_\ell - 1}.$$

The parameter  $k$  can be chosen to control how confident we are that  $V_\ell \leq \hat{V}_\ell$  for a small number of samples  $M_\ell^o$ . Larger values of  $k$  produce more reliable, conservative bounds.

To estimate  $E_\ell = |p_\ell - q_\ell|$  we construct biased estimates for  $p_\ell - q_\ell$  and  $q_\ell - p_\ell$  separately. First, in the case where  $p_\ell > q_\ell$ , we wish to estimate of  $p_\ell - q_\ell$ . In doing so, we impose a Dirichlet( $d_\ell, j + 1, 1$ ) prior on  $p_\ell, q_\ell$  with density

$$\tilde{\rho}_\ell(p, q) = (1 - p - q)^{d_\ell - 1} p^j.$$

Given the observations  $\{\Delta_\ell \mathbb{H}^m\}_{m=1}^{M_\ell^o}$  the posterior distribution becomes a Dirichlet( $d_\ell, \nu_\ell^+ + j + 1, \nu_\ell^- + 1$ ) distribution with maximum a posteriori estimator given by

$$\frac{\nu_\ell^+ - \nu_\ell^- + j}{M_\ell^o + j + d_\ell - 1}.$$

Under the prior distribution we have

$$\mathbb{E}_{\tilde{\rho}_\ell}[p_\ell - q_\ell] = \frac{j}{j + d_\ell + 2},$$

so enforcing the condition

$$\mathbb{E}_{\tilde{\rho}_\ell}[p_\ell - q_\ell] \geq e_0 2^{-\alpha_\ell} \approx |p_\ell - q_\ell|$$

requires

$$d_\ell \leq j(e_0^{-1} 2^{\alpha_\ell} - 1) - 2.$$

In the case where  $q_\ell > p_\ell$  one can repeat this process by interchanging the roles of  $p$  and  $q$ . Taking the maximum of the two resulting estimators gives

$$\hat{E}_\ell := \frac{|\nu_\ell^+ - \nu_\ell^-| + j}{M_\ell^o + j + d_\ell - 1}.$$

For the experiments presented here we choose  $k = j = 1$ .

### 4.3 Results

Figure 1 plots various statistics of the multilevel correction terms  $\Delta_\ell \mathbb{H}$  for each method. Figure 2 further plots the total work required to compute  $\eta$  to accuracy TOL, against TOL for each method. We discuss the results below:

#### Work $C_\ell$

The top left plot in Figure 1 shows the expected work  $C_\ell$  of generating a single sample of  $\Delta_\ell \mathbb{H}$ . By construction, the non-adaptive samplers have  $\mathcal{O}(2^{\gamma_\ell})$  cost (with some variability owed to the random cost of the unbiased MLMC estimators). On the other hand, the adaptive sampler has expected cost  $\mathbb{E}[2^{\ell + \eta_\ell}] = \mathcal{O}(2^\ell)$  as discussed in Section 2.2.

### Variance $V_\ell$

The top right plot in Figure 1 shows the variance  $V_\ell = \text{Var}[\Delta_\ell \mathbb{H}]$  as a function of  $\ell$ . Following the discussion in Section 2.2, the non-adaptive samplers have  $V_\ell = \mathcal{O}(N_\ell^{-1/2}) = \mathcal{O}(2^{-\gamma\ell/2})$ . On the other hand, the adaptive sampler has  $V_\ell = \mathcal{O}(N_\ell^{-1}) = \mathcal{O}(2^{-\ell})$ . Combined with an expected sampling cost of order  $2^\ell$ , this improves the convergence rate of the adaptive MLMC over the non-adaptive estimators.

### Kurtosis $\kappa_\ell$

The bottom left plot in Figure 1 shows the kurtosis of  $\Delta_\ell \mathbb{H}$ ,  $\kappa_\ell$ , against  $\ell$ . We observe  $\kappa_\ell = \mathcal{O}(V_\ell^{-1})$  for all methods. As mentioned above, this reduces the accuracy of standard estimates for the overall error made by MLMC, especially when  $L$  is large. This motivates the use of the methods discussed in Section 4.2 to improve the robustness of MLMC.

### Optimal Starting Point

The bottom right plot in Figure 1 shows the ratio  $R_{\ell_0}$  between a two-level and single-level estimator starting at level  $\ell_0$  as discussed in Section 4.2. To ensure MLMC performs optimally, one should start from a level  $\ell_0$  for which  $R_{\ell_0} < 1$ .

### Total Work

In Figure 2, we see a reduction in cost by a factor of around 7 times when using adaptive opposed non-adaptive MLMC when computing  $\eta$  to 2 significant figures. We also see that our results closely match the theoretical complexities  $\mathcal{O}(\text{TOL}^{-5/2})$  and  $\mathcal{O}(\text{TOL}^{-2}(\log \text{TOL})^2)$  stated for non-adaptive and adaptive sampling, respectively in Section 2.2. A reference line shows the theoretical  $\mathcal{O}(\text{TOL}^{-5})$  complexity discussed in Section 1 for traditional Monte Carlo without any of the multilevel techniques in Section 2. This illustrates savings of several orders of magnitude at the smallest computed error tolerance by using the adaptive hierarchical MLMC approach compared to using standard Monte Carlo.

## 4.4 Extension to Larger Portfolios

The experiment above is based on a toy problem consisting of a single stock and counterparty. This setup is enough to test the methods of the preceding sections in the context of risk estimation of the CVA. However, it is important to stress that these methods can be readily extended to more realistic problems, with many counterparties and a large market to simulate. There are several additional considerations than can be useful when considering such problems:




Figure 1: Statistics of the correction terms  $\Delta_\ell \mathbb{H}$  plotted against  $\ell$  for the MLMC estimators using both adaptive and non-adaptive sampling.




Figure 2: Total work required to compute the probability of large loss due to changes in the CVA multiplied by  $TOL^2$  against  $TOL$ , normalised by the true value of  $\eta$ .

- (*Randomized subsampling of counterparties.*) For a large number of counterparties ( $K \gg 1$  in (11)) it can be more efficient to write the sum

$$\sum_{k=1}^K \Lambda^{(k)} = \mathbb{E}[\Lambda^{(k^*)} q_{k^*}^{-1}],$$

where  $k^*$  is drawn from a discrete probability distribution on  $\{1, \dots, k\}$ . The expectation  $\mathbb{E}[\Lambda^{k^*} q_{k^*}^{-1}]$  may then be approximated using a Monte Carlo average over  $k^*$ . As a result, we need only consider a single counterparty with each sample of the inner expectation rather than evaluate each of  $\{\Lambda^{(k)}\}_{k=1}^K$  for each sample. This approach, known as randomised sub-sampling [15], can reduce the cost of the MLMC estimator when  $K$  is sufficiently large.

- (*Simulation of market at large number of intermediate points.*) When there are many counterparties with different default times and/or many options per counterparty with differing maturities, it may be necessary to evaluate the stock process at many intermediate time points. It is possible to modify the Milstein discretisation to ensure the underlying Brownian path is sampled at each intermediate point. However, when the number of intermediate points is large, this can add a significant cost to the computation of the unbiased MLMC. One possible alternative is to interpolate the intermediate points using an antithetic approach similar to that used in Section 3.5.2, by sampling an antithetic pair  $S_t^{f,\pm}$  at the fine level which follow  $\pm$  an underlying Brownian path between each pair of coarse time points. If the coarse path is  $S_t^c$  and the Brownian path is sampled at evenly spaced timepoints over  $[0, T]$ , numerical experiments indicate that using linear interpolation the multilevel correction term

$$\Delta_\ell^{\text{ant}} S_t := \frac{1}{2}(S_t^{f,+} + S_t^{f,-}) - S_t^c$$

satisfies the necessary condition for unbiased MLMC in Assumption 2.1 for all values  $t \in [0, T]$ . Analysis of such an approach is left to future work.

## 5 Conclusion

Due to large costs induced by the approximation of market factors combined with several nested Monte Carlo averages, accurate estimation of xVA risk measures is often considered infeasible using traditional Monte Carlo methods. By combining recent developments in MLMC for nested simulation with variance reduction techniques, this paper illustrates an efficient framework for accurate simulation of these risk measures at greatly reduced costs. In the context of the CVA-VaR problem, we are able to reduce the cost of achieving accuracy TOL from  $\mathcal{O}(\text{TOL}^{-5})$  for traditional Monte Carlo to  $\mathcal{O}(\text{TOL}^{-2}(\log \text{TOL})^2)$  using MLMC with an adaptive number of inner samples of the loss. This results

in computational savings of several orders of magnitude for realistic problems. Paired with the capabilities of high performance computing, this shows that Monte Carlo methods are a viable option for efficient and accurate simulation of xVA risk measures.

Future aspects of this work involve the use of stochastic approximation techniques to compute the Value-at-Risk as in Section 2.3, along with an application to larger portfolios as discussed in Section 4.4. An additional area of future research concerns multilevel Quasi-Monte Carlo methods for the approximation of (1). Such methods are used in [30] to construct an estimator with  $\mathcal{O}(\text{TOL}^{-2}(\log \text{TOL})^2)$  cost without adaptive sampling techniques. When combined with the adaptive methods, we can potentially construct an estimator with even better costs. For problems with similar convergence properties [19], multilevel Quasi-Monte Carlo estimators have been shown to have  $\mathcal{O}(\text{TOL}^{-1})$  cost to achieve accuracy TOL.

### Acknowledgments

J. Spence was supported by EPSRC grant EP/S023291/1.

### References

- [1] C. Bayer, C. B. Hammouda, and R. Tempone. Multilevel Monte Carlo combined with numerical smoothing for robust and efficient option pricing and density estimation. *arXiv preprint arXiv:2003.05708*, 2020.
- [2] D. Brigo, M. Morini, and A. Pallavicini. *Counterparty credit risk, collateral and funding: with pricing cases for all asset classes*. Wiley Finance Series. John Wiley and Sons, 2013.
- [3] M. Broadie, Y. Du, and C. Moallemi. Efficient risk estimation via nested sequential simulation. *Management Science*, 57(6):1172–1194, 2011.
- [4] K. Bujok, B. Hambly, and C. Reisinger. Multilevel simulation of functionals of Bernoulli random variables with application to basket credit derivatives. *Methodology and Computing in Applied Probability*, 17(3):579–604, 2015.
- [5] K. Cliffe, M. Giles, R. Scheichl, and A. Teckentrup. Multilevel Monte Carlo methods and applications to elliptic PDEs with random coefficients. *Computing and Visualization in Science*, 14(1):3–15, 2011.
- [6] N. Collier, A.-L. Haji-Ali, F. Nobile, E. von Schwerin, and R. Tempone. A continuation multilevel Monte Carlo algorithm. *BIT Numerical Mathematics*, 55(2):399–432, 2015.
- [7] S. Dereich. General multilevel adaptations for stochastic approximation algorithms II: CLTs. *Stochastic Processes and their Applications*, 132:226–260, 2021.
- [8] S. Dereich and T. Müller-Gronbach. General multilevel adaptations for stochastic approximation algorithms of Robbins–Monro and Polyak–Ruppert type. *Numerische Mathematik*, 142(2):279–328, 2019.

- [9] T. Dodwell, S. Kynaston, R. Butler, R. Haftka, N. H. Kim, and R. Scheichl. Multilevel Monte Carlo simulations of composite structures with uncertain manufacturing defects. *Probabilistic Engineering Mechanics*, 63:103–116, 2021.
- [10] D. Elfverson, F. Hellman, and A. Målqvist. A multilevel Monte Carlo method for computing failure probabilities. *SIAM/ASA Journal on Uncertainty Quantification*, 4(1):312–330, 2016.
- [11] N. Frikha. Multi-level stochastic approximation algorithms. *The Annals of Applied Probability*, 26(2):933–985, 2016.
- [12] M. Giles. Multilevel Monte Carlo path simulation. *Operations Research*, 56(3):607–617, 2008.
- [13] M. Giles. Multilevel Monte Carlo methods. *Acta Numerica*, 24:259–328, 2015.
- [14] M. Giles and A.-L. Haji-Ali. Multilevel nested simulation for efficient risk estimation. *SIAM/ASA Journal on Uncertainty Quantification*, 7(2):497–525, 2019.
- [15] M. Giles and A.-L. Haji-Ali. Sub-sampling and other considerations for efficient risk estimation in large portfolios. *Journal of Computational Finance*, 26(1):113–140, 2022.
- [16] M. Giles, D. Higham, and X. Mao. Analysing multilevel Monte Carlo for options with non-globally Lipschitz payoff. *Finance and Stochastics*, 13(3):403–413, 2009.
- [17] M. Giles, T. Nagapetyan, and K. Ritter. Multilevel Monte Carlo approximation of distribution functions and densities. *SIAM/ASA Journal on Uncertainty Quantification*, 3(1):267–295, 2015.
- [18] M. Giles and L. Szpruch. Antithetic multilevel Monte Carlo estimation for multi-dimensional SDEs without Lévy area simulation. *Annals of Applied Probability*, 24(4):1585–1620, 2014.
- [19] M. Giles and B. Waterhouse. Multilevel quasi-Monte Carlo path simulation. In *Advanced Financial Modelling*, Radon Series on Computational and Applied Mathematics, pages 165–181. De Gruyter, 2009.
- [20] P. Glasserman. *Monte Carlo methods in Financial Engineering*, volume 53 of *Stochastic Modelling and Applied Probability*. Springer New York, 2003.
- [21] M. Gordy and S. Juneja. Nested simulation in portfolio risk measurement. *Management Science*, 56(10):1833–1848, 2010.
- [22] J. Gregory. *The xVA Challenge: Counterparty Risk, Funding, Collateral, Capital and Initial Margin*. John Wiley and Sons, fourth edition, 2020.
- [23] A.-L. Haji-Ali, J. Spence, and A. Teckentrup. Adaptive multilevel Monte Carlo for probabilities. *SIAM Journal on Numerical Analysis*, 60(4):2125–2149, 2022.
- [24] P. Kloeden and E. Platen. *Numerical Solution of Stochastic Differential Equations*, volume 23 of *Stochastic Modelling and Applied Probability*. Springer Berlin, Heidelberg, third edition, 1999.
- [25] H. Kushner. Stochastic approximation: a survey. *WIREs Computational Statistics*, 2(1):87–96, 2010.
- [26] H. Kushner and G. Yin. *Stochastic Approximation and Recursive Algorithms and Applications*, volume 35 of *Stochastic Modelling and Applied Probability*. Springer New York, second edition, 2003.

- [27] M. Pykhtin. Model foundations of the Basel III standardised CVA charge. *Risk*, 25(7):60–66, 2012.
- [28] C.-H. Rhee and P. Glynn. Unbiased estimation with square root convergence for SDE models. *Operations Research*, 63(5):1026–1043, 2015.
- [29] H. Robbins and M. S. A stochastic approximation method. *The Annals of Mathematical Statistics*, 22(3):400–407, 1951.
- [30] Z. Xu, Z. He, and X. Wang. Efficient risk estimation via nested multilevel quasi-Monte Carlo simulation. *arXiv preprint arXiv:2011.11898*, 2020.

## A Antithetic Randomised Multilevel Monte Carlo Estimator

Here we provide a sketch proof that the antithetic estimator (5), satisfies Assumption 2.1 with  $\beta = 3/2$ . We provide a proof only for

$$f(x, t_1) = \sum_{j=1}^J c_j(t_1) \max(0, x - d_j(t_1)), \quad (17)$$

for continuous functions  $c_j(t_1)$  and  $d_j(t_1)$  and  $1 \leq j \leq J$ . The loss function  $\Lambda$  occurring in the CVA-VaR problem can be written of the form (17) provided one treats the delta control variate as in Section 3.7. For simplicity, we assume  $t_1 \in (t_0, T)$  is fixed and consider  $f(x, t_1) = f(x)$ . An extension to the case where  $t_1 = \tau$  is random, as in Section 3, is discussed in Remark A.3.

We make the following assumption on the random variable  $\Delta_{\tilde{\ell}} g(X_{t_1, \ell}, W_{t_1, T}) p_{\tilde{\ell}}^{-1}$ , introduced in Section 2.1.2, which is used in Theorem A.2 below. This assumption is comparable to those made in [14, 15].

**Assumption A.1.** *Let  $\delta(X_{t_1, \ell}) := \min_{1 \leq k \leq K} |\mathbb{E}[g(X_T) | X_{t_1} = X_{t_1, \ell}] - d_k|$ , where  $d_k$  are the points where  $f$  has a discontinuous derivative. Then, for all  $\ell$ ,  $\delta(X_{t_1, \ell})$  admits a density  $\tilde{\rho}_\ell$ . There exists constants  $\delta_0, \bar{\rho}_0 > 0$ , independent of  $\ell$ , such that  $x < \delta_0$  implies  $\tilde{\rho}_\ell(x) < \bar{\rho}_0$  for all  $\ell$ . Moreover, the third centralised moments of  $\Delta_{\tilde{\ell}} g(X_{t_1, \ell}, W_{t_1, T}) p_{\tilde{\ell}}^{-1}$  satisfy the following conditions:*

$$\begin{aligned} \sup_{\ell \geq 0} \mathbb{E} \left[ \left| \Delta_{\tilde{\ell}} g(X_{t_1, \ell}, W_{t_1, T}) p_{\tilde{\ell}}^{-1} - \mathbb{E}[g(X_T) | X_{t_1} = X_{t_1, \ell}] \right|^3 \right] &< \infty \\ \sup_{\delta(x) < \delta_0} \mathbb{E} \left[ \left| \Delta_{\tilde{\ell}} g(X_{t_1}, W_{t_1, T}) p_{\tilde{\ell}}^{-1} - \mathbb{E}[g(X_T) | X_{t_1}] \right|^3 \middle| X_{t_1} = x \right] &< \infty. \end{aligned}$$

The assumption that  $\delta(X_{t_1, \ell})$  has a bounded density is used in [14, 23] and controls the probability of sampling (an approximation of)  $X_{t_1}$  such that  $\mathbb{E}[g(X_T) | X_{t_1}]$  falls near a discontinuity in the derivative of  $f$ . Recall from Section 2.1.1 that  $\tilde{\ell}$  is a random integer with mass function  $\mathbb{P}[\tilde{\ell} = k] = p_k \propto 2^{-\zeta k}$



for fixed  $\zeta$  chosen between 1 and  $3/2$ , which ensures that

$$\mathbb{E} \left[ \left| \Delta_{\bar{\ell}} g(X_{t_1}, W_{t_1, T}) p_{\bar{\ell}}^{-1} \right|^3 \middle| X_{t_1} \right] < \infty,$$

for all  $X_{t_1}$ . The assumptions on the third centralised moment then follow.

The main result of this section is expressed in the following theorem:

**Theorem A.2.** *Assume that  $f$  is of the form (17), with Lipschitz constant  $L_f$ , and that  $g$  is either Lipschitz or twice differentiable with bounded second derivative. Assume the coefficients of the underlying SDE, and their derivatives, satisfy growth and Lipschitz conditions as in [24, Theorem 10.3.5]. Let  $1 < \zeta < 3/2$  with  $\mathbb{P}[\bar{\ell} = k] \propto 2^{-\zeta k}$  and assume Assumption A.1 holds. Then, there exists  $A_0 > 0$ , independent of  $\ell$ , with*

$$\mathbb{E} \left[ (\Delta_{\ell}^{\text{ant}} f)^2 \right] \leq A_0 2^{-3\ell/2}.$$

*Proof.* Throughout this proof we assume  $A_i$  are constants independent of  $\ell, k$ . We can express

$$\Delta_{\ell}^{\text{ant}} f = \Delta_{\ell}^{(1)} f + \Delta_{\ell}^{(2)} f,$$

where

$$\begin{aligned} \Delta_{\ell}^{(1)} f &= f(E_{\ell, f}^i(X_{t_1, \ell})) - f(E_{\ell, f}^i(X_{t_1, \ell-1})), \\ \Delta_{\ell}^{(2)} f &= f(E_{\ell, f}^i(X_{t_1, \ell-1})) - \frac{1}{2} \left( f(E_{\ell-1, c, 0}^i(X_{t_1, \ell-1})) + f(E_{\ell-1, c, 1}^i(X_{t_1, \ell-1})) \right). \end{aligned}$$

That is,  $\Delta_{\ell}^{(1)} f$  captures the error made from using Milstein discretisation with step sizes  $h_{\ell}$  and  $h_{\ell-1}$ , and  $\Delta_{\ell}^{(2)} f$  captures the error made from using Monte Carlo estimates at varying levels of accuracy. It follows that

$$\mathbb{E} \left[ (\Delta_{\ell}^{\text{ant}} f)^2 \right] \leq 2\mathbb{E} \left[ (\Delta_{\ell}^{(1)} f)^2 \right] + 2\mathbb{E} \left[ (\Delta_{\ell}^{(2)} f)^2 \right]. \quad (18)$$

At this point, we introduce the following notation:

- $X_{T, \ell, k}$  denotes the Milstein approximation of  $X_T$  using step-sizes  $h_{\ell}^m$  on  $[t_0, t_1]$  and  $h_k^i$  on  $[t_1, T]$ . We allow the cases  $\ell, k = \infty$  to indicate exact simulation on the intervals  $[t_0, t_1]$  or  $[t_1, T]$ , respectively.
- $g_{\ell, k} = g(X_{T, \ell, k})$ .
- For  $\Delta_k g(X_{t_1}, W_{t_1, T})$  as in Section 2.1.2, define the double difference

$$\begin{aligned} \Delta_{\ell} \Delta_k g &\equiv \Delta_{\ell} \Delta_k g(X_{t_1, \ell}, X_{t_1, \ell-1}, W_{t_1, T}) \\ &:= \Delta_k g(X_{t_1, \ell}, W_{t_1, T}) - \Delta_k g(X_{t_1, \ell-1}, W_{t_1, T}) \\ &= g_{\ell, k} - g_{\ell, k-1} - g_{\ell-1, k} + g_{\ell-1, k-1}, \end{aligned} \quad (19)$$

where the random variables  $X_{t_1, \ell}, X_{t_1, \ell-1}$  are determined by the same Brownian path  $W_{0, t_1}$ .

- For  $E_{\ell,f}^i$  as in Section 2.1.2, define the average

$$\begin{aligned}\overline{\Delta\Delta g} &= E_{\ell,f}^i(X_{t_1,\ell}) - E_{\ell,f}^i(X_{t_1,\ell-1}) \\ &= \frac{1}{M_\ell^i} \sum_{m=1}^{M_\ell^i} \Delta_\ell \Delta_{\tilde{\ell}(m)} g(X_{t_1,\ell}, X_{t_1,\ell-1}, W_{t_1,T}^{(m)}) p_{\tilde{\ell}(m)}^{-1},\end{aligned}$$

where the samples  $\Delta_\ell \Delta_{\tilde{\ell}(m)} g(X_{t_1,\ell}, X_{t_1,\ell-1}, W_{t_1,T}^{(m)}) p_{\tilde{\ell}(m)}^{-1}$  are i.i.d samples of  $\Delta_\ell \Delta_{\tilde{\ell}} g p_{\tilde{\ell}}^{-1}$  given  $X_{t_1,\ell}$  and  $X_{t_1,\ell-1}$ .

Note that if  $\bar{Y}$  is an average of  $M$  i.i.d. samples of a random variable  $Y$  given  $Z$ , then

$$\mathbb{E}\left[(\bar{Y})^2 \mid Z\right] = (\mathbb{E}[Y \mid Z])^2 + \frac{1}{M} \text{Var}[Y \mid Z] \leq (\mathbb{E}[Y \mid Z])^2 + \frac{1}{M} \mathbb{E}[Y^2 \mid Z],$$

and hence

$$\mathbb{E}\left[(\bar{Y})^2\right] \leq \mathbb{E}\left[(\mathbb{E}[Y \mid Z])^2\right] + \frac{1}{M} \mathbb{E}[Y^2].$$

Combining this with the fact that  $f$  is  $L_f$ -Lipschitz we have

$$\begin{aligned}\mathbb{E}\left[\left(\Delta_\ell^{(1)} f\right)^2\right] &\leq L_f^2 \mathbb{E}\left[\left(\overline{\Delta\Delta g}\right)^2\right] \\ &\leq L_f^2 \left(\mathbb{E}\left[\left(\mathbb{E}\left[\Delta_\ell \Delta_{\tilde{\ell}} g p_{\tilde{\ell}}^{-1} \mid X_{t_1,\ell}, X_{t_1,\ell-1}\right]\right)^2\right] + \frac{1}{M_\ell^i} \mathbb{E}\left[\left(\Delta_\ell \Delta_{\tilde{\ell}} g p_{\tilde{\ell}}^{-1}\right)^2\right]\right).\end{aligned}\quad (20)$$

For the first term, recall that

$$\mathbb{E}[\Delta_{\tilde{\ell}} g(X_{t_1,\ell}, W_{t_1,T}) p_{\tilde{\ell}}^{-1} \mid X_{t_1,\ell}] = \mathbb{E}[g_{\ell,\infty} \mid X_{t_1,\ell}].$$

Therefore,

$$\begin{aligned}&\mathbb{E}\left[\Delta_\ell \Delta_{\tilde{\ell}} g p_{\tilde{\ell}}^{-1} \mid X_{t_1,\ell}, X_{t_1,\ell-1}\right] \\ &= \mathbb{E}\left[\left(\Delta_{\tilde{\ell}} g(X_{t_1,\ell}, W_{t_1,T}) - \Delta_{\tilde{\ell}} g(X_{t_1,\ell-1}, W_{t_1,T})\right) p_{\tilde{\ell}}^{-1} \mid X_{t_1,\ell}, X_{t_1,\ell-1}\right] \\ &= \mathbb{E}[g_{\ell,\infty} - g_{\ell-1,\infty} \mid X_{t_1,\ell}, X_{t_1,\ell-1}].\end{aligned}$$

In particular, by Jensen's inequality and the tower property

$$\begin{aligned}\mathbb{E}\left[\left(\mathbb{E}\left[\Delta_\ell \Delta_{\tilde{\ell}} g p_{\tilde{\ell}}^{-1} \mid X_{t_1,\ell}, X_{t_1,\ell-1}\right]\right)^2\right] &\leq \mathbb{E}\left[(g_{\ell,\infty} - g_{\ell-1,\infty})^2\right] \\ &\leq A_1 2^{-2\ell},\end{aligned}$$

where we use the strong order of convergence of the Milstein scheme and the regularity conditions on  $g$  to obtain the final bound.

For the second term in (20), we have

$$\mathbb{E} \left[ \left( \Delta_\ell \Delta_{\bar{\ell}} g p_{\bar{\ell}}^{-1} \right)^2 \right] = \sum_{k=0}^{\infty} p_k^{-1} \mathbb{E} \left[ \left( \Delta_\ell \Delta_k g \right)^2 \right]. \quad (21)$$

When  $k < \ell$ , under the Lipschitz or smoothness conditions on  $g$  and by the strong order of convergence for Milstein [24], we have

$$\begin{aligned} \mathbb{E} \left[ \left( \Delta_\ell \Delta_k g \right)^2 \right] &\leq 2 \sum_{j=k-1}^k \mathbb{E} \left[ \left( g_{\ell,j} - g_{\ell-1,j} \right)^2 \right] \\ &\leq A_2 2^{-2\ell}. \end{aligned}$$

Likewise, when  $\ell < k$

$$\begin{aligned} \mathbb{E} \left[ \left( \Delta_\ell \Delta_k g \right)^2 \right] &\leq 2 \sum_{j=\ell-1}^{\ell} \mathbb{E} \left[ \left( g_{j,k} - g_{j,k-1} \right)^2 \right] \\ &\leq A_3 2^{-2k}. \end{aligned}$$

Hence, from (21),

$$\mathbb{E} \left[ \left( \Delta_\ell \Delta_{\bar{\ell}} g p_{\bar{\ell}}^{-1} \right)^2 \right] \leq A_4 2^{-(2-\zeta)\ell},$$

and, from (20), we have

$$\mathbb{E} \left[ \left( \Delta_\ell^{(1)} f \right)^2 \right] \leq L_f^2 \left( A_1 2^{-2\ell} + (M_0^i)^{-1} A_4 2^{-(3-\zeta)\ell} \right), \quad (22)$$

Now consider the second term in (18). Under Assumption A.1, a slight modification to the proof of [14, Theorem 4.1] gives  $\mathbb{E} \left[ \left( \Delta_\ell^{(2)} f \right)^2 \right] \leq A_5 (M_\ell^i)^{-3/2} = A_5 (M_0^i)^{-3/2} 2^{-3\ell/2}$ . Combining this fact with (18) and (22) proves the result.  $\square$

**Remark A.3.** *The proof of Theorem A.2 holds for all  $t_1 \in [t_0, T]$ , with constant  $A_0 = A_0(t_1)$ . By taking the maximum of  $A_0(t_1)$  over this interval, the result can be extended to the case where  $t_1 = \tau$  is a random variable taking values in  $[t_0, T]$ . This extends the results to the CVA problem considered in Section 3.*

**Remark A.4.** *It is possible to add another term  $f_0(x)$  to the right hand side of (17), where  $f_0$  is twice differentiable with bounded second derivative. The result of Theorem A.2 can be extended to this case under the same assumptions. However, since this is not required for the CVA problem, we omit a proof here.*

**Remark A.5.** *The preceding analysis can be repeated with  $\zeta = 3/2$ . Then, Assumption A.1 only holds with  $3 - \varepsilon$  centralised moments of  $\Delta_{\bar{\ell}} g p_{\bar{\ell}}^{-1}$ , for any  $\varepsilon > 0$ . In this case, the analysis can be adjusted to show that for any  $\varepsilon > 0$  we have*

$$\mathbb{E} \left[ \left( \Delta_\ell^{\text{ant}} f \right)^2 \right] = \mathcal{O}(2^{-3\ell/2+\varepsilon\ell}),$$

*and the discussion in this paper is unchanged.*