Efficient Risk Estimation for the Credit Valuation Adjustment

Michael B. Giles[∗] Abdul-Lateef Haji-Ali† Jonathan Spence‡

Abstract

AMS Subject Classication: 65C05, 62P05

Keywords: Multilevel Monte Carlo, Nested simulation, Risk estimation

The valuation of over-the-counter derivatives is subject to a series of valuation adjustments known as xVA, which pose additional risks for financial institutions. Associated risk measures, such as the value-at-risk of an underlying valuation adjustment, play an important role in managing these risks. Monte Carlo methods are often regarded as inefficient for computing such measures. As an example, we consider the value-at-risk of the Credit Valuation Adjustment (CVA-VaR), which can be expressed using a triple nested expectation. Traditional Monte Carlo methods are often inefficient at handling several nested expectations. Utilising recent developments in multilevel nested simulation for probabilities, we construct a hierarchical estimator of the CVA-VaR which reduces the computational complexity by several orders of magnitude compared to standard Monte Carlo.

1 Introduction

Financial derivatives traded on over-the-counter markets are subject to a series of valuation adjustments, collectively known as x-valuation adjustments (xVA) [\[8,](#page-25-0) [32,](#page-26-0) [33\]](#page-26-1), reflecting the value of external risk factors. The presence of xVA components in valuation formulae pose an additional source of risk to financial organisations: Fluctuations in external risk factors can adversely affect the adjusted value of a financial portfolio. A common feature of xVA calculations and associated risk measures is a recursive or nested dependency on market factors, which poses difficulties for numerical computation. This paper develops efficient hierarchical estimators for such problems, with an application presented for a risk estimate involving the credit valuation adjustment (CVA), which accounts for the possibility of a counterparty defaulting and failing to meet future payoffs. To mitigate the possibility of fluctuations in the CVA resulting in significant losses to a portfolio holder, the Basel Committee on Banking Supervision proposed a capital charge, which is based on a CVA value-at-risk formula over a short risk horizon $H \ll 1$ [\[4,](#page-25-1) [8,](#page-25-0) [43\]](#page-27-0). At a sufficiently high confidence level $\varphi \in (0,1)$, the (unilateral) CVA capital charge owed to a single counterparty can be expressed as the value $\lambda_{\varphi} \in \mathbb{R}$ solving [\[2,](#page-25-2)[8\]](#page-25-0)

$$
\varphi = \mathsf{P}\left[\frac{\text{CVA}_{H}}{B_{H}} - \text{CVA}_{0} > \lambda_{\varphi}\right],\tag{1a}
$$

$$
CVA_t = B_t \mathsf{E}^{\mathsf{Q}} \big[\, \chi_{t \le \tau \le T} \mathsf{LGD} \, B_\tau^{-1} \max \big\{ \mathsf{E}^{\mathsf{Q}} \big[\, \pi_{\tau,T} \, | \, \mathcal{G}_\tau \, \big], 0 \big\} \, \big| \, \mathcal{G}_t \, \big],\tag{1b}
$$

where X_A denotes the characteristic function of the event A, taking value 1 if A is true and 0 otherwise, and the remaining terms appearing in the above formula are defined as follows: Stochastic future payoffs accrued from default time τ to maturity T, discounted to time τ , are denoted by $\pi_{\tau,T}$

[∗]University of Oxford [\(mike.giles@maths.ox.ac.uk\)](mailto::mike.giles@maths.ox.ac.uk).

[†]Heriot-Watt University [\(a.hajiali@hw.ac.uk\)](mailto::a.hajiali@hw.ac.uk).

[‡]Corresponding author. University of Edinburgh [\(j.spence@ed.ac.uk\)](mailto::j.spence@ed.ac.uk).

and assumed to be determined by \mathbb{R}^d -valued assets $S = (S_t)_{t\geq 0}$ which have dynamics modelled through an SDE of the form

$$
dS_t = \mu(t, S_t)dt + \varsigma(t, S_t)dW_t,
$$
\n(2)

under a physical market measure, denoted P, with drift $\mu : [0, T] \times \mathbb{R}^d \to \mathbb{R}^d$, volatility $\varsigma : [0, T] \times$ $\mathbb{R}^d \to \mathbb{R}^{d \times m}$ and where $W = (W_t)_{t \geq 0}$ is a d'-dimensional Brownian motion adapted to a market filtration $\mathcal{F} := (\mathcal{F}_t)_{t \geq 0}$. B_t represents a discounting factor at time t, such that at time t the discounting factor is \overline{B}_t^{-1} , where $B_0 = 1$. We assume the existence of a risk-neutral measure Q, which is used for arbitrage-free pricing of derivatives, such that the discounted assets S_t/B_t are a W_t^{Q} -martingale, where $W^{\mathsf{Q}} = (W_t^{\mathsf{Q}})_{t \geq 0}$ is a Brownian motion under Q. Since P represents the true dynamics of the market, the risk measure [\(1a\)](#page-0-0) is computed under P . However, since CVA_t represents a valuation adjustment, it is priced under Q. The random future time of counterparty default is represented by τ , which is a stopping time with respect to an extended filtration $\mathcal{G} := (\mathcal{G}_t)_{t>0}$ such that $\mathcal{G}_t \supset \mathcal{F}_t$ for all $t \geq 0$, incorporating credit risk factors not present in the market filtration F. The term $\max\{E^Q[\pi_{\tau,T} | \mathcal{G}_{\tau}],0\}$ is the positive part of the (discounted) risk-neutral portfolio valuation at the default time τ and is referred to as the exposure at default, representing the amount the portfolio holder stands to lose at the time of counterparty default. Only the positive part of the portfolio value is considered in the CVA formulae since the portfolio holder must still typically repay any negative values, for example to creditors, in the event of counterparty default. The loss given default, LGD, is a \mathcal{G}_{τ} -measurable random variable taking values in $(0,1)$ and represents the proportion of expected future cash-flows which are lost in the event of counterparty default.

Due, for example, to complex market dynamics [\(2\)](#page-1-0), it is typically not possible to sample the assets S, respectively the payoffs $\pi_{\tau,T}$, exactly. Often, accurate approximations of the term φ [\(1a\)](#page-0-0) are obtained using expensive time-discretisations to approximate the process [\(2\)](#page-1-0). Furthermore, due to approximation of the repeatedly nested expectations defining $CVA_H, CVA₀$ and φ , the compounding cost of combining time-discrete approximations of the SDE [\(2\)](#page-1-0) within nested averages leads to impractically large costs involved in approximating [\(1a\)](#page-0-0).

Motivated by the problem of approximating the CVA capital charge [\(1a\)](#page-0-0), this paper considers efficient methods to evaluate the distribution of random variables which can be expressed as a sequence of nested expectations. Abstractly, for $\varphi \in (0,1)$ and $\lambda_{\varphi} \in \mathbb{R}$, we consider problems taking the form

$$
\varphi = P[U_0(Z) > \lambda_{\varphi}]
$$

\n
$$
U_0(z) = E[f(U_1(Y)) | Z = z]
$$

\n
$$
U_1(y) = E[X | Y = y],
$$
\n(3)

where $f : \mathbb{R} \to \mathbb{R}$ is Lipschitz continuous, but not necessarily everywhere differentiable, and X, Y, Z are random variables, with X taking values in $\mathbb R$ and Y, Z taking values in $\mathbb R^d$. In particular, we do not require that the variables X, Y, Z, $U_0(Z)$ and $U_1(Y)$ can be sampled directly, but rather through a hierarchy of increasingly accurate approximations. Letting c_H be an \mathbb{R}^{d_c} random variable denoting the value at the horizon H of all relevant \mathcal{G}_H -measurable credit risk factors, influencing the default times τ , the CVA problem [\(1a\)](#page-0-0) can be expressed in the form [\(3\)](#page-1-1), with

- $Z = (S_H, c_H)$ is simulated under P and \mathcal{G}_H -measurable, taking values in \mathbb{R}^{d+d_c} ,
- $Y = (S_\tau, \tau)$ is simulated under Q and G_τ -measurable, taking values in \mathbb{R}^{d+1} , (4)
- $X = \text{LGD}\, \chi_{H \leq \tau \leq T} B_{\tau}^{-1} \pi_{\tau,T}$ is simulated under Q and \mathcal{G}_T -measurable, taking values in R,

each depending on the solution to an SDE of the form [\(2\)](#page-1-0). The connection between the problem [\(1a\)](#page-0-0) and the general equation [\(3\)](#page-1-1) is discussed further in Section [3.2.](#page-13-0) When considering the general problem [\(3\)](#page-1-1), we assume each expectation and probability is computed with respect to a single, arbitrary measure, P. Alternatively, when considering the CVA problem [\(1\)](#page-0-1), we explicitly track the dependency of each term on the physical measure, P, and the risk-neutral measure, Q. We focus primarily on the problem of approximating the probability of large loss φ given a fixed λ_{φ} . Such methods can then be combined with stochastic root finding techniques to find the quantile λ_{φ} corresponding to a fixed confidence level φ [\[13,](#page-25-3) [15,](#page-25-4) [16,](#page-25-5) [19\]](#page-26-2).

Monte Carlo estimators to approximate probabilities containing a single nested expectation of the form $P[E[X|Y] > \lambda_{\varphi}]$ are considered in [\[9,](#page-25-6)[31\]](#page-26-3), with multilevel extensions considered in [\[24,](#page-26-4) [26,](#page-26-5) [34\]](#page-26-6). Stochastic approximation methods to compute the corresponding value-at-risk λ_{φ} for such problems are considered in [\[3,](#page-25-7)[13,](#page-25-3)[14,](#page-25-8)[19\]](#page-26-2). An application of MLMC methods to problems defined by arbitrarily many nested Lipschitz transformations of conditional expectations is considered in [\[47,](#page-27-1)[49\]](#page-27-2) for problems arising in optimal control and non-European option pricing. Within the CVA framework, multilevel methods have previously been employed in [\[36\]](#page-26-7) to the problem of calibrating CVA model parameters to observed market data. The key contributions of this paper are as follows:

- A novel combination of multilevel Monte Carlo (MLMC) methods for nested expectations and probabilities [\[24,](#page-26-4)[34\]](#page-26-6) to compute φ in [\(3\)](#page-1-1) is analysed in Section [2.](#page-2-0) Under general assumptions on the underlying variables, Theorem [2.6](#page-9-0) proves that the proposed estimator approximates φ with root mean square error ε with order $\varepsilon^{-2-\delta}$ computational cost, for any $\delta > 0$. This is contrasted with an order ε^{-5} computational cost using standard Monte Carlo techniques for the same problem.
- Section [3](#page-11-0) discusses an implementation of the hierarchical estimator for the CVA risk problem [\(1a\)](#page-0-0). Section [4](#page-15-0) leverages variance reduction techniques from the literature to further improve the efficiency of the estimator. Extensions of the model to include credit risk mitigation factors such as the posting of collateral and to portfolios with more than a single counterparty are discussed in Sections [3.3](#page-14-0) and [4.4.](#page-20-0)
- A numerical study to support theoretical results is conducted in Section [5.](#page-21-0)

2 Nested Multilevel Monte Carlo Estimator

In this section we consider the construction of efficient MLMC estimators for the general problem (3) defined for some random variables X, Y, Z . We begin by outlining key assumptions on (approximations) of the underlying variables in Section [2.1,](#page-2-1) before considering a single-level Monte Carlo estimator in Section [2.2](#page-3-0) and multilevel Monte Carlo estimator in Section [2.3.](#page-5-0)

2.1 Setup and Assumptions

In the context of financial risk estimation, the variable Z in [\(3\)](#page-1-1) typically represents random market fluctuations over a short time horizon. As such, it is often possible to obtain sufficiently accurate samples of Z with negligible cost. To simplify the discussion, we therefore assume that Z can be sampled exactly, while emphasising that the methods which follow can be extended naturally to include approximation of Z. Given samples of Z, we assume that X and Y can be approximated to sufficient accuracy, but with increasing cost as the error converges to zero: For $\ell, k \geq 0$, we denote approximations $Y_{\ell}(Z) = Y_{\ell} \approx Y(Z)$ and $X_k(Y) \approx X(Y)$, where the notation $X(Y)$ represents the random variable X conditioned on the value of Y. In particular, we impose the following conditions on (approximations of) X and Y :

Assumption 2.1. For all $\ell, k \geq 0$ and $q \geq 2$, there exists a constant $c > 0$, independent of ℓ and k such that

$$
\operatorname{Cost}(X_k(Y)) \le c2^k, \qquad \operatorname{Cost}(Y_\ell(Z)) \le c2^\ell, \n\mathsf{E}[|X_k(Y_\ell) - X(Y_\ell)|^q] \le c2^{-qk}, \qquad \qquad \mathsf{E}[\|Y_\ell - Y\|^q] \le c2^{-q\ell}, \n\mathsf{E}[X_k(Y_\ell) - X(Y)]| \le c2^{-\min\{\ell, k\}}, \qquad \|\mathsf{E}[Y_\ell - Y]\| \le c2^{-\ell}.
$$

When the variables X and Y take the form (4) depending on the solution of an SDE (2) and are approximated using Milstein discretisation with step size $h_{\ell} \propto 2^{-\ell}$ at level ℓ , Assumption [2.1](#page-3-1) holds, assuming $\pi_{\tau,T}$ is a Lipschitz functional of the assets S, whenever the coefficients μ and ς of the SDE are Lipschitz continuous, twice differentiable and, combined with their first and (mixed) second order derivatives, satisfy linear growth conditions as specified in $[40,$ Theorem 10.3.5. The order of convergence specified by this assumption enables the use of randomised MLMC techniques [\[23](#page-26-9)[,44](#page-27-3)[,45\]](#page-27-4) in Section [2.3](#page-5-0) to sample unbiased estimators of $U_1(Y)$ and $U_0(Z)$ with finite cost and variance, see Proposition [2.3.](#page-6-0)

We additionally impose the following condition on the function f within (3) .

Assumption 2.2. The function $f : \mathbb{R} \to \mathbb{R}$ is L_f -Lipschitz on \mathbb{R} , and twice differentiable on $\mathbb{R} \backslash \Gamma$ for some $\Gamma \subset \mathbb{R}$. The second derivative of f is uniformly bounded by $L'_f > 0$. That is,

$$
\sup_{u \in \mathbb{R} \backslash \Gamma} |f''(u)| \le L_f'.\tag{5}
$$

Moreover, there is $\bar{\delta}, \bar{\rho} > 0$ such that $0 < x \leq \bar{\delta}$ implies

$$
\mathsf{P}\bigg[\inf_{u\in\Gamma}|U_1(Y)-u|\leq x\bigg]\leq \bar{\rho}\,x,\tag{6}
$$

where $\bar{\delta}$ and $\bar{\rho}$ are independent of x.

Similar conditions to Assumption [2.2](#page-3-2) are imposed in [\[7,](#page-25-9)[28\]](#page-26-10) to approximate averages of Lipschitz functionals, f, of an underlying quantity of interest using antithetic MLMC differences based on a Taylor expansion of f. This methodology is used in Section [2.3](#page-5-0) to approximate $U_0(Z)$. The global bound on the second derivative (5) can be used to bound a second-order Taylor expansion of f on regions where the function is smooth. The additional condition (6) allows f to be non-continuously differentiable on a set Γ, which is essential when considering the CVA in [\(1b\)](#page-0-2), provided we can linearly bound the probability that $U_1(Y)$ lies a distance x from Γ .

2.2 Nested Monte Carlo Estimate

To motivate our work and as a baseline for comparison, we begin by considering an estimator for the quantity φ defined by [\(3\)](#page-1-1) using a hierarchy of single-level Monte Carlo averages. In particular, due to the recursively nested structure of the problem and compounding approximation costs through several nested Monte Carlo estimates, we will see that accurate computation of φ is expensive. We construct the estimator iteratively, starting from the innermost conditional expectation and working outward.

For a fixed sample of Y, a Monte Carlo average for the innermost conditional expectation $U_1(Y)$, given Y, at level $\mathbf{k} = (k_0, k_1) \in \mathbb{N}^2$ is

$$
\widehat{U}_{1,\mathbf{k}}(Y) := \frac{1}{N_{k_0}} \sum_{n_1=1}^{N_{k_0}} X_{k_1}^{(n_1)}(Y),
$$

where we set $N_{k_0} = N_0 2^{k_0}$ and $\{X_{k_1}^{(n_1)}\}$ $\binom{(n_1)}{k_1}$ $\binom{N_{1,k_0}}{n_1-1}$ are independent samples of $X_{k_1}(Y)$ given Y. At approximation level k_1 , the cost of sampling X_{k_1} is of order 2^{k_1} under Assumption [2.1.](#page-3-1) Hence, the sampling cost of $\hat{U}_{1,\boldsymbol{k}}(Y)$ is of order $N_{k_0} \text{Cost}(X_{\ell}(Y)) \propto 2^{k_0+k_1}$.

Similarly, a (biased) Monte Carlo average for the conditional expectation $U_0(Z)$, given Z, at level $\ell = (\ell_0, \ell_1) \in \mathbb{N}^2$ is then given by

$$
\widehat{U}_{0,\ell,{\bf k}}(Z) \coloneqq \frac{1}{N_{\ell_0}} \sum_{n_0=1}^{N_{\ell_0}} f\Big(\widehat{U}_{1,{\bf k}}^{(n_0)}\Big(Y_{\ell_1}^{(n_0)}(Z)\Big)\Big),\,
$$

where $\{Y_{\ell_1}^{(n_0)}\}$ $\mathcal{U}_{\ell_1}^{(n_0)}(Z)\}_{n_0=1}^{N_{\ell_0}}$ and $\{\widehat{U}_{1,\boldsymbol{k}}^{(n_0)}\left(Y_{\ell_1}^{(n_0)}\right)$ $\binom{n_0}{\ell_1}(Z)$ $\big\}^{N_{\ell_0}}_{n_0=1}$ are conditionally independent samples of $Y_{\ell_1}(Z)$ and $\hat{U}_{1,k}(Y(Z))$ given Z. Since samples of $Y_{\ell_1}(Z)$ can be generated with cost of order 2^{ℓ_1} under Assumption [2.1,](#page-3-1) it follows that

$$
Cost(\widehat{U}_{0,\ell,k}(Z)) = \mathcal{O}\Big(N_{\ell_0}\Big(\mathrm{Cost}(\widehat{U}_{1,\boldsymbol{k}}(Y)) + \mathrm{Cost}(Y_{\ell_1}(Z))\Big)\Big) = \mathcal{O}(2^{\ell_0 + k_0 + k_1} + 2^{\ell_0 + \ell_1}).
$$
\n(7)

Using the above nested Monte Carlo estimator for U_0 , a nested Monte Carlo estimate for $P[U_0(Z) > \lambda_{\varphi}]$ is given by

$$
\varphi_{M,\boldsymbol{\ell},\boldsymbol{k}} = \frac{1}{M} \sum_{m=1}^{M} \chi_{\widehat{U}_{0,\boldsymbol{\ell},\boldsymbol{k}}^{(m)}(Z^{(m)}) > \lambda_{\varphi}},
$$

where $\{(Z^{(m)}, \widehat{U}_{0,\ell,k}^{(m)}(Z^{(m)}))\}_{m=1}^M$ are independent samples of $(Z, \widehat{U}_{0,\ell,k}(Z))$, respectively. The sampling cost of this nested estimate is $Cost(\varphi_{M,\ell,k}) = \mathcal{O}(M(2^{\ell_0+k_0+k_1}+2^{\ell_0+\ell_1}))$. Meanwhile, the mean square error of the approximation $\varphi_{M,\ell,k}$ can be decomposed into a bias and statistical error through $\mathsf{E}[(\varphi - \varphi_{M,\ell,k})^2] = |\mathsf{E}[\varphi - \varphi_{M,\ell,k}]|^2 + \mathsf{Var}[\varphi_{M,\ell,k}]$. By following a similar argument to [\[10,](#page-25-10) Proposition 1], under Assumption [2.1](#page-3-1) and assuming that the joint distribution of $U_0(Z)$ and $U_{0,\ell,k}(Z)$ admits a density $\rho(u,\hat{u})$ which is twice differentiable in u for all ℓ, k , one can show that $|E[\varphi - \varphi_{M,\ell,\mathbf{k}}]|$ is of order $2^{-\min\{\ell_0,\ell_1,k_0,k_1\}}$. The hierarchical estimator discussed in the following section does not require a bound on the estimation bias of this form and therefore we do not include a rigorous proof of this bound. Furthermore, it can be shown that $Var[\varphi_{M,\ell,k}] = \mathcal{O}(M^{-1}).$ Therefore, to ensure $\mathsf{E}[(\varphi - \varphi_{M,\ell,\mathbf{k}})^2] \leq \varepsilon^2$, we require $M = \mathcal{O}(\varepsilon^{-2})$, and $2^{-\min\{\ell_0,\ell_1,k_0,k_1\}} = \mathcal{O}(\varepsilon)$. Hence, the cost of attaining a root mean square error ε for the estimator $\varphi_{M,\ell,k}$ is of order ε^{-5} . This scaling of computational cost makes it extremely expensive to attain sufficiently accurate estimates of φ using $\varphi_{M,\ell,k}$, particularly for the root finding problem of computing λ_{φ} in [\(3\)](#page-1-1), which requires computing several estimates of $\varphi_{M,\ell,k}$ with different loss thresholds λ_{φ} . This motivates the development of more advanced estimators of φ , with a computational cost which scales more efficiently with the designated error tolerance ε .

2.3 Nested Unbiased MLMC Estimate

To improve the computational cost, we first consider the use of recursive randomised MLMC esti-mation [\[41,](#page-27-5) [44,](#page-27-3) [45\]](#page-27-4) to construct an estimator of $U_0(Z)$ with a cost which is lower than that in [\(7\)](#page-4-0). We begin by considering the innermost expectation $U_1(Y)$. Define the multilevel differences, for a given Y , by

$$
\Delta_k X(Y) := \begin{cases} X_k(Y) - X_{k-1}(Y) & k \ge 1 \\ X_0(Y) & k = 0, \end{cases}
$$

so that under Assumption [2.1](#page-3-1) we have $U_1(Y) = \mathsf{E}[\sum_{k=0}^{\infty} \Delta_k X(Y) | Y]$. Then, for $\zeta_1 > 1$, define a random non-negative integer κ_1 , which is independent of $\{X_k\}_{k=0}^{\infty}$, satisfying $p_{1,k} := \mathsf{P}[\kappa_1 = k] =$ $(1 - 2^{-\zeta_1}) 2^{-\zeta_1 k}$. It follows that $[26, 45]$ $[26, 45]$

$$
U_1(Y) = \mathsf{E}\left[\sum_{k=0}^{\infty} \Delta_k X(Y) \middle| Y \right]
$$

= $\mathsf{E}\left[\sum_{k=0}^{\infty} \left(\Delta_k X(Y) p_{1,k}^{-1}\right) p_{1,k} \middle| Y \right]$
= $\mathsf{E}\left[\Delta_{\kappa_1} X(Y) p_{1,\kappa_1}^{-1} \middle| Y \right],$ (8)

where p_{1,κ_1} denotes the probability mass function $p_{1,k}$ evaluated at the random variable κ_1 . An unbiased Monte Carlo estimate of $U_1(Y)$ is thus given by

$$
\widehat{U}_{1,\ell}^{\mathrm{(UB)}}(Y) = \frac{1}{N_{1,\ell}} \sum_{n=1}^{N_{1,\ell}} \Delta_{\kappa_1^{(n)}} X^{(n)}(Y) \, p_{1,\kappa_1^{(n)}}^{-1},
$$

where $N_{1,\ell} = N_{1,0} 2^{\ell}$ and $\{(\kappa_1^{(n)}, \Delta_{\kappa_1^{(n)}} X^{(n)}(Y))\}_{n=1}^{N_{1,\ell}}$ are conditionally independent samples of $(\kappa_1, \Delta_{\kappa_1^{(n)}} X(Y))$, given Y. Under the convergence conditions in Assumption [2.1,](#page-3-1) it follows that $\widehat{U}_{1,\ell}^{(\text{UB})}(Y)$ has finite variance and an expected sampling cost of order $N_{1,\ell}$ provided $1 < \zeta_1 < 2$ [\[26,](#page-26-5) [45\]](#page-27-4).

Following the approach in [\[7,](#page-25-9) [10,](#page-25-10) [24\]](#page-26-4), we use the unbiased Monte Carlo estimates of $U_1(Y)$ to construct an antithetic multilevel difference for the term $f(U_1(Y))$ appearing within $U_0(Z)$. Specifically, we define

$$
\Delta_{\ell}^{(\text{UB})} f(Y_{\ell}, Y_{\ell-1}) = f\left(\hat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell})\right) - \frac{1}{2} \sum_{i=0}^{1} f\left(\hat{U}_{1,\ell-1}^{(\text{UB},c,i)}(Y_{\ell-1})\right),\tag{9}
$$

where $\widehat{U}_{1,\ell-1}^{(\text{UB},c,i)}(Y_{\ell-1})$ are conditionally independent samples of $\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell-1})$, given $Y_{\ell-1}$, for $i=0,1$, defined by

$$
\widehat{U}_{1,\ell-1}^{(\text{UB},c,i)}(Y_{\ell-1}) = \frac{1}{N_{1,\ell-1}} \sum_{n=1}^{N_{1,\ell-1}} \Delta_{\kappa_1^{(n)}} X^{(n+iN_{\ell-1})}(Y_{\ell-1}) p_{1,\kappa_1^{(n)}}^{-1},
$$

where the samples $\Delta_{\kappa_1^{(n)}} X^{(n)}(Y_{\ell-1})$ should be correlated to $\Delta_{\kappa_1^{(n)}} X^{(n)}(Y_{\ell})$ which are used to compute $\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell})$, in the sense of Assumption [2.1.](#page-3-1) In particular, for a given k, the \mathcal{L}^q -error between both terms can be bounded by

$$
\mathsf{E}\Big[\Big|\Delta_k X^{(n)}(Y_\ell) - \Delta_k X^{(n)}(Y_{\ell-1})\Big|^q\Big] \n= \mathsf{E}\Big[\Big|X_k^{(n)}(Y_\ell) - X_{k-1}^{(n)}(Y_\ell) - X_k^{(n)}(Y_{\ell-1}) + X_{k-1}^{(n)}(Y_{\ell-1})\Big|^q\Big] \n\leq a 2^{-q \max\{\ell, k\}},
$$

where we use Jensen's inequality to bound

$$
\mathsf{E}\Big[\left|\Delta_k X^{(n)}(Y_\ell) - \Delta_k X^{(n)}(Y_{\ell-1})\right|^q\Big] \leq 2^{q-1} \bigg(\mathsf{E}\Big[\left|X_k^{(n)}(Y_\ell) - X_{k-1}^{(n)}(Y_\ell)\right|^q\Big] + \mathsf{E}\Big[\left|X_k^{(n)}(Y_{\ell-1}) + X_{k-1}^{(n)}(Y_{\ell-1})\right|^q\Big]\bigg),
$$

when $k \geq \ell$, and again to bound

$$
\begin{aligned} \mathsf{E}\Big[\left|\Delta_k X^{(n)}(Y_\ell)-\Delta_k X^{(n)}(Y_{\ell-1})\right|^q\Big] &\leq 2^{q-1}\bigg(\mathsf{E}\Big[\left|X_k^{(n)}(Y_\ell)-X_k^{(n)}(Y_{\ell-1})\right|^q\Big] \\ &\qquad\qquad +\mathsf{E}\Big[\left|X_{k-1}^{(n)}(Y_\ell)-X_{k-1}^{(n)}(Y_{\ell-1})\right|^q\Big]\bigg), \end{aligned}
$$

when $\ell > k$, where the former term is bounded by $c2^{-k}$ and the latter by $c2^{-\ell}$ under Assumption [2.1.](#page-3-1) In particular, on the event $Y_{\ell} = Y_{\ell-1} = y$, it follows that

$$
\widehat{U}_{1,\ell}^{(\text{UB})}(y) - \frac{1}{2} \sum_{i=0}^{1} \widehat{U}_{1,\ell-1}^{(\text{UB},c,i)}(y) = 0.
$$

Therefore, on a linearisation of f, it follows that $\Delta_{\ell}^{\text{(UB)}} f(Y_{\ell}, Y_{\ell-1})$ is dominated by the approximation error between Y_{ℓ} and $Y_{\ell-1}$. This fact can be used to prove Lemma [A.1](#page-27-6) in the appendix, which can be used to establish the convergence of the moments of $\Delta_{\ell}^{\text{(UB)}} f(Y_{\ell}, Y_{\ell-1})$.

Randomised MLMC techniques can be applied recursively to create an unbiased Monte Carlo estimator of $U_0(Z)$, based on a randomisation of the approximation level ℓ used for the antithetic differences $\Delta_{\ell}^{\text{(UB)}} f(Y_{\ell}, Y_{\ell-1})$. Let κ_0 be a random, non-negative, integer with mass function $p_{0,\ell}$:= $P[\kappa_0 = \ell] = (1 - 2^{-\zeta_0})2^{-\zeta_0 \ell}$. Then, conditioned on Z, consider the random variable

$$
\Delta(Z) := \Delta_{\kappa_0}^{(\text{UB})} f\!\left(Y_{\kappa_0}(Z), Y_{\kappa_0 - 1}(Z)\right) p_{0,\kappa_0}^{-1},\tag{10}
$$

where p_{0,κ_0} denotes the probability mass $p_{0,k}$ evaluated at the random variable κ_0 . By a similar calculation to [\(8\)](#page-5-1), $\Delta(Z)$ satisfies $U_0(Z) = E[\Delta(Z) | Z]$. In particular, we can consider the unbiased Monte Carlo estimate of $U_0(Z)$ defined by

$$
\widehat{U}_{0,\ell}^{(\text{UB})}(Z) = \frac{1}{N_{0,\ell}} \sum_{n=1}^{N_{0,\ell}} \Delta^{(n)}(Z),\tag{11}
$$

for an increasing sequence of step sizes $N_{0,\ell} := N_{0,0} 2^{\gamma \ell}$, with $\gamma > 0$ fixed. The following result, which is proved in Appendix [A,](#page-27-7) can be used to bound the sampling cost and variance of a sample from [\(11\)](#page-6-1), with fixed values of ζ_0 and ζ_1 .

Proposition 2.3. Let Assumptions [2.1](#page-3-1) and [2.2](#page-3-2) hold and consider the variable $\Delta(Z)$ defined by [\(10\)](#page-6-2), where we fix $\zeta_0 = 17/16$ and $\zeta_1 = 3/2$. Then, $\Delta(Z)$ has finite expected sampling cost and variance. Moreover, $E[|\Delta(Z)|^{q_0}] < \infty$ for all $q_0 < 23/11$.

As a consequence of Proposition [2.3,](#page-6-0) for the same values of ζ_0, ζ_1 and q_0 , it follows that the Monte Carlo estimate [\(11\)](#page-6-1) has a sampling cost of order $N_{0,\ell}$ and finite q_0^{th} moment.

Remark 2.4. By the linearity of expectation, the approach in this section can be applied naturally in the case where $U_0(Z)$ or $U_1(Y)$ are replaced with a linear combination of expectations of the same form, each satisfying Assumptions [2.1](#page-3-1) and [2.2.](#page-3-2) This fact will be useful when considering the CVA problem in Section [4.](#page-15-0)

More generally, Assumption [2.1](#page-3-1) could be generalised to include Euler-Maruyama type convergence rates by relaxing the convergence rates $2^{-q\ell}$ and 2^{-qk} to $2^{-q\ell/2}$ and $2^{-qk/2}$ therein, however the conclusion of Proposition [2.3](#page-6-0) will no longer hold true. In this setting, it is possible to construct a hierarchy of nested biased MLMC estimates for $U_0(Z)$ and $U_1(Y)$ [\[46\]](#page-27-8). An application of such an approach to the CVA risk problem [\(1\)](#page-0-1) is left to future work.

2.4 Adaptive MLMC for Probabilities

Using the estimator [\(10\)](#page-6-2), the problem [\(3\)](#page-1-1) can be reduced to

$$
\varphi = P[U_0(Z) > \lambda_{\varphi}] = E[\chi_{U_0(Z) > \lambda_{\varphi}}],
$$
\n
$$
U_0(Z) = E[\Delta(Z) | Z],
$$
\n
$$
(12)
$$

where $\Delta(Z)$ can be sampled with finite expected sampling cost. At level $\ell \geq 0$, we consider the Monte Carlo estimate $\widehat{U}_{0,\ell}^{(\mathrm{UB})}(Z)$ defined in [\(11\)](#page-6-1). Since Assumption [2.2](#page-3-2) does not hold for the discontinuous function $\chi_{x>\lambda_{\varphi}}$, the methods and analysis of the previous section are not applicable to construct an unbiased estimator for φ which has both finite sampling cost and variance. Instead, we consider a truncated (biased) MLMC estimate [\[20\]](#page-26-11) defined using the multilevel differences [\[24\]](#page-26-4)

$$
\Delta_{\text{ind},\ell} := \begin{cases} \chi_{\widehat{U}_{0,0}^{(\text{UB})}(Z) > \lambda_{\varphi}} & \ell = 0\\ \chi_{\widehat{U}_{0,\ell}^{(\text{UB})}(Z) > \lambda_{\varphi}} - \chi_{\widehat{U}_{0,\ell-1}^{(\text{UB})}(Z) > \lambda_{\varphi}} & \ell > 0, \end{cases}
$$
(13)

through

$$
\varphi \approx \varphi_{\mathbf{M}_{L},L}^{(\text{MLMC})} \coloneqq \sum_{\ell=0}^{L} \frac{1}{M_{\ell}} \sum_{m=1}^{M_{\ell}} \Delta_{\text{ind},\ell}^{(m)}, \tag{14}
$$

for a truncation level $L \in \mathbb{N}$ and inner sample sizes $M_L = (M_0, \ldots, M_L) \in \mathbb{N}^{L+1}$ and where $\{\Delta_{\mathrm{ind},\ell}^{(m)}\}_{m=1}^{M_{\ell}}$ are independent samples of $\Delta_{\mathrm{ind},\ell}$ for $0 \leq \ell \leq L$. By the linearity of expectation, $\varphi_{\mathbf{M}_L,L}^{(\text{MLMC})}$ is an unbiased estimate of $P[\hat{U}_{0,L}^{(\text{UB})}(Z) > \lambda_{\varphi}] \approx \varphi$. Furthermore, for large ℓ , the variance of $\Delta_{\text{ind},\ell}$ becomes small, requiring fewer samples at costly levels $\ell \gg 1$ to control the variance of the estimator $\varphi_{\mathbf{M}_{L},L}^{(\text{MLMC})}$. From standard MLMC complexity analysis [\[11,](#page-25-11) [21\]](#page-26-12), since the cost of sampling $\Delta_{\text{ind},\ell}$ is of order $N_{0,\ell}$ by Proposition [2.3,](#page-6-0) for a specified root mean square error of ε it is optimal to allocate $\mathbf{M}_{L} = \mathbf{M}_{L}^{*}$, defined component-wise by $M_{\ell}^{*} \propto \varepsilon^{-2} \sqrt{V_{\ell} N_{0,\ell}^{-1}} \sum_{k=0}^{L} \sqrt{V_{k} N_{0,k}}$, where $V_{\ell} \coloneqq \text{Var}[\Delta_{\text{ind},\ell}]$. For this choice of sample sizes, the cost of the MLMC estimate is

$$
Cost\left(\varphi_{\mathbf{M}_{L}^{\star},L}^{(\text{MLMC})};\varepsilon\right) \propto \varepsilon^{-2} \left(\sum_{\ell=0}^{L} \sqrt{V_{\ell} N_{0,\ell}}\right)^{2},\tag{15}
$$

where the truncation level L should be chosen such that the bias $|\mathsf{E}[\varphi-\varphi_{\mathbf{M}_L,L}^{(\text{MLMC})}]| \leq \varepsilon/2$. From [\[31,](#page-26-3) Proposition 1, under the assumption that the joint distribution of $U_0(Z)$ and $\widehat{U}_{0,\ell}^{(\text{UB})}(Z)$ exists and is twice differentiable in $U_0(Z)$, it follows that $|E[\Delta_{ind,\ell}]| = O(N_{0,\ell}^{-1})$. In particular, to meet a specified root mean square error tolerance, ε , one should scale L such that $2^{-L} = \mathcal{O}(\varepsilon)$. In practice, both the bias $|E[\Delta_{ind,\ell}]|$ and variances V_{ℓ} must be approximated from the available samples of $\Delta_{\text{ind},\ell}$ in order to estimate the optimal truncation level L and \mathbf{M}^\star_L .

To study the asymptotic behaviour of this estimator, we impose an additional condition on the approximations of $U_0(Z)$ [\[34,](#page-26-6) Assumptions 1.3 and 1.4]. Within this assumption and in what follows, we introduce an abstract sequence of positive-valued random variables $\{\sigma_\ell\}_{\ell \in \mathbb{N}}$, which can be used to improve the concentration properties of the normalized statistic $\sigma_{\ell}^{-1} \widehat{U}_{0,\ell}^{(\text{UB})}(Z)$.

Assumption 2.5. For some $p > 2$ and a sequence $\{\sigma_\ell\}_{\ell \in \mathbb{N}}$ of positive-valued random variables, the pth centralized moment of $\widehat{U}_{0,\ell}^{(\text{UB})}(Z)$, normalized by $\sigma_{\ell} N_{0,\ell}^{-1/2}$, is finite. That is,

$$
\lim_{\ell_0 \to \infty} \sup_{\ell > \ell_0} \mathsf{E}\left[\left|\frac{\widehat{U}_{0,\ell}^{(\text{UB})}(Z) - U_0(Z)}{\sigma_\ell N_{0,\ell}^{-1/2}}\right|^p\right] < \infty. \tag{16}
$$

Moreover, there exists $\delta_0, \rho_0 > 0$ such that

$$
\sup_{\substack{\ell \geq 0 \\ 0 < x \leq \delta_0}} \mathsf{P}\left[\left|\frac{\widehat{U}_{0,\ell}^{(\text{UB})}(Z) - \lambda_{\varphi}}{\sigma_{\ell}}\right| < x\right] \leq \rho_0 x. \tag{17}
$$

Within Assumption [2.5,](#page-8-0) the condition [\(16\)](#page-8-1) controls the \mathcal{L}^p -error, normalized by σ_{ℓ} , made in approximating the nested expectation $U_0(Z)$ by the unbiased MLMC estimate [\(11\)](#page-6-1). Under the conditions and result of Proposition [2.3,](#page-6-0) it follows that this condition holds for any constant $\sigma_\ell \equiv$ $\sigma \in \mathbb{R}$ and any $p < 23/11$. Alternatively, one can consider setting σ_{ℓ}^2 to be the sample variance of the samples used to compute $\widehat{U}_{0,\ell}^{(\mathrm{UB})}(Z)$,

$$
\sigma_{\ell}^2 := \frac{1}{N_{0,\ell}} \sum_{n=1}^{N_{0,\ell}} \left(\Delta^{(n)}(Z) - \widehat{U}_{0,\ell}^{(\text{UB})}(Z) \right)^2.
$$
 (18)

In this context, the random variable $(\widehat{U}_{0,\ell}^{(\text{UB})}(Z) - U_0(Z))\sigma_{\ell}^{-1}N_{0,\ell}^{1/2}$ is equivalent to Student's test statistic for the samples ${\{\Delta^{(n)}(Z)\}}_{n=1}^{N_{0,\ell}}$. The samples $\Delta(Z)$ satisfy the hypothesis of the central limit theorem from Proposition [2.3](#page-6-0) and, consequently, one can leverage results from the theory of self-normalising sums $\left[42\right]$ to analyse the condition [\(16\)](#page-8-1). In particular, under the assumption that there exists ρ_0 , $\lambda > 0$ such that

$$
\sup_{\delta \in \mathbb{R}, z \in \mathbb{R}^d} \mathsf{P}\bigg[\left| \Delta(Z) - U_0(Z) - \delta \right| \leq |\delta| x \, \bigg| \, Z = z \bigg] < \rho_0 x^\lambda,\tag{19}
$$

it follows from [\[39,](#page-26-13) Theorem 6.2] that [\(16\)](#page-8-1) holds for any $p < \infty$. In addition, the bound [\(17\)](#page-8-2) controls the probability of sampling an approximation for $U_0(Z)$ which lies close to the discontinuity at λ_{φ} . Under Assumption [2.5,](#page-8-0) it follows from [\[34,](#page-26-6) Proposition 2.3, Corollary 2.10] that V_{ℓ} is of order $N_{0,\ell}^{-p/2(p+1)}$ and therefore $\text{Cost}(\varphi_{\mathbf{M}_{L},L}^{(\text{MLMC})};\varepsilon)$, for $N_{0,L}^{-1} = \mathcal{O}(\varepsilon)$, is of order $\varepsilon^{-3+p/2(p+1)}$. While the hierarchical approach improves upon the order ε^{-5} cost discussed in Section [2.2,](#page-3-0) the cost still scales worse than order ε^{-2} , which would be observed using standard Monte Carlo simulation if one could sample $U_0(Z)$ exactly with finite cost. The increased cost for the estimator $\varphi_{\mathbf{M}_L,L}^{(\text{MLMC})}$ can be attributed to the fact that an order 1 error in $\Delta_{ind,\ell}$ is made whenever a pair of samples $\hat{U}_{0,\ell}^{(\text{UB})}(Z^{(\ell,m)})$, $\hat{U}_{0,\ell-1}^{(\text{UB})}(Z^{(\ell,m)})$ lie on opposite sides of λ_{φ} , inflating the variances V_{ℓ} [\[24,](#page-26-4)[34\]](#page-26-6). Solutions to this problem are well studied within the MLMC literature [\[5,](#page-25-12) [6,](#page-25-13) [22,](#page-26-14) [25,](#page-26-15) [27,](#page-26-16) [48\]](#page-27-10).

In the present work, we consider adaptive sampling $[9,17,18,24,26]$ $[9,17,18,24,26]$ $[9,17,18,24,26]$ $[9,17,18,24,26]$ $[9,17,18,24,26]$ by refining approximations of $U_0(Z)$ which are relatively close to the discontinuity λ_{φ} , thereby reducing the chance that $\Delta_{\text{ind},\ell}$ is non-zero. Approximations of $U_0(Z)$ which lie relatively far from λ_{φ} do not require significant refinement, allowing us to control the computational cost. Specifically, we introduce a random refinement parameter η_{ℓ} , taking values in $\{0,\ldots,\ell\}$, to the discretisation $\widehat{U}_{0,\ell+\eta_{\ell}}^{(\text{UB})}(Z)$. For $k \in$ $\{0,\ldots,\ell\}$ and $1 < R < 2$, the value of η_{ℓ} is chosen according to the relation [\[24,](#page-26-4)[34\]](#page-26-6)

$$
\eta_{\ell} = k \iff \left\{ \left| \frac{\widehat{U}_{0,\ell+m}^{(\text{UB})}(Z) - \lambda_{\varphi}}{\sigma_{\ell+m}} \right| \le N_{0,\ell}^{(1-R)/R} N_{0,m}^{-1/R} \quad \text{for all } 0 \le m \le k-1
$$
\n
$$
\left| \frac{\widehat{U}_{0,\ell+k}^{(\text{UB})}(Z) - \lambda_{\varphi}}{\sigma_{\ell+k}} \right| > N_{0,\ell}^{(1-R)/R} N_{0,k}^{-1/R} \quad \text{if } k < \ell,
$$
\n
$$
(20)
$$

where $\{\sigma_\ell\}_{\ell \in \mathbb{N}}$ is as in Assumption [2.5.](#page-8-0) The value of η_ℓ increases according to uncertainty in the sign of $U_0(Z) - \lambda_{\varphi}$. The parameter $R \in (1,2)$ controls the strictness of the refinement procedure. Larger values of R lead to less strict refinements, with less cost but greater chance of error. When σ_{ℓ} is chosen to be the sample standard deviation as in [\(18\)](#page-8-3), the refinements outlined in [\(20\)](#page-9-1) can be thought of as conducting a one-sample t-test for the null hypothesis $U_0(Z) = \lambda_{\varphi}$. The procedure for generating adaptive Monte Carlo approximations for $U_0(Z)$ is outlined in Algorithm [1,](#page-10-0) which is based on [\[34,](#page-26-6) Algorithm 1]. An adaptive MLMC estimate of φ is constructed as in [\(14\)](#page-7-0) using the multilevel differences

$$
\Delta_{\text{ind},\ell} := \chi_{\widehat{U}_{0,\ell+\eta_{\ell}}^{(\text{UB})}(Z) > \lambda_{\varphi}} - \chi_{\widehat{U}_{0,\ell-1+\eta_{\ell-1}}^{(\text{UB})}(Z) > \lambda_{\varphi}},\tag{21}
$$

when $\ell > 0$. The key convergence properties of the adaptive hierarchical estimator is summarised in the following result, which is proved in Appendix [A.](#page-27-7)

Theorem 2.6. Let Assumption [2.5](#page-8-0) hold for some $p \geq 2$ and random variables σ_{ℓ} which can be sampled with order 2^{ℓ} cost. The MLMC estimator $\varphi_{\mathbf{M}_L,L}^{(\text{MLMC})}$ defined through [\(14\)](#page-7-0) and [\(21\)](#page-9-2), with adaptive refinements obtained through Algorithm [1](#page-10-0) with $1 < R < 2$ satisfies

$$
\text{Var}[\Delta_{\text{ind},\ell}] = \mathcal{O}(N_{0,\ell}^{-p/(p+2)}) \quad \text{and} \quad \mathsf{E}[N_{0,\ell+\eta_\ell}] = \mathcal{O}(N_{0,\ell}).
$$

Consequently, the cost of sampling $\varphi_{\mathbf{M}_L^{\star},L}^{(\text{MLMC})}$ within a specified root mean square error ε using optimal sample sizes defined by \mathbf{M}^{\star}_L can be characterized as follows:

• Under the additional assumption that $|\mathsf{E}[\Delta_{\mathrm{ind},\ell}]| = \mathcal{O}(N_{0,\ell}^{-1})$, the cost of obtaining a root mean square error ε satisfies $Cost(\varphi_{\mathbf{M}^*_L,L}^{(\mathrm{MLMC})}; \varepsilon) = \mathcal{O}(\varepsilon^{-2(p+3)/(p+2)})$.

• Otherwise, the cost can be bounded by $Cost(\varphi_{\mathbf{M}_{L},L}^{(\mathrm{MLMC})};\varepsilon) = \mathcal{O}(\varepsilon^{-2(p+2)/p}).$

Remark 2.7. Theorem [2.6](#page-9-0) implies the asymptotic cost of computing $\varphi_{\mathbf{M}_{L}^{*},L}^{(\text{MLMC})}$ to an error tolerance ε improves as $p \to \infty$ within Assumption [2.5.](#page-8-0) Proposition [2.3](#page-6-0) shows that the conditions hold with constants $\sigma_\ell \equiv \sigma$ for any $p < 23/11$. Alternatively, choosing σ_ℓ to be the sample standard deviation [\(18\)](#page-8-3) and under the condition [\(19\)](#page-8-4), the hypothesis of Theorem [2.6](#page-9-0) holds for any $p < \infty$. Taking p to be arbitrarily large, one obtains $\text{Cost}(\varphi_{\mathbf{M}_{L}^{*},L}^{(\text{MLMC})};\varepsilon) = \mathcal{O}(\varepsilon^{-2-\delta})$ for any $\delta > 0$.

2.4.1 Optimal Starting Level

The proof of Theorem [2.6](#page-9-0) uses asymptotic results on $\text{Var}[\Delta_{ind,\ell}]$ and $\text{E}[N_{0,\ell+\eta_\ell}]$ to bound the cost of the MLMC estimator [\(14\)](#page-7-0) with adaptive differences [\(21\)](#page-9-2). In practice, it is possible that the differences $\Delta_{\text{ind},\ell}$ exhibit pre-asymptotic behaviour at small levels $\ell \approx 0$. In particular, defining $\Delta_{\text{ind},\ell}^f := \chi_{\widehat{U}_{0,\ell+\eta_\ell}^{(\text{UB})}(Z) > \lambda_\varphi}$ to represent the finest approximation in [\(21\)](#page-9-2), it is important that the ratio

$$
R_{\ell_0} := \frac{\sqrt{\textsf{Var}[\, \Delta^f_{\text{ind},\ell_0} \,] N_{0,\ell_0}} + \sqrt{\textsf{Var}[\, \Delta_{\text{ind},\ell_0+1} \,] N_{0,\ell_0+1}}}{\sqrt{\textsf{Var}[\, \Delta^f_{\text{ind},\ell_0+1} \,] N_{\ell_0+1}}},
$$

is less than 1, otherwise using [\(15\)](#page-8-5) we can see it is more efficient to start the MLMC estimate from level $\ell_0 + 1$. In [\[24\]](#page-26-4), the authors suggest choosing ℓ_0 a-priori such that $R_{\ell_0} \leq 1$. Note that for the adaptive refinements defined by Algorithm [1,](#page-10-0) increasing ℓ_0 is not the equivalent to increasing the base sample size, $N_{0,0}$, unlike a non-adaptive algorithm.

2.4.2 Robust Error Estimation

Robust evaluation of the MLMC estimators [\(14\)](#page-7-0) requires accurate estimates of the error terms $V_{\ell} := \text{Var}[\Delta_{\text{ind},\ell}]$ and $E_{\ell} := |E[\Delta_{\text{ind},\ell}]|$ to approximate the optimal maximum level L and number of samples M_L . Since $|\Delta_{ind,\ell}|$ is a Bernoulli random variable, reliable Monte Carlo estimates for V_{ℓ} require $\mathcal{O}(\mathsf{Kurtosis}[\Delta_{\text{ind},\ell}]) = \mathcal{O}(V_{\ell}^{-1})$ samples, contradicting the fact that MLMC aims to reduce the number of samples required at fine levels and forcing a trade-off between efficiency and robustness of the estimator. In [\[48\]](#page-27-10), the authors approximate $\chi_{>0}$ using a sequence of smooth logistic sigmoid functions which depend on ℓ and are capable of reducing Kurtosis[$\Delta_{ind,\ell}$] by a constant factor. Alternatively, in [\[12\]](#page-25-16) the authors assume $V_{\ell} \approx a_0 2^{-\beta \ell}, E_{\ell} \approx a_1 2^{-\alpha \ell}$ for general MLMC problems and use Bayesian estimates for the parameters a_0, a_1, α, β based on the observed samples at levels $\ell = 0, \ldots, L$. Alternatively, in [\[18\]](#page-25-15), the authors consider a Bayesian approach which aims to estimate the probability that $\Delta_{ind,\ell}$ is non-zero. An extension of this Bayesian approach is outlined in Appendix [B.](#page-32-0)

3 Model for the Credit Valuation Adjustment

We now apply the hierarchical estimator [\(14\)](#page-7-0) constructed in Section [2](#page-2-0) to the problem of computing the probability of large loss due to fluctuations in the CVA [\(1a\)](#page-0-0). Consider a financial market with d assets $(S_t)_{0\leq t\leq T}$ with dynamics modelled by the Brownian SDE [\(2\)](#page-1-0) under a physical measure P and which are adapted to the filtration \mathcal{F} . The discounting factor is modelled by the dynamics $dB_t = r_t B_t dt$, with $B_0 = 1$, where the process $r := (r_t)_{t\geq 0}$ is adapted to the filtration \mathcal{F} , with r_t taking values in \mathbb{R}_+ for all $t \geq 0$. To simplify the presentation, it is assumed that B_t can be sampled directly. However, the slightly more general case where B_t requires approximate sampling and r_t follows a Brownian SDE can be tackled naturally by setting $\tilde{S}_t := (S_t, r_t, B_t)$ for all $t \geq 0$. We assume the existence of a risk-neutral measure, Q, under which the assets evolve under the risk-neutral SDE

$$
dS_t = r_t S_t dt + \varsigma(t, S_t) dW_t^Q,
$$
\n(22)

where $W^{\mathsf{Q}} := (W_t^{\mathsf{Q}})_{t \geq 0}$ is a d'-dimensional Q-Brownian motion.

Consider a contract between a bank and a counterparty, which has a single payoff occurring at maturity T, given by $\pi(S_T)$ for some $\pi : \mathbb{R}^d \to \mathbb{R}$. We emphasise here that the restriction of the portfolio to a single payoff and counterparty is made solely with the intention of simplifying the following discussion and apply equally to any number of counterparties, see Section [4.4.](#page-20-0) Counterparty credit risk is modelled through the stopping time τ with respect to the expanded filtration G that incorporates additional credit risk factors. To account for potential counterparty default, the value of the banks position at time t is adjusted by the amount CVA_t defined in [\(1b\)](#page-0-2), with discounted payoffs $\pi_{\tau,T} = B_{\tau} B_T^{-1} \pi(S_T)$, given S_{τ} , for $\tau \leq T$. Letting $V_t^{\pi} = B_{\tau} \mathsf{E}^{\mathsf{Q}}[B_T^{-1} \pi(S_T) | S_t]$ represent the risk-neutral value of the portfolio at time $t \in [0, T]$, the adjusted value of the banks position, accounting for counterparty credit risk, is

$$
V_t^{(\text{CVA}, \pi)} = V_t^{\pi} \chi_{\tau \ge t} - \text{CVA}_t. \tag{23}
$$

For the purposes of this work, we assume that the bank itself carries no risk of default. In settings where the bank carries a non-negligible default risk, a debit valuation adjustment (DVA) should also be considered in the adjusted valuation [\(23\)](#page-11-1) [\[33\]](#page-26-1), which is defined in the same manner as CVA but with the roles of the bank and counterparty reversed and therefore the methods discussed herein are equally applicable to the DVA.

Random losses for the adjusted valuation occurring by the short risk horizon $H \ll T$ can be expressed as

$$
\Lambda_H^{(\text{CVA}, \pi)} \coloneqq \Lambda_{\tau < H}^{(\text{CVA}, \pi)} + \Lambda_{H \leq \tau}^{(\text{CVA}, \pi)},
$$

where $\Lambda_{\tau \leq H}^{(\text{CVA}, \pi)}$ and $\Lambda_{H \leq \tau}^{(\text{CVA}, \pi)}$ reflect separately the nature of losses depending on whether τ occurs before or after H. When τ falls before H, losses in the valuation are given by the difference in (adjusted) valuations at time 0 and time τ . If default occurs before H, the losses can be expressed as

$$
\Lambda_{\tau < H}^{\rm (CVA, \pi)} \coloneqq \chi_{\tau < H} \Big(V_0^{\rm (CVA, \pi)} - B_\tau^{-1} V_\tau^{\rm (CVA, \pi)} \Big),
$$

whereas random losses in the event $\tau \geq H$ take the form

$$
\Lambda_{H \leq \tau}^{(\text{CVA}, \pi)} := \chi_{H \leq \tau} \left(V_0^{(\text{CVA}, \pi)} - B_H^{-1} V_H^{(\text{CVA}, \pi)} \right)
$$
\n
$$
= \underbrace{\chi_{H \leq \tau} \left(V_0^{\pi} - B_H^{-1} V_H^{\pi} \right)}_{=: \Lambda_{\text{market}}^{(\text{CVA}, \pi)}} + \underbrace{\chi_{H \leq \tau} \left(B_H^{-1} \text{CVA}_H - \text{CVA}_0 \right)}_{=: \Lambda_{\text{credit}}^{(\text{CVA}, \pi)}}.
$$

In the above expression, the component $\Lambda_{\text{market}}^{(\text{CVA}, \pi)}$ captures the market risk associated with fluctuations in the risk-neutral valuation of the banks position due to random fluctuations in the assets by the risk horizon H. Monte Carlo and hierarchical estimators for market risk measures of such losses have been widely studied [\[9,](#page-25-6)[24,](#page-26-4)[26,](#page-26-5)[29,](#page-26-17)[31\]](#page-26-3). Credit risk factors are captured by terms $\Lambda_{\rm credit}^{({\rm CVA}, \pi)}$ and $\Lambda_{\tau < H}^{(\text{CVA}, \pi)}$, each considering a subtly different aspect of counterparty credit risk:

- $\Lambda_{\text{credit}}^{(\text{CVA}, \pi)}$ captures the risk posed by fluctuations in the credit worthiness of the counterparty impacting the valuation adjustment over the interval $[0, H]$ and thereby reducing the adjusted value of the banks position.
- $\Lambda_{\tau \leq H}^{(\text{CVA}, \pi)}$ captures the risk posed by imminent counterparty default leading to losses before the risk horizon H.

In practice, under the physical measure which captures the underlying risk factors, $P[\tau < H] \approx 0$ since $H \ll T$ and the physical probability that the counterparty will default over a small period of time is small for counterparties with reasonable credit worthiness. On the other hand, default probabilities are often inflated under the risk-neutral measure when compared to the physical measure [\[33,](#page-26-1) Chapter 12], [\[37\]](#page-26-18). It is therefore more likely that small fluctuations in the credit worthiness of the counterparty will lead to significant losses in the adjusted valuation of the portfolio through the CVA, as captured by $\Lambda_{\text{credit}}^{(\text{CVA}, \pi)}$. Consequently, in what follows we assume that $P[\tau < H] = 0$, and focus on the losses $\Lambda_{\text{credit}}^{(\text{CVA}, \pi)}$ which are then equal to $B_H^{-1} CVA_H - CVA_0$ almost surely under P. Our goal is then to construct an efficient estimator for the probability φ in [\(1a\)](#page-0-0), that the losses $\Lambda_{\text{credit}}^{(\text{CVA}, \pi)}$ exceed a given threshold λ_{φ} at the risk horizon $H \ll T$.

3.1 Model for the Default Times

Following the approach in [\[8,](#page-25-0) Section 3.3] and [\[33,](#page-26-1) Chapter 12], we assume an intensity model for the default times under the risk-neutral measure. Specifically, for a positive-valued process $h = (h_t)_{t>0}$, adapted to \mathcal{G} , representing a hazard rate, assume that under the risk-neutral measure $Q[\tau \leq t] = 1 - \exp(-\int_{s=0}^{t} h_s ds)$ for all $t \geq 0$. Then, the instantaneous probability of default at time t is h_t dt. A common model for the hazard rates is $h_t = LGD^{-1}c_t$, which depends on the loss given default and the counterparties credit spread, c_t , which is a positive-valued, \mathcal{G}_t measurable, random variable representing the difference in return between a counterparty bond and the risk-neutral bond B_t [\[33,](#page-26-1) Section 12.2.3]. A large credit spreads implies the counterparty bond has high associated risk, which is a sign of poor credit worthiness and large default probabilities. Following [\[43\]](#page-27-0), we assume that the credit spread is constant under the risk-neutral measure Q. Then, for any fixed $v \geq 0$, we have:

$$
Q[v \le \tau \le t + v | \tau \ge v, c_v] = 1 - \exp\left(-\frac{c_v t}{\text{LGD}}\right).
$$

In particular, conditioned on $\tau \geq v$, the random variable $(\tau - v)$ follows an Exponential (c_v/LGD) distribution.

As discussed above, we assume that $P[\tau < H] = 0$. Following [\[43\]](#page-27-0), to model random fluctuations in the creditworthiness of the counterparty over the short risk horizon, we assume that the credit spread follows a geometric Brownian motion under P. In particular, assume

$$
\mathrm{d}c_t = \sigma^{\text{(cs)}} c_t \mathrm{d}W_t^{\text{(cs,P)}},\tag{24}
$$

where $c_0 > 0$ can be inferred from market data, $\sigma^{(cs)}$ is a constant representing the volatility of the credit spread and $\{W_t^{(cs,P)}\}_{0 \le t \le T}$ is a P-Brownian motion. Large values of the product $\sigma^{(cs)}c_0$ imply that there is high initial volatility in the credit spread. This can lead to significant worsening of the credit worthiness of the counterparty by the risk horizon H , causing large losses due to fluctuations in the CVA. In [\[43,](#page-27-0) Table A], the product $\sigma^{(cs)}c_0$ is directly related to the credit rating of the counterparty to compute the standardised CVA formula.

To summarise, the above model for default times relies on two assumptions: Under the physical measure, the physical probability of counterparty default before H is negligible, but the credit spread fluctuates according to a geometric Brownian motion. As a result, risks posed by changing credit worthiness of the counterparty are captured by the dynamics (24) of c_t under P. On the other hand, under the risk-neutral measure, conditioned on the credit spread c_v at time $v \geq 0$ and the event $\tau \geq v$, the credit spread is fixed and the random variable $(\tau - v)$ follows an Exponential (c_v/LGD) distribution. Important to what follows, this model allows us to sample random default times τ under the risk-neutral measure, conditioned on a given risk scenario, exactly. More general riskneutral models for the default time are discussed in [\[8,](#page-25-0) [33\]](#page-26-1), which do not rely on constant credit spreads under Q. However, such models can lead to difficulties in sampling the default times which are beyond the scope of algorithms presented in this paper, and are left to future work. Nevertheless, as remarked in [\[43\]](#page-27-0), the assumption of constant credit spreads is practical since the information used to model credit spreads is often sparsely distributed in time.

3.2 Formulation as a Nested Expectation

Under the assumption that no default occurs under the physical measure and for the default time model in Section [3.1,](#page-12-0) the (discounted) CVA at time t is equivalent to

$$
\frac{\text{CVA}_{t}}{B_{t}} = \mathsf{E}^{\mathsf{Q}} \Big[\, \chi_{\tau_{c_{t}} \leq T} \, \text{LGD} \, \text{max} \Big\{ \mathsf{E}^{\mathsf{Q}} \Big[\, B_{T}^{-1} \pi(S_{T}) \, \Big| \, S_{\tau_{c_{t}}}^{t, S_{t}^{p}}, \tau_{c_{t}} \, \Big], 0 \Big\} \, \Big| \, S_{t}^{p}, c_{t} \, \Big],
$$

where we introduce the following notation:

- S_t^{P} is superscripted by P to indicate it is driven by the P-Brownian motion W through the dynamics [\(2\)](#page-1-0).
- τ_{c_t} is indexed to highlight the dependency of the default times on the credit spread c_t . In the setup of Section [3.1,](#page-12-0) $\tau_{c_t} - t$ is an Exponential(c_t/LGD) random variable.
- For $t \le u \le T$ and $x \in \mathbb{R}^d$, $S_u^{t,x}$ represents the asset value S_u given $S_t^{t,x} = x$.

In particular, defining $f(u) = \max\{u, 0\}$, and $U_1(s, \tau) = \text{LGD}\, \chi_{\tau \leq T} \mathsf{E}^{\mathsf{Q}}[B_T^{-1} \pi(S_T) | S_\tau = s, \tau]$, we have

$$
\frac{\text{CVA}_{t}}{B_{t}} = \mathsf{E}^{\mathsf{Q}} \Big[f\Big(U_1\Big(S_{\tau_{c_t}}^{t, S_t^{\mathsf{P}}}, \tau_{c_t}\Big)\Big) \Big| S_t^{\mathsf{P}}, c_t \Big],
$$

such that the problem [\(1a\)](#page-0-0) becomes

$$
\varphi = \mathsf{P}\big[U_0(S_H^{\mathsf{P}}, c_H) > \lambda_\varphi \big],\tag{25a}
$$

$$
U_0(s,c) \coloneqq \mathsf{E}^{\mathsf{Q}} \Big[\, f\Big(U_1\Big(S_{\tau_{c_H}}^{H,S_H^{\mathsf{P}}},\tau_{c_H}\Big) \Big) \, \Big| \, S_H^{\mathsf{P}} = s, c_H = c \Big] - \text{CVA}_0,\tag{25b}
$$

where $S_{\tau_{c_H}}^{H,S_H^{\mathcal{P}}}$ represents the risky asset values at default, first simulated under the physical measure up to the horizon H, and then simulated up to the default time τ_{c_H} using the risk-neutral dynamics [\(22\)](#page-11-2) driven by the Q-Brownian motion W^{Q} . In contrast, $S_{\tau_{c_0}}^{0,S_0}$ is simulated entirely under the measure Q up to the default time τ_{c_0} , which depends only on the initial credit spread c_0 . This problem can be expressed in the general form [\(3\)](#page-1-1), where

$$
Z = (S_H^{\mathsf{P}}, c_H) \text{ is } \mathcal{G}_H\text{-measurable, taking values in } \mathbb{R}^{d+1}
$$

\n
$$
Y = \left(S_{\tau_{c_H}}^{H, S_H^{\mathsf{P}}}, \tau_{c_H}\right) \text{ is } \mathcal{G}_{\tau_{c_H}}\text{-measurable, taking values in } \mathbb{R}^{d+1}
$$

\n
$$
X = \text{LGD}\,\chi_{\tau_{c_H} \leq T} B_T^{-1} \pi(S_T^{H, S_H^{\mathsf{P}}}) \text{ is } \mathcal{G}_T\text{-measurable, taking values in } \mathbb{R}.
$$
 (26)

The variable X contains a discontinuous function of the default time τ_{c_H} . In the current setting, this has no influence on the complexity of the MLMC estimation of φ (equivalently λ_{φ}) since the current framework samples the default times exactly. More generally, this discontinuity can be removed by using importance sampling techniques, as we will discuss in Section [4.2,](#page-18-0) to ensure the sampled default times always fall before T . Within the computation [\(25\)](#page-14-1), CVA₀ is independent of the generated risk scenario S_H^{P} and c_H and can be approximated a-priori outside of the nested expectation over the risk factors. In Section [4,](#page-15-0) we show that by including $CVA₀$ as part of the inner computation we can leverage dependencies between S_t^{0, S_0} and $S_t^{H, S_H^{\mathbb{P}}}$ to reduce the variance of the estimator.

3.3 Collateralization

A more realistic model for losses arising due to CVA should incorporate credit risk mitigation factors, with the aim of lowering the risk faced by institutions due to counterparty credit risk. In recent years, the initial and variational margins have become essential in reducing the exposure of the bank to its counterparties [\[32,](#page-26-0) [33\]](#page-26-1). Assuming credit risk arises only through the counterparty, the variational margin requires the counterparty to post funds to a segregated account (free from default-risk) at a sequence of discrete time-points to cover the cost of their exposure to the bank. The initial margin is then based on a (conditional) Value-at-Risk formula on the exposure evaluated at each posting time of the variational margin to account for fluctuations which arise between posting times. Computing variational adjustments therefore requires evaluating additional nested expectations defining the exposure faced to the counterparty at the posting time immediately prior to default. On the other hand, initial margin computation requires a nested value-at-risk calculation, given the market state at the posting time prior to counterparty default. Defined in full generality, inclusion of nested risk measures will require, for example, nested stochastic approximation schemes to approximate the initial margin. Such methods fall beyond the scope of this paper and are left as an interesting avenue for future work. In the particular case where the assets S evolve according to a geometric Brownian motion, the initial margin can be related to the solution of a McKean backward stochastic differential equation [\[1,](#page-25-17) [35\]](#page-26-19). In [\[7\]](#page-25-9), the authors use this approach to codmpute the initial margin by approximating a pair of nested expectations using an antithetic MLMC estimator as discussed in Section [2.2.](#page-3-0) Defined in this manner, the initial margin computation will add an extra layer of nested expectations when computing the CVA. The additional layer of nested expectations can be handled by extending the estimator in Section [2.2](#page-3-0) to handle an additional nested expectation, which is possible using MLMC techniques for repeatedly nested expectations as discussed in [\[46,](#page-27-8) [47,](#page-27-1) [49\]](#page-27-2).

4 Approximation of the Capital Charge

With the aim of reducing the variance of $U_0(S_H^{\mathsf{P}}, c_H)$ in [\(25\)](#page-14-1), we include the computation of CVA₀ within the conditional average over S_H^{P}, c_H through

$$
U_0(S_H^{\mathbf{P}}, c_H) = \mathsf{E}\Big[\,\Phi\Big(U_1\Big(S_{\tau_{c_H}}^{H,S_H^{\mathbf{P}}}, \tau_{c_H}\Big), U_1\Big(S_{\tau_{c_0}}^{0,S_0}, \tau_{c_0}\Big)\Big)\,\Big|\, c_H, S_H^{\mathbf{P}}\,\Big],
$$

where we define the function $\Phi(u, v) := f(u) - f(v)$ and, by an abuse of notation, refer to the random variable

$$
\Phi := \Phi \Big(U_1 \Big(S_{\tau_{c_H}}^{H, S_H^{\mathcal{P}}}, \tau_{c_H} \Big), U_1 \Big(S_{\tau_{c_0}}^{0, S_0}, \tau_{c_0} \Big) \Big). \tag{27}
$$

The assets S^{H,S_H^P} and S^{0,S_0} can be approximated under the risk-neutral measure Q using the Milstein scheme [\[40\]](#page-26-8) and the methods of Section [2](#page-2-0) can be applied directly, using the linearity of expectation to handle the difference between both averages U_1 within [\(27\)](#page-15-1), to compute φ . There are several aspects of the problem in this form which impede efficient multilevel estimation:

- The variance of the inner samples Φ within [\(25\)](#page-14-1), is inflated by the two different sources of risk:
	- (Credit Risk). The default times τ_{c_H} and τ_{c_0} occur with different rates and it is difficult to correlate a pair of samples under each rate effectively. It is unlikely that independent samples of τ_{c_H} and τ_{c_0} will both occur before maturity T.
	- (Market Risk). The processes S^{H,S_H^P} and S^{0,S_0} evolve under the measures P and Q, respectively, on the interval $[0, H]$.
- Under the risk-neutral measure, the events $\tau_{c_H} \leq T$ and $\tau_{c_0} \leq T$ are rare, leading to instability within the MLMC estimate of φ .

In Section [4.1,](#page-16-0) several variance reduction techniques are used to replace Φ with a variable Φ^{III} satisfying $\mathsf{E}[\Phi^{\text{III}} | S_H^{\mathsf{P}}, c_H] = \mathsf{E}[\Phi | S_H^{\mathsf{P}}, c_H],$ but with $\mathsf{Var}[\Phi^{\text{III}} | S_H^{\mathsf{P}}, c_H] = \mathcal{O}(H)$, constituting a significant reduction in variance. In Section [4.2,](#page-18-0) importance sampling techniques are used to sample default times from the conditional distribution of the counterparty defaulting before T , resolving instability issues from rare event sampling.

4.1 Variance Reduction

We begin by using a change of measure to remove the need to sample τ_{c_H} under a different rate from τ_{c_0} . For the exponentially distributed default times in Section [3.1](#page-12-0) and any $F : \mathbb{R}_+ \to \mathbb{R}$, we have

$$
\begin{split} \mathsf{E}^{\mathsf{Q}}[F(\tau_{c_H})|c_H] &= \int_H^\infty F(t) \frac{c_H}{\mathrm{LGD}} e^{-c_H(t-H)/\mathrm{LGD}} \mathrm{d}t \\ &= \frac{c_H}{c_0} e^{c_H H/\mathrm{LGD}} \int_0^\infty \chi_{t \geq H} F(t) \frac{c_0}{\mathrm{LGD}} e^{-(c_H - c_0)t/\mathrm{LGD}} e^{-c_0 t/\mathrm{LGD}} \mathrm{d}t \\ &= \mathsf{E}^{\mathsf{Q}} \Big[\chi_{\tau_{c_0} \geq H} g_{\tau_{c_0}}(c_H) e^{c_H H/\mathrm{LGD}} F(\tau_{c_0}) \Big| c_H \Big], \end{split}
$$

where we define $g_t(x) := c_0^{-1} x e^{-(x-c_0)t/\text{LGD}}$. In particular,

$$
\begin{split} &\mathsf{E}^\mathsf{Q}\Big[\,f\Big(U_1\Big(S^{H,S^{\mathsf{P}}}_{\tau_{c_H}} , \tau_{c_H}\Big)\Big)\,\Big|\,S^{\mathsf{P}}_H , c_H\,\Big] \\ &=\mathsf{E}^\mathsf{Q}\Big[\,\chi_{\tau_{c_0}\ge H} g_{\tau_{c_0}}(c_H) e^{c_H H/\,\mathrm{LGD}}f\Big(U_1\Big(S^{H,S^{\mathsf{P}}}_{\tau_{c_0}} , \tau_{c_0}\Big)\Big)\,\Big|\,S^{\mathsf{P}}_H , c_H\,\Big], \end{split}
$$

where the change of measure moves the dependency on c_H to the factor $g_{\tau_{c_0}}(c_H)e^{c_H H/LGD}$, and simulates default time using τ_{c_0} opposed to τ_{c_H} , eliminating the need to sample τ_{c_H} . Consequently, we henceforth express $\tau_{c_0} \equiv \tau$. The above discussion then motivates replacing Φ as in [\(27\)](#page-15-1) with

$$
\Phi^{\mathcal{I}} \coloneqq \chi_{\tau \geq H} \left(g_{\tau}(c_H) e^{c_H H / \text{LGD}} \underbrace{\left(f \left(U_1 \left(S_{\tau}^{H, S_H^{\mathcal{P}}}, \tau \right) \right) - f \left(U_1 \left(S_{\tau}^{0, S_0}, \tau \right) \right) \right)}_{\text{Market risk}} + \underbrace{\left(g_{\tau}(c_H) e^{c_H H / \text{LGD}} - 1 \right) f \left(U_1 \left(S_{\tau}^{0, S_0}, \tau \right) \right) \right)},
$$
\n(28)

whenever $\tau \geq H$. By construction $\mathsf{E}^{\mathsf{Q}}[\Phi^{\mathrm{I}} | S_H^{\mathsf{P}}, c_H] = \mathsf{E}^{\mathsf{Q}}[\chi_{\tau_{c_0} \geq H} \Phi | S_H^{\mathsf{P}}, c_H]$. The component of Φ reflecting default before H , under Q , is

$$
\chi_{\tau < H} \Phi = -\chi_{\tau < H} f(U_1(S_{\tau}^{0, S_0}, \tau)).
$$

Since this is independent of the risk factors at the horizon H , we can cancel the bias induced by considering only $\tau \geq H$ in Φ^{I} by setting

$$
\lambda_{\varphi}^{\mathrm{I}} \coloneqq \lambda_{\varphi} + \mathsf{E}^{\mathsf{Q}} \big[\, \chi_{\tau < H} f\big(U_1 \big(S_{\tau}^{0, S_0}, \tau \big) \big) \, \big],
$$

which be pre-computed offline at insignificant cost compared with the cost of computing φ . By the change of measure from τ_{c_H} to τ above, it follows that $\mathsf{E}^{\mathsf{Q}}[\Phi - \lambda_{\varphi} | S_H^{\mathsf{P}}, c_H] = \mathsf{E}^{\mathsf{Q}}[\Phi^{\mathrm{I}} - \lambda_{\varphi}^{\mathrm{I}} | S_H^{\mathsf{P}}, c_H]$. We consider variance reduction for the credit and market risk factors in turn.

4.1.1 Credit Risk

In [\[43\]](#page-27-0), it is shown that the standardised CVA capital charge can be derived by linearising the term $g_{\tau}(c_H)e^{c_H}$ as a function of c_H . We use this approach here to construct a control variate that reduces the variance of Φ . In particular, since $g_{\tau}(c_0) = 1$, we have

$$
g_{\tau}(c_H)e^{c_HH/LGD} - 1 = g_{\tau}(c_H)(1 + \mathcal{O}(H)) - g_{\tau}(c_0)
$$

$$
= \underbrace{(c_H - c_0) \left(\frac{1}{c_0} - \frac{\tau}{LGD}\right)}_{\mathcal{O}(\sqrt{H})} + \mathcal{O}(H), \tag{29}
$$

where the difference $c_H - c_0$ is of order \sqrt{H} since the process $(c_t)_{t\geq 0}$ evolves under P according to the dynamics [\(24\)](#page-13-1) with an underlying P-Brownian motion. To remove this leading order term, we set

$$
\Phi^{II} := \Phi^I - \chi_{\tau \ge H}(c_H - c_0) \Psi_\tau(S^{0, S_0}_\tau),
$$

where we introduce the control variate

$$
\Psi_{\tau}(s) := \left(\frac{1}{c_0} - \frac{\tau}{\text{LGD}}\right) f(U_1(s, \tau)).
$$

To ensure the control variate introduces no bias, we set

$$
\lambda_{\varphi}^{\text{II}} \coloneqq \lambda_{\varphi}^{\text{I}} - (c_H - c_0) \mathsf{E}^{\mathsf{Q}} \big[\Psi_{\tau} (S_{\tau}^{0, S_0}) \big],
$$

so that again $\mathsf{E}^{\mathsf{Q}}[\Phi^{\mathrm{II}} - \lambda_{\varphi}^{\mathrm{II}} | S_H^{\mathsf{P}}, c_H] = \mathsf{E}^{\mathsf{Q}}[\Phi - \lambda_{\varphi} | S_H^{\mathsf{P}}, c_H]$. Since $\Psi_\tau(S_\tau^{0, S_0})$ does not depend on the risk factors, $E^{\mathsf{Q}}[\Psi_{\tau}(S_{\tau}^{0,S_0})]$ can be computed offline at negligible cost. Consequently, $\lambda_{\varphi}^{\text{II}}$ may be computed with bounded cost for each risk scenario c_H .

4.1.2 Market Risk

The market risk component of Φ is primarily composed of the term

$$
f\left(U_1\left(S_{\tau}^{H,S_H^{\mathcal{P}}},\tau\right)\right)-f\left(U_1\left(S_{\tau}^{0,S_0},\tau\right)\right)=\bar{f}\left(S_{\tau}^{H,S_H^{\mathcal{P}}}\right)-\bar{f}\left(S_{\tau}^{0,S_0}\right),
$$

where we introduce the functions $\bar{f} : \mathbb{R}^d \to \mathbb{R}$, dropping the dependency of \bar{f} on τ , to simplify notation in what follows. As with the credit risk, we focus on the case where $H \leq \tau \leq T$. Following the techniques in [\[26,](#page-26-5) Section 3.2], this term can be decomposed into

$$
\bar{f}(S_{\tau}^{H,S_H^{\mathcal{P}}}) - \bar{f}(S_{\tau}^{0,S_0}) = (\bar{f}(S_{\tau}^{H,S_H^{\mathcal{P}}}) - \bar{f}(S_{\tau}^{0,S_H^{\mathcal{P}}})) + (\bar{f}(S_{\tau}^{0,S_H^{\mathcal{P}}}) - \bar{f}(S_{\tau}^{0,S_0}))
$$
\n
$$
= (S_{\tau}^{H,S_H^{\mathcal{P}}} - S_{\tau}^{0,S_H^{\mathcal{P}}})^{\mathsf{T}} \nabla \bar{f}(S_{\tau}^{0,S_H^{\mathcal{P}}}) + (S_H^{\mathcal{P}} - S_0)^{\mathsf{T}} \nabla_{S_0} \bar{f}(S_{\tau}^{0,S_0}) + \mathcal{O}(H),
$$

using a Taylor expansion in the final step. Both of the terms $S_{\tau}^{H,S_H^P} - S_{\tau}^{0,S_H^P}$ and $S_H^P - S_0$ are of order \sqrt{H} since they involve simulating the assets under different Brownian motions along the interval $[0, H]$. The former term can eliminated by utilising an antithetic pair of risk-neutral paths $S^{0, S_0, \pm}$, each of which follows the Q-Brownian path $\{\pm W_t^Q\}_{0 \leq t \leq H}$ up to the risk horizon. After the risk horizon, each of the processes S^{H,S_H^P} and $S^{0,S_0,\pm}$ should follow the same Brownian path ${W_t^{\mathbf{Q}}}_{H \leq t \leq \tau}$ up to the default time. It then follows for the antithetic variate [\[26,](#page-26-5) [29\]](#page-26-17)

$$
\begin{split} &\bar{f}\left(S_{\tau}^{H,S_{H}^{0}}\right)-\frac{1}{2}\big(\bar{f}\big(S_{\tau}^{0,S_{0},+}\big)+\bar{f}\big(S_{\tau}^{0,S_{0},-}\big)\big) \\ &=\frac{1}{2}\big(S_{H}^{0}-S_{0}\big)^{\intercal}\big(\nabla_{S_{0}}\bar{f}\big(S_{\tau}^{0,S_{0},+}\big)+\nabla_{S_{0}}\bar{f}\big(S_{\tau}^{0,S_{0},-}\big)\big)+\mathcal{O}(H). \end{split}
$$

By subtracting the term

$$
\frac{1}{2}\left(S_H^{\mathsf{P}}-S_0\right)^{\mathsf{T}}\left(\nabla_{S_0}\bar{f}\left(S_\tau^{0,S_0,+}\right)+\nabla_{S_0}\bar{f}\left(S_\tau^{0,S_0,-}\right)\right),\,
$$

one obtains samples which are of $\mathcal{O}(H)$, constituting a significant improvement to the variance. The components $\nabla_{S_0} \cdot \bar{f}$ are referred to in financial settings as the delta control variate [\[29\]](#page-26-17).

Following the above approach and using the decomposition in [\(28\)](#page-16-1) motivates the term

$$
\Phi^{\text{III}} := \chi_{\tau \ge H} \bigg(g_{\tau}(c_H) e^{c_H H / \text{LGD}} f \bigg(U_1 \bigg(S_{\tau}^{H, S_H^{\mathbf{P}}}, \tau \bigg) \bigg) \n- \frac{1}{2} \big(f \big(U_1 \big(S_{\tau}^{0, S_0, +}, \tau \big) \big) + f \big(U_1 \big(S_{\tau}^{0, S_0, -}, \tau \big) \big) \big) \n- \frac{1}{2} (c_H - c_0) \big(\Psi_{\tau} (S_{\tau}^{0, S_0, +}) + \Psi_{\tau} (S_{\tau}^{0, S_0, -}) \big) \n- \frac{1}{2} \big(S_H^{H, S_H^{\mathbf{P}}} - S_0 \big)^{\mathsf{T}} \big(\nabla_{S_0} f \big(U_1 \big(S_{\tau}^{0, S_0, +}, \tau \big) \big) + \nabla_{S_0} f \big(U_1 \big(S_{\tau}^{0, S_0, -}, \tau \big) \big) \big) \bigg),
$$

As before, to compensate for the bias induced by the delta control variate, we replace $\lambda_{\varphi}^{\text{II}}$ with

$$
\lambda_{\varphi}^{\text{III}} := \lambda_{\varphi}^{\text{II}} - \left(S_{H}^{\text{P}} - S_0\right)^{\intercal} \mathsf{E}^{\text{Q}} \big[\nabla_{S_0} f\big(U_1\big(S_{\tau}^{0, S_0}, \tau\big)\big)\big]
$$

so that $\mathsf{E}^{\mathsf{Q}}[\Phi^{\text{III}}] S_H^{\mathsf{P}}, c_H] - \lambda_{\varphi}^{\text{III}} = \mathsf{E}^{\mathsf{Q}}[\Phi] S_H^{\mathsf{P}}, c_H] - \lambda_{\varphi}$. Since all of the expectations appearing in the definition of $\lambda_{\varphi}^{\text{III}}$ are independent of S_H^{P} and c_H , they may be pre-computed offline at insignificant cost.

4.2 Importance Sampling

Observe that the CVA loss, Φ^{III} , is non-zero only when $H \leq \tau \leq T$. For credible counterparties this is a rare event, making it costly to simulate a single instance of default. To address this issue note that, defining $\tilde{p} := \mathsf{P}[H \leq \tau \leq T]$, we have

$$
\mathsf{E}^{\mathsf{Q}}\big[\,\Phi^{\mathrm{III}}\,\big|\,S_{H}^{\mathsf{P}},c_{H}\,\big]-\lambda_{\varphi}^{\mathrm{III}}=\mathsf{E}^{\mathsf{Q}}\bigg[\,\Phi^{\mathrm{III}}\tilde{p}\bigg|S_{H}^{\mathsf{P}},c_{H},H\leq\tau\leq T\,\bigg]-\lambda_{\varphi}^{\mathrm{III}}.\tag{30}
$$

Under the model for default times in Section [3.1,](#page-12-0) τ is an Exponential(c_0 /LGD) random variable under the risk-neutral measure Q. Hence, for $H \le t \le T$, we have

$$
Q[\tau \le t | H \le \tau \le T] = \frac{Q[H \le \tau \le t]}{Q[H \le \tau \le T]} = \frac{e^{-c_0 H / \text{LGD}} - e^{-c_0 t / \text{LGD}}}{e^{-c_0 H / \text{LGD}} - e^{-c_0 T / \text{LGD}}}.
$$

By inverting this equation for t , we can sample directly from the conditional distribution using inverse transform sampling. This reduces the kurtosis of samples from the CVA loss by ensuring default always occurs.

The above approach relies explicitly on the model for default times described in Section [3.1.](#page-12-0) More complex models of the default time will require more advanced importance sampling techniques to achieve a similar result.

4.3 Hierarchical MLMC Estimates of the Loss

To summarize, combining the techniques above, we have expressed the problem [\(25\)](#page-14-1) in the form

$$
\varphi = \mathsf{P}\big[\,\mathsf{E}^\mathsf{Q}\big[\,\Phi\,\big|\,S_H^\mathsf{P}, c_H\,\big] > \lambda_\varphi\,\big] \\ = \mathsf{P}\big[\,\mathsf{E}^\mathsf{Q}\big[\,\Phi^\mathrm{III}\tilde{p} - \lambda_\varphi^\mathrm{III}\,\big|\,S_H^\mathsf{P}, c_H, H \leq \tau \leq T\,\big] > 0\,\big],
$$

where Φ^{III} depends only on the default times $\tau = \tau_{c_0}$ and has significantly reduced variance compared with Φ as in [\(27\)](#page-15-1). Furthermore, by sampling from the conditional distribution $H \leq \tau \leq T$ we are able to sample directly from the conditional distribution of $\tau \in [H, T]$. To counteract the bias induced replacing Φ with Φ^{III} , we set

$$
\lambda_{\varphi}^{\text{III}} = \lambda_{\varphi} + \mathsf{E}^{\mathsf{Q}} \big[\chi_{\tau < H} f(U_1(S_{\tau}^{0, S_0}, \tau)) \big] - (c_H - c_0) \mathsf{E}^{\mathsf{Q}} \big[\Psi_{\tau}(S_{\tau}^{0, S_0}) \big] - (S_H^{\mathsf{P}} - S_0)^{\mathsf{T}} \mathsf{E}^{\mathsf{Q}} \big[\nabla_{S_0} f(U_1(S_{\tau}^{0, S_0}, \tau)) \big],
$$

where each expectation can be pre-computed offline at insignificant cost using, for example, (multilevel) Monte Carlo sampling, since they do not depend on the generated risk variables. Since the variable Φ^{III} is a functions of the nested expectations $U_1(s,\tau)$ for $s \in \{S_{\tau}^{0,S_0,\pm}, S_{\tau}^{H,S_H^{\text{P}}}\},\$ the techniques of Section [2.3](#page-5-0) can be used to construct antithetic correction terms $\Delta_{\ell}^{(\text{ant})} \Phi^{\text{III}}$ satisfying

$$
\mathsf{E}^{\mathsf{Q}}\!\left[\left.\Phi^{\mathrm{III}}\tilde{p} - \lambda^\mathrm{III}_\varphi\,\right|S_H^{\mathsf{P}}, c_H,H\leq \tau\leq T \right] = \mathsf{E}^{\mathsf{Q}}\!\left[\left.\sum_{\ell=0}^\infty\Delta_\ell^{(\mathrm{ant})}\Phi^{\mathrm{III}}\,\right|S_H^{\mathsf{P}}, c_H,H\leq \tau\leq T \right].
$$

The antithetic differences $\Delta_{\ell}^{(ant)}\Phi^{III}$ combine nested (unbiased) MLMC estimation of the expectations $U_1(s,\tau)$ with (Euler-Maruyama or Milstein) approximations of the underlying asset processes. Note that for the delta control variates used in Φ^{III} , we have

$$
\nabla_{S_0} f(U_1(S_{\tau}^{0,S_0,\pm}, \tau)) = \chi_{\tau \leq T} \operatorname{LGD}(\nabla_{S_0} \max\{U_1(S_{\tau}^{0,S_0,\pm}, \tau), 0\}),
$$

which is a discontinuous function of the nested expectation $U_1(S^{0, S_0, \pm}_{\tau}, \tau)$. Consequently, the delta control variates do not satisfy the conditions of Assumption [2.2,](#page-3-2) and the analysis in Section [2.3](#page-5-0) cannot be applied directly. One method of balancing the variance reduction properties of the delta control variate with the faster convergence of antithetic MLMC for Lipschitz continuous observables is to consider the delta control variates at a single level [\[26\]](#page-26-5). Defining

$$
\tilde{\Phi}^{III} = \Phi^{III} + \frac{1}{2} \Big(S_H^{H,S_H^{\mathcal{P}}} - S_0 \Big)^{\mathsf{T}} \big(\nabla_{S_0} f \big(U_1 \big(S_\tau^{0,S_0,+}, \tau \big) \big) + \nabla_{S_0} f \big(U_1 \big(S_\tau^{0,S_0,-}, \tau \big) \big) \big),
$$

to be equivalent to Φ^{III} but without the delta control variates, this motivates considering

$$
\tilde{\Delta}^{\rm (ant)}_\ell \Phi^{\rm III} = \begin{cases} \Delta_0 \Phi^{\rm III} & \ell=0 \\ \Delta_\ell \tilde{\Phi}^{\rm III} & \ell>0. \end{cases}
$$

To remove the bias induced from including the delta control variates only at level zero, one should ensure the approximation of $\mathsf{E}^{\mathsf{Q}}\big[\nabla_{S_0} f(U_1(S^{0,S_0}_{\tau},\tau))\big]$ within $\lambda_{\varphi}^{\text{III}}$ only contains a level zero of the approximation of S_{τ}^{0,S_0} and $U_1(\cdot)$. Then

$$
\mathsf{E}^{\mathsf{Q}}\!\left[\left.\Phi^{\mathrm{III}}\tilde{p} - \lambda^\mathrm{III}_\varphi\,\right|S_H^{\mathsf{P}}, c_H, H\leq \tau\leq T \right] = \mathsf{E}^{\mathsf{Q}}\!\left[\left.\sum_{\ell=0}^\infty\tilde{\Delta}_\ell^{(\mathrm{ant})}\Phi^{\mathrm{III}}\,\right|S_H^{\mathsf{P}}, c_H, H\leq \tau\leq T \right],
$$

can be used as in Section [2](#page-2-0) to construct an (adaptive) MLMC estimate of φ . Since Section 2 assumed that the variable Z in [\(3\)](#page-1-1) can be sampled directly, we assume that S_H^{P} can be sampled exactly, which is a reasonable assumption provided $H \ll T$. If H represents a significant time interval for which approximation of S_H^{P} is costly, one can incorporate multilevel approximations of $Z := S_H^{\mathsf{P}}$ within the methods of Section [2.](#page-2-0) In this case, Assumptions [2.1](#page-3-1) and [2.5](#page-8-0) should be extended naturally to include approximations $Z_{\ell} \approx Z$.

4.4 Extension to Multiple Counterparties

 \mathbf{v}

To simplify the preceding discussion, we have considered only the case where the contract occurs between the bank and a single counterparty. Typically, the CVA capital charge considers large portfolios containing many options with $N_{cp} \geq 1$ counterparties. In this case, the adjustment made to the valuation of the banks portfolio is

$$
CVA_t := \sum_{n=1}^{N_{\rm cp}} CVA_t^{(n)},
$$

where $\text{CVA}_{t}^{(n)}$ denotes the valuation adjustment accounting for the banks exposure to counterparty n. In this approach, the credit spread at time $t \geq 0$ becomes $c_t = (c_t^{(n)})_{n=1}^{N_{\text{cp}}}$ and loss given default will differ for each counterparty. The CVA capital charge becomes the value λ_{φ} solving

$$
\varphi = P \left[\sum_{n=0}^{N_{cp}} \left(B_H^{-1} \text{CVA}_H^{(n)} - \text{CVA}_0^{(n)} \right) > \lambda_{\varphi} \right]
$$

=
$$
P \left[\sum_{n=0}^{N_{cp}} \mathsf{E}^{\mathsf{Q}} \left[\Phi^{(n)} \, \middle| \, S_H^{\mathsf{P}}, c_0^{(n)}, c_H^{(n)} \right] > \lambda_{\varphi} \right]
$$
 (31)

where $\Phi^{(n)}$ is an extension of [\(27\)](#page-15-1) for the counterparty n. While we focused on the case $N_{\rm cp} = 1$ until now, the discussion applies in the extended case where $N_{cp} > 1$. We conclude this section by considering some additional points worth considering when $N_{cp} \gg 1$.

• (Randomized subsampling of counterparties.) For a large number of counterparties ($N_{\rm cp} \gg 1$ in [\(31\)](#page-20-1)) it can be more efficient to write the sum

$$
\sum_{n=1}^{N_{\rm cp}} \mathsf{E}^{\mathsf{Q}} \Big[\Phi^{(n)} \, \Big| \, S_H^{\mathsf{P}}, c_0^{(n)}, c_H^{(n)} \Big] = \mathsf{E}^{\mathsf{Q}} \Big[\Phi^{(\kappa)} q_{\kappa}^{-1} \, \Big| \, S_H^{\mathsf{P}}, c_0^{(\kappa)}, c_H^{(\kappa)} \Big],
$$

where κ is drawn from a discrete probability distribution on $\{1, \ldots, N_{cp}\}$ with mass function q_k . As a result, we need only consider a single counterparty with each sample of the inner expectation rather than evaluate each of $\{\Phi^{(n)}\}_{n=1}^{N_{\text{cp}}}$ for each sample. This approach, known as randomised sub-sampling [\[26\]](#page-26-5), can reduce the cost of the MLMC estimator for large values of $N_{\rm cp}$.

• (Simulation of market at large number of intermediate points.) When there are many counterparties with different default times and/or many options per counterparty with differing maturities, it may be necessary to sample the assets at many intermediate time-points. In this case, it may be beneficial to use a linear interpolant to sample the market over discrete trajectories of the asset prices, opposed to generating discrete samples of the market at each default time. By combining the interpolant with an antithetic Milstein scheme, as in [\[28,](#page-26-10) Theorem 4.13], one may do this without affecting the convergence rates required within Assumption [2.1.](#page-3-1)

5 Numerical Experiments

To illustrate the preceding methods, we consider the estimation of the probability of a large loss, φ in [\(25\)](#page-14-1), for a model CVA portfolio consisting of a single stock process and a single counterparty outlined in Section [5.1](#page-21-1)[1](#page-21-2) .

5.1 Simple Synthetic Portfolio

The underlying contract between the bank and counterparty is given by a linear combination of two European call options with payoff

$$
\pi(s) = c_0(s - K_0)^+ + c_1(s - K_1)^+,
$$

where K_0, K_1 are the strike price and the constants c_0, c_1 are determined by a pre-defined value of $B_0 \mathsf{E}^{\mathsf{Q}}[B_T^{-1} \pi(S_T)]$ and the delta-neutral condition $\partial_{S_0} \mathsf{E}^{\mathsf{Q}}[B_T^{-1} \pi(S_T)] = 0$. In particular, the delta-neutral condition implies that c_0 and c_1 are of opposite sign. Importantly, this ensures that the payoff π can be negative and thus the exposure to the counterparty can be zero, which retains the need to compute a nested expectation to approximate the CVA.

We assume there is a single asset following a Geometric Brownian Motion under the Black-Scholes market assumptions with fixed interest rate $r = 1\%$ so that $B_t = e^{t/100}$. Then, under P and Q , respectively, the dynamics of the assets S are

$$
dS_t = \mu S_t dt + \varsigma S_t dW_t^P,
$$

\n
$$
dS_t = rS_t dt + \varsigma S_t dW_t^Q.
$$
\n(32)

We take $T = 1$ year, as well as $S_0 = 1$ and $\mu = \varsigma = 0.1$. To test the methods in greater generality, we ignore the existing analytic formulae for the value and stock processes, resorting instead to Milstein simulation of S and Monte Carlo estimation of the exposure at default. Specifically, we use the nested unbiased MLMC estimators [\(14\)](#page-7-0) to construct an estimator of φ as in Section [2.](#page-2-0)

¹The experiments are performed using Python, the accompanying code may be found at <https://github.com/JSpence97/mlcva>.

Following the model for default times in Section [3.1,](#page-12-0) we take $c_0 = 5\%$ and $\sigma^{(cs)} = 0.8\%/c_0$, corresponding to an A-rated counterparty according to [\[43,](#page-27-0) Table A]. The probability of large loss [\(25\)](#page-14-1) is computed with risk horizon $H = 10$ days and a fixed loss threshold $\lambda_{\varphi} = 5 \times 10^{-4}$, giving $\varphi \approx 2\%.$

5.2 MLMC Setup

Simulation of the outer expectation in [\(25\)](#page-14-1) under P is performed using the methods in Section [2.3.](#page-5-0) In particular, we obtain the MLMC estimate [\(14\)](#page-7-0) which uses the multilevel correction terms

$$
\Delta_{\mathrm{ind},\ell} = \chi_{\widehat{U}(\mathrm{UB})_{0,\ell+\eta_\ell}(S_H^{\mathbf{p}}, c_H)>\lambda_\varphi} - \chi_{\widehat{U}(\mathrm{UB})_{0,\ell-1+\eta_{\ell-1}}(S_H^{\mathbf{p}}, c_H)>\lambda_\varphi},
$$

where $U_0(S_H^{\mathsf{P}}, c_H)$, defined by [\(25\)](#page-14-1), is approximated by the unbiased estimates $\widehat{U}_{0,\ell}^{(\text{UB})}$ defined through [\(11\)](#page-6-1) using $N_{0,\ell}$ inner unbiased MLMC samples and random levels scaled by the parameters ζ_0 and ζ_1 as in Proposition [2.3.](#page-6-0) We use the techniques summarised in Section [4.3](#page-19-0) to reduce the variance of samples within U_0 . To compute the probability, we use adaptive sampling with η_ℓ chosen according to Algorithm [1](#page-10-0) with $N_{0,\ell} = N_{0,0} 2^{\ell}$ samples per level, where the refinement parameter R is chosen to be 1.95 and the refinements use a normalising factor σ_{ℓ} given by the corresponding conditional sample standard deviation [\(18\)](#page-8-3). To illustrate the effectiveness of Algorithm [1,](#page-10-0) we further consider non-adaptive sampling, taking $\eta_{\ell} \equiv 0$, with $N_{0,\ell} = N_{0,0} 2^{\gamma \ell}$ for $\gamma = 1,2$. At level $\ell = 0$ we use $N_{0,0} = 8$ samples per level. Under Assumptions [2.1](#page-3-1) and [2.5,](#page-8-0) which hold for Milstein discretisation of the SDE [\(32\)](#page-21-3) by [\[40,](#page-26-8) Theorem 10.3.5], and provided the condition [\(19\)](#page-8-4) holds for the variables in [\(26\)](#page-14-2), it follows from Theorem [2.6](#page-9-0) and Remark [2.7](#page-10-1) that $\text{Var}[\Delta_{\text{ind},\ell}]$ is of order $N_{0,\ell}^{-1/2+\delta}$ and order $N_{0,\ell}^{-1+\delta}$ for any $\delta > 0$ when using non-adaptive and adaptive sampling, respectively.

For each MLMC estimate, we approximate the optimal starting level of MLMC using the iterative procedure in Section [2.4.1.](#page-10-2) Furthermore, as discussed in Section [2.4.2,](#page-11-3) MLMC level estimates of the probability φ exhibit a high kurtosis. To obtain more stable variance and error estimates, we use a biased Bayesian approach discussed in Appendix [B.](#page-32-0)

5.3 Results

Figure [1](#page-23-0) plots various statistics of the multilevel correction terms $\Delta_{ind,\ell}$ for each method. Figure [2](#page-24-0) further plots the total work required to compute φ to accuracy ε , against ε for each method. We discuss the results below: The top left plot in Figure [1](#page-23-0) shows the expected number of Gaussian and exponential random variables required to generate a single sample of $\Delta_{ind,\ell}$. By construction, since the cost of sampling the unbiased MLMC difference [\(10\)](#page-6-2) is bounded by Proposition [2.3,](#page-6-0) the nonadaptive samplers have $\mathcal{O}(2^{\gamma_{\ell}})$ cost (with some variability due to the random cost of the unbiased MLMC estimators). On the other hand, the adaptive sampler has expected cost $E[2^{\ell+\eta_\ell}] = \mathcal{O}(2^{\ell})$ as expected from Theorem [2.6.](#page-9-0) The top right plot in Figure [1](#page-23-0) shows the variance $V_{\ell} = \text{Var}[\Delta_{\text{ind},\ell}]$ as a function of ℓ . In agreement with taking the limit $p \to \infty$ in Assumption [2.5](#page-8-0) and Theorem [2.6,](#page-9-0) the non-adaptive samplers have $V_{\ell} = \mathcal{O}(N_{0,\ell}^{-1/2}) = \mathcal{O}(2^{-\gamma_{\ell}/2})$. On the other hand, the adaptive sampler has $V_{\ell} = \mathcal{O}(N_{0,\ell}^{-1}) = \mathcal{O}(2^{-\ell})$. Combined with an expected sampling cost of order 2^{ℓ} , this improves the convergence rate of the adaptive MLMC over the non-adaptive estimators. The bottom left plot in Figure [1](#page-23-0) shows the kurtosis of $\Delta_{ind,\ell}$ against ℓ . We observe Kurtosis $[\Delta_{ind,\ell}] = \mathcal{O}(V_{\ell}^{-1})$ for all methods. As mentioned above, this reduces the accuracy of standard estimates for the overall error made by MLMC, especially when L is large. This highlights the necessity to use more stable

Figure 1: Statistics of the correction terms $\Delta_{ind,\ell}$ plotted against ℓ for the MLMC estimators using both adaptive and non-adaptive sampling. From top left to bottom right, the plots show the expected cost, C_{ℓ} , of generating a single sample of $\Delta_{ind,\ell}$, computed as the number of independent Gaussian and Exponential random variables simulated, the variance $V_{\ell} = \text{Var}[\Delta_{\text{ind},\ell}]$ and the kurtosis Kurtosis $[\Delta_{ind,\ell}]$, each versus ℓ . The bottom right plot shows the ratio R_{ℓ_0} in Section [2.4.1](#page-10-2) versus ℓ_0 , which is used to determine a starting level for the MLMC estimates.

estimates of the error as discussed in Section [2.4](#page-7-1) to improve the overall stability of MLMC. The bottom right plot in Figure [1](#page-23-0) shows the ratio R_{ℓ_0} defined in Section [2.4.1](#page-10-2) against ℓ_0 . To ensure MLMC performs optimally, one should start from a level ℓ_0 for which $R_{\ell_0} < 1$.

In Figure [2,](#page-24-0) we see a reduction in cost by a factor of around 7 times when using adaptive opposed to non-adaptive MLMC when computing φ to 2 significant figures. We also see that our results closely match the complexities $\mathcal{O}(\varepsilon^{-5/2})$ for non-adaptive sampling and $\mathcal{O}(\varepsilon^{-2}(\log \varepsilon)^2)$ for adaptive sampling discussed in Section [2.4,](#page-7-1) where the factor $(\log \varepsilon)^2 = o(\varepsilon^{-\delta})$ for any $\delta > 0$ is used to model the limiting convergence rate from Theorem [2.6](#page-9-0) as $p \to \infty$. A reference line shows the theoretical $\mathcal{O}(\varepsilon^{-5})$ complexity discussed in Section [2.2](#page-3-0) for single-level nested Monte Carlo in Section [2.](#page-2-0) This illustrates savings of several orders of magnitude at the smallest computed error tolerance by using the adaptive hierarchical estimator compared to using standard Monte Carlo.

Figure 2: Total work required to compute the probability of large loss due to changes in the CVA multiplied by ε^2 versus ε , normalised by the true value of φ .

6 Conclusion

Due to large costs induced by the approximation of market factors combined with several nested Monte Carlo averages, accurate estimation of credit risk measures is often considered infeasible using traditional Monte Carlo methods. By combining recent developments in MLMC for nested simulation with variance reduction techniques, this paper illustrates an efficient framework for accurate simulation of these risk measures at greatly reduced costs. In the context of the CVA-VaR problem, we are able to reduce the cost of achieving accuracy ε from $\mathcal{O}(\varepsilon^{-5})$ for traditional nested Monte Carlo to $\mathcal{O}(\varepsilon^{-2-\delta})$, for any $\delta > 0$, using MLMC with an adaptive number of inner samples of the loss. This results in computational savings of several orders of magnitude for realistic problems.

Future directions of this work include numerical studies for larger portfolios, containing a greater number of counterparties and the posting of collateral as discussed in Sections [3.3](#page-14-0) and [4.4,](#page-20-0) and a combination of the methods in this work with multilevel stochastic approximation techniques [\[13,](#page-25-3) [16,](#page-25-5) [19\]](#page-26-2) to compute the value-at-risk λ_{φ} associated with a fixed confidence level φ . Further analysis of the numerical estimator is required to handle non-linear nested market factors such as the initial margin and also to include more general models for default which cannot be simulated directly, and will require more advanced importance sampling methods. In more general models for default, an important concept to consider is that of wrong way risk [\[30,](#page-26-20)[33,](#page-26-1)[38\]](#page-26-21), where the probability of counterparty default is positively correlated to the risk-neutral valuation of the portfolio, leading to greater credit risk.

Acknowledgements

The authors wish to express thanks for helpful comments from two anonymous reviewers on an earlier draft of this paper.

JS was supported by the EPSRC Centre for Doctoral Training in Mathematical Modelling, Analysis and Computation (MAC-MIGS) funded by the UK Engineering and Physical Sciences Research Council (grant EP/S023291/1), Heriot-Watt University and the University of Edinburgh.

References

- [1] Agarwal, A., De Marco, S., Gobet, E., L´opez-Salas, J. G., Noubiagain, F., and Zhou, A. Numerical approximations of McKean anticipative backward stochastic differential equations arising in initial margin requirements. ESAIM: ProcS, 65:1-26, 2019.
- [2] J. Ahlberg. Credit value adjustment. Masters thesis, available at Lund University [\(link\)](http://lup.lub.lu.se/student-papers/record/4057708), 2013.
- [3] O. Bardou, N. Frikha, and G. Pagés. Computing VaR and CVaR using stochastic approximation and adaptive unconstrained importance sampling. Monte Carlo Methods and Applications, 15(3):173–210, 2009.
- [4] Basel Committee on Banking Supervision. Basel III: A global regulatory framework for more resilient banks and banking systems. https://www.bis.org/publ/bcbs189_dec2010.htm, 2010.
- [5] C. Bayer, C. B. Hammouda, and R. Tempone. Multilevel Monte Carlo combined with numerical smoothing for robust and efficient option pricing and density estimation. arXiv preprint [arXiv:2003.05708](https://arxiv.org/abs/2003.05708), 2023.
- [6] C. Bayer, C. B. Hammouda, and R. Tempone. Numerical smoothing with hierarchical adaptive sparse grids and quasi-Monte Carlo methods for efficient option pricing. Quantitative Finance, 23(2):209–227, 2023.
- [7] F. Bourgey, S. De Marco, E. Gobet, and A. Zhou. Multilevel Monte Carlo methods and lower-upper bounds in initial margin computations. Monte Carlo Methods and Applications, 26(2):131–161, 2020.
- [8] D. Brigo, M. Morini, and A. Pallavicini. Counterparty credit risk, collateral and funding: with pricing cases for all asset classes. Wiley Finance Series. John Wiley and Sons, 2013.
- [9] M. Broadie, Y. Du, and C. Moallemi. Efficient risk estimation via nested sequential simulation. Management Science, 57(6):1172–1194, 2011.
- [10] K. Bujok, B. Hambly, and C. Reisinger. Multilevel simulation of functionals of Bernoulli random variables with application to basket credit derivatives. Methodology and Computing in Applied Probability, 17(3):579–604, 2015.
- [11] K. Cliffe, M. Giles, R. Scheichl, and A. Teckentrup. Multilevel Monte Carlo methods and applications to elliptic PDEs with random coefficients. Computing and Visualization in Science, 14(1):3–15, 2011.
- [12] N. Collier, A.-L. Haji-Ali, F. Nobile, E. von Schwerin, and R. Tempone. A continuation multilevel Monte Carlo algorithm. BIT Numerical Mathematics, 55(2):399–432, 2015.
- [13] S. Crépey, N. Frikha, and A. Louzi. A multilevel stochastic approximation algorithm for value-at-risk and expected shortfall estimation. arXiv preprint [arXiv:2304.01207](https://arxiv.org/abs/2304.01207), 2023.
- [14] S. Crépey, N. Frikha, A. Louzi, and G. Pagès. Asymptotic error analysis of multilevel stochastic approximations for the value-at-risk and expected shortfall. arXiv preprint [arXiv:2311.15333](https://arxiv.org/abs/2311.15333), 2023.
- [15] S. Dereich. General multilevel adaptations for stochastic approximation algorithms II: CLTs. Stochastic Processes and their Applications, 132:226–260, 2021.
- [16] S. Dereich and T. M¨uller-Gronbach. General multilevel adaptations for stochastic approximation algorithms of Robbins–Monro and Polyak–Ruppert type. Numerische Mathematik, 142(2):279–328, 2019.
- [17] T. Dodwell, S. Kynaston, R. Butler, R. Haftka, N. H. Kim, and R. Scheichl. Multilevel Monte Carlo simulations of composite structures with uncertain manufacturing defects. Probabilistic Engineering Mechanics, 63:103–116, 2021.
- [18] D. Elfverson, F. Hellman, and A. Målqvist. A multilevel Monte Carlo method for computing failure probabilities. SIAM/ASA Journal on Uncertainty Quantification, 4(1):312–330, 2016.
- [19] N. Frikha. Multi-level stochastic approximation algorithms. The Annals of Applied Probability, 26(2):933–985, 2016.
- [20] M. Giles. Multilevel Monte Carlo path simulation. Operations Research, 56(3):607–617, 2008.
- [21] M. Giles. Multilevel Monte Carlo methods. Acta Numerica, 24:259–328, 2015.
- [22] M. Giles. MLMC techniques for discontinuous functions. Monte Carlo and Quasi-Monte Carlo Methods 2022. Springer, to appear, 2024.
- [23] M. Giles and T. Goda. Decision-making under uncertainty: using MLMC for efficient estimation of EVPPI. Statistics and Computing, 29(4):739–751, 2019.
- [24] M. Giles and A.-L. Haji-Ali. Multilevel nested simulation for efficient risk estimation. SIAM/ASA Journal on Uncertainty Quantification, 7(2):497–525, 2019.
- [25] M. Giles and A.-L. Haji-Ali. Multilevel path branching for digital options. arXiv pre-print: 2209.03017, 2022.
- [26] M. Giles and A.-L. Haji-Ali. Sub-sampling and other considerations for efficient risk estimation in large portfolios. Journal of Computational Finance, 26(1):113–140, 2022.
- [27] M. Giles, T. Nagapetyan, and K. Ritter. Multilevel Monte Carlo approximation of distribution functions and densities. SIAM/ASA Journal on Uncertainty Quantification, 3(1):267–295, 2015.
- [28] M. Giles and L. Szpruch. Antithetic multilevel Monte Carlo estimation for multi-dimensional SDEs without Lévy area simulation. Annals of Applied Probability, 24(4):1585–1620, 2014.
- [29] P. Glasserman. Monte Carlo methods in Financial Engineering, volume 53 of Stochastic Modelling and Applied Probability. Springer New York, 2003.
- [30] P. Glasserman and L. Yang. Bounding wrong-way risk in CVA calculation. Columbia Business School Research Paper No. 15-59, Available at SSRN: *[https: // ssrn. com/ abstract= 2607649](https://ssrn.com/abstract=2607649) [http: // dx. doi. org/ 10. 2139/ ssrn. 2607649](http://dx.doi.org/10.2139/ssrn.2607649)* , 2015.
- [31] M. Gordy and S. Juneja. Nested simulation in portfolio risk measurement. Management Science, 56(10):1833–1848, 2010.
- [32] A. Green. XVA: Credit, Funding and Capital Valuation Adjustments. The Wiley Finance Series. Wiley, Newark, 2015.
- [33] J. Gregory. The xVA Challenge: Counterparty Risk, Funding, Collateral, Capital and Initial Margin. John Wiley and Sons, fourth edition, 2020.
- [34] A.-L. Haji-Ali, J. Spence, and A. Teckentrup. Adaptive multilevel Monte Carlo for probabilities. SIAM Journal on Numerical Analysis, 60(4):2125–2149, 2022.
- [35] P. Henry-Labordére. Deep primal-dual algorithm for BSDEs: Applications of machine learning to CVA and IM. Available at SSRN: https://ssrn.com/abstract=3071506 *[http: // dx. doi. org/ 10. 2139/ ssrn. 3071506](http://dx.doi.org/10.2139/ssrn.3071506)* , 2017.
- [36] M. Hofer and P. Karlsson. Efficient calibration for CVA using multi-level Monte Carlo. Available at SSRN: *[https: // ssrn. com/ abstract= 2776932](https://ssrn.com/abstract=2776932)* , 2017.
- [37] J. Hull, M. Predescu, and A. White. Bond prices, default probabilities and risk premiums. Journal of Credit Risk, 1(2):53–60, 2005.
- [38] J. Hull and A. White. CVA and wrong-way risk. Financial Analysts Journal, $68(5):58-69$, 2012.
- [39] F. Jonsson. On the heavy-tailedness of Student's t-statistic. Bernoulli, 17(1):276–289, 2011.
- [40] P. Kloeden and E. Platen. Numerical Solution of Stochastic Differential Equations, volume 23 of Stochastic Modelling and Applied Probability. Springer Berlin, Heidelberg, third edition, 1999.
- [41] D. McLeish. A general method for debiasing a Monte Carlo estimator. Monte Carlo Methods and Applications, 17(4):301–315, 2011.
- [42] V. Peña, T. Lai, and Q. Shao. Self-Normalized Processes: Limit Theory and Statistical Applications. Probability and Its Applications. Springer Berlin Heidelberg, 2008.
- [43] M. Pykhtin. Model foundations of the Basel III standardised CVA charge. Risk, 25(7):60–66, 2012.
- [44] C.-H. Rhee and P. Glynn. A new approach to unbiased estimation for SDEs. In Proceedings of the 2012 Winter Simulation Conference. IEEE, 2012.
- [45] C.-H. Rhee and P. Glynn. Unbiased estimation with square root convergence for SDE models. Operations Research, 63(5):1026–1043, 2015.
- [46] J. Spence. Hierarchical and Adaptive Methods for Accurate and Efficient Risk Estimation. PhD thesis, Heriot-Watt University, 2023.
- [47] Y. Syed and G. Wang. Optimal randomized multilevel Monte Carlo for repeatedly nested expectations. Proceedings of the 40th International Conference on Machine Learning, PLMR, 202:33343–33364, 2023.
- [48] Z. Xu, Z. He, and X. Wang. Efficient risk estimation via nested multilevel quasi-Monte Carlo simulation. Journal of Computational and Applied Mathematics, 443:115745, 2024.
- [49] Z. Zhou, G. Wang, J. Blanchet, and P. Glynn. Unbiased optimal stopping via the MUSE. Stochastic Processes and their Applications, 166:104088, 2023.

A Proofs

To prove Proposition [2.3,](#page-6-0) we rely on the following general result to bound moments of antithetic multilevel differences of the form [\(9\)](#page-5-2).

 $\bold{Lemma \; A.1. \; Let \;} U^{(0)}_{1,\ell}.$ $_{1,\ell-1}^{(0)}$ and $U_{1,\ell}^{(1)}$ $1,_{1,\ell-1}$ be conditionally independent approximations of $U_1(Y)$ at level $\ell-1$, with finite q_1^{th} moment for some $q_1 \geq 2$. Let $U_{1,\ell} := (U_{1,\ell-1}^{(0)} + U_{1,\ell}^{(1)})$ $\binom{1}{1,\ell-1}/2$ be an approximation to $U_1(Y)$ at level ℓ . For an $f : \mathbb{R} \to \mathbb{R}$ satisfying Assumption [2.2,](#page-3-2) there is a constant $c > 0$, independent of ℓ , such that for all $2 \le q_0 \le q_1$ and $\ell \ge 0$ we have

$$
\mathsf{E}\bigg[\bigg|f(U_{1,\ell}) - \frac{1}{2}\Big(f(U_{1,\ell-1}^{(0)}) + f(U_{1,\ell-1}^{(1)})\Big)\bigg|^{q_0}\bigg] \le c\bigg(\mathsf{E}\bigg[\left|U_{1,\ell-1}^{(0)} - U_1(Y)\right|^{q_1}\bigg]^{\min\{2q_0/q_1,1\}} + \mathsf{E}[|U_{1,\ell} - U_1(Y)|^{q_1}|^{(q_0+1)/(q_1+1)}\bigg).
$$

Comment. The proof of Lemma [A.1](#page-27-6) follows similar steps to [\[24,](#page-26-4) Theorem 4.1], [\[7,](#page-25-9) Proposition 5], [\[23,](#page-26-9) Theorem 3] and [\[28,](#page-26-10) Theorem 5.2]. A slightly more general version of this result, which is not specific to unbiased approximations of a nested conditional expectation may be found in $/46$, Corollary 2.6]. Since the complexity of MLMC estimates depends on the variance of the multilevel differences, several existing results assume $q_0 = 2$. However, when estimating the triply nested expectation [\(3\)](#page-1-1), since Assumption [2.5](#page-8-0) requires more than convergence in the second moment, Propo-sition [2.3](#page-6-0) includes the asymptotic behaviour of $2 \le q_0 \le q_1$ moments of the multilevel differences. A similar approach for an arbitrary number of nested expectations may be found in $\ket{49}$. Theorem 1] and $\frac{1}{47}$, Theorems 2.2 and 2.4].

Proof. Throughout this proof, consider

$$
\Delta_{\ell} f := f\left(\frac{U_{1,\ell-1}^{(0)} + U_{1,\ell-1}^{(1)}}{2}\right) - \frac{1}{2} \sum_{i=0}^{1} f\left(U_{1,\ell-1}^{(i)}\right).
$$

Following the same steps as [\[28,](#page-26-10) Theorem 5.2], for some $\psi > 0$ consider the indicator functions

$$
E_0 := \chi_{\inf_{u \in \Gamma} |U_1(Y) - u| > \psi}
$$

\n
$$
E_1 := \chi_{|U_{1, \ell-1}^{(0)} - U_1(Y)| \le \psi/2}
$$

\n
$$
E_2 := \chi_{|U_{1, \ell-1}^{(0)} - U_{1, \ell-1}^{(1)}| \le \psi/2}
$$

and let $E_j^c := 1 - E_j$ for $j = 0, 1, 2$. Then,

$$
\mathsf{E}[\left|\Delta_{\ell}f\right|^{q_0}]\leq \mathsf{E}[\left|\Delta_{\ell}f\right|^{p}E_0E_1E_2]+\sum_{j=0}^{2}\mathsf{E}\left[\left|\Delta_{\ell}f\right|^{p}E_j^{c}\right].
$$

For the first term, note that $E_0E_1E_2 = 1$ implies that f is twice differentiable on an open interval of radius ψ centred on $U_1(Y)$ and containing both $U_{1,\ell}^{(0)}$ $U_{1,\ell-1}^{(0)}$ and $U_{1,\ell}^{(1)}$. $1, \ell-1$. In particular, since $U_{1,\ell} =$ $(U_{1,\ell-1}^{(0)} + U_{1,\ell}^{(1)}$ $\sum_{i,\ell=1}^{(1)}$ /2 lies in this interval, and since $\Delta_{\ell} f = 0$ on a linearisation of f, one can use the Lipschitz property of f and a second order Taylor expansion to conclude that when $E_0E_1E_2 = 1$, the following two inequalities hold

$$
|\Delta_{\ell} f| \le L_f \left| U_{1,\ell-1}^{(0)} - U_{1,\ell-1}^{(1)} \right|
$$

$$
|\Delta_{\ell} f| \le L'_f \left| U_{1,\ell-1}^{(0)} - U_{1,\ell-1}^{(1)} \right|^2.
$$

By taking appropriate powers of the above expansions, with $v = \max\{0, (2q_0 - q_1)/q_0\}$, for $q_0 \leq q_1$ it follows that

$$
\begin{split} \mathsf{E}[\,|\Delta_{\ell}f|^{q_0}E_0E_1E_2\,] &\leq 2^{q_1-1}(L_f)^{vq_0}(L_f')^{(1-v)q_0}\mathsf{E}\bigg[\,\Big|U_{1,\ell-1}^{(0)}-U_1(Y)\Big|^{min\{2q_0,q_1\}}\bigg] \\ &\leq 2^{q_1-1}(L_f)^{vq_0}(L_f')^{(1-v)q_0}\mathsf{E}\bigg[\,\Big|U_{1,\ell-1}^{(0)}-U_1(Y)\Big|^{q_1}\bigg]^{min\{2q_0/q_1,1\}}, \end{split}
$$

using Hölder's inequality in the final line.

For the remaining terms, since f is L_f -Lipschitz, it follows for $j = 0, 1, 2$ that

$$
\mathsf{E}\big[\,|\Delta_{\ell}f|^{q_0}E_j^c\,\big] \le (L_f)^{q_0}\mathsf{E}\Big[\,\Big|U_{1,\ell-1}^{(0)} - U_{1,\ell-1}^{(1)}\Big|^{q_0}E_j^c\,\Big] \newline \le (L_f)^{q_0}2^{q_0-1}\mathsf{E}\Big[\,\Big|U_{1,\ell-1}^{(0)} - U_1(Y)\Big|^{q_1}\,\Big]^{q_0/q_1}\mathsf{E}\big[\,E_j^c\,\big]^{(q_1-q_0)/q_1},
$$

using Hölder and Jensen's inequalities in the final line for $q_0 \leq q_1$. For $j = 0$, we have

$$
\mathsf{E}[E_0^c] = \mathsf{P}\bigg[\inf_{u \in \Gamma} |U_1(Y) - u| \le \psi\bigg] \le \bar{\rho}\psi
$$

for any $\psi < \bar{\delta}$ under Assumption [2.2.](#page-3-2) For $j = 1$ we have,

$$
\mathsf{E}[E_1^c] = \mathsf{E}\Big[\chi_{|U_{1,\ell-1}^{(0)} - U_1(Y)| > \psi/2}\Big] \leq 2^{q_1} \psi^{-q_1} \mathsf{E}\Big[\Big|U_{1,\ell-1}^{(0)} - U_1(Y)\Big|^{q_1}\Big].
$$

Similar steps show that

$$
\begin{aligned} \mathsf{E}[\, E_2^c \,] \leq 2^{q_1} \psi^{-q_1} \mathsf{E}\Big[\, \Big| U_{1, \ell - 1}^{(0)} - U_{1, \ell - 1}^{(1)} \Big|^{q_1} \, \Big] \\ \leq 2^{q_1} 2^{q_1 - 1} \psi^{-q_1} \mathsf{E}\Big[\, \Big| U_{1, \ell - 1}^{(0)} - U_1(Y) \Big|^{q_1} \, \Big] \end{aligned}
$$

using Jensen's inequality in the final line. To make the above terms of the same order, one should take

$$
\psi = \min\left\{1, \frac{\bar{\delta}}{\sup_{\ell \geq 0} |U_{1,\ell-1}^{(0)} - U_1(Y)|}\right\} \mathsf{E}\left[\left|U_{1,\ell-1}^{(0)} - U_1(Y)\right|^{q_1}\right]^{1/(q_1+1)}.
$$

Hence, there is $b > 0$ independent of ℓ such that for $j = 0, 1, 2$

$$
\mathsf{E}\big[\left|\Delta_{\ell}f\right|^{p}E_{j}^{c}\big] \leq b\mathsf{E}\big[\left|U_{1,\ell-1}^{(0)} - U_{1}(Y)\right|^{q_{1}}\big]^{q_{0}/q_{1}}\mathsf{E}\big[\left|U_{1,\ell-1}^{(0)} - U_{1}(Y)\right|^{q_{1}}\big]^{(q_{1}-q_{0})/q_{1}(q_{1}+1)}
$$
\n
$$
= b\mathsf{E}\big[\left|U_{1,\ell-1}^{(0)} - U_{1}(Y)\right|^{q_{1}}\big]^{(q_{0}+1)/(q_{1}+1)},
$$

which completes the result.

Proof of Proposition [2.3.](#page-6-0) Throughout this proof, we denote by $c > 0$ an arbitrary constant which is independent of ℓ and k . We begin by proving that $\Delta(Z)$ has finite expected sampling cost. First, by Assumption [2.1,](#page-3-1) we know that the cost of sampling $\Delta_k X(Y)$ is bounded by $c2^k$. Therefore

$$
E[Cost(\Delta_{\kappa_1} X(Y))] = \sum_{k=0}^{\infty} Cost(\Delta_k X(Y))p_{1,k}
$$

$$
\leq c \sum_{k=0}^{\infty} 2^{(1-\zeta_1)k},
$$

which is finite using the geometric decay of $p_{1,k}$ since $1 < \zeta_1 = 3/2$ by hypothesis. Hence,

$$
\mathsf{E}\Big[\operatorname{Cost}\Big(\widehat{U}_{1,\ell}^{\mathrm{(UB)}}(Y_\ell)\Big)\,\Big]\propto 2^\ell,
$$

from which it follows from Assumption [2.1](#page-3-1) and similar steps as above that

$$
\mathsf{E}[\mathrm{Cost}(\Delta(Z))] \le c \sum_{\ell=0}^{\infty} 2^{(1-\zeta_0)\ell},
$$

which is finite since it is assumed that $\zeta_0 = 17/16 > 1$.

 \Box

It remains to provide bounds on the moments of $\Delta(Z)$. We will show, for the random variable $\Delta_\ell^{\rm (UB)}$ $\ell^{(0)}$ $f(Y_{\ell}, Y_{\ell-1})$ defined in [\(9\)](#page-5-2), that for fixed $\ell \geq 1$ and $2 \leq q_0 < 3$, we have

$$
\mathsf{E}\left[\left|\Delta_{\ell}^{(\text{UB})}f(Y_{\ell}, Y_{\ell-1})\right|^{q_0}\right] \le c2^{-3(q_0+1)\ell/8 + \delta\ell},\tag{33}
$$

for any $\delta > 0$. We split a proof of [\(33\)](#page-30-0) into three steps:

1. We first show that, for $2 \le q_1 < 3$,

$$
\mathsf{E}\left[\left|\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell}) - U_1(Y)\right|^{q_1}\right] \le c2^{-q_1\ell/2}.\tag{34}
$$

2. Secondly, for $2 \le q_0 \le q_1$, we prove

$$
\mathsf{E}\Big[\Big|\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell}) - \widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell-1})\Big|^{q_0}\,\Big] \le c2^{-3\ell/2}.\tag{35}
$$

3. Use Lemma [A.1](#page-27-6) to bound moments of $\Delta_{\ell}^{\text{(UB)}} f$.

Step 1. For $q_1 < 3$, it follows from the definition of κ_1 with $\zeta_1 = 3/2$ that

$$
\mathsf{E}\Big[\left|\Delta_{\kappa_1}X(Y_{\ell})\,p_{1,\kappa_1}^{-1}\right|^{q_1}\Big] = \sum_{k=0}^{\infty} p_{1,k}^{1-q_1} \mathsf{E}\big[\left|\Delta_k X(Y_{\ell})\right|^{q_1}\big] \n\leq p_{1,0}^{1-q_1} c \sum_{k=0}^{\infty} 2^{3(q_1-1)k/2} 2^{-q_1 k} < \infty.
$$

Therefore, using Assumption [2.1,](#page-3-1) Jensen's inequality, the tower property and the discrete Burkholder-Davis-Gundy inequality following similar steps to [\[24,](#page-26-4) Lemma 2.5], we obtain the bound

$$
\begin{aligned} & \mathsf{E}\Big[\left|\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell}) - U_1(Y)\right|^{q_1}\Big] \\ &\leq 2^{q_1 - 1} \Big(\mathsf{E}\Big[\left|\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell}) - U_1(Y_{\ell})\right|^{q_1}\Big] + \mathsf{E}[\left|U_1(Y_{\ell}) - U_1(Y)\right|^{q_1}]\Big) \\ &\leq c2^{-q_1\ell/2}.\end{aligned}
$$

Step 2. For $\ell \geq 1$, denote the double difference

$$
\Delta_{\ell} \Delta_k X(Y) := \Delta_k X(Y_{\ell}) - \Delta_k X(Y_{\ell-1})
$$

= $X_k(Y_{\ell}) - X_{k-1}(Y_{\ell}) - X_k(Y_{\ell}) + X_{k-1}(Y_{\ell})$

and let

$$
\overline{\Delta_{\ell}\Delta_{\kappa_1}X(Y)}:=\Delta_{\kappa_1}X(Y)\,p_{1,\kappa_1}^{-1}-\mathsf{E}\big[\,\Delta_{\ell}\Delta_{\kappa_1}X(Y)\,p_{1,\kappa_1}^{-1}\,\big|\,Y_{\ell},Y_{\ell-1}\,\big]
$$

represent the centralised random double differences with mean zero. Under Assumption [2.1](#page-3-1) it follows that

$$
\mathsf{E}[\left|\Delta_{\ell}\Delta_{k}X\right|^{q_{0}}] \leq c2^{-q_{0}\max\{\ell,k\}}.\tag{36}
$$

Using the triangle and discrete Burkholder-Davis-Gundy inequalities, since $\overline{\Delta_{\ell} \Delta_{\kappa_1} X(Y)}$ has mean zero, following similar steps to $[24, \text{Lemma } 2.5]$, we have

$$
\label{eq:4.13} \begin{aligned} &\mathsf{E}\Big[\left|\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell})-\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell-1})\right|^{q_{0}}\Big]\\ &\leq \mathsf{E}\Bigg[\left|\frac{1}{N_{1,\ell}}\sum_{n=1}^{N_{1,\ell}}\Delta_{\ell}\Delta_{\kappa_{1}^{(n)}}X^{(n)}(Y)\right|^{q_{0}}\Bigg]\\ &\leq 2^{q_{0}-1}\Bigg(\mathsf{E}\Bigg[\left|\frac{1}{N_{1,\ell}}\sum_{n=1}^{N_{1,\ell}}\overline{\Delta_{\ell}\Delta_{\kappa_{1}^{(n)}}X^{(n)}(Y)}\right|^{q_{0}}\Bigg]+ \mathsf{E}\Big[\left|\mathsf{E}\big[\Delta_{\ell}\Delta_{\kappa_{1}}X(Y)\,p^{-1}_{1,\kappa_{1}}\,\big|\,Y_{\ell},Y_{\ell-1}\,\big]\right|^{q_{0}}\Bigg]\Bigg)\\ &\leq c_{q_{0}}\Big(N_{1,\ell}^{-q_{0}/2}\mathsf{E}\Big[\left|\Delta_{\ell}\Delta_{\kappa_{1}}X(Y)\,p^{-1}_{1,\kappa_{1}}\right|^{q_{0}}\Big]+ \mathsf{E}\Big[\left|\mathsf{E}\big[\Delta_{\ell}\Delta_{\kappa_{1}}X(Y)\,p^{-1}_{1,\kappa_{1}}\,\big|\,Y_{\ell},Y_{\ell-1}\,\big]\right|^{q_{0}}\Big]\Bigg), \end{aligned}
$$

where the constant c_{q0} depends only on q_0 . The latter term can be bounded using [\(8\)](#page-5-1), Jensen's inequality and the tower property to obtain

$$
\mathsf{E}\Big[\left|\mathsf{E}\big[\Delta_{\ell}\Delta_{\kappa_{1}}X(Y)\,p_{1,\kappa_{1}}^{-1}\,\big|\,Y_{\ell},Y_{\ell-1}\,\big]\right|^{q_{0}}\Big] = \mathsf{E}\big[\left|\mathsf{E}\big[X(Y_{\ell})-X(Y_{\ell-1})\,\big|\,Y_{\ell},Y_{\ell-1}\,\big]\right|^{q_{0}}\big] \\\leq \mathsf{E}\big[\left|X(Y_{\ell})-X(Y_{\ell-1})\right|^{q_{0}}\big] \\\leq c2^{-q_{0}\ell},
$$

where we used Assumption [2.1](#page-3-1) in the final line. For the remaining term, by the definition of κ_1 with $\zeta_1 = 3/2$ and using [\(36\)](#page-30-1), we have

$$
\mathsf{E}\Big[\left|\Delta_{\ell}\Delta_{\kappa_1}X(Y)\,p_{1,\kappa_1}^{-1}\right]^{q_0}\Big] = \sum_{k=0}^{\infty} p_{1,k}^{1-q_0} \mathsf{E}\big[\left|\Delta_{\ell}\Delta_k X(Y)\right]^{q_0}\big] \n\leq p_{1,0}^{1-q_0} c \sum_{k=0}^{\infty} 2^{3(q_0-1)k/2} 2^{-q_0 \max\{\ell,k\}} \n\leq c 2^{(q_0-3)\ell/2},
$$

where we use the fact that $q_0 < 3$ so that the geometric series is convergent.

The previous results, with $N_{1,\ell} \propto 2^{\ell}$ are enough to show that

$$
\mathsf{E}\Big[\Big|\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell}) - \widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell-1})\Big|^{q_0}\,\Big] \leq c2^{-3\ell/2} + 2^{-q_0\ell} \leq c2^{-3\ell/2},
$$

where the last line follows since $q_0 \geq 2$.

Step 3. Using the fact that f satisfies Assumption [2.2,](#page-3-2) it follows from Lemma [A.1,](#page-27-6) [\(34\)](#page-30-2) and

[\(35\)](#page-30-3) that for $\ell \geq 1$ and $2 \leq q_0 \leq q_1$ we have

$$
\mathsf{E}\Big[\Big|\Delta_{\ell}^{(\text{UB})}f(Y_{\ell}, Y_{\ell-1})\Big|^{q_{0}}\Big] \leq c\mathsf{E}\Big[\Big|\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell}) - \widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell-1})\Big|^{q_{1}}\Big]^{min\{2q_{0}/q_{1},1\}} + c\mathsf{E}\Big[\Big|\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell}) - U_{1}(Y)\Big|^{q_{1}}\Big]^{(q_{0}+1)/(q_{1}+1)} + c\mathsf{E}\Big[\Big|\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell}) - U_{1}(Y)\Big|^{q_{1}}\Big]^{(q_{0}+1)/(q_{1}+1)} + c\mathsf{E}\Big[\Big|\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell}) - U_{1}(Y)\Big|^{q_{1}}\Big]^{(q_{0}+1)/(q_{1}+1)} + c\mathsf{E}\Big[\Big|\widehat{U}_{1,\ell}^{(\text{UB})}(Y_{\ell}) - U_{1}(Y)\Big|^{q_{1}}\Big]^{(q_{0}+1)/(q_{1}+1)}
$$

$$
\leq c\Big(2^{-\min\{2q_{0},q_{1}\}\ell/2} + 2^{-q_{1}(q_{0}+1)\ell/2(q_{1}+1)}\Big)
$$

for any $\delta > 0$, where the last line follows from taking q_1 sufficiently close to 3, completing the desired bound on $E[|\Delta_{\ell}^{\text{(UB)}}|]$ $\int_{\ell}^{(UB)} f(Y_{\ell},Y_{\ell-1})|^{q_0}$].

It follows from the definition of κ_0 , with $\zeta_0 = 17/16$, that

$$
\mathsf{E}[|\Delta(Z)|^{q_0}] \le c \sum_{\ell=0}^{\infty} 2^{17(q_0-1)\ell/16} 2^{-3(q_0+1)\ell/8 + \delta\ell},
$$

which is finite provided

$$
\frac{17}{16}(q_0-1) < \frac{3(q_0+1)-\delta}{8}.
$$

By taking $\delta > 0$ small enough, the above relation holds for any $2 \le q_0 < 23/11$, completing the proof proof.

Proof of Theorem [2.6.](#page-9-0) In [\[34,](#page-26-6) Proposition 3.1 and Lemma 3.2] it is shown that under Assump-tion [2.5](#page-8-0) for $\Delta_{\text{ind},\ell}$ defined by [\(21\)](#page-9-2), with refinement parameter $1 < R < 2$, we have $\mathsf{E}[\Delta_{\text{ind},\ell}^2] = \mathcal{O}(N_{0,\ell}^{-p/(p+2)})$ and $\mathsf{E}[N_{0,\ell+\eta_\ell}] = \mathcal{O}(N_{0,\ell}),$ with the former result implying the bound on $V_\ell = \mathsf{Var}[\Delta_{\mathrm{ind},\ell}].$ Furthermore, $\mathsf{E}[\text{Cost}(\Delta_{\text{ind},\ell})] = N_{0,\ell}$, hence the total computational cost satisfies [\(15\)](#page-8-5). Substituting the asymptotic form for V_{ℓ} , we get $Cost(\varphi_{\mathbf{M}_{L}^{*},L}^{(\text{MLMC})};\varepsilon) = \mathcal{O}(\varepsilon^{-2}(\sum_{\ell=0}^{L} N_{0,\ell}^{1/(p+2)})^2)$. If $|E[\Delta_{\text{ind},\ell}]| = \mathcal{O}(N_{0,\ell}^{-1})$ then it suffices from standard MLMC complexity theory to take L such that $N_{0,L}^{-1} = \mathcal{O}(\varepsilon)$, providing the desired bound. In the absence of such a bound, one can use $|\mathsf{E}[\Delta_{\text{ind},\ell}]| \leq \mathsf{E}[\Delta_{\text{ind},\ell}^2]^{1/2} = \mathcal{O}(N_{0,\ell}^{-p/2(p+2)})$. In this case, one should take L such that $N_{0,L}^{-1} =$ $\mathcal{O}(\varepsilon^{p/2(p+2)})$, which provides the alternative bound on the cost. \Box

B Bayesian Estimates of the Error

In this appendix, we describe an approach for obtaining more robust estimates of the error and variance estimate in MLMC [\(14\)](#page-7-0). We describe a Bayesian approach, based on a similar method in [\[18\]](#page-25-15). Throughout this section, let $V_{\ell} = \text{Var}[\Delta_{ind,\ell}]$ and $E_{\ell} = |E[\Delta_{ind,\ell}]|$, where $\Delta_{ind,\ell}$ represent either the non-adaptive multilevel differences defined through [\(13\)](#page-7-2) or the adaptive differences [\(21\)](#page-9-2). For $\ell \geq 0$ there exists $p_{\ell}, q_{\ell} \in [0, 1]$ such that the multilevel correction terms $\Delta_{ind,\ell}$ take the form

$$
\Delta_{\text{ind}, \ell} = \begin{cases} 1 & \text{probability } p_{\ell} \\ -1 & \text{probability } q_{\ell} \\ 0 & \text{probability } 1 - p_{\ell} - q_{\ell}. \end{cases}
$$

In particular,

$$
V_{\ell} \leq p_{\ell} + q_{\ell}, \quad E_{\ell} = |p_{\ell} - q_{\ell}|.
$$

When ℓ is large, Theorem [2.6](#page-9-0) implies that both $p_{\ell}, q_{\ell} \ll 1$. Heuristically, when the number of samples M_{ℓ} of $\Delta_{\text{ind},\ell}$ is small, it is extremely unlikely we will observe a non-zero sample, creating unstable approximations to V_{ℓ} and E_{ℓ} . The idea is to impose a Bayesian prior on p_{ℓ} and q_{ℓ} which incorporates knowledge of the theoretical convergence rates of V_{ℓ} and E_{ℓ} such that when the number of samples M_{ℓ} is low, we may rely more heavily on the theoretical rates than on the observations, and vice-versa as M_{ℓ} becomes large.

Specifically, assume that we know a relationship of the form $V_{\ell} \approx a_0 2^{-\beta \ell}$, where the constant β can be obtained a-priori given Theorem [2.6](#page-9-0) or estimated from data and a_0 may be reasonably estimated using information from coarse levels where samples of $\Delta_{ind,\ell}$ are inexpensive. We impose a Dirichlet prior density on the quantities p_{ℓ}, q_{ℓ}

$$
\pi_{0,\ell}(p,q) = (1-p-q)^{c_{\ell}-1} p^A q^A,
$$

for constants $c_{\ell}, A > 0$. For the robustness of the MLMC estimator, it is preferable to over-estimate opposed to under-estimate V_{ℓ} . Therefore, we require the condition $\mathsf{E}_{\pi_{0,\ell}}[p_{\ell}+q_{\ell}] \geq V_{\ell}$, which is true provided $c_\ell \leq 2(A+1)(V_\ell^{-1}-1)$. Hence, using $V_\ell \approx a_0 2^{-\beta \ell}$, a reasonable conservative estimate is $c_{\ell} = 2A(a_0^{-1}2^{\beta \ell} - 1)$. Given observations $\{\Delta_{\text{ind},m}\}_{m=1}^{M_{\ell}}$, define

$$
\nu_{\ell}^{\pm}\coloneqq \sum_{m=1}^{M_{\ell}}\chi_{\Delta_{\mathrm{ind},\ell}^{(m)}=\pm 1}.
$$

Then, the maximum-a-posteriori estimate of $p_{\ell} + q_{\ell}$ is

$$
\hat{V}_{\ell} := \frac{\nu_{\ell}^{+} + \nu_{\ell}^{-} + 2A}{M_{\ell} + 2Aa_0^{-1}2^{\beta \ell} - 1}.
$$

The parameter A can be chosen to control how confident we are that $V_{\ell} \leq \hat{V}_{\ell}$ for a small number of samples M_{ℓ} . Larger values of A produce more reliable, conservative bounds.

To estimate $E_{\ell} = |p_{\ell} - q_{\ell}|$ we construct biased estimates for $p_{\ell} - q_{\ell}$ and $q_{\ell} - p_{\ell}$ separately. First, in the case where $p_{\ell} > q_{\ell}$, we wish to estimate of $p_{\ell} - q_{\ell}$. In doing so, we impose a Dirichlet $(d_{\ell}, B+1, 1)$ prior on p_ℓ, q_ℓ with density

$$
\overline{\pi_{0,\ell}}(p,q) = (1 - p - q)^{d_{\ell}-1} p^{B},
$$

for constants d_{ℓ} and B. Given the observations $\{\Delta_{ind,\ell}^{(m)}\}_{m=1}^{M_{\ell}}$ the posterior distribution becomes a Dirichlet $(d_{\ell}, \nu_{\ell}^{+} + B + 1, \nu_{\ell}^{-} + 1)$ distribution with maximum a posteriori estimator given by

$$
\frac{\nu_{\ell}^{+} - \nu_{\ell}^{-} + B}{M_{\ell} + B + d_{\ell} - 1}
$$

.

Under the prior distribution we have

$$
E_{\overline{\pi_{0,\ell}}}[p_{\ell} - q_{\ell}] = \frac{B}{B + d_{\ell} + 2},\tag{37}
$$

thus enforcing the condition

$$
\mathsf{E}_{\overline{\pi_{0,\ell}}}[p_\ell - q_\ell] \ge e_0 2^{-\alpha \ell} \approx |p_\ell - q_\ell|
$$

requires

$$
d_{\ell} \le B(e_0^{-1}2^{\alpha \ell} - 1) - 2. \tag{38}
$$

In the case where $q_{\ell} > p_{\ell}$ one can repeat this process by interchanging the roles of p_{ℓ} and q_{ℓ} in [\(37\)](#page-33-0). Taking the maximum of the two resulting estimators, with equality in [\(38\)](#page-34-0), yields

$$
\hat{E}_{\ell} := \frac{|\nu_{\ell}^{+} - \nu_{\ell}^{-}| + B}{M_{\ell} + Be_0^{-1}2^{\alpha \ell} - 3}.
$$

This method is used for the numerical experiments in Section [5,](#page-21-0) where we set the constants $A =$ $B=1$ therein.