

# On some multiplicative properties of large difference sets

I.D. Shkredov

Annotation.

*In our paper we study multiplicative properties of difference sets  $A - A$  for large sets  $A \subseteq \mathbb{Z}/q\mathbb{Z}$  in the case of composite  $q$ . We obtain a quantitative version of a result of A. Fish about the structure of the product sets  $(A - A)(A - A)$ . Also, we show that the multiplicative covering number of any difference set is always small.*

## 1 Introduction

The landmark question about solvability of equations of the form  $f(x_1, \dots, x_n) = 0$ , where  $f \in \mathbb{Z}[x_1, \dots, x_n]$  and the variables  $x_j \in X_j$  belong to some “large” but unspecified sets  $X_j$  of the prime field  $\mathbb{F}_q$  was firstly posed, probably, in [11]. Interesting in its own right the problem has a clear connection with the sum–product phenomenon [20] due to the fact that as a rule the polynomial  $f$  includes both the addition and the multiplication. This theme becomes rather popular last years, e.g., see [7]–[11], [15], [18] and many other papers.

The question about a partial resolution of some specific equations  $f(x_1, \dots, x_n) = 0$  in large subrings of rings  $\mathbb{Z}_q := \mathbb{Z}/(q\mathbb{Z})$  for composite  $q$  was firstly considered by A. Fish in [5] (nevertheless, let us remark that a similar problem was formulated in [8, Problem 5]). In particular, in [5, Corollary 1.2] A. Fish considered the polynomial  $f(x_1, x_2, x_3, x_4) = (x_1 - x_2)(x_3 - x_4)$  and proved the following result.

**Theorem 1** *Let  $q$  be a positive integer,  $A, B \subset \mathbb{Z}_q$  be sets,  $|A| = \alpha q$ ,  $|B| = \beta q$ , and suppose that  $\alpha \geq \beta$ . Then there is  $d|q$  with*

$$d \leq F(\beta) \tag{1}$$

and such that

$$d \cdot \mathbb{Z}_q \subseteq (A - A)(B - B). \tag{2}$$

Here we use the following standard notation [20], namely, given two sets  $A, B \subset \mathbb{Z}_q$ , define the *sumset* of  $A$  and  $B$  as

$$A + B := \{a + b : a \in A, b \in B\}.$$

In a similar way we define the *difference sets*  $A - A$ , the *higher sumsets*, e.g.,  $2A - A$  is  $A + A - A$  and, further, the *products sets*

$$AB := \{ab : a \in A, b \in B\}$$

the *higher product sets* and so on. Finally, if  $A \subseteq \mathbb{Z}_q$  and  $\lambda \in \mathbb{Z}_q$ , then we write

$$\lambda \cdot A = \{\lambda a : a \in A\}.$$

It is easy to see that in this generality one cannot have  $\mathbb{Z}_q = (A - A)(B - B)$  in inclusion (2) for all sets  $A, B$  and thus we indeed need this additional (but small) divisor  $d$ . In contrary, for prime  $q$  the divisor  $d$  can be omitted and the questions of this type were studied in [9] and [18]. In paper [5] the dependence on  $F(\beta)$  was triple exponential on  $\beta^{-1}$ . Using a series of other methods, we improve and generalize the last result in several directions. The signs  $\ll$  and  $\gg$  below are the usual Vinogradov symbols.

**Theorem 2** *Let  $q$  be a positive integer,  $A, B \subset \mathbb{Z}_q$  be sets,  $|A| = \alpha q$ ,  $|B| = \beta q$ , and suppose that  $\alpha \geq \beta$ . Then there is  $d|q$  with*

$$d \ll \exp(C\beta^{-4}), \quad (3)$$

where  $C > 0$  is an absolute constant and such that

$$d \cdot \mathbb{Z}_q \subseteq (A - A)(B - B). \quad (4)$$

In [5] the author posed a series of questions in much more general form, as well as for other polynomials  $f(x_1, \dots, x_n)$ . Using different approaches we partially resolve some of them, see Sections 3 and 4. In particular, we have deal with the equation

$$(a_1 - b_1)(a_2 - b_2) \equiv \lambda \pmod{q}, \quad (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B},$$

and

$$(a_1 - b_1)^2 - (a_2 - b_2)^2 \equiv \lambda \pmod{q}, \quad (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B}$$

for rather general two-dimensional sets  $\mathcal{A}, \mathcal{B} \subseteq \mathbb{Z}_q \times \mathbb{Z}_q$  and composite numbers  $q$  with some restrictions on its prime divisors, see Theorems 7, 10 and Theorem 12 below. As an example, we formulate a part of Theorem 7.

**Theorem 3** *Let  $q$  be a squarefree number,  $\mathcal{A}, \mathcal{B} \subseteq \mathbb{Z}_q^2$  be sets,  $|\mathcal{A}| = \alpha q^2$ ,  $|\mathcal{B}| = \beta q^2$ , and suppose that  $\alpha \geq \beta$ . Then*

$$d \cdot \mathbb{Z}_q^* \subseteq \{(a_1 - b_1)(a_2 - b_2) : (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B}\} \quad (5)$$

with

$$d \ll \exp(O(\omega(q) \log \omega(q) - \log \beta)).$$

In particular, for  $\mathcal{A} = \mathcal{B}$  one has with the same  $d$  that

$$d \cdot \mathbb{Z}_q \subseteq \{(a_1 - b_1)(a_2 - b_2) : (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B}\}. \quad (6)$$

Our another result may be interesting in itself (even in the case of prime  $q$ ) due to it gives a new necessary condition for a set to be a difference set but moreover, in addition, it yields another proof of Theorem 1 (see Theorems 16, 20 below).

**Theorem 4** *Let  $q$  be a positive integer,  $A \subseteq \mathbb{Z}_q$  be a set,  $|A| = \alpha q$ . Suppose that the least prime factor of  $q$  greater than  $2\alpha^{-1} + 3$ . Then there is  $X \subseteq \mathbb{Z}_q$  such that*

$$|X| \leq \frac{1}{\alpha} + 1,$$

and

$$X(A - A) = \mathbb{Z}_q.$$

The equation  $X(A - A) = \mathbb{Z}_q$  for a set  $X \subseteq \mathbb{Z}_q$  induces a coloring of  $\mathbb{Z}_q$  via suitable subsets of our difference set  $A - A$ . Hence Theorem 4 gives us a new connection between coloring problems and difference sets. Finally, our result and the Ruzsa covering lemma (see inclusion (50) below) show that for any set  $A \subseteq \mathbb{Z}_q$ ,  $|A| \gg q$ , where  $q$  is a prime number, say, the set  $A - A$  is a syndetic set in both multiplicative and additive ways.

Let us say a few words about the notation. Having a positive integer  $q$  we denote by  $\omega(q)$  the total number of prime divisors of  $q$  and by  $\tau(q)$  the number of all divisors. Let  $\varphi(q)$  be the Euler function. We use the same capital letter to denote a set  $A \subseteq \mathbb{Z}_q$  and its characteristic function  $A : \mathbb{Z}_q \rightarrow \{0, 1\}$ . If  $\mathcal{R}$  is a ring, then we write  $\mathcal{R}^*$  for the group of all inverse elements of  $\mathcal{R}$ . Let  $e_q(x) = e^{2\pi i x/q}$  and let us denote by  $[n]$  the set  $\{1, 2, \dots, n\}$ . All logarithms are to base 2.

The author thanks Alexander Fish for very useful remarks (especially Remark 17 below) and comments.

## 2 An effective version of Fish's theorem

Having a positive integer  $n$  and a set  $A \subseteq \mathbb{Z}_q \times \dots \times \mathbb{Z}_q = \mathbb{Z}_q^n$  (or just  $A \subset \mathbb{Z}^n$ ), as well as a divisor  $q_*|q$ , we write

$$\pi_{q_*}(A) = \{(a_1 \pmod{q_*}, \dots, a_n \pmod{q_*}) : (a_1, \dots, a_n) \in A\} \subseteq \mathbb{Z}_{q_*}^n.$$

We need a simple regularization result similar to [2, Lemma 2.1].

**Lemma 5** *Let  $\delta, \varepsilon \in (0, 1)$ ,  $M \geq 2$  be real numbers,  $n$  be a positive integer, and  $A \subset \mathbb{Z}_q^n$  be a set,  $|A| = \delta q^n$ . Then there is  $q_*|q$ , and a set  $A_* \subseteq A$ ,  $|\pi_{q/q_*}(A_*)| = 1$  such that  $q_* = \frac{q}{q_1 \dots q_s}$ ,  $M \leq q_j \leq \delta^{-\varepsilon^{-1}}$ ,  $s$  is the least number with  $\delta M^{\varepsilon s} > 1$  and for all  $\tilde{q}|q_*$ ,  $\tilde{q} \geq M$  one has*

$$\max_{\xi \in \mathbb{Z}_{\tilde{q}}^n} |A_* \cap \pi_{\tilde{q}}^{-1}(\xi)| \leq \frac{|A_*|}{\tilde{q}^{1-\varepsilon}}. \quad (7)$$

*Proof.* Suppose not. Then for a certain  $\xi \in \mathbb{Z}_q^n$  and  $q_1|q$ ,  $q_1 \geq M$  we find  $A' := A \cap \pi_{q_1}^{-1}(\xi)$  with  $|A'| \geq \frac{|A|}{q_1^{1-\varepsilon}}$ . Clearly,  $|\pi_{q_1}(A')| = 1$  and the density of  $A'$  in the appropriate shift of  $\mathbb{Z}_{q/q_1}^n$  is at least  $\delta q_1^\varepsilon \geq \delta M^\varepsilon$ . Hence applying the same procedure to the set  $A'$  and to the new module  $q/q_1$ , we see that our algorithm must stop after at most  $s$  steps. Notice that condition (7) holds

automatically if  $\tilde{q} \geq \delta^{-\varepsilon^{-1}}$  and hence at the final step of our procedure we find a set  $A_* \subset A$ ,  $|\pi_{q/q_*}(A_*)| = 1$ , having all required properties. This completes the proof.  $\square$

Now we are ready to obtain the main result of this section, which implies Theorem 2 from the introduction. Our proof uses the Fourier analysis (its standard facts can be found in [20], say) and classical estimates for the Kloosterman sums. Having a group  $\mathbf{G}$ , we define for any function  $f : \mathbf{G} \rightarrow \mathbb{C}$  and a representation  $\rho \in \widehat{\mathbf{G}}$  the Fourier transform of  $f$  at  $\rho$  by the formula

$$\widehat{f}(\rho) = \sum_{g \in \mathbf{G}} f(g)\rho(g). \quad (8)$$

**Theorem 6** *Let  $q$  be a positive integer,  $A, B \subset \mathbb{Z}_q$  be sets,  $|A| = \alpha q$ ,  $|B| = \beta q$ , and suppose that  $\alpha \geq \beta$ . Then there is  $d|q$  with*

$$d \ll \exp(C\beta^{-4}), \quad (9)$$

where  $C > 0$  is an absolute constant and such that

$$d \cdot \mathbb{Z}_q \subseteq (A - A)(B - B). \quad (10)$$

In addition

$$d \ll \beta^{-O(\omega(q))}. \quad (11)$$

*Proof.* Let  $q = p_1^{\rho_1} \dots p_t^{\rho_t}$ , where  $p_j$  are different primes,  $p_1 < \dots < p_t$ . Also, let  $M \geq 2$ ,  $\varepsilon \in (0, 1)$  be parameters, which we will choose later. First of all, we remove all divisors less than  $M$  from  $q$ . More precisely, for any  $p_j$ ,  $j \in [t]$  let  $\gamma_j \leq \rho_j$  be the maximal nonnegative integer such that  $p_j^{\gamma_j} \leq M$ . Clearly,  $\gamma_1 \geq \gamma_2 \geq \dots \geq \gamma_t \geq 0$  and let  $t_0 \leq t$  be the maximal  $j$  with  $\gamma_j \neq 0$ . Thus  $t_0 \leq \pi(M)$ . Now we define

$$Q_1 := \prod_{j=1}^{t_0} p_j^{\gamma_j} \leq M^{t_0} \leq \min\{M^{\pi(M)}, M^{\omega(q)}\} \quad (12)$$

and take  $A_1 \subseteq A$  such that a shift of  $A_1$  belongs to  $\mathbb{Z}_{q/Q_1}$  and has density at least  $\alpha$ . In particular,  $|\pi_{Q_1}(A_1)| = 1$  and of course such a shift exists by the Dirichlet principle. Similarly, we can do the same with the set  $B$  so as not to lose the density. Secondly, we apply Lemma 2 with  $n = 1$ ,  $A = A_1$  to regularize the set  $A_1$  and find a set  $A_* \subseteq A_1$  and a module  $q_*$  that satisfies (7) and all other restrictions. Again, using the Dirichlet principle, we take  $B_* \subseteq B$  such that the density of  $B$  does not decrease. Let  $\lambda \in \mathbb{Z}_{q_*}$  be an arbitrary number and we first suppose that  $\lambda \in \mathbb{Z}_{q_*}^*$ . To prove  $\lambda \in (A - A)(B - B)$  it is enough to show that  $\lambda \in (A_* - A_*)(B_* - B_*)$  or, equivalently, in terms of the Fourier transform it suffices obtain the inequality

$$\frac{|A_*|^2 |B_*|^2}{q_*} > \frac{1}{q_*} \sum_{r \neq 0} |\widehat{B}_*(r)|^2 \cdot \sum_{a_1, a_2 \in A_*} e_{q_*} \left( \frac{\lambda r}{a_1 - a_2} \right) := \sigma. \quad (13)$$

Now clearly,

$$\sigma \leq \frac{1}{q_*} \sum_{q_2 | q_*, q_2 > 1} \sum_{z \in \mathbb{Z}_{q_2}^*} |\widehat{B}_*(z q_* q_2^{-1})|^2 \left| \sum_{a_1, a_2 \in A_*} e_{q_2} \left( \frac{\lambda z}{a_1 - a_2} \right) \right|. \quad (14)$$

In terms of the Kloosterman sums

$$K_q(\lambda, r) := \sum_{x \in \mathbb{Z}_q^*} e_q \left( \frac{\lambda}{x} + rx \right)$$

and the density function

$$\eta_{q_2}(\xi) := |\{a \in A_* : a \equiv \xi \pmod{q_2}\}| \quad (15)$$

one has (recall that  $\lambda \in \mathbb{Z}_{q_*}^*$  and  $z \in \mathbb{Z}_{q_2}^*$ )

$$\begin{aligned} \sum_{a_1, a_2 \in A_*} e_{q_2} \left( \frac{\lambda z}{a_1 - a_2} \right) &= \sum_{\xi_1, \xi_2 \in \mathbb{Z}_{q_2}} \eta_{q_2}(\xi_1) \eta_{q_2}(\xi_2) e_{q_2} \left( \frac{\lambda z}{\xi_1 - \xi_2} \right) = q_2^{-1} \sum_{\xi \in \mathbb{Z}_{q_2}} |\widehat{\eta}_{q_2}(\xi)|^2 K_{q_2}(\lambda z, \xi) \\ &\leq 2\sqrt{q_2} \tau(q_2) \|\eta_{q_2}\|_2^2. \end{aligned} \quad (16)$$

In the last line we have applied the well-known bound for the Kloosterman sum and the Parseval identity. Now to estimate  $\|\eta_{q_2}\|_2^2$  we use the regularity property of  $A_*$  and derive

$$\|\eta_{q_2}\|_2^2 \leq \|\eta_{q_2}\|_\infty \|\eta_{q_2}\|_1 \leq \frac{|A_*|^2}{q_2^{1-\varepsilon}}. \quad (17)$$

Further let us obtain a lower bound for divisors  $q_2$ . Since  $|\pi_{Q_1}(A_1)| = 1$ , it follows that for all  $q_1 | Q_1$ , we have

$$\frac{1}{q_1} \sum_{\xi \in \mathbb{Z}_{q_1}} |\widehat{\eta}_{q_1}(\xi)|^2 K_{q_1}(\lambda z, \xi) = \frac{|A_*|^2}{q_1} \sum_{\xi \in \mathbb{Z}_{q_1}} K_{q_1}(\lambda z, \xi) = 0.$$

Thus one can see that summations in (14) is taken over  $q_2 \geq M$ . Choosing  $\varepsilon = 1/4$ , say, and using the last fact, we get in view of the Parseval identity that

$$\sigma \ll M^{-1/4} \frac{|A_*|^2}{q_*} \sum_{q_2 | q_*, q_2 > 1} \sum_{z \in \mathbb{Z}_{q_2}^*} |\widehat{B}_*(z q_* q_2^{-1})|^2 \leq M^{-1/4} |A_*|^2 |B_*|.$$

Returning to (13), we obtain a contradiction provided  $|B_*| \gg q_* M^{-1/4}$ . In other words, we have for a certain  $s \geq 0$ ,  $\alpha M^{s/4} > 1$  that

$$|B| M^{s/4} \ll q M^{-1/4}$$

and this implies  $M \ll \beta^{-4}$ . Thus in view of our restriction to the divisors of  $q_*$ , the condition  $\alpha \geq \beta$ , the first bound for  $Q_1$  from (12), and the bound for  $s$ , which follows from Lemma 2, we get

$$d \ll M^{\pi(M)} \exp(O(\log^2(1/\alpha)/\log(1/\beta))) \ll \exp(O(\beta^{-4}))$$

as required.

Now let  $\lambda \in \mathbb{Z}_{q_*}$  be an arbitrary element. Write  $\lambda = q' \lambda'$ , where  $q' | q$  and  $\lambda' \in \mathbb{Z}_{q_*/q'}^*$ . Using the Dirichlet principle, choose a subset of  $B' \subseteq B_*$  of density at least  $\beta$  such that all elements of

a shift of  $B'$  are divisible by  $q'$ . Then our inclusion can be rewritten as  $\lambda' \in (A_* - A_*)(B' - B')$  modulo  $q_*/q'$  and we can apply the arguments above replacing module  $q_*$  to  $q_*/q'$ .

To obtain (11) we use the second bound for  $Q_1$  from (12) and derive as above

$$d \ll M^{\omega(q)} \exp(O(\log(1/\beta))) \ll \exp(O(\omega(q) \log(1/\beta))).$$

This completes the proof.  $\square$

As one can see from the proof of Theorem 6 that the constant four in (9) can be decreased to  $2 + o(1)$  but we leave such calculations to the interested reader.

### 3 On the general case

In [5, Problem 2] A. Fish considered a more general two-dimensional case (actually, in his paper he had to deal with even more general dynamical setting) and formulated the following problem.

**Problem.** *Let  $q$  be a positive number and  $\mathcal{A}, \mathcal{B} \subseteq \mathbb{Z}_q^2$  be sets,  $|\mathcal{A}| = \alpha q^2$ ,  $|\mathcal{B}| = \beta q^2$ , and suppose that  $\alpha \geq \beta$ . Prove that in the case  $\mathcal{A} = \mathcal{B}$  for a certain function  $F$  there is  $d|q$  such that  $d \leq F(\beta)$  and*

$$d \cdot \mathbb{Z}_q \subseteq \{(a_1 - b_1)(a_2 - b_2) : (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B}\}, \quad (18)$$

provided  $\beta$  is sufficiently large.

In this section we study the number  $N_{\mathcal{A}, \mathcal{B}}(\lambda)$  of the solutions to the equation

$$(a_1 - b_1)(a_2 - b_2) \equiv \lambda \pmod{q}, \quad (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B}. \quad (19)$$

and give a partial answer to the problem above. We consider the squarefree case for simplicity and emphasize one more time that our sets  $\mathcal{A}, \mathcal{B}$  are arbitrary (in the case of Cartesian products and squarefree  $q$  one can apply other methods, see [10]). Also, in the case of prime  $q$  we obtain a result of Vinh-type [21], see asymptotic formula (22) below.

**Theorem 7** *Let  $q$  be a squarefree number,  $\mathcal{A}, \mathcal{B} \subseteq \mathbb{Z}_q^2$  be sets,  $|\mathcal{A}| = \alpha q^2$ ,  $|\mathcal{B}| = \beta q^2$ , and suppose that  $\alpha \geq \beta$ . Then*

$$d \cdot \mathbb{Z}_q^* \subseteq \{(a_1 - b_1)(a_2 - b_2) : (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B}\} \quad (20)$$

with

$$d \ll \exp(O(\omega(q) \log \omega(q) - \log \beta)).$$

In particular, for  $\mathcal{A} = \mathcal{B}$  one has with the same  $d$  that

$$d \cdot \mathbb{Z}_q \subseteq \{(a_1 - b_1)(a_2 - b_2) : (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B}\}. \quad (21)$$

In the case when  $q$  is a prime number, we have

$$\left| N_{\mathcal{A}, \mathcal{B}}(\lambda) - \frac{|\mathcal{A}||\mathcal{B}|}{q} \right| < 4q^{7/8} \sqrt{|\mathcal{A}||\mathcal{B}|}. \quad (22)$$

In particular, equality (18) holds for  $|\mathcal{A}||\mathcal{B}| \geq 16q^{15/4}$  and  $d = 1$ .

Proof. We start with (20). The proof follows the arguments of the proof of Theorem 6 and thus we use the notation from this result. In particular, writing  $q = p_1^{\rho_1} \dots p_t^{\rho_t}$ ,  $t = \omega(q)$  with  $\rho_j = 1$ ,  $j \in [t]$  we define  $Q_1 = \prod_{j=1}^s p_j$  such that (12) holds and further we take  $\lambda \in \mathbb{Z}_q^*$ . The only difference is that one should use Lemma 2 with  $n = 2$  to regularize the two-dimensional set  $\mathcal{A}$  and let  $\varepsilon = 1/4$ . For a moment we assume that  $M \geq 100t^2$ , say, and we will choose the parameter  $M$  later. Finally, with some abuse of the notation we do not use new letters  $\mathcal{A}_*, \mathcal{B}_*, q_*$  below but the old ones  $\mathcal{A}, \mathcal{B}$  and  $q$  (in other words, one can think that  $\mathcal{A}$  is a regularized set already). Also, we utilize the fact that  $\mathbb{Z}_q = \mathbb{Z}_{p_1^{\rho_1}} \times \dots \times \mathbb{Z}_{p_t^{\rho_t}} = \mathbb{Z}_{p_1} \times \dots \times \mathbb{Z}_{p_t}$  thanks the Chinese remainder theorem.

Now for  $a = (a_1, a_2)$  and  $b = (b_1, b_2)$  let us write  $I(a, b) = 1$  if the pair  $a, b$  satisfies (19) and  $I(a, b) = 0$  otherwise. Then clearly,

$$N_{\mathcal{A}, \mathcal{B}}(\lambda) = \sum_{a \in \mathcal{A}, b \in \mathcal{B}} I(a, b). \quad (23)$$

Without loosing of the generality we assume that  $\lambda = 1$ . Obviously,  $I(a, b) = I(b, a)$  and we can rewrite the matrix  $I(a, b)$  as  $I(a, b) = \sum_{j=1}^{q^2} \mu_j u_j(a) \bar{u}_j(b)$ , where  $\mu_j$  are eigenvalues and  $u_j(x)$  are correspondent normalized eigenfunctions of  $I$ . One can easily check that  $u_1(x) = q^{-1}(1, \dots, 1)$ ,  $\|u_1\|_2 = 1$  and  $\mu_1 = |\mathbb{Z}_q^*| = \varphi(q)$ . Writing  $I'(a, b) = I(a, b) - \mu_1 u_1(a) \bar{u}_1(b)$ , we obtain

$$N_{\mathcal{A}, \mathcal{B}}(\lambda) - \frac{|\mathcal{A}| |\mathcal{B}| \varphi(q)}{q^2} = \sum_{a \in \mathcal{A}, b \in \mathcal{B}} I'(a, b) := N'_{\mathcal{A}, \mathcal{B}}(\lambda). \quad (24)$$

By the Cauchy–Schwarz inequality, we get

$$N'_{\mathcal{A}, \mathcal{B}}(\lambda)^2 \leq |\mathcal{B}| \sum_{a, a' \in \mathcal{A}} \sum_b I'(a, b) I'(a', b) = |\mathcal{B}| \sum_{a, a' \in \mathcal{A}} (I')^2(a, a') := |\mathcal{B}| \cdot \sigma. \quad (25)$$

Here  $(I')^2$  is the second power of the matrix  $I'$ . Similarly,  $I^2(a, a') = \sum_b I(a, b) I(a', b)$  and the last quantity coincides with the number of the solutions to the equation

$$a_2 - a'_2 = \frac{a'_1 - a_1}{(a_1 + x)(a'_1 + x)}, \quad (26)$$

where  $b = (x, y)$ ,  $a = (a_1, a_2)$  and  $a' = (a'_1, a'_2)$ . Assume that  $a \neq a'$  and rewrite our equation (26) as

$$x^2 + (a_1 + a'_1)x + a_1 a'_1 + \frac{a_1 - a'_1}{a_2 - a'_2} = 0, \quad (27)$$

and its discriminant is  $D'(a, a') := (a_1 - a'_1)(a_2 - a'_2)^{-1}[(a_1 - a'_1)(a_2 - a'_2) - 4]$ . Notice that if  $a = a'$ , then we have  $\varphi(q)$  solutions to equation (26). By  $\chi_p$  denote the Legendre symbol modulo a prime  $p$  and let  $\chi_0$  be the main character (modulo  $p$ ). We have the identity  $\chi_p(x^{-1}) = \chi_p(x)$ ,  $x \in \mathbb{Z}_p^*$  and hence  $\chi_p(D'(a, a')) = \chi_p((a_1 - a'_1)(a_2 - a'_2)[(a_1 - a'_1)(a_2 - a'_2) - 4]) := \chi_p(D(a, a'))$ . In view of the Chinese remainder theorem, and our choice of the regularized set  $\mathcal{A}$  one has

$$I^2(a, a') = \prod_{j=s+1}^t (\chi_{p_j}(D(a, a')) + \chi_0(D(a, a')) + (p_j - 1)\delta_{p_j}(a_1 - a'_1, a_2 - a'_2)) \quad (28)$$

$$= \mathcal{E}(a, a') + \prod_{j=s+1}^t (\chi_0(D(a, a')) + (p_j - 1)\delta_{p_j}(a_1 - a'_1, a_2 - a'_2)) = \mathcal{E}(a, a') + \mathcal{E}'(a, a'), \quad (29)$$

where for a positive integer  $m$  we have put  $\delta_m(z, w) = 1$  if  $z \equiv w \equiv 0 \pmod{m}$ , and zero otherwise. Equivalently, writing  $T$  for the segment  $[s+1, t]$ , one has

$$\begin{aligned} \mathcal{E}(a, a') &= \sum_{\emptyset \neq S \subseteq T} \prod_{j \notin S} (\chi_0(D(a, a')) + (p_j - 1)\delta_{p_j}(a_1 - a'_1, a_2 - a'_2)) \cdot \prod_{j \in S} \chi_{p_j}(D(a, a')) \\ &= \sum_{\emptyset \neq S \subseteq T} \prod_{j \notin S} w_{p_j}(a, a') \cdot \prod_{j \in S} \chi_{p_j}(D(a, a')). \end{aligned}$$

Notice that  $\mathcal{E}(a, a) = 0$ . From (28), (29), it follows that  $\mathcal{E}u_1 = 0$ . Indeed, we know that  $I^2u_1 = \mu_1^2u_1 = \varphi^2(q)u_1$  and

$$\sum_a \prod_{j=s+1}^t (\chi_0(D(a, a')) + (p_j - 1)\delta_{p_j}(a_1 - a'_1, a_2 - a'_2)) \quad (30)$$

$$= \prod_{j=s+1}^t \left( \sum_{z, w \in \mathbb{Z}_{p_j}} \chi_0((zw)^2 - 4zw) + p_j - 1 \right) \quad (31)$$

$$= \prod_{j=s+1}^t \left( (p_j - 1) \sum_{z \in \mathbb{Z}_{p_j}} \chi_0(z^2 - 4z) + p_j - 1 \right) = \prod_{j=s+1}^t (p_j - 1)^2 = \varphi^2(q). \quad (32)$$

Hence in very deed  $\mathcal{E}u_1 = 0$  and thus

$$\sigma = \langle (I')^2 \mathcal{A}, \mathcal{A} \rangle = \langle (I')^2 f_{\mathcal{A}}, f_{\mathcal{A}} \rangle = \langle I^2 f_{\mathcal{A}}, f_{\mathcal{A}} \rangle = \langle \mathcal{E} \mathcal{A}, \mathcal{A} \rangle + \langle \mathcal{E}' f_{\mathcal{A}}, f_{\mathcal{A}} \rangle, \quad (33)$$

where  $f_{\mathcal{A}}(a) = \mathcal{A}(a) - \langle \mathcal{A}, u_1 \rangle u_1(a)$ ,  $\sum_a f_{\mathcal{A}}(a) = 0$ . Let us estimate the term  $r := \langle \mathcal{E}' f_{\mathcal{A}}, f_{\mathcal{A}} \rangle$  rather roughly. Since the function  $f_{\mathcal{A}}$  is orthogonal to  $u_1$  and  $\|f_{\mathcal{A}}\|_{\infty} \leq 1$ , it follows that

$$\begin{aligned} |r| &= \left| \sum_{a, a'} f_{\mathcal{A}}(a) f_{\mathcal{A}}(a') \prod_{j=s+1}^t (1 - \delta_{p_j}(D(a, a')) + (p_j - 1)\delta_{p_j}(a_1 - a'_1, a_2 - a'_2)) \right| \\ &\leq \sum_{\emptyset \neq S \subseteq T} \left| \sum_{a, a'} f_{\mathcal{A}}(a) f_{\mathcal{A}}(a') \prod_{j \in S} (-\delta_{p_j}(D(a, a')) + (p_j - 1)\delta_{p_j}(a_1 - a'_1, a_2 - a'_2)) \right| \\ &\leq 2|\mathcal{A}|q^2 \sum_{n=1}^{t-s} \sum_{S \subseteq T, |S|=n} \prod_{j \in S} \left( \frac{3}{p_j} + \frac{p_j - 1}{p_j^2} \right) \leq 2|\mathcal{A}|q^2 \sum_{n=1}^{t-s} \binom{t-s}{n} \left( \frac{4}{M} \right)^n \\ &\leq 10|\mathcal{A}|q^2 t M^{-1}. \end{aligned} \quad (34)$$



Now returning to the definition of the operator  $\mathcal{E}(a, a')$ , recalling estimate (25) and using the Cauchy–Schwarz inequality, we obtain

$$\begin{aligned} \sigma^2 \leq |\mathcal{A}| \sum_{a, a' \in \mathcal{A}} \sum_{x, y} \sum_{\emptyset \neq S_1, S_2 \subseteq T} \prod_{i \in S_1, j \in S_2} \chi_{p_i}(D((x, y), (a_1, a_2))) \chi_{p_j}(D((x, y), (a'_1, a'_2))) \\ \prod_{i \notin S_1, j \notin S_2} w_{p_i}((x, y), (a_1, a_2)) w_{p_j}((x, y), (a'_1, a'_2)). \end{aligned} \quad (35)$$

The term with  $a \equiv a' \pmod{q}$  gives us a contribution at most  $4^t |\mathcal{A}| q^2$  into the last sum (see (30)–(32) to estimate  $\|w_{p_j}\|_1$  for  $j \notin S$  and use the trivial fact that  $\|\chi_p\|_\infty \leq 1$  to bound the rest). Now let  $a \not\equiv a' \pmod{q}$  but  $a \equiv a' \pmod{q_*}$  with maximal  $q_* | q$ . Thus  $q_* \neq q$  and  $Q_1 | q_*$ . We can write  $q_* = q_*(W) = Q_1 \prod_{j \in W} p_j$  for a certain (possibly empty) set  $W \subseteq T$ . Let us say that all primes  $p$  such that  $p | (q/q_*)$  (that is,  $p | q$  and  $p \notin W$ ) are *good*. In particular, for all good primes  $p$  one has  $p > M$ . Now for a good prime  $p$  the sum above  $\sum_{x, y \pmod{\mathbb{Z}_p}} \chi_p(D(x, y), (a_1, a_2))$  (or, analogously, the sum  $\sum_{x, y \pmod{\mathbb{Z}_p}} \chi_p(D(x, y), (a'_1, a'_2))$ ) is either at most  $3p^{3/2}$  by Weil, or the sum over  $y$  is  $p$  if  $\frac{2}{x-a_1} + a_2 = \frac{2}{x-a'_1} + a'_2$  modulo  $p$ . The last equation is nontrivial one by our choice of  $p$  hence it has at most two solutions and thus in any case the sum over  $x, y \pmod{\mathbb{Z}_p}$  is at most  $3p^{3/2} < 3p^2/\sqrt{M}$ . Further we split the sets  $S_1, S_2$  as  $S_1 = S_1^* \sqcup G_1$ ,  $S_2 = S_2^* \sqcup G_2$ , where (possibly empty) sets  $G_1, G_2$  correspond to good primes and the sets  $S_1^* \subseteq W$ ,  $S_2^* \subseteq W$  correspond to the divisors of  $q_*(W)$ . Since  $S_1, S_2 \neq \emptyset$ , it follows that either  $G_1 \cup G_2 \neq \emptyset$  or  $S_1^*, S_2^* \neq \emptyset$ . Using the notation as in (15), namely,

$$\eta_{\tilde{q}}(\xi) := |\{a \in \mathcal{A} : a \equiv \xi \pmod{\tilde{q}}\}|, \quad \tilde{q} | q, \xi \in \mathbb{Z}_{\tilde{q}}^2, \quad (36)$$

we see that the number of pairs  $a \equiv a' \pmod{\tilde{q}}$  is exactly  $\|\eta_{\tilde{q}}\|_2^2$  for any  $\tilde{q} | q$  and one can use bound (17) to estimate the last quantity. Now recalling inequality (7) and splitting sum (35) according the case  $W \neq \emptyset$  or not, we get

$$\begin{aligned} \sigma^2 |\mathcal{A}|^{-1} \leq \\ 4^t |\mathcal{A}| q^2 + q^2 \sum_{\emptyset \neq W \subseteq T} \sum_{a, a' \in \mathcal{A}, a \equiv a' \pmod{q(W)}} 4^{|W|} + q^2 \sum_{a, a' \in \mathcal{A}} \sum_{n+m \geq 1} \binom{t-s}{n} \binom{t-s}{m} \left(\frac{3}{\sqrt{M}}\right)^{n+m} \\ \leq 4^t |\mathcal{A}| q^2 + q^2 |\mathcal{A}|^2 \sum_{\emptyset \neq W \subseteq T} 4^{|W|} M^{-3|W|/4} + 4q^2 |\mathcal{A}|^2 t M^{-1/2} \ll q^2 |\mathcal{A}|^2 t M^{-1/2}. \end{aligned} \quad (37)$$

Using (23), (24), (25), (34) and the Cauchy–Schwarz inequality, we get

$$N_{\mathcal{A}, \mathcal{B}}(\lambda) - \frac{|\mathcal{A}| |\mathcal{B}| \varphi(q)}{q^2} \ll (tM^{-1/2})^{1/4} \cdot |\mathcal{A}|^{3/4} \sqrt{|\mathcal{B}|q} + (tM^{-1})^{1/2} \cdot \sqrt{|\mathcal{A}| |\mathcal{B}|} q. \quad (38)$$

We have  $\varphi(q) \gg q/\log t$  and hence after some calculations we see that  $N_{\mathcal{A}, \mathcal{B}}(\lambda) > 0$  provided  $M \gg t^2 \beta^{-6} \log^8 t$ . As in Theorem 6, one has  $Q_1 \leq M^t$  and thus

$$d \ll \exp(t \log M) = \exp(O(t \log t - \log \beta)).$$

In the case of prime  $q$  the argument is even simpler because one do not need the regularization, the second term in (37) plus the quantity  $r$  is negligible, see estimate (34). Finally, let  $\mathcal{A} = \mathcal{B}$  and if  $\lambda \notin \mathbb{Z}_q^*$ , then write  $\lambda = q'\lambda'$ , where  $q'|q$  and  $\lambda' \in \mathbb{Z}_{q^*/q'}^*$ . Using the Dirichlet principle, choose a subset of  $\mathcal{A}' \subseteq \mathcal{A}$  of density at least  $\alpha$  such that  $|\pi_{q'}(\mathcal{A}')| = 1$ . Then the required inclusion (21) can be rewritten as

$$\lambda' \in \{(a_1 - b_1)(a_2 - b_2) : (a_1, a_2), (b_1, b_2) \in \mathcal{A}\}$$

and we can apply the arguments above replacing  $q_*$  to  $q^*/q'$ . This completes the proof.  $\square$

**Remark 8** *Of course, inclusion (21) does not hold for  $\mathcal{A} \neq \mathcal{B}$ , just take  $\mathcal{A} = (d \cdot \mathbb{Z}_q) \times (d \cdot \mathbb{Z}_q)$  and  $\mathcal{B} = (d \cdot \mathbb{Z}_q + 1) \times (d \cdot \mathbb{Z}_q + 1)$  for an arbitrary  $d|q$ ,  $1 < d \ll 1$ . Also, the author thinks that the error term in (22) can be improved but this weaker bound is enough for us to resolve our equation for sets of positive densities.*

**Remark 9** *The attentive reader may be alerted that we have two different main terms in (13) and in (38). Nevertheless, they are asymptotically the same due to the fact that in (38) our parameter  $M$  depends on growing quantity  $\omega(q)$ .*

Similarly, we obtain an affirmative answer to [5, Problem 1] in the case of squarefree  $q$ . By  $M_{\mathcal{A}, \mathcal{B}}(\lambda)$  denote the number of the solutions to the equation

$$(a_1 - b_1)^2 - (a_2 - b_2)^2 \equiv \lambda \pmod{q}, \quad (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B}. \quad (39)$$

**Theorem 10** *Let  $q$  be a squarefree number,  $\mathcal{A}, \mathcal{B} \subseteq \mathbb{F}_q^2$  be sets,  $|\mathcal{A}| = \alpha q^2$ ,  $|\mathcal{B}| = \beta q^2$ , and suppose that  $\alpha \geq \beta$ . Then*

$$d\mathbb{Z}_q^* \subseteq \{(a_1 - b_1)^2 - (a_2 - b_2)^2 : (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B}\} \quad (40)$$

with

$$d \ll \exp(O(\omega(q) \log \omega(q) - \log \beta)).$$

In particular, for  $\mathcal{A} = \mathcal{B}$  one has with the same  $d$  that

$$d\mathbb{Z}_q \subseteq \{(a_1 - b_1)^2 - (a_2 - b_2)^2 : (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B}\}. \quad (41)$$

In the case when  $q$  is a prime number one has

$$M_{\mathcal{A}, \mathcal{B}}(\lambda) - \frac{|\mathcal{A}||\mathcal{B}|}{q} < 4q^{7/8} \sqrt{|\mathcal{A}||\mathcal{B}|}. \quad (42)$$

**Proof.** The argument differs from the proof of Theorem 7 in some unimportant details only, so we use the notation from the former result. Indeed, for  $a = (a_1, a_2)$  and  $b = (b_1, b_2)$  we write  $\tilde{I}(a, b) = 1$  if the pair  $a, b$  satisfies (39) and  $\tilde{I}(a, b) = 0$  otherwise. Calculating  $\tilde{I}^2(a, a')$ , we arrive to the equation

$$a_1^2 - (a_1')^2 + 2(a_1' - a_1)x - a_2^2 + (a_2')^2 + 2(a_2 - a_2')y = 0 \quad (43)$$

and hence we can find  $x$  via  $y$  or  $y$  via  $x$ , provided  $a \not\equiv a' \pmod{q}$ . Assuming that  $a'_2 \neq a_2$ , say, we derive

$$y = \frac{(a'_1)^2 - a_1^2 + a_2^2 - (a'_2)^2}{2(a_2 - a'_2)} + \frac{a_1 - a'_1}{a_2 - a'_2} \cdot x = s + tx,$$

and hence substituting the last expression into (39) and computing the discriminant  $\tilde{D}(a, a')$  (without loss of the generality, we put  $\lambda = 1$ ), one obtains

$$\begin{aligned} \tilde{D}(a, a') &= (t(a_2 - s) - a_1)^2 + (1 - t)^2(1 + (a_2 - s)^2 - a_1^2) \\ &= 2t(t - 1)(a_2 - s)^2 - 2a_1t(a_2 - s) + (1 - t)^2(1 - a_1^2) + (a_2 - s)^2 + a_1^2. \end{aligned} \quad (44)$$

As in the proof of Theorem 7 we consider  $\mathcal{E}(a, a')$ , take good primes and so on. The first eigenvalue  $\mu_1$  equals the number of the solutions to the equation  $x^2 - y^2 \equiv 1 \pmod{q}$ , that is,  $\varphi(q)$  again. Also,  $\tilde{I}^2(a, a) = \mu_1$  and for  $a \neq a'$  the quantity  $\tilde{I}^2(a, a')$  expressed exactly as in (28) (with another discriminant  $\tilde{D}$ , of course) and thus one can check that  $\mathcal{E}u_1$  vanishes making calculations as in (30)—(32). Further as in Theorem 7 we apply the standard Weil bound to estimate the sum of characters. For any good prime  $p$  it gives us a nontrivial bound of the form  $O(p^{3/2}) = O(p^2/\sqrt{M})$  and hence we obtain (40) and thus (41) by the same argument as at the end of Theorem 7 (one can check or see below that all obtained varieties are non-degenerated). Finally, to get (42) we need to estimate

$$\sum_{a, a' \in \mathcal{A}} \sum_{x, y} \chi_q(\tilde{D}((x, y), (a_1, a_2))) \chi_q(\tilde{D}((x, y), (a'_1, a'_2)))$$

and by the Weil estimate it is at most  $20q^{3/2}$ , say, excluding the case  $\tilde{D}((x, y), (a_1, a_2))$  is proportional to  $\tilde{D}((x, y), (a'_1, a'_2))$ . In particular, it means that the coefficients of these polynomials are proportional ones and using (44) and comparing the coefficients before the highest degrees in  $x$ , say, we get  $\frac{a_2 - 2a_1 - y}{(y - a_2)^4} = \frac{a'_2 - 2a'_1 - y}{(y - a'_2)^4}$ . Again, thanks to  $a \neq a'$  we see that this equation is nontrivial one and hence it has at most four solutions. It follows that our sum is at most  $4q$  in this case. Thus as in (37), (38) we have

$$M_{\mathcal{A}, \mathcal{B}}(\lambda) - \frac{|\mathcal{A}||\mathcal{B}|\varphi(q)}{q^2} \leq 3(q^{3/2}|\mathcal{A}|)^{1/4} \sqrt{|\mathcal{A}||\mathcal{B}|} \leq 3q^{7/8} \sqrt{|\mathcal{A}||\mathcal{B}|}.$$

This completes the proof.  $\square$

**Remark 11** *We have used a direct way of the proof of Theorem 10 above, another approach is to notice that  $\tilde{I}(a, b) = I(ga, gb)$ , where the linear transformation  $g$  is given by the formula  $g(x, y) = (x + y, x - y)$ . After that one can apply Theorem 7 with the sets  $g^{-1}(\mathcal{A})$ ,  $g^{-1}(\mathcal{B})$ .*

## 4 On an application of group actions

In this section we discuss another approach to results of Fish-type, namely, we consider an intermediate situation between Theorems 6 and 7: our set  $\mathcal{A} \subseteq \mathbb{Z}_q^2$  is an arbitrary but the set  $\mathcal{B} \subseteq \mathbb{Z}_q^2$  is a Cartesian product. In this case one can deal with rather general  $q$  (and not just

squarefree). For simplicity, we do not do any regularization as in the previous section immediately assuming that all prime factors of  $q$  are large.

In the proof we follow the methods from [3] and [16].

**Theorem 12** *Let  $q$  be a positive odd integer, and  $\mathcal{A}, \mathcal{B} \subseteq \mathbb{Z}_q^2$  be sets,  $|\mathcal{A}| = \delta q^2$ ,  $\mathcal{B} = A \times B$ ,  $|A| = \alpha q$ ,  $|B| = \beta q$ . Suppose that all prime divisors of  $q$  are at least  $M$ , where*

$$M \geq C_1 \tau(q) \delta^{-2} (\sqrt{\alpha\beta} (\log \log q)^{-1})^{-C_2},$$

and  $C_1, C_2 > 0$  are absolute constants. Then

$$\mathbb{Z}_q^* \subseteq \{(a_1 - b_1)(a_2 - b_2) : (a_1, a_2) \in \mathcal{A}, (b_1, b_2) \in \mathcal{B}\}. \quad (45)$$

*Proof.* Let  $q = p_1^{\rho_1} \dots p_t^{\rho_t}$ , where  $p_j$  are different odd primes and  $\rho_j$  are positive integers. By our assumption  $p_j \geq M$  for all  $j \in [t]$ . Without losing of the generality one can take  $\lambda = -1$  in formula (19). Recall that  $\mathrm{SL}_2(\mathbb{Z}_q)$  acts on  $\mathbb{Z}_q$  via Möbius transformations:  $x \rightarrow gx = \frac{ax+b}{cx+d}$ , where  $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  (for composite  $q$  the equivalence is taken over  $\mathbb{Z}_q^*$ , of course). Since  $\mathcal{B} = A \times B$  we can rewrite our equation (19) as

$$a = gb, \quad a \in A, b \in B, g \in G, \quad (46)$$

where  $G \subset \mathrm{SL}_2(\mathbb{Z}_q)$  is the set of matrices of the form

$$g = \begin{pmatrix} -\alpha & \alpha\beta + 1 \\ -1 & \beta \end{pmatrix}, \quad (\alpha, \beta) \in \mathcal{A},$$

see [16, Section 5] or just make a direct calculation. Clearly,  $|G| = |\mathcal{A}|$ . Further by [16, Lemma 15] the *multiplicative energy*  $\mathbf{E}(G)$  of the set  $G$ , that is,

$$\mathbf{E}(G) = |\{(g_1, g_2, g_3, g_4) \in G \times G \times G \times G : g_1 g_2^{-1} = g_3 g_4^{-1}\}|$$

coincides with the number of the solutions to the system

$$\beta_1 - \beta_2 = \beta_3 - \beta_4 := s, \quad s(\alpha_1 - \alpha_3) = s(\alpha_2 - \alpha_4) = 0, \quad \alpha_1 - \alpha_2 - \alpha_1 \alpha_2 s = \alpha_3 - \alpha_4 - \alpha_3 \alpha_4 s,$$

where  $(\alpha_i, \beta_i) \in \mathcal{A}$ ,  $i \in [4]$ . Let  $s = ds'$ , where  $d$  is a divisor of  $q$  and  $s'$  is coprime to  $q$ . Taking  $(\alpha_1, \beta_1), (\alpha_4, \beta_4) \in \mathcal{A}$ , we find  $\beta_2, \beta_3$  from the first equation and  $\alpha_2, \alpha_3$  modulo  $q/d$  from the second one. Also, using  $\alpha_3$  we can reconstruct  $\alpha_2$  from the third equation, provided  $d > 1$ . In other words, for fixed  $d$  there are  $q/d$  possibilities for  $s'$  and  $d$  possibilities for  $\alpha_3$ . Finally, if  $d = 1$ , then we have at most  $q|G|^2$  solutions. Thus we obtain the bound

$$\mathbf{E}(G) \leq |G|^2 \sum_{d|q} \frac{q}{d} \cdot d \leq \tau(q) q |G|^2. \quad (47)$$

Now let us say a few words about representations of the group  $\mathrm{SL}_2(\mathbb{Z}_q)$ , see [3, Sections 7, 8]. First of all, for any irreducible representation  $\rho_q$  of  $\mathrm{SL}_2(\mathbb{Z}_q)$  we have  $\rho = \rho_q = \rho_{p_1^{e_1}} \otimes \cdots \otimes \rho_{p_t^{e_t}}$  and hence it is sufficient to understand the representation theory for  $\mathrm{SL}_2(\mathbb{Z}_{p^n})$ , where  $p$  is a prime number and  $n$  is a positive integer. Now by [3, Lemma 7.1] we know that for any odd prime the dimension of any faithful irreducible representation of  $\mathrm{SL}_2(\mathbb{Z}_{p^n})$  is at least  $2^{-1}p^{n-2}(p-1)(p+1)$ . For an arbitrary  $r \leq n$  we can consider the natural projection  $\pi_r : \mathrm{SL}_2(\mathbb{Z}_{p^n}) \rightarrow \mathrm{SL}_2(\mathbb{Z}_{p^r})$  and let  $H_r = \mathrm{Ker} \pi_r$ . One can show that the set  $\{H_r\}_{r \leq n}$  gives all normal subgroups of  $\mathrm{SL}_2(\mathbb{Z}_{p^n})$  and hence any nonfaithful irreducible representation arises as a faithful irreducible representation of  $\mathrm{SL}_2(\mathbb{Z}_{p^r})$  for a certain  $r < n$ . Anyway, we see that the multiplicity (dimension)  $d_\rho$  of any nontrivial irreducible representation  $\rho$  of  $\mathrm{SL}_2(\mathbb{Z}_{p^n})$  is at least  $p/3 \geq M/3$ .

Applying estimate (47), using the formula for  $\mathbf{E}(G)$  via the representations and taking into account the obtained lower bound for the multiplicities of the representations, we get

$$\frac{M \|\widehat{G}\|_{op}^4}{3|\mathrm{SL}_2(\mathbb{Z}_q)|} \leq \frac{1}{|\mathrm{SL}_2(\mathbb{Z}_q)|} \sum_{\rho} d_\rho \|\widehat{G}(\rho) \widehat{G}^*(\rho)\|^2 = \mathbf{E}(G) \leq \tau(q)q|G|^2,$$

and hence

$$\|\widehat{G}\|_{op} \leq |G| \cdot \left( \frac{3\tau(q)}{M\delta^2} \right)^{1/4} := \frac{|G|}{K}, \quad (48)$$

where by  $\|\widehat{G}\|_{op}$  we have denoted the maximum of the operator norm of matrices  $\widehat{G}(\rho)$  for all nontrivial representations  $\rho$  and  $\|\cdot\|$  is the usual Hilbert–Schmidt norm. Thanks to our choice of  $M$  one can see that bound (48) is nontrivial, that is,  $K > 1$ . Returning to (46) and using the standard scheme, e.g., see [16, Lemma 13, Section 5 and Section 6], we obtain

$$N_{\mathcal{A}, \mathcal{B}}(\lambda) - \frac{|A||B||G|\varphi(q)}{q^2} \leq \sqrt{|A||B|}|G|q^{-1/k},$$

where  $k \sim \log q / \log K$ . Hence  $N_{\mathcal{A}, \mathcal{B}}(\lambda) > 0$ , provided  $K \gg (\sqrt{\alpha\beta}(\log \log q)^{-1})^{-O(1)}$ . The last condition is equivalent to  $M \gg \tau(q)\delta^{-2}(\sqrt{\alpha\beta}(\log \log q)^{-1})^{-O(1)}$ . This completes the proof.  $\square$

## 5 On the covering numbers of difference sets

Let us recall the definition of the covering number of a set, e.g., see [1].

**Definition 13** *Let  $\mathbf{G}$  be a finite abelian group with the group operation  $+$ , and let  $A \subseteq \mathbf{G}$  be a set. We write*

$$\mathrm{cov}^+(A) = \mathrm{cov}(A) = \min\{|X| : X \subseteq \mathbf{G}, A + X = \mathbf{G}\}$$

*and the quantity  $\mathrm{cov}^+(A)$  is called the (additive) covering number of  $A$ .*

Having a finite ring  $\mathcal{R}$  with two operations  $+$ ,  $\times$  we underline which covering number we use, writing  $\mathrm{cov}^+$  or  $\mathrm{cov}^\times$ . It is known [1, Corollary 3.2] that for any set  $A \subseteq \mathbf{G}$  one has  $\mathrm{cov}^+(A) = O\left(\frac{|\mathbf{G}|}{|A|} \log |A|\right)$  and the last bound is tight. In this section we study difference sets  $A - A$ ,  $A \subseteq \mathbb{Z}_q$  and show that  $\mathrm{cov}^\times(A - A)$  is always small. First of all, let us make a remark about a connection between  $\mathrm{cov}^+$  and  $\mathrm{cov}^\times$  in a ring  $\mathcal{R}$ .

**Proposition 14** *Let  $\mathcal{R}$  be a finite ring, and  $S \subseteq \mathcal{R}$  be a set. Then*

$$\text{cov}^\times(S - S) \leq \text{cov}^+(S), \quad (49)$$

*provided all numbers  $1, \dots, \text{cov}^+(S)$  belong to  $\mathcal{R}^*$ .*

*Proof.* Let  $S + X = \mathbb{Z}_q$  and  $|X| = \text{cov}^+(S) := k$ . For any  $g \in \mathbb{Z}_q$  consider  $kg$ , where  $j = 0, 1, \dots, k$ . By the pigeonhole principle there are different  $j_1 \neq j_2$  such that  $j_1g \in S + x$  and  $j_2g \in S + x$  with the same  $x \in X$ . It implies that  $(j_1 - j_2)g \in S - S$  and hence  $g \in (j_1 - j_2)^{-1}(S - S)$ , provided  $(j_1 - j_2)^{-1} \in \mathcal{R}^*$ . It remains to notice that  $[-k, k]^{-1} \cdot (S - S) = [k]^{-1} \cdot (S - S)$ . This completes the proof.  $\square$

By the well-known consequence of the Ruzsa covering lemma [20, Section 2.4], we have for any finite group  $\mathbf{G}$  and a set  $A \subseteq \mathbf{G}$  that for a certain set  $Z \subseteq \mathbf{G}$  one has

$$\mathbf{G} \subseteq A - A + Z, \quad |Z| \leq |\mathbf{G}|/|A|. \quad (50)$$

In particular, it means that  $\text{cov}^+(A - A) \leq |\mathbf{G}|/|A|$ . Thus Proposition 14 gives us

**Corollary 15** *Let  $\mathcal{R}$  be a finite ring,  $A \subseteq \mathcal{R}$  be a set, and  $|A| = \alpha|\mathcal{R}|$ . Then*

$$\text{cov}^\times(2A - 2A) \leq \alpha^{-1},$$

*provided all numbers  $1, \dots, [\alpha^{-1}]$  belong to  $\mathcal{R}^*$ .*

Using the same method, one can estimate the multiplicative covering number of a Bohr set in  $\mathbb{Z}_p$  ( $p$  is a prime number)

$$\mathcal{B}(\Gamma, \varepsilon) = \{x \in \mathbb{Z}_p : \|x\gamma/p\| \leq \varepsilon, \forall \gamma \in \Gamma\} \quad \varepsilon \in (0, 1], \quad \Gamma \subseteq \mathbb{Z}_p,$$

namely, we have

$$\text{cov}^\times(\mathcal{B}(\Gamma, \varepsilon)) \leq \varepsilon^{-|\Gamma|}.$$

It is interesting to decrease the number of summands in Corollary 15. To this end let us obtain the main result of this section.

**Theorem 16** *Let  $q$  be a positive integer,  $A \subseteq \mathbb{Z}_q$  be a set,  $|A| = \alpha q$ . Suppose that the least prime factor of  $q$  greater than  $2\alpha^{-1} + 3$ . Then*

$$\text{cov}^\times(A - A) \leq \frac{1}{\alpha} + 1. \quad (51)$$

*More concretely,  $[k_*]^{-1} \cdot (A - A) = \mathbb{Z}_q$  for a certain  $k_* \leq \alpha^{-1} + 1$ .*

**Proof.** Let  $p_1$  be the least prime factor of  $q$ . By our assumption we know that  $p_1 \geq 2\alpha^{-1} + 3$ . Write  $p_1 = 2k + 1$  and take  $\Lambda = \{0, 1, \dots, k_*\}$ , where  $\lceil \alpha^{-1} - 1 \rceil + 1 = k_* \leq k$ . Then one has  $Y := (\Lambda - \Lambda) \setminus \{0\} \subseteq \mathbb{Z}_q^*$ . First of all, consider  $n \in \mathbb{Z}_q^*$  and form the set  $n \cdot \Lambda + A$ . Since  $|\Lambda||A| = (k_* + 1)\alpha q > q$ , it follows that there are different  $\lambda_1, \lambda_2 \in \Lambda$  such that

$$n\lambda_1 + a_1 \equiv n\lambda_2 + a_2 \pmod{q},$$

where  $a_1, a_2 \in A$  and  $a_1 \neq a_2$ . Hence  $n \in Y^{-1}(A - A)$  and thus  $\mathbb{Z}_q^* \subseteq Y^{-1}(A - A)$ . Also, notice that as in Proposition 14 one has  $Y^{-1}(A - A) = [k_*]^{-1} \cdot (A - A)$ .

Now let  $n = n'q_1$ , where  $q_1|q$  and  $n'$  is coprime to  $q$ . By the pigeonhole principle there is  $B \subseteq \mathbb{Z}_{q/q_1}$  and  $s \in \mathbb{Z}_q$  such that  $q_1B + s \subseteq A$  and the density of  $B$  in  $\mathbb{Z}_{q/q_1}$  is at least  $\alpha$ . In particular, we have  $q_1(B - B) \subseteq A - A$ . By the same argument as above one has  $n' \equiv y^{-1}(b_1 - b_2) \pmod{q/q_1}$ , where  $y \in Y$  and  $b_1, b_2 \in B$ . It follows that  $n \equiv y^{-1}(a_1 - a_2) \pmod{q}$  as required. Thus we have proved that  $[k_*]^{-1}(A - A) = \mathbb{Z}_q$  and hence  $\text{cov}^\times(A - A) \leq k_* \leq \alpha^{-1} + 1$ . This completes the proof.  $\square$

**Remark 17** *After the paper was written the author was informed by A. Fish that Theorem 16 holds in greater generality, namely, for any measure preserving system the same is true for the set of return times of a set of positive measure.*

Theorem 16 implies a consequence about the multiplicative covering numbers of the intersections of difference sets in the spirit of paper [19], see [19, Theorems 1,3].

**Corollary 18** *Let  $q$  be a positive integer, and  $A_1, \dots, A_k \subseteq \mathbb{Z}_q$  be sets,  $|A_i| = \alpha_i q$ ,  $i \in [k]$ . Suppose that the least prime factor of  $q$  greater than  $2(\alpha_1 \dots \alpha_k)^{-1} + 3$ . Then*

$$\text{cov}^\times \left( \bigcap_{i=1}^k (A_i - A_i) \right) \leq \frac{1}{\alpha_1 \dots \alpha_k} + 1. \quad (52)$$

**Proof.** Put  $A_{\vec{s}} = A_1 \cap (A_2 - s_1) \cap \dots \cap (A_k - s_{k-1})$ , where  $\vec{s} = (s_1, \dots, s_{k-1}) \in \mathbb{Z}_q^{k-1}$ . We have  $\sum_{\vec{s}} |A_{\vec{s}}| = |A_1| \dots |A_k|$  and hence there is  $\vec{s}_*$  such that  $|A_{\vec{s}_*}| \geq \alpha_1 \dots \alpha_k q$ . Clearly, for any  $\vec{s}$  one has

$$A_{\vec{s}} - A_{\vec{s}_*} \subseteq \bigcap_{i=1}^k (A_i - A_i).$$

Applying Theorem 16 with  $A = A_{\vec{s}_*}$ , we obtain bound (52). This completes the proof.  $\square$

As we have seen before Corollary 15 and Theorem 16 give us some bounds for the multiplicative covering numbers of difference sets. On the other hand, one can see that Theorem 16 does not hold for, say, nonzero shifts of Bohr sets, for the sumsets  $A + A$ , for the higher sumsets  $nA$ ,  $n > 2$  and so on. Indeed, consider the following

**Example.** *Let  $p$  be a prime number and  $S = [p/3, 2p/3)$  or  $S = \pm[p/6, p/3)$  to make  $S$  symmetric. Then the equation  $a + b \equiv c \pmod{p}$  has no solutions in  $a, b, c \in S$ . Further, we*

have  $|S| \gg p$  but it is easy to see that  $\text{cov}^\times(S)$  is unbounded. Indeed, if  $SX = \mathbb{Z}_p$  for a set  $X$  with  $|X| = O(1)$ , then we obtain a coloring of  $\mathbb{Z}_p$  with a finite number of colors and every color has no solutions to our equation  $a + b \equiv c \pmod{p}$ . It gives us a contradiction with the famous Schur theorem, see [13] (actually, it implies  $\text{cov}^\times(S) \gg \log p / \log \log p$ ).

In particular, we see that  $\text{cov}^\times(X + s)$  can be much larger than  $\text{cov}^\times(X)$  for a set  $X$  and a nonzero  $s$ .

Proposition 14 implies that any syndetic (i.e. having bounded gaps between its consecutive elements) set  $S \subseteq \mathbb{F}_p$ ,  $|S| \gg p$  has  $\text{cov}^\times(S - S) = O(1)$ . On the other hand, thanks to inclusion (50) any set of the form  $A - A$ , where  $A \subseteq \mathbb{F}_p$ ,  $|A| \gg p$  is syndetic (with the gap depending on  $A$  but not just on  $p/|A|$ , of course). Thus it is natural to ask about a generalization of Theorem 16 to the family of syndetic sets. Nevertheless, taking  $S = \{1 + kM\}_{k \in [(p-1)/M]}$ ,  $M \geq 4$  and  $p \equiv 2 \pmod{M}$ , say, we see that  $S$  is a syndetic set and  $S$  has no solutions to the equation  $a + b \equiv c \pmod{p}$ . Thus as in the example above we see that  $\text{cov}^\times(S)$  is unbounded.

**Remark 19** A dual form of Theorem 16 has no place, namely, there is a set  $A \subseteq \mathbb{Z}_p$ ,  $|A| \gg p$  such that  $\text{cov}^+(A/A) \gg \log p$ . In other words,  $\text{cov}^+(A/A)$  is close to the maximal possible value. To see this just take  $A$  to be the set of all quadratic residues, e.g., see [12, Proposition 14].

Finally, let us give another proof of a variant of Theorem 1 via our covering Theorem 16. Notice that the number  $d$  below can be a non-divisor of  $q$ .

**Theorem 20** Let  $q$  be a positive integer,  $A, B \subset \mathbb{Z}_q$  be sets,  $|A| = \alpha q$ ,  $|B| = \beta q$ , and let us assume that  $\alpha \geq \beta$ . Suppose that the least prime factor of  $q$  greater than  $2\beta^{-1} + 3$ . Then there is  $d \neq 0$  with

$$d \leq \alpha^{-\beta^{-1}-1}, \quad (53)$$

and such that

$$d \cdot \mathbb{Z}_q \subseteq (A - A)(B - B). \quad (54)$$

**Proof.** Applying Theorem 16 with  $A = B$ , we find a set  $X \subseteq \mathbb{Z}_q$ ,  $n := |X| \leq \beta^{-1} + 1$  such that  $X(B - B) = \mathbb{Z}_q$ . Let  $X = \{x_1, \dots, x_n\}$  and  $\vec{x} = (x_1, \dots, x_n) \in \mathbb{Z}_q^n$ . Considering the collection of the sets  $A^n + j \cdot \vec{x} \subseteq \mathbb{Z}_q^n$ ,  $j \geq 1$ , we see that there is  $0 < d \leq \alpha^{-n}$  with  $d \cdot X \subseteq A - A$ . Hence

$$(A - A)(B - B) \supseteq d \cdot X(B - B) \supseteq d \cdot \mathbb{Z}_q$$

as required. It remains to notice that

$$d \leq \alpha^{-n} \leq \alpha^{-\beta^{-1}-1}.$$

This completes the proof. □



## 6 Concluding remarks

Let us discuss other approaches to Theorem 1. First of all, recall the well-known Furstenberg's result [6].

**Theorem 21** *Let  $n$  be a positive integer,  $\delta \in (0, 1]$  be a real number, and  $S$  be a set of size  $n$ . Then for all sufficiently large  $N \geq N(\delta, n)$  an arbitrary set  $A \subseteq [N] \times [N]$ ,  $|A| \geq \delta N^2$  contains the set  $\alpha + \beta \cdot S$  for some  $\alpha$  and  $\beta \neq 0$ .*

Quantitative bounds for  $N(\delta, n)$  from Theorem 21 can be found in [14].

**Corollary 22** *Let  $q$  be a prime number,  $\mathcal{A} \subseteq \mathbb{F}_q^2$ ,  $|\mathcal{A}| = \delta q^2$  and  $A, B \subseteq \mathbb{F}_q$ ,  $|A| = \alpha_* q$ ,  $|B| = \beta_* q$ . Then there is a decreasing positive function  $\varphi$  such that if  $\min\{\alpha_*, \beta_*, \delta\} \geq \varphi(q)$ , then formula (18) takes place for  $\mathcal{B} = A \times B$ , any  $\lambda \in \mathbb{Z}_q^*$  and  $d = 1$ .*

**Proof.** Take  $S = S_1 = [k] \times [k]$  or  $S = S_2 = \{(2j, 2j) : j \in [k]\}$  for a certain positive integer  $k$ . Applying Theorem 21 with  $n = |S|$  and  $A = \mathcal{A}$ , we see that for some  $\alpha, \beta \neq 0$  the following holds  $\alpha + \beta \cdot S \subseteq \mathcal{A}$  and hence to solve (18) with  $d = 1$  it is sufficiently to find for any  $\lambda \in \mathbb{Z}_q^*$  some elements  $a \in \beta^{-1}(A - \alpha)$ ,  $b \in \beta^{-1}(B - \alpha)$  and  $(t_1, t_2) \in S$  such that

$$(t_1 - a)(t_2 - b) \equiv \lambda \pmod{q}.$$

If for a certain absolute constant  $C > 0$  one has  $k \gg \min^{-C}\{\alpha_*, \beta_*, \delta\}$ , then for  $S = S_2$  the last equation has a solution thanks to the famous Bourgain–Gamburd machine [4] (see details in [17], say) and for  $S = S_1$  (actually, for any dense subset of  $S_1$ ) the latter fact was obtained in [17, Theorem 3]. This completes the proof.  $\square$

The author does not know how to obtain Corollary 22 for composite  $q$  because there is no control over divisors of  $\beta$  in Theorem 21. It would be interesting to say something about prime factors of the dilation  $\beta$ .

We finish this section with a problem (it is interesting in its own right from a combinatorial point of view), which potentially gives another proof of Corollary 22 thanks to [17, Theorem 3].

**Problem.** *Let  $n$  be a positive integer,  $\delta, \kappa \in (0, 1]$  be real numbers. Then for all sufficiently large  $N \geq N(\delta, \kappa, n)$  an arbitrary set  $A \subseteq [N] \times [N]$ ,  $|A| \geq \delta N^2$  contains the set  $\alpha + \beta \cdot S$  for some  $\alpha$  and  $\beta$ , where  $S \subseteq [n] \times [n]$  is any set of size  $n^{1+\kappa}$ .*

Of course, some estimates on  $N(\delta, \kappa, n)$  follow from Theorem 21 but maybe it is possible to obtain a better bound.

## References

- [1] B. Bollobás, S. Janson, and O. Riordan. On covering by translates of a set. *Random Structures & Algorithms*, 38(1-2):33–67, 2011.

- [2] J. Bourgain. The sum-product theorem in  $\mathbb{Z}_q$  with  $q$  arbitrary. *Journal d'Analyse Mathématique*, 106(1):1–93, 2008.
- [3] J. Bourgain and A. Gamburd. Expansion and random walks in  $\mathrm{SL}_d(\mathbb{Z}/p^n\mathbb{Z})$ : I. *Journal of the European Mathematical Society*, 10(4):987–1011, 2008.
- [4] J. Bourgain and A. Gamburd. Uniform expansion bounds for cayley graphs of  $\mathrm{SL}_2(\mathbb{F}_p)$ . *Annals of Mathematics*, pages 625–642, 2008.
- [5] A. Fish. On product of difference sets for sets of positive density. *Proceedings of the American Mathematical Society*, 146(8):3449–3453, 2018.
- [6] H. Furstenberg. *Recurrence in ergodic theory and combinatorial number theory*, volume 10. Princeton University Press, 2014.
- [7] K. Gyarmati and A. Sárközy. Equations in finite fields with restricted solution sets. I (Character sums). *Acta Mathematica Hungarica*, 118(1-2):129–148, 2008.
- [8] K. Gyarmati and A. Sárközy. Equations in finite fields with restricted solution sets. II (Algebraic equations). *Acta Mathematica Hungarica*, 119(3):259–280, 2008.
- [9] D. Hart, A. Iosevich, and J. Solymosi. Sum-product estimates in finite fields via Kloosterman sums. *International Mathematics Research Notices*, 2007(9):rnm007–rnm007, 2007.
- [10] P. P. Pach. Ramsey type results on the solvability of certain equation in  $\mathbb{Z}_m$ . *Annual Volume 2013*, 13:41, 2013.
- [11] A. Sárközy. On sums and products of residues modulo  $p$ . *Acta Arithmetica*, 4(118):403–409, 2005.
- [12] T. Schoen and I. D. Shkredov. Higher moments of convolutions. *J. Number Theory*, 133(5):1693–1737, 2013.
- [13] I. Schur. Über die kongruenz  $x^m + y^m \equiv z^m \pmod{p}$ , jber. *Deutsch. Math. Verein*, 25:114–116, 1916.
- [14] S. Shelah. Primitive recursive bounds for van der waerden numbers. *Journal of the American Mathematical Society*, 1(3):683–697, 1988.
- [15] I. D. Shkredov. On monochromatic solutions of some nonlinear equations in  $\mathbb{Z}/p\mathbb{Z}$ . *Mat. Zametki*, 88(4):625–634, 2010.
- [16] I. D. Shkredov. Modular hyperbolas and bilinear forms of Kloosterman sums. *Journal of Number Theory*, 220:182–211, 2021.
- [17] I. D. Shkredov. On a girth-free variant of the Bourgain–Gamburd machine. *arXiv:2111.05751v1*, 2022.
- [18] I. E. Shparlinski. On the solvability of bilinear equations in finite fields. *Glasgow Mathematical Journal*, 50(3):523–529, 2008.

- [19] C. L. Stewart and R. Tijdeman. *On density-difference sets of sets of integers*, pages 701–710. Birkhäuser Basel, Basel, 1983.
- [20] T. Tao and V. Vu. *Additive combinatorics*, volume 105 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2006.
- [21] L. A. Vinh. The Szemerédi–Trotter type theorem and the sum-product estimate in finite fields. *European Journal of Combinatorics*, 32(8):1177–1181, 2011.