

Sequencings in Semidirect Products via the Polynomial Method

S. Costa¹, S. Della Fiore² and M. A. Ollis³

¹DICATAM, Sez. Matematica, Università degli Studi di Brescia,
Via Branze 43, I 25123 Brescia, Italy

²DI, Università degli Studi di Salerno, Via Papa Giovanni Paolo II 132,
I 84084 Fisciano, Italy,

³Emerson College, 120 Boylston Street, Boston, MA 02116, USA

Abstract

The *partial sums* of a sequence $\mathbf{x} = x_1, x_2, \dots, x_k$ of distinct non-identity elements of a group (G, \cdot) are $s_0 = id_G$ and $s_j = \prod_{i=1}^j x_i$ for $0 < j \leq k$. If the partial sums are all different then \mathbf{x} is a *linear sequencing* and if the partial sums are all different when $|i - j| \leq t$ then \mathbf{x} is a *t-weak sequencing*. We investigate these notions of sequenceability in semidirect products using the polynomial method. We show that every subset of order k of the non-identity elements of the dihedral group of order $2m$ has a linear sequencing when $k \leq 12$ and either $m > 3$ is prime or every prime factor of m is larger than $k!$, unless s_k is unavoidably the identity; that every subset of order k of a non-abelian group of order three times a prime has a linear sequencing when $5 < k \leq 10$, unless s_k is unavoidably the identity; and that if the order of a group is pe then all sufficiently large subsets of the non-identity elements are *t-weakly sequenceable* when $p > 3$ is prime, $e \leq 3$ and $t \leq 6$.

1 Introduction

Let (G, \cdot) be a group. We continue the investigation of when the elements of a subset of G may be ordered in such a way that their partial sums are distinct. We set up a general framework for investigating semidirect product groups and look in detail at the dihedral groups of order twice a prime and non-abelian groups of order three times a prime.

Definition 1.1. Let S be a subset of the non-identity elements of a group G with $|S| = k$. If x_1, x_2, \dots, x_k is an ordering of the elements of S , define the *j*th *partial sum* of the ordering by $s_0 = id_G$ and $s_j = \prod_{i=1}^j x_i$ for $0 < j \leq k$. The ordering of the subset S is a *linear sequencing* if the partial sums s_0, s_1, \dots, s_k are all different. It is a *cyclic sequencing* if the partial sums are different with the exception that $s_0 = s_k = id_G$. If S has a linear or cyclic sequencing then it is *sequenceable*.

2010 *Mathematics subject classification.* 05C25, 05C38, 05D40

Keywords and phrases. Combinatorial Nullstellensatz, Sequenceability.

If every subset of the non-identity elements of a group is sequenceable then the group is *strongly sequenceable*. Combining several results and conjectures concerning subsets of abelian groups dating back to 1961 [8], we have the conjecture that all abelian groups are strongly sequenceable. See [2, 5, 18] for details of the history, motivation and current state of this conjecture.

The question for non-abelian groups is much more widely open. In [6, 16] small counterexamples are presented to show that the potential conjecture that all non-abelian groups are strongly sequenceable is false.

Keedwell's Conjecture [12] is that if S contains all the non-identity elements of a non-abelian group G , then a linear sequencing exists if and only if $|G| \geq 10$. This is known to hold for many groups, including dihedral groups [11, 13], soluble groups with a single involution [3], at least one group of every odd order at which a non-abelian group exists [17] and all groups of order at most 255 [15].

Let S be a subset of size k of $G \setminus \{id_G\}$ whose elements can be ordered so that their total sum is not the identity. Then S is known to have a linear sequencing in the following situations [16]:

- $k \leq 4$ and S does not have one of two specific structures,
- G is dihedral of order $2m$ for prime $m > 3$ and $k \leq 9$,
- G is dihedral of order $2m$ for even or prime $m > 4$ and $k = 2m - 2$.

In this paper we make two contributions to finding linear sequencings in non-abelian groups. First, we improve the second item above, showing that it holds up to $k = 12$ and also for m whose prime factors are each larger than $k!$. (The result that allows for composite m with large prime factors in fact applies more widely than dihedral groups; see Section 3.) The second contribution concerns non-abelian groups of order three times a prime p ; there is at most one such group for any p and we denote it G_{3p} . In this case we show that subsets S of size $5 < k \leq 10$ are sequenceable and, further, that we can always find a linear sequencing for these subsets when the sum of all the elements is not unavoidably the identity.

In [4] a generalization of sequenceability is introduced.

Definition 1.2. Let S be a subset of the non-identity elements of a group G with $|S| = k$ and let x_1, x_2, \dots, x_k be an ordering of the elements of S with partial sums s_0, s_1, \dots, s_k . If $s_i \neq s_j$ for any distinct pair of indices i, j with $|i - j| \leq t$ then the ordering is a *t-weak sequencing* of S and S is *t-weakly sequenceable*.

Observe that a k -weak sequencing is exactly a linear sequencing and that a $(k - 1)$ -weak sequencing is either a linear or cyclic sequencing.

In [4] it is shown that a subset of non-identity elements of the cyclic group \mathbb{Z}_p of prime order p is t -weak sequenceable when $t \leq 6$ and, with the additional condition that there are no pairs of the form $\{x, -x\}$ in the subset, when $t = 7$. We extend the methodology to apply to groups G that are not cyclic of prime order, showing that if $|G| = pe$ then all sufficiently large subsets of $G \setminus \{id_G\}$ are t -weakly sequenceable when $p > 3$ is prime, $e \leq 3$ and $t \leq 6$.

In the next section we describe the polynomial method and how it can be implemented for groups that are semidirect products. In Subsection 2.1 and Section 3 we apply the

method to obtain the results on sequenceability and in Section 4 we apply it to obtain the results on weak sequenceability.

2 Polynomial Method in Semidirect Products

In this section, we apply a method that relies on the Non-Vanishing Corollary of the Combinatorial Nullstellensatz, see [1, 14]. Given a prime p (in the following p will be always assumed to be a prime), this corollary allows us to obtain a nonzero point to suitable polynomials on \mathbb{Z}_p derived starting from the ones defined in [10] and in [16].

Theorem 2.1. (Non-Vanishing Corollary) *Let \mathbb{F} be a finite field, and let $f(x_1, x_2, \dots, x_k)$ be a polynomial in $\mathbb{F}[x_1, x_2, \dots, x_k]$. Suppose the degree $\deg(f)$ of f is $\sum_{i=1}^k \gamma_i$, where each γ_i is a nonnegative integer, and suppose the coefficient of $\prod_{i=1}^k x_i^{\gamma_i}$ in f is nonzero. If C_1, C_2, \dots, C_k are subsets of \mathbb{F} with $|C_i| > \gamma_i$, then there are $c_1 \in C_1, \dots, c_k \in C_k$ such that $f(c_1, c_2, \dots, c_k) \neq 0$.*

In the notation of the Non-Vanishing Corollary, we call the monomial $x_1^{|C_1|-1} \dots x_k^{|C_k|-1}$ the *bounding monomial*. The corollary can be rephrased as requiring the polynomial to include a monomial of maximum degree that divides the bounding monomial (where by “include” we mean that it has a nonzero coefficient).

To use the Non-Vanishing Corollary we require a polynomial for which the nonzeros correspond to successful solutions to the case of the problem under consideration. Here, we will consider subsets S of a group (G, \cdot_G) that is the semidirect product of $(\mathbb{Z}_p, +)$ and another group (H, \cdot_H) . For such groups we will use the notation $G = \mathbb{Z}_p \rtimes_{\varphi} H$ for some group homomorphism $\varphi : H \rightarrow \text{Aut}(\mathbb{Z}_p)$.

Remark 2.2. These kind of semidirect products define a wide class of groups. Our hypotheses are, indeed, equivalent to consider groups (G, \cdot) that admit an isomorphic copy of $(\mathbb{Z}_p, +)$ as a normal subgroup N and whose cosets are of the form $N \cdot id, N \cdot c_1, \dots, N \cdot c_{e-1}$ where $e = |G/N|$ and $\{c_0 = id, \dots, c_{e-1}\}$ is a subgroup of G .

Now revisit the polynomial method introduced in [16] for dihedral groups and in [5] for groups of type $\mathbb{Z}_p \times H$. To avoid confusion, from now on, the symbol $+$ will be used only between elements of \mathbb{Z}_p . Let $\pi_2 : \mathbb{Z}_p \rtimes_{\varphi} H \rightarrow H$ be the projection map that picks out the second coordinate of an element and for a subset $S \subseteq (\mathbb{Z}_p \rtimes_{\varphi} H) \setminus \{(0, 0)\}$ let $\pi_2(S)$ be the multiset $\{\pi_2(s) : s \in S\}$. Also, the generic element of $\mathbb{Z}_p \rtimes_{\varphi} H$ will be indicated by $x \cdot_G a$ (or $y \cdot b$) implying that $x \in \mathbb{Z}_p$ and $a \in H$. We define the *type* of S to be the sequence $\boldsymbol{\lambda} = (\lambda_0, \dots, \lambda_{e-1})$, where λ_i is the number of times that i appears in $\pi_2(S)$.

Let T be a multiset of elements from $H \setminus \{0\}$ with size k . Let $\mathbf{a} = (a_1, \dots, a_k)$ be an arrangement of the elements of T with partial sums $\mathbf{b} = (b_0, b_1, \dots, b_k)$. If, for each i , the element i appears at most r times in \mathbf{b} , then \mathbf{a} is a *quotient sequencing* of T with respect to r . In our setting, given some $S \subseteq (\mathbb{Z}_p \rtimes_{\varphi} H) \setminus \{(0, 0)\}$ for which we wish to find a sequencing, we shall be interested in quotient sequencings of $\pi_2(S)$ with respect to p .

Given $S \subseteq (\mathbb{Z}_p \rtimes_{\varphi} H) \setminus \{(0, 0)\}$ we first construct a quotient sequencing of $\pi_2(S)$ with respect to p . We then use the polynomial method to show that there is a sequencing for S that projects elementwise onto that quotient sequencing. Given such a quotient

sequencing $\mathbf{a} = (a_1, \dots, a_k)$ with partial sums $\mathbf{b} = (b_0, b_1, \dots, b_k)$, let

$$\mathbf{x}_\mathbf{a} = (x_1 \cdot_G a_1, x_2 \cdot_G a_2, \dots, x_k \cdot_G a_k)$$

be a putative arrangement of the elements of S with partial sums

$$\mathbf{y}_\mathbf{a} = (s_0, \dots, s_k) = (y_0 \cdot_G b_0, y_1 \cdot_G b_1, \dots, y_k \cdot_G b_k).$$

We will need the following lemma:

Lemma 2.3. *Let (G, \cdot_G) be a group of type $\mathbb{Z}_p \rtimes_\varphi H$. Then, considering $x_1 \cdot_G a_1$ and $x_2 \cdot_G a_2$ in G , there exists $\varphi_{a_1} \in \mathbb{Z}_p$ such that:*

$$(x_1 \cdot_G a_1) \cdot_G (x_2 \cdot_G a_2) = (x_1 + (\varphi_{a_1} x_2)) \cdot_G (a_1 \cdot_H a_2)$$

where $\varphi_{a_1} x_2$ represent the product between the elements of \mathbb{Z}_p , φ_{a_1} and x_2 .

Proof. Due to the definition, we have that

$$(x_1 \cdot_G a_1) \cdot_G (x_2 \cdot_G a_2) = (x_1 + \varphi(a_1)(x_2)) \cdot_G (a_1 \cdot_H a_2).$$

Here the map

$$x \rightarrow \varphi(a_1)(x)$$

is an automorphism of $(\mathbb{Z}_p, +)$. It is well known that the automorphism of $(\mathbb{Z}_p, +)$ are the multiplications and hence there exists $\varphi_{a_1} \in \mathbb{Z}_p$ such that

$$\varphi(a_1)(x) = \varphi_{a_1} x.$$

Hence we have that

$$(x_1 + \varphi(a_1)(x_2)) \cdot_G (a_1 \cdot_H a_2) = (x_1 + (\varphi_{a_1} x_2)) \cdot_G (a_1 \cdot_H a_2)$$

which implies the thesis. □

From now on, when it is clear from the context, the index G or H in the products will be omitted.

Here we note that, if $j > i$, $s_i = s_j$ only when

$$(x_{i+1} \cdot a_{i+1}) \cdot (x_{i+2} \cdot a_{i+2}) \cdot \dots \cdot (x_j \cdot a_j) = id.$$

Applying Lemma 2.3, this equality can be written as

$$(x_{i+1} + \varphi_{a_{i+1}} x_{i+2}) \cdot (a_{i+1} \cdot a_{i+2}) \cdot \dots \cdot (x_j \cdot a_j) = id$$

and, reiterating it, as

$$(x_{i+1} + \varphi_{a_{i+1}} x_{i+2} + \varphi_{a_{i+1} \cdot a_{i+2}} x_{i+3} + \dots + \varphi_{a_{i+1} \dots a_{j-1}} x_j) \cdot (a_{i+1} \cdot a_{i+2} \cdot \dots \cdot a_j) = id.$$

This implies that

$$a_{i+1} \cdot a_{i+2} \cdot \dots \cdot a_j = id$$

which is $b_i = b_j$, and

$$x_{i+1} + \varphi_{a_{i+1}}x_{i+2} + \varphi_{a_{i+1}a_{i+2}}x_{i+3} + \cdots + \varphi_{a_{i+1}\dots a_{j-1}}x_j = 0.$$

Hence, reasoning as in [5], we can consider the following polynomial in the variables x_1, x_2, \dots, x_k

$$p_{\mathbf{a}} = \prod_{\substack{1 \leq i < j \leq k \\ a_i = a_j}} (x_j - x_i) \prod_{\substack{0 \leq i < j \leq k \\ b_i = b_j \\ j \neq i+1}} (x_{i+1} + \varphi_{a_{i+1}}x_{i+2} + \varphi_{a_{i+1}a_{i+2}}x_{i+3} + \cdots + \varphi_{a_{i+1}\dots a_{j-1}}x_j) \quad (1)$$

where the variable x_i ranges over the values $\{c : c \cdot a_i \in S\}$ for each i . Note that, if G is non-abelian, there is no reason to exclude the pair $(0, k)$ in the second product of Equation (1). Indeed we can even ask that $s_0 \neq s_k$. Now, it is clear that, given $S = \{x_1 \cdot a_1, x_2 \cdot a_2, \dots, x_k \cdot a_k\} \subseteq G \setminus \{0\}$ of size k , $p_{\mathbf{a}}(x_1, \dots, x_k) \neq 0$ only if the sequence $(x_1 \cdot a_1, x_2 \cdot a_2, \dots, x_k \cdot a_k)$ is a sequencing for the set S .

2.1 Computational Results

The method introduced in the previous paragraph generalizes that used in [16] to investigate the problem for dihedral groups. We recall the definition of these groups.

Definition 2.4. The dihedral group D_{2n} is the semidirect product $\mathbb{Z}_n \rtimes_{\varphi} \mathbb{Z}_2$ where

$$\varphi(0) = id \text{ and } \varphi(1)(x) = -x.$$

In the following example, we see how the coefficients of Equation (1) behave for such groups.

Example 2.5. *Example Let p be a prime, $p > 3$, and suppose $S \subseteq D_{2p} \setminus \{0\}$ with $|S| = 5$ and of type $(3, 2)$. The sequence $\mathbf{a} = (0, 1, 0, 0, 1)$ has partial sums $(0, 0, 1, 1, 1, 0)$ and so is a quotient sequencing of $\pi_2(S)$ with respect to p . We desire a sequencing of S of the form*

$$(x_1 \cdot 0, x_2 \cdot 1, x_3 \cdot 0, x_4 \cdot 0, x_5 \cdot 1).$$

The polynomial is

$$p_{\mathbf{a}} = (x_3 - x_1)(x_4 - x_1)(x_5 - x_2)(x_4 - x_3)(x_1 + x_2 - x_3 - x_4 - x_5)(x_2 - x_3 - x_4 - x_5)(x_3 + x_4).$$

To apply the Non-Vanishing Corollary we need a monomial of this polynomial which divides the bounding monomial $x_1^2 x_2 x_3^2 x_4^2 x_5$ with a nonzero coefficient. One such is $x_1^2 x_2 x_3^2 x_4 x_5^2$, which has coefficient 6. Hence whenever S has this form it has a sequencing.

Another class of groups that we will consider in this discussion is that of G_{3p} .

Definition 2.6. Given $p \equiv 1 \pmod{6}$ and r such that $r^3 \equiv 1 \pmod{p}$ and $r \not\equiv 1 \pmod{p}$, G_{3p} is defined as $\mathbb{Z}_p \rtimes_{\varphi} \mathbb{Z}_3$ where

$$\varphi(i)(x) = r^i x \text{ for any } i \in [0, 1, 2].$$

In case $r \equiv 1 \pmod{p}$, using the same construction we would obtain the direct product $\mathbb{Z}_p \times \mathbb{Z}_3$.

In the following example, we see how the coefficients of Equation (1) behave for such groups.

Example 2.7. *Example Let p be a prime, $p > 3$, and suppose $S \subseteq G_{3p} \setminus \{0\}$ with $|S| = 5$ and of type $(1, 3, 1)$. The sequence $\mathbf{a} = (1, 1, 0, 2, 1)$ has partial sums $(0, 1, 2, 2, 1, 2)$ and so is a quotient sequencing of $\pi_2(S)$ with respect to p . We desire a sequencing of S of the form*

$$(x_1 \cdot 1, x_2 \cdot 1, x_3 \cdot 0, x_4 \cdot 2, x_5 \cdot 1).$$

The polynomial is

$$p_{\mathbf{a}} = (x_2 - x_1)(x_5 - x_1)(x_5 - x_2)(x_2 + rx_3 + rx_4)(x_3 + x_4 + r^2x_5)(x_4 + r^2x_5).$$

To apply the Non-Vanishing Corollary we need a monomial of this polynomial which divides the bounding monomial $x_1^2x_2^2x_5^2$ with a nonzero coefficient. One such is $x_1^2x_2^2x_5^2$, which has coefficient $-r^4 = -r$. Hence whenever S has this form it has a sequencing.

We are now ready to prove the main result of this section.

Theorem 2.8. *Let $n = pe$ with p prime and let G be a group of size n . Then subsets S of size k of G are sequenceable in the following cases:*

1. $k \neq 4, 5$ such that $k \leq 10$ and $e \in \{1, 2, 3\}$;
2. $k \leq 12$, $p > 3$ and $e \in \{1, 2\}$.

Proof. The case $e = 1$ follows from the results of [10] and [5]. Since $\mathbb{Z}_p \rtimes_{\varphi} \{0\}$ is isomorphic to \mathbb{Z}_p , from the same papers, it follows the sequenceability of subsets S of D_{2p} (resp. G_{3p}) whose type is $(k, 0)$ (resp. $(k, 0, 0)$).

Note that a group of size $2p$ is either $\mathbb{Z}_p \times \mathbb{Z}_2$ or the dihedral group D_{2p} . Analogously, a group of size $3p$ is either $\mathbb{Z}_p \times \mathbb{Z}_3$ or G_{3p} for some r . Since the direct product case has already been considered in [5], we consider here just the cases of D_{2p} and G_{3p} . Also, for the dihedral groups, we can suppose that $k \geq 10$ since by [16] we know that subsets S of size $k \leq 9$ in D_{2p} (with $p > 3$) are sequenceable. We also know from [16, Lemma 3.1] that subsets $S \subseteq D_{2p}$ of size k and of type $(0, k)$ are sequenceable for every k . Then, in all of the other cases stated in the theorem, we can use the Non-Vanishing Corollary for suitable quotient sequences and nonzero monomial (in Tables 1 and 2 below we report the extreme cases). These results are obtained using the Python framework SageMath [19].

Here, when considering G_{3p} , for each type we provide a set of coefficients whose expressions depend on r . We have checked that, for any prime $p \equiv 1 \pmod{6}$, at least one of these coefficients is nonzero proceeding as follows. The coefficients can be written in the form $ar + b$ for some a, b whose expression do not depend on p . If $a \not\equiv 0 \pmod{p}$, the coefficient is zero if and only if $r \equiv -b \cdot a^{-1} \pmod{p}$. Recalling that r satisfies $r^2 + r + 1 \equiv 0 \pmod{p}$ we have that $a^{-2}(b^2 - ab + a^2) \equiv 0 \pmod{p}$ and hence,

$$b^2 - ab + a^2 \equiv 0 \pmod{p}. \tag{2}$$

On the other hand, if $a \equiv 0 \pmod{p}$ the coefficient is zero only when also b is zero modulo p and hence Equation (2) is satisfied also in this case. Since Equation (2) is satisfied for finitely many p , one can check that, for any prime p , at least one of the coefficients presented in Table 2 is nonzero. \square

Remark 2.9. Since in Equation (1) we have imposed also $s_0 \neq s_k$, we have that in case $G = D_{2p}$ (resp. $G = G_{3p}$), S is not of type $(k, 0)$ (resp. $(k, 0, 0)$) and k satisfies the hypotheses of the previous theorem, then S admits a linear sequencing.

This implies that, under the hypothesis of Theorem 2.8, S admits a linear sequencing unless s_k is unavoidably the identity.

The following Tables 1 and 2 contain the required monomials and their coefficients for the proof of Theorem 2.8 in the cases $(G, k) = (D_{2p}, 12)$ and $(G, k) = (G_{3p}, 10)$.

Table 1: Monomials and their coefficients for the proof of Theorem 2.8 in the case $|S| = 12$ and $G = D_{2p}$.

λ	\mathbf{a}	deg	monomial/s	coefficient/s
(11, 1)	(0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0)	71	$x_2^3 x_3^5 x_4^7 x_5^9 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^{10} x_{11}^{10} x_{12}^{10}$ $x_2^3 x_3^5 x_4^8 x_5^9 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^{10} x_{11}^{10} x_{12}^{10}$	$2 \cdot 3 \cdot 7^2 \cdot 131$ $2 \cdot 5 \cdot 7 \cdot 593$
(10, 2)	(0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 1)	64	$x_2^4 x_3^5 x_4^9 x_5^8 x_6 x_7 x_8^9 x_9^9 x_{10}^9 x_{11}^9 x_{12}$ $x_2^4 x_3^5 x_4^8 x_5^9 x_6 x_7 x_8^8 x_9^9 x_{10}^9 x_{11}^9 x_{12}$	$-2^7 \cdot 3 \cdot 29 \cdot 41$ $-2^5 \cdot 3 \cdot 101 \cdot 227$
(9, 3)	(0, 0, 0, 0, 1, 1, 1, 0, 0, 0, 0, 0)	57	$x_2^2 x_3^8 x_4^8 x_5^2 x_6^2 x_7^2 x_8^8 x_9^8 x_{10}^8 x_{11}^8 x_{12}^8$ $x_2^2 x_3^7 x_4^8 x_5^2 x_6^2 x_7^2 x_8^8 x_9^8 x_{10}^8 x_{11}^8 x_{12}^8$	$-2^3 \cdot 3^3 \cdot 5 \cdot 31$ $-2^2 \cdot 3^2 \cdot 61 \cdot 73$
(8, 4)	(0, 0, 0, 0, 1, 1, 1, 0, 0, 0, 0, 1)	54	$x_2^6 x_3^7 x_4^7 x_5^3 x_6^3 x_7^3 x_8^7 x_9^7 x_{10}^7 x_{11}^3 x_{12}^3$ $x_2^6 x_3^7 x_4^7 x_5^3 x_6^3 x_7^3 x_8^6 x_9^7 x_{10}^7 x_{11}^3 x_{12}^3$	$2^6 \cdot 7 \cdot 7669$ $-2^6 \cdot 11 \cdot 28591$
(7, 5)	(0, 0, 0, 1, 1, 1, 1, 1, 0, 0, 0, 0)	51	$x_2^6 x_3^6 x_4^4 x_5^4 x_6^4 x_7^4 x_8^6 x_9^6 x_{10}^6 x_{11}^6 x_{12}^6$ $x_1^6 x_3^6 x_4^4 x_5^4 x_6^4 x_7^4 x_8^6 x_9^6 x_{10}^6 x_{11}^6 x_{12}^6$	$-2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 401$ $-2^2 \cdot 5^2 \cdot 179 \cdot 389$
(6, 6)	(0, 0, 0, 1, 1, 1, 1, 1, 0, 0, 0, 1)	52	$x_1^2 x_2^4 x_3^5 x_4^5 x_5^5 x_6^5 x_7^5 x_8^5 x_9^5 x_{10}^5 x_{11}^5 x_{12}^5$ $x_1^3 x_2^3 x_3^5 x_4^5 x_5^5 x_6^5 x_7^5 x_8^5 x_9^5 x_{10}^5 x_{11}^5 x_{12}^5$	$2 \cdot 3^3 \cdot 181 \cdot 481373$ $-2 \cdot 3^2 \cdot 13 \cdot 22399121$
(4, 8)	(0, 0, 1, 1, 1, 1, 1, 1, 1, 0, 0, 1)	58	$x_2^3 x_3^4 x_4^5 x_5^7 x_6^7 x_7^7 x_8^7 x_9^3 x_{10}^3 x_{11}^7 x_{12}^7$ $x_2^3 x_3^4 x_4^6 x_5^8 x_6^7 x_7^7 x_8^7 x_9^3 x_{10}^3 x_{11}^7 x_{12}^7$	$2^4 \cdot 3 \cdot 79 \cdot 179 \cdot 18979$ $-2^4 \cdot 3^2 \cdot 5 \cdot 109 \cdot 119839$
(3, 9)	(0, 1, 1, 1, 1, 1, 1, 1, 1, 1, 0, 0)	63	$x_1^2 x_2^4 x_3^6 x_4^8 x_5^8 x_6^8 x_7^8 x_8^8 x_9^8 x_{10}^2 x_{11}^2 x_{12}^2$ $x_1^2 x_2^5 x_3^6 x_4^8 x_5^8 x_6^8 x_7^8 x_8^8 x_9^8 x_{10}^2 x_{11}^2 x_{12}^2$	$3 \cdot 62701 \cdot 131111$ $-2 \cdot 3 \cdot 5 \cdot 13 \cdot 17 \cdot 3699679$
(2, 10)	(0, 1, 1, 1, 1, 1, 1, 1, 1, 1, 0, 1)	72	$x_1 x_2^4 x_3^4 x_4^8 x_5^9 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^9 x_{11}^9 x_{12}^9$ $x_1 x_2^4 x_3^5 x_4^8 x_5^9 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^9 x_{11}^9 x_{12}^9$	$2 \cdot 5 \cdot 71 \cdot 1427 \cdot 1033463$ $-2 \cdot 3^2 \cdot 7 \cdot 23 \cdot 215947813$
(1, 11)	(1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 0)	80	$x_2^5 x_3^5 x_4^7 x_5^7 x_6^8 x_7^8 x_8^{10} x_9^{10} x_{10}^{10} x_{11}^{10} x_{12}^{10}$ $x_2^4 x_3^6 x_4^7 x_5^8 x_6^8 x_7^8 x_8^{10} x_9^{10} x_{10}^{10} x_{11}^{10} x_{12}^{10}$	$2^3 \cdot 7 \cdot 101 \cdot 81083329$ $-2^5 \cdot 5 \cdot 31 \cdot 3557 \cdot 21647$

Table 2: Monomials and their coefficients for the proof of Theorem 2.8 in the case $|S| = 10$ and $G = G_{3p}$.

λ	\mathbf{a}	deg	monomial/s	coefficient/s
(9, 1, 0)	(0, 0, 0, 0, 0, 1, 0, 0, 0, 0)	52	$x_2^2 x_3^4 x_4^7 x_5^8 x_7^8 x_8^8 x_9^8 x_{10}^8$	-2^2
(8, 2, 0)	(0, 0, 0, 1, 0, 0, 0, 1, 0, 0)	36	$x_2^2 x_3^4 x_5^3 x_6^6 x_7^7 x_8^6 x_9^6 x_{10}^7$	-2
(8, 1, 1)	(0, 0, 0, 0, 1, 0, 0, 0, 0, 2)	45	$x_1 x_2^4 x_3^6 x_4^7 x_5^6 x_6^7 x_7^7 x_8^7 x_9^7$ $x_1^2 x_2^3 x_3^6 x_4^7 x_5^6 x_6^7 x_7^7 x_8^7 x_9^7$	$-1791r + 573$ $-235r - 746$

λ	\mathbf{a}	deg	monomial/s	coefficient/s
(7, 3, 0)	(0, 0, 1, 0, 0, 0, 1, 0, 0, 1)	32	$x_1^2 x_2^3 x_4^2 x_5^5 x_6^6 x_7 x_8^5 x_9^6 x_{10}^2$ $x_1^2 x_2^2 x_3^4 x_5^5 x_6^6 x_7 x_8^5 x_9^6 x_{10}^2$	$14r + 73$ $12r + 21$
(7, 2, 1)	(0, 0, 0, 1, 0, 0, 1, 0, 0, 2)	30	$x_1 x_2^2 x_3^4 x_4 x_5^5 x_6^5 x_7 x_8^5 x_9^6$	1
(6, 4, 0)	(0, 0, 1, 0, 0, 1, 0, 0, 1, 1)	30	$x_2^2 x_3^2 x_4^4 x_5^5 x_6^2 x_7^4 x_8^5 x_9^3 x_{10}^3$ $x_1 x_2^2 x_3^2 x_4^3 x_5^5 x_6^2 x_7^4 x_8^5 x_9^3 x_{10}^3$	$-1168r + 72$ $-436r - 1078$
(6, 3, 1)	(0, 0, 0, 1, 0, 0, 1, 0, 2, 1)	27	$x_2^2 x_3^5 x_4 x_5^5 x_6^2 x_7^5 x_8^2 x_{10}$	$r + 1$
(6, 2, 2)	(0, 0, 1, 0, 0, 1, 0, 0, 2, 2)	26	$x_1^3 x_2^3 x_3 x_4^3 x_5^4 x_6 x_7^4 x_8^5 x_9 x_{10}$ $x_1^2 x_2^5 x_3^4 x_5^4 x_6 x_7^4 x_8^5 x_9 x_{10}$	$10r - 64$ $155r + 583$
(5, 5, 0)	(0, 0, 1, 0, 0, 1, 0, 1, 1, 1)	30	$x_2^2 x_3 x_4^3 x_5^4 x_6^4 x_7^4 x_8^4 x_9^4 x_{10}$ $x_1 x_2^2 x_3^2 x_4^2 x_5^4 x_6^4 x_7^4 x_8^4 x_9^4 x_{10}$	$1965r + 2715$ $385r - 190$
(5, 4, 1)	(0, 0, 1, 0, 0, 1, 0, 2, 1, 1)	26	$x_1 x_2^4 x_3 x_4^3 x_5^4 x_6^3 x_7^4 x_8^3 x_9^3 x_{10}$ $x_1^2 x_2^3 x_3 x_4^3 x_5^4 x_6^3 x_7^4 x_8^3 x_9^3 x_{10}$	$30r + 156$ $42r + 72$
(5, 3, 2)	(0, 0, 0, 1, 0, 1, 0, 2, 1, 2)	24	$x_1^4 x_2^4 x_3^3 x_4^2 x_5^4 x_6^2 x_7^2 x_8 x_9^2$	2
(4, 6, 0)	(0, 1, 0, 0, 1, 0, 1, 1, 1, 1)	32	$x_1^3 x_2^5 x_3^3 x_4^3 x_5^5 x_6^3 x_7^4 x_8^4 x_9 x_{10}$ $x_1^2 x_2^5 x_3^3 x_4^3 x_5^5 x_6^3 x_7^5 x_8^4 x_9 x_{10}$	$-11377r + 11374$ $-21587r - 27358$
(4, 5, 1)	(0, 0, 1, 0, 1, 0, 1, 1, 1, 2)	27	$x_2^2 x_3^3 x_4^3 x_5^4 x_6^3 x_7^4 x_8^4 x_9^4$ $x_1^2 x_2^2 x_3^3 x_4^3 x_5^3 x_6^3 x_7^3 x_8^4 x_9^4$	$-1286r + 68$ $-333r - 119$
(4, 4, 2)	(0, 0, 0, 1, 0, 1, 2, 1, 2, 1)	24	$x_1^2 x_2^2 x_3^3 x_4^3 x_5^3 x_6^3 x_7 x_8^3 x_9 x_{10}^3$	$-2r$
(4, 3, 3)	(0, 0, 1, 0, 1, 0, 2, 1, 2, 2)	23	$x_1^2 x_3^2 x_4^2 x_5^3 x_6^2 x_7^2 x_8^2 x_9^2 x_{10}^2$ $x_1^3 x_2^2 x_3^2 x_4^2 x_5^3 x_6^2 x_7^2 x_8^2 x_9^2 x_{10}^2$	$-1452r + 1686$ $-759r - 348$
(3, 7, 0)	(0, 1, 0, 1, 0, 1, 1, 1, 1, 1)	36	$x_2 x_3^2 x_4 x_5^2 x_6^6 x_7^6 x_8^6 x_9^6 x_{10}^6$ $x_2 x_3^2 x_4^2 x_5^4 x_6^5 x_7^6 x_8^6 x_9^6 x_{10}^6$	$-19616r + 331264$ $802469r - 381850$
(3, 6, 1)	(0, 0, 1, 0, 1, 1, 1, 1, 2, 1)	30	$x_1^2 x_2 x_3^5 x_4^2 x_5^5 x_6^5 x_7^5 x_8^4 x_{10}$ $x_1^2 x_2 x_3^3 x_4^2 x_5^5 x_6^5 x_7^5 x_8^4 x_{10}^3$	$3396r - 230$ $133r + 344$
(3, 5, 2)	(0, 0, 1, 0, 1, 2, 1, 2, 1, 1)	26	$x_1^2 x_2^2 x_3^2 x_4^2 x_5^4 x_6 x_7^4 x_8 x_9^4 x_{10}^4$ $x_1^2 x_2^2 x_3^2 x_4^2 x_5^3 x_6 x_7^4 x_8 x_9^4 x_{10}^4$	$-3r + 24$ $21r + 21$
(3, 4, 3)	(0, 0, 1, 0, 1, 1, 2, 2, 1, 2)	24	$x_1^2 x_2^2 x_3^3 x_4^2 x_5^3 x_6^3 x_7^2 x_8^2 x_9^3 x_{10}^2$	$1358r + 1812$
	(0, 0, 1, 1, 0, 1, 2, 2, 1, 2)	24	$x_1^2 x_2^2 x_3^3 x_4^2 x_5^3 x_6^3 x_7^2 x_8^2 x_9^3 x_{10}^2$	$-1742r - 1035$
(2, 8, 0)	(0, 1, 0, 1, 1, 1, 1, 1, 1, 1)	42	$x_3 x_4^2 x_5^4 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}^7$ $x_3 x_4^2 x_5^5 x_6^6 x_7^7 x_8^7 x_9^7 x_{10}^7$	$65374r + 67999$ $-160333r - 7057$
(2, 7, 1)	(0, 1, 0, 1, 1, 1, 1, 2, 1, 1)	35	$x_1 x_3 x_4^3 x_5^6 x_6^6 x_7^6 x_8^6 x_{10}^6$ $x_1 x_3 x_4^4 x_5^6 x_6^6 x_7^6 x_8^6 x_{10}^6$	$17861r - 27263$ $12821r + 26143$
(2, 6, 2)	(2, 0, 1, 1, 1, 1, 1, 2, 1, 0)	30	$x_3^3 x_4^5 x_5^5 x_6^5 x_7^5 x_8^5 x_9^5 x_{10}$ $x_3^4 x_4^4 x_5^5 x_6^5 x_7^5 x_8^5 x_9^5 x_{10}$	$-292r + 246$ $85r + 126$

λ	\mathbf{a}	deg	monomial/s	coefficient/s
(2, 5, 3)	(2, 1, 1, 1, 0, 2, 2, 1, 0, 1)	27	$x_1^2 x_2^4 x_3^4 x_4^4 x_5^2 x_6^2 x_7^4 x_8 x_9 x_{10}^4$	17651r - 23546
			$x_1^2 x_2^3 x_3^4 x_4^4 x_5 x_6^2 x_7^2 x_8^4 x_9 x_{10}^4$	-10860r - 152465
(2, 4, 4)	(0, 1, 0, 1, 1, 2, 2, 1, 2, 2)	26	$x_1 x_2^3 x_3 x_4^3 x_5^3 x_6^3 x_7^3 x_8^3 x_9^3 x_{10}^3$	136392r + 247068
	(0, 1, 1, 0, 1, 2, 2, 1, 2, 2)	26	$x_1 x_2^3 x_3^3 x_4 x_5^3 x_6^3 x_7^3 x_8^3 x_9^3 x_{10}^3$	-175716r - 167652
(1, 9, 0)	(1, 0, 1, 1, 1, 1, 1, 1, 1, 1)	50	$x_2^2 x_3^4 x_4^5 x_5^7 x_6^8 x_7^8 x_8^8 x_9^8 x_{10}^8$	992906r + 1427608
			$x_3^3 x_4^3 x_5^5 x_6^7 x_7^8 x_8^8 x_9^8 x_{10}^8$	-1815167r - 733348
(1, 8, 1)	(0, 1, 1, 1, 1, 1, 1, 1, 1, 2)	42	$x_2 x_3^2 x_4^4 x_5^7 x_6^7 x_7^7 x_8^7 x_9^7$	63854r - 6907
			$x_2 x_3^2 x_4^5 x_5^6 x_6^7 x_7^7 x_8^7 x_9^7$	38518r + 68818
(1, 7, 2)	(0, 1, 1, 1, 1, 1, 1, 2, 2, 1)	36	$x_3^4 x_4^6 x_5^6 x_6^6 x_7^6 x_8 x_9 x_{10}^6$	-16124r - 27940
			$x_3^5 x_4^5 x_5^6 x_6^6 x_7^6 x_8 x_9 x_{10}^6$	3718r + 13449
(1, 6, 3)	(1, 1, 0, 1, 1, 1, 2, 1, 2, 2)	32	$x_1 x_2^5 x_3^5 x_4^5 x_5^2 x_6^2 x_7^2 x_8^2 x_9^2 x_{10}^2$	31065r + 30039
			$x_1^2 x_2^4 x_3^5 x_4^5 x_5^2 x_6^2 x_7^2 x_8^2 x_9^2 x_{10}^2$	1764r + 9147
(1, 5, 4)	(0, 1, 1, 1, 1, 1, 2, 2, 2, 2)	30	$x_2^4 x_3^4 x_4^4 x_5^4 x_6^3 x_7^3 x_8^3 x_9^3 x_{10}^3$	211692r + 376028
			$x_2^2 x_3^4 x_4^4 x_5^4 x_6^3 x_7^3 x_8^3 x_9^3 x_{10}^3$	78700r - 731650
(0, 10, 0)	(1, 1, 1, 1, 1, 1, 1, 1, 1, 1)	60	$x_1^2 x_2^2 x_3^4 x_4^7 x_5^9 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^9$	-250444r - 433547
			$x_1^2 x_2^2 x_3^5 x_4^6 x_5^9 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^9$	1040607r + 497898
(0, 9, 1)	(1, 1, 1, 1, 1, 2, 1, 1, 1, 1)	51	$x_2^2 x_3^3 x_4^6 x_5^8 x_6^8 x_7^8 x_8^8 x_9^8 x_{10}^8$	21957r - 14546
			$x_2^2 x_3^4 x_4^5 x_5^8 x_6^8 x_7^8 x_8^8 x_9^8 x_{10}^8$	37855r + 47642
(0, 8, 2)	(1, 1, 1, 1, 1, 1, 1, 1, 2, 2)	44	$x_2^3 x_3^4 x_4^7 x_5^7 x_6^7 x_7^7 x_8^7 x_9 x_{10}$	1698588r + 406932
			$x_2^3 x_3^5 x_4^6 x_5^7 x_6^7 x_7^7 x_8^7 x_9 x_{10}$	201528r + 1213386
(0, 7, 3)	(2, 1, 1, 1, 1, 1, 1, 1, 2, 2)	39	$x_1 x_2^2 x_3^2 x_4^6 x_5^6 x_6^6 x_7^6 x_8^6 x_9^2 x_{10}^2$	824103r + 258808
			$x_1 x_2^2 x_3^3 x_4^5 x_5^6 x_6^6 x_7^6 x_8^6 x_9^2 x_{10}^2$	89629r + 972050
(0, 6, 4)	(2, 1, 2, 1, 1, 1, 1, 1, 2, 2)	38	$x_1 x_2^3 x_3^3 x_4^5 x_5^5 x_6^5 x_7^5 x_8^5 x_9^3 x_{10}^3$	1390432r + 1536910
			$x_2^4 x_3^3 x_4^5 x_5^5 x_6^5 x_7^5 x_8^5 x_9^3 x_{10}^3$	-792953r - 108625
(0, 5, 5)	(1, 1, 1, 1, 1, 2, 2, 2, 2, 2)	35	$x_1 x_2^2 x_3^4 x_4^4 x_5^4 x_6^4 x_7^4 x_8^4 x_9^4 x_{10}^4$	-710283r + 3022047
			$x_1^2 x_2^2 x_3^3 x_4^4 x_5^4 x_6^4 x_7^4 x_8^4 x_9^4 x_{10}^4$	2272296r + 657732

3 To Infinity and Back

In this section we apply the method of [7] (see also [5]) to generalize the sequenceability results here obtained to a more general class of semidirect products.

First, given a subset S of a group G , we define the set

$$\Upsilon(S) = S \cup \Delta(S).$$

Then we can adapt [7, Lemma 2.1] as

Lemma 3.1. *Let G_1 and G_2 be groups. Given a subset S of $G_1 \setminus \{0_{G_1}\}$ of size k , suppose there exists a homomorphism $\phi : G_1 \rightarrow G_2$ such that $\ker(\phi) \cap \Upsilon(S) = \emptyset$. Then $\phi(S)$*

is a subset of $G_2 \setminus \{0_{G_2}\}$ of size k and the subset S is sequenceable whenever $\phi(S)$ is sequenceable.

In the following, we consider groups of the form $(G \rtimes_{\varphi} H, \cdot)$ where $(G, +)$ is abelian, $H = \{h_0 = 0_H, h_1, \dots, h_{e-1}\}$ is finite, $(H, \cdot_H)^1$ is not necessarily abelian and where φ is a group homomorphism $\varphi : H \rightarrow \text{Aut}(G)$. We will also assume that the homomorphism φ satisfies the following property:

- (*) given $h \in H$, $\varphi(h)$ is either the identity or $\varphi(h)(x) = -x$ for any $x \in G$.

Note that the latter is an automorphism of G because G is abelian.

Remark 3.2. Two classes of semidirect product that satisfy property (*) are the generalized dihedral groups and the standard direct product. Indeed:

- 1) The generalized dihedral group $Dih(G)$ can be defined as $G \rtimes_{\varphi} \mathbb{Z}_2$ where

$$\varphi(0) = id \text{ and } \varphi(1)(x) = -x.$$

- 2) Any direct product $G \times H$ can be defined as $G \rtimes_{\varphi} H$ where

$$\varphi(h) = id \text{ for any } h \in H.$$

Given a multiset S of elements of $G \rtimes_{\varphi} H$, we can define the *type* of S also when G is a generic group. It is the sequence $\boldsymbol{\lambda} = (\lambda_0, \dots, \lambda_{e-1})$, where λ_i is the number of times that h_i appears in $\pi_2(S)$ and π_2 is the projection over H . Note that two multisets S_1 of $G_1 \rtimes_{\varphi} H$ and S_2 of $G_2 \rtimes_{\varphi} H$ can have the same type even though G_1 is not equal to G_2 .

Given a semidirect product $G_1 \rtimes_{\varphi} H$, we can define the corresponding product $G_2 \rtimes_{\varphi'} H$ where $\varphi'(h)$ is the identity on G_2 (resp $\varphi'(h)(x) = -x$ for any $x \in G_2$) if and only if $\varphi(h)$ is the identity on G_1 (resp $\varphi(h)(x) = -x$ for any $x \in G_1$). For simplicity, we will denote, with abuse of notation, by $G_2 \rtimes_{\varphi} H$ this semidirect product.

Proposition 3.3. *Let H be a finite group, let φ satisfy property (*), and suppose that, for infinitely many primes p , any subset of a given type $\boldsymbol{\lambda} = (\lambda_0, \dots, \lambda_{e-1})$ of $\mathbb{Z}_p \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_p \rtimes_{\varphi} H}\}$ is sequenceable. Then also any subset of type $\boldsymbol{\lambda}$ of $\mathbb{Z} \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z} \rtimes_{\varphi} H}\}$ is sequenceable.*

Proof. Consider a subset S of $\mathbb{Z} \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z} \rtimes_{\varphi} H}\}$ of type $\boldsymbol{\lambda}$. Let $p > \max_{z \in \Upsilon(S)} |\pi_{\mathbb{Z}}(z)|$ be a prime such that any subset of $\mathbb{Z}_p \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_p \rtimes_{\varphi} H}\}$ of type $\boldsymbol{\lambda}$ is sequenceable.

Here we note that, since $\mathbb{Z} \rtimes_{\varphi} H$ and $\mathbb{Z}_p \rtimes_{\varphi} H$ are defined with corresponding maps φ , the map $\pi_p \times id : \mathbb{Z} \rtimes_{\varphi} H \rightarrow \mathbb{Z}_p \rtimes_{\varphi} H$ is a group homomorphism. Indeed in this case we have that

$$\begin{aligned} (\pi_p \times id)(x_1, k_1) \cdot (\pi_p \times id)(x_2, k_2) &= (x_1 \pmod{p}, k_1) \cdot (x_2 \pmod{p}, k_2) = \\ &= ((x_1 + \varphi(k_1)(x_2)) \pmod{p}, k_1 \cdot_H k_2) = (\pi_p \times id)[(x_1, k_1) \cdot (x_2, k_2)]. \end{aligned}$$

Then $\Upsilon(S)$ and the kernel of the map $\pi_p \times id : \mathbb{Z} \rtimes_{\varphi} H \rightarrow \mathbb{Z}_p \rtimes_{\varphi} H$ are disjoint sets. Therefore, since $\pi_p \times id$ is the identity on H and S is of type $\boldsymbol{\lambda}$, we have $(\pi_p \times id)(S)$ also is of type $\boldsymbol{\lambda}$. It then follows from Lemma 3.1 that S also is sequenceable. \square

¹Also in this section, when it is clear from the context, the index H will be omitted.

Remark 3.4. In the previous proposition, we can see why property (*) is needed in this discussion. Indeed we need that $\varphi(a)$ is a group automorphism of \mathbb{Z} and hence it maps 1 in a generator of \mathbb{Z} . It follows that $\varphi(a)(1) \in \{1, -1\}$ or, equivalently, φ satisfies property (*).

We now consider the free abelian group \mathbb{Z}^n of rank n . We obtain the following proposition whose proof is omitted since it is the same as Proposition 4.3 of [5].

Proposition 3.5. *Let H be a finite group, let φ satisfy property (*), and suppose that any subset of a given type $\lambda = (\lambda_0, \dots, \lambda_{e-1})$ of $\mathbb{Z} \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z} \rtimes_{\varphi} H}\}$ is sequenceable. Then also any subset of type λ of $\mathbb{Z}^n \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}^n \rtimes_{\varphi} H}\}$ (for any $n \geq 2$) is sequenceable.*

From the previous proposition, we deduce this result. Also in this case we omit the proof since it is the same as Theorem 4.4 of [5]

Theorem 3.6. *Let H be a finite group, let φ satisfy property (*), and suppose that, for infinitely many primes p , any subset of a given type $\lambda = (\lambda_0, \dots, \lambda_{e-1})$ of $\mathbb{Z}_p \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_p \rtimes_{\varphi} H}\}$ is sequenceable. Then also any subset of $G \rtimes_{\varphi} H \setminus \{0_{G \rtimes_{\varphi} H}\}$ of type λ is sequenceable provided that G is torsion-free and abelian.*

Corollary 3.7. *Let H be a finite group, let φ satisfy property (*), k be a positive integer, and suppose that, for infinitely many primes p , any subset of size k of $\mathbb{Z}_p \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_p \rtimes_{\varphi} H}\}$ is sequenceable. Then also any subset of size k of $G \rtimes_{\varphi} H \setminus \{0_{G \rtimes_{\varphi} H}\}$ is sequenceable provided that G is torsion-free and abelian.*

Given an element g of an abelian group G , we denote by $o(g)$ the cardinality of the cyclic subgroup $\langle g \rangle$ generated by g . Furthermore, we set

$$\vartheta(G) = \min_{0_G \neq g \in G} o(g).$$

As a consequence of Theorem 3.6, and with essentially the same proof of Theorem 4.6 of [5], we can prove the following result.

Theorem 3.8. *Let H be a finite group, let φ satisfy property (*), and suppose that, for infinitely many primes p , any subset of $\mathbb{Z}_p \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_p \rtimes_{\varphi} H}\}$ of a given type $\lambda = (\lambda_0, \dots, \lambda_{e-1})$ is sequenceable. Then there exists a positive integer $N(\lambda)$ such that any subset of type λ of $G \rtimes_{\varphi} H \setminus \{0_{G \rtimes_{\varphi} H}\}$ is sequenceable provided that G is abelian and $\vartheta(G) > N(\lambda)$.*

Corollary 3.9. *Let H be a finite group, let φ satisfy property (*), k be a positive integer, and suppose that, for infinitely many primes p , any subset of size k of $\mathbb{Z}_p \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_p \rtimes_{\varphi} H}\}$ is sequenceable. Then, there exists a positive integer $N(k)$ such that any subset of size k of $G \rtimes_{\varphi} H \setminus \{0_{G \rtimes_{\varphi} H}\}$ is sequenceable provided that G is abelian and $\vartheta(G) > N(k)$.*

Therefore, as a consequence of Theorem 2.8, and recalling that the only exceptions to that theorem, in case of dihedral groups and $k \leq 12$, are for $k = 4$ and $p = 3$ (see Theorem 3.4 of [16]), we can prove our first asymptotic result.

Theorem 3.10. *Let G be a group such that $\vartheta(G) > N(k)$. Then any subset S of $Dih(G)$ such that $|S| \leq 12$ is sequenceable.*

Remark 3.11. Because of Remark 2.9, with essentially the same proof of Theorem 3.10, we can prove that if S is a subset of $Dih(G)$ not of type $(k, 0)$, $k \leq 12$ and $\vartheta(G) > N(k)$, then S admits a linear sequencing.

This implies that, under the hypothesis of Theorem 3.10, S admits a linear sequencing unless s_k is unavoidably the identity.

3.1 An Explicit Upper Bound

Now, in case $G = \mathbb{Z}_m$, and adapting again the procedure of [5], we provide an explicit upper bound on the numbers $N(\lambda)$ and $N(k)$. First, we need to recall some elementary facts about the solutions of a linear system over fields and commutative rings.

Theorem 3.12 (Corollary of Cramer's Theorem). *Let*

$$\begin{cases} m_{1,1}x_1 + \cdots + m_{1,k}x_k = b_1 \\ \cdots \\ m_{l,1}x_1 + \cdots + m_{l,k}x_k = b_l \end{cases} \quad (3)$$

be a linear system over a commutative ring R whose associated matrix $M = (m_{i,j})$ is an $l \times k$ matrix over R . Let (x_1, \dots, x_k) be a solution of the system and let M' be a $k' \times k'$ square, nonsingular (i.e. $\det(M')$ is invertible in R), submatrix of M . We assume, without loss of generality, that M' is obtained by considering the first k' rows and the first k' columns of M . Then x_1, \dots, x_k satisfy the following relation

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_{k'} \end{bmatrix} = \begin{bmatrix} m_{1,1} & m_{1,2} & m_{1,3} & \cdots & m_{1,k'} \\ m_{2,1} & m_{2,2} & m_{2,3} & \cdots & m_{2,k'} \\ \vdots & \vdots & \vdots & & \vdots \\ m_{k',1} & m_{k',2} & m_{k',3} & \cdots & m_{k',k'} \end{bmatrix}^{-1} \begin{bmatrix} b_1 - (m_{1,k'+1}x_{k'+1} + \cdots + m_{1,k}x_k) \\ b_2 - (m_{2,k'+1}x_{k'+1} + \cdots + m_{2,k}x_k) \\ \vdots \\ b_{k'} - (m_{k',k'+1}x_{k'+1} + \cdots + m_{k',k}x_k) \end{bmatrix}.$$

Equivalently, set

$$M'_j := \begin{bmatrix} m_{1,1} & m_{1,2} & \cdots & m_{1,j-1} & b_1 - (m_{1,k'+1}x_{k'+1} + \cdots + m_{1,k}x_k) & \cdots & m_{1,k'} \\ m_{2,1} & m_{2,2} & \cdots & m_{2,j-1} & b_2 - (m_{2,k'+1}x_{k'+1} + \cdots + m_{2,k}x_k) & \cdots & m_{2,k'} \\ \vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\ m_{k',1} & m_{k',2} & \cdots & m_{k',j-1} & b_{k'} - (m_{k',k'+1}x_{k'+1} + \cdots + m_{k',k}x_k) & \cdots & m_{k',k'} \end{bmatrix}$$

we have that, for any $j \in [1, k']$

$$x_j = \frac{\det(M'_j)}{\det(M')}. \quad (4)$$

Now assume that R is a field and that the system (3) admits at least one solution. Then, given a maximal square nonsingular submatrix M' of M , and fixed $x_{k'+1}, \dots, x_k$ in R , we have that x_1, x_2, \dots, x_k are a solution of the system (3) if and only if $x_1, \dots, x_{k'}$ satisfy the relation (4).

Remark 3.13. In case the system (3) of Theorem 3.12 is homogeneous (i.e. $b_1 = b_2 = \dots = b_l = 0$), due to the Laplace expansion of the determinant, $\det(M'_j)$ can be written as a linear combination of $x_{k'+1}, \dots, x_k$. More precisely, denoting by $M'_{i,j}$ the matrix obtained by deleting the i -th row and the j -th column of M' , we have that

$$\det(M'_j) = \sum_{i=1}^{k'} (-1)^{j+i+1} \det(M'_{i,j}) (m_{i,k'+1} x_{k'+1} + \dots + m_{i,k} x_k). \quad (5)$$

Now we are ready to provide, in case of groups of type $\mathbb{Z}_m \rtimes_{\varphi} H$ (where φ satisfies property (*)), a quantitative version of Theorem 3.8. In the following proof, developing the notations of the previous sections, given a set S of size k and an ordering $\mathbf{x} = (x_1, \dots, x_k)$ of S , we denote by $s_i(\mathbf{x})$ the partial sum $x_1 \cdot x_2 \dots x_{i-1} \cdot x_i$ where $s_0(\mathbf{x})$ is always zero.

Theorem 3.14. *Let H be a finite group, let φ satisfy property (*), and suppose that, for infinitely many primes p , any subset of a given type $\lambda = (\lambda_0, \dots, \lambda_{e-1})$ of $\mathbb{Z}_p \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_p \rtimes_{\varphi} H}\}$ is sequenceable. Then also any subset of type λ of $\mathbb{Z}_m \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_m \rtimes_{\varphi} H}\}$ is sequenceable provided that the prime factors of m are all greater than $k!$ where $k = \lambda_0 + \lambda_1 + \dots + \lambda_{e-1}$.*

Proof. Let us assume, by contradiction, that there exists a subset $S = \{x_1, \dots, x_k\}$ of type λ in $\mathbb{Z}_m \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_m \rtimes_{\varphi} H}\}$ that is not sequenceable and all the prime factors of m are larger than $k!$. Here S would be a set of type λ whose elements do not admit an ordering with different partial sums.

This implies that for any possible permutation $\omega \in \text{Sym}(k)$, considered the ordering

$$\mathbf{x}_{\omega} = (x_{\omega(1)}, x_{\omega(2)}, \dots, x_{\omega(k)}),$$

there exist h and t , with $h, t \in [0, k]$, $\{h, t\} \neq \{0, k\}$ and $h \neq t$, such that $s_h(\mathbf{x}_{\omega}) = s_t(\mathbf{x}_{\omega})$, that is, assuming $h < t$,

$$x_{\omega(h+1)} \cdot x_{\omega(h+2)} \dots x_{\omega(t-1)} \cdot x_{\omega(t)} = 0. \quad (6)$$

Therefore we would have a solution $(\pi_{\mathbb{Z}_m}(x_1), \dots, \pi_{\mathbb{Z}_m}(x_k))^T$ to the system of equations derived from (6) by considering the projection over \mathbb{Z}_m . In the following, for simplicity, we will denote $\pi_{\mathbb{Z}_m}(x_j)$ by x_j^1 and, using this notation, we are considering the system:

$$x_{\omega(h+1)}^1 + \varphi_{\pi_H(x_{\omega(h+1)})} x_{\omega(h+2)}^1 + \dots + \varphi_{\pi_H(x_{\omega(h+1)}) \dots \pi_H(x_{\omega(t-1)})} x_{\omega(t)}^1 = 0 \quad (7)$$

where, using the notation of the previous sections, and since φ satisfies property (*), we set $\varphi(a)(x) = \varphi_a x$ for some constant $\varphi_a \in \{1, -1\}$. Denoted by M the $(k!) \times k$ matrix of the coefficients of this system, we note that, since φ satisfies property (*), the coefficients of M belong to $\{-1, 0, 1\}$. It follows that the determinant of any square submatrix of M (that is big at most $k \times k$) is a sum of at most $k!$ terms that belong to ± 1 . Therefore, if all the prime factors of m are larger than $k!$, a square submatrix M' of M is nonsingular in \mathbb{Z}_m if and only if it is nonsingular in \mathbb{Q} .

Following the proof of Theorem 4.12 of [5] and assuming the existence a subset $S = \{x_1, \dots, x_k\}$ of type λ in $\mathbb{Z}_m \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_m \rtimes_{\varphi} H}\}$ that is not sequenceable, we find a subset

of the same type in $\mathbb{Q} \rtimes_{\varphi} H \setminus \{0_{\mathbb{Q} \rtimes_{\varphi} H}\}$ that is also not sequenceable. However, because of Theorem 3.6 and since \mathbb{Q} is a torsion-free abelian group, any subset \tilde{S} of type λ of $G \rtimes_{\varphi} H \setminus \{0_{G \rtimes_{\varphi} H}\}$ is sequenceable. But this is a contradiction and hence the thesis is verified. \square

Corollary 3.15. *Let H be a finite group, let φ satisfy property $(*)$, k be a positive integer, and suppose that, for infinitely many primes p , any subset of size k of $\mathbb{Z}_p \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_p \rtimes_{\varphi} H}\}$ is sequenceable. Then also any subset of size k of $\mathbb{Z}_m \rtimes_{\varphi} H \setminus \{0_{\mathbb{Z}_m \rtimes_{\varphi} H}\}$ is sequenceable provided that the prime factors of m are all greater than $k!$.*

If we consider direct products of type $\mathbb{Z}_m \times H$, we can improve the result of Theorem 3.14. Indeed, with essentially the same proof of Theorem 22 of [5], we can state the following.

Theorem 3.16. *Let H be a finite group, k be a positive integer, and suppose that, for infinitely many primes p , any subset of size k of $\mathbb{Z}_p \times H \setminus \{0_{\mathbb{Z}_p \times H}\}$ is sequenceable. Then also any subset of size k of $\mathbb{Z}_m \times H \setminus \{0_{\mathbb{Z}_m \times H}\}$ is sequenceable provided that the prime factors of m are all greater than $k!/2$.*

As regards dihedral groups, as a consequence of Theorem 2.8, we can prove Theorem 3.17.

Theorem 3.17. *Let m be such that all its prime factors are larger than $k!$. Then any subset S of D_{2m} such that $|S| \leq 12$ is sequenceable.*

Remark 3.18. Because of Remark 2.9, with essentially the same proof of Theorem 3.17, we can prove that if S is a subset of D_{2m} not of type $(k, 0)$, $k \leq 12$ and all the prime factors of m are larger than $k!$, then S admits a linear sequencing.

This implies that, under the hypothesis of Theorem 3.17, S admits a linear sequencing unless s_k is unavoidably the identity.

4 Weak Sequenceability

In this section we consider again subsets S of a group (G, \cdot) of the form $G = \mathbb{Z}_p \rtimes_{\varphi} H$ which is the semidirect product of $(\mathbb{Z}_p, +)$ and another group (H, \cdot) . As in the previous section we indicate by $x \cdot a$ (or by $y \cdot b$), where $x \in \mathbb{Z}_p$ and $a \in H$, a generic element of G . We investigate the t -weak sequenceability of subsets of semidirect products. Recall that in addition to requiring that $x_i - x_j \neq 0$ for $1 \leq i < j \leq k$ and $a_i = a_j$, we seek an ordering having no two of its partial sums s_i, s_j equal for $1 \leq i < j \leq k$ and $|i - j| \leq t$. Hence, modifying the expression of Equation (1), we define, for $t < k$, the following polynomial

$$q_{\mathbf{a}} = \prod_{\substack{1 \leq i < j \leq k \\ a_i = a_j}} (x_j - x_i) \prod_{\substack{0 \leq i < j \leq k \\ b_i = b_j \\ j - i \leq t \\ j \neq i + 1}} (x_{i+1} + \varphi_{a_{i+1}} x_{i+2} + \varphi_{a_{i+1} \cdot a_{i+2}} x_{i+3} + \cdots + \varphi_{a_{i+1} \cdots a_{j-1}} x_j). \quad (8)$$

In this case we have that a set $S = \{x_1 \cdot a_1, x_2 \cdot a_2, \dots, x_k \cdot a_k\} \subseteq G \setminus \{0\}$ of size k is t -weak sequenceable if there exists an ordering $\mathbf{x}_{\mathbf{a}} = (x_1 \cdot a_1, x_2 \cdot a_2, \dots, x_k \cdot a_k)$ of its elements such that $q_{\mathbf{a}}(x_1, \dots, x_k) \neq 0$.

Now, given a set $S = \{x_1 \cdot a_1, x_2 \cdot a_2, \dots, x_k \cdot a_k\}$ of k elements, the idea is to fix, *a priori*, the first h elements $(x_1 \cdot a_1, x_2 \cdot a_2, \dots, x_h \cdot a_h)$, where h is not too big, of the ordering in such a way that no of its partial sums s_i, s_j are equal when $|i - j| \leq t$ and $1 \leq i < j \leq h$. This can be expressed by requiring that $q_{\mathbf{a}'}(x_1, \dots, x_h) \neq 0$ where $\mathbf{a}' = (a_1, \dots, a_h)$ and we show that this can be done due the following Proposition whose proof is inspired by that of Proposition 2.2 of [4].

Proposition 4.1. *Let $S = \{x_1 \cdot a_1, x_2 \cdot a_2, \dots, x_k \cdot a_k\} \subseteq G \setminus \{0\}$ be a set of size k and let h and t be positive integers such that $k - h \geq 2(t - 1)$ and $h \geq t - 1$. Denoted by \bar{a} the element that appears the most in $\pi_2(S)$, we assume that at least $2(t - 1)$ elements of S have \bar{a} as second coordinate. Then there exists an ordering of h -elements of S that we denote, up to relabeling, with $(x_1 \cdot a_1, x_2 \cdot a_2, \dots, x_h \cdot a_h)$, such that*

- (a) $q_{\mathbf{a}'}(x_1, \dots, x_h) \neq 0$ where $\mathbf{a}' = (a_1, \dots, a_h)$;
- (b) $a_h, a_{h-1}, \dots, a_{h-t+2}$ are all equals to \bar{a} ;
- (c) at least $t - 1$ elements of $\{x_{h+1} \cdot a_{h+1}, \dots, x_k \cdot a_k\}$ have \bar{a} as second coordinate.

Proof. Here we choose a set S' of $2(t - 1)$ elements that have \bar{a} as the second coordinate, and we set $S'' := S \setminus S'$. Now we prove that we can choose, inductively on ℓ , $(x_1 \cdot a_1, x_2 \cdot a_2, \dots, x_{h-t} \cdot a_{h-t})$ in S'' and $(x_{h-t+1} \cdot a_{h-t+1}, \dots, x_h \cdot a_h)$ in S' such that, for any $\ell \leq h$, we have $q_{(a_1, \dots, a_\ell)}(x_1, \dots, x_\ell) \neq 0$. Then, properties (b) and (c) will be guaranteed by our choice for $x_1 \cdot a_1, \dots, x_h \cdot a_h$.

BASE CASE: Let $\ell = 1$. Since $q_{(a)}(x) = 1$ for any t and for any $x \cdot a$ of both S' and S'' , the statement is realized for $\ell = 1$.

INDUCTIVE CASE: Let us assume the statement for $\ell \in \{1, \dots, m\}$ and let us prove it for $\ell = m + 1$ where $m + 1 \leq h$. Since the statement is true for $\ell = m$, there exists an m -tuple $(x_1 \cdot a_1, \dots, x_m \cdot a_m)$ such that

$$q_{(a_1, \dots, a_m)}(x_1, \dots, x_m) \neq 0.$$

Given $x \cdot a \in G$, we set $s_m := (x_1 \cdot a_1) \cdots (x_m \cdot a_m) = y_m \cdot b_m$ and we define y and s such that $s = s_m \cdot (x \cdot a) = y \cdot b$ where $b = b_m \cdot a$. Then we have

$$\begin{aligned} & \frac{q_{(a_1, \dots, a_m, a)}(x_1, \dots, x_m, x)}{q_{(a_1, \dots, a_m)}(x_1, \dots, x_m)} = \\ &= \prod_{1 \leq i < m+1, a=a_i} (x - x_i) \prod_{\substack{0 \leq i < m \\ m+1-i \leq t, b_i=b}} (x_{i+1} + \varphi_{a_{i+1}} x_{i+2} + \varphi_{a_{i+1} \cdot a_{i+2}} x_{i+3} + \cdots + \varphi_{a_{i+1} \dots a_m} x). \end{aligned}$$

Here, any element $x \cdot a$ of $S \setminus \{x_1 \cdot a_1, \dots, x_m \cdot a_m\}$ satisfies $\prod_{1 \leq i < m+1} (x - x_i) \neq 0$ if $a = a_i$. Hence, to have $\frac{q_{(a_1, \dots, a_m)}(x_1, \dots, x_m, x)}{q_{(a_1, \dots, a_m, a)}(x_1, \dots, x_m)} \neq 0$, it suffice to find x such that

$$\prod_{\max(0, m+1-t) \leq i < m, b_i=b} (x_{i+1} + \varphi_{a_{i+1}} x_{i+2} + \varphi_{a_{i+1} \cdot a_{i+2}} x_{i+3} + \cdots + \varphi_{a_{i+1} \dots a_m} x) \neq 0.$$

Note that there is at most one $x \cdot a \in S \setminus \{x_1 \cdot a_1, \dots, x_m \cdot a_m\}$ such that

$$(x_{i+1} + \varphi_{a_{i+1}} x_{i+2} + \varphi_{a_{i+1} \cdot a_{i+2}} x_{i+3} + \cdots + \varphi_{a_{i+1} \dots a_m} x) = 0$$

and $b_i = b$. Indeed this would imply that $s_i = s$ in G that is $(x_{i+1} \cdot a_{i+1}) \cdot \dots \cdot (x_m \cdot a_m) \cdot (x \cdot a) = 0$ which is satisfied by exactly one element $x \cdot a$ of G . Since those relations are at most $t - 1$, we have at most $t - 1$ values $x \cdot a$ in $S \setminus \{x_1 \cdot a_1, \dots, x_m \cdot a_m\}$ such that

$$\prod_{\max(0, m+1-t) \leq i < m, b_i = b} (x_{i+1} + \varphi_{a_{i+1}} x_{i+2} + \varphi_{a_{i+1} \cdot a_{i+2}} x_{i+3} + \dots + \varphi_{a_{i+1} \dots a_m} x) = 0.$$

Here we have that either $h - t + 1 > m$ or $h - t + 1 \leq m$. In the first case, we have that

$$|S'' \setminus \{x_1 \cdot a_1, \dots, x_m \cdot a_m\}| = k - 2(t - 1) - m \geq t + 1$$

where the inequality holds since $k - 2(t - 1) - m = (k - m) - 2(t - 1)$ and $k - m \geq k - h \geq 2(t - 1)$. It follows that there exists $x \cdot a \in S'' \setminus \{x_1 \cdot a_1, \dots, x_m \cdot a_m\}$ such that

$$\prod_{\max(0, m+1-t) \leq i < m, b_i = b} (x_{i+1} + \varphi_{a_{i+1}} x_{i+2} + \varphi_{a_{i+1} \cdot a_{i+2}} x_{i+3} + \dots + \varphi_{a_{i+1} \dots a_m} x) \neq 0.$$

In the second case, we have that

$$|S' \setminus \{x_1 \cdot a_1, \dots, x_m \cdot a_m\}| = 2(t - 1) - (m - (h - t + 1)) \geq t$$

where the last inequality holds since $m \leq h - 1$ and hence $2(t - 1) - (m - (h - t + 1)) \geq 2(t - 1) - (h - 1 - (h - t + 1)) = t$. It follows that also in this case, there exists $x \cdot a \in S' \setminus \{x_1 \cdot a_1, \dots, x_m \cdot a_m\}$ such that

$$\prod_{\max(0, m+1-t) \leq i < m, b_i = b} (x_{i+1} + \varphi_{a_{i+1}} x_{i+2} + \varphi_{a_{i+1} \cdot a_{i+2}} x_{i+3} + \dots + \varphi_{a_{i+1} \dots a_m} x) \neq 0.$$

In both cases, we can find $x_{m+1} \cdot a_{m+1}$ that satisfies property (b) and (c) and which it is such that

$$\frac{q_{(a_1, \dots, a_m, a_{m+1})}(x_1, \dots, x_m, x_{m+1})}{q_{(a_1, \dots, a_m)}(x_1, \dots, x_m)} = \prod_{1 \leq i < m+1, a_{m+1} = a_i} (x_{m+1} - x_i).$$

$$\cdot \prod_{\max(0, m+1-t) \leq i < m, b_i = b_{m+1}} (x_{i+1} + \varphi_{a_{i+1}} x_{i+2} + \varphi_{a_{i+1} \cdot a_{i+2}} x_{i+3} + \dots + \varphi_{a_{i+1} \dots a_m} x_{m+1}) \neq 0.$$

Since \mathbb{Z}_p is a field and due to the inductive hypothesis $q_{(a_1, \dots, a_m, a)}(x_1, \dots, x_m) \neq 0$, we also have that

$$\frac{q_{(a_1, \dots, a_m, a_{m+1})}(x_1, \dots, x_m, x_{m+1})}{q_{(a_1, \dots, a_m)}(x_1, \dots, x_m)} \cdot q_{(a_1, \dots, a_m)}(x_1, \dots, x_m) = q_{(a_1, \dots, a_m, a_{m+1})}(x_1, \dots, x_{m+1}) \neq 0 \quad (9)$$

which completes the proof. \square

Remark 4.2. Note that the hypothesis that at least $2(t - 1)$ elements of S has the same second coordinate is always satisfied when

$$k > (2t - 3) \frac{|G|}{p} = (2t - 3)|H|.$$

This means that the approach of Proposition 4.1 can be always tried assuming that k is big enough. For smaller values of k one could fix the first elements differently, as done for example in Proposition 2.2 of [4], but, since then one should consider a huge number of cases (see also the following Remark 4.3), we consider explicitly only this asymptotic case.

In the following we assume that we have fixed, according to Proposition 4.1,

$$\{x_1 \cdot a_1, x_2 \cdot a_2, \dots, x_h \cdot a_h\} \subseteq S$$

that satisfies properties (a), (b) and (c) of such proposition. We note that every $x \cdot a \in S \setminus \{x_1 \cdot a_1, x_2 \cdot a_2, \dots, x_h \cdot a_h\}$ is such that $x - x_i \neq 0$ for any $i \in \{1, \dots, h\}$ such that $a = a_i$. Therefore, if \mathbf{a} is a quotient sequencing that extends \mathbf{a}'' , it is left to find a nonzero point for the polynomial

$$\frac{q_{\mathbf{a}}(x_1, \dots, x_h, x_{h+1}, \dots, x_k)}{q_{\mathbf{a}'}(x_1, \dots, x_h) \prod_{1 \leq i \leq h < j \leq k, a_j = a_i} (x_j - x_i)}.$$

Since the free variables are now x_{h+1}, \dots, x_k , we set $\ell := k - h$ and $z_i := x_{i+h}$; here the constrain $k - h \geq 2(t - 1)$ of Proposition 4.1 becomes $\ell \geq 2(t - 1)$. Then we denote by $h_{\mathbf{a}, \ell}$ the polynomial

$$h_{\mathbf{a}, \ell}(z_1, \dots, z_{\ell}) := \frac{q_{\mathbf{a}}(x_1, \dots, x_h, z_1, \dots, z_{\ell})}{q_{\mathbf{a}'}(x_1, \dots, x_h) \prod_{1 \leq i \leq 1 < j \leq \ell, a_j = a_i} (z_j - x_i)}. \quad (10)$$

Since we are also assuming that $k - \ell \geq t - 1$, that is, $k - (t - 1) \geq \ell \geq 2(t - 1)$, denoting by \mathbf{a}'' the quotient sequencing given by the last ℓ elements of \mathbf{a} , we obtain the following expression

$$\begin{aligned} h_{\mathbf{a}, \ell}(z_1, \dots, z_{\ell}) &= q_{\mathbf{a}''}(z_1, \dots, z_{\ell}) \cdot \\ &\prod_{\substack{0 \leq i \leq t-1; \\ 1 \leq j \leq t-i-1; \\ b_{k-\ell-i-1} = b_{(k-\ell)+j}}} (x_{k-\ell-i} + \varphi_{\bar{a}} x_{k-\ell-i+1} + \dots + \varphi_{\bar{a}^i} x_{k-\ell} + \varphi_{\bar{a}^{i+1}} z_1 + \dots + \varphi_{\bar{a}^{i+1} \cdot a_{k-\ell+1} \dots a_{k-\ell+j-1}} z_j). \end{aligned} \quad (11)$$

Now we aim to apply the Non-Vanishing Corollary (of the Combinatorial Nullstellensatz) to the polynomial $h_{\mathbf{a}, \ell}$. For this purpose, it is enough to consider the terms of $h_{\mathbf{a}, \ell}$ of maximal degree in the variables z_1, \dots, z_{ℓ} that are the ones where no x_i appears. We denote by $r_{\mathbf{a}, \ell}$ the polynomial given by those terms, that is

$$\begin{aligned} r_{\mathbf{a}, \ell} &:= q_{\mathbf{a}''}(z_1, \dots, z_{\ell}) \cdot \\ &\prod_{\substack{0 \leq i \leq t-1; \\ 1 \leq j \leq t-i-1; \\ b_{k-\ell-i-1} = b_{(k-\ell)+j}}} (\varphi_{\bar{a}^{i+1}} z_1 + \varphi_{\bar{a}^{i+1} \cdot a_{k-\ell+1}} z_2 + \dots + \varphi_{\bar{a}^{i+1} \cdot a_{k-\ell+1} \dots a_{k-\ell+j-1}} z_j). \end{aligned} \quad (12)$$

Since $b_{k-\ell-i-1} = b_{(k-\ell)+j}$ if and only if

$$a_{k-\ell-i} \cdot a_{k-\ell-1} \cdot \dots \cdot a_{k-\ell} \cdot a_{k-\ell+1} \cdot \dots \cdot a_{k-\ell+j} = id$$

and $a_{k-\ell-i} = a_{k-\ell-1} = \dots = a_{k-\ell} = \bar{a}$, we can state the following, simple but very powerful, remark.

Remark 4.3. Since we are assuming that $a_h, a_{h-1}, \dots, a_{h-t+1}$ are all equals to \bar{a} , the expression of $r_{\mathbf{a},\ell}$ does not depend on all the quotient sequence \mathbf{a} but only by \mathbf{a}'' and \bar{a} . Here property *b* of Proposition 4.1 is crucial since, otherwise, given \mathbf{a}'' , we would have $|H|^{t-1}$ possibilities for the polynomial $r_{\mathbf{a},\ell}(z_1, \dots, z_\ell)$ according to the sequence $a_h, a_{h-1}, \dots, a_{h-t+1}$.

In the following, we just denote this polynomial by $r_{\mathbf{a}'',\bar{a},\ell}$.

Indeed Remark 4.3 means that, after these manipulations, we are left to consider a polynomial that does not depend on $k = |S|$, and hence we have chances to get a result that is very general on k .

To apply the Non-Vanishing Corollary, we also need to find, for any quotient sequencing \mathbf{a}'' of length ℓ , a nonzero coefficient in $r_{\mathbf{a}'',\bar{a},\ell}$ that divides the bounding monomial. Moreover, due to property *c* of Proposition 4.1, we may assume that \bar{a} appears at least $t - 1$ times in \mathbf{a}'' .

4.1 Computational Results

As done for the sequenceability problem in the more general setting, we consider here groups of type D_{2p} and G_{3p} . Also, since the case $G = \mathbb{Z}_p \times \mathbb{Z}_e$ has not been considered in [4], we consider these groups here.

Example 4.4. *Example* Let $t = 4$, $\ell = 2(t - 1) = 6$ and $\bar{a} = 0$. Let p be a prime and suppose $S \subseteq D_{2p} \setminus \{0\}$ with $|S| = k \geq 11$ and suppose that S has at least 6 elements with coset \bar{a} . Let $S' = S \setminus \{x_1 \cdot a_1, \dots, x_{k-\ell} \cdot a_{k-\ell}\}$ be of type $(3, 3)$. The sequence $\mathbf{a}'' = (1, 0, 1, 0, 1, 0)$ has partial sums $(0, 1, 1, 0, 0, 1, 1)$.

By Proposition 4.1, we desire a sequencing of S' of the form

$$(z_1 \cdot 1, z_2 \cdot 0, z_3 \cdot 1, z_4 \cdot 0, z_5 \cdot 1, z_6 \cdot 0).$$

The polynomial we need to study is the one defined in (12). Hence we get

$$\begin{aligned} r_{\mathbf{a}'',\bar{a},\ell} &= (z_1 - z_2 - z_3)(z_3 - z_1)(z_5 - z_1)(z_4 - z_2)(z_6 - z_2)(z_5 - z_3)(z_6 - z_4) \\ &(z_1 - z_2 - z_3)(z_1 - z_2 - z_3 + z_4)(z_2 + z_3 - z_4 - z_5)(z_3 - z_4 - z_5)(z_3 - z_4 - z_5 + z_6) \end{aligned}$$

To apply the Non-Vanishing Corollary we need a monomial of this polynomial which divides the bounding monomial $z_1^2 z_2^2 z_3^2 z_4^2 z_5^2 z_6^2$ with a nonzero coefficient. One such is $z_1^2 z_2^2 z_3^2 z_4^2 z_5^2 z_6^2$, which has coefficient -12 . Hence whenever S has this form it is t -weakly sequenceable.

We are now ready to prove the main result of this section.

Theorem 4.5. *Let $n = pe$ with $p > 3$ prime and let G be a group of size n . Then subsets S of size k of G are t -weakly sequenceable whenever $e \in \{1, 2, 3\}$, k is large enough and $t \leq 6$.*

Proof. The case $e = 1$ follows from the results of [4].

As already noted in the proof of Theorem 2.8, a group of size $2p$ is either $\mathbb{Z}_p \times \mathbb{Z}_2$ or the dihedral group D_{2p} and a group of size $3p$ is either $\mathbb{Z}_p \times \mathbb{Z}_3$ or G_{3p} for some value of r . Here, due to Remark 4.2, if $k \geq (2t - 3)e + 1$, S admits a coset whose size is at least

$2(t-1)$ and hence, due to Proposition 4.1, we can apply Non-Vanishing Corollary to the polynomial of Equation (12) for suitable quotient sequences and nonzero monomial (see Tables 3, 4, 5 and 6 below). These results are obtained using SageMath [19].

Here we note that the coefficients presented in Table 6 can be zero only for finitely many primes p (see also the discussion in the proof of Theorem 2.8). On the other hand, since $3p > k$, if k is large enough all these coefficients must be nonzero. \square

The following Tables 3, 4, 5 and 6 contain the required monomials and their coefficients for the proof of Theorem 4.5 in the cases $G \in \{\mathbb{Z}_p \times \mathbb{Z}_2, \mathbb{Z}_p \times \mathbb{Z}_3, D_{2p}, G_{3p}\}$.

Table 3: Monomials and their coefficients sufficient for the proof of Theorem 4.5 in the case $t = 6$ and $G = \mathbb{Z}_p \times \mathbb{Z}_2$.

\bar{a}	λ	\mathbf{a}''	deg	monomial/s	coefficient/s
0	(5, 5)	(0, 1, 0, 0, 1, 1, 1, 0, 0, 1)	40	$x_1^4 x_2^4 x_3^4 x_4^4 x_5^4 x_6^4 x_7^4 x_8^4 x_9^4 x_{10}^4$	$-5 \cdot 467$
0	(6, 4)	(1, 0, 0, 0, 1, 0, 1, 0, 1, 0)	38	$x_2^4 x_3^5 x_4^5 x_5^3 x_6^5 x_7^3 x_8^5 x_9^3 x_{10}^5$ $x_1 x_2^3 x_3^5 x_4^5 x_5^3 x_6^5 x_7^3 x_8^5 x_9^3 x_{10}^5$	-1063 $2^4 \cdot 503$
0	(7, 3)	(1, 0, 0, 0, 1, 0, 0, 0, 1, 0)	36	$x_3^2 x_4^6 x_5^2 x_6^6 x_7^6 x_8^6 x_9^2 x_{10}^6$	2
0	(8, 2)	(1, 0, 0, 0, 1, 0, 0, 0, 0, 0)	45	$x_3^2 x_4^7 x_5 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}^7$	-2^2
0	(9, 1)	(0, 0, 0, 0, 1, 0, 0, 0, 0, 0)	66	$x_1^5 x_2^7 x_3^7 x_4^8 x_6^7 x_7^8 x_8^8 x_9^8 x_{10}^8$	$-2^5 \cdot 3$
1	(4, 6)	(0, 1, 1, 1, 0, 0, 1, 0, 1, 1)	42	$x_1^3 x_2^5 x_3^5 x_4^5 x_5^3 x_6^3 x_7^5 x_8^3 x_9^5 x_{10}^5$	-1187
1	(3, 7)	(1, 0, 1, 1, 1, 1, 1, 0, 1, 0)	47	$x_1^5 x_2^2 x_3^6 x_4^6 x_5^6 x_6^6 x_7^6 x_8^2 x_9^6 x_{10}^2$ $x_1^6 x_2 x_3^6 x_4^6 x_5^6 x_6^6 x_7^6 x_8^2 x_9^6 x_{10}^2$	$2 \cdot 3^2 \cdot 1511$ $23 \cdot 499$
1	(2, 8)	(1, 0, 1, 1, 1, 1, 1, 1, 1, 0)	54	$x_1^4 x_2 x_3^6 x_4^7 x_5^7 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}$ $x_1^5 x_2 x_3^5 x_4^7 x_5^7 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}$	$-2^5 \cdot 5^2$ -4337
1	(1, 9)	(1, 0, 1, 1, 1, 1, 1, 1, 1, 1)	62	$x_1^5 x_3^3 x_4^6 x_5^8 x_6^8 x_7^8 x_8^8 x_9^8 x_{10}^8$	-2
1	(0, 10)	(1, 1, 1, 1, 1, 1, 1, 1, 1, 1)	75	$x_1^3 x_2^4 x_3^5 x_4^9 x_5^9 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^9$ $x_1^3 x_2^5 x_3^5 x_4^8 x_5^9 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^9$	$-2^4 \cdot 5^2$ $-7 \cdot 853$

Table 4: Monomials and their coefficients sufficient for the proof of Theorem 4.5 in the case $t = 6$ and $G = \mathbb{Z}_p \times \mathbb{Z}_3$.

\bar{a}	λ	\mathbf{a}''	deg	monomial/s	coefficient/s
0	(5, 3, 2)	(1, 0, 0, 1, 0, 0, 0, 2, 1, 2)	25	$x_1^2 x_2 x_3^4 x_4^2 x_5^4 x_6^4 x_7^4 x_8^2 x_9 x_{10}$	$2 \cdot 3^2$
0	(5, 4, 1)	(1, 0, 0, 1, 0, 0, 0, 1, 1, 2)	22	$x_3^4 x_4 x_5^3 x_6^4 x_7^3 x_8^3 x_9^3$	1
0	(5, 5, 0)	(0, 1, 0, 1, 0, 0, 0, 1, 1, 1)	33	$x_1^2 x_2 x_3^4 x_4^4 x_5^4 x_6^4 x_7^4 x_8^2 x_9^4 x_{10}^4$	$3 \cdot 5^2$
0	(6, 2, 2)	(1, 0, 0, 1, 0, 0, 0, 0, 2, 2)	25	$x_3^3 x_4 x_5^4 x_6^5 x_7^5 x_8^5 x_9 x_{10}$	-1
0	(6, 3, 1)	(1, 0, 0, 1, 0, 0, 0, 0, 1, 2)	30	$x_2 x_3^5 x_4^2 x_5^5 x_6^5 x_7^5 x_8^5 x_9^2$	-2^4
0	(6, 4, 0)	(1, 0, 0, 1, 0, 0, 0, 0, 1, 1)	28	$x_3^3 x_4 x_5^4 x_6^5 x_7^5 x_8^2 x_9^3 x_{10}^3$	-1

\bar{a}	λ	\mathbf{a}''	deg	monomial/s	coefficient/s
0	(7, 2, 1)	(1, 0, 0, 1, 0, 0, 0, 0, 0, 2)	33	$x_3^3 x_4 x_5^5 x_6^6 x_7^6 x_8^6 x_9^6$	-2^2
0	(7, 3, 0)	(1, 0, 0, 1, 0, 0, 0, 0, 0, 1)	35	$x_3^3 x_4 x_5^5 x_6^6 x_7^6 x_8^6 x_9^6 x_{10}^2$	-2^2
0	(8, 1, 1)	(1, 0, 0, 0, 2, 0, 0, 0, 0, 0)	44	$x_3^2 x_4^7 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}^7$	-2^2
0	(8, 2, 0)	(1, 0, 0, 0, 1, 0, 0, 0, 0, 0)	42	$x_3^2 x_4^5 x_5 x_6^6 x_7^7 x_8^7 x_9^7 x_{10}^7$	-2^2
0	(9, 1, 0)	(0, 0, 0, 1, 0, 0, 0, 0, 0, 0)	66	$x_1^5 x_2^6 x_3^8 x_5^7 x_6^8 x_7^8 x_8^8 x_9^8 x_{10}^8$	$2^2 \cdot 7$
1	(3, 5, 2)	(0, 1, 1, 0, 1, 1, 1, 2, 0, 2)	27	$x_1 x_2^4 x_3^4 x_4^2 x_5^4 x_6^4 x_7^4 x_8 x_9^2 x_{10}$ $x_1^2 x_2^3 x_3^4 x_4^2 x_5^4 x_6^4 x_7^4 x_8 x_9^2 x_{10}$	-73 -233
1	(2, 5, 3)	(0, 1, 0, 1, 2, 1, 1, 1, 2, 2)	28	$x_1 x_2^4 x_3 x_4^4 x_5^2 x_6^4 x_7^4 x_8^4 x_9^2 x_{10}^2$	7
1	(4, 5, 1)	(0, 1, 1, 0, 1, 1, 1, 0, 0, 2)	31	$x_1^2 x_2^4 x_3^4 x_4^3 x_5^4 x_6^4 x_7^4 x_8^3 x_9^3$ $x_1^3 x_2^3 x_3^4 x_4^3 x_5^4 x_6^4 x_7^4 x_8^3 x_9^3$	$-5 \cdot 43$ $-3 \cdot 13 \cdot 17$
1	(1, 5, 4)	(1, 2, 2, 0, 2, 2, 1, 1, 1, 1)	32	$x_1^4 x_2^3 x_3^3 x_4^3 x_5^3 x_6^4 x_7^4 x_8^4 x_9^4 x_{10}^4$	-7
1	(0, 5, 5)	(1, 2, 1, 2, 1, 1, 1, 2, 2, 2)	38	$x_1^3 x_2^4 x_3^4 x_4^4 x_5^4 x_6^3 x_7^4 x_8^4 x_9^4 x_{10}^4$	$-2 \cdot 3 \cdot 5$
1	(2, 6, 2)	(0, 1, 0, 1, 1, 2, 2, 1, 1, 1)	30	$x_1 x_2^2 x_3 x_4^4 x_5^5 x_6 x_7 x_8^5 x_9^5 x_{10}^5$	-5
1	(3, 6, 1)	(0, 1, 0, 1, 1, 0, 2, 1, 1, 1)	32	$x_1 x_2^2 x_3^2 x_4^5 x_5^5 x_6^2 x_8^5 x_9^5 x_{10}^5$	2^2
1	(1, 6, 3)	(0, 1, 0, 1, 1, 0, 2, 1, 1, 1)	32	$x_1 x_2^2 x_3^2 x_4^5 x_5^5 x_6^2 x_8^5 x_9^5 x_{10}^5$	2^2
1	(4, 6, 0)	(0, 1, 1, 0, 1, 1, 1, 1, 0, 0)	34	$x_1 x_2 x_3^3 x_4^3 x_5^5 x_6^5 x_7^5 x_8^3 x_9^3 x_{10}^3$	$-2 \cdot 3$
1	(0, 6, 4)	(2, 1, 1, 2, 1, 1, 1, 1, 2, 2)	37	$x_1^2 x_2^2 x_3^4 x_4^3 x_5^5 x_6^5 x_7^5 x_8^3 x_9^3 x_{10}^3$	2^2
1	(2, 7, 1)	(0, 1, 1, 0, 1, 1, 1, 1, 1, 2)	35	$x_1 x_2 x_3^3 x_4 x_5^5 x_6^6 x_7^6 x_8^6 x_9^6$	-2
1	(1, 7, 2)	(1, 0, 1, 2, 1, 1, 1, 1, 1, 2)	40	$x_1^3 x_3^5 x_4 x_5^6 x_6^6 x_7^6 x_8^6 x_9^6 x_{10}$	3^2
1	(3, 7, 0)	(0, 1, 1, 0, 1, 1, 1, 1, 1, 0)	36	$x_1 x_2 x_3^2 x_4^2 x_5^4 x_6^6 x_7^6 x_8^6 x_9^6 x_{10}^2$	1
1	(0, 7, 3)	(2, 1, 1, 2, 1, 1, 1, 1, 1, 2)	40	$x_1^2 x_2^2 x_3^3 x_4^2 x_5^5 x_6^6 x_7^6 x_8^6 x_9^6 x_{10}^2$	-2^2
1	(1, 8, 1)	(1, 0, 1, 1, 1, 1, 1, 1, 1, 2)	43	$x_1^3 x_3^3 x_4^3 x_5^6 x_6^7 x_7^7 x_8^7 x_9^7$	1
1	(2, 8, 0)	(1, 0, 1, 1, 1, 1, 1, 1, 1, 0)	43	$x_1^2 x_2 x_3^3 x_4^3 x_5^5 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}$	-1
1	(0, 8, 2)	(1, 2, 1, 1, 1, 1, 1, 1, 1, 2)	47	$x_1^3 x_2 x_3^4 x_4^3 x_5^7 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}$	2^2
1	(1, 9, 0)	(1, 0, 1, 1, 1, 1, 1, 1, 1, 1)	51	$x_1^3 x_2^2 x_3^4 x_4^5 x_5^8 x_6^8 x_7^8 x_8^8 x_9^8 x_{10}$	2
1	(0, 9, 1)	(1, 2, 1, 1, 1, 1, 1, 1, 1, 1)	54	$x_1^2 x_2^2 x_3^3 x_4^3 x_5^5 x_6^6 x_7^6 x_8^9 x_9^9 x_{10}^9$	2
1	(0, 10, 0)	(1, 1, 1, 1, 1, 1, 1, 1, 1, 1)	65	$x_1^3 x_2^4 x_3^4 x_4^4 x_5^6 x_6^8 x_7^8 x_8^9 x_9^9 x_{10}^9$	$2 \cdot 5$

\bar{a}	λ	\mathbf{a}''	deg	monomial/s	coefficient/s
2	(3, 2, 5)	(0, 2, 2, 0, 2, 2, 2, 1, 0, 1)	27	$x_1x_2^4x_3^4x_4^2x_5^4x_6^4x_7^4x_8x_9^2x_{10}$ $x_1^2x_2^3x_3^4x_4^2x_5^4x_6^4x_7^4x_8x_9^2x_{10}$	-73 -233
2	(2, 3, 5)	(0, 2, 0, 2, 1, 2, 2, 2, 1, 1)	28	$x_1x_2^4x_3x_4^4x_5^2x_6^4x_7^4x_8^4x_9^2x_{10}^2$	7
2	(4, 1, 5)	(0, 2, 2, 0, 2, 2, 2, 0, 0, 1)	31	$x_1^2x_2^4x_3^4x_4^3x_5^4x_6^4x_7^4x_8^3x_9^3$ $x_1^3x_2^3x_3^4x_4^3x_5^4x_6^4x_7^4x_8^3x_9^3$	-5 · 43 -3 · 13 · 17
2	(1, 4, 5)	(2, 1, 1, 0, 1, 1, 2, 2, 2, 2)	32	$x_1^4x_2^3x_3^3x_4^3x_5^3x_6^4x_7^4x_8^4x_9^4x_{10}$	-7
2	(5, 0, 5)	(0, 2, 0, 2, 0, 0, 0, 2, 2, 2)	33	$x_1^2x_2x_3^4x_4^4x_5^4x_6^4x_7^4x_8^2x_9^4x_{10}^4$	$3 \cdot 5^2$
2	(2, 2, 6)	(0, 2, 0, 2, 2, 1, 1, 2, 2, 2)	30	$x_1x_2^2x_3x_4^4x_5^5x_6x_7x_8^5x_9^5x_{10}^5$	-5
2	(3, 1, 6)	(0, 2, 0, 2, 2, 0, 1, 2, 2, 2)	32	$x_1x_2^2x_3^2x_4^5x_5^5x_6^2x_8^5x_9^5x_{10}^5$	2^2
2	(1, 3, 6)	(0, 2, 0, 2, 2, 0, 1, 2, 2, 2)	32	$x_1x_2^2x_3^2x_4^5x_5^5x_6^2x_8^5x_9^5x_{10}^5$	2^2
2	(4, 0, 6)	(0, 2, 2, 0, 2, 2, 2, 2, 0, 0)	34	$x_1x_2x_3^3x_4^3x_5^5x_6^5x_7^5x_8^3x_9^3x_{10}^3$	-2 · 3
2	(0, 4, 6)	(1, 2, 2, 1, 2, 2, 2, 2, 1, 1)	37	$x_1^2x_2^2x_3^4x_4^3x_5^5x_6^5x_7^5x_8^3x_9^3x_{10}^3$	2^2
2	(2, 1, 7)	(0, 2, 2, 0, 2, 2, 2, 2, 2, 1)	35	$x_1x_2x_3^3x_4x_5^6x_6^6x_7^6x_8^6x_9^6$	-2
2	(1, 2, 7)	(2, 0, 2, 1, 2, 2, 2, 2, 2, 1)	40	$x_1^3x_3^5x_4x_5^6x_6^6x_7^6x_8^6x_9^6x_{10}$	3^2
2	(3, 0, 7)	(0, 2, 2, 0, 2, 2, 2, 2, 2, 0)	36	$x_1x_2x_3^2x_4^2x_5^4x_6^6x_7^6x_8^6x_9^6x_{10}^2$	1
2	(0, 3, 7)	(1, 2, 2, 1, 2, 2, 2, 2, 2, 1)	40	$x_1^2x_2^2x_3^3x_4^2x_5^6x_6^6x_7^6x_8^6x_9^6x_{10}^2$	- 2^2
2	(1, 1, 8)	(2, 0, 2, 2, 2, 2, 2, 2, 2, 1)	43	$x_1^3x_3^3x_4^3x_5^6x_6^7x_7^7x_8^7x_9^7$	1
2	(2, 0, 8)	(2, 0, 2, 2, 2, 2, 2, 2, 2, 0)	43	$x_1^2x_2x_3^3x_4^3x_5^5x_6^7x_7^7x_8^7x_9^7x_{10}$	-1
2	(0, 2, 8)	(2, 1, 2, 2, 2, 2, 2, 2, 2, 1)	47	$x_1^3x_2x_3^4x_4^3x_5^7x_6^7x_7^7x_8^7x_9^7x_{10}$	2^2
2	(1, 0, 9)	(2, 0, 2, 2, 2, 2, 2, 2, 2, 2)	51	$x_1^3x_2^2x_3^4x_4^5x_5^5x_6^8x_7^8x_8^8x_9^8x_{10}$	2
2	(0, 1, 9)	(2, 1, 2, 2, 2, 2, 2, 2, 2, 2)	54	$x_1^2x_2^2x_3^3x_4^3x_5^6x_6^6x_7^9x_8^9x_9^9x_{10}$	2
2	(0, 0, 10)	(2, 2, 2, 2, 2, 2, 2, 2, 2, 2)	65	$x_1^3x_2^4x_3^4x_4^4x_5^6x_6^8x_7^9x_8^9x_9^9x_{10}$	$2 \cdot 5$

Table 5: Monomials and their coefficients sufficient for the proof of Theorem 4.5 in the case $t = 6$ and $G = D_{2p}$.

\bar{a}	λ	\mathbf{a}''	deg	monomial/s	coefficient/s
0	(5, 5)	(1, 0, 0, 0, 1, 1, 1, 0, 1, 0)	34	$x_2^2x_3^4x_4^4x_5^4x_6^4x_7^4x_8^4x_9^4x_{10}$ $x_2^3x_3^3x_4^4x_5^4x_6^4x_7^4x_8^4x_9^4x_{10}$	$2^6 \cdot 5 \cdot 13$ $2^4 \cdot 17 \cdot 23$
0	(6, 4)	(1, 0, 0, 0, 1, 0, 1, 0, 1, 0)	38	$x_2^4x_3^5x_4^5x_5^3x_6^5x_7^3x_8^3x_9^5x_{10}$ $x_1x_2^3x_3^5x_4^5x_5^3x_6^5x_7^3x_8^3x_9^5x_{10}$	-3 · 41543 $2^2 \cdot 3 \cdot 31 \cdot 617$

\bar{a}	λ	\mathbf{a}''	deg	monomial/s	coefficient/s
0	(7, 3)	(1, 0, 0, 0, 1, 0, 0, 0, 1, 0)	36	$x_3^2 x_4^6 x_5^2 x_6^6 x_7^6 x_8^2 x_9^6 x_{10}^6$	$-2^3 \cdot 3$
0	(8, 2)	(1, 0, 0, 0, 1, 0, 0, 0, 0, 0)	45	$x_3^2 x_4^7 x_5 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}^7$	-2^2
0	(9, 1)	(0, 0, 0, 0, 1, 0, 0, 0, 0, 0)	66	$x_1^5 x_2^7 x_3^7 x_4^8 x_6^7 x_7^8 x_8^8 x_9^8 x_{10}^8$	$-2^5 \cdot 3$
1	(5, 5)	(1, 0, 1, 1, 1, 0, 0, 0, 1, 0)	39	$x_1^3 x_2^4 x_3^4 x_4^4 x_5^4 x_6^4 x_7^4 x_8^4 x_9^4 x_{10}^4$ $x_1^4 x_2^3 x_3^4 x_4^4 x_5^4 x_6^4 x_7^4 x_8^4 x_9^4 x_{10}^4$	$43 \cdot 97$ $-2 \cdot 3^3 \cdot 257$
1	(4, 6)	(0, 1, 1, 1, 1, 1, 0, 0, 1, 0)	39	$x_1^2 x_2^3 x_3^5 x_4^5 x_5^5 x_6^5 x_7^3 x_8^3 x_9^3 x_{10}^3$ $x_1^3 x_2^2 x_3^5 x_4^5 x_5^5 x_6^5 x_7^3 x_8^3 x_9^3 x_{10}^3$	$-11 \cdot 23 \cdot 61$ $-7^2 \cdot 953$
1	(3, 7)	(1, 0, 1, 1, 1, 1, 1, 0, 1, 0)	43	$x_1^3 x_2^4 x_3^6 x_4^6 x_5^6 x_6^6 x_7^2 x_8^2 x_9^2 x_{10}^2$ $x_1^4 x_2^3 x_3^6 x_4^6 x_5^6 x_6^6 x_7^2 x_8^2 x_9^2 x_{10}^2$	$-5 \cdot 23 \cdot 61$ $2 \cdot 32117$
1	(2, 8)	(1, 0, 1, 1, 1, 1, 1, 1, 1, 0)	48	$x_1^4 x_2 x_3^4 x_4^7 x_5^7 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}^7$ $x_1^4 x_2 x_3^3 x_4^5 x_6^6 x_7^7 x_8^7 x_9^7 x_{10}^7$	$-2^2 \cdot 5 \cdot 61$ $2^4 \cdot 3^2 \cdot 29$
1	(1, 9)	(1, 0, 1, 1, 1, 1, 1, 1, 1, 1)	55	$x_1^5 x_3^3 x_4^4 x_5^7 x_6^8 x_7^8 x_8^8 x_9^8 x_{10}^8$ $x_1^5 x_3^3 x_4^4 x_5^6 x_6^8 x_7^8 x_8^8 x_9^8 x_{10}^8$	$2 \cdot 3 \cdot 61$ $-2^4 \cdot 83$
1	(0, 10)	(1, 1, 1, 1, 1, 1, 1, 1, 1, 1)	66	$x_1^4 x_2^4 x_3^8 x_4^9 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^9$ $x_1^4 x_2^4 x_3^9 x_4^8 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^9$	$-2 \cdot 457$ $-2^8 \cdot 3^2$

Table 6: Monomials and their coefficients sufficient for the proof of Theorem 4.5 in the case $t = 6$ and $G = G_{3p}$.

\bar{a}	λ	\mathbf{a}''	deg	monomial/s	coefficient/s
0	(5, 3, 2)	(1, 0, 0, 1, 0, 0, 0, 2, 1, 2)	25	$x_1^2 x_2^4 x_3^4 x_4^2 x_5^2 x_6^4 x_7^4 x_8^2 x_{10}^2$	$-r - 1$
0	(5, 4, 1)	(1, 0, 0, 1, 0, 0, 0, 1, 1, 2)	22	$x_1^3 x_2^4 x_3^3 x_4^3 x_5 x_6^2 x_7^4 x_8 x_9$	1
0	(5, 5, 0)	(0, 1, 0, 1, 0, 0, 0, 1, 1, 1)	33	$x_1^2 x_2^2 x_3^2 x_4^3 x_5^4 x_6^4 x_7^4 x_8^4 x_9^4 x_{10}^4$	$124r + 294$
0	(6, 2, 2)	(1, 0, 0, 1, 0, 0, 0, 0, 2, 2)	25	$x_1 x_2^5 x_3^2 x_4 x_5^3 x_6^2 x_7^5 x_8^5 x_{10}$	1
0	(6, 3, 1)	(1, 0, 0, 1, 0, 0, 0, 0, 1, 2)	30	$x_1^2 x_2^3 x_3^4 x_4 x_5^5 x_6^5 x_7^4 x_8^4 x_9^2$	1
0	(6, 4, 0)	(1, 0, 0, 1, 0, 0, 0, 0, 1, 1)	28	$x_2^2 x_3^3 x_4 x_5^4 x_6^4 x_7^5 x_8^4 x_9^2 x_{10}^3$	2
0	(7, 2, 1)	(1, 0, 0, 1, 0, 0, 0, 0, 0, 2)	33	$x_3^3 x_4 x_5^6 x_6^5 x_7^6 x_8^6 x_9^6$	2
0	(7, 3, 0)	(1, 0, 0, 1, 0, 0, 0, 0, 0, 1)	35	$x_3^3 x_4 x_5^6 x_6^6 x_7^6 x_8^5 x_9^6 x_{10}^2$	2
0	(8, 1, 1)	(1, 0, 0, 0, 2, 0, 0, 0, 0, 0)	44	$x_2^3 x_3^4 x_4^2 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}^7$	$12r + 12$
0	(8, 2, 0)	(1, 0, 0, 0, 1, 0, 0, 0, 0, 0)	42	$x_2^3 x_3^4 x_4^4 x_5 x_6^5 x_7^5 x_8^7 x_9^7 x_{10}^6$	5
0	(9, 1, 0)	(0, 0, 0, 1, 0, 0, 0, 0, 0, 0)	66	$x_1^5 x_2^7 x_3^8 x_5^8 x_6^7 x_7^8 x_8^8 x_9^8 x_{10}^8$	$2^2 \cdot 5^2$
1	(3, 5, 2)	(0, 1, 1, 0, 1, 1, 1, 2, 0, 2)	27	$x_1 x_2^4 x_3^4 x_4^2 x_5^4 x_6^4 x_7^4 x_8 x_9^2 x_{10}$	$-3055r - 1872$
1	(2, 5, 3)	(0, 1, 1, 0, 1, 1, 1, 2, 2, 2)	28	$x_1 x_2^4 x_3^4 x_4 x_5^4 x_6^4 x_7^4 x_8^2 x_9^2 x_{10}^2$	$3993r + 2619$

\bar{a}	λ	\mathbf{a}''	deg	monomial/s	coefficient/s
1	(4, 5, 1)	(0, 1, 1, 0, 1, 1, 1, 0, 0, 2)	31	$x_1^2 x_2^4 x_3^4 x_4^3 x_5^4 x_6^4 x_7^4 x_8^3 x_9^3$	$3130r + 21$
1	(1, 5, 4)	(1, 2, 2, 1, 2, 2, 1, 1, 1, 0)	32	$x_1^4 x_2^3 x_3^3 x_4^4 x_5^3 x_6^3 x_7^4 x_8^4 x_9^4$	$-4043r - 1625$
1	(0, 5, 5)	(1, 2, 1, 2, 1, 1, 1, 2, 2, 2)	38	$x_1^3 x_2^3 x_3^4 x_4^4 x_5^4 x_6^4 x_7^4 x_8^4 x_9^4 x_{10}^4$	$7128r - 12852$
1	(2, 6, 2)	(0, 1, 0, 1, 1, 2, 2, 1, 1, 1)	30	$x_1 x_2^3 x_3 x_4^3 x_5^5 x_6 x_7 x_8^5 x_9^5 x_{10}^5$	$-424r - 595$
1	(3, 6, 1)	(0, 1, 0, 1, 1, 0, 2, 1, 1, 1)	36	$x_1^2 x_2^5 x_3^2 x_4^5 x_5^5 x_6^2 x_8^5 x_9^5 x_{10}^5$	$-566r + 1653$
1	(1, 6, 3)	(1, 0, 1, 1, 2, 2, 2, 1, 1, 1)	34	$x_1^4 x_3^4 x_4^5 x_5^2 x_6^2 x_7^2 x_8^5 x_9^5 x_{10}^5$	$61364r + 25951$
1	(4, 6, 0)	(0, 1, 1, 0, 1, 1, 1, 1, 0, 0)	34	$x_1^3 x_2^3 x_3^3 x_4^3 x_5^3 x_6^3 x_7^5 x_8^5 x_9^3 x_{10}^3$	$-1272r + 653$
1	(0, 6, 4)	(2, 1, 1, 2, 1, 1, 1, 1, 2, 2)	37	$x_1^3 x_2^3 x_3^3 x_4^4 x_5^5 x_6^5 x_7^5 x_8^5 x_9^3 x_{10}^3$	$-5321r - 5949$
1	(2, 7, 1)	(0, 1, 1, 0, 1, 1, 1, 1, 1, 2)	35	$x_1 x_2^3 x_3^3 x_4 x_5^3 x_6^6 x_7^6 x_8^6 x_9^6$	$-291r + 1$
1	(1, 7, 2)	(1, 0, 1, 2, 1, 1, 1, 1, 1, 2)	40	$x_1^4 x_3^4 x_4 x_5^6 x_6^6 x_7^6 x_8^6 x_9^6 x_{10}$	$11711r + 1287$
1	(3, 7, 0)	(0, 1, 1, 0, 1, 1, 1, 1, 1, 0)	36	$x_1^2 x_2^3 x_3^3 x_4^2 x_5^3 x_6^3 x_8^6 x_9^6 x_{10}^2$	$5839r + 1451$
1	(0, 7, 3)	(2, 1, 1, 2, 1, 1, 1, 1, 1, 2)	40	$x_1^2 x_2^3 x_3^3 x_4^2 x_5^6 x_6^6 x_7^6 x_8^6 x_9^6 x_{10}^2$	$-1365r - 1577$
1	(1, 8, 1)	(1, 0, 1, 1, 1, 1, 1, 1, 1, 2)	43	$x_1^4 x_3^4 x_4^4 x_5^6 x_6^6 x_7^7 x_8^7 x_9^7$	$239r - 1811$
1	(2, 8, 0)	(1, 0, 1, 1, 1, 1, 1, 1, 1, 0)	43	$x_1^4 x_2 x_3^4 x_4^4 x_5^4 x_6^4 x_7^7 x_8^7 x_9^7 x_{10}$	$-10919r - 9004$
1	(0, 8, 2)	(1, 2, 1, 1, 1, 1, 1, 1, 1, 2)	47	$x_1^4 x_2 x_3^4 x_4^4 x_5^5 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}$	$5278r + 39848$
1	(1, 9, 0)	(1, 0, 1, 1, 1, 1, 1, 1, 1, 1)	51	$x_1^4 x_3^4 x_4^4 x_5^5 x_6^5 x_7^8 x_8^8 x_9^8 x_{10}^8$	$-12212r - 8422$
1	(0, 9, 1)	(1, 2, 1, 1, 1, 1, 1, 1, 1, 1)	54	$x_1^4 x_3^4 x_4^4 x_5^6 x_6^6 x_7^8 x_8^8 x_9^8 x_{10}^8$	$-82561r - 26061$
1	(0, 10, 0)	(1, 1, 1, 1, 1, 1, 1, 1, 1, 1)	65	$x_1^2 x_2^4 x_3^4 x_4^4 x_5^6 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^9$	$7028r + 8686$
2	(3, 2, 5)	(0, 2, 2, 0, 2, 2, 2, 1, 0, 1)	27	$x_1 x_2^4 x_3^4 x_4^2 x_5^4 x_6^4 x_7^4 x_8 x_9^2 x_{10}$	$3055r + 1183$
2	(2, 3, 5)	(0, 2, 2, 0, 2, 2, 2, 1, 1, 1)	28	$x_1 x_2^4 x_3^4 x_4 x_5^4 x_6^4 x_7^4 x_8^2 x_9^2 x_{10}^2$	$-3993r - 1374$
2	(4, 1, 5)	(0, 2, 2, 0, 2, 2, 2, 0, 0, 1)	31	$x_1^2 x_2^4 x_3^4 x_4^3 x_5^4 x_6^4 x_7^4 x_8^3 x_9^3$	$-3130r - 3109$
2	(1, 4, 5)	(2, 1, 1, 2, 1, 1, 2, 2, 2, 0)	32	$x_1^4 x_2^3 x_3^3 x_4^4 x_5^3 x_6^3 x_7^4 x_8^4 x_9^4$	$4043r + 2418$
2	(5, 0, 5)	(0, 2, 0, 2, 0, 0, 0, 2, 2, 2)	33	$x_1^2 x_2^2 x_3^2 x_4^3 x_5^4 x_6^4 x_7^4 x_8^4 x_9^4 x_{10}^4$	$-124r + 170$
2	(2, 2, 6)	(0, 2, 0, 2, 2, 1, 1, 2, 2, 2)	30	$x_1 x_2^3 x_3 x_4^3 x_5^5 x_6 x_7 x_8^5 x_9^5 x_{10}^5$	$424r - 171$
2	(3, 1, 6)	(0, 2, 0, 2, 2, 0, 1, 2, 2, 2)	36	$x_1^2 x_2^5 x_3^2 x_4^5 x_5^5 x_6^2 x_8^5 x_9^5 x_{10}^5$	$566r + 2219$
2	(1, 3, 6)	(2, 0, 2, 2, 1, 1, 1, 2, 2, 2)	34	$x_1^4 x_3^4 x_4^5 x_5^2 x_6^2 x_7^2 x_8^5 x_9^5 x_{10}^5$	$-61364r - 35413$

\bar{a}	λ	\mathbf{a}''	deg	monomial/s	coefficient/s
2	(4, 0, 6)	(0, 2, 2, 0, 2, 2, 2, 2, 0, 0)	34	$x_1^3 x_2^3 x_3^3 x_4^3 x_5^3 x_6^3 x_7^5 x_8^5 x_9^3 x_{10}^3$	$1272r + 1925$
2	(0, 4, 6)	(1, 2, 2, 1, 2, 2, 2, 2, 1, 1)	37	$x_1^3 x_2^3 x_3^3 x_4^3 x_5^4 x_6^5 x_7^5 x_8^5 x_9^3 x_{10}^3$	$5321r - 628$
2	(2, 1, 7)	(0, 2, 2, 0, 2, 2, 2, 2, 2, 1)	35	$x_1 x_2^3 x_3^3 x_4 x_5^3 x_6^6 x_7^6 x_8^6 x_9^6$	$291r + 29$
2	(1, 2, 7)	(2, 0, 2, 1, 2, 2, 2, 2, 2, 1)	40	$x_1^4 x_2^4 x_3 x_4 x_5^6 x_6^6 x_7^6 x_8^6 x_9^6 x_{10}$	$-11711r - 10424$
2	(3, 0, 7)	(0, 2, 2, 0, 2, 2, 2, 2, 2, 0)	36	$x_1^2 x_2^3 x_3^3 x_4^2 x_5^3 x_6^3 x_7^6 x_8^6 x_9^6 x_{10}^2$	$-5839r - 4388$
2	(0, 3, 7)	(1, 2, 2, 1, 2, 2, 2, 2, 2, 1)	40	$x_1^2 x_2^3 x_3^3 x_4^2 x_5^4 x_6^6 x_7^6 x_8^6 x_9^6 x_{10}^2$	$1365r - 212$
2	(1, 1, 8)	(2, 0, 2, 2, 2, 2, 2, 2, 2, 1)	43	$x_1^4 x_2^4 x_3^4 x_4^4 x_5^6 x_6^7 x_7^7 x_8^7 x_9^7$	$-239r - 2050$
2	(2, 0, 8)	(2, 0, 2, 2, 2, 2, 2, 2, 2, 0)	43	$x_1^4 x_2 x_3^4 x_4^4 x_5^4 x_6^4 x_7^7 x_8^7 x_9^7 x_{10}$	$10919r + 1915$
2	(0, 2, 8)	(2, 1, 2, 2, 2, 2, 2, 2, 2, 1)	47	$x_1^4 x_2 x_3^4 x_4^4 x_5^5 x_6^7 x_7^7 x_8^7 x_9^7 x_{10}$	$-5278r + 34570$
2	(1, 0, 9)	(2, 0, 2, 2, 2, 2, 2, 2, 2, 2)	51	$x_1^4 x_2^4 x_3^4 x_4^5 x_5^5 x_6^5 x_7^8 x_8^8 x_9^8 x_{10}^8$	$12212r + 3790$
2	(0, 1, 9)	(2, 1, 2, 2, 2, 2, 2, 2, 2, 2)	54	$x_1^4 x_2^4 x_3^4 x_4^6 x_5^6 x_6^6 x_7^8 x_8^8 x_9^8 x_{10}^8$	$82561r + 56500$
2	(0, 0, 10)	(2, 2, 2, 2, 2, 2, 2, 2, 2, 2)	65	$x_1^2 x_2^4 x_3^4 x_4^4 x_5^6 x_6^9 x_7^9 x_8^9 x_9^9 x_{10}^9$	$-7028r + 1658$

References

- [1] N. Alon, Combinatorial Nullstellensatz, *Combin. Probab. Comput.* **8** (1999) 7–29.
- [2] B. Alspach and G. Liversidge, On strongly sequenceable abelian groups, *Art Discrete Appl. Math.* **3** (2020) 19pp.
- [3] B. A. Anderson and E. C. Ihrig, Every finite solvable group with a unique element of order two, except the quaternion group, has a symmetric sequencing, *J. Combin. Des.* **1** (1993) 3–14.
- [4] S. Costa and S. Della Fiore, Weak sequenceability in cyclic groups, *J. Combin. Des.* **30** (2022) 735–751.
- [5] S. Costa, S. Della Fiore, M. A. Ollis and S. Z. Rovner-Frydman, On Sequences in Cyclic Groups with Distinct Partial Sums, *Electron. J. Combin.* **29** (2022) #P3.33.
- [6] S. Costa, F. Morini, A. Pasotti and M. A. Pellegrini, A problem on partial sums in abelian groups, *Discrete Math.* **341** (2018) 705–712.
- [7] S. Costa and M. A. Pellegrini, Some new results about a conjecture by Brian Alspach, *Archiv der Mathematik*, **115** (2020) 479–488.
- [8] B. Gordon, Sequences in groups with distinct partial products, *Pacific J. Math.* **11** (1961) 1309–1313.

- [9] R. L. Graham, On sums of integers taken from a fixed sequence, *Proceedings, Washington State University Conference on Number Theory* (1971) 22–40.
- [10] J. Hicks, M. A. Ollis, and J. R. Schmitt, Distinct partial sums in cyclic groups: polynomial method and constructive approaches, *J. Combin. Des.* **27** (2019) 369–385.
- [11] J. Isbell, Sequencing certain dihedral groups, *Discrete Math.* **85** (1990), 323–328.
- [12] A. D. Keedwell, On the sequenceability of non-abelian groups of order pq , *Discrete Math.* **37** (1981) 203–216.
- [13] P. Li, Sequencing the dihedral groups D_{4k} , *Discrete Math.* **175** (1997) 271–276.
- [14] M. Michałek, A short proof of Combinatorial Nullstellensatz, *Amer. Math. Monthly* **117** (2010) (821–823).
- [15] M. A. Ollis, Terraces for small groups, *J. Combin. Math. Comput.* **108** (2019) 231–244.
- [16] M. A. Ollis, Sequences in dihedral groups with distinct partial products, *Australas. J. Combin.* **78** (2020) 35–60.
- [17] M. A. Ollis and C. R. Tripp, The spectrum of group-based complete latin squares, *Electron. J. Combin.* **26** #3.15 (2019) 13pp.
- [18] A. Pasotti and J. H. Dinitz, A survey of Heffter arrays, [arXiv:2209.13879](https://arxiv.org/abs/2209.13879).
- [19] Sage Developers, Sage Mathematics Software (Version 9.5.0), The SAGE Group, 2022, <http://www.sagemath.org>.