

Hopf algebraic structures on mixed graphs

Loïc Foissy

Univ. Littoral Côte d'Opale, UR 2597 LMPA, Laboratoire de Mathématiques Pures et Appliquées
Joseph Liouville F-62100 Calais, France.
Email: foissy@univ-littoral.fr

Abstract

We introduce two coproducts on mixed graphs (that is to say graphs with both oriented and unoriented edges), the first one by separation of the vertices into two parts, and the second one given by contraction and extractions of subgraphs. We show that, with the disjoint union product, this gives a double bialgebra, that is to say that the first coproduct makes it a Hopf algebra in the category of right comodules over the second coproduct.

This structures implies the existence of a unique polynomial invariant on mixed graphs compatible with the product and both coproducts: we prove that it is the (strong) chromatic polynomial of Beck, Bogart and Pham. Using the action of the monoid of characters, we relate it to the weak chromatic polynomial, as well to Ehrhart polynomials and to a polynomial invariants related to linear extensions. As applications, we give an algebraic proof of the link between the values of the strong chromatic polynomial at negative values and acyclic orientations (a result due to Beck, Blado, Crawford, Jean-Louis and Young) and obtain a combinatorial description of the antipode of the Hopf algebra of mixed graphs.

Keywords. double bialgebra; mixed graph; chromatic polynomial; Ehrhart polynomial

AMS classification. 16T05 16T30 05C15 05C25

Contents

1	Species, twisted bialgebras and contractions	5
1.1	Species and twisted bialgebras	5
1.2	Contraction-extraction coproducts	8
2	The species of mixed graphs	9
2.1	Mixed graphs	9
2.2	The twisted bialgebra of mixed graphs	11
2.3	Contraction-extraction on mixed graphs	12
2.4	Cofreeness of the coalgebra $\mathcal{F}[\mathbf{G}]$	16
3	Sub-objects and quotients of mixed graphs	17
3.1	Simple and oriented graphs	17
3.2	Acyclic mixed graphs and finite topologies	17
3.3	Totally acyclic graphs	21
4	Applications	22
4.1	Three polynomial invariants	22
4.2	Invertible characters	24

5	From mixed graphs to acyclic oriented graphs	29
5.1	A double bialgebra epimorphism	29
5.2	Erhart polynomials for mixed graphs	31
6	Applications to characters on mixed graphs	34
6.1	Weak chromatic polynomial	34
6.2	The character ν_W	36
7	Appendix: number of mixed graphs up to isomorphism	37

Introduction

Mixed graphs are graphs with both unoriented (which we will simply called edges) and oriented edges (which we will called arcs). They are used for example to study scheduling problems with disjunctive (represented by unoriented edges) and precedence (represented by oriented edges) constraints [18, 18, 12]. In this context, a notion of admissible coloring, similar to the notion used for classical graphs, give a solution of the scheduling problem represented by the mixed graph. These admissible colorings can be counted according to the number of colors, which gives a chromatic polynomial [5, 4]. The aim of this text is the study of this chromatic polynomial for mixed graphs in the context of double bialgebras, as this has been done for graphs in [9] and for posets and finite topologies in [7]. A double bialgebra is a family (A, m, Δ, δ) such that:

- (A, m, δ) is a bialgebra.
- (A, m, Δ) is a bialgebra in the category of right comodules over (A, m, δ) , with the coaction given by δ itself.

In particular, δ and Δ satisfies the following compatibility:

$$(\Delta \otimes \text{Id}) \circ \delta = (\text{Id} \otimes \text{Id} \otimes m) \circ (\text{Id} \otimes c \otimes \text{Id}) \circ (\delta \otimes \delta) \circ \Delta,$$

where $c : A \otimes A \longrightarrow A \otimes A$ is the usual flip. The counit ε_Δ of Δ satisfies the following compatibility with δ : for any $a \in A$,

$$(\varepsilon_\Delta \otimes \text{Id}) \circ \delta(x) = \varepsilon_\Delta(x)1_A.$$

A simple example of such an object is $\mathbb{K}[X]$, with its usual algebra structures and its multiplicative coproducts defined by

$$\Delta(X) = X \otimes 1 + 1 \otimes X, \quad \delta(X) = X \otimes X.$$

We proved in [7, 9] the following results: if (A, m, Δ, δ) is a double bialgebra, under a condition of connectedness of (A, m, Δ) ,

- There exists a unique double bialgebra morphism $P_0 : (A, m, \Delta, \delta) \longrightarrow (\mathbb{K}[X], m, \Delta, \delta)$, which can be explicitly described with iterations of the coproduct Δ of A and with the counit ε_δ of its coproduct δ .
- Let us denote by $\text{Char}(A)$ the set of characters of A . It inherits a convolution product \star , dual to δ . We denote by $\text{End}_B(A, \mathbb{K}[X])$ the set of bialgebra morphisms from (A, m, Δ) to $(\mathbb{K}[X], m, \Delta)$. Then [10, Proposition 2.5], the following defines an action of the monoid $(\text{Char}(A), \star)$ on $\text{End}_B(A, \mathbb{K}[X])$:

$$\leftarrow : \begin{cases} \text{End}_B(A, \mathbb{K}[X]) \times \text{Char}(A) & \longrightarrow & \text{End}_B(A, \mathbb{K}[X]) \\ (\phi, \lambda) & \longmapsto & \phi \leftarrow \lambda = (\phi \otimes \lambda) \circ \delta. \end{cases}$$

Moreover, the two following maps are bijection, inverse one from the other:

$$\left\{ \begin{array}{l} \text{End}_B(A, \mathbb{K}[X]) \longrightarrow \text{Char}(A) \\ \phi \longmapsto \epsilon_\delta \circ \phi, \end{array} \right. \quad \theta : \left\{ \begin{array}{l} \text{Char}(A) \longrightarrow \text{End}_B(A, \mathbb{K}[X]) \\ \lambda \longmapsto P_0 \leftarrow \lambda, \end{array} \right. \quad (1)$$

where here ϵ_δ is the counit of $(\mathbb{K}[X], m, \delta)$, which send any $P \in \mathbb{K}[X]$ onto $P(1)$.

- When (A, m, Δ) is a graded and connected bialgebra, then under a technical condition on δ , the set of homogeneous bialgebra morphisms from (A, m, Δ) to $(\mathbb{K}[X], m, \Delta)$ is in bijection with the dual of the homogeneous component A_1 of A of degree 1.
- If (A, m, Δ) is a Hopf algebra, then its antipode is given by

$$S = (\epsilon_\delta^{*-1} \otimes \text{Id}) \circ \delta,$$

where ϵ_δ^{*-1} is the inverse of the counit ϵ_δ of δ for the convolution product $*$ associated to Δ . Moreover, for any $a \in A$,

$$\epsilon_\delta^{*-1}(a) = P_0(a)(-1).$$

We apply here these results on mixed graphs. In the second section, we define a structure of double bialgebras on mixed graphs. We work in the frame of species and use the formalism built in [11] of contraction-extraction coproduct, which is shortly described in the first section. We first give in Proposition 2.4 to the species of mixed graphs \mathbf{G} a bialgebra structure in the category of species (what is commonly called a twisted bialgebra structure), and then a contraction-extraction coproduct in Proposition 2.6. Consequently, applying the bosonic Fock functor [1], we obtain a double bialgebra of mixed graphs $\mathcal{F}[\mathbf{G}]$, and more generally, for any commutative and cocommutative bialgebra V , a double bialgebra $\mathcal{F}_V[\mathbf{G}]$ of mixed graphs which vertices are decorated by elements of V . Using Loday and Ronco's rigidity theorem, we prove that $(\mathcal{F}_V[\mathbf{G}], \Delta)$ is a cofree coalgebra (Corollary 2.9).

We study certain sub-objects and quotients of \mathbf{G} in the third section. Obviously, simple (i.e., unoriented) graphs and oriented graphs define twisted double subbialgebras of \mathbf{G} , denoted respectively by \mathbf{G}_s and \mathbf{G}_o (Proposition 2.1). Applying the bosonic Fock functor to \mathbf{G}_s , we obtain again the double bialgebra of graphs of [9, 11]. We also consider the subspecies of acyclic mixed graphs \mathbf{G}_{aco} , which turns out to be stable under the product and the first coproduct Δ , but not on the second one. However, quotienting by non acyclic mixed graphs, \mathbf{G}_{ac} can be seen as a twisted double bialgebra, quotient of \mathbf{G} (Proposition 2.5). It contains a twisted double subbialgebra of oriented acyclic mixed graphs \mathbf{G}_{aco} , which has itself for quotient the twisted double bialgebra of topologies of [7] (Proposition 2.6).

The fourth part is devoted to polynomial invariants of mixed graphs, that is to say bialgebra morphisms from the bialgebra of mixed graphs $(\mathcal{F}[\mathbf{G}], m, \Delta)$ to $(\mathbb{K}[X], m, \Delta)$. We first describe the unique polynomial invariant compatible with the second coalgebraic structure of mixed graphs: it turns out to be the strong chromatic polynomial P_{chr_S} of [5], see Proposition 4.2. In other words, for any mixed graph G , for any $n \geq 1$, $P_{chr_S}(G)(n)$ is the number of n -valid colorings of G , that is to say maps c from the set of vertices $V(G)$ of G to $\{1, \dots, n\}$ such that, for any pair of vertices x, y of G ,

- If x and y are related by an edge of G , then $c(x) \neq c(y)$.
- If x and y are related by an arc of G , then $c(x) < c(y)$.

A notion of weak valid coloring is also defined in [5]. A weak n -valid coloring is a map c from the set of vertices $V(G)$ of G to $\{1, \dots, n\}$ such that, for any pair of vertices x, y of G ,

- If x and y are related by an edge of G , then $c(x) \neq c(y)$.
- If x and y are related by an arc of G , then $c(x) \leq c(y)$.

The polynomial counting the number of weak n -valid colorings of G is denoted by $P_{chr_W}(G)$. Using the action of the monoid of characters described earlier and the character of $\mathcal{F}[\mathbf{G}]$ defined by

$$\lambda_W(G) = \begin{cases} 1 & \text{if } G \text{ is an oriented graph,} \\ 0 & \text{otherwise,} \end{cases}$$

we obtain that $P_{chr_W} = P_{chr_S} \leftarrow \lambda_W$, which implies that P_{chr_W} is a bialgebra morphism from $(\mathcal{F}[\mathbf{G}], m, \Delta)$ to $(\mathbb{K}[X], m, \Delta)$, see Corollary 4.4. Finally, using the correspondence between homogeneous polynomial invariants and elements of $\mathcal{F}[\mathbf{G}]_1$, we construct a homogeneous bialgebra morphism P_0 from $(\mathcal{F}[\mathbf{G}], m, \Delta)$ to $(\mathbb{K}[X], m, \Delta)$, related to the number of linear extensions (Corollary 4.5) and to a character λ_0 . When V is a commutative and cocommutative bialgebra, similar results are given for $\mathcal{F}_V[\mathbf{G}]$, where $\mathbb{K}[X]$ is replaced by a double bialgebra of quasi-shuffles, see Theorem ?? and Corollary ?. After the determination of invertible characters of $\mathcal{F}[\mathbf{G}]$ for the convolution product \star dual to δ , we prove that both characters λ_0 and λ_W are invertible, which allows to express P_{chr_S} in terms of P_{chr_W} or P_0 , with the help of certain characters ν_W and μ_W (Proposition 4.11). This allows to give a formula for the leading monomial of $P_{chr_W}(G)$ and $P_{chr_S}(G)$ in Corollary 4.12, with coefficients (in the case of $P_{chr_S}(G)$) related to Murua's coefficients [16].

In the fifth section, we give an algebraic proof of the result [5], which gives a combinatorial interpretation of $P_{chr_S}(G)(-1)$ in terms of acyclic orientations. We firstly introduce a surjective double bialgebra morphism Θ from $\mathcal{F}[\mathbf{G}]$ to the double bialgebra of acyclic oriented mixed graphs $\mathcal{F}[\mathbf{G}_{aco}]$ in Theorem 5.2. The unicity of the double bialgebra morphism to $\mathbb{K}[X]$ immediately implies for example that the strong chromatic polynomial of a graph G is the sum of the strong chromatic polynomial of all its acyclic orientations (Corollary 5.3). Introducing two one-parameter families of characters of $\mathcal{F}[\mathbf{G}]$ in Proposition 5.4, we introduce two new polynomial invariants, which turn out to be Ehrhart polynomials and satisfy a duality principle on acyclic mixed graphs (Corollary 5.7). Mixing this duality principle with the morphism Θ , we obtain a new proof that for any mixed graph, $P_{chr_S}(G)(-1)$ is the number of acyclic orientations of G , up to a sign (Corollary 5.9). This allows to give a formula for the antipode of $(\mathcal{F}[\mathbf{G}], m, \Delta)$ involving the number of acyclic orientations of G , see Corollary 5.10.

The last section is devoted to combinatorial interpretations of special characters of mixed graphs. We give an algebraic proof of a result of [5] in an algebraic way (Proposition 6.1 and Corollary 6.2) about values of the weak chromatic polynomial on negative integers (for totally mixed graphs only), and we give a combinatorial interpretation of $\nu_W(G)$ when G is a simple graph or an oriented graph (Proposition 6.3), where ν_W is the inverse of λ_W for the convolution product \star .

The paper ends with an appendix giving formulas for the number of N -decorated mixed, oriented or simple graphs with a fixed number of vertices. These numbers are the dimensions of the homogeneous components of $\mathcal{F}_V[\mathbf{G}]$, when V is finite-dimensional.

Acknowledgments. The author acknowledges support from the grant ANR-20-CE40-0007 *Combinatoire Algébrique, Résurgence, Probabilités Libres et Opérades*.

Notations 0.1. 1. We denote by \mathbb{K} a commutative field of characteristic zero. Any vector space in this field will be taken over \mathbb{K} .

2. For any $n \in \mathbb{N}$, we denote by $[n]$ the set $\{1, \dots, n\}$. In particular, $[0] = \emptyset$.

3. If (C, Δ) is a (coassociative but not necessarily counitary) coalgebra, we denote by $\Delta^{(n)}$

the n -th iterated coproduct of C : $\Delta^{(1)} = \Delta$ and if $n \geq 2$,

$$\Delta^{(n)} = \left(\Delta \otimes \text{Id}^{\otimes(n-1)} \right) \circ \Delta^{(n-1)} : C \longrightarrow C^{\otimes(n+1)}.$$

4. If (B, m, Δ) is a bialgebra of unit 1_B and of counit ε_B , let us denote by $B_+ = \text{Ker}(\varepsilon_B)$ its augmentation ideal. We define a coproduct on B_+ by

$$\forall x \in B_+, \quad \tilde{\Delta}(x) = \Delta(x) - x \otimes 1_B - 1_B \otimes x.$$

Then $(B_+, \tilde{\Delta})$ is a coassociative (not necessarily counitary) coalgebra.

1 Species, twisted bialgebras and contractions

1.1 Species and twisted bialgebras

Recall that a species [13, 14] is a functor from the category of finite sets with bijections, to the category of vector spaces. We shall use the following notations: if \mathbf{P} is a species,

- For any finite set X , the vector space associated to X by \mathbf{P} is denoted by $\mathbf{P}[X]$. We will shortly write $\mathbf{P}[n]$ instead of $\mathbf{P}[[n]]$ for any $n \in \mathbb{N}$.
- For any bijection $\sigma : X \longrightarrow Y$ between two finite sets, the linear map associated to σ by \mathbf{P} is denoted by $\mathbf{P}[\sigma] : \mathbf{P}[X] \longrightarrow \mathbf{P}[Y]$.

With these notations, if $\sigma : X \longrightarrow Y$ and $\tau : Y \longrightarrow Z$ are two bijections between finite sets, then $\mathbf{P}[\tau \circ \sigma] = \mathbf{P}[\tau] \circ \mathbf{P}[\sigma]$, and for any finite set X , $\mathbf{P}[\text{Id}_X] = \text{Id}_{\mathbf{P}[X]}$.

If \mathbf{P} and \mathbf{Q} are two species, a morphism $f : \mathbf{P} \longrightarrow \mathbf{Q}$ is a natural transformation from \mathbf{P} to \mathbf{Q} . In other terms, for any finite X , $f_X : \mathbf{P}[X] \longrightarrow \mathbf{Q}[X]$ is a linear map such that for any bijection $\sigma : X \longrightarrow Y$ between two finite sets, the following diagram commutes:

$$\begin{array}{ccc} \mathbf{P}[X] & \xrightarrow{\mathbf{P}[\sigma]} & \mathbf{P}[Y] \\ f_X \downarrow & & \downarrow f_Y \\ \mathbf{Q}[X] & \xrightarrow{\mathbf{Q}[\sigma]} & \mathbf{Q}[Y] \end{array}$$

The Cauchy tensor product of species is denoted by \otimes : if \mathbf{P} and \mathbf{Q} are two species, for any finite set X ,

$$\mathbf{P} \otimes \mathbf{Q}[X] = \bigoplus_{X=Y \sqcup Z} \mathbf{P}[Y] \otimes \mathbf{Q}[Z].$$

If $\sigma : X \longrightarrow X'$ is a bijection between two finite sets, then

$$\mathbf{P} \otimes \mathbf{Q}[\sigma] = \bigoplus_{X=Y \sqcup Z} \mathbf{P}[\sigma|_Y] \otimes \mathbf{Q}[\sigma|_Z].$$

With this tensor product, the category of species is symmetric monoidal, with the flip defined from $\mathbf{P} \otimes \mathbf{Q}$ to $\mathbf{Q} \otimes \mathbf{P}$ by

$$c_{\mathbf{P}[X], \mathbf{P}[Y]} = (c_{\mathbf{P} \otimes \mathbf{Q}})_{\mathbf{P}[X] \otimes \mathbf{Q}[Y]} = \begin{cases} \mathbf{P}[X] \otimes \mathbf{Q}[Y] & \longrightarrow & \mathbf{Q}[Y] \otimes \mathbf{P}[X] \\ x \otimes y & \longmapsto & y \otimes x. \end{cases}$$

A twisted algebra (resp. coalgebra, bialgebra) is an algebra (resp. coalgebra, bialgebra) in this symmetric monoidal category. Let us now precise what this means. A twisted algebra \mathbf{P} comes with maps $m_{X,Y} : \mathbf{P}[X] \otimes \mathbf{P}[Y] \longrightarrow \mathbf{P}[X \sqcup Y]$ for any finite sets X and Y , such that:

- (m is a species morphism). For any bijections $\sigma : X \rightarrow X'$ and $\tau : Y \rightarrow Y'$, where (X, X') and (Y, Y') are pairs of disjoint sets, the following diagram commutes:

$$\begin{array}{ccc} \mathbf{P}[X] \otimes \mathbf{P}[Y] & \xrightarrow{m_{X,Y}} & \mathbf{P}[X \sqcup Y] \\ \mathbf{P}[\sigma] \otimes \mathbf{P}[\tau] \downarrow & & \downarrow \mathbf{P}[\sigma \sqcup \tau] \\ \mathbf{P}[X'] \otimes \mathbf{P}[Y'] & \xrightarrow{m_{X',Y'}} & \mathbf{P}[X' \sqcup Y'] \end{array}$$

where

$$\sigma \sqcup \tau : \begin{cases} X \sqcup Y & \rightarrow & X' \sqcup Y' \\ x \in X & \rightarrow & \sigma(x), \\ y \in Y & \rightarrow & \tau(y). \end{cases}$$

- (Associativity of m). For any disjoint finite sets X, Y, Z , the following diagram commutes:

$$\begin{array}{ccc} \mathbf{P}[X] \otimes \mathbf{P}[Y] \otimes \mathbf{P}[Z] & \xrightarrow{m_{X,Y} \otimes \text{Id}_{\mathbf{P}[Z]}} & \mathbf{P}[X \sqcup Y] \otimes \mathbf{P}[Z] \\ \text{Id}_{\mathbf{P}[X]} \otimes m_{Y,Z} \downarrow & & \downarrow m_{X \sqcup Y, Z} \\ \mathbf{P}[X] \otimes \mathbf{P}[Y \sqcup Z] & \xrightarrow{m_{X, Y \sqcup Z}} & \mathbf{P}[X \sqcup Y \sqcup Z] \end{array}$$

- There exists an element $1_{\mathbf{P}} \in \mathbf{P}[\emptyset]$ such that for any finite set X , for any $x \in \mathbf{P}[X]$,

$$m_{\emptyset, X}(1_{\mathbf{P}} \otimes x) = m_{X, \emptyset}(x \otimes 1_{\mathbf{P}}) = x.$$

- (Commutativity of m). We shall say that \mathbf{P} is commutative if, for any finite disjoint sets X, Y , in $\mathbf{P}[X \sqcup Y]$,

$$\forall x \in \mathbf{P}[X], \forall y \in \mathbf{P}[Y], \quad m_{X,Y}(x \otimes y) = m_{Y,X}(y \otimes x).$$

A twisted coalgebra \mathbf{P} comes with maps $\Delta_{X,Y} : \mathbf{P}[X \sqcup Y] \rightarrow \mathbf{P}[X] \otimes \mathbf{P}[Y]$ for any disjoint finite sets X and Y , such that:

- (Δ is a species morphism). For any bijections $\sigma : X \rightarrow X'$ and $\tau : Y \rightarrow Y'$, where (X, Y) and (X', Y') are pairs of disjoint finite sets, the following diagram commutes:

$$\begin{array}{ccc} \mathbf{P}[X \sqcup Y] & \xrightarrow{\Delta_{X,Y}} & \mathbf{P}[X] \otimes \mathbf{P}[Y] \\ \mathbf{P}[\sigma \sqcup \tau] \downarrow & & \downarrow \mathbf{P}[\sigma] \otimes \mathbf{P}[\tau] \\ \mathbf{P}[X' \sqcup Y'] & \xrightarrow{\Delta_{X',Y'}} & \mathbf{P}[X'] \otimes \mathbf{P}[Y'] \end{array}$$

- (Coassociativity of Δ). For any disjoint finite sets X, Y, Z , the following diagram commutes:

$$\begin{array}{ccc} \mathbf{P}[X \sqcup Y \sqcup Z] & \xrightarrow{\Delta_{X \sqcup Y, Z}} & \mathbf{P}[X \sqcup Y] \otimes \mathbf{P}[Z] \\ \Delta_{X \sqcup Y, Z} \downarrow & & \downarrow \Delta_{X,Y} \otimes \text{Id}_{\mathbf{P}[Z]} \\ \mathbf{P}[X] \otimes \mathbf{P}[Y \sqcup Z] & \xrightarrow{\text{Id}_{\mathbf{P}[X]} \otimes \Delta_{Y,Z}} & \mathbf{P}[X] \otimes \mathbf{P}[Y] \otimes \mathbf{P}[Z] \end{array}$$

- (Counit). There exists a linear map $\varepsilon_{\Delta} : \mathbf{P}[\emptyset] \rightarrow \mathbb{K}$ such that for any finite set X , the following diagram commutes:

$$\begin{array}{ccccc} \mathbf{P}[\emptyset] \otimes \mathbf{P}[X] & \xleftarrow{\Delta_{\emptyset, X}} & \mathbf{P}[X] & \xrightarrow{\Delta_{X, \emptyset}} & \mathbf{P}[X] \otimes \mathbf{P}[\emptyset] \\ & \searrow \varepsilon_{\Delta} \otimes \text{Id}_{\mathbf{P}[X]} & \downarrow \text{Id}_{\mathbf{P}[X]} & \swarrow \text{Id}_{\mathbf{P}[X]} \otimes \varepsilon_{\Delta} & \\ & & \mathbf{P}[X] & & \end{array}$$

- We shall say that \mathbf{P} is cocommutative if for any disjoint finite sets X, Y ,

$$\Delta_{Y,X} = c_{\mathbf{P}[X], \mathbf{P}[Y]} \circ \Delta_{X,Y}.$$

A twisted bialgebra is both a twisted coalgebra and a twisted bialgebra, such that the counit and the coproduct are algebra morphisms. In other words, if X is a finite set and $X = I \sqcup J = I' \sqcup J'$, the following diagram commutes:

$$\begin{array}{ccc} \mathbf{P}[I'] \otimes \mathbf{P}[J'] & \xrightarrow{m_{I',J'}} & \mathbf{P}[I' \sqcup J'] = \mathbf{P}[I \sqcup J] \\ \Delta_{I' \cap I, I' \cap J} \otimes \Delta_{J' \cap I, J' \cap J} \downarrow & & \downarrow \Delta_{I,J} \\ \mathbf{P}[I' \cap I] \otimes \mathbf{P}[I' \cap J] \otimes \mathbf{P}[J' \cap I] \otimes \mathbf{P}[J' \cap J] & & \\ \text{Id}_{\mathbf{P}[I' \cap I]} \otimes c_{\mathbf{P}[I' \cap J], \mathbf{P}[J' \cap I]} \otimes \text{Id}_{\mathbf{P}[J' \cap J]} \downarrow & & \\ \mathbf{P}[I' \cap I] \otimes \mathbf{P}[J' \cap I] \otimes \mathbf{P}[I' \cap J] \otimes \mathbf{P}[J' \cap J] & \xrightarrow{m_{I' \cap I, J' \cap I} \otimes m_{I' \cap J, J' \cap J}} & \mathbf{P}[I] \otimes \mathbf{P}[J] \end{array}$$

or equivalently

$$\begin{aligned} \Delta_{I,J} \circ m_{I',J'} &= (m_{I' \cap I, J' \cap I} \otimes m_{I' \cap J, J' \cap J}) \circ (\text{Id}_{\mathbf{P}[I \cap I]} \otimes c_{\mathbf{P}[I' \cap J], \mathbf{P}[J' \cap I]} \otimes \text{Id}_{\mathbf{P}[J' \cap J]}) \\ &\quad \circ (\Delta_{I' \cap I, I' \cap J} \otimes \Delta_{J' \cap I, J' \cap J}). \end{aligned}$$

The compatibility between the coproduct and the unit is written as $\Delta_{\mathbf{P}}(1_{\mathbf{P}}) = 1_{\mathbf{P}} \otimes 1_{\mathbf{P}}$ and the compatibility between the counit and the product is equivalent to

$$\forall x, y \in \mathbf{P}[\emptyset], \quad \varepsilon_{\Delta}(xy) = \varepsilon_{\Delta}(x)\varepsilon_{\Delta}(y).$$

Let V be a vector space. The V -colored Fock functor \mathcal{F}_V , defined in [11, Definition 3.2], sends any species \mathbf{P} to

$$\begin{aligned} \mathcal{F}_V[\mathbf{P}] &= \bigoplus_{n=0}^{\infty} \text{coInv}(V^{\otimes n} \otimes \mathbf{P}[n]) \\ &= \bigoplus_{n=0}^{\infty} \frac{V^{\otimes n} \otimes \mathbf{P}[n]}{\text{Vect}(v_1 \dots v_n \otimes \mathbf{P}[\sigma](p) - v_{\sigma(1)} \dots v_{\sigma(n)} \otimes p \mid \sigma \in \mathfrak{S}_n, p \in \mathbf{P}[n], v_1, \dots, v_n \in V)} \\ &= V^{\otimes n} \otimes_{\mathfrak{S}_n} \mathbf{P}[n]. \end{aligned}$$

If $f : \mathbf{P} \rightarrow \mathbf{Q}$ is a species morphism, then

$$\mathcal{F}_V(f) : \begin{cases} \mathcal{F}_V[\mathbf{P}] & \mapsto \mathcal{F}_V[\mathbf{Q}] \\ v_1 \dots v_n \otimes p & \mapsto v_1 \dots v_n \otimes f(p). \end{cases}$$

Note that \mathcal{F}_V is a functor of symmetric monoidal categories, that is to say it sends the Cauchy tensor product of two species to the tensor products of the associated vector spaces, and that it is compatible with the flip. Consequently, it sends a twisted algebra, coalgebra or bialgebra to an algebra, coalgebra or bialgebra.

Remark 1.1. When $V = \mathbb{K}$, we obtain the bosonic Fock functor of [1]:

$$\mathcal{F}[\mathbf{P}] = \bigoplus_{n=0}^{\infty} \text{coInv}(\mathbf{P}[n]) = \bigoplus_{n=0}^{\infty} \frac{\mathbf{P}[n]}{\text{Vect}(\mathbf{P}[\sigma](p) - p \mid \sigma \in \mathfrak{S}_n, p \in \mathbf{P}[n])}.$$

1.2 Contraction-extraction coproducts

Let us now recall the contraction-extraction coproduct of [11]. We shall need the following notations:

Notations 1.1. The species **Com** is defined by $\mathbf{Com}[X] = \mathbb{K}$ for any finite set X and $\mathbf{Com}[\sigma] = \text{Id}_{\mathbb{K}}$ for any bijection σ between two finite sets.

Notations 1.2. 1. Let X be a finite set. We denote by $\mathcal{E}[X]$ the set of equivalences on X .

2. For any bijection $\sigma : X \rightarrow Y$ between two finite sets, for any $\sim \in \mathcal{E}[X]$, we define an equivalence \sim_σ on Y by

$$\forall y, y' \in Y, \quad y \sim_\sigma y' \iff \sigma^{-1}(y) \sim \sigma^{-1}(y').$$

This defines a bijection from $\mathcal{E}[X]$ to $\mathcal{E}[Y]$. Moreover, σ induces a bijection between X/\sim and Y/\sim_σ , sending $C \in X/\sim$ to $\sigma(C) \in Y/\sim_\sigma$.

3. The set $\mathcal{E}[X]$ is partially ordered by the refinement order:

$$\forall \sim, \sim' \in \mathcal{E}[X], \quad \sim \leq \sim' \iff (\forall x, y \in X, x \sim' y \implies x \sim y),$$

that is to say, $\sim \leq \sim'$ if the classes of \sim are unions of classes of \sim' .

4. If $\sim' \in \mathcal{E}[X]$, there is a natural bijection from $\mathcal{E}[X/\sim']$ to $\{\sim \in \mathcal{E}[X], \sim \leq \sim'\}$. From now, we identify $\mathcal{E}[X/\sim']$ and $\{\sim \in \mathcal{E}[X], \sim \leq \sim'\}$, as well as $(X/\sim')/\simeq$ and X/\sim through these bijections.

Let (\mathbf{P}, m, Δ) be a twisted bialgebra. A contraction-extraction coproduct on \mathbf{P} is a family δ of maps such that for any finite set X , for any $\sim \in \mathcal{E}[X]$, $\delta_\sim : \mathbf{P}[X] \rightarrow \mathbf{P}[X/\sim] \otimes \mathbf{P}[X]$, with the following conditions:

- (Compatibility with the species structure [11, Definition 2.2, second item]). For any bijection $\sigma : X \rightarrow Y$ between two finite sets, for any $\sim \in \mathcal{E}[X]$, the following diagram commutes:

$$\begin{array}{ccc} \mathbf{P}[X] & \xrightarrow{\delta_\sim} & \mathbf{P}[X/\sim] \otimes \mathbf{P}[X] \\ \mathbf{P}[\sigma] \downarrow & & \downarrow \mathbf{P}[\sigma/\sim] \otimes \mathbf{P}[\sigma] \\ \mathbf{P}[Y] & \xrightarrow{\delta_{\sim_\sigma}} & \mathbf{P}[Y/\sim_\sigma] \otimes \mathbf{P}[Y] \end{array}$$

- (Coassociativity of δ [11, Definition 2.2, third item]). If X is a finite set and $\sim \leq \sim' \in \mathcal{E}[X]$, the following diagram commutes:

$$\begin{array}{ccc} \mathbf{P}[X] & \xrightarrow{\delta_{\sim'}} & \mathbf{P}[X/\sim'] \otimes \mathbf{P}[X] \\ \delta_\sim \downarrow & & \downarrow \delta_\sim \otimes \text{Id} \\ \mathbf{P}[X/\sim] \otimes \mathbf{P}[X] & \xrightarrow{\text{Id} \otimes \delta_{\sim'}} & \mathbf{P}[X/\sim] \otimes \mathbf{P}[X/\sim'] \otimes \mathbf{P}[X] \end{array}$$

Moreover, if $\sim, \sim' \in \mathcal{E}[X]$, such that we do not have $\sim \leq \sim'$, then $(\text{Id} \otimes \delta_{\sim'}) \circ \delta_\sim = 0$.

- (Counit [11, Definition 2.2, fourth item]). There exists a twisted algebra morphism $\epsilon_\delta : \mathbf{P} \rightarrow \mathbf{Com}$ such that for any finite set X , for any $\sim \in \mathcal{E}[X]$,

$$(\text{Id} \otimes \epsilon_{\delta_X}) \circ \delta_\sim = \begin{cases} \text{Id}_{\mathbf{P}[X]} & \text{if } \sim \text{ is the equality of } X, \\ 0 & \text{otherwise,} \end{cases}$$

and

$$\sum_{\sim \in \mathcal{E}[X]} (\epsilon_{\delta_{X/\sim}} \otimes \text{Id}) \circ \delta_\sim = \text{Id}_{\mathbf{P}[X]}.$$

- (Compatibility with the algebraic structure [11, Proposition 2.4]). For any finite set X , for any $\sim \in \mathcal{E}[X]$, putting $\sim_X = \sim \cap X^2$ and $\sim_Y = \sim \cap Y^2$,

$$\delta_{\sim} \circ m_{X,Y} = \begin{cases} (m_{X/\sim_X, Y/\sim_Y} \otimes m_{X,Y}) \circ (\text{Id} \otimes c \otimes \text{Id}) \circ (\delta_{\sim_X} \otimes \delta_{\sim_Y}) & \text{if } \sim = \sim_X \sqcup \sim_Y, \\ 0 & \text{otherwise,} \end{cases}$$

Moreover, $\delta_{\sim_{\emptyset}}(1_{\mathbf{P}}) = 1_{\mathbf{P}} \otimes 1_{\mathbf{P}}$, where \sim_{\emptyset} is the unique equivalence on \emptyset .

- (Compatibility with the coalgebraic structure [11, Proposition 2.4]). For any finite set $X \sqcup Y$, for any $\sim \in \mathcal{E}[X \sqcup Y]$, putting $\sim_X = \sim \cap X^2$ and $\sim_Y = \sim \cap Y^2$,

$$(\Delta_{X/\sim_X, Y/\sim_Y} \otimes \text{Id}) \circ \delta_{\sim} = m_{1,3,24} \circ (\delta_{\sim_X} \otimes \delta_{\sim_Y}) \circ \Delta_{X,Y} \text{ if } \sim = \sim_X \sqcup \sim_Y.$$

Moreover, for any $x \in \mathbf{P}[\emptyset]$,

$$(\varepsilon_{\Delta} \otimes \text{Id}) \circ \delta_{\sim_{\emptyset}}(x) = \varepsilon_{\Delta}(x) 1_{\mathbf{P}}.$$

The following result is proved in [11, Proposition 3.7]:

Proposition 1.1. *Let (V, \cdot, δ_V) be a counitary, not necessarily unitary bialgebra, and let (\mathbf{P}, m, Δ) be a twisted bialgebra with a contraction-extraction coproduct δ . Then $\mathcal{F}_V[\mathbf{P}]$ is a double bialgebra, with the product and the two coproducts defined by the following:*

$$\begin{aligned} m(\overline{v_1 \dots v_m \otimes p} \otimes \overline{w_1 \dots w_n \otimes q}) &= \overline{\mathbf{P}[\sigma_{m,n}](m(p \otimes q)) \otimes v_1 \dots v_m w_1 \dots w_n}, \\ \Delta(\overline{v_1 \dots v_n \otimes p}) &= \sum_{I \subseteq [n]} \left(\overline{\prod_{i \in I} v_i \otimes \mathbf{P}[\sigma_I](p_I^{(1)})} \right) \otimes \left(\overline{\prod_{i \notin I} v_i \otimes \mathbf{P}[\sigma_{[n] \setminus I}](p_I^{(2)})} \right), \\ \delta(\overline{v_1 \dots v_n \otimes p}) &= \sum_{\sim \in \mathcal{E}[n]} \left(\overline{\prod_{C \text{ class of } \sim} \prod_{i \in C} v'_i} \right) \otimes p'_{\sim} \otimes \overline{v''_1 \dots v''_n \otimes p''_{\sim}}, \end{aligned}$$

where $\sigma_{m,n} : [m] \sqcup [n] \rightarrow [m+n]$ is the bijection sending the elements of $[m]$ to themselves and any element $k \in [n]$ to $k+m$, $\sigma_I : I \rightarrow [|I|]$ is the unique increasing bijection (where I is any subset of $[n]$), and, with Sweedler's notations $\Delta_{I, [n] \setminus I}(p) = p_I^{(1)} \otimes p_I^{(2)}$, $\delta_{\sim}(p) = p'_{\sim} \otimes p''_{\sim}$ and $\delta_V(v) = v' \otimes v''$.

2 The species of mixed graphs

2.1 Mixed graphs

Definition 2.1. 1. A mixed graph is a triple $G = (V(G), E(G), A(G))$ where:

- $V(G)$ is a finite set, called the set of vertices of G ,
- $E(G)$ is a subset of $\{\{x, y\} \subseteq V(G), x \neq y\}$, called the set of edges of G ,
- $A(G)$ is a subset of $\{(x, y) \in V(G)^2, x \neq y\}$, called the set of arcs of G ,


such that, for any $x, y \in V(G)$, with $x \neq y$,

$$\begin{aligned} \{x, y\} \in E(G) &\implies (x, y) \notin A(G) \text{ and } (y, x) \notin A(G), \\ (x, y) \in A(G) &\implies \{x, y\} \notin E(G). \end{aligned}$$

2. For any finite set X , we denote by $\mathcal{G}[X]$ the set of mixed graphs G such that $V(G) = X$. The vector space generated by $\mathcal{G}[X]$ is denoted by $\mathbf{G}[X]$. This defines a species \mathbf{G} of mixed graphs.

3. A mixed graph is an oriented graph if $E(G) = \emptyset$: this defines a subset of $\mathcal{G}[X]$ denoted by $\mathcal{G}_o[X]$ for any finite set X , and a subspecies of \mathbf{G} denoted by \mathbf{G}_o .
4. A mixed graph G is a simple graph if $A(G) = \emptyset$: this defines a subset of $\mathcal{G}[X]$ denoted by $\mathcal{G}_s[X]$ for any finite set X , and a subspecies of \mathbf{G} denoted by \mathbf{G}_s .

Example 2.1. There are five elements in $\mathcal{G}[2]$, which we graphically represent on the right:



Moreover,

$$\mathcal{G}_o[2] = \{G_1, G_3, G_4, G_5\}, \quad \mathcal{G}_s[2] = \{G_1, G_2\}.$$

Remark 2.1. In a mixed graph G , for any pair of vertices $\{x, y\}$ of G , there are five possibilities to define edges or arcs between x and y . Hence, if X is of cardinality n , then

$$|\mathcal{G}[X]| = 5^{\frac{n(n-1)}{2}}.$$

Similarly,

$$|\mathcal{G}_o[X]| = 4^{\frac{n(n-1)}{2}}, \quad |\mathcal{G}_s[X]| = 2^{\frac{n(n-1)}{2}}.$$

These gives respectively sequences A109345, A053763 and A006125 of the OEIS [17].

Notations 2.1. Let G be a mixed graph and $x, y \in V(G)$. We shall write $x \xrightarrow{G} y$ if $(x, y) \in A(G)$ and $x \xrightarrow{G} y$ if $\{x, y\} \in E(G)$.

Definition 2.2. Let G be a mixed graph.

1. An oriented path in G is a finite sequence $P = (x_0, \dots, x_n)$ of vertices of G such that for any $i \in \{0, \dots, n-1\}$, $x_i \xrightarrow{G} x_{i+1}$. The vertices x_0 and x_n are respectively the beginning and the end of P .
2. A mixed path in G is a finite sequence $P = (x_0, \dots, x_n)$ of vertices of G such that for any $i \in \{0, \dots, n-1\}$, $x_i \xrightarrow{G} x_{i+1}$ or $x_i \xrightarrow{G} x_{i+1}$. The vertices x_0 and x_n are the extremities of P .
3. A path in G is a finite sequence $P = (x_0, \dots, x_n)$ of vertices of G such that for any $i \in \{0, \dots, n-1\}$, $x_i \xrightarrow{G} x_{i+1}$ or $x_{i+1} \xrightarrow{G} x_i$ or $x_i \xrightarrow{G} x_{i+1}$. The vertices x_0 and x_n are the extremities of P . The mixed graph G is connected if for any vertices $x, y \in V(G)$, there exists a path of extremities x and y .

Example 2.2. Let us consider the mixed graphs of Example 2.1. The connected ones are G_2, G_3, G_4 and G_5 .

2.2 The twisted bialgebra of mixed graphs

We now define a product and a coproduct on the species \mathbf{G} . Let X, Y be two finite sets, $G \in \mathcal{G}[X]$ and $H \in \mathcal{G}[Y]$. The mixed graph $GH \in \mathcal{G}[X \sqcup Y]$, disjoint union of G and H , is defined by

$$V(GH) = V(G) \sqcup V(H), \quad E(GH) = E(G) \sqcup E(H), \quad A(GH) = A(G) \sqcup A(H).$$

This product is bilinearly extended to \mathbf{G} . It is clearly associative, commutative and has a unit, which is the empty graph $1 \in \mathcal{G}[\emptyset]$. With this product, \mathbf{G} is a commutative twisted algebra. Note that \mathbf{G}_s and \mathbf{G}_o are subalgebras of \mathbf{G} .

Definition 2.3. Let G be a mixed graph and let $I \subseteq V(G)$.

1. The mixed graph $G|_I$ is defined by

$$\begin{aligned} V(G|_I) &= I, \\ E(G|_I) &= \{\{x, y\} \in E(G) \mid x, y \in I\}, \\ A(G|_I) &= \{(x, y) \in A(G) \mid x, y \in I\}. \end{aligned}$$

2. We shall say that I is an ideal of G if

$$\forall x, y \in V(G), \quad (x \in I \text{ and } x \xrightarrow{G} y) \implies (y \in I).$$

Proposition 2.4. We define a coproduct Δ on the species $\mathbf{G}[X]$ in the following way: for any finite sets X and Y , for any mixed graph $G \in \mathcal{G}[X \sqcup Y]$,

$$\Delta_{X,Y}(G) = \begin{cases} G|_X \otimes G|_Y & \text{if } Y \text{ is an ideal of } G, \\ 0 & \text{otherwise.} \end{cases}$$

Together with the product defined earlier, this coproduct makes \mathbf{G} a twisted bialgebra.

Proof. Let us first prove the coassociativity of Δ . Let X, Y and Z be disjoint finite sets and $G \in \mathcal{G}[X \sqcup Y \sqcup Z]$. Then

$$\begin{aligned} (\Delta_{X,Y} \otimes \text{Id}) \circ \Delta_{X \sqcup Y, Z}(G) &= \begin{cases} G|_X \otimes G|_Y \otimes G|_Z \\ \quad \text{if } Z \text{ is an ideal of } G \text{ and } Y \text{ is an ideal of } G|_{X \sqcup Y}, \\ 0 & \text{otherwise.} \end{cases} \\ (\text{Id} \otimes \Delta_{Y,Z}) \circ \Delta_{X, Y \sqcup Z}(G) &= \begin{cases} G|_X \otimes G|_Y \otimes G|_Z \\ \quad \text{if } Y \sqcup Z \text{ is an ideal of } G \text{ and } Z \text{ is an ideal of } G|_{Y \sqcup Z}, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Moreover,

$$\begin{aligned} &Z \text{ is an ideal of } G \text{ and } Y \text{ is an ideal of } G|_{X \sqcup Y} \\ &\iff \left(\forall (x, y) \in V(G)^2, x \xrightarrow{G} y \implies (x, y) \notin (Y \times X) \sqcup (Z \times X) \sqcup (Z \times Y) \right) \\ &\iff Y \sqcup Z \text{ is an ideal of } G \text{ and } Z \text{ is an ideal of } G|_{Y \sqcup Z}, \end{aligned}$$

so $(\Delta_{X,Y} \otimes \text{Id}) \circ \Delta_{X \sqcup Y, Z} = (\text{Id} \otimes \Delta_{Y,Z}) \circ \Delta_{X, Y \sqcup Z}$. As for any graph G , \emptyset and $V(G)$ are ideals of G ,

$$\Delta_{V(G), \emptyset}(G) = G \otimes 1, \quad \Delta_{\emptyset, V(G)}(G) = 1 \otimes G.$$

So Δ is counitary, and the counit is given by

$$\varepsilon_\Delta : \begin{cases} \mathbf{G}[\emptyset] & \longrightarrow \mathbb{K} \\ 1 \text{ (empty graph)} & \longrightarrow 1. \end{cases}$$

Let $G \in \mathcal{G}[X]$, $H \in \mathcal{G}[Y]$ and X', Y' be sets such that $X \sqcup Y = X' \sqcup Y'$. In GH , there is no arc between any element of X and any element of Y , nor between any element of Y and any element of X . Hence, the ideals of GH are of the form $I \sqcup J$, where I is an ideal of G and J is an ideal of H . Therefore,

$$\begin{aligned} \Delta_{X',Y'}(GH) &= \begin{cases} G|_{X \cap X'} H|_{Y \cap X'} \otimes G|_{X \cap Y'} H|_{Y \cap Y'} \\ \quad \text{if } Y' \cap X \text{ is an ideal of } G \text{ and } Y' \cap Y \text{ is an ideal of } H, \\ 0 \text{ otherwise} \end{cases} \\ &= \Delta_{X \cap X', X \cap Y'}(G) \Delta_{Y \cap X', Y \cap Y'}(H). \end{aligned}$$

Moreover, $\Delta_{\emptyset, \emptyset}(1) = 1 \otimes 1$. So Δ is a morphism of twisted algebras. \square

Example 2.3. With the notations of Example 2.1 (or Example ?? below):

	$\Delta_{\{1,2\}, \emptyset}$	$\Delta_{\{1\}, \{2\}}$	$\Delta_{\{2\}, \{1\}}$	$\Delta_{\emptyset, \{1,2\}}$
G_1	$G_1 \otimes 1$	$\textcircled{1} \otimes \textcircled{2}$	$\textcircled{2} \otimes \textcircled{1}$	$1 \otimes G_1$
G_2	$G_2 \otimes 1$	$\textcircled{1} \otimes \textcircled{2}$	$\textcircled{2} \otimes \textcircled{1}$	$1 \otimes G_2$
G_3	$G_3 \otimes 1$	$\textcircled{1} \otimes \textcircled{2}$	0	$1 \otimes G_3$
G_4	$G_4 \otimes 1$	0	$\textcircled{2} \otimes \textcircled{1}$	$1 \otimes G_4$
G_5	$G_5 \otimes 1$	0	0	$1 \otimes G_5$

2.3 Contraction-extraction on mixed graphs

Definition 2.5. Let $G \in \mathcal{G}[X]$ and $\sim \in \mathcal{E}[X]$.

1. We define a mixed graph $G | \sim \in \mathcal{G}[X]$ by

$$\begin{aligned} V(G | \sim) &= V(G), \\ E(G | \sim) &= \{\{x, y\} \in E(G) \mid x \sim y\}, \\ A(G | \sim) &= \{(x, y) \in A(G) \mid x \sim y\}. \end{aligned}$$

In other words, $G | \sim$ is obtained from G by deleting all the edges or arcs which extremities are not equivalent; or equivalently, $G | \sim$ is the disjoint union of the restrictions of G to the equivalence classes of \sim .

2. We define a mixed graph $G / \sim \in \mathcal{G}[X / \sim]$ by

$$\begin{aligned} V(G / \sim) &= X / \sim, \\ E(G / \sim) &= \{\{\text{cl}_\sim(x), \text{cl}_\sim(y)\} \mid \{x, y\} \in E(G), (x, y) \notin A(G), (y, x) \notin A(G), \text{cl}_\sim(x) \neq \text{cl}_\sim(y)\}, \\ A(G / \sim) &= \{(\text{cl}_\sim(x), \text{cl}_\sim(y)) \mid (x, y) \in A(G), \text{cl}_\sim(x) \neq \text{cl}_\sim(y)\}. \end{aligned}$$

In other words, G / \sim is obtained from G by identifying the vertices according to \sim , then deleting the loops created in the process and the redundant edges, giving priority to the oriented ones.

3. We shall say that $\sim \in \mathcal{E}^c[G]$ if for any equivalence class C of \sim , $G|_C$ is connected.

Proposition 2.6. *We define a contraction-extraction coproduct δ on \mathbf{G} by the following: for any finite set X , for any $\sim \in \mathcal{E}[X]$, for any $G \in \mathcal{G}[X]$,*

$$\delta_{\sim}(G) = \begin{cases} G/\sim \otimes G|\sim & \text{if } \sim \in \mathcal{E}^c[G], \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The compatibility of δ with the species structure is clear. Let us prove the coassociativity of δ . Let X be a finite set, $\sim, \sim' \in \mathcal{E}[X]$ and $G \in \mathcal{G}[X]$.

If $\sim \leq \sim'$, let us prove that $\sim \in \mathcal{E}^c[G/\sim']$ and $\sim' \in \mathcal{E}^c[G]$ if, and only if, $\sim' \in \mathcal{E}^c[G|\sim]$ and $\sim \in \mathcal{E}^c[G]$.

\implies . Let C' be a class of \sim' . As $\sim' \in \mathcal{E}^c[G]$, it is a connected subgraph of G . Moreover, as $\sim \leq \sim'$, all its elements are in the same class of \sim , so $G|_{C'} = (G|\sim)|_{C'}$: as a consequence, $(G|\sim)|_{C'}$ is connected, so $\sim' \in \mathcal{E}^c[G|\sim]$. Let C be a class of \sim , and $x, y \in C$. As $\sim \in \mathcal{E}^c[G/\sim']$, it is connected in G/\sim' : there exists a path in G/\sim' from $\text{cl}_{\sim'}(x)$ to $\text{cl}_{\sim'}(y)$. Moreover, as $\sim' \in \mathcal{E}^c[G]$, each $\text{cl}_{\sim'}(z)$ is a connected subgraph of G , so there is a path from x to y in G : $\sim \in \mathcal{E}^c[G]$.

\impliedby . Let C be a class of \sim . As $\sim \in \mathcal{E}^c[G]$, any of its class is a connected subgraph of G , so by contraction is a connected subgraph of G/\sim' : $\sim \in \mathcal{E}^c[G/\sim']$. Let C' be a class of \sim' . As $\sim' \in \mathcal{E}^c[G|\sim]$, it is a connected subgraph of $G|\sim$, so also of G : $\sim' \in \mathcal{E}^c[G]$.

As a conclusion,

$$\begin{aligned} (\delta_{\sim} \otimes \text{Id}) \circ \delta_{\sim'}(G) &= \begin{cases} (G/\sim')/\sim \otimes (G/\sim')|\sim \otimes G|\sim' & \text{if } \sim \in \mathcal{E}^c[G/\sim'] \text{ and } \sim' \in \mathcal{E}^c[G], \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} G/\sim \otimes (G|\sim)/\sim' \otimes (G|\sim)|\sim' & \text{if } \sim' \in \mathcal{E}^c[G|\sim] \text{ and } \sim \in \mathcal{E}^c[G], \\ 0 & \text{otherwise} \end{cases} \\ &= (\text{Id} \otimes \delta_{\sim'}) \circ \delta_{\sim}(G). \end{aligned}$$

If we do not have $\sim \leq \sim'$, then at least one class C of \sim intersects two classes of \sim' , so intersects two connected components of $|\sim'$: we obtain that $\sim \notin \mathcal{E}^c[G|\sim']$. So $\delta_{\sim}(G|\sim') = 0$ and finally $(\text{Id} \otimes \delta_{\sim}) \circ \delta_{\sim'}(G) = 0$.

Let us now study the counit. We define a species morphism $\epsilon_{\delta} : \mathbf{G} \longrightarrow \mathbf{Com}$ by the following: if $G \in \mathcal{G}[X]$,

$$\epsilon_{\delta}[X](G) = \begin{cases} 1 & \text{if } E(G) = A(G) = \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

Let $G \in \mathcal{G}[X]$ and $\sim \in \mathcal{E}[X]$. If \sim is the equality of X , then $\sim \in \mathcal{E}^c[G]$, $G/\sim = G$ and $G|\sim$ as no edge, so $(\text{Id} \otimes \epsilon_{\delta}[X])(G) = G$. Otherwise, either $G \notin \mathcal{E}^c[G]$ or at least one class of \sim contains an edge or an arc, so $\epsilon_{\delta}[X](G|\sim) = 0$. In both cases, $(\text{Id} \otimes \epsilon_{\delta}[X])(G) = 0$.

Let $\sim \in \mathcal{E}^c[G]$, such that $E(G/\sim) = E(G|\sim) = \emptyset$. If two vertices of G are related by an edge of an arc, there are necessarily equivalent, so any connected component of G is included in a single class of \sim . As the classes of \sim are connected, \sim is the relation \sim_c which classes are the connected components of G . Moreover, G/\sim_c has no edge nor arc, and $G|\sim_c = G$. Therefore,

$$\begin{aligned} \sum_{\sim \in \mathcal{E}[X]} (\epsilon_{\delta}[X/\sim] \otimes \text{Id}) \circ \delta_{\sim}(G) &= \sum_{\sim \in \mathcal{E}^c[G]} (\epsilon_{\delta}[X/\sim] \otimes \text{Id}) \circ \delta_{\sim}(G) \\ &= (\epsilon_{\delta}[X/\sim] \otimes \text{Id}) \circ \delta_{\sim_c}(G) \\ &= G|\sim_c \\ &= G. \end{aligned}$$

Let us prove the compatibility of δ with the algebraic structure. Obviously, $\delta_{\sim_\emptyset}(1) = 1 \otimes 1$. Let X and Y be two finite sets, $\sim \in \mathcal{E}[X \sqcup Y]$, $G \in \mathcal{G}[X]$ and $H \in \mathcal{G}[Y]$. If $\sim \neq \sim_X \sqcup \sim_Y$, at least one class C of \sim intersects both X and Y , so is not connected in $GH = m_{X,Y}(G \otimes H)$. Therefore, $\sim \notin \mathcal{E}^c[GH]$ and

$$\delta_{\sim} \circ m_{X,Y}(G \otimes H) = 0.$$

Let us assume that $\sim = \sim_X \sqcup \sim_Y$. Then $\sim \in \mathcal{E}^c[GH]$ if, and only if, $\sim_X \in \mathcal{E}^c[G]$ and $\sim_Y \in \mathcal{E}^c[H]$, as the connected components of GH are the connected components of G and of H . If so, $(GH)/\sim = (G/\sim_X)(H/\sim_Y)$ and $(GH)|\sim = (G|\sim_X)(H|\sim_Y)$. Therefore,

$$\begin{aligned} \delta_{\sim} \circ m_{X,Y}(G \otimes H) &= \begin{cases} (GH)/\sim \otimes (GH)|\sim & \text{if } \sim \in \mathcal{E}^c[GH], \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} (G/\sim_X)(H/\sim_Y) \otimes (G|\sim_X)(H|\sim_Y) & \text{if } \sim_X \in \mathcal{E}^c[G] \text{ and } \sim_Y \in \mathcal{E}^c[H], \\ 0 & \text{otherwise} \end{cases} \\ &= (m_{X/\sim_X, Y/\sim_Y} \otimes m_{X,Y}) \circ (\text{Id} \otimes c \otimes \text{Id}) \circ (\delta_{\sim_X} \otimes \delta_{\sim_Y})(G \otimes H). \end{aligned}$$

Let us finally prove the compatibility of δ with the coalgebraic structure. Obviously,

$$(\varepsilon_{\Delta} \otimes \text{Id}) \circ \delta_{\sim_\emptyset}(1) = 1 = \varepsilon_{\Delta}(1)1.$$

Let X and Y be two finite sets, $\sim_X \in \mathcal{E}[X]$, $\sim_Y \in \mathcal{E}[Y]$ and $G \in \mathcal{G}[X]$. We put $\sim = \sim_X \sqcup \sim_Y$.

$$\begin{aligned} (\Delta_{X/\sim_X, Y/\sim_Y} \otimes \text{Id}) \circ \delta_{\sim}(G) &= \begin{cases} (G/\sim)_{|X/\sim_X} \otimes (G/\sim)_{|Y/\sim_Y} & \text{if } \sim \in \mathcal{E}^c[G] \text{ and } Y/\sim_Y \text{ is an ideal of } G/\sim, \\ 0 & \text{otherwise,} \end{cases} \\ m_{1,3,24} \circ (\delta_{\sim_X} \otimes \delta_{\sim_Y}) \circ \Delta_{X,Y}(G) &= \begin{cases} (G|_X)/\sim_X \otimes (G|_Y)/\sim_Y & \text{if } Y \text{ is an ideal of } G, \sim_X \in \mathcal{E}^c[G|_X] \text{ and } \sim_Y \in \mathcal{E}^c[G|_Y], \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

Let us prove that $\sim \in \mathcal{E}^c[G]$ and Y/\sim_Y is an ideal of G/\sim if, and only if, Y is an ideal of G , $\sim_X \in \mathcal{E}^c[G|_X]$ and $\sim_Y \in \mathcal{E}^c[G|_Y]$.

\implies . Let $y \in Y$ and $z \in X \sqcup Y$ such that $x \xrightarrow{G} y$. Then either $\text{cl}_{\sim}(y) = \text{cl}_{\sim}(z)$ or $\text{cl}_{\sim}(y) \xrightarrow{G/\sim} \text{cl}_{\sim}(z)$. As Y/\sim_Y is an ideal of G/\sim , in both cases $z \in Y$. As $\sim = \sim_X \sqcup \sim_Y$, its classes are the classes of \sim_X and \sim_Y , and are connected by hypothesis. So $\sim_X \in \mathcal{E}^c[G|_X]$ and $\sim_Y \in \mathcal{E}^c[G|_Y]$.

\impliedby . As $\sim = \sim_X \sqcup \sim_Y$, its classes are the classes of \sim_X and \sim_Y , which are connected by hypothesis. Hence, $\sim \in \mathcal{E}^c[G]$. Let $\text{cl}_{\sim}(y) \in Y/\sim_Y$ and $\text{cl}_{\sim}(z) \in [X \sqcup Y]/\sim$, such that $\text{cl}_{\sim}(y) \xrightarrow{G/\sim} \text{cl}_{\sim}(z)$. There exist $y', z' \in X \sqcup Y$ such that $y \sim y'$, $z \sim z'$ and $y' \xrightarrow{G} z'$. As $\sim = \sim_X \sqcup \sim_Y$, $y' \in Y$. As Y is an ideal of G , necessarily $z' \in Y$. As $\sim = \sim_X \sqcup \sim_Y$, $z' \in Y$ and finally $\text{cl}_{\sim}(z') \in Y/\sim_Y$.

Moreover,

$$(G/\sim)_{|X/\sim_X} = (G|_X)/\sim_X, \quad (G/\sim)_{|Y/\sim_Y} = (G|_Y)/\sim_Y,$$

which finally proves the compatibility between δ and Δ . \square

As a consequence, by Proposition 1.1, for any vector space V , we obtain a graded bialgebra $\mathcal{F}_V[\mathbf{G}]$. This is the vector space of mixed graphs which vertices are decorated by elements of V , any graph being linear in any of its decorations: these objects will be called V -linearly decorated graphs. For example, if $v_1, v_2, w_1, w_2 \in V$ and $\lambda_1, \lambda_2, \mu_1, \mu_2 \in \mathbb{K}$, in $\mathcal{F}_V[\mathbf{G}]$, if $v = \lambda_1 v_1 + \lambda_2 v_2$ and $w = \mu_1 w_1 + \mu_2 w_2$,

$$\begin{aligned} \text{Diagram}(v, w) &= \lambda_1 \mu_1 \text{Diagram}(v_1, w_1) + \lambda_2 \mu_1 \text{Diagram}(v_2, w_1) \\ &+ \lambda_1 \mu_2 \text{Diagram}(v_1, w_2) + \lambda_2 \mu_2 \text{Diagram}(v_2, w_2). \end{aligned}$$

If \mathcal{B} is a basis of V , a basis of $\mathcal{F}_V[\mathbf{G}]$ is the set of mixed graphs which vertices are decorated by elements of \mathcal{B} . The product is the disjoint union. For any V -linearly decorated graph G ,

$$\Delta(G) = \sum_{I \text{ ideal of } G} G_{|V(G)\setminus I} \otimes G_{|I}.$$

For example, if $v, w \in V$,

$$\begin{aligned} \Delta((v) \circlearrowleft (w)) &= (v) \circlearrowleft (w) \otimes 1 + 1 \otimes (v) \circlearrowleft (w) + (v) \otimes (w) + (w) \otimes (v), \\ \Delta((v) \text{---} (w)) &= (v) \text{---} (w) \otimes 1 + 1 \otimes (v) \text{---} (w) + (v) \otimes (w) + (w) \otimes (v), \\ \Delta((v) \text{---} (w)) &= (v) \text{---} (w) \otimes 1 + 1 \otimes (v) \text{---} (w) + (v) \otimes (w), \\ \Delta((v) \rightleftarrows (w)) &= (v) \rightleftarrows (w) \otimes 1 + 1 \otimes (v) \rightleftarrows (w). \end{aligned}$$

The counit ε_Δ sends any mixed graph $G \neq 1$ to 0. If (V, \cdot, Δ) is a not necessarily unitary, commutative and cocommutative bialgebra, then $\mathcal{F}_V[\mathbf{G}]$ inherits a second coproduct δ : if G is a V -linearly decorated graph,

$$\delta(G) = \sum_{\sim \in \mathcal{E}^c[G]} G / \sim \otimes G | \sim,$$

where the vertices of $G / \sim \otimes G | \sim$ are decorated in the following way: denoting by $d_G(x)$ the decoration of the vertex $x \in V(G)$, any vertex $\text{cl}_\sim(x)$ of G / \sim is decorated by the products of elements $d_G(y)'$, where $y \in \text{cl}_\sim(x)$, whereas the vertex $x \in V(G | \sim) = V(G)$ is decorated by $d_G(x)''$, and everything being extended by multilinearity of each decoration. The counit ε_δ is given on any mixed graph G by

$$\varepsilon_\delta(G) = \begin{cases} \prod_{x \in V(G)} \varepsilon_V \circ d_G(x) & \text{if } A(G) = E(G) = \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

This construction is functorial in V .

Example 2.4. If $v, w \in V$, with Sweedler's notation $\delta_V(u) = u' \otimes u''$,

$$\begin{aligned} \delta((v) \circlearrowleft (w)) &= (v') \circlearrowleft (w') \otimes (v'') \circlearrowleft (w''), \\ \delta((v) \text{---} (w)) &= (v') \text{---} (w'') \otimes (v'') \circlearrowleft (w'') + (v' \cdot w') \otimes (v'') \text{---} (w''), \\ \delta((v) \text{---} (w)) &= (v') \text{---} (w') \otimes (v'') \circlearrowleft (w'') + (v' \cdot w') \otimes (v'') \text{---} (w''), \\ \delta((v) \rightleftarrows (w)) &= (v') \rightleftarrows (w') \otimes (v'') \circlearrowleft (w'') + (v' \cdot w') \otimes (v'') \rightleftarrows (w''). \end{aligned}$$

Remark 2.2. We shall often work with $V = \mathbb{K}$, with its usual bialgebraic structure defined by $\delta_{\mathbb{K}}(1) = 1 \otimes 1$. We shall then identify any V -decorated mixed graph which any vertex is decorated by 1 with the underlying mixed graph. The double bialgebra $\mathcal{F}_V[\mathbf{G}]$ is identified with $\mathcal{F}[\mathbf{G}]$ and

has for basis the set of (isomorphism classes of) mixed graphs. The coproduct simplifies:

$$\begin{aligned}
\Delta(\bullet \bullet) &= \bullet \bullet \otimes 1 + 1 \otimes \bullet \bullet + \bullet \otimes \bullet + \bullet \otimes \bullet, \\
\Delta(\bullet \text{---} \bullet) &= \bullet \text{---} \bullet \otimes 1 + 1 \otimes \bullet \text{---} \bullet + \bullet \otimes \bullet + \bullet \otimes \bullet, \\
\Delta(\bullet \longrightarrow \bullet) &= \bullet \longrightarrow \bullet \otimes 1 + 1 \otimes \bullet \longrightarrow \bullet + \bullet \otimes \bullet, \\
\Delta(\bullet \overset{\curvearrowright}{\longleftarrow} \bullet) &= \bullet \overset{\curvearrowright}{\longleftarrow} \bullet \otimes 1 + 1 \otimes \bullet \overset{\curvearrowright}{\longleftarrow} \bullet; \\
\\
\delta(\bullet \bullet) &= \bullet \bullet \otimes \bullet \bullet, \\
\delta(\bullet \text{---} \bullet) &= \bullet \text{---} \bullet \otimes \bullet \bullet + \bullet \otimes \bullet \text{---} \bullet, \\
\delta(\bullet \longrightarrow \bullet) &= \bullet \longrightarrow \bullet \otimes \bullet \bullet + \bullet \otimes \bullet \longrightarrow \bullet, \\
\delta(\bullet \overset{\curvearrowright}{\longleftarrow} \bullet) &= \bullet \overset{\curvearrowright}{\longleftarrow} \bullet \otimes \bullet \bullet + \bullet \otimes \bullet \overset{\curvearrowright}{\longleftarrow} \bullet.
\end{aligned}$$

Remark 2.3. If V is finite-dimensional, then, considering the number of vertices, $\mathcal{F}_V[\mathbf{G}]$ is a graded bialgebra, which homogeneous components are finite-dimensional. The dimension of the homogeneous component of degree n of $\mathcal{F}_V[\mathbf{G}]$ is given by the number of isomorphism classes of mixed graphs which vertices are decorated by elements of $[N]$, where $N = \dim(V)$, see the Appendix for more details on these numbers.

Proposition 2.7. *Let V be a (non necessarily unitary) commutative and cocommutative bialgebra. For any linearly V -decorated mixed graph G , we denote by $d_G : V(G) \rightarrow V$ the decoration map of G and by \overline{G} the underlying mixed graph. Then the following map is a double bialgebra morphism:*

$$\Theta_V : \begin{cases} \mathcal{F}_V[\mathbf{G}] & \rightarrow \mathcal{F}[\mathbf{G}] \\ G & \mapsto \left(\prod_{x \in V(G)} \epsilon_V \circ d_G(x) \right) \overline{G}. \end{cases}$$

Proof. The counit $\epsilon_V : V \rightarrow \mathbb{K}$ is a bialgebra map. By functoriality (in V), Θ_V is a double bialgebra morphism. \square

2.4 Cofreeness of the coalgebra $\mathcal{F}[\mathbf{G}]$

Let us define a second product on $\mathcal{F}[\mathbf{G}]$.

Proposition 2.8. *Let G and H be two mixed graphs. The mixed graph $G \curvearrowright H$ is defined by*

$$\begin{aligned}
V(G \curvearrowright H) &= V(G) \sqcup V(H), \\
E(G \curvearrowright H) &= E(G) \sqcup E(H), \\
A(G \curvearrowright H) &= A(G) \sqcup A(H) \sqcup (V(G) \times V(H)).
\end{aligned}$$

This product is bilinearly extended to $\mathcal{F}[\mathbf{G}]$. Then $(\mathcal{F}[\mathbf{G}], \curvearrowright, \Delta)$ is a unital infinitesimal bialgebra in the sense of [15, Definition 2.1].

Proof. As we already know that Δ is coassociative and unitary, it remains to prove that:

1. \curvearrowright is associative and unitary.
 2. For any $x, y \in \mathcal{F}[\mathbf{G}]$, $\Delta(x \curvearrowright y) = (x \otimes 1) \curvearrowright \Delta(y) + \Delta(x) \curvearrowright (1 \otimes y) - x \otimes y$.
1. Let G, H, K be three mixed graphs. Then

$$\begin{aligned}
V((G \curvearrowright H) \curvearrowright K) &= V(G \curvearrowright (H \curvearrowright K)) = V(G) \sqcup V(H) \sqcup V(K), \\
E((G \curvearrowright H) \curvearrowright K) &= E(G \curvearrowright (H \curvearrowright K)) = E(G) \sqcup E(H) \sqcup E(K), \\
A((G \curvearrowright H) \curvearrowright K) &= A(G \curvearrowright (H \curvearrowright K)) = A(G) \sqcup A(H) \sqcup A(K) \sqcup (V(G) \times V(H)) \\
&\quad \sqcup (V(G) \times V(K)) \sqcup (V(H) \times V(K)),
\end{aligned}$$

so $(G \curvearrowright H) \curvearrowright K = G \curvearrowright (H \curvearrowright K)$. Therefore, \curvearrowright is associative. The unit is the empty mixed graph 1.

2. Let G, H be two mixed graph. As there is an arc from any vertex of G to any vertex of H in $G \curvearrowright H$, the ideals of $G \curvearrowright H$ are:

- $I \sqcup V(H)$ where I is an ideal of G . For such an ideal,

$$(G \curvearrowright H)_{|I \sqcup V(H)} = G_{|I} \curvearrowright H, \quad (G \curvearrowright H)_{|V(G \curvearrowright H) \setminus (I \sqcup V(H))} = G_{|V(G) \setminus I}.$$

- Ideals J of H . For such an ideal,

$$(G \curvearrowright H)_{|J} = H_{|J}, \quad (G \curvearrowright H)_{|V(G \curvearrowright H) \setminus J} = G \curvearrowright H_{|V(H) \setminus J}.$$

Note that the ideal $V(H)$ appears twice in this list, for $I = \emptyset$ and $J = V(H)$. Therefore,

$$\begin{aligned} \Delta(G \curvearrowright H) &= \sum_{I \text{ ideal of } G} G_{|V(G) \setminus I} \otimes G_{|I} \curvearrowright H + \sum_{J \text{ ideal of } G} G \curvearrowright H_{|V(H) \setminus J} \otimes H_{|J} - G \otimes H \\ &= \Delta(G) \curvearrowright (1 \otimes H) + (G \otimes 1) \curvearrowright \Delta(H) - G \otimes H. \quad \square \end{aligned}$$

From [15, Theorem 2.6]:

Corollary 2.9. *The coalgebra $(\mathcal{F}[\mathbf{G}], \Delta)$ is isomorphic to the coalgebra $T(\text{Prim}(\mathcal{F}[\mathbf{G}]))$ with the deconcatenation coproduct.*

Remark 2.4. The same proof can be adapted to any $\mathcal{F}_V[\mathbf{G}]$.

3 Sub-objects and quotients of mixed graphs

3.1 Simple and oriented graphs

Proposition 3.1. \mathbf{G}_s and \mathbf{G}_o are twisted subbialgebras of (\mathbf{G}, m, Δ) and are stable under the contraction-extraction coproduct δ .

Proof. If G and H are simple graphs, then GH is a simple graph. If G is a simple graph, then all its subgraphs are also simple graphs. Moreover, if $\sim \in \mathcal{E}^c[G]$, then G/\sim and $G|\sim$ are also simple graphs. The proof is similar for oriented graphs. \square

Corollary 3.2. *For any vector space V , $\mathcal{F}_V[\mathbf{G}_s]$ is a subbialgebra of $\mathcal{F}_V[\mathbf{G}]$ and $\mathcal{F}_V[\mathbf{G}_o]$ is a subbialgebra of $\mathcal{F}_V[\mathbf{G}]$. For any (non necessarily unitary) commutative and cocommutative bialgebra V , $\mathcal{F}_V[\mathbf{G}_s]$ is a double subbialgebra of $\mathcal{F}_V[\mathbf{G}]$ and $\mathcal{F}_V[\mathbf{G}_o]$ is a double subbialgebra of $\mathcal{F}_V[\mathbf{G}]$.*

In particular, $\mathcal{F}_{\mathbb{K}}[\mathbf{G}_s] = \mathcal{F}[\mathbf{G}_s]$ is the double bialgebra of graphs of [8, 9] and $\mathcal{F}_{\mathbb{K}}[\mathbf{G}_o] = \mathcal{F}[\mathbf{G}_o]$ is the double bialgebra of [6].

3.2 Acyclic mixed graphs and finite topologies

Let us recall the following definition:

Definition 3.3. *Let X be a finite set.*

1. *A topology on X is a subset \mathcal{O} of the set of subsets of X such that:*

- \emptyset and X belong to \mathcal{O} .
- If $O_1, O_2 \in \mathcal{O}$, then $O_1 \cap O_2 \in \mathcal{O}$ and $O_1 \cup O_2 \in \mathcal{O}$.

2. The set of topologies on X is denoted by $\mathcal{Topo}[X]$ and the space generated by $\mathcal{Topo}[X]$ is denoted by $\mathbf{Topo}[X]$. This defines a species \mathbf{Topo} .

3. A topology \mathcal{O} is T_0 if for any $x, y \in X$, with $x \neq y$, there exists $O \in \mathcal{O}$ such that $(x \in O$ and $y \notin O)$ or $(x \notin O$ and $y \in O)$.

Example 3.1. Let $G \in \mathcal{G}[X]$ be a mixed graph. We denote by \mathcal{O}_G the set of ideals of G .

Let us prove this reformulation of Alexandroff's theorem [2]:

Lemma 3.4. *Let X be a finite set and \mathcal{O} be a topology on X . There exists an oriented graph G such that $\mathcal{O}_G = \mathcal{O}$.*

Proof. Let $\mathcal{O} \in \mathcal{Topo}[X]$. We define a relation \leq on X by the following: for any $x, y \in X$, $x \leq y$ if any $O \in \mathcal{O}$ containing x also contains y . We then define an oriented graph G by $V(G) = X$ and

$$E(G) = \{(x, y) \in X^2 \mid x \neq y \text{ and } x \leq y\}.$$

Let us prove that $\mathcal{O}_G = \mathcal{O}$. Let $O \in \mathcal{O}$, $x \in O$ and $y \in X$ such that $x \xrightarrow{G} y$. Then $x \leq y$: by definition of \leq , $y \in O$. So $O \in \mathcal{O}_G$. Therefore, $\mathcal{O} \subseteq \mathcal{O}_G$. Let $O \in \mathcal{O}_G$. As O is an ideal of G ,

$$\begin{aligned} O &= \{y \in X \mid \exists x \in O, x \leq y\} \\ &= \bigcup_{x \in O} \{y \in X \mid x \leq y\} \\ &= \bigcup_{x \in X} \left(\bigcap_{O' \in \mathcal{O}, x \in O'} O' \right). \end{aligned}$$

As \mathcal{O} is a topology, $O \in \mathcal{O}$, so $O = \mathcal{O}_G$. □

Proposition 3.5. *The species \mathbf{Topo} is equipped with a twisted bialgebra structure by the following:*

- For any finite sets X, Y , for any $(\mathcal{O}_X, \mathcal{O}_Y) \in \mathcal{Topo}[X] \times \mathcal{Topo}[Y]$,

$$m_{X,Y}(\mathcal{O}_X \times \mathcal{O}_Y) = \{I \sqcup J, I \in \mathcal{O}_X, J \in \mathcal{O}_Y\}.$$

- For any finite sets X, Y , for any $\mathcal{O} \in \mathcal{Topo}[X \sqcup Y]$,

$$\Delta_{X,Y}(\mathcal{O}) = \begin{cases} \mathcal{O}|_X \otimes \mathcal{O}|_Y & \text{if } Y \in \mathcal{O}, \\ 0 & \text{otherwise,} \end{cases}$$

where

$$\mathcal{O}|_X = \{X \cap O \mid O \in \mathcal{O}\}, \quad \mathcal{O}|_Y = \{Y \cap O \mid O \in \mathcal{O}\}.$$

Moreover, the following map is a surjective morphism of twisted bialgebras:

$$\Upsilon : \begin{cases} \mathbf{G} & \longrightarrow \mathbf{Topo} \\ G \in \mathcal{G}[X] & \longrightarrow \mathcal{O}_G \in \mathcal{Topo}[X]. \end{cases}$$

Proof. The map Υ is clearly a species morphism. By Lemma 3.4, it is surjective. Let G, G', H, H' be graphs such that $\mathcal{O}_G = \mathcal{O}_{G'}$ and $\mathcal{O}_H = \mathcal{O}_{H'}$. Then

$$\mathcal{O}_{GH} = \mathcal{O}_G \mathcal{O}_H = \mathcal{O}_{G'} \mathcal{O}_{H'} = \mathcal{O}_{G'H'}.$$

Therefore, the product of \mathbf{G} is compatible with the products of \mathbf{G} and \mathbf{Topo} .

For any graph $G \in \mathcal{G}[X]$ and for any $Y \subset X$, $(\mathcal{O}_G)|_Y = \mathcal{O}_{G|_Y}$. This implies that ν is compatible with the coproducts of \mathbf{G} and \mathbf{Topo} . As Υ is surjective and \mathbf{G} is a twisted bialgebra, \mathbf{Topo} is also a twisted bialgebra. □

This map Υ is not compatible with the contraction-extraction coproduct: for example, if



then $\mathcal{O}_{G_1} = \mathcal{O}_{G_2} = \{\{1, 2, 3\}, \{2, 3\}, \{2\}, \emptyset\}$. Let us denote by \sim the equivalence with classes $\{1, 2\}$ and $\{3\}$. Then


$$\delta_{\sim}(G) = \begin{array}{c} \textcircled{1, 2} \\ \leftarrow \quad \rightarrow \\ \textcircled{3} \end{array} \otimes \textcircled{1} \text{---} \textcircled{2} \textcircled{3}, \quad \delta_{\sim}(G') = 0.$$

In order to obtain a second coproduct, we have to restrict ourselves to acyclic mixed graphs:

Proposition 3.6. *Let G be a mixed graph. We shall say that G is acyclic if does not contain any oriented path $x_0 \xrightarrow{G} \dots \xrightarrow{G} x_n$ with $x_0 = x_n$ and $n \geq 2$. Acyclic mixed graphs and acyclic oriented graphs form two twisted subbialgebras of (\mathbf{G}, m, Δ) .*

Proof. Obviously, if G and H are acyclic mixed graphs, then GH is acyclic; if G is acyclic and $I \subseteq V(G)$, then $G|_I$ is also acyclic. Therefore, \mathbf{G}_{ac} is a twisted subbialgebra of \mathbf{G} and \mathbf{G}_{aco} is a twisted subbialgebra of \mathbf{G}_o . \square

Remark 3.1. \mathbf{G}_{ac} is not stable under δ . For example, let us consider the following acyclic oriented graph:



Let us denote by \sim the equivalence with classes $\{1, 2\}$ and $\{3\}$. Then

$$\delta_{\sim}(G) = \begin{array}{c} \textcircled{1, 2} \\ \leftarrow \quad \rightarrow \\ \textcircled{3} \end{array} \otimes \textcircled{1} \text{---} \textcircled{2} \textcircled{3}.$$

Proposition 3.7. *A contraction-extraction coproduct on \mathbf{G}_{ac} is defined by the following: for any acyclic graph $G \in \mathcal{G}_{ac}[X]$, for any $\sim \in \mathcal{E}[X]$,*

$$\delta_{\sim}(G) = \begin{cases} G / \sim \otimes G | \sim & \text{if } \sim \in \mathcal{E}^{c,ac}[G], \\ 0 & \text{otherwise.} \end{cases}$$

where $\mathcal{E}^{c,ac}[G]$ the set of equivalences on $V(G)$ such that the classes of G are connected and G / \sim is acyclic. Moreover, the following map is a surjective morphisms of twisted bialgebras, compatible with the contraction-extraction coproducts:

$$\varpi_0 : \begin{cases} \mathbf{G} & \longrightarrow \mathbf{G}_{ac} \\ G \in \mathcal{G}[X] & \longmapsto \begin{cases} G & \text{if } G \text{ is acyclic,} \\ 0 & \text{otherwise.} \end{cases} \end{cases} \quad (2)$$

Proof. Let \mathbf{I} be the subspecies of \mathbf{G} of non acyclic mixed graphs. If G is a non acyclic mixed graph, then for any mixed graph H , GH is not acyclic: \mathbf{I} is an ideal. If I is an ideal of G , if it contains a vertex on a cycle of G , then it contains all the vertices of the cycle: therefore, $G|_I$ or $G_{V(G) \setminus I}$ is not acyclic, which proves that \mathbf{I} is a coideal for Δ . Let $\sim \in \mathcal{E}[G]$. Let us consider a cycle C of G . If all the vertices of C are equivalent for \sim , then $G | \sim$ contains a cycle, so is not acyclic. Otherwise, G / \sim contains a cycle: \mathbf{I} is a coideal for δ . Identifying the species \mathbf{G}/\mathbf{I} and \mathbf{G}_{ac} via ϖ_0 , \mathbf{G}_{ac} inherits a contraction-extraction coproduct δ , which is precisely the one defined in this proposition. \square

Similarly, restricting ϖ_0 to \mathbf{G}_o , its image \mathbf{G}_{aco} inherits a contraction-extraction coproduct δ , as a sub-quotient of \mathbf{G} . The image of acyclic graphs by Υ is given by T_0 -topologies:

Definition 3.8. *Let X be a finite set and \mathcal{O} a topology on X . We shall say that \mathcal{O} is T_0 if for any $x \neq y \in X$, there exists $O \in \mathcal{O}$ such that $(x \in O \text{ and } y \notin O)$ or $(x \notin O \text{ and } y \in O)$. This defines a subset $\mathcal{Topo}_{T_0}[X]$ of $\mathcal{Topo}[X]$ subspecies \mathbf{Topo}_{T_0} of \mathbf{Topo} .*

Lemma 3.9. *Let X be a finite set and $G \in \mathcal{G}[X]$. The topology \mathcal{O}_G is T_0 if, and only if, G is acyclic. $x_0 \xrightarrow{G} \dots \xrightarrow{G} x_n$ with $x_0 = x_n$ and $n \geq 2$.*

Proof. Let us assume that G has a cycle $x_0 \xrightarrow{G} \dots \xrightarrow{G} x_n$ with $x_0 = x_n$ and $n \geq 2$. Then any ideal of G containing one of the x_i 's contains all the x_i 's, so \mathcal{O}_G is not T_0 . Let us assume that G is acyclic. Let $x \neq y \in X$. We consider

$$\begin{aligned} O_x &= \{z \in X \mid \exists k \in \mathbb{N}, \exists x_1, \dots, x_k \in X, x \xrightarrow{G} x_1 \xrightarrow{G} \dots \xrightarrow{G} x_k \xrightarrow{G} z\}, \\ O_y &= \{z \in X \mid \exists k \in \mathbb{N}, \exists y_1, \dots, y_k \in X, y \xrightarrow{G} y_1 \xrightarrow{G} \dots \xrightarrow{G} y_k \xrightarrow{G} z\}. \end{aligned}$$

Both O_x and O_y are ideals of G , so belong to \mathcal{O} . Moreover, for $k = 0$, $x \in O_x$ and $y \in O_y$. If $x \in O_y$ and $y \in O_x$, then there exists $k, l \in \mathbb{N}$ and $x_1, \dots, x_k, y_1, \dots, y_l \in X$ such that

$$: x \xrightarrow{G} x_1 \dots \xrightarrow{G} x_k \xrightarrow{G} y \xrightarrow{G} y_1 \dots \xrightarrow{G} y_l \xrightarrow{G} x,$$

which contradicts the acyclicity of G . So $y \notin O_x$ or $x \notin O_y$: \mathcal{O}_G is T_0 . \square

Lemma 3.10. *Let G be an oriented graph. Then \mathcal{O}_G is connected if and only if G is connected.*

Proof. Let us assume that \mathcal{O}_G is not connected. Let $O_1, O_2 \in \mathcal{O}_G$, both nonempty, such that $V(G) = O_1 \sqcup O_2$. If $x \xrightarrow{G} y$, with $x \in O_1$, then as O_1 is an ideal, $y \in O_1$. Consequently, there is no arc from a vertex of O_1 to a vertex of O_2 . Symmetrically, there is no arc from a vertex of O_2 to a vertex of O_1 . So G is not connected. Let us assume that G is not connected. We can write $V(G) = O_1 \sqcup O_2$, such that there is no arc from a vertex of O_1 to a vertex of O_2 , nor from a vertex of O_2 to a vertex of O_1 . Consequently, $O_1, O_2 \in \mathcal{O}_G$, so \mathcal{O}_G is not connected. \square

Proposition 3.11. *There exists a unique product, a unique coproduct and a unique contraction-extraction coproduct on \mathbf{Topo}_{T_0} making the map $\Upsilon|_{\mathbf{G}_{aco}} : \mathbf{G}_{aco} \longrightarrow \mathbf{Topo}_{T_0}$ a morphism of twisted bialgebras, compatible with the contraction-extraction coproduct.*

Proof. By Lemma 3.9, $\Upsilon(\mathbf{G}_{aco}) = \mathbf{Topo}_{T_0}$, which gives the unicity of the contraction-extraction coproduct on \mathbf{Topo}_{T_0} compatible with Υ . The product and the coproduct on \mathbf{Topo}_{T_0} are obviously the restriction of the product and of the coproduct on finite topologies.

Let X be a finite set, $G, G' \in \mathbf{G}_{aco}[X]$ such that $\Upsilon(G) = \Upsilon(G')$, and let $\sim \in \mathcal{E}[X]$. Obviously, $\Upsilon(G|\sim) = \Upsilon(G'|\sim)$. Moreover,

$$\mathcal{O}_{G/\sim} = \{\pi_{\sim}(O) \mid O \in \mathcal{O}_G\} = \{\pi_{\sim}(O) \mid O \in \mathcal{O}_{G'}\} = \mathcal{O}_{G'/\sim},$$

where $\pi_{\sim} : X \longrightarrow X/\sim$ is the canonical surjection. So $\Upsilon(G/\sim) = \Upsilon(G'/\sim)$. Let us now prove that $\mathcal{E}^{c,ac}[G] = \mathcal{E}^{c,ac}[G']$. If $\sim \in \mathcal{E}^{c,ac}[G]$, then G/\sim is acyclic, so $\Upsilon(G/\sim)$ is T_0 , by Lemma 3.9, so $\Upsilon(G'/\sim) = \Upsilon(G/\sim)$ is T_0 and G'/\sim is acyclic. Moreover, by Lemma 3.10, the connected components of $\Upsilon(G|\sim) = \Upsilon(G'|\sim)$ are the connected components of the oriented graph $G|\sim$, that is to say the classes of \sim as $\sim \in \mathcal{E}^{c,ac}[G]$. Consequently, if C is a class of \sim , $(\mathcal{O}_{G'})|_C$ is connected: $\sim \in \mathcal{E}^{c,ac}[G']$. By symmetry, we obtain $\mathcal{E}^{c,ac}[G] = \mathcal{E}^{c,ac}[G']$.

As a consequence, for any $\sim \in \mathcal{E}[X]$,

$$\begin{aligned} (\Upsilon \otimes \Upsilon) \circ \delta_{\sim}(G) &= \begin{cases} \Upsilon(G/\sim) \otimes \Upsilon(G|\sim) & \text{if } \sim \in \mathcal{E}^{c,ac}[G], \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} \Upsilon(G'/\sim) \otimes \Upsilon(G'|\sim) & \text{if } \sim \in \mathcal{E}^{c,ac}[G'], \\ 0 & \text{otherwise} \end{cases} \\ &= (\Upsilon \otimes \Upsilon) \circ \delta_{\sim}(G). \end{aligned}$$

Consequently, \mathbf{Topo}_{T_0} inherits a contraction-extraction coproduct as a quotient of \mathbf{G}_{aco} . \square

Remark 3.2. Let X be a finite set and \mathcal{O} be a topology on X . For any $x, y \in X$, we shall say that $x \sim_{\mathcal{O}} y$ if any $O \in \mathcal{O}$ containing x or y contains both x and y . This defines an equivalence $\sim_{\mathcal{O}}$ on X . Moreover, $X/\sim_{\mathcal{O}}$ inherits from \mathcal{O} a topology, which turns out to be T_0 . In other words, we obtain that $\mathbf{Topo}[X]$ and $\bigoplus_{\sim \in \mathcal{E}[X]} \mathbf{Topo}_{T_0}[X/\sim]$ are isomorphic, which gives a species isomorphism between \mathbf{Topo} and $\mathbf{Topo}_{T_0} \circ \mathbf{Com}$, where here \circ is the composition of species. The double twisted bialgebra structure which we obtain in this way is described in [8]. Applying Aguiar and Mahajan's bosonic Fock functor [1], we obtain the double algebra of finite topologies of [7].

3.3 Totally acyclic graphs


Definition 3.12. Let G be a mixed graph. We shall say that it is totally acyclic if does not contain any mixed path (x_0, \dots, x_n) , with $x_0 = x_n$ and $n \geq 2$. Totally acyclic graphs form a subspecies \mathbf{G}_{tac} of \mathbf{G} .

Note that totally acyclic mixed graphs are simply called acyclic in [5].


Proposition 3.13. \mathbf{G}_{tac} is a twisted subbialgebra of \mathbf{G} .

Proof. If G and H are totally acyclic graphs, then GH is totally acyclic. So \mathbf{G}_{tac} is a twisted subalgebra of \mathbf{G} . Let G be a totally acyclic mixed graph and $I \subseteq V(G)$. As G does not contain any mixed cycle, so does $G|_I$: $G|_I$ is totally acyclic. As a conclusion, \mathbf{G}_{tac} is a twisted subcoalgebra of \mathbf{G} . \square

Consequently, for any vector space V , $\mathcal{F}_V[\mathbf{G}_{tac}]$ is a subbialgebra of $(\mathcal{F}_V[\mathbf{G}], m, \Delta)$. The subspecies \mathbf{G}_{tac} is not stable under δ . For example, considering the mixed graph



which is totally acyclic, the equivalence relation \sim which classes are $\{x, y\}$, $\{z\}$ and $\{t\}$ belongs to $\mathcal{E}^c[G]$ (in fact, even to $\mathcal{E}^{c,ac}[G]$), and



is clearly not totally acyclic.

4 Applications

4.1 Three polynomial invariants

Let (V, \cdot, δ_V) be a non necessarily unitary, commutative bialgebra. From [10, Theorem 3.9], there exists a unique morphism ϕ_1 of double bialgebras from $(\mathcal{F}_V[\mathbf{G}], m, \Delta, \delta)$ onto $(\mathbb{K}[X], m, \Delta, \delta)$ where the two coproducts of $\mathbb{K}[X]$ are defined by

$$\Delta(X) = X \otimes 1 + 1 \otimes X, \quad \delta(X) = X \otimes X.$$

Let us determine ϕ_1 , firstly when $V = \mathbb{K}$. Let $G \in \mathcal{G}[X]$, nonempty. Then, still by [10, Theorem 3.9],

$$\phi_1(G) = \sum_{k=0}^{\infty} \epsilon_{\delta}^{\otimes(k-1)} \circ \tilde{\Delta}^{(k-1)}(G) H_k(X),$$

where H_k is the k -th Hilbert polynomial:

$$H_k(X) = \frac{X(X-1)\dots(X-k+1)}{k!}.$$

Notations 4.1. For any $k \geq 1$, for any mixed graph G , we denote by $L_k(G)$ the set of surjections $c : V(G) \rightarrow [k]$ such that

$$\forall x, y \in V(G), \quad x \xrightarrow{G} y \implies c(x) \leq c(y).$$

By definition of the coproduct Δ , for any mixed graph G with $n \geq 1$ vertices,

$$\tilde{\Delta}^{(k-1)}(G) = \sum_{c \in L_k(G)} G|_{c^{-1}(1)} \otimes \dots \otimes G|_{c^{-1}(k)}.$$

This leads to the following definition:

Definition 4.1. *Let G be a mixed graph.*

1. *A valid coloring of G is a map $c : V(G) \rightarrow \mathbb{N}_{>0}$ such that*

$$\forall x, y \in V(G), \quad \begin{cases} x \xrightarrow{G} y & \implies c(x) < c(y), \\ x \xrightarrow{G} y & \implies c(x) \neq c(y). \end{cases}$$

2. *A valid coloring c of G is packed if $c(V(G)) = [\max(c)]$. The set of valid packed colorings of G is denoted by $\text{VPC}(G)$.*

Thanks to this definition, we obtain, for any V -linearly decorated mixed graph G ,

$$\epsilon_{\delta}^{\otimes(k-1)} \circ \tilde{\Delta}^{(k-1)}(G) = |\{c \in \text{VPC}(G), |\max(c) = k\}|,$$

and finally:

Proposition 4.2. *The unique morphism of double bialgebras from $\mathcal{F}[\mathbf{G}]$ to $\mathbb{K}[X]$ is given on any mixed graph G by*

$$P_{chr_S}(G) = \sum_{c \in \text{VPC}(G)} H_{\max(c)}.$$

Consequently, if $N \in \mathbb{N}$, $P_{chr_S}(G)(N)$ is the number of valid colorings c such that $\max(c) \leq N$: we recover the (strong) chromatic polynomial $P_{chr_S}(G)$ of [5].

Remark 4.1. If V is a non necessarily unitary, commutative bialgebra, the unique double bialgebra morphism from $(\mathcal{F}_V[\mathbf{G}], m, \Delta, \delta)$ to $(\mathbb{K}[X], m, \Delta, \delta)$ is $P_{chr_S} \circ \Theta_V$ (which is indeed a double bialgebra morphism by composition). It sends any V -linearly decorated mixed graph G to

$$P_{chr_S} \circ \Theta_V(G) = \left(\prod_{x \in V(G)} \epsilon_V \circ d_G(x) \right) P_{chr_S}(\overline{G}),$$

where d_G is the decoration map of G and \overline{G} the underlying mixed graph.

Let us now recover the weak chromatic polynomial of [5].

Definition 4.3. *Let G be a mixed graph.*

1. *A weak valid coloring of G is a map $c : V(G) \rightarrow \mathbb{N}_{>0}$ such that*

$$\begin{aligned} \forall x, y \in V(G), \quad & x \xrightarrow{G} y \implies c(x) \leq c(y), \\ & x \xrightarrow{G} y \implies c(x) \neq c(y). \end{aligned}$$

2. *A weak valid coloring c of G is packed if $c(V(G)) = [\max(c)]$. The set of weak valid packed colorings of G is denoted by $\text{WVPC}(G)$.*

We are going to use the action \leftarrow of the monoid of characters on the set of morphisms and the map θ given in (1). Let $\lambda_W : \mathcal{F}[G] \rightarrow \mathbb{K}$ defined on any mixed graph G by

$$\lambda_W(G) = \begin{cases} 1 & \text{if } E(G) = \emptyset, \\ 0 & \text{otherwise.} \end{cases} \quad (3)$$

This is obviously a character. We consider $\theta(\lambda_W) = P_{chr_S} \leftarrow \lambda_W = P_{chr_W}$. This is a bialgebra morphism from $(\mathcal{F}[\mathbf{G}], m, \Delta)$ to $(\mathbb{K}[X], m, \Delta)$. For any mixed graph G ,

$$P_{chr_W}(G) = \sum_{c \in \text{WVPC}(G)} H_{\max(c)}.$$

Consequently, if $N \in \mathbb{N}$, $P_{chr_W}(G)(N)$ is the number of weak valid colorings c such that $\max(c) \leq N$: we recover the weak chromatic polynomial of [5]. Therefore:

Corollary 4.4. $P_{chr_W} = P_{chr_S} \leftarrow \lambda_W : (\mathcal{F}[\mathbf{G}], m, \Delta) \rightarrow (\mathbb{K}[X], m, \Delta)$ is a Hopf algebra morphism.

Remark 4.2. Let $G \in \mathcal{G}[X]$. We denote by $\mathcal{E}_W^c[G]$ the set of equivalences $\sim \in \mathcal{E}^c[G]$ such that

$$\forall x, y \in V(G), \quad x \xrightarrow{G} y \implies x \not\sim y.$$


Then, for any mixed graph G ,

$$P_{chr_W}(G) = \sum_{\sim \in \mathcal{E}^c[G]} \lambda_W(G | \sim) P_{chr_S}(G / \sim) = \sum_{\sim \in \mathcal{E}_W^c[G]} P_{chr_S}(G / \sim).$$

Example 4.1. For $n \geq 3$, let G_n be the following mixed graph:

$$V(G_n) = [n], \quad E(G_n) = \{\{1, n\}\}, \quad A(G_n) = \{(i, i+1) \mid i \in [n-1]\}.$$

In other terms,



Weak valid colorings of G_n are non-decreasing maps $c : [n] \longrightarrow \mathbb{N}_{>0}$, such that $c(1) \neq c(n)$. Therefore,

$$P_{chr_W}(G_n) = \frac{X(X+1)\dots(X+n-1)}{n!} - X.$$

Valid colorings of G_n are strictly increasing maps $c : [n] \longrightarrow \mathbb{N}_{>0}$. Therefore,

$$P_{chr_S}(G_n) = \frac{X(X-1)\dots(X-n+1)}{n!}.$$

With the help of [10, Propositions 3.10 and 5.2], we now define a homogeneous morphism $P_0 : \mathcal{F}[\mathbf{G}] \longrightarrow \mathbb{K}[X]$ with the help of the element $\mu \in \mathcal{F}[\mathbf{G}]_1^*$ defined by

$$\mu(\bullet) = 1,$$

where \bullet is the unique (up to an isomorphism) mixed graph with only one vertex. Then, if G is a mixed graph,

$$\mu^{\otimes k} \circ \tilde{\Delta}^{(k-1)}(G) = \begin{cases} |L_k(G)| & \text{if } k = |V(G)|, \\ 0 & \text{otherwise.} \end{cases}$$

We denote by $\ell(G)$ the cardinality of $L_n(G)$, that is to say the number of bijections $c : V(G) \longrightarrow [n]$ such that

$$\forall x, y \in V(G), \quad x \xrightarrow{G} y \implies c(x) < c(y),$$

and finally:

Corollary 4.5. *For any mixed graph G , we put*

$$\lambda_0(G) = \frac{\ell(G)}{|V(G)|!}, \quad P_0(G) = \lambda_0(G)X^{|V(G)|}.$$

Then λ_0 is a character of $\mathcal{F}[\mathbf{G}]$ and $P_0 : (\mathcal{F}[\mathbf{G}], m, \Delta) \longrightarrow (\mathbb{K}[X], m, \Delta)$ is a bialgebra morphism.

For any graph G , $P_0(G)(1) = \lambda_0(G)$. From [10, Corollary 3.11], $P_0 = P_{chr_S} \leftarrow \lambda_0$. Therefore:

Corollary 4.6. *For any mixed graph G with n vertices,*

$$\ell(G)X^n = \sum_{\sim \in \mathcal{E}^c[G]} \ell(G|\sim)P_{chr_S}(G/\sim).$$

4.2 Invertible characters

Let us fix a non unitary, commutative and cocommutative bialgebra (V, \cdot, δ_V) . The product of the dual algebra is denoted by \star_V ; its unit is the counit ϵ_V . Let us now study the monoid of characters of $(\mathcal{F}_V[\mathbf{G}], m, \delta)$, which product is denoted by \star , and in particular let us look for its invertible. We shall use the following lemma:

Lemma 4.7. *Let (B, m, δ) be a graded bialgebra. In particular, its homogeneous component of degree 0 is a subbialgebra. Let λ be a character of B . Then λ is an invertible character of B if, and only if, its restriction λ_0 to B_0 is invertible in the algebra B_0^* . In the particular case where B_0 is generated by a family $(x_i)_{i \in I}$ of group-like elements, λ is an invertible character if, and only if, $\lambda(x_i) \neq 0$ for any $i \in I$.*

Proof. We shall denote by π_k the canonical projection on B_k for any $k \in \mathbb{N}$. We put

$$\rho_L = (\pi_0 \otimes \text{Id}) \circ \delta, \quad \rho_R = (\pi \otimes \pi_0) \circ \delta.$$

As $\pi_0 : B \rightarrow B_0$ is a bialgebra map, (B, ρ_L, ρ_R) is a B_0 -bicomodule. For any $x \in B_n$ with $n \geq 1$, we put

$$\delta'(x) = \delta(x) - \rho_L(x) - \rho_R(x).$$

By homogeneity of δ ,

$$\delta(x) = \sum_{i=0}^n (\pi_i \otimes \pi_{n-i}) \circ \delta(x) = \rho_L(x) + \underbrace{\sum_{i=1}^{n-1} (\pi_i \otimes \pi_{n-i}) \circ \delta(x)}_{\delta'(x)} + \rho_R(x).$$

\implies . Let us denote by μ the inverse of λ in the monoid of characters of B . We put $\mu_0 = \mu|_{B_0}$. For any $x \in B_0$,

$$\lambda_0 * \mu_0(x) = (\lambda \otimes \mu) \circ \delta_0(x) = (\lambda \otimes \mu) \circ \delta(x) = \lambda * \mu(x) = \epsilon(x).$$

Similarly, $\mu_0 * \lambda_0(x) = \epsilon(x)$, so λ_0 is invertible in B_0^* .

\longleftarrow . Let us define $\mu_n : B_n \rightarrow \mathbb{K}$ for any n , such that if $x \in B_n$,

$$\left(\lambda \otimes \sum_{i=0}^n \mu_i \right) \circ \delta(x) = \epsilon(x).$$

We proceed by induction on n . If $n = 0$, we us take μ_0 the inverse of λ_0 in B_0^* . Let us assume $n \geq 1$ and μ_0, \dots, μ_{n-1} defined. We first define $\nu : B_n \rightarrow \mathbb{K}$ by

$$\forall x \in B_n, \quad \nu(x) = \epsilon(x) - \left(\lambda \otimes \sum_{i=0}^{n-1} \mu_i \right) (\delta'(x) + \rho_R(x)).$$

This is well-defined, as $\delta'(x) + \rho(x) \in B \otimes \bigoplus_{i=0}^{n-1} B_i$. We then put $\mu_n = (\mu_0 \otimes \nu) \circ \rho_L : B_n \rightarrow \mathbb{K}$.

As $\rho_L = (\pi_0 \otimes \text{Id}) \circ \delta$, by homogeneity of δ , $\rho_L(B_n) \subseteq B_0 \otimes B_n$ and μ_n is well-defined. For any $x \in B_n$,

$$\begin{aligned} \left(\lambda \otimes \sum_{i=0}^n \mu_i \right) \circ \delta(x) &= (\lambda \otimes \mu_n) \circ \rho_L(x) + \epsilon(x) - \nu(x) \\ &= (\lambda \otimes \mu \circ \nu) \circ (\text{Id} \otimes \rho_L) \circ \rho_L(x) + \epsilon(x) - \nu(x) \\ &= (\lambda \otimes \mu \circ \nu) \circ (\delta_0 \otimes \text{Id}) \circ \rho_L(x) + \epsilon(x) - \nu(x) \\ &= ((\lambda_0 * \mu_0) \circ \nu) \circ \rho_L(x) + \epsilon(x) - \nu(x) \\ &= (\epsilon_0 \circ \nu) \circ \rho_L(x) + \epsilon(x) - \mu(x) \\ &= \nu(x) + \epsilon(x) - \mu(x) \\ &= \epsilon(x). \end{aligned}$$

Considering $\mu = \sum_{i=0}^{\infty} \mu_i \in B^*$, by construction $\lambda * \mu = \epsilon$. Similarly, we can define $\mu' \in B^*$ such that $\mu' * \lambda = \epsilon$. Then, as the convolution product $*$ is associative, $\mu' = \mu$ and λ is invertible in B^* . Let us now prove that μ is a character. We work in the algebra $(B \otimes B)^*$, which convolution

product is also denoted by $*$. For any $x, y \in B$, with Sweedler's notation $\delta(z) = \sum_z z^{(1)} \otimes z^{(2)}$ for any $z \in B$,

$$\begin{aligned} (\mu \circ m) * (\lambda \circ m)(x \otimes y) &= \sum_x \sum_y \mu(x^{(1)}y^{(1)})\lambda(x^{(1)}y^{(1)}) \\ &= \sum_{xy} \mu((xy)^{(1)})\lambda((xy)^{(2)}) \\ &= \epsilon(xy) \\ &= \epsilon(x)\epsilon(y) \\ &= \epsilon_{B \otimes B}(x \otimes y). \end{aligned}$$

Similarly, $(\lambda \circ m) * (\mu \circ m) = \epsilon_{B \otimes B}$, so, as λ is a character,

$$\mu \circ m = (\lambda \circ m)^{*^{-1}} = (\lambda \otimes \lambda)^{*^{-1}} = \mu \otimes \mu.$$

So μ is indeed a character of B .

Let us now consider the particular case where B_0 is generated by a family $(x_i)_{i \in I}$ of group-like elements.

\implies . If λ is an invertible character, denoting its inverse by μ , for any $i \in I$,

$$\lambda * \mu(x_i) = \epsilon(x_i) = 1 = \lambda(x_i)\mu(x_i),$$

so $\lambda(x_i) \neq 0$.

\impliedby . Let us assume that λ is a character of B such that $\lambda(x_i) \neq 0$ for any $i \in I$. In order to prove that λ is an invertible character, it is enough to prove that λ_0 is invertible in B_0^* . By hypothesis, B_0 has a basis $(y_j)_{j \in J}$ of noncommutative monomials in $(x_i)_{i \in I}$. By multiplicativity, for any $j \in J$, y_j is group-like and $\lambda(y_j) \neq 0$. We then define $\mu_0 \in B_0^*$ by

$$\forall j \in J, \quad \mu_0(y_j) = \frac{1}{\lambda(y_j)}.$$

Then for any $j \in J$,

$$\lambda_0 * \mu_0(y_j) = \mu_0 * \lambda_0(y_j) = \lambda_0(y_j)\mu_0(y_j) = 1 = \epsilon(y_j),$$

so λ_0 is invertible in B_0^* . □

In order to use this lemma, let us introduce a gradation of $(\mathcal{F}_V[\mathbf{G}], m, \delta)$.

Proposition 4.8. *For any V -linearly decorated mixed graph G , we denote by $\text{cc}(G)$ the number of connected components of G and we put*

$$\text{deg}(G) = |V(G)| - \text{cc}(G).$$

This defines a gradation of the bialgebra $(\mathcal{F}_V[\mathbf{G}], m, \delta)$.

Proof. Note that for any graph G , $\text{deg}(G) \geq 0$. Let G and H be two V -linearly decorated mixed graphs. Then

$$|V(GH)| = |V(G)| + |V(H)|, \quad \text{cc}(GH) = \text{cc}(G) + \text{cc}(H),$$

so $\text{deg}(GH) = \text{deg}(G) + \text{deg}(H)$. Let G be a V -linearly decorated mixed graph and $\sim \in \mathcal{E}^c[G]$. We denote by $\text{cl}(\sim)$ the number of equivalence classes of \sim . As $\sim \in \mathcal{E}^c[G]$,

$$|V(G|\sim)| = |V(G)|, \quad \text{cc}(G|\sim) = \text{cl}(\sim).$$

Moreover, the connected components of G/\sim are the contractions of the connected components of G , so

$$|V(G/\sim)| = \text{cl}(\sim), \quad \text{cc}(G/\sim) = \text{cc}(G).$$

We obtain that $\deg(G/\sim) + \deg(G|\sim) = |V(G)| - \text{cl}(\sim) + \text{cl}(\sim) - \text{cc}(G) = \deg(G)$. So $(\mathcal{F}_V[\mathbf{G}], m, \delta)$ is graded. \square

For any graph G , $\deg(G) = 0$ if, and only if, $E(G) = A(G) = \emptyset$. The subalgebra $\mathcal{F}_V[\mathbf{G}]_{\deg=0}$ of elements of degree 0 is the symmetric algebra generated by elements $\bigcirc v$, with $v \in V$. The coproduct of such an element is given by the coproduct of V ,

$$\delta(\bigcirc v) = \bigcirc v' \otimes \bigcirc v''.$$

Proposition 4.9. *Let $\lambda \in \text{Char}(\mathcal{F}_V[\mathbf{G}])$. We define a map $\lambda_V \in V^*$ by*

$$\forall v \in V, \quad \lambda_V(v) = \lambda(\bigcirc v).$$

Then λ is invertible in $(\text{Char}(\mathcal{F}_V[\mathbf{G}]), \star)$ if, and only if, λ_V is invertible in (V^, \star_V) .*

Proof. \implies . Let us assume that λ is an invertible character. Denoting by μ its inverse, $\mu|_V$ provides an inverse of λ_V in V^* .

\impliedby . Let us assume that λ_V is invertible in V^* . By Lemma 4.7, it is enough to prove that λ_0 is invertible in the algebra $\mathcal{F}_V[\mathbf{G}]_0$. By construction of the graduation, $\mathcal{F}_V[\mathbf{G}]_0$ is the symmetric algebra generated by V . Extending multiplicatively the inverse of λ_V to $\mathcal{F}_V[\mathbf{G}]_0$, we obtain an inverse of λ_0 . \square

In the particular case where $V = \mathbb{K}$:

Corollary 4.10. *Let λ be a character of $\mathcal{F}[\mathbf{G}]$. It is invertible in the monoid $(\text{Char}(\mathcal{F}[\mathbf{G}]), \star)$ if, and only if,*

$$\lambda(\bullet) \neq 0.$$

Proof. This is implied by Lemma 4.7, with the family of group-like elements reduced to \bullet . \square

In particular, λ_W and λ_0 are invertible. Their inverses are denoted respectively by ν_W and μ_S . We also put $\mu_W = \mu_S \star \lambda_W$. Then, as $P_{chr_W} = P_{chr_S} \leftarrow \lambda_W$ and $P_0 = P_{chr_S} \leftarrow \lambda_0$, we obtain

$$P_{chr_S} = P_{chr_W} \leftarrow \nu_W, \quad P_{chr_S} = P_0 \leftarrow \mu_S, \quad P_{chr_W} = P_0 \leftarrow \mu_W.$$

Proposition 4.11. *For any mixed graph G ,*

$$P_{chr_S}(G) = \sum_{\sim \in \mathcal{E}^c[G]} \lambda_0(G/\sim) \mu_S(G|\sim) X^{\text{cl}(\sim)} = \sum_{\sim \in \mathcal{E}^c[G]} \nu_W(G|\sim) P_{chr_W}(G/\sim),$$

$$P_{chr_W}(G) = \sum_{\sim \in \mathcal{E}^c[G]} \lambda_0(G/\sim) \mu_W(G|\sim) X^{\text{cl}(\sim)},$$

where $\text{cl}(\sim)$ is the number of classes of \sim .

Corollary 4.12. *If G is a connected mixed graph, then $\mu_S(G)$ is the coefficient of X in $P_{chr_S}(G)$ whereas $\mu_W(G)$ is the coefficient of X in $P_{chr_W}(G)$. Moreover, $\lambda_0(G)$ is the coefficient of $X^{|V(G)|}$ in both $P_{chr_S}(G)$ and $P_{chr_W}(G)$.*

Proof. As G is connected, the unique element \sim of $\mathcal{E}^c[G]$ with $\text{cl}(\sim) = 1$ is \sim_L , which has for only class $V(G)$. So the coefficient of X in $P_{chr_S}(G)$ is, as λ_0 and ϵ coincide on $\mathcal{F}_V[\mathbf{G}]_{\text{deg}=0}$, equal to

$$\lambda_0(G/\sim_L)\mu_S(G|\sim_L) = (\lambda_0 \otimes \mu_S) \circ \rho_L(G) = (\epsilon \otimes \mu_S) \circ \rho_L(G) = \mu_S(G).$$

Similarly, the unique equivalence $\sim \in \mathcal{E}^c[G]$ such that $\text{cl}(\sim) = |V(G)|$ is \sim_R , which has for classes the connected components the singletons (or in other words, \sim_R is the equality of $V(G)$). So the coefficient of $X^{|V(G)|}$ in $P_{chr_S}(G)$ is, as μ_S and ϵ coincide on $\mathcal{F}_V[\mathbf{G}]_{\text{deg}=0}$ equal to

$$\lambda_0(G/\sim_R)\mu_S(G|\sim_R) = (\lambda_0 \otimes \mu_S) \circ \rho_R(G) = (\lambda_0 \otimes \epsilon) \circ \rho_R(G) = \lambda_0(G).$$

The proof is similar for the weak chromatic polynomial. \square

Remark 4.3. One can define an infinitesimal character μ'_S by the following: for any mixed graph G ,

$$\mu'_S(G) = \begin{cases} \mu_S(G) & \text{if } G \text{ is connected,} \\ 0 & \text{otherwise.} \end{cases}$$

This infinitesimal character is equal to $\ln(\epsilon_\delta)$ and is studied in [10, Proposition 4.1]. It is closely related to the eulerian idempotent. Consequently, as the coefficient of X in $H_k(X)$ is $\frac{(-1)^{k+1}}{k}$ if $k \geq 1$, we obtain that for any connected mixed graph G ,

$$\mu_S(G) = \sum_{c \in \text{VPC}(G)} \frac{(1)^{\max(c)+1}}{\max(c)} = \sum_{k=1}^{\infty} |\{c \in \text{VPC}(G) \mid \max(c) = k\}| \frac{(-1)^{k+1}}{k}.$$

When G is a rooted tree, we recover Murua's coefficients [3, 16], which appeared in the analysis of the continuous Baker–Campbell–Hausdorff problem.

Example 4.2.

G	$P_{chr_S}(G)$	$P_{chr_W}(G)$	$\lambda_0(G)$	$\mu_S(G)$	$\mu_W(G)$	$\nu_W(G)$
	$X(X-1)$	$X(X-1)$	1	-1	-1	0
	$\frac{X(X-1)}{2}$	$\frac{X(X+1)}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	-1
	0	X	0	0	1	-1
	$X(X-1)^2$	$X(X-1)^2$	1	1	1	0
	$\frac{X(X-1)^2}{2}$	$\frac{X(X+1)(X-1)}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	0
	$\frac{X(X-1)^2}{2}$	$\frac{X(X+1)(X-1)}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	0
	0	$X(X-1)$	0	0	-1	0
	$\frac{X(X-1)(X-2)}{6}$	$\frac{X(X+1)(X+2)}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{3}$	1
	$\frac{X(2X-1)(X-1)}{6}$	$\frac{X(2X+1)(X+1)}{6}$	$\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{6}$	1
	0	$\frac{X(X+1)}{2}$	0	0	$\frac{1}{2}$	1
	0	$\frac{X(X+1)}{2}$	0	0	$\frac{1}{2}$	1
	0	X	0	0	1	1

Example 4.3. Let us consider again the graph G_n of Example 4.1. We obtain

$$\mu_S(G_n) = \frac{(-1)^n}{n!}, \quad \mu_W(G_n) = \frac{(n-1)!}{n!} - 1 = -\frac{n-1}{n}, \quad \lambda_0(G_n) = \frac{1}{n!}.$$

As an example of application:

Proposition 4.13. 1. $P_{chr_S} \circ \varpi_0 = P_{chr_S}$.

2. Let G be an acyclic mixed graph. Then $P_{chr_S}(G)$ is of degree $|V(G)|$ and its leading term is $\ell(G)$.

Proof. 1. Note that $P_{chr_S} \circ \varpi_0$ and P_{chr_S} are bialgebra morphisms from $\mathcal{F}[\mathbf{G}]$ to $\mathbb{K}[X]$. Moreover,

$$\epsilon_\delta \circ P_{chr_S} \circ \varpi_0 = \epsilon_\delta \circ \varpi_0 = \epsilon_\delta = \epsilon_\delta \circ P_{chr_S}.$$

By unicity, $P_{chr_S} \circ \varpi_0 = P_{chr_S}$.

2. Let G be an acyclic graph. As $P_{chr_S} = P_0 \leftarrow \mu_S$,

$$P_{chr_S}(G) = \sum_{\sim \in \mathcal{E}_c[G]} \mu_S(G | \sim) \ell(G / \sim) X^{\text{cl}(\sim)},$$

which implies that $\deg(P_{chr_S}(G)) \leq |V(G)|$. If \sim is the equality of $V(G)$, then $G | \sim$ is a graph with no edge, so $\mu_S(G | \sim) = 1$. We obtain that

$$P_{chr_S}(G) = \ell(G) X^{|V(G)|} + \text{terms of degree } < |V(G)|.$$

As G has no cycle, $\ell(G) \neq 0$, so $\deg(P_{chr_S}(G)) = |V(G)|$. □

5 From mixed graphs to acyclic oriented graphs

5.1 A double bialgebra epimorphism

Definition 5.1. Let G a mixed graph. An orientation of G is an oriented graph H with $V(G) = V(H)$ and $A(H) = A(G) \sqcup E'$, where E' is a set of arcs in bijection with $E(G)$, through a bijection respecting the extremities. Such an orientation of G is acyclic if H is an acyclic oriented graph. We denote by $\mathcal{O}_{ac}(G)$ the set of acyclic orientations of G .

Theorem 5.2. Let G be a mixed graph. We put

$$\Theta(G) = \sum_{G' \in \mathcal{O}_{ac}(G)} G',$$

with the usual convention that this sum is 0 if $\mathcal{O}_{ac}(G)$ is empty. Then Θ is a double twisted bialgebra morphism from \mathbf{G} to \mathbf{G}_{aco} .

Proof. Firstly, Θ is indeed a species morphism from \mathbf{G} to \mathbf{G}_{aco} . Let G and H be two mixed graphs, G' and H' be orientations of G and H . Then $G'H'$ is an acyclic orientation of GH if, and only if, G' and H' are acyclic orientations of G and H . This implies directly that $\Theta(GH) = \Theta(G)\Theta(H)$. Therefore, Θ is a twisted algebra morphism.

Let $G \in \mathbf{G}[I \sqcup J]$ be a mixed graph. If J is not an ideal of G , then $\Delta_{I,J}(G) = 0$. Moreover, for any orientation G' of G , J is not an ideal of G' , so $\Delta_{I,J}(G') = 0$. Hence, in this case,

$$(\Theta \otimes \Theta) \circ \Delta_{I,J}(G) = \Delta_{I,J} \circ \Theta(G) = 0.$$

We now assume that J is an ideal of G . Then

$$\begin{aligned} (\Theta \otimes \Theta) \circ \Delta_{I,J}(G) &= \sum_{(G', G'') \in \mathcal{O}_{ac}(G|_I) \times \mathcal{O}_{ac}(G|_J)} G' \otimes G'', \\ \Delta_{I,J} \circ \Theta(G) &= \sum_{\substack{H \in \mathcal{O}_{ac}(G), \\ J \text{ ideal of } H}} H|_I \otimes H|_J. \end{aligned}$$

We put

$$A = \mathcal{O}_{ac}(G|_I) \times \mathcal{O}_{ac}(G|_J), \quad B = \{H \in \mathcal{O}_{ac}(G) \mid J \text{ ideal of } H\},$$

and we consider the map

$$\psi : \begin{cases} B & \longrightarrow A \\ H & \longrightarrow (H|_I, H|_J). \end{cases}$$

This is obviously well-defined. We now consider the map $\psi' : A \longrightarrow B$, sending any pair (G', G'') to an orientation $\psi(G', G'') = H$ of G defined in this way: for any edge $\{x, y\}$ of G ,

- If $x, y \in I$, then orient this edge as in G' .
- If $x, y \in J$, then orient this edge as in G'' .
- If $x \in I$ and $y \in J$, then orient this edge from x to y .
- If $x \in J$ and $y \in I$, then orient this edge from y to x .

As there is no arc in H from J to I , J is an ideal of H . Moreover, as G' and G'' are acyclic, H is acyclic: ψ' is well-defined. We immediately obtain that $\psi' \circ \psi = \text{Id}_B$ and $\psi \circ \psi' = \text{Id}_A$, so ψ and ψ' are bijections. Therefore,

$$(\Theta \otimes \Theta) \circ \Delta_{I,J}(G) = \sum_{(G', G'') \in A} G' \otimes G'' = \sum_{H \in B} H|_I \otimes H|_J = \Delta_{I,J} \circ \Theta(G).$$

So $\Theta : (\mathbf{G}, m, \Delta) \longrightarrow (\mathbf{G}_{aco}, m, \Delta)$ is a twisted bialgebra morphism.

Let $G \in \mathbf{G}[I \sqcup J]$ be a mixed graph and $\sim \in \mathcal{E}(G)$. If H is an orientation of G , as the paths in G and H are the same, $\sim \in \mathcal{E}_\sim(G)$ if, and only if, $\sim \in \mathcal{E}_\sim(H)$. Hence, if $\sim \notin \mathcal{E}_c(G)$,

$$\delta_\sim \circ \Theta(G) = (\Theta \otimes \Theta) \circ \delta_\sim(G) = 0.$$

Let us now assume that $\sim \in \mathcal{E}_c(G)$. Then

$$\begin{aligned} \delta_\sim \circ \Theta(G) &= \sum_{\substack{H \in \mathcal{O}_{ac}(G), \\ H/\sim \text{ acyclic}}} H/\sim \otimes H|\sim, \\ (\Theta \otimes \Theta) \circ \delta_\sim(G) &= \sum_{(G', G'') \in \mathcal{O}_{ac}(G/\sim) \times \mathcal{O}_{ac}(G|\sim)} G' \otimes G''. \end{aligned}$$

We put

$$C = \mathcal{O}_{ac}(G/\sim) \times \mathcal{O}_{ac}(G|\sim), \quad D = \{H \in \mathcal{O}_{ac}(G) \mid H/\sim \text{ acyclic}\}.$$

If $H \in D$, then H/\sim is an acyclic orientation of G/\sim by definition of D and $H|\sim$ is an acyclic orientation of $G|\sim$ by restriction. This defines a map

$$\phi : \begin{cases} C & \longrightarrow D \\ H & \longrightarrow (H/\sim, H|\sim). \end{cases}$$

Let us now consider $(G', G'') \in C$. We define an orientation of G in the following way: if $\{x, y\}$ is an edge of G ,

- If $x \sim y$, then $\{x, y\}$ is an edge of $G|\sim$: orient it as in G'' .
- Otherwise, $\{\bar{x}, \bar{y}\}$ is an edge or an arc of G/\sim : orient $\{x, y\}$ as $\{\bar{x}, \bar{y}\}$ in G' : as G' is acyclic, this is unambiguous.

Note that $H/\sim = G'$ and $H|\sim = G''$ by construction. Moreover, this is an acyclic orientation of G : if $x_1 \rightarrow \dots \rightarrow x_k \rightarrow x_1$ is a cycle in H , as G' is acyclic, necessarily $x_1 \sim \dots \sim x_k$, so this is a cycle in G'' : as G'' is acyclic, this is not possible. Moreover, $H/\sim = G'$ is acyclic, so this defines a map $\phi' : D \rightarrow C$ such that $\phi \circ \phi' = \text{Id}_D$.

Let $H \in C$. We put $H' = \phi' \circ \phi(H)$. Let (x, y) be an arc of H . If $x \sim y$, then (x, y) is an arc of $H|\sim$, so is an arc of H' . Otherwise, (\bar{x}, \bar{y}) is an arc of $H/\sim = H'/\sim$. If (y, x) is an arc of H' , then $\bar{x} \rightarrow \bar{y} \rightarrow \bar{x}$ is a cycle in H'/\sim , so H/\sim is not acyclic: this is a contradiction. So (x, y) is an arc of H' . Therefore, H and H' have the same arcs, so are equal. We proved that $\phi' \circ \phi = \text{Id}_C$, so ϕ is a bijection. We obtain

$$\delta_\sim \circ \Theta(G) = \sum_{H \in D} H/\sim \otimes H|\sim = \sum_{(G', G'') \in C} G' \otimes G'' = (\Theta \otimes \Theta) \circ \delta_\sim(G).$$

So Θ is compatible with δ . It is obviously compatible with the unit and both counits ε_Δ and ε_δ . \square

Corollary 5.3. *For any mixed graph G ,*

$$P_{chr_S}(G) = \sum_{H \in \mathcal{O}_{ac}(G)} P_{chr_S}(H).$$

Proof. By functoriality, $\mathcal{F}[\Theta] : \mathcal{F}[\mathbf{G}] \rightarrow \mathcal{F}[\mathbf{G}_{aco}]$ is a double bialgebra morphism. By composition, $P_{chr_S} \circ \mathcal{F}[\Theta] : \mathcal{F}[\mathbf{G}] \rightarrow \mathbb{K}[X]$ is a double bialgebra morphism. By unicity of such a morphism,

$$P_{chr_S} \circ \mathcal{F}[\Theta] = P_{chr_S}. \quad \square$$

5.2 Erhart polynomials for mixed graphs

Proposition 5.4. *Let $q \in \mathbb{K}$. The two following maps are characters of $\mathcal{F}[\mathbf{G}]$:*

$$\text{ehr}_{str}^{(q)} : \begin{cases} \mathcal{F}[\mathbf{G}] & \longrightarrow \mathbb{K} \\ G & \longmapsto \begin{cases} q^{|V(G)|} & \text{if } A(G) = \emptyset, \\ 0 & \text{otherwise,} \end{cases} \end{cases} \quad \text{ehr}^{(q)} : \begin{cases} \mathcal{F}[\mathbf{G}] & \longrightarrow \mathbb{K} \\ G & \longmapsto q^{|V(G)|}, \end{cases}$$

with the convention $q^0 = 1$ even if $q = 0$. Let $\text{Ehr}_{str}^{(q)}$ and by $\text{Ehr}^{(q)}$ be Hopf algebra morphisms from $(\mathcal{F}[\mathbf{G}], m, \Delta)$ to $(\mathbb{K}[X], m, \Delta)$ given by

$$\begin{aligned} \text{Ehr}_{str}^{(q)} &= \theta \left(\text{ehr}_{str}^{(q)} \right) = P_{chr_S} \leftarrow \text{ehr}_{str}^{(q)}, \\ \text{Ehr}^{(q)} &= \theta \left(\text{ehr}^{(q)} \right) = P_{chr_S} \leftarrow \text{ehr}^{(q)}. \end{aligned}$$

where θ is defined in (1).

$$\varepsilon_\delta \circ \text{Ehr}_{str}^{(q)} = \text{ehr}_{str}^{(q)}, \quad \varepsilon_\delta \circ \text{Ehr}^{(q)} = \text{ehr}^{(q)}.$$

Then, for any $n \in \mathbb{N}$, for any mixed graph G ,

$$\begin{aligned} \text{Ehr}_{str}^{(q)}(G)(n) &= q^{|V(G)|} |\{f : V(G) \rightarrow [n] \mid \forall x, y \in V(G), x \xrightarrow{G} y \implies f(x) < f(y)\}|, \\ \text{Ehr}^{(q)}(G)(n) &= q^{|V(G)|} |\{f : V(G) \rightarrow [n] \mid \forall x, y \in V(G), x \xrightarrow{G} y \implies f(x) \leq f(y)\}|. \end{aligned}$$

Moreover, $\text{Ehr}_{str}^{(q)} \circ \varpi_0 = \text{Ehr}_{str}^{(q)}$ (recall that ϖ_0 is defined in (2)).

From now, we shall write simply ehr_{str} , ehr , Ehr_{str} and Ehr for $\text{ehr}_{str}^{(1)}$, $\text{ehr}^{(1)}$, $\text{Ehr}_{str}^{(1)}$ and $\text{Ehr}^{(1)}$.

Proof. The maps $\text{ehr}^{(q)}$ and $\text{ehr}_{str}^{(q)}$ are obviously characters of $\mathcal{F}[\mathbf{G}]$. Let G be a mixed graph. For $k \geq 1$, we denote by $S_k(G)$ the set of surjective maps $f : V(G) \rightarrow [k]$ such that for any $x, y \in V(G)$,

$$x \xrightarrow{G} y \implies f(x) \leq f(y).$$

Then

$$\begin{aligned} \text{Ehr}_{str}^{(q)}(G) &= \sum_{k=1}^{\infty} \sum_{f \in S_k(G)} \text{ehr}_{str}^{(q)}(G|_{f^{-1}(1)}) \cdots \text{ehr}_{str}^{(q)}(G|_{f^{-1}(k)}) H_k(X) \\ &= \sum_{k=1}^{\infty} \sum_{f \in S_k(G)} q^{|f^{-1}(1)|} \text{ehr}_{str}^{(1)}(G|_{f^{-1}(1)}) \cdots q^{|f^{-1}(k)|} \text{ehr}_{str}^{(1)}(G|_{f^{-1}(k)}) H_k(X) \\ &= q^{|V(G)|} \sum_{k=1}^{\infty} \sum_{f \in S_k(G)} \text{ehr}_{str}^{(1)}(G|_{f^{-1}(1)}) \cdots \text{ehr}_{str}^{(1)}(G|_{f^{-1}(k)}) H_k(X) \\ &= q^{|V(G)|} \text{Ehr}_{str}(G), \end{aligned}$$

Similarly, $\text{Ehr}^{(q)}(G) = q^{|V(G)|} \text{Ehr}(G)$. We now study Ehr_{str} and Ehr . Let G be a mixed graph.

$$\text{Ehr}(G) = \sum_{k=1}^{\infty} \sum_{f \in S_k(G)} H_k(X),$$

which gives the announced result. For $k \geq 1$, we denote by $S'_k(G)$ the set of surjective maps $f : V(G) \rightarrow [k]$ such that for any $x, y \in V(G)$,

$$x \xrightarrow{G} y \implies f(x) < f(y).$$

By definition of ehr_{str} ,

$$\text{Ehr}_{str}(G) = \sum_{k=1}^{\infty} \sum_{f \in S_k(G)} \text{ehr}_{str}(G|_{f^{-1}(1)}) \cdots \text{ehr}_{str}(G|_{f^{-1}(k)}) H_k(X) = \sum_{k=1}^{\infty} \sum_{f \in S'_k(G)} H_k(X),$$

which implies the announced result.

Let us prove that $\text{Ehr}_{str}^{(q)} \circ \varpi_0 = \text{Ehr}_{str}^{(q)}$. For this, it is enough to prove that $\epsilon_\delta \circ \text{Ehr}_{str}^{(q)} \circ \varpi_0 = \epsilon_\delta \circ \text{Ehr}_{str}^{(q)}$, that is to say $\text{ehr}_{str}^{(q)} \circ \varpi_0 = \text{ehr}_{str}^{(q)}$. Let G be a graph. If G is not acyclic, then $\text{ehr}_{str}^{(q)} \circ \varpi_0(G) = 0$. Moreover, necessarily $A(G) \neq \emptyset$, so $\text{ehr}_{str}^{(q)} = 0$. Otherwise, $\varpi_0(G) = G$ and $\text{ehr}_{str}^{(q)} \circ \varpi_0(G) = \text{ehr}_{str}^{(q)}(G)$. \square

Remark 5.1. By [10, Corollary 3.12],

$$\text{Ehr}_{str} = P_{chr_S} \leftarrow \text{ehr}_{str}, \quad \text{Ehr} = P_{chr_S} \leftarrow \text{ehr}.$$

Notations 5.1. Let G be a mixed graph. We denote by $S(G)$ the set of vertices $y \in V(G)$ such that there exists no $e \in A(G)$ such that y is the final vertex of e (set of *sources* of G) and by $W(G)$ the set of vertices $x \in V(G)$ such that there exists no $e \in A(G)$ such that x is the initial vertex of e (set of *wells* of G).

Proposition 5.5. *Let $q, q' \in \mathbb{K}$. For any mixed graph G , denoting by $*$ the convolution product associated to Δ ,*

$$\begin{aligned} \text{ehr}_{str}^{(q)} * \text{ehr}_{str}^{(q')}(G) &= q^{|V(G) \setminus S(G)|} (q + q')^{|S(G)|}, \\ \text{ehr}_{str}^{(q)} * \text{ehr}_{str}^{(q')}(G) &= q^{|V(G) \setminus W(G)|} (q + q')^{|W(G)|}. \end{aligned}$$

In particular, if $S(G) \neq \emptyset$ and $W(G) \neq \emptyset$,

$$\text{ehr}_{str}^{(-q)} * \text{ehr}_{str}^{(q)}(G) = \text{ehr}_{str}^{(q)} * \text{ehr}_{str}^{(-q)}(G) = 0 = \varepsilon_\Delta(G).$$

Proof. Indeed,

$$\text{ehr}_{str}^{(q)} * \text{ehr}^{(q')}(G) = \sum q^{|I_1|} q'^{|I_2|},$$

where the sum is over all partitions $V(G) = I_1 \sqcup I_2$ such that if $x \xrightarrow{G} y$ in $V(G)$, then $(x, y) \in (I_1 \times I_2) \cup I_2^2$, that is to say such that $I_1 \subseteq S(G)$. Hence,

$$\begin{aligned} \text{ehr}_{str}^{(q)} * \text{ehr}^{(q')}(G) &= \sum_{I_1 \subseteq S(G)} q^{|I_1|} q'^{|V(G) \setminus I_1|} \\ &= q^{|V(G) \setminus S(G)|} \sum_{I_1 \subseteq S(G)} q^{|I_1|} q'^{|S(G) \setminus I_1|} \\ &= q^{|V(G) \setminus S(G)|} (q + q')^{|S(G)|}. \end{aligned}$$

The proof is similar for $\text{ehr}_{str}^{(q)} * \text{ehr}^{(q')}(G)$, replacing sources by wells. \square

Corollary 5.6. *The inverse (for the convolution product $*$) of the restriction of $\text{ehr}^{(q)}$ to $\mathcal{F}[\mathbf{G}_{ac}]$ is the restriction of $\text{ehr}_{str}^{(-q)}$.*

Proof. Let G be an acyclic mixed graph. Obviously, if $G = 1$, then

$$\text{ehr}_{str}^{(-q)} * \text{ehr}^{(q)}(G)1 = \varepsilon_{\Delta}(G).$$

Otherwise, as G is acyclic, then $S(G) \neq \emptyset$ and $W(G) \neq \emptyset$. We can conclude with Proposition 5.5. \square

Let us now prove the duality principle for Ehrhart polynomials:

Corollary 5.7. *Let G be an acyclic mixed graph. Then*

$$\text{Ehr}_{str}(G)(-X) = (-1)^{|V(G)|} \text{Ehr}(G)(X).$$

Proof. We denote by $S_{\mathbf{G}_{ac}}$ the antipode of $(\mathcal{F}[\mathbf{G}_{ac}], m, \Delta)$ and by S the antipode of $(\mathbb{K}[X], m, \Delta)$. In particular, for any $P \in \mathbb{K}[X]$, $S(P(X)) = P(-X)$. As $\text{Ehr} : (\mathcal{F}[\mathbf{G}_{ac}], m, \Delta) \longrightarrow (\mathbb{K}[X], m, \Delta)$ is a Hopf algebra morphism,

$$\text{Ehr}_{str}(G)(-X) = S \circ \text{Ehr}_{str}(G) = \text{Ehr}_{str} \circ S_{\mathbf{G}_{ac}}(G).$$

Therefore, by Corollary 5.6,

$$\begin{aligned} \text{Ehr}_{str}(G)(-1) &= S \circ \text{Ehr}_{str}(G)(1) \\ &= \text{Ehr}_{str} \circ S_{\mathbf{G}_{ac}}(G)(1) \\ &= \text{ehr}_{str} \circ S_{\mathbf{G}_{ac}}(G) \\ &= \text{ehr}_{str}^{*-1}(G) \\ &= \text{ehr}^{(-1)}(G). \end{aligned}$$

This implies that $\text{Ehr}_{str} \circ S$ is the Hopf algebra morphism $P_{chr_S} \leftarrow \text{ehr}^{(-1)}$:

$$\begin{aligned} \text{Ehr}_{str} \circ S(G) &= (P_{chr_S} \otimes \text{ehr}^{(-1)}) \circ \delta(G) \\ &= (-1)^{|V(G)|} (P_{chr_S} \otimes \text{ehr}) \circ \delta(G) \\ &= (-1)^{|V(G)|} \text{Ehr}(G)(X). \end{aligned} \quad \square$$

Corollary 5.8. *Let G be an acyclic oriented graph. Then*

$$\begin{aligned} \text{Ehr}_{str}(G) &= P_{chr_S}(G), & P_{chr_S}(G)(-1) &= (-1)^{|V(G)|}, \\ \text{Ehr}(G) &= P_{chr_W}(G), & P_{chr_W}(G)(-1) &= (-1)^{|V(G)|} \varepsilon_{\delta}(G). \end{aligned}$$

Proof. We work in $\mathcal{F}[\mathbf{G}]$. For any oriented acyclic graph H ,

$$\text{ehr}_{str}(H) = \epsilon_\delta(H) = \begin{cases} 1 & \text{if } A(H) = \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

Hence, $\text{Ehr}_{str}(G) = P_{chr_S}(G)$: the restriction of Ehr to $\mathcal{F}[\mathbf{G}_{aco}]$ is P_{chr_S} . Moreover,

$$\text{Ehr}_{str}(G)(-1) = (-1)^{|V(G)|} \text{Ehr}(G)(1) = (-1)^{|V(G)|} \text{ehr}(G) = (-1)^{|V(G)|}.$$

For any oriented acyclic graph H , $\text{ehr}(H) = 1 = \lambda_W(H)$, so $\text{Ehr}(G) = P_{chr_W}(G)$. Moreover,

$$\text{Ehr}(G)(-1) = (-1)^{|V(G)|} \text{Ehr}_{str}(G)(1) = (-1)^{|V(G)|} \text{ehr}_{str}(G) = (-1)^{|V(G)|} \epsilon_\delta(G). \quad \square$$

From Corollary 5.3, we obtain another proof of the following result, obtained in a different way in [5, Theorem 3]:

Corollary 5.9. *Let G be a mixed graph. Then*

$$P_{chr_S}(G)(-1) = (-1)^{|V(G)|} |\mathcal{O}_{ac}(G)|.$$

Proof. Indeed,

$$P_{chr_S}(G)(-1) = \sum_{H \in \mathcal{O}_{ac}(G)} P_{chr_S}(H)(-1) = \sum_{H \in \mathcal{O}_{ac}(G)} (-1)^{|V(H)|} = (-1)^{|V(G)|} |\mathcal{O}_{ac}(G)|. \quad \square$$

Remark 5.2. We recover the classical result on the chromatic polynomial when this corollary is applied to graphs [19].

From [10, Corollary 2.3]:

Corollary 5.10. *Denoting by S the antipode of $(\mathcal{F}[\mathbf{G}], m, \Delta)$, for any mixed graph G ,*

$$S(G) = \sum_{\sim \in \mathcal{E}^c[G]} (-1)^{\text{cl}(\sim)} |\mathcal{O}_{ac}(G/\sim)| (G|\sim).$$

6 Applications to characters on mixed graphs

6.1 Weak chromatic polynomial

Proposition 6.1. *Let G be a totally acyclic mixed graph. Then*

$$P_{chr_W}(G)(-1) = \begin{cases} 0 & \text{if } A(G) \neq \emptyset, \\ (-1)^{|V(G)|} |\mathcal{O}_{ac}(G)| & \text{otherwise.} \end{cases}$$

Proof. If $A(G) = \emptyset$, then $P_{chr_W}(G) = P_{chr_S}(G)$, and the result comes from Corollary 5.9. Let us assume that $A(G) \neq \emptyset$. We proceed by induction on $|E(G)|$. If $E(G) = \emptyset$, then by definition, $P_{chr_W}(G) = \text{Ehr}(G)$. By the duality principle for Ehrhart polynomials (Corollary 5.7),

$$P_{chr_W}(G)(-1) = (-1)^{|V(G)|} \text{Ehr}_{str}(G)(1) = 0,$$

as $A(G) \neq \emptyset$. Let us assume the result for all acyclic graph H such that $|E(G)| > |E(H)|$ and $A(H) \neq \emptyset$. Let e be an edge of G . We denote respectively by G/e and by $G \setminus e$ the mixed graph obtained from G by contraction of the edge e respectively by deleting the edge e . From [5, Proposition 6],

$$P_{chr_W}(G)(-1) = P_{chr_W}(G \setminus e)(-1) - P_{chr_W}(G/e)(-1).$$

Moreover, $G \setminus e$ and G/e are mixed graph with at least one arc and strictly less edges than G . Moreover, as G is totally acyclic, G/e and $G \setminus e$ are acyclic: we deduce that $P_{chr_W}(G \setminus e)(-1) = P_{chr_W}(G/e)(-1) = 0$. Hence, $P_{chr_W}(G)(-1) = 0$. \square

Remark 6.1. If G is not totally acyclic, no interpretation of $P_{chr_W}(G)(-1)$, and even of its sign, is known. For example, if G_n is the graph of Example 4.1, then $P_{chr_W}(G_n)(-1) = 1$ for any $n \geq 2$.

We recover the interpretation of [4] of the values of the weak chromatic polynomial at negative values:

Corollary 6.2. *Let G be a totally acyclic mixed graph and $k \in \mathbb{N}$. Then $(-1)^{|V(G)|} P_{chr_W}(G)(-k)$ is the number of pairs (H, f) such that:*

- H is an acyclic orientation of G .
- f is a k -coloring of G compatible with H , that is,

$$\forall x, y \in V(G), \quad \begin{cases} x \xrightarrow{G} y & \implies f(x) < f(y), \\ x \xrightarrow{H} y & \implies f(x) \leq f(y). \end{cases}$$

Proof. By compatibility of P_{chr_W} with the coproduct Δ ,

$$P_{chr_W}(G)(-k) = \sum_{\substack{f: V(G) \rightarrow [k], \\ x \xrightarrow{G} y \implies f(x) \leq f(y)}} P_{chr_W}(G|_{f^{-1}(1)})(-1) \dots P_{chr_W}(G|_{f^{-1}(k)})(-1).$$

By the Proposition 6.1, if $G|_{f^{-1}(i)}$ has an arc, then $P_{chr_W}(G|_{f^{-1}(i)})(-1) = 0$. Therefore,

$$\begin{aligned} P_{chr_W}(G)(-k) &= \sum_{\substack{f: V(G) \rightarrow [k], \\ x \xrightarrow{G} y \implies f(x) < f(y)}} P_{chr_W}(G|_{f^{-1}(1)})(-1) \dots P_{chr_W}(G|_{f^{-1}(k)})(-1) \\ &= \sum_{\substack{f: V(G) \rightarrow [k], \\ x \xrightarrow{G} y \implies f(x) < f(y)}} (-1)^{|f^{-1}(1)| + \dots + |f^{-1}(k)|} \prod_{i=1}^k |\mathcal{O}_{ac}(G|_{f^{-1}(i)})| \\ &= (-1)^{|V(G)|} \sum_{\substack{f: V(G) \rightarrow [k], \\ x \xrightarrow{G} y \implies f(x) < f(y)}} \prod_{i=1}^k |\mathcal{O}_{ac}(G|_{f^{-1}(i)})|. \end{aligned}$$

We consider

$$\begin{aligned} A_k &= \{(f, H_1, \dots, H_k) \mid f: V(G) \rightarrow [k], H_i \in \mathcal{O}_{ac}(G|_{f^{-1}(i)})\}, \\ B_k &= \{(H, f) \mid H \in \mathcal{O}_{ac}(G), f: V(G) \rightarrow [k] \text{ compatible with } H\}. \end{aligned}$$

The map $\phi: B_k \rightarrow A_k$ which send (H, f) to $(f, H|_{f^{-1}(1)}, \dots, H|_{f^{-1}(k)})$ is well-defined. If $\phi(H, f) = \phi(H', f')$, then $f = f'$. Moreover, if $\{x, y\} \in E(G)$:

- If $f(x) = f(y) = i$, then $\{x, y\}$ is oriented in the same way in $H|_{f^{-1}(i)}$ and in $H'|_{f^{-1}(i)}$, as these oriented graphs are equal. So $\{x, y\}$ is oriented in the same way in H and in H' .
- If $f(x) < f(y)$, as f is compatible with H and in H' , then $(x, y) \in E(H)$ and $(x, y) \in E(H')$.
- If $f(x) > f(y)$, as f is compatible with H and in H' , then $(y, x) \in E(H)$ and $(y, x) \in E(H')$.

Therefore, $H = H'$: ϕ is injective. Let $(f, H_1, \dots, H_k) \in A_k$. We define an orientation of G by the following: if $\{x, y\} \in E(G)$,

- if $f(x) = f(y) = i$, we keep the orientation of this edge in H_i .
- If $f(x) < f(y)$, we orient this edge in (x, y) in H .

- If $f(x) > f(y)$, we orient this edge in (y, x) in H .

By construction, for any $i \in [k]$, $H_{|f^{-1}(i)} = H_i$. Moreover, f is compatible with H , by construction. Consequently, f is constant on any cycle of H . As the oriented graphs H_i are acyclic, H is acyclic. We obtain that $(H, f) \in B_k$ and $\phi(H, f) = (f, H_1, \dots, H_k)$. Finally,

$$P_{chr_W}(G)(-k) = (-1)^{|V(G)|} |A_k| = (-1)^{|V(G)|} |B_k|. \quad \square$$

6.2 The character ν_W

Recall that ν_W is the inverse of λ_W (defined in (3)) for the convolution \star associated to δ .

Proposition 6.3. 1. For any simple graph G with at least one edge, $\nu_W(G) = 0$.

2. For any oriented graph G , $\nu_W(G) = (-1)^{|V(G)| + cc(G)}$, where $cc(G)$ is the number of connected components of G .

Proof. 1. We denote by λ_W^s the restriction of λ_W to $\mathcal{F}[\mathbf{G}_s]$. For any simple graph G , $\lambda_W(G) = \epsilon_\delta(G)$, so $\lambda_W^s = \epsilon_{\delta|_{\mathcal{F}_V[\mathbf{G}_s]}}$. As $\mathcal{F}[\mathbf{G}_s]$ is a double subbialgebra of $\mathcal{F}[\mathbf{G}]$,

$$(\lambda_W^s)^{\star^{-1}} = (\epsilon_{\delta|_{\mathcal{F}[\mathbf{G}_s]}})^{\star^{-1}} = (\epsilon_\delta^{\star^{-1}})|_{\mathcal{F}[\mathbf{G}_s]} = \epsilon_{\delta|_{\mathcal{F}[\mathbf{G}_s]}}.$$

2. Let λ be a character of $\mathcal{F}[\mathbf{G}_o]$, such that $\lambda(\bullet) \neq 0$. Then λ is invertible for the convolution product \star associated to δ : its inverse is denoted by μ . Denoting by S the antipode of $(\mathcal{F}[\mathbf{G}_o], m, \Delta)$, let us prove that $\lambda \circ S$ is invertible for \star and that its inverse is $\mu \star (\epsilon_\delta \circ S)$. Firstly, $\lambda \circ S(\bullet) = -\lambda(\bullet) \neq 0$, so $\lambda \circ S$ is invertible. Moreover,

$$\begin{aligned} (\lambda \circ S) \star \mu \star (\epsilon_\delta \circ S) &= (\lambda \otimes \mu \circ \epsilon_\delta) \circ (S \otimes \text{Id} \otimes S) \circ (\delta \otimes \text{Id}) \circ \delta \\ &= (\lambda \otimes \mu \circ \epsilon_\delta) \circ (\text{Id} \otimes \text{Id} \otimes S) \circ (\delta \otimes \text{Id}) \circ \delta \circ S \\ &= (\lambda \otimes \mu \circ \epsilon_\delta) \circ (\delta \otimes \text{Id}) \circ (\text{Id} \otimes S) \circ \delta \circ S \\ &= (\lambda \star \mu \otimes \epsilon_\delta) \circ (\text{Id} \otimes S) \circ \delta \circ S \\ &= \epsilon_\delta \otimes \epsilon_\delta \circ (\text{Id} \otimes S) \circ \delta \circ S \\ &= \epsilon_\delta \circ S^2 \\ &= \epsilon_\delta. \end{aligned}$$

We used for the second equality that $(S \otimes \text{Id}) \circ \delta = \delta \circ S$ (see [10, Proposition 2.1]) and, for the last equality, that $S^2 = \text{Id}$, as $\mathcal{F}[\mathbf{G}]$ is commutative. So $\mu \star (\epsilon_\delta \circ S) = (\lambda \circ S)^{\star^{-1}}$.

In the particular case were $\lambda = \epsilon_\delta$, then $\mu = \epsilon_\delta$ and we obtain that $(\epsilon_\delta \circ S)^{\star^{-1}} = \epsilon_\delta \circ S$.

Let G be an oriented graph. By definition of the weak Ehrhart polynomial and by the duality principle for Ehrhart polynomial (Corollary 5.7),

$$\begin{aligned} \lambda_W(G) &= 1 \\ &= \text{Ehr}_W(G)(1) \\ &= (-1)^{|V(G)|} \text{Ehr}(G)(-1) \\ &= (-1)^{|V(G)|} S \circ \text{Ehr}(G)(1) \\ &= (-1)^{|V(G)|} \text{Ehr}(S(G))(1) \\ &= (-1)^{|V(G)|} \epsilon_\delta \circ S(G). \end{aligned}$$

We now consider the three characters of $\mathcal{F}[\mathbf{G}_o]$ defined on any oriented graph G by

$$\lambda(G) = \epsilon_\delta \circ S(G), \quad \mu(G) = (-1)^{|V(G)|} \lambda(G), \quad \nu(G) = (-1)^{cc(G)} \lambda(G).$$

We already proved that $\lambda \star \lambda = \epsilon_\delta$. For any graph G ,

$$\begin{aligned} \mu \star \nu(G) &= \sum_{\sim \in \mathcal{E}^c[G]} (-1)^{|V(G/\sim)| + \text{cc}(G/\sim)} \lambda(G/\sim) \lambda(G|\sim) \\ &= \sum_{\sim \in \mathcal{E}^c[G]} (-1)^{2\text{cl}(\sim)} \lambda(G/\sim) \lambda(G|\sim) \\ &= \sum_{\sim \in \mathcal{E}^c[G]} \lambda(G/\sim) \lambda(G|\sim) \\ &= \lambda \star \lambda(G) \\ &= \epsilon_\delta(G). \end{aligned}$$

Therefore, $\nu = \mu^{\star -1}$. As $\mu = \lambda_W$, we obtain that $\nu = \nu_W$ and, for any oriented graph G ,

$$\nu_W(G) = (-1)^{\text{cc}(G)} \lambda(G) = (-1)^{\text{cc}(G) + |V(G)|} \lambda_W(G) = (-1)^{\text{cc}(G) + |V(G)|}. \quad \square$$

No interpretation of $\nu_W(G)$ is known in general. For example:

Proposition 6.4. *Let G_n be the mixed graph of Example 4.1, with the convention $G_2 = \bullet \longrightarrow \bullet$. For any $n \geq 2$, $\nu_W(G_n) = (-1)^{n-1}(n-1)$.*

Proof. For any $n \geq 2$,

$$\nu_W * \lambda_W(G_n) = \sum_{\sim \in \mathcal{E}^c[G_n]} \nu_W(G/\sim) \lambda_W(G|\sim) = \epsilon_\delta(G_n) = 0.$$

By definition of λ_W , for any $\sim \in \mathcal{E}^c[G_n]$, $\lambda_W(G|\sim) = 0$ if, and only if, $1 \sim n$. Therefore, the contributing terms corresponds to the equivalences which classes are intervals of $[n]$, at the exception of the one with only one class. For such an equivalence \sim , G/\sim is isomorphic to $G_{\text{cl}(\sim)}$. We obtain that if $n \geq 3$,

$$\sum_{k=2}^n \sum_{\substack{i_1 + \dots + i_k = n, \\ i_1, \dots, i_k \geq 1}} \nu_W(G_k) = 0.$$

A direct computation shows that $\lambda_W(G_2) = -1$. Summing, we obtain in the ring of formal series $\mathbb{Q}[[X]]$ that

$$\sum_{k=2}^{\infty} \sum_{\substack{i_1 + \dots + i_k = n, \\ i_1, \dots, i_k \geq 1}} \nu_W(G_k) X^{i_1 + \dots + i_n} = \sum_{k=2}^{\infty} \nu_W(G_k) \left(\frac{X}{1-X} \right)^k = -X^2.$$

Substituting $\frac{X}{1+X}$ to X , we obtain

$$\sum_{n=2}^{\infty} \nu_W(G_n) X^n = -\frac{X^2}{(1+X)^2} = \sum_{n=2}^{\infty} (-1)^{n+1}(n-1) X^n. \quad \square$$

7 Appendix: number of mixed graphs up to isomorphism

Proposition 7.1. *Let $N \in \mathbb{N}$, nonzero. For any $n \in N$, we denote respectively by $G_m(N, n)$, $G_o(N, n)$ and $G_s(N, n)$ the number of isomorphism classes of mixed, oriented, simple graphs with n vertices decorated by elements of $[N]$. Then, for $t \in \{m, o, s\}$,*

$$G_t(N, n) = \sum_{\substack{c_1, \dots, c_n \geq 0, \\ 1c_1 + \dots + nc_n = n}} \frac{N^{c_1 + \dots + c_n}}{1^{c_1} \dots n^{c_n} c_1! \dots c_n!} A_t^{1 \leq k < l \leq n} \sum_{1 \leq k < l \leq n} c_k c_l (k \wedge l) + \sum_{1 \leq k \leq n} \frac{c_k (kc_k - 1)}{2} \left(\frac{B_t}{\sqrt{A_t}} \right)^{1 \leq k \leq n/2} \sum c_{2k},$$

with

t	A_t	B_t
m	5	3
o	4	2
s	2	2

Proof. We put

$$\begin{aligned} F_m &= \{(0, 0), (0, \rightarrow), (\rightarrow, 0), (\rightarrow, \rightarrow), (-, -)\}, \\ F_o &= \{(0, 0), (0, \rightarrow), (\rightarrow, 0), (\rightarrow, \rightarrow)\}, \\ F_s &= \{(0, 0), (-, -)\}. \end{aligned}$$

These sets give the possibilities of edges or arcs between two vertices of a graph of the considered graphs. For any $t \in \{m, o, s\}$, F_t is given an involution ι , given by the usual flip. Note that A_t is the cardinal of F_t and that B_t is the number of elements of F_t invariant under ι . Then, a graph G of type t , which set of vertices is $[n]$, together with a map from $V(G)$ to $[N]$, can be seen as a pair (d_V, d_E) , where $d_V : [n] \rightarrow [N]$ is a map and $d_E = \mathcal{P}_2([n]) \rightarrow F_t$ is a map which makes explicit the edge and arc situation between two vertices i, j , with $i < j$. The symmetric group \mathfrak{S}_n acts on these pairs (d_V, d_E) , in the following sense: for any $\sigma \in \mathfrak{S}_n$, putting $\sigma \cdot (d_V, d_E) = (\sigma \cdot d_V, \sigma \cdot d_E)$,

- $\sigma \cdot d_V = d_V \circ \sigma^{-1}$.
- For any $i, j \in [n]$, with $i < j$,

$$\sigma \cdot d_E(\{i, j\}) = \begin{cases} d_E(\{\sigma^{-1}(i), \sigma^{-1}(j)\}) & \text{if } \sigma^{-1}(i) < \sigma^{-1}(j), \\ \iota \circ d_E(\{\sigma^{-1}(i), \sigma^{-1}(j)\}) & \text{if } \sigma^{-1}(i) > \sigma^{-1}(j). \end{cases}$$

The number of isomorphism classes $G_t(N, n)$ we are looking for is the number of orbits of this action. Using Burnside's formula,

$$G_t(N, n) = \frac{1}{n!} \text{Fix}(\sigma),$$

where $\text{Fix}(\sigma)$ is the number of pairs (d_V, d_E) invariant under the action of σ .

Let $\sigma \in \mathfrak{S}_n$. Then $\sigma \cdot (d_V, d_E) = (d_V, d_E)$ if, and only if:

- For any i, j in the same σ -orbit, $d_V(i) = d_V(j)$. The number of σ -orbits is denoted by $n(\sigma)$.
- For any orbit of the action of σ on pairs $\{i, j\}$, with $i < j$, such that there exists $k \in \mathbb{N}$ with $\sigma^k(i) = j$ and $\sigma^k(j) = i$, d_E is constant on this orbit and its values is invariant under ι . Such an orbit will be called an orbit of type B . The number of orbits of type B is denoted by $n_B(\sigma)$.
- For any orbit of the action of σ on pairs $\{i, j\}$, with $i < j$, such that for any $k \in \mathbb{N}$, $(\sigma^k(i), \sigma^k(j)) \neq (j, i)$, d_E is constant on this orbit. Such an orbit will be called an orbit of type A . The number of orbits of type A is denoted by $n_A(\sigma)$.

Consequently,

$$\text{Fix}(\sigma) = N^{n(\sigma)} A^{n_A(\sigma)} B^{n_B(\sigma)}.$$

Let us decompose σ into cycles with disjoint supports. For any $i \in [N]$, the number of cycles of length i in this decomposition is denoted by c_i . Then $1c_1 + \dots + nc_n = n$. Then $n(\sigma) = c_1 + \dots + c_n$.

Let $\{i, j\}$ be a pair such that i, j belong to two different cycles of σ . As $\sigma^k(i)$ belongs to the cycle containing i for any $k \in \mathbb{N}$, $\sigma^k(i) \neq j$. So the orbit of $\{i, j\}$ is of type A. Moreover, denoting by k and l the lengths of the cycle containing i and j , this orbit is of cardinality $k \vee l$. Hence, these two cycles give $\frac{kl}{k \vee l} = k \wedge l$ orbits of type A. Summing over all possible choices of disjoint cycles of σ , we obtain

$$\sum_{1 \leq k < l \leq n} c_k c_l (k \wedge l) + \sum_{1 \leq k \leq n} \frac{c_k (c_k - 1)}{2} k$$

orbits of type A.

Let $\{i, j\}$ be a pair such that i, j belong to the same cycle of σ . The length of this cycle is denoted by l : for a given cycle, this gives $\frac{l(l-1)}{2}$ pairs. Moreover, for any $k \in \mathbb{N}$,

$$\{\sigma^k(i), \sigma^k(j)\} = \{i, j\} \iff \begin{cases} \sigma^k(i) = i \\ \sigma^k(j) = j \end{cases} \quad \text{or} \quad \begin{cases} \sigma^k(i) = j \\ \sigma^k(j) = i. \end{cases}$$

Moreover, $\sigma^k(i) = i$ if and only if, k is a multiple of l . If $\sigma^k(i) = j$ and $\sigma^k(j) = i$, then $\sigma^{2k}(i) = i$, so $2k$ is a multiple of l . We obtain two cases:

- If l is odd, then If $\sigma^k(i) = j$ and $\sigma^k(j) = i$, $l \mid 2k$ and, as l is odd, $l \mid k$. In other words,

$$\{\sigma^k(i), \sigma^k(j)\} = \{i, j\} \iff l \mid k.$$

As a consequence, all the orbits contained in this cycle are of type A, and of cardinality l . This gives $\frac{l-1}{2}$ orbits of type A.

- If l is even, we obtain an orbit of type B: if the cycle is (i_1, \dots, i_{2p}) , this orbit is

$$\{\{i_1, i_{p+1}\}, \dots, \{i_p, i_{2p}\}\}.$$

It contains $p = \frac{l}{2}$ elements. All the other orbits included in this cycle have cardinality l , so we obtain $\frac{1}{l} \left(\frac{l(l-1)}{2} - \frac{l}{2} \right) = \frac{l-2}{2}$ orbits of type A.

Summing over all possible choices of cycles, we obtain $c_2 + c_4 + \dots$ orbits of type B and

$$\sum_{\substack{1 \leq k \leq n, \\ k \text{ odd}}} c_k \frac{k-1}{2} + \sum_{\substack{1 \leq k \leq n, \\ k \text{ even}}} c_k \frac{k-2}{2}.$$

Finally,

$$\begin{aligned} n_B(\sigma) &= \sum_{1 \leq k \leq n/2} c_{2k}, \\ n_A(\sigma) &= \sum_{1 \leq k < l \leq n} c_k c_l (k \wedge l) + \sum_{1 \leq k \leq n} \frac{c_k (c_k - 1)}{2} k + \sum_{\substack{1 \leq k \leq n, \\ k \text{ odd}}} c_k \frac{k-1}{2} + \sum_{\substack{1 \leq k \leq n, \\ k \text{ even}}} c_k \frac{k-2}{2} \\ &= \sum_{1 \leq k < l \leq n} c_k c_l (k \wedge l) + \sum_{1 \leq k \leq n} \frac{(k c_k - 1) c_k}{2} - \frac{1}{2} \sum_{1 \leq k \leq n/2} c_{2k}, \end{aligned}$$

which finally gives the announced formula, as there are $1^{c_1} \dots 1^{c_n} c_1! \dots c_n!$ permutations which decomposition in cycles with disjoint supports is given by c_i cycles of length i for any $i \in [n]$. \square

Example 7.1. Note for a given n , $G_t(N, n)$ is a polynomial in N , A_t and B_t .

$$G_t(N, 1) = N,$$

$$G_t(N, 2) = \frac{(A_t N + B_t)N}{2},$$

$$G_t(N, 3) = \frac{(A_t^2 N^2 + 3B_t N + 2)A_t N}{6},$$

$$G_t(N, 4) = \frac{(A_t^5 N^3 + 6A_t^2 B_t N^2 + 8A_t N + 6B_t)A_t N}{24}.$$

These array gives the first values of $G_t(N, n)$. For mixed graphs,

$n \setminus N$	1	2	3	4	5	6
1	1	2	3	4	5	6
2	4	13	27	46	70	99
3	30	200	635	1460	2800	4780
4	785	11320	55605	173265	419550	865335
5	86130	2673260	20113890	84385520	256856275	638050530
6	43053850	2733053500	31051529575	174249075200	664212533500	1982349763225

The second row is entry A147875 of the OEIS [17]. For oriented graphs,

$n \setminus N$	1	2	3	4	5	6
1	1	2	3	4	5	6
2	3	10	21	36	55	78
3	16	104	328	752	1440	2456
4	218	3044	14814	45960	111010	228588
5	9608	291968	2183400	9133760	27755016	68869824
6	1540944	96928992	1098209328	6154473664	23441457680	69924880288

The second row is entry A014105 of the OEIS, whereas the first and second columns are entries A000273 and A000595. For simple graphs,

$n \setminus N$	1	2	3	4	5	6
1	1	2	3	4	5	6
2	2	6	12	20	30	42
3	4	20	56	120	220	364
4	11	90	357	996	2255	4446
5	34	544	3258	12208	34754	82608
6	156	5096	47324	241520	871580	2510424

The second, third and fourth rows are respectively entries A002378, A002492 and A199394 of the OEIS, whereas the first and second columns are entries A000088 and A000666.

References

- [1] Marcelo Aguiar and Swapneel Mahajan, *Monoidal functors, species and Hopf algebras*, CRM Monograph Series, vol. 29, American Mathematical Society, Providence, RI, 2010, With forewords by Kenneth Brown and Stephen Chase and André Joyal.
- [2] P. Alexandroff, *Diskrete Räume.*, Rec. Math. Moscou, n. Ser. **2** (1937), 501–519 (German).
- [3] Octavio Arizmendi and Adrian Celestino, *Monotone cumulant-moment formula and Schröder trees*, SIGMA, Symmetry Integrability Geom. Methods Appl. **18** (2022), paper 073, 22 (English).

- [4] Matthias Beck, Daniel Blado, Joseph Crawford, Taïna Jean-Louis, and Michael Young, *On weak chromatic polynomials of mixed graphs*, *Graphs Comb.* **31** (2015), no. 1, 91–98 (English).
- [5] Matthias Beck, Tristram Bogart, and Tu Pham, *Enumeration of Golomb rulers and acyclic orientations of mixed graphs*, *Electron. J. Comb.* **19** (2012), no. 3, research paper p42, 13 (English).
- [6] Damien Calaque, Kurusch Ebrahimi-Fard, and Dominique Manchon, *Two interacting Hopf algebras of trees: a Hopf-algebraic approach to composition and substitution of B-series*, *Adv. in Appl. Math.* **47** (2011), no. 2, 282–308.
- [7] Loïc Foissy, *Commutative and non-commutative bialgebras of quasi-posets and applications to Ehrhart polynomials*, *Adv. Pure Appl. Math.* **10** (2019), no. 1, 27–63.
- [8] ———, *Twisted bialgebras, cofreeness and cointeraction*, arXiv:1905.10199, 2019.
- [9] ———, *Chromatic polynomials and bialgebras of graphs*, *Int. Electron. J. Algebra* **30** (2021), 116–167.
- [10] ———, *Bialgebras in cointeraction, the antipode and the eulerian idempotent*, arXiv:2201.11974, 2022.
- [11] ———, *Contractions and extractions on twisted bialgebras and coloured Fock functors*, arXiv:2301.09447, 2023.
- [12] Pierre Hansen, Julio Kuplinsky, and Dominique de Werra, *Mixed graph colorings*, *Math. Methods Oper. Res.* **45** (1997), no. 1, 145–160 (English).
- [13] André Joyal, *Une théorie combinatoire des séries formelles*, *Adv. in Math.* **42** (1981), no. 1, 1–82.
- [14] ———, *Foncteurs analytiques et espèces de structures*, *Combinatoire énumérative (Montreal, Que., 1985/Quebec, Que., 1985)*, *Lecture Notes in Math.*, vol. 1234, Springer, Berlin, 1986, pp. 126–159.
- [15] Jean-Louis Loday and María Ronco, *On the structure of cofree Hopf algebras*, *J. Reine Angew. Math.* **592** (2006), 123–155.
- [16] A. Murua, *The Hopf algebra of rooted trees, free Lie algebras, and Lie series*, *Found. Comput. Math.* **6** (2006), no. 4, 387–426 (English).
- [17] N. J. A. Sloane, *The on-line encyclopedia of integer sequences*, <https://oeis.org/>.
- [18] Yuri N. Sotskov, Vjacheslav S. Tanaev, and Frank Werner, *Scheduling problems and mixed graphs colorings*, *Optimization* **51** (2002), no. 3, 597–624.
- [19] Richard P. Stanley, *Acyclic orientations of graphs*, *Discrete Math.* **5** (1973), 171–178.