

FULLY COMPLEMENTARY HIGHER DIMENSIONAL PARTITIONS

FLORIAN SCHREIER-AIGNER

ABSTRACT. We introduce a symmetry class for higher dimensional partitions - *fully complementary higher dimensional partitions* (FCPs) - and prove a formula for their generating function. By studying symmetry classes of FCPs in dimension 2, we define variations of the classical symmetry classes for plane partitions. As a by-product we obtain conjectures for three new symmetry classes of plane partitions and prove that another new symmetry class, namely *quasi transpose complementary plane partitions* are equinumerous to symmetric plane partitions.

1. INTRODUCTION

A *plane partition* π is an array $(\pi_{i,j})$ of non-negative integers with all but finitely many entries equal to 0, which is weakly decreasing along rows and columns, i.e., $\pi_{i,j} \geq \pi_{i+1,j}$ and $\pi_{i,j} \geq \pi_{i,j+1}$; see Figure 1 (left) for an example. MacMahon [12] introduced them at the end of the 19th century as two dimensional generalisations of ordinary partitions and proved in [14] two enumeration results: He showed that the generating function of plane partitions is given by

$$\sum_{\pi} q^{|\pi|} = \prod_{i \geq 1} \frac{1}{(1 - q^i)^i}, \quad (1.1)$$

where the sum is over all plane partitions and $|\pi|$ is defined as the sum of the entries of π . A plane partition π is said to be *contained in an (a, b, c) -box* if the entries of π are at most c and $\pi_{i,j} \neq 0$ implies $i \leq a$ and $j \leq b$. The plane partition in Figure 1 is contained in a $(3, 4, 4)$ -box or any box of larger size. MacMahon showed that the weighted enumeration of plane partitions inside an (a, b, c) -box is given by

$$\sum_{\pi} q^{|\pi|} = \prod_{i=1}^a \prod_{j=1}^b \prod_{k=1}^c \frac{1 - q^{i+j+k-1}}{1 - q^{i+j+k-2}}, \quad (1.2)$$

where the sum is over all plane partitions contained in an (a, b, c) -box.

While plane partitions were already introduced at the end of the 19th century, they came into the focus of the combinatorics community mainly in the second half of the last century. One major goal was to prove the enumeration formulas for the ten symmetry classes of plane partitions. These classes are defined via combinations of the three operations *reflection*, *rotation* and *complementation* which are best defined by viewing plane partitions as lozenge tilings: First, we represent a plane partition π as stacks of unit cubes by placing $\pi_{i,j}$ unit cubes at the position (i, j) , see Figure 1 (middle). By further displaying the shape of the (a, b, c) -box in which we regard the plane partition in and forgetting the shading of the cubes, we obtain a lozenge tiling of a hexagon with side lengths a, b, c, a, b, c , see Figure 1 (right). Interestingly, this was first observed by David and Tomei [7] in 1989. The operation *reflection* is defined as vertical reflection of the lozenge tiling, *rotation* as rotation by 120 degrees and *complementation* as rotation by 180 degree. While MacMahon [13] already considered plane partitions invariant under reflection, the operation completion was first described by Mills, Robbins and Rumsey [15] in 1986. A systematic study of the

Florian Schreier-Aigner acknowledges the financial support from the Austrian Science Foundation FWF, grant J 4387.




FIGURE 1. A plane partition contained in a $(3, 4, 4)$ -box on the left, its graphical representation as stacks of unit cubes (middle) and the associated lozenge tiling (right).

ten symmetry classes which are defined through these operations was initiated by Stanley [18, 19] and finished in 2011 by Koutschan, Kauers and Zeilberger [9]. For a more detailed overview see [11].

Already in 1916, MacMahon [14] introduced a further generalisation of partitions to arbitrary dimension, namely *higher-dimensional partitions*. A *d -dimensional partition* π is an array (π_{i_1, \dots, i_d}) of non-negative integers with all but finitely many entries equal to 0, such that $\pi_{\mathbf{i}} \geq \pi_{\mathbf{i}+e_k}$ for all indices $\mathbf{i} = (i_1, \dots, i_d)$ and $1 \leq k \leq d$, where e_k denotes the k -th unit vector. We say that π is contained in an (n_1, \dots, n_{d+1}) -box if all entries are at most n_{d+1} and $\pi_{\mathbf{i}} > 0$ implies that $i_j \leq n_j$ for all $1 \leq j \leq d$. Contrary to dimension 1 (partitions) and dimension 2 (plane partitions), there are hardly any results known for higher-dimensional partitions in dimension 3 or higher. MacMahon conjectured a generating formula for each dimension d but it was disproved by Atkin, Bratley, Macdonald and McKay [2] in 1967; see also [8]. Only recently the first enumeration result for higher-dimensional partitions was presented by Amanov and Yeliussizov [1]. They were able to “correct” MacMahons formula and showed that

$$\sum_{\pi} t^{\text{cor}(\pi)} q^{|\pi|_{\text{ch}}} = \prod_{i \geq 1} (1 - tq^i)^{-\binom{i+d-2}{d-1}}, \quad (1.3)$$

where the sum is over all d -dimensional partitions, and cor and $|\cdot|_{\text{ch}}$ are certain statistics defined in [1, Section 4 and 5].

In this paper we introduce a new symmetry class for plane partitions, namely *quarter complementary plane partitions* (QCPPs), which can be generalised immediately to higher-dimensional partitions. Instead of presenting the definition for QCPPs (it follows from the corresponding definition for higher-dimensional partitions in Section 2.2) we aim to convey the geometric intuition of this symmetry class next.

Let $\pi = (\pi_{i,j})$ be a plane partition inside an (a, b, c) -box and define by $\pi' = (\pi'_{i,j})$ the “dual partition” of π inside the (a, b, c) -box. Geometrically, we think of the dual partition as stacks of unit cubes hanging from the ceiling of the (a, b, c) -box instead of standing at its floor, where the first stack is positioned at the corner with coordinates (a, b, c) instead of the corner with coordinates $(1, 1, 1)$, see Figure 2 (left) for a sketch. It is not difficult to see, compare for example with [11, Section 6], that π is *self-complementary* if π and π' fill the (a, b, c) -box without overlap when regarded as stacks of unit cubes. In the array perspective this means $\pi_{i,j} + \pi'_{i,j} = c$ for all $1 \leq i \leq a$ and $1 \leq j \leq b$. We can now generalise this idea. Let C be a set of corners of the (a, b, c) -box. We define a partition π to be *C -complementary* if we can fill the (a, b, c) -box without overlap by copies of π placed at the corners of C similar to before, i.e., if a copy of π is placed at a corner of the form $(*, *, c)$ then its stacks of unit cubes hang from the ceiling




FIGURE 2. The labels of the corners of an (a, b, c) -box (middle), the corners where a copy of π is placed for self-complementary plane partitions (left), and the corners where copies of π are placed for quarter complementary plane partitions (right). The colour and lengths of the arrows indicate the orientation of the copies.

of the box instead of standing on its floor. It is immediate that such a π can only exist if $|C| \in \{1, 2, 4, 8\}$. We can ignore the cases $|C| = 1$ and $|C| = 8$ since they are trivial.

If $|C| = 2$ and the two corners are on a common face of the box, it is not difficult to see that π is determined along the direction orthogonal to this face. Up to rotation there is only one configuration of the two corners such that they do not share a common face: $C = \{(1, 1, 1), (a, b, c)\}$. The C -complementary plane partition in this case are self-complementary plane partitions. Similarly, if two corners in C are the endpoints of an edge of the box, then π is determined along the direction of this edge. The only configuration of four corners for which no pair of them are endpoints of an edge is up to rotation $C = \{(1, 1, 1), (a, b, 1), (1, b, c), (a, 1, c)\}$, see Figure 2 (right) for a sketch. We call the C -complementary plane partitions in this case *quarter complementary plane partitions*.

In Section 2 we generalise this geometric approach to higher dimensions by using *higher dimensional Ferrers diagrams*. Translating the obtained criteria on Ferrers diagrams back to the array description for higher-dimensional partitions, we obtain in Lemma 2.1 the definition of *d -dimensional fully complementary partitions* (FCPs). In Proposition 2.2 we describe the recursive structure of FCPs which implies immediately our main result.

Theorem 1.1. *Let $\mathbf{x} = (x_1, \dots, x_{d+1})$, $\mathbf{n} = (n_1, \dots, n_{d+1}) \in \mathbb{N}_{>0}^{d+1}$ and denote by $\text{FCP}(\mathbf{n})$ the set of fully complementary partitions inside a $(2n_1, \dots, 2n_{d+1})$ -box. Then*

$$\sum_{\mathbf{n} \in \mathbb{N}^{d+1}} |\text{FCP}(\mathbf{n})| \mathbf{x}^{\mathbf{n}} = \frac{\prod_{i=1}^{d+1} x_i \left(\sum_{i=1}^{d+1} (x_i^{-1} + dx_i) - \sum_{1 \leq i, j \leq d+1} x_i x_j^{-1} \right)}{\left(1 - \sum_{i=1}^{d+1} x_i \right) \prod_{i=1}^{d+1} (1 - x_i)}. \quad (1.4)$$

The recursive structure of FCPs can be further used to construct a bijection between d -dimensional FCPs and lattice paths in the positive $d + 1$ -dimensional orthant starting from any integer point in the interior of its d -dimensional boundary.

In Section 3 we consider the “typical” symmetry classes for plane partitions restricted to 2-dimensional FCPs, i.e. quarter complementary plane partitions. It turns out that there exists at most one symmetric QCPP in an (a, b, c) -box and that a QCPP can be neither cyclically symmetric nor transpose-complementary. By introducing two variations, namely *quasi symmetric* and *quasi transpose-complementary*, we are able to show enumerative results for the corresponding symmetry classes and combinations thereof for QCPPs, see Proposition 3.1 to Proposition 3.4. It is an immediate question if the enumeration of plane partitions under these variations of symmetry classes have closed expressions. In Section 4 we consider these new symmetry classes and similar generalisations. We present

three conjectures, the corresponding data generated in computer experiments is given in Appendix A, and the proof of the next result.

Theorem 1.2. *A plane partition π inside an (n, n, c) -box is called **quasi transpose-complementary** if $\pi_{i,j} + \pi_{n+1-j, n+1-i} = c$ holds for all $1 \leq i, j \leq n$ with $i \neq n+1-j$. The number of quasi transpose-complementary plane partitions inside an (n, n, c) -box is equal to the number of symmetric plane partitions inside an (n, n, c) -box.*

We present different proofs of the above theorem and how it is related to known results on plane partitions, lozenge tilings and perfect matchings respectively. For odd c we relate quasi transpose-complementary plane partitions to transpose-complementary plane partitions and obtain immediately the following numerical connection between symmetric and transpose-complementary plane partitions which seems to be new

$$2^{n-1} \text{TCPP}(n, n, 2c) = \text{SPP}(n-1, n-1, 2c+1), \quad (1.5)$$

where $\text{TCPP}(n, n, 2c)$ denotes the number of transpose-complementary plane partitions inside an $(n, n, 2c)$ -box and $\text{SPP}(n, n, c)$ denotes the number of symmetric plane partitions inside an (n, n, c) -box. Since the proof of the above theorem is computational, it is still an open question if one can find a bijection between symmetric plane partitions and quasi transpose-complementary plane partitions.

2. FULLY COMPLEMENTARY PARTITIONS

2.1. Fully complementary Ferrers diagrams. A *d -dimensional Ferrers diagram* λ is a finite subset of $(\mathbb{N}_{>0})^{d+1}$ such that $(x_1, \dots, x_{d+1}) \in \lambda$ implies $(y_1, \dots, y_{d+1}) \in \lambda$ whenever $1 \leq y_i \leq x_i$ for all $1 \leq i \leq d+1$. Equivalently λ is an order ideal in the poset $(\mathbb{N}_{>0})^{d+1}$ where the order relation is component-wise the order relation of the integers. For a positive integer n , we define $[n] = \{1, \dots, n\}$. We say that λ is *contained* in an (n_1, \dots, n_{d+1}) -box for positive integers n_1, \dots, n_{d+1} , if λ is a subset of $[n_1] \times \dots \times [n_{d+1}]$. It is immediate that the map

$$\lambda \mapsto \pi(\lambda) = (|\{(i_1, \dots, i_d, k) \in \lambda\}|)_{i_1, \dots, i_d}$$

is a bijection between d -dimensional Ferrers diagrams and d -dimensional partitions which respects the property of being contained in an (n_1, \dots, n_{d+1}) -box. Therefore, we identify for the remainder of this paper a d -dimensional Ferrers diagram with the d -dimensional partition it is mapped to.

Let $\mathbf{n} = (n_1, \dots, n_{d+1})$ be a sequence of positive integers and $I \subseteq [d+1]$. We define the bijection $\rho_{I, \mathbf{n}}$ from $[2n_1] \times \dots \times [2n_{d+1}]$ onto itself as

$$\rho_{I, \mathbf{n}}(x_1, \dots, x_{d+1}) := \left(\begin{cases} x_i & i \notin I, \\ 2n_i + 1 - x_i & i \in I, \end{cases} \right)_{1 \leq i \leq d+1}.$$

If \mathbf{n} is clear from the context we will omit it and write ρ_I instead of $\rho_{I, \mathbf{n}}$. We can use the map ρ_I to rephrase the definition of *quarter complementary plane partitions*. Let λ be a 2-dimensional Ferrers diagram contained in a $(2n_1, 2n_2, 2n_3)$ -box. Then $\rho_{\{1,2\}}(\lambda)$ corresponds to the copy of λ placed in the corner $(2n_1, 2n_2, 1)$ of the box, $\rho_{\{1,3\}}(\lambda)$ to the copy of λ placed in the corner $(2n_1, 1, 2n_3)$ and $\rho_{\{2,3\}}(\lambda)$ to the copy placed in the corner $(1, 2n_2, 2n_3)$. Hence λ corresponds to a QCPP if $\lambda, \rho_{\{1,2\}}(\lambda), \rho_{\{1,3\}}(\lambda), \rho_{\{2,3\}}(\lambda)$ have pairwise empty intersection and their union is equal to $[2n_1] \times [2n_2] \times [2n_3]$. We extend this definition to any dimension d .

Definition 2.1. Let n_1, \dots, n_{d+1} be positive integers. A d -dimensional Ferrers diagram λ is called *fully complementary* inside a $(2n_1, \dots, 2n_{d+1})$ -box if for all pairs of subsets

$I, J \subseteq [d+1]$ of even size holds $\rho_I(\lambda) \cap \rho_J(\lambda) = \emptyset$ and

$$\bigcup_{\substack{I \subseteq [d+1] \\ |I| \text{ even}}} \rho_I(\lambda) = [2n_1] \times \dots \times [2n_{d+1}].$$

It is easy to see, that $\rho_I(\lambda) \cap \rho_J(\lambda) = \emptyset$ holds for all subsets $I, J \subseteq [d+1]$ of even size exactly if $\lambda \cap \rho_I(\lambda) = \emptyset$ holds for all subsets $I \subseteq [d+1]$ of even size.

The 3-dimensional Ferrers diagram $\lambda = \{(1, 1, 1, 1), (2, 1, 1, 1)\}$ is fully complementary inside a $(2, 2, 2, 2)$ -box. This can be seen easily by calculating the according images under ρ_I for even sized I which are given by

$$\begin{aligned} \rho_{\{1,2\}}(\lambda) &= \{(2, 2, 1, 1), (1, 2, 1, 1)\}, & \rho_{\{2,3\}}(\lambda) &= \{(1, 2, 2, 1), (2, 2, 2, 1)\}, \\ \rho_{\{1,3\}}(\lambda) &= \{(2, 1, 2, 1), (1, 1, 2, 1)\}, & \rho_{\{2,4\}}(\lambda) &= \{(1, 2, 1, 2), (2, 2, 1, 2)\}, \\ \rho_{\{1,4\}}(\lambda) &= \{(2, 1, 1, 2), (1, 1, 1, 2)\}, & \rho_{\{3,4\}}(\lambda) &= \{(1, 1, 2, 2), (2, 1, 2, 2)\}, \\ \rho_{\{1,2,3,4\}}(\lambda) &= \{(2, 2, 2, 2), (1, 2, 2, 2)\}. \end{aligned}$$

There are three further Ferrers diagrams which are fully complementary inside $(2, 2, 2, 2)$, namely $\{(1, 1, 1, 1), (1, 2, 1, 1)\}$, $\{(1, 1, 1, 1), (1, 1, 2, 1)\}$ and $\{(1, 1, 1, 1), (1, 1, 1, 2)\}$.

In the above definition of fully complementary we restricted ourselves to boxes with even side lengths. The definition can be extended to any side lengths, however, as we see next, either there are no such Ferrers diagrams, or we obtain a set of Ferrers diagram which is in bijection to one where the box has only even side lengths. First assume that there exist $k < l$ such that the k -th and l -th side lengths of the box are given by $2n_k + 1$ and $2n_l + 1$ respectively and that λ is a Ferrers diagram which is fully complementary inside this box. Let us regard the point $P = (1, \dots, 1, n_k, 1, \dots, 1, n_l, 1, \dots, 1)$ whose components are 1 except for the k -th and l -th component. Then there exists an $I \subset [d+1]$ of even size such that $P \in \rho_{I, \hat{\mathbf{n}}}(\lambda)$, where $\hat{\mathbf{n}} = (\hat{n}_1, \dots, \hat{n}_{d+1})$ and \hat{n}_i is half of the side length of the box in the direction of i -th standard vector. Let I' be the set $I' = (I \setminus \{k, l\}) \cup (\{k, l\} \setminus I)$. It is immediate that $|I'|$ is even and that $P \in \rho_{I', \hat{\mathbf{n}}}(\lambda)$ which is a contradiction. Hence there exists no fully complementary Ferrers diagram inside a box with at least two odd side lengths.

Now let all side lengths of the box be even with the exception of one side length. Without loss of generality we can assume that the box is a $(2n_1, \dots, 2n_d, 2n_{d+1} + 1)$ -box. For a fully complementary partition π in this box, we see that a point of the form $(i_1, \dots, i_d, n_{d+1} + 1)$ is exactly in π if $i_j \leq n_j$ for all $1 \leq j \leq d$. The map

$$\Psi : \lambda \mapsto \{(i_1, \dots, i_d, i_{d+1}) \in \lambda : i_{d+1} \leq n_{d+1}\} \cup \{(i_1, \dots, i_d, i_{d+1} - 1) \in \lambda : i_{d+1} > n_{d+1} + 1\},$$

is a surjection from Ferrers diagrams inside a $(2n_1, \dots, 2n_d, 2n_{d+1} + 1)$ -box to Ferrers diagrams inside a $(2n_1, \dots, 2n_d, 2n_{d+1})$ -box. It is not difficult to see, that the map Ψ commutes with ρ_I for each $I \subseteq [d+1]$, i.e.,

$$\Psi \circ \rho_{I, (n_1, \dots, n_d, n_{d+1} + \frac{1}{2})} = \rho_{I, (n_1, \dots, n_{d+1})} \circ \Psi.$$

Hence Ψ maps fully complementary Ferrers diagrams inside a $(2n_1, \dots, 2n_d, 2n_{d+1} + 1)$ -box bijectively to those inside a $(2n_1, \dots, 2n_{d+1})$ -box.

2.2. The generating function of FCPs. We call a d -dimensional partition π *fully complementary* inside a $(2n_1, \dots, 2n_{d+1})$ -box if its corresponding Ferrers diagram is fully complementary in this box. The fully complementary partitions inside the $(2, 2, 2, 2)$ -box

are shown next where we write $(\pi_{i,j,1})$ in the top row and $(\pi_{i,j,2})$ in the row below.

$$\begin{array}{cc} 1 & 1 & 1 & 0 & 1 & 0 & 2 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array}$$

For $\mathbf{n} = (n_1, \dots, n_{d+1})$ and a subset $I \subseteq [d]$ we define the map $\gamma_{I,\mathbf{n}}$ as

$$\gamma_{I,\mathbf{n}}(\pi) = \left(\pi_{\rho_{I,\mathbf{n}}(i_1, \dots, i_n)} \right)_{i_1, \dots, i_n}.$$

Again we omit the subscript \mathbf{n} and write γ_I whenever \mathbf{n} is clear from context. The next lemma rephrases the conditions of being fully complementary directly for a d -dimensional partition.

Lemma 2.1. *Let n_1, \dots, n_{d+1} be positive integers. A d -dimensional partition π is fully complementary inside a $(2n_1, \dots, 2n_{d+1})$ -box if and only if*

$$\pi_{i_1, \dots, i_d} \cdot \gamma_J(\pi)_{i_1, \dots, i_d} = 0, \quad (2.1)$$

$$\sum_{I \subseteq [d]} \gamma_I(\pi)_{i_1, \dots, i_d} = 2n_{d+1}, \quad (2.2)$$

for all non-empty subsets $J \subseteq [d]$ of even size and for all $(i_1, \dots, i_d) \in [2n_1] \times \dots \times [2n_d]$.

Proof. Let λ be a fully complementary Ferrers diagram inside a $(2n_1, \dots, 2n_{d+1})$ -box and $(\pi_{i_1, \dots, i_d})_{i_1, \dots, i_d}$ the corresponding d -dimensional partition. For $I \subseteq [d]$ we have

$$\gamma_I(\pi)_{i_1, \dots, i_d} = \begin{cases} |\{(i_1, \dots, i_d, k) \in \rho_I(\lambda) : k \in \mathbb{N}_{>0}\}| & |I| \text{ is even,} \\ |\{(i_1, \dots, i_d, k) \in \rho_{I \cup \{d+1\}}(\lambda) : k \in \mathbb{N}_{>0}\}| & |I| \text{ is odd.} \end{cases}$$

By definition the intersection $\lambda \cap \rho_I(\lambda)$ is empty for even sized I exactly if $\pi_{i_1, \dots, i_d} \cdot \gamma_I(\pi)_{i_1, \dots, i_d} = 0$ for all $(i_1, \dots, i_d) \in [2n_1] \times \dots \times [2n_d]$. For odd sized $I \subseteq [d]$, the intersection $\lambda \cap \rho_{I \cup \{d+1\}}(\lambda)$ is empty if and only if $\pi_{i_1, \dots, i_d} + \gamma_I(\pi)_{i_1, \dots, i_d} \leq 2n_{d+1}$ for all $(i_1, \dots, i_d) \in [2n_1] \times \dots \times [2n_d]$. Finally, the union of all $\rho_J(\lambda)$ with $J \subseteq [d+1]$ of even size is exactly $[2n_1] \times \dots \times [2n_{d+1}]$ if and only if (2.2) is satisfied for all $(i_1, \dots, i_d) \in [2n_1] \times \dots \times [2n_d]$. \square

Denote by $\text{FCP}(n_1, \dots, n_{d+1})$ the set of fully complementary partitions inside a $(2n_1, \dots, 2n_{d+1})$ -box. For $1 \leq k \leq d$ we define the map¹ $\varphi_k : \text{FCP}(n_1, \dots, n_{d+1}) \rightarrow \text{FCP}(n_1, \dots, n_k + 1, \dots, n_{d+1})$ as

$$\varphi_k(\pi)_{i_1, \dots, i_d} = \begin{cases} \pi_{i_1, \dots, i_d} & i_k \leq n_k, \\ n_{d+1} & i_k \in \{n_k + 1, n_k + 2\} \text{ and } i_j \leq n_j \text{ for all } 1 \leq j \neq k \leq d, \\ \pi_{i_1, \dots, i_k - 2, \dots, i_d} & i_k > n_k + 2, \\ 0 & \text{otherwise,} \end{cases}$$

and the map $\varphi_{d+1} : \text{FCP}(n_1, \dots, n_{d+1}) \rightarrow \text{FCP}(n_1, \dots, n_d, n_{d+1} + 1)$ as

$$\varphi_{d+1}(\pi)_{i_1, \dots, i_d} = \begin{cases} \pi_{i_1, \dots, i_d} + 2 & i_j \leq n_j \text{ for all } 1 \leq j \leq d, \\ \pi_{i_1, \dots, i_d} & \text{otherwise.} \end{cases}$$

¹Formally, we have for each $\mathbf{n} = (n_1, \dots, n_{d+1})$ a different map φ_k . However since \mathbf{n} will always be clear from the context, we do not include \mathbf{n} in the notation.

It is not difficult to see, that these maps are well-defined. Let π be the fully complementary partition inside the $(4, 4, 4)$ -box displayed next.

$$\begin{array}{cccc} 4 & 2 & 2 & 0 \\ 3 & 2 & 2 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array}$$

The images of π under the maps φ_1, φ_2 or φ_3 respectively are given below.

$$\begin{array}{ccc} \begin{array}{cccc} 4 & 2 & 2 & 0 \\ 3 & 2 & 2 & 0 \\ 2 & 2 & 0 & 0 \\ 2 & 2 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} & \begin{array}{ccccccc} 4 & 2 & 2 & 2 & 2 & 0 \\ 3 & 2 & 2 & 2 & 2 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} & \begin{array}{cccc} 6 & 4 & 2 & 0 \\ 5 & 4 & 2 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \end{array}$$

As we see in a moment, it is useful to extend the definition of fully complementary partitions to “empty boxes”. In particular we define $\text{FCP}(n_1, \dots, n_{d+1})$ to consist of the “empty array” in case that one n_{k_0} is equal to 0 and all other n_i are positive. We extend the map φ_k to these sets, where φ_k is the identity (mapping the empty array onto the empty array) if $k \neq k_0$ and mapping the empty array to

$$\left(\begin{array}{cc} \left\{ \begin{array}{l} n_{d+1} \\ 0 \end{array} \right. & \begin{array}{l} i_j \leq n_j \text{ for all } 1 \leq j \neq k_0 \leq d, \\ \text{otherwise,} \end{array} \end{array} \right)_{i_1, \dots, i_d},$$

if $k = k_0 \neq d + 1$ and to

$$\left(\begin{array}{cc} \left\{ \begin{array}{l} 2 \\ 0 \end{array} \right. & \begin{array}{l} i_j \leq n_j \text{ for all } 1 \leq j \leq d, \\ \text{otherwise,} \end{array} \end{array} \right)_{i_1, \dots, i_d},$$

if $k = k_0 = d + 1$. The maps φ_k allow us to prove the following recursive structure for FCPs.

Proposition 2.2. *Let $\mathbf{n} = (n_1, \dots, n_{d+1})$ be a sequence of positive integers. Then $\text{FCP}(\mathbf{n})$ is equal to the disjoint union*

$$\text{FCP}(\mathbf{n}) = \bigcup_{1 \leq k \leq d+1} \varphi_k(\text{FCP}(\mathbf{n} - e_k)), \quad (2.3)$$

where e_k is the k -th unit vector.

Proof. First we prove that the images of the φ_k are disjoint. Let $k < l$ be integers with $\varphi_k(\text{FCP}(\mathbf{n} - e_k)) \cap \varphi_l(\text{FCP}(\mathbf{n} - e_l)) \neq \emptyset$ and let π be an element of this intersection. First, we assume that $l = d + 1$. Since $\pi \in \varphi_k(\text{FCP}(\mathbf{n} - e_k))$ we have by definition $\pi_{n_1, \dots, n_d} = n_{d+1}$. On the other hand, $\pi \in \varphi_{d+1}(\text{FCP}(\mathbf{n} - e_{d+1}))$ implies $\pi_{n_1, \dots, n_d} \geq n_{d+1} + 1$ which is a contradiction. Now let $l < d + 1$. By definition of φ_k, φ_l we have

$$\pi_{\mathbf{n}+e_k} = \pi_{\mathbf{n}+e_l} = n_{d+1}.$$

This implies

$$\pi_{\mathbf{n}+e_k} \cdot (\gamma_{\{k,l\}}(\pi))_{\mathbf{n}+e_k} = \pi_{\mathbf{n}+e_k} \cdot \pi_{\mathbf{n}+e_l} = n_{d+1}^2 \neq 0,$$

which contradicts (2.1). Hence the union in (2.3) is disjoint.

Let $\pi \in \text{FCP}(\mathbf{n})$. It is easy to see that $\pi_{\mathbf{n}} \geq n_{d+1}$. If $\pi_{\mathbf{n}} \geq n_{d+1} + 1$, then $\pi \in \varphi_{d+1}(\text{FCP}(\mathbf{n} - e_{d+1}))$. Hence let us assume that $\pi_{\mathbf{n}} = n_{d+1}$. By (2.1), $\gamma_I(\pi)_{\mathbf{n}} = 0$ for all $I \subset [d]$ of even size. By applying any γ_I for an odd sized $I \subseteq [d]$ to (2.1), we see that there is at most one I of odd size for which $\gamma_I(\pi)_{\mathbf{n}}$ is non-zero. Hence by (2.2) there exists exactly one $I \subseteq [d]$ with $\pi_{\mathbf{n}} + \gamma_I(\pi)_{\mathbf{n}} = 2n_{d+1}$ which implies $\gamma_I(\pi)_{\mathbf{n}} = \pi_{\rho_I(\mathbf{n})} = n_{d+1}$. Let

$k \in I$ and define $\widehat{\mathbf{n}} = (1, \dots, n_k, \dots, 1)$ as the vector whose entries are 1 except on position k , where the entry is n_k and $\widehat{\mathbf{i}} = (i_1, \dots, i_d)$ as the vector with $i_k = n_k$ and $i_j \leq n_j$ for all $1 \leq j \neq k \leq d$. Since π is a d -dimensional partition, we have the following inequalities.

$$\begin{array}{ccccccc} \pi_{\widehat{\mathbf{n}}} & \geq & \pi_{\widehat{\mathbf{i}}} & \geq & \pi_{\mathbf{n}} & = & n_{d+1} \\ \vee & & \vee & & \vee & & \parallel \\ \pi_{\widehat{\mathbf{n}}+e_k} & \geq & \pi_{\widehat{\mathbf{i}}+e_k} & \geq & \pi_{\mathbf{n}+e_k} & \geq & \pi_{\rho_I(\mathbf{n})} \end{array}$$

Since $\pi_{\rho_{\{k\}}(\mathbf{n})} = \pi_{\mathbf{n}+e_k} > 0$ and there exists only one non-empty I with $\pi_{\rho_I(\mathbf{n})} \neq 0$, this implies that $I = \{k\}$ and hence $\pi_{\mathbf{n}+e_k} = n_{d+1}$. Furthermore we have $\pi_{\widehat{\mathbf{n}}} + \pi_{\widehat{\mathbf{n}}+e_k} \geq \pi_{\mathbf{n}} + \pi_{\mathbf{n}+e_k} = 2n_{d+1}$ by the above inequalities and $\pi_{\widehat{\mathbf{n}}} + \pi_{\widehat{\mathbf{n}}+e_k} \leq 2n_{d+1}$ by (2.2) since $\gamma_{\{k\}}(\pi)_{\widehat{\mathbf{n}}} = \pi_{\widehat{\mathbf{n}}+e_k}$. This implies $\pi_{\widehat{\mathbf{n}}} = \pi_{\widehat{\mathbf{n}}+e_k} = n_{d+1}$ and hence $\pi_{\widehat{\mathbf{i}}} = \pi_{\widehat{\mathbf{i}}+e_k} = n_{d+1}$ for all $\widehat{\mathbf{i}}$ defined as before. Denote for $l \neq k$ by $\widehat{\mathbf{n}}_l$ the vector with all components equal to 1 except of the k -th and l -th component which are n_k or n_l respectively. Since $\rho_{\{k,l\}}(\widehat{\mathbf{n}}_l) = \widehat{\mathbf{n}}_l + e_k + e_l$ and $\rho_{\{k,l\}}(\widehat{\mathbf{n}}_l + e_k) = \widehat{\mathbf{n}}_l + e_l$ it follows from (2.1) that $\pi_{\widehat{\mathbf{n}}+e_l} = \pi_{\widehat{\mathbf{n}}+e_l+e_k} = 0$ and hence $\pi_{i_1, \dots, i_d} = 0$ for all (i_1, \dots, i_d) such that $i_k \in \{n_k, n_k + 1\}$ and there exists an $i_j > n_j$. Denote by π' the array obtained by deleting all entries for which the i -th component of the index is either n_k or $n_k + 1$. It is not difficult to verify that $\pi' \in \text{FCP}(\mathbf{n} - e_k)$ and $\pi = \varphi_k(\pi')$ which proves the claim. \square

As we see next, Theorem 1.1 is now a direct consequence of the above proposition.

Proof of Theorem 1.1. Let us denote by $Z(\mathbf{x})$ the left hand side of (1.4), i.e., $Z(\mathbf{x}) = \sum_{\mathbf{n} \in \mathbb{N}^{d+1}} |\text{FCP}(\mathbf{n})| \mathbf{x}^{\mathbf{n}}$. Denote further by $Z_j(\mathbf{x}) = \sum_{\mathbf{n}} |\text{FCP}(\mathbf{n})| \mathbf{x}^{\mathbf{n}}$, where the sum is over all $\mathbf{n} \in \mathbb{N}^{d+1}$ where the j -th component is 0 and all the other components are positive. It is immediate that

$$Z_j(\mathbf{x}) = \prod_{\substack{1 \leq i \leq d+1 \\ i \neq j}} \frac{x_i}{1 - x_i},$$

since each of the $\text{FCP}(\mathbf{n})$ in the sum has exactly one element, namely the “empty array”. Using Proposition 2.2, we rewrite $Z(\mathbf{x})$ as

$$\begin{aligned} Z(\mathbf{x}) &= \sum_{\mathbf{n} \in \mathbb{N}_{>0}^{d+1}} |\text{FCP}(\mathbf{n})| \mathbf{x}^{\mathbf{n}} + \sum_{j=1}^{d+1} Z_j(\mathbf{x}) = \sum_{\mathbf{n} \in \mathbb{N}_{>0}^{d+1}} \sum_{i=1}^{d+1} |\varphi_i^{-1}(\text{FCP}(\mathbf{n}))| \mathbf{x}^{\mathbf{n}} + \sum_{j=1}^{d+1} Z_j(\mathbf{x}) \\ &= \sum_{i=1}^{d+1} x_i \left(\sum_{\mathbf{n} \in \mathbb{N}_{>0}^{d+1}} |\text{FCP}(\mathbf{n} - e_i)| \mathbf{x}^{\mathbf{n} - e_i} \right) + \sum_{j=1}^{d+1} Z_j(\mathbf{x}) \\ &= \sum_{i=1}^{d+1} x_i \left(Z(\mathbf{x}) - \sum_{\substack{1 \leq j \leq d+1 \\ j \neq i}} Z_j(\mathbf{x}) \right) + \sum_{j=1}^{d+1} Z_j(\mathbf{x}) \\ &= Z(\mathbf{x}) \sum_{i=1}^{d+1} x_i + \sum_{j=1}^{d+1} Z_j(\mathbf{x}) \left(1 - \sum_{\substack{1 \leq i \leq d+1 \\ i \neq j}} x_i \right). \end{aligned}$$

By bringing all $Z(\mathbf{x})$ terms on one side and using the explicit formula for $Z_j(\mathbf{x})$ we obtain the assertion. \square

Remark 2.3. Let $\mathbf{n} = (n_1, \dots, n_{d+1}) \in \mathbb{N}_{>0}^{d+1}$ be given. By applying Proposition 2.2 iteratively, we see that each $\pi \in \text{FCP}(\mathbf{n})$ can be uniquely written as $\varphi_{i_k} \circ \dots \circ \varphi_{i_1}(\sigma)$

where σ is an “empty array” inside an appropriate “empty box” with dimension $\mathbf{n}' = (n'_1, \dots, n'_{d+1})$ such that $\varphi_{i_1}(\sigma)$ is a non empty array. We map π to the lattice path starting at \mathbf{n}' and ending at \mathbf{n} whose j -th step is e_{i_j} . This yields a bijection between FCPs inside a $(2n_1, \dots, 2n_{d+1})$ -box and lattice paths inside the positive $(d+1)$ -dimensional orthant starting from any integer point on its d -dimensional boundary and ending at \mathbf{n} with step set $\{e_1, \dots, e_{d+1}\}$ such that all coordinates are positive after the first step. Below we show the construction for an FCP inside a $(6, 4, 4)$ -box which is mapped to the lattice path from $(1, 1, 0)$ to $(3, 2, 2)$ with steps $(e_3, e_1, e_3, e_2, e_1)$.

$$\begin{array}{cccc}
4 & 2 & 2 & 0 \\
3 & 2 & 2 & 0 \\
2 & 2 & 0 & 0 \\
2 & 2 & 0 & 0 \\
1 & 0 & 0 & 0 \\
0 & 0 & 0 & 0
\end{array}
\begin{array}{c}
\longleftarrow \varphi_1 \\
\longleftarrow \varphi_2 \\
\longleftarrow \varphi_3 \\
\longleftarrow \varphi_1 \\
\longleftarrow \varphi_3
\end{array}
\begin{array}{cccc}
4 & 2 & 2 & 0 \\
3 & 2 & 2 & 0 \\
1 & 0 & 0 & 0 \\
0 & 0 & 0 & 0
\end{array}
\begin{array}{c}
\longleftarrow \varphi_2 \\
\longleftarrow \varphi_3 \\
\longleftarrow \varphi_1 \\
\longleftarrow \varphi_3
\end{array}
\begin{array}{cccc}
4 & 0 & & 2 & 0 \\
3 & 0 & & 1 & 0 \\
1 & 0 & & 1 & 0 \\
0 & 0 & & 0 & 0
\end{array}
\begin{array}{c}
\longleftarrow \varphi_3 \\
\longleftarrow \varphi_1 \\
\longleftarrow \varphi_3
\end{array}
\begin{array}{cccc}
2 & 0 & & \emptyset \\
0 & 0 & &
\end{array}$$

3. SYMMETRY CLASSES OF QCPPS

3.1. (Quasi) symmetric QCPPs. Remember that a plane partition π is quarter complementary inside a $(2a, 2b, 2c)$ -box if for all $1 \leq i \leq 2a$ and $1 \leq j \leq 2b$ we have

$$\pi_{i,j} \cdot \pi_{2a+1-i, 2b+1-j} = 0, \quad (3.1)$$

and for $\pi_{i,j} > 0$ exactly one of the following equations holds:

$$\pi_{i,j} + \pi_{2a+1-i, j} = 2c, \quad \text{or} \quad \pi_{i,j} + \pi_{i, 2b+1-j} = 2c. \quad (3.2)$$

Regarded as a regular plane partition, π is called symmetric if $a = b$ and $\pi_{i,j} = \pi_{j,i}$ for all $1 \leq i, j \leq 2a$. By (3.1) we see that π can only be symmetric if the entries on its anti-diagonal are 0, i.e., $\pi_{i, 2a+1-i} = 0$ for $1 \leq i \leq 2a$. Together with (3.2) this implies $\pi_{a,a} = 2c$ and hence that the only symmetric quarter complementary plane partition is

$$\pi = \left(\begin{array}{cc} \left\{ \begin{array}{l} 2c \\ 0 \end{array} \right. & \begin{array}{l} i, j \leq a, \\ \text{otherwise,} \end{array} \end{array} \right)_{1 \leq i, j \leq 2a}.$$

In order to obtain more interesting objects we omit the symmetry condition on the anti-diagonal and call a quarter complementary plane partition π *quasi symmetric* if $\pi_{i,j} = \pi_{j,i}$ for all $1 \leq i, j \leq 2a$ and $i \neq 2a+1-j$. Denote by $\text{QS}(a, c)$ the set of quarter complementary plane partitions inside a $(2a, 2a, 2c)$ -box which are quasi symmetric.

Proposition 3.1. *Let a, c be positive integers. Then $\text{QS}(a, c)$ is equal to*

$$\text{QS}(a, c) = \varphi_1 \circ \varphi_2(\text{QS}(a-1, c)) \dot{\cup} \varphi_2 \circ \varphi_1(\text{QS}(a-1, c)) \dot{\cup} \varphi_3(\text{QS}(a, c-1)) \quad (3.3)$$

Proof. Let $\pi \in \text{QS}(a, c)$. By Proposition 2.2 π is either in $\varphi_3(\text{FCP}(a, a, c-1))$, $\varphi_1(\text{FCP}(a-1, a, c))$ or $\varphi_2(\text{FCP}(a, a-1, c))$. In the first case $\varphi_3^{-1}(\pi)$ is obviously quasi symmetric again. Assume $\pi \in \varphi_1(\text{FCP}(a-1, a, c))$. By the definition of φ_1 we have $\pi_{a,j} = \pi_{a+1,j} = c$ if $j \leq a$ and $\pi_{a,j} = \pi_{a+1,j} = 0$ if $j > a$. By the quasi symmetry of π this implies $\pi_{j,a} = \pi_{j,a+1} = c$ for $j < a$ and $\pi_{j,a} = \pi_{j,a+1} = 0$ for $j > a+1$. Therefore we obtain $\varphi_1^{-1}(\pi) \in \varphi_2(\text{FCP}(a-1, a-1, c))$. It is not difficult to check that $\varphi_1^{-1}(\varphi_2^{-1}(\pi))$ is again quasi symmetric. The last case follows analogously. \square

By defining $\text{QS}(a, c)$ to consist of the “empty array” if either a or c is equal to 0, we obtain immediately the following corollary.

Corollary 3.2. *The generating function for quasi symmetric quarter complementary plane partitions is given by*

$$\sum_{a,c \geq 0} |\text{QS}(a, c)| x^a y^c = \frac{x + y - 2x^2 - xy}{(1-x)(1-2x-y)}.$$

3.2. Cyclically symmetric QCPPs. A plane partition π is called *cyclically symmetric* if a point (i, j, k) in its Ferrers diagram implies that (j, k, i) is also in its Ferrers diagram. As we see next, there is no cyclically symmetric quarter complementary plane partition. Let π be a cyclically symmetric quarter complementary Ferrers diagram inside a $(2a, 2a, 2a)$ -box. It is not difficult to see, that being quarter complementary implies that one of the points $(1, 1, 2a)$, $(1, 2a, 1)$ or $(2a, 1, 1)$ has to be part of π , and hence all of them since π is cyclically symmetric. The set $\rho_{\{1,2\}}(\pi)$ therefore contains the points $(2a, 1, 1)$ and $(1, 2a, 1)$ which is a contradiction to $\pi \cap \rho_{\{1,2\}}(\pi) = \emptyset$.

Contrary to the previous subsection, we did not find an “interesting” generalisation of cyclically symmetric to “quasi cyclically symmetric” for which we can deduce either a result or a conjecture in the case of QCPPs.

3.3. Self- and transpose-complementary QCPPs. In order to study self- or transpose-complementary QCPPs, we need to specify the box we want to consider the complementation in. For a QCPP inside a $(2a, 2b, 2c)$ -box, the possible boxes for complementation are a $(2a, 2b, c)$ -, a $(2a, b, 2c)$ - and an $(a, 2b, 2c)$ -box. For symmetry reasons it suffices to consider QCPPs which are self- or transpose-complementary inside a $(2a, 2b, c)$ -box. We call a QCPP π inside a $(2a, 2b, 2c)$ -box *self-complementary* if $\pi_{i,j} + \pi_{2a+1-i, 2b+1-j} = c$ for all $1 \leq i \leq 2a$ and $1 \leq j \leq 2b$, and *quasi transpose-complementary*² if $a = b$ and $\pi_{i,j} + \pi_{2a+1-j, 2a+1-i} = c$ for all $1 \leq i, j \leq 2a$ with $i \neq 2a + 1 - j$. We obtain the following enumeration results.

Proposition 3.3. *The number of self-complementary QCPPs inside a $(2a, 2b, 2c)$ -box is $\binom{a+b}{a}$.*

Proof. The condition $\pi_{i,j} + \pi_{2a+1-i, 2b+1-j} = c$ implies that all entries are at most c . Together with (3.2) this implies that all entries are either 0 or c . Hence by Proposition 2.2, π is either of the form $\varphi_1(\sigma)$ or $\varphi_2(\sigma)$ for an appropriate QCPP σ . It is easy to check that σ is again self-complementary. The assertion follows now by induction on $a + b$. \square

Proposition 3.4. *A QCPP inside a $(2a, 2a, 2c)$ -box is quasi transpose-complementary if it is quasi symmetric and self-complementary. The number of quasi transpose-complementary QCPPs inside a $(2a, 2a, 2c)$ -box is 2^a .*

Proof. The condition $\pi_{i,j} + \pi_{2a+1-j, 2a+1-i} = c$ together with (3.2) implies that all entries of π are either 0 or c . Since π is quarter complementary inside a $(2a, 2a, 2c)$ -box, the sum over all entries must be $2a^2c$. Hence exactly $2a^2$ entries are equal to c and $2a^2$ entries are equal to 0. For $i \leq a$ exactly one of the equations $\pi_{i,i} + \pi_{i, 2a+1-i} = 2c$ or $\pi_{i,i} + \pi_{2a+1-i, i} = 2c$ holds by (2.2). Hence exactly half of the entries on the anti-diagonal are equal to c . For $i \neq 2a + 1 - j$ we therefore have $\pi_{2a+1-j, 2a+1-i} = c$ exactly if $\pi_{i,j} = 0$, which is by (3.1) equivalent to $\pi_{2a+1-i, 2a+1-j} = c$. This implies that π is quasi symmetric and therefore also self-complementary. It is immediate that a quasi symmetric, self-complementary QCPP is also quasi transpose-complementary.

By the proof of Proposition 3.1 and Proposition 3.3 each quasi transpose-complementary QCPP π is of the form $\varphi_1 \circ \varphi_2(\sigma)$ or $\varphi_2 \circ \varphi_1(\sigma)$, where σ is a quasi transpose-complementary QCPP inside a $(2(a-1), 2(a-1), 2c)$ -box. The assertion follows now by induction on a . \square

²It is immediate that there are no transpose-complementary QCPPs since the condition $\pi_{i,j} + \pi_{2a+1-j, 2a+1-i} = c$ on the anti-diagonal contradicts (2.1). Similar to quasi symmetric, we therefore exclude the condition on the anti-diagonal to obtain interesting objects.

4. QUASI SYMMETRY CLASSES OF PLANE PARTITIONS

In the previous section we introduced variations of two symmetry classes for quarter complementary plane partitions. The aim of this section is to consider these and similar symmetry classes for plane partitions. The following three conjectures were found by computer experiments. We have added the according data as well as explicit guessed enumeration formulas for small values of one of the parameters in the appendix.

Conjecture 4.1. *Let us denote by $\text{qspp}(a, c)$ the number of quasi symmetric plane partitions inside an (a, a, c) -box. Then,*

$$\text{qspp}(a, c - a) = \begin{cases} c \binom{c+a-1}{2a-1} p_a(c) & a \text{ is even,} \\ \binom{c+a-1}{2a-1} p_a(c) & a \text{ is odd,} \end{cases} \quad (4.1)$$

where $p_a(c)$ is an irreducible polynomial in $\mathbb{Q}[c]$ that is even, i.e., $p_a(c) = p_a(-c)$. Further the common denominator of the coefficients of $p_a(c)$ is a product of “small primes”.

Conjecture 4.2. *We call a plane partition π inside an (a, a, c) -box **quasi transpose complementary of second kind (QTC2)**, if π is transpose complementary except along the diagonal, i.e., $\pi_{i,j} + \pi_{a+1-j, a+1-i} = c$ for all $1 \leq i, j \leq a$ with $i \neq j$. Then for $a \geq 2$, the number $\text{qtcpp}_2(a, c)$ of QTC2 plane partitions inside an (a, a, c) -box is given by*

$$\text{qtcpp}_2(a, c - \frac{a}{2}) = \begin{cases} c \binom{c+\frac{a}{2}-1}{\frac{a}{2}-1} p_a(c) & a \text{ is even,} \\ \binom{c+\frac{a}{2}-1}{\frac{a}{2}-1} p_a(c) & a \text{ is odd,} \end{cases} \quad (4.2)$$

where $p_a(c)$ is an irreducible polynomial in $\mathbb{Q}[c]$ that is even. Further the common denominator of the coefficients of $p_a(c)$ is a product of “small primes”.

Conjecture 4.3. *Denote by $\text{qtcspp}_2(a, c)$ the number of symmetric QTC2 plane partitions. Then for $a \geq 2$,*

$$\text{qtcspp}_2(a, c - \frac{a}{2}) = \begin{cases} c \binom{c+\frac{a}{2}-1}{\frac{a}{2}-1} p_a(c) & a \text{ is even,} \\ \binom{c+\frac{a}{2}-1}{\frac{a}{2}-1} p_a(c) & a \text{ is odd,} \end{cases} \quad (4.3)$$

where $p_a(c)$ is an irreducible polynomial in $\mathbb{Q}[c]$ that is even. Further the common denominator of the coefficients of $p_a(c)$ is a product of “small primes”.

4.1. Proof of Theorem 1.2. Let π be a quasi transpose complementary plane partition (QTCPP) inside an (n, n, c) -box. By definition we have $\pi_{n-j,j} \geq \pi_{n-j+1,j} \geq \pi_{n-j+1,j+1} = c - \pi_{n-j,j}$ for each $1 \leq j \leq n-1$ and equivalently $\pi_{n-j+1,j+1} = c - \pi_{n-j,j} \leq c - \pi_{n-j+1,j} \leq \pi_{n-j,j}$. Hence π stays a QTCPP if we replace its diagonal entries $\pi_{n-j+1,j}$ by $\max(\pi_{n-j+1,j}, c - \pi_{n-j+1,j})$ for all $1 \leq j \leq n$. We denote the resulting QTCPP by $\hat{\pi}$. Denote by $d(\hat{\pi})$ the number of anti-diagonal entries which are equal to $\hat{c} = \lfloor \frac{c}{2} \rfloor$ and define the weight $\omega(\hat{\pi})$ as

$$\omega(\hat{\pi}) = \begin{cases} 2^n & c \text{ is odd,} \\ 2^{n-d(\hat{\pi})} & c \text{ is even.} \end{cases}$$

Since $|\{\pi_{n-j+1,j}, c - \pi_{n-j+1,j}\}| = 1$ implies that c is even and $\pi_{n-j+1,j} = \hat{c}$, it is clear that there are $\omega(\hat{\pi})$ many QTCPPs mapping to $\hat{\pi}$ by the above map. Hence the number of QTCPPs is equal to the weighted enumeration of QTCPPs $\hat{\pi}$ whose anti-diagonal entries are at least $\frac{c}{2}$. In the following we present two (and a half) proofs for the weighted enumeration of these $\hat{\pi}$.

For odd c , each $\hat{\pi}$ corresponds to a plane partition with entries at most \hat{c} for which the i -th row from top has at most $n+1-i$ positive entries. The number of these plane partitions can be found in [16, Corollary 4.1], compare also with [4, Corollary 4.1]. For even c , each $\hat{\pi}$ corresponds to a lozenge tiling in the “top half” of an hexagon with side lengths n, n, c, n, n, c , where the bottom of the region is a zig-zag shape directly below the

centre line of the hexagon, and each \diamond lozenge at the bottom of the region is weighted by $\frac{1}{2}$, see Figure 3 (right) for an example. The number of these tilings is given in [4, Corollary 4.3]. The assertion follows in both cases by using the explicit formulas from [4, 16].

Denote by $\text{qtcpp}(n, n, c)$ the number of QTCPPs inside an (n, n, c) -box and denote by $M(R)$ the number of perfect matchings of a region R . We can rephrase the above observations as

$$\text{qtcpp}(n, n, 2c) = 2^n M(P'_{n,n,c}), \quad \text{qtcpp}(n, n, 2c + 1) = 2^n M(P_{n,n,c}),$$

where the Regions $P_{n,n,c}$ is defined in [4, Section 4, Fig. 5] and $P'_{n,n,c}$ is defined in [4, Section 4, Fig. 10]. On the one side we have by Ciucus factorization theorem for graphs with reflective symmetry [3] the identity

$$\text{PP}(a, a, 2c) = 2^n M(P'_{n,n,c})M(P_{n-1,n-1,c}),$$

where $\text{PP}(n, n, 2c)$ denotes the number of plane partitions inside an $(n, n, 2c)$ -box (compare for example with [3] or [4, Equation (4.16)]). This implies the assertion for even c if we know it for odd c and vice versa by using the explicit formulas for the number of (symmetric) plane partitions. Further it is well known, that $2^n M(P'_{n,n,c}) = \text{SPP}(n, n, 2c)$, see [5, Equation (5.1)] or [6, Equation (2.6)] and that $M(P_{n-1,n-1,c}) = \text{TCPP}(n, n, 2c)$, where $\text{TCPP}(n, n, 2c)$ is the number of transpose-complementary plane partitions inside an $(n, n, 2c)$ -box (see for example [3, Section 6]). Combining the above and the assertion (which is already proved above) we obtain immediately the numerical connection between SPPs and TCPPs stated in (1.5).

Finally we present another proof using non-intersecting lattice paths. We regard $\hat{\pi}$ as a lozenge tiling as above and draw n lattice paths in the following way. Each path ends at the top right boundary of the hexagon, uses the steps $\begin{smallmatrix} \blacksquare \\ \blacksquare \end{smallmatrix}$ or $\begin{smallmatrix} \blacktriangle \\ \blacktriangle \end{smallmatrix}$ and the i -th path from left has length $n + 1 - i + \hat{c}$; see Figure 3 (left) for an example. By straightening the paths, we obtain non-intersecting lattice paths starting at $A_i = (2i, -i)$ and ending at $E_i = (n + 1 + i, \hat{c} - i)$ with north-steps $(0, 1)$ and east-steps $(1, 0)$, see Figure 3 (right). For odd c , each family of paths has the same weight, namely 2^n . Hence the weighted enumeration is therefore by the Lindström-Gessel-Viennot Theorem equal to

$$2^n \det_{1 \leq i, j \leq n} \left(\binom{n + \hat{c} + 1 - i}{n + 1 + j - 2i} \right). \quad (4.4)$$

For even c we see that an anti-diagonal entry $\pi_{n+1-j,j} = \hat{c}$ corresponds to the j -th path starting with an east-step. Define the points $B_i^0 = (2i + 1, -i)$ and $B_i^1 = (2i, -i + 1)$ which are reached from A_i by an east-step or a north-step respectively. By deleting the first step of each path, we obtain a family of non-intersecting lattice paths starting from either B_i^0 or B_i^1 , where the weight is given by 2 to the power of times we start at B_i^1 . Using again the Lindström-Gessel-Viennot Theorem we obtain for the weighted enumeration

$$\begin{aligned} \sum_{(b_1, \dots, b_n) \in \{0,1\}^n} \det_{1 \leq i, j \leq n} \left(2^{b_i} \binom{n + \hat{c} - i}{n + j - 2i + b_i} \right) \\ = \det_{1 \leq i, j \leq n} \left(\binom{n + \hat{c} - i}{n + j - 2i} + 2 \binom{n + \hat{c} - i}{n + j - 2i + 1} \right), \end{aligned} \quad (4.5)$$

where we used the multilinearity of the determinant in the last step. Both determinants could be evaluated by guessing the corresponding LU decomposition and using the Pfaff-Saalschütz-summation formula, see for example [17, (2.3.1.3); Appendix (III.2)]; we omit




FIGURE 3. A QTCPP $\hat{\pi}$ inside a $(5, 5, 6)$ -box whose anti-diagonal entries are at least 3 (left) and its corresponding lattice path configuration (right).

however the details since there is a simpler and more elegant solution as follows. First we rewrite the determinant in (4.5) as

$$\det_{1 \leq i, j \leq n} \left(\binom{n + \hat{c} - i}{n + j - 2i} (n + 2\hat{c} - j + 1) \right). \quad (4.6)$$

Then both determinants are special cases of determinant evaluation [10, Equation (3.13)]

$$\det_{1 \leq i, j \leq n} \left(\binom{BL_i + A}{L_i + j} \right) = \frac{\prod_{1 \leq i < j \leq n} (L_i - L_j)}{\prod_{i=1}^n (L_i + n)!} \prod_{i=1}^n \frac{(BL_i + A)!}{((B-1)L_i + A - 1)!} \prod_{i=1}^n (A - Bi + 1)_{i-1},$$

by setting $L_i = n + 1 - 2i$, $B = \frac{1}{2}$ and $A = \hat{c} + \frac{n+1}{2}$ in the case of (4.4) and $L_i = n - 2i$, $B = \frac{1}{2}$ and $A = \hat{c} + \frac{n}{2}$ in the case of (4.6).

5. ACKNOWLEDGEMENT

The author thanks Ilse Fischer and Christian Krattenthaler for helpful discussions.

APPENDIX A. DATA FROM THE COMPUTER EXPERIMENTS

A.1. Quasi symmetric plane partitions. The values for the number $\text{qspp}(a, c)$ of quasi symmetric plane partitions inside an (a, a, c) -box with $a \leq 6$ and $c \leq 10$ are shown in the next table.

$c \setminus a$	1	2	3	4	5	6
0	1	1	1	1	1	1
1	2	6	12	32	60	164
2	3	20	69	400	1312	7952
3	4	50	272	3052	16572	200956
4	5	105	846	16932	145428	3284589
5	6	196	2232	74868	979068	38963092
6	7	336	5214	278928	5376673	360346984
7	8	540	11088	908336	25100880	2727638524
8	9	825	21879	2653001	102593290	17499041992
9	10	1210	40612	7081776	375222392	97667820784
10	11	1716	71643	17524416	1248707892	483901238656

Using the above values, we conjecture the following formulas for $\text{qspp}(a, c)$ for $1 \leq a \leq 6$.

$$\text{qspp}(1, c-1) = \binom{c}{1},$$

$$\text{qspp}(2, c-2) = c \binom{c+1}{3},$$

$$\text{qspp}(3, c-3) = \binom{c+2}{5} \frac{c^2 - 2}{7},$$

$$\text{qspp}(4, c-4) = c \binom{c+3}{7} \frac{41c^4 - 229c^2 - 892}{23760},$$

$$\text{qspp}(5, c-5) = \binom{c+4}{9} \frac{683c^8 - 8206c^6 - 14473c^4 - 310644c^2 + 756000}{122522400},$$

$$\text{qspp}(6, c-6) = c \binom{c+5}{11} \frac{1}{161911881331200} (56381c^{12} - 1850347c^{10} + 11282865c^8 - 28759181c^6 - 1859025278c^4 + 20697349128c^2 + 194655992832).$$

A.2. Quasi transpose complementary plane partitions of second kind. For the number $\text{qtcp}_2(a, c)$ of QTC2 plane partitions inside an (a, a, c) -box with $a \leq 6$ and $c \leq 10$, we have the following values.

$c \setminus a$	1	2	3	4	5	6
0	1	1	1	1	1	1
1	2	4	7	24	62	216
2	3	9	26	224	1210	12177
3	4	16	70	1280	12819	314624
4	5	25	155	5361	91694	4860048
5	6	36	301	18088	496796	51955744
6	7	49	532	52032	2185860	420545536
7	8	64	876	132408	8177634	2735918368
8	9	81	1365	305745	26861407	14918043569
9	10	100	2035	652432	79299714	–
10	11	121	2926	1304160	214133686	–

We conjecture the following explicit formulas for $qtcpp_2(a, c)$ for $1 \leq a \leq 6$.

$$\begin{aligned} qtcpp_2(1, c - \frac{1}{2}) &= \binom{c + \frac{1}{2}}{1}, \\ qtcpp_2(2, c - 1) &= c \binom{c}{1}, \\ qtcpp_2(3, c - \frac{1}{2}) &= \binom{c + \frac{1}{2}}{2} \frac{4c^2 + 3}{12}, \\ qtcpp_2(4, c - 2) &= c \binom{c + 1}{3} \frac{5c^4 + 19c^2 - 16}{280}, \\ qtcpp_2(5, c - \frac{1}{2}) &= \binom{c + \frac{3}{2}}{4} \frac{54016c^8 + 522240c^6 - 1322016c^4 + 80480c^2 + 883575}{159667200}, \\ qtcpp_2(6, c - 3) &= c \binom{c + 2}{5} \frac{1}{256505356800} (73325c^{12} + 1648357c^{10} - 12312285c^8 \\ &\quad + 29029591c^6 + 201378740c^4 - 876526848c^2 + 395435520). \end{aligned}$$

A.3. Quasi transpose complementary symmetric plane partitions of second kind. Computer experiments yield the following values for the number $qtcspp_2(a, c)$ of symmetric QTC2 plane partitions inside an (a, a, c) -box with $a \leq 6$ and $c \leq 10$.

$c \setminus a$	1	2	3	4	5	6
0	1	1	1	1	1	1
1	2	4	5	12	18	40
2	3	9	14	68	142	625
3	4	16	30	260	723	5728
4	5	25	55	777	2782	36876
5	6	36	91	1960	8796	184224
6	7	49	140	4368	24036	759708
7	8	64	204	8856	58674	2695200
8	9	81	285	16665	130911	8468889
9	10	100	385	29524	271414	–
10	11	121	506	49764	529386	–

For $a \leq 6$ we conjecture the following formulas for $qtcspp_2(a, c)$.

$$\begin{aligned} qtcspp_2(1, c) &= c + 1, \\ qtcspp_2(2, c - 1) &= c \binom{c}{1}, \\ qtcspp_2(3, c - \frac{3}{2}) &= \binom{c + \frac{1}{2}}{2} \frac{2c}{3}, \\ qtcspp_2(4, c - 2) &= c \binom{c + 1}{3} \frac{c^2 + 1}{10}, \\ qtcspp_2(5, c - \frac{5}{2}) &= \binom{c + \frac{3}{2}}{4} \frac{144c^4 + 248c^2 - 455}{6720}, \\ qtcspp_2(6, c - 3) &= c \binom{c + 2}{5} \frac{3c^6 + 15c^4 - 58c^2 + 200}{9240}, \end{aligned}$$

REFERENCES

- [1] A. Amanov and D. Yeliussizov. MacMahon's statistics on higher-dimensional partitions. *arXiv:2009.00592*, 2020.

- [2] A. O. L. Atkin, P. Bratley, I. G. Macdonald, and J. K. S. McKay. Some computations for m -dimensional partitions. *Proc. Cambridge Philos. Soc.*, 63:1097–1100, 1967. doi:10.1017/s0305004100042171.
- [3] M. Ciucu. Enumeration of perfect matchings in graphs with reflective symmetry. *J. Combin. Theory Ser. A*, 77(1):67–97, 1997. doi:10.1006/jcta.1996.2725.
- [4] M. Ciucu and I. Fischer. Proof of two conjectures of Ciucu and Krattenthaler on the enumeration of lozenge tilings of hexagons with cut off corners. *J. Combin. Theory Ser. A*, 133:228–250, 2015. doi:10.1016/j.jcta.2015.02.008.
- [5] M. Ciucu and C. Krattenthaler. A factorization theorem for classical group characters, with applications to plane partitions and rhombus tilings. In I. S. Kotsireas and E. V. Zima, editors, *Advances in combinatorial mathematics: Proceedings of the Waterloo Workshop in Computer Algebra 2008*, pages 39–60. Springer, Berlin, 2010. doi:10.1007/978-3-642-03562-3.
- [6] M. Ciucu and C. Krattenthaler. A factorization theorem for lozenge tilings of a hexagon with triangular holes. *Trans. Amer. Math. Soc.*, 369(5):3655–3672, 2017. doi:10.1090/tran/7047.
- [7] G. David and C. Tomei. The problem of the calissons. *Amer. Math. Monthly*, 96(5):429–431, 1989. doi:10.2307/2325150.
- [8] D. E. Knuth. A note on solid partitions. *Math. Comp.*, 24:955–961, 1970. doi:10.2307/2004628.
- [9] C. Koutschan, M. Kauers, and D. Zeilberger. Proof of George Andrews’s and David Robbins’s q -TSP conjecture. *Proc. Natl. Acad. Sci. USA*, 108(6):2196–2199, 2011. doi:10.1073/pnas.1019186108.
- [10] C. Krattenthaler. Advanced determinant calculus. *Sém. Lothar. Combin.*, 42:Art. B 42q, 67 pp., 1999.
- [11] C. Krattenthaler. *The mathematical legacy of Richard P. Stanley*, chapter Plane partitions in the work of Richard Stanley and his school, pages 231–261. Amer. Math. Soc., Providence, RI, 2016. doi:10.1090/mbk/100.
- [12] P. A. MacMahon. Memoir on the theory of the partition of numbers, I. *Lond. Phil. Trans. (A)*, 187:619–673, 1897.
- [13] P. A. MacMahon. Partitions of numbers whose graphs possess symmetry. *Trans. Cambridge Philos. Soc.*, 17:149–170, 1899.
- [14] P. A. MacMahon. *Combinatory Analysis, vol. 2*. Cambridge University Press, 1916; reprinted by Chelsea, New York, 1960.
- [15] W. H. Mills, D. P. Robbins, and H. Rumsey Jr. Self-complementary totally symmetric plane partitions. *J. Combin. Theory Ser. A*, 42(2):277–292, 1986. doi:10.1016/0097-3165(86)90098-1.
- [16] R. A. Proctor. Odd symplectic groups. *Invent. Math.*, 92(2):307–332, 1988. doi:10.1007/BF01404455.
- [17] L. J. Slater. *Generalized hypergeometric functions*. Cambridge University Press, Cambridge, 1966.
- [18] R. P. Stanley. A baker’s dozen of conjectures concerning plane partitions. In *Combinatoire énumérative*, volume 1234 of *Lecture Notes in Math.*, pages 285–293. Springer, Berlin, 1986. doi:10.1007/BFb0072521.
- [19] R. P. Stanley. Symmetries of plane partitions. *J. Combin. Theory Ser. A*, 43(1):103–113, 1986. Erratum 44:310, 1987. doi:10.1016/0097-3165(86)90028-2.

UNIVERSITY OF VIENNA, AUSTRIA

URL: <https://homepage.univie.ac.at/florian.schreier-aigner>