

NILPOTENT GROUPS WHOSE DIFFERENCE GRAPHS HAVE POSITIVE GENUS

PARVEEN, JITENDER KUMAR*

ABSTRACT. The power graph of a finite group G is a simple undirected graph with vertex set G and two vertices are adjacent if one is a power of the other. The enhanced power graph of a finite group G is a simple undirected graph whose vertex set is the group G and two vertices a and b are adjacent if there exists $c \in G$ such that both a and b are powers of c . In this paper, we study the difference graph $\mathcal{D}(G)$ of a finite group G which is the difference of the enhanced power graph and the power graph of G with all isolated vertices removed. We characterize all the finite nilpotent groups G such that the genus (or cross-cap) of the difference graph $\mathcal{D}(G)$ is at most 2.

1. HISTORICAL BACKGROUND AND MAIN RESULT

There are number of graphs attached to groups, in particular: Cayley graphs, commuting graphs, power graphs, enhanced power graph etc. These graphs have been studied extensively by various researchers because of their vast applications (see [16, 19, 20, 22]). The *power graph* $\mathcal{P}(G)$ of a finite group G is a simple undirected graph with vertex set G such that two vertices a and b are adjacent if one is a power of the other. Kelarev and Quinn [21] introduced the concept of directed power graph. Topological graph theory is mainly concerned with embedding of a graph into a surface without edge crossings. Its applications lie in electronic printing circuits where the purpose is to embed a circuit, that is, the graph on a circuit board (the surface), without two connections crossing each other, resulting in a short circuit. The problem of determining the genus of a graph is NP-hard [35]. Mirzargar *et al.* [31] classified all the finite groups with planar power graphs. Further, Doostabadi *et al.* [11], characterized the finite groups whose power graphs are of (non)orientable genus one. Then all the finite groups with (non)orientable genus two power graphs have been characterized in [28]. The undirected power graphs of groups have been studied in other aspects, see [7, 9, 13, 24, 26] and references therein. The *commuting graph* $\Delta(G)$ of a group G is a simple graph with vertex set G and two distinct vertices x, y are adjacent if $xy = yx$. Clearly, all the central elements are the dominating vertices in $\Delta(G)$. Moreover, $\Delta(G)$ has been investigated by various researchers by taking its vertex set as non-central elements. Afkhami *et al.* [2] characterized all the finite groups whose commuting and noncommuting graphs are planar, projective planar and of genus one, respectively. Results on the commuting graph associated to groups can be found in [6, 15, 17, 25, 29, 38] and references therein. Aalipour *et al.* [1] characterize the finite group G such that the power graph $\mathcal{P}(G)$ and the commuting graph $\Delta(G)$ are not equal and hence they introduced a new graph between power graph and commuting graph, called enhanced power graph. The *enhanced power graph* $\mathcal{P}_E(G)$ of a finite group G is a simple undirected graph with vertex set G and two vertices x and y are adjacent if $x, y \in \langle z \rangle$ for some $z \in G$. Equivalently, two vertices x and y are adjacent in $\mathcal{P}_E(G)$ if and only if $\langle x, y \rangle$ is a cyclic subgroup of G . The clique number of enhanced power graph of an arbitrary group G was obtained by Aalipour *et al.* [1] in terms of orders of elements of G . Bera and Bhuniya [3] proved that $\mathcal{P}_E(G)$ is planar if and only if order of each element of G is at most 4. However, all the finite groups having genus one (or two) enhanced power graphs have not been classified so far. For a detailed list of results and open problems on enhanced power graphs of groups, we refer the reader to [27].

From the above definitions, it is easy to observe that the power graph is a spanning subgraph of the enhanced power graph. Also, the enhanced power graph is a spanning subgraph of the commuting graph. Consequently, Aalipour *et al.* [1, Question 42] motivated the researchers to study the connectedness of the difference graph of the commuting graph and power graph of a group G i.e. the graph with vertex set G in which x and y are adjacent if they commute but neither is a power of the other. Further, Cameron [8] discussed some developments on the difference graph $\Delta(G) - \mathcal{P}(G)$. Moreover, some results on the difference graph of the commuting graph and the enhanced power graph of a group have been discussed in [8]. Motivated by the work of [8], Biswas *et al.* [5] studied the difference graph $\mathcal{D}(G) = \mathcal{P}_E(G) - \mathcal{P}(G)$ of enhanced power graph and power graph of a finite group G with all isolated vertices

2020 *Mathematics Subject Classification.* 05C25.

Key words and phrases. Enhanced power graph, power graph, nilpotent groups, genus and cross-cap of a graph.

* Corresponding author.

removed. For certain group classes, the connectedness and perfectness of $\mathcal{D}(G)$ has been investigated in [5]. Together with the planarity, various forbidden graph classes of $\mathcal{D}(G)$ have been studied in [33]. The purpose of this article is to classify all the finite nilpotent groups such that the difference graph $\mathcal{D}(G)$ is of genus (or cross-cap) at most two. If G is a p -group, then it is well known that the power graph and enhanced power graph of G are equal. Thus, $\mathcal{D}(G)$ is a null graph, whenever G is a p -group. Before providing our main result of this paper, for a finite group G , first we define

- G satisfies the condition \mathcal{C}_1 , if $G \cong P \times \mathbb{Z}_3$, where P is a 2-group with exponent 4. Moreover, P contains two maximal cyclic subgroups H and K of order 4 such that $|H \cap K| = 2$, and the intersection of any other pair of maximal cyclic subgroups of P is trivial.
- G satisfies the condition \mathcal{C}_2 , if $G \cong P \times \mathbb{Z}_3$, where P is a 2-group with exponent 4. Moreover, P contains four maximal cyclic subgroups H_1, H_2, H_3 and H_4 of order 4 such that $|H_1 \cap H_2| = |H_3 \cap H_4| = 2$, and the intersection of any other pair of maximal cyclic subgroups of P is trivial.
- G satisfies the condition \mathcal{C}_3 , if $G \cong P \times \mathbb{Z}_3$, where P is a 2-group with exponent 4. Moreover, P contains three maximal cyclic subgroups H_1, H_2 and H_3 of order 4 such that $|H_1 \cap H_2 \cap H_3| = 2$, and the intersection of any other pair of maximal cyclic subgroups of P is trivial.

For $1 \leq i \leq 3$, if the group G satisfies the condition \mathcal{C}_i , then we write it by the group \mathcal{G}_i . The main result of this paper is as follows.

Theorem 1.1. *Let G be a nilpotent group which is not a p -group and let $\gamma(\mathcal{D}(G))$ and $\bar{\gamma}(\mathcal{D}(G))$ be the genus and cross-cap of $\mathcal{D}(G)$, respectively. Then*

- (i) $\gamma(\mathcal{D}(G)) = 1$ if and only if G is isomorphic to one of the following groups:

$$\mathbb{Z}_{18}, \mathbb{Z}_{20}, \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_5, \mathbb{Z}_{28}, \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_7, \mathcal{G}_1.$$

- (ii) $\gamma(\mathcal{D}(G)) = 2$ if and only if G is isomorphic to one of the following groups:

$$\mathbb{Z}_{35}, \mathbb{Z}_4 \times \mathbb{Z}_3 \times \mathbb{Z}_3, \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3, \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_{11}, \mathbb{Z}_{44}, \mathcal{G}_2, \mathcal{G}_3.$$

- (iii) $\bar{\gamma}(\mathcal{D}(G)) = 1$ if and only if G is isomorphic to \mathbb{Z}_{20} or $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_5$.

- (iv) $\bar{\gamma}(\mathcal{D}(G)) = 2$ if and only if G is isomorphic to one of the groups: $\mathbb{Z}_{18}, \mathbb{Z}_{28}, \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_7, \mathcal{G}_1$.

2. PRELIMINARIES

In this section, we recall the necessary definitions, results which we need in the sequel of this paper. We also fix our notations in this section. Let G be a group. The order of an element $x \in G$ is denoted by $o(x)$ and we denote $\pi_G = \{o(g) : g \in G\}$. By $\langle x, y \rangle$, we mean the subgroup of G generated by x and y . The *exponent* $\exp(G)$ of a finite group G is defined as the least common multiple of the orders of all the elements of G . For $d \in \pi_G$, C_d denotes the number of cyclic subgroups of order d in G . A cyclic subgroup of a group G is called a *maximal cyclic subgroup* if it is not properly contained in any cyclic subgroup of G other than itself. Note that if G is a cyclic group, then G is the only maximal cyclic subgroup of G . A finite group G is called a *p -group* if $|G| = p^\alpha$ for some prime p . A finite group G is said to be an EPPO-group if the order of each element of G is of prime power. Otherwise, G is called a non-EPPO-group. The following results are useful for later use.

Lemma 2.1. *Let G be a finite p -group with exponent p^2 . Then either G has exactly one cyclic subgroup of order p^2 or G contains at least two cyclic subgroups M and N of order p^2 such that $|M \cap N| = p$.*

Proof. First note that $|Z(G)| \geq p$. If G has exactly one cyclic subgroup of order p^2 , then there is nothing to prove. We may now suppose that G has two cyclic subgroups M and N of order p^2 . Let $x \in Z(G)$ such that $o(x) = p$. If $x \in M \cap N$, then $|M \cap N| = p$. Now assume that $x \notin M$ and $M = \langle y \rangle$. We claim that $o(xy) = p^2$. Clearly, $(xy)^{p^2} = x^{p^2} y^{p^2} = e$. Consequently, $o(xy) | p^2$. Thus, $o(xy) \in \{1, p, p^2\}$. If $o(xy) = 1$, then $x = y^{-1}$; a contradiction. If $o(xy) = p$, then $x^p y^p = e y^p = e$, again a contradiction. It follows that $o(xy) = p^2$. If $\langle xy \rangle = \langle y \rangle$, then $xy = y^k$ for some positive integer k and so $x = y^{k-1}$, which is not possible. Thus, $M' = \langle xy \rangle$ is a cyclic subgroup of order p^2 in G . Moreover, $M \cap M' = \{e, y^p, y^{2p}, \dots, y^{(p-1)p}\}$. Thus, the result holds. \square

Theorem 2.2. [4, 18, 23, 30] *Let G be a finite p -group of exponent p^k . Assume that G is not cyclic for an odd prime p , and for $p = 2$ it is neither cyclic nor of maximal class. Then*

- (i) $C_p \equiv 1 + p \pmod{p^2}$.
- (ii) $p \mid C_{p^i}$ for every $2 \leq i \leq k$.

Corollary 2.3. [34] *Let G be a finite p -group of exponent p^k . Then $C_{p^i} = 1$, for some $1 \leq i \leq k$, if and only if one of the following occurs:*

- (1) $G \cong \mathbb{Z}_{p^k}$ and $C_{p^j} = 1$ for all $1 \leq j \leq k$, or
- (2) $p = 2$ and G is isomorphic to one of the following 2-groups:

(i) *dihedral 2-group*

$$\mathbb{D}_{2^{k+1}} = \langle x, y : x^{2^k} = 1, y^2 = 1, y^{-1}xy = x^{-1} \rangle, \quad (k \geq 1)$$

where $C_2 = 1 + 2^k$ and $C_{2^j} = 1$ for all $(2 \leq j \leq k)$.

(ii) *generalized quaternion 2-group*

$$\mathbb{Q}_{2^{k+1}} = \langle x, y : x^{2^k} = 1, x^{2^{k-1}} = y^2, y^{-1}xy = x^{-1} \rangle, \quad (k \geq 2)$$

where $C_4 = 1 + 2^{k-1}$ and $C_{2^j} = 1$ for all $1 \leq j \leq k$ and $j \neq 2$.

(iii) *semi-dihedral 2-group*

$$\mathbb{SD}_{2^{k+1}} = \langle x, y : x^{2^k} = 1, y^2 = 1, y^{-1}xy = x^{-1+2^{k-1}} \rangle, \quad (k \geq 3)$$

where $C_2 = 1 + 2^{k-1}$, $C_4 = 1 + 2^{k-2}$ and $C_{2^j} = 1$ for all $3 \leq j \leq k$.

In view of Theorem 2.2 and Corollary 2.3, we have the following lemma:

Lemma 2.4. *Let G be a finite p -group with exponent p^2 and G contains exactly one cyclic subgroup of order p^2 . Then the following holds:*

- (i) *If $p = 2$, then G is isomorphic to \mathbb{Z}_4 or \mathbb{D}_8 .*
- (ii) *If $p > 2$, then G is isomorphic to \mathbb{Z}_{p^2} .*

Theorem 2.5. [12] *Let G be a finite group. Then the following statements are equivalent:*

- (i) G is a nilpotent group.
- (ii) Every Sylow subgroup of G is normal.
- (iii) G is the direct product of its Sylow subgroups.
- (iv) For $x, y \in G$, x and y commute whenever $o(x)$ and $o(y)$ are relatively primes.

Lemma 2.6. [32, Lemma 2.5] *Let G be a finite nilpotent group and $x, y \in G$ be such that $o(x) = s$ and $o(y) = t$. Then there exists an element $z \in G$ such that $o(z) = \text{lcm}(s, t)$.*

Now we recall the necessary graph theoretic definitions and notions from [14, 36]. A *graph* Γ consists of a vertex set $V(\Gamma)$ and an edge set $E(\Gamma)$, where an edge is an unordered pair of distinct vertices of Γ . If $\{u_1, u_2\}$ is an edge then, we say that u_1 is *adjacent* to u_2 and write it as $u_1 \sim u_2$. Otherwise, we denote by $u_1 \not\sim u_2$. An edge $\{u, v\}$ in a graph Γ is called a *loop* if $u = v$. A graph with no loops or multiple edges is called a *simple* graph. In this paper, we are considering only simple graphs. Let Γ be a graph. If both the sets $V(\Gamma)$ and $E(\Gamma)$ are empty, then Γ is called a *null graph*. A *subgraph* of Γ is a graph Γ' such that $V(\Gamma') \subseteq V(\Gamma)$ and $E(\Gamma') \subseteq E(\Gamma)$. A subgraph Γ' of Γ is an *induced subgraph* if two vertices of $V(\Gamma')$ are adjacent in Γ' if and only if they are adjacent in Γ . A graph is called *complete* if every pair of distinct vertices are adjacent. The complete graph on n vertices is denoted by K_n . A graph Γ is said to be k -partite if the vertex set of Γ can be partitioned into k subsets such that no two vertices in the same subset being adjacent. If $k = 2$, then Γ is called a *bipartite* graph. A *complete k -partite* graph, denoted by K_{n_1, n_2, \dots, n_k} , is a k -partite graph having its parts sizes n_1, n_2, \dots, n_k such that every vertex in each part is adjacent to all the vertices of all other parts of K_{n_1, n_2, \dots, n_k} . A *walk* λ in a graph Γ from the vertex u to the vertex v is a sequence of vertices $u = u_1, u_2, \dots, u_m = v$ ($m > 1$) such that $u_i \sim u_{i+1}$ for every $i \in \{1, 2, \dots, m-1\}$. If no edge is repeated in λ , then it is called a *trail* in Γ . If no vertex is repeated in λ , then it is called a *path* in Γ . A graph Γ is *connected* if every pair of vertices has a path in Γ . Otherwise, Γ is *disconnected*. A graph Γ is *planar* if it can be drawn on a plane without edge crossing. A planar graph is said to be *outerplanar* if it can be drawn in the plane such that all its vertices lie on the outer face. A compact connected topological space such that each point has a neighbourhood homeomorphic to an open disc is called a *surface*. A graph is said to be *embeddable* on a topological surface if it can be drawn on the surface without edge crossing. Let \mathbb{S}_g be an orientable surface with g handles, where g is a non-negative integer. The genus $\gamma(\Gamma)$ of a graph Γ , is the minimum integer g such that the graph can be embedded in \mathbb{S}_g , i.e. the graph Γ can be drawn into the surface \mathbb{S}_g with no edge crossing. Note that the graphs having genus 0 are planar, and the graphs having genus one are toroidal. Let \mathbb{N}_k be the non-orientable surface formed by connected sum of k projective planes, that is, \mathbb{N}_k is a non-orientable surface with k cross-cap. The *cross-cap* $\bar{\gamma}(\Gamma)$ of a graph

Γ , is the minimum non-negative integer k such that Γ can be embedded in \mathbb{N}_k . For instance, a graph Γ is planar if $\bar{\gamma}(\Gamma) = 0$ and, Γ is *projective-planar* if $\bar{\gamma}(\Gamma) = 1$. The following results are useful for later use.

Theorem 2.7. [36] *A graph Γ is bipartite if and only if it has no odd cycle.*

Theorem 2.8. [37] *The genus and cross-cap of the complete graphs K_n and $K_{m,n}$ are given below:*

- (i) $\gamma(K_n) = \left\lceil \frac{(n-3)(n-4)}{12} \right\rceil, n \geq 3.$
- (ii) $\gamma(K_{m,n}) = \left\lceil \frac{(m-2)(n-2)}{4} \right\rceil, m, n \geq 2.$
- (iii) $\bar{\gamma}(K_n) = \left\lceil \frac{(n-3)(n-4)}{6} \right\rceil, n \geq 3, n \neq 7; \bar{\gamma}(K_7) = 3.$
- (iv) $\bar{\gamma}(K_{m,n}) = \left\lceil \frac{(m-2)(n-2)}{2} \right\rceil, m, n \geq 2.$

Remark 2.9. Let x and y be two elements of a finite group G such that neither $o(x)|o(y)$ nor $o(y)|o(x)$. Then $x \approx y$ in $\mathcal{P}(G)$. The converse is also true if x and y belong to the same cyclic subgroup of G .

Proposition 2.10. [5, Proposition 2.1] *Let G be a non-trivial finite group and $g \in G$ be a non-identity element. Then $g \notin V(\mathcal{D}(G))$ if and only if either $\langle g \rangle$ is a maximal cyclic subgroup of G , or every cyclic subgroup of G containing g has prime-power order.*

Proposition 2.11. [5, Proposition 4.1] *Let G be a finite nilpotent group and x, y be two non-identity elements of G such that $\gcd(o(x), o(y)) = 1$, then $x \sim y$ in $\mathcal{D}(G)$.*

Proposition 2.12. [33, Corollary 3.6] *Let G be a finite nilpotent group and x, y be two non-identity elements of a Sylow subgroup of G . Then $x \approx y$ in $\mathcal{D}(G)$.*

Lemma 2.13. [5, Lemma 4.4] *Let G be a group and let H be a non-EPPO-subgroup of G . Then $\mathcal{D}(H)$ is an induced subgraph of $\mathcal{D}(G)$.*

Theorem 2.14. [33, Theorem 4.7] *Let G be a finite nilpotent group which is not a p -group. Then $\mathcal{D}(G)$ is planar if and only if G is isomorphic to one of the following groups:*

$$\mathbb{Z}_{12}, \mathbb{D}_8 \times \mathbb{Z}_3, \mathbb{Z}_2 \times \mathbb{Z}_2 \times \cdots \times \mathbb{Z}_2 \times \mathbb{Z}_3, \mathbb{Z}_2 \times Q_1, \mathbb{Z}_3 \times Q_2, \text{ where}$$

Q_i is a q_i -group of prime exponent q_i such that $q_1 > 2$ and $q_2 > 3$.

3. PROOF OF THE MAIN RESULT

In this section, we provide the proof of our main result. Let $G = P_1 P_2 \cdots P_r$, where P_i ($1 \leq i \leq r$) is a Sylow p_i -subgroup of G and $|P_i| = p_i^{\alpha_i}$, be a nilpotent group. For $x \in G$, there exists a unique element $x_i \in P_i$ for each $i \in [r] = \{1, 2, \dots, r\}$ such that $x = x_1 x_2 \cdots x_r$. Since $(x_1, x_2, \dots, x_r) \mapsto x_1 x_2 \cdots x_r$ is a group isomorphism from $P_1 \times P_2 \times \cdots \times P_r$ to $P_1 P_2 \cdots P_r$. Thus, instead of $P_1 P_2 \cdots P_r$, we sometimes write $P_1 \times P_2 \times \cdots \times P_r$ without referring to it. In order to prove Theorem 1.1, the following lemmas are useful.

Lemma 3.1. *Let $G = P_1 P_2 \cdots P_r$ ($r \geq 2$) be a finite nilpotent group. Then the subgraph of $\mathcal{D}(G)$ induced by the set $(P_1 \cup P_2 \cup \cdots \cup P_r) \setminus \{e\}$ is isomorphic to complete r -partite graph $K_{|P_1|-1, |P_2|-1, \dots, |P_r|-1}$.*

Proof. The result holds by Propositions 2.11 and 2.12. □

Lemma 3.2. *Let $G \cong P_1 \times P_2$ be a finite nilpotent group, where $\exp(P_i) = p_i$ for each $i \in \{1, 2\}$. Then $\mathcal{D}(G)$ is a complete bipartite graph isomorphic to $K_{|P_1|-1, |P_2|-1}$.*

Proof. First note that $\pi_G = \{1, p_1, p_2, p_1 p_2\}$ and the identity element of G does not belong to $V(\mathcal{D}(G))$. By Proposition 2.10, the elements of the order $p_1 p_2$ do not belong to $V(\mathcal{D}(G))$. Thus, $V(\mathcal{D}(G))$ contains the elements of orders p_1 and p_2 only. Consider the sets $A = \{x \in V(\mathcal{D}(G)) : o(x) = p_1\}$ and $B = \{y \in V(\mathcal{D}(G)) : o(y) = p_2\}$. Clearly, A and B forms a partition of $V(\mathcal{D}(G))$. By Propositions 2.11 and 2.12, $\mathcal{D}(G)$ is a complete bipartite graph which is isomorphic to $K_{|P_1|-1, |P_2|-1}$. This completes our proof. □

Lemma 3.3. *Let $G = P_1 P_2 \cdots P_r$ be a finite nilpotent group. If $r \geq 3$, then $\gamma(\mathcal{D}(G)) \geq 3$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 6$.*

Proof. Let $G = P_1 P_2 \cdots P_r$ be a nilpotent group such that $|P_i| = p_i^{\alpha_i}$ and $p_j < p_{j+1}$ for $j \in [r-1]$. We prove our result in the following two cases:

Case-1: $r \geq 4$. By Lemma 3.1, the subgraph of $\mathcal{D}(G)$ induced by the set $(P_1 \cup P_2 \cup \cdots \cup P_r) \setminus \{e\}$ contains a subgraph isomorphic to $K_{1,2,4,6}$. Note that $K_{1,2,4,6}$ has a subgraph which is isomorphic to $K_{7,6}$. By Theorem 2.8, we get $\gamma(\mathcal{D}(G)) \geq 5$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 10$.

Case-2: $r = 3$. In this case, we have $p_3 \geq 5$. Now, we discuss the following subcases:

Subcase-2.1: $p_3 = 5$. Clearly, $p_1 = 2$ and $p_2 = 3$. By Lemma 2.6, G has an element x such that $\langle x \rangle \cong \mathbb{Z}_{30}$. Note that \mathbb{Z}_{30} has 4 elements of order 10 and 8 elements of order 15. Suppose $x_i, y_j \in \langle x \rangle$ such that $o(x_i) = 10$ and $o(y_j) = 15$ for every $i \in [4], j \in [8]$. Note that $x_i \sim y_j$ in $\mathcal{P}_E(G)$, but they are not adjacent in $\mathcal{P}(G)$ (see Remark 2.9). It follows that for each $i \in [4]$ and $j \in [8]$, we have $x_i \sim y_j$ in $\mathcal{D}(G)$. Consequently, the subgraph of $\mathcal{D}(G)$ induced by the set $\{x_1, x_2, x_3, x_4, y_1, \dots, y_8\}$ is isomorphic to $K_{4,8}$. Thus, $\gamma(\mathcal{D}(G)) \geq 3$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 6$.

Subcase-2.2: $p_3 \geq 7$. In this subcase G has at least 1 element of order p_1 , 2 elements of order p_2 , 2 elements of order $p_1 p_2$ and 6 elements of order p_3 . By Proposition 2.11, the elements of order p_1, p_2 and $p_1 p_2$ will be adjacent to each element of order p_3 . Consequently, $\mathcal{D}(G)$ contains a subgraph isomorphic to $K_{5,6}$ and so $\gamma(\mathcal{D}(G)) \geq 3$, $\bar{\gamma}(\mathcal{D}(G)) \geq 6$. \square

Now we prove our main result of this manuscript.

Proof of Theorem 1.1. Let $G = P_1 P_2 \cdots P_r$ be a nilpotent group such that $|P_i| = p_i^{\alpha_i}$ and $p_j < p_{j+1}$ for $j \in [r-1]$. First, suppose that $\gamma(\mathcal{D}(G)) \leq 2$ and $\bar{\gamma}(\mathcal{D}(G)) \leq 2$. By Lemma 3.3, we obtain $r = 2$. Thus, $G = P_1 \times P_2$. We prove our result through the following cases:

Case-1: $p_1 \geq 7$. It follows that $p_2 \geq 11$. By Lemma 3.1, the subgraph of $\mathcal{D}(G)$ induced by the set $(P_1 \cup P_2) \setminus \{e\}$ has a subgraph isomorphic to $K_{6,10}$. Consequently, $\gamma(\mathcal{D}(G)) \geq 8$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 16$.

Case-2: $p_1 = 5$. Then we must have $p_2 \geq 7$.

Subcase-2.1: $|P_1| = 5$ and $|P_2| = p_2$. Then $G \cong \mathbb{Z}_5 \times \mathbb{Z}_{p_2}$. By Lemma 3.2, $\mathcal{D}(G) \cong K_{4,p_2-1}$. If $p_2 = 7$, then $\gamma(\mathcal{D}(\mathbb{Z}_{35})) = 2$ and $\bar{\gamma}(\mathcal{D}(\mathbb{Z}_{35})) = 4$. If $p_2 \geq 11$, then $\gamma(\mathcal{D}(G)) \geq 4$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 8$.

Subcase-2.2: $|P_1| = 5^\alpha$ and $|P_2| = p_2^\beta$, where both α and β are not equal to 1. If $\alpha \geq 2$, then the graph induced by the set $(P_1 \cup P_2) \setminus \{e\}$ has a subgraph isomorphic to $K_{24,6}$. Consequently, $\gamma(\mathcal{D}(G)) \geq 22$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 44$. Similarly, if $\alpha = 1$ and $\beta \geq 2$, then we get $\gamma(\mathcal{D}(G)) \geq 23$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 46$.

Case-3: $p_1 = 3$. Clearly, $p_2 \geq 5$.

Subcase-3.1: $|P_1| = 3$. Note that $\exp(P_2) = p_2^\alpha$, where $\alpha \geq 1$. If $\alpha = 1$, then by Theorem 2.14, the graph $\mathcal{D}(G)$ is planar. Consequently, $\gamma(\mathcal{D}(G)) = 0 = \bar{\gamma}(\mathcal{D}(G))$. Now we assume that $\alpha \geq 2$. Let $x \in P_1$ such that $o(x) = 3$ and $y \in P_2$ such that $o(y) = p_2^2$. Notice that $\langle xy \rangle$ is a cyclic subgroup of order $3p_2^2$ in G . Consider the sets $S = \{z \in \langle xy \rangle : o(z) = 3p_2\}$ and $T = \{z' \in \langle xy \rangle : o(z') = p_2^2\}$. Let $x' \in S$ and $y' \in T$. Clearly, $x' \sim y'$ in $\mathcal{P}_E(G)$. Also, neither $o(x')|o(y')$ nor $o(y')|o(x')$. By Remark 2.9, $x' \approx y'$ in $\mathcal{P}(G)$. Thus, $x' \sim y'$ in $\mathcal{D}(G)$. Consequently, $\mathcal{D}(G)$ contains a subgraph isomorphic to $K_{|S|,|T|}$. Since $p_2 \geq 5$, we obtain $|S| \geq 8$ and $|T| \geq 20$. Thus, $\gamma(\mathcal{D}(G)) \geq 27$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 54$.

Subcase-3.2: $|P_1| = 3^\alpha$, where $\alpha \geq 2$. Notice that P_1 has at least 8 non-identity elements and P_2 has at least 4 non-identity elements. By Lemma 3.1, the graph induced by $(P_1 \cup P_2) \setminus \{e\}$ contains a subgraph isomorphic to $K_{8,4}$. Consequently, $\gamma(\mathcal{D}(G)) \geq 3$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 6$.

Case-4: $p_1 = 2$. Now, we have the following possible subcases.



Subcase-4.1.1: $|P_1| = 2$ and $\exp(P_2) = p_2$. By Theorem 2.14, the graph $\mathcal{D}(G)$ is planar. Consequently, $\gamma(\mathcal{D}(G)) = 0 = \bar{\gamma}(\mathcal{D}(G))$.

Subcase-4.1.2: $|P_1| = 2, |P_2| = 3^\alpha$, where $\alpha \geq 2$ and $\exp(P_2) = 9$. In view of Lemma 2.1, we have the following two further subcases:

Subcase-4.1.2(a): P_2 contains exactly one cyclic subgroup of order 9. By Lemma 2.4, we get $G \cong \mathbb{Z}_{18}$. Observe that the subgraph of $\mathcal{D}(\mathbb{Z}_{18})$ induced by the set $\{3, 9, 15, 2, 4, 8, 10, 14, 16\}$ is isomorphic to $K_{3,6}$. Thus, $\gamma(\mathcal{D}(G)) \geq 1$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 2$. A genus 1 and cross-cap 2 drawings of $\mathcal{D}(\mathbb{Z}_{18})$ are given in Figures 1 and 2, respectively.

Subcase-4.1.2(b): P_2 contains two cyclic subgroups H and K of order 9 such that $|H \cap K| = 3$. Let $P_1 = \langle x \rangle$, $H = \langle y \rangle$ and $K = \langle z \rangle$. Suppose $x_1, x_2 \in H \cap K$ such that $o(x_1) = o(x_2) = 3$. Then

$$P_1 H = \{e, x, x_1, x_2, y, y^2, y^4, y^5, y^7, y^8, x x_1, x x_2, x y, x y^2, x y^4, x y^5, x y^7, x y^8\}, \text{ and}$$

FIGURE 1. Embedding of $\mathcal{D}(\mathbb{Z}_{18})$ in \mathbb{S}_1 .FIGURE 2. Embedding of $\mathcal{D}(\mathbb{Z}_{18})$ in \mathbb{N}_2 .

$$P_1K = \{e, x, x_1, x_2, z, z^2, z^4, z^5, z^7, z^8, xx_1, xx_2, xz, xz^2, xz^4, xz^5, xz^7, xz^8\}$$

are maximal cyclic subgroups of order 18 (see [10, Lemma 2.11]). Consider the sets $S = \{x, xx_1, xx_2\}$ and $T = \{y, y^2, y^4, y^5, y^7, y^8, z, z^2, z^4, z^5, z^7, z^8\}$. Let $x' \in S$ and $y' \in T$. Notice that $x' \sim y'$ in $\mathcal{P}_E(G)$. Also, neither $o(x')|o(y')$ nor $o(y')|o(x')$. It follows that $x' \approx y'$ in $\mathcal{P}(G)$ and so $x' \sim y'$ in $\mathcal{D}(G)$. Thus, the subgraph induced by $S \cup T$ has a subgraph isomorphic to $K_{3,12}$. It implies that $\gamma(\mathcal{D}(G)) \geq 3$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 5$.

Subcase-4.1.3: $|P_1| = 2$ and $\exp(P_2) = 3^\alpha$, where $\alpha \geq 3$. Then there exists an element $y \in P_2$ such that $o(y) = 27$. Let $P_1 = \langle x \rangle$. Observe that $\langle xy \rangle$ is a cyclic subgroup of order 54 in G . Consider the sets $S = \{s \in \langle xy \rangle : o(s) = 18\}$ and $T = \{t \in \langle xy \rangle : o(t) = 27\}$. Suppose $x' \in S$ and $y' \in T$. Observe that $x' \sim y'$ in $\mathcal{P}_E(G)$ and $x' \approx y'$ in $\mathcal{P}(G)$ (cf. Remark 2.9). It follows that $x' \sim y'$ in $\mathcal{D}(G)$. Thus, $\mathcal{D}(G)$ has a subgraph isomorphic to $K_{|S|,|T|}$. Since $|S| = 6$ and $|T| = 18$, we obtain $\gamma(\mathcal{D}(G)) \geq 16$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 32$.

Subcase-4.1.4: $|P_1| = 2$ and $\exp(P_2) = p_2^\alpha$, where $p_2 \geq 5$, $\alpha \geq 2$. Then there exists an element $y \in P_2$ such that $o(y) = p_2^2$. Let $P_1 = \langle x \rangle$. Observe that $\langle xy \rangle$ is a cyclic subgroup of order $2p_2^2$ in G . Consider the sets $S = \{s \in \langle xy \rangle : o(s) = 2p_2\}$ and $T = \{t \in \langle xy \rangle : o(t) = p_2^2\}$. Similar to Subcase 4.1.3, we obtain a subgraph of $\mathcal{D}(G)$ which is isomorphic to $K_{|S|,|T|}$. Since $p_2 \geq 5$, we have $|S| \geq 4$ and $|T| \geq 20$. It follows that $\gamma(\mathcal{D}(G)) \geq 9$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 18$.

Subcase-4.2.1: $|P_1| = 4$ and $|P_2| = 3$. Then either $G \cong \mathbb{Z}_4 \times \mathbb{Z}_3$ or $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3$. By Theorem 2.14, in both of these cases, $\mathcal{D}(G)$ is a planar graph. Consequently, $\gamma(\mathcal{D}(G)) = 0 = \bar{\gamma}(\mathcal{D}(G))$.

Subcase-4.2.2: $|P_1| = 4$ and $|P_2| = 5$. Then either $G \cong \mathbb{Z}_4 \times \mathbb{Z}_5 \cong \mathbb{Z}_{20}$ or $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_5$. If $G \cong \mathbb{Z}_{20}$, then by Theorem 2.14, $\gamma(\mathcal{D}(G)) \geq 1$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 1$. A genus 1 and cross-cap 1 drawings of $\mathcal{D}(\mathbb{Z}_{20})$ are given in Figures 3 and 4, respectively.




FIGURE 3. Embedding of $\mathcal{D}(\mathbb{Z}_{20})$ in \mathbb{S}_1 .






FIGURE 4. Embedding of $\mathcal{D}(\mathbb{Z}_{20})$ in \mathbb{N}_1 .

If $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_5$, then by Lemma 2.1, $\mathcal{D}(G) \cong K_{3,4}$. Consequently, $\gamma(\mathcal{D}(G)) = 1$ and $\bar{\gamma}(\mathcal{D}(G)) = 1$.

Subcase-4.2.3: $|P_1| = 4$ and $|P_2| = 7$. Then G is isomorphic to $\mathbb{Z}_4 \times \mathbb{Z}_7$ or $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_7$. If $G \cong \mathbb{Z}_4 \times \mathbb{Z}_7$, then $\mathcal{D}(G)$ contains a subgraph which is isomorphic to $K_{3,6}$ (see Lemma 3.2). Consequently, $\gamma(\mathcal{D}(G)) \geq 1$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 2$. A genus 1 and cross-cap 2 drawings of $\mathcal{D}(\mathbb{Z}_{28})$ are given in Figure 5 and 6, respectively.

FIGURE 5. Embedding of $\mathcal{D}(\mathbb{Z}_{28})$ in \mathbb{S}_1 .FIGURE 6. Embedding of $\mathcal{D}(\mathbb{Z}_{28})$ in \mathbb{N}_2 .

If $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_7$, then by Lemma 3.2, $\mathcal{D}(G) \cong K_{3,6}$. It follows that $\gamma(\mathcal{D}(G)) = 1$ and $\bar{\gamma}(\mathcal{D}(G)) = 2$.

Subcase-4.2.4: $|P_1| = 4$ and $|P_2| = 9$. Then either $P_1 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ or $P_1 \cong \mathbb{Z}_4$. Also, P_2 is isomorphic to $\mathbb{Z}_3 \times \mathbb{Z}_3$ or \mathbb{Z}_9 . Consequently, G is isomorphic to one of the groups: \mathbb{Z}_{36} , $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_9$, $\mathbb{Z}_4 \times \mathbb{Z}_3 \times \mathbb{Z}_3$, $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3$. If $G \cong \mathbb{Z}_{36}$, then $\mathcal{P}_E(G)$ is a complete graph. Consider the set $S = \{x \in G : o(x) = 9\}$ and $T = \{y \in G : o(y) \in \{2, 4, 6\}\}$. Let $x' \in S$ and $y' \in T$. Clearly, $x' \sim y'$ in $\mathcal{P}_E(G)$ and by Remark 2.9, $x' \approx y'$ in $\mathcal{P}(G)$. Thus, $x' \sim y'$ in $\mathcal{D}(G)$ and so $\mathcal{D}(G)$ contains a subgraph which is isomorphic to $K_{|S|,|T|}$. Since $|S| = 6$ and $|T| = 5$, we obtain $\gamma(\mathcal{D}(G)) \geq 3$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 6$.

If $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_9$, then note that G has exactly 3 maximal cyclic subgroups $H_1 = \langle(1, 0, 1)\rangle$, $H_2 = \langle(0, 1, 1)\rangle$ and $H_3 = \langle(1, 1, 1)\rangle$. Consider the set $S = \{x \in G : o(x) = 9\}$ and $T = \{y \in G : o(y) \in \{2, 6\}\}$. Then observe that $S \subseteq (H_1 \cap H_2 \cap H_3)$, $|S| = 6$ and $|T| = 9$. Suppose $x' \in S$ and $y' \in T$ are arbitrary elements. Then $x', y' \in H_i$ for some $i \in \{1, 2, 3\}$. It follows that $x' \sim y'$ in $\mathcal{P}_E(G)$. Consequently, $x' \sim y'$ in $\mathcal{D}(G)$ (cf. Remark 2.9). Thus, $\mathcal{D}(G)$ contains a subgraph isomorphic to $K_{6,9}$. Hence, $\gamma(\mathcal{D}(G)) \geq 7$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 14$.

If $G \cong \mathbb{Z}_4 \times \mathbb{Z}_3 \times \mathbb{Z}_3$, then by Lemma 3.1, $\mathcal{D}(G)$ contains a subgraph which is isomorphic to $K_{3,8}$. Thus, $\gamma(\mathcal{D}(G)) \geq 2$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 3$. A genus 2 drawing of $\mathcal{D}(G)$ is given in Figure 7.

If $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3$ then by Lemma 3.2, we have $\mathcal{D}(G) \cong K_{3,8}$. Consequently, $\gamma(\mathcal{D}(G)) = 2$ and $\bar{\gamma}(\mathcal{D}(G)) = 3$.

Subcase-4.2.5: $|P_1| = 4$ and $|P_2| = 11$. Thus, either $G \cong \mathbb{Z}_4 \times \mathbb{Z}_{11} \cong \mathbb{Z}_{44}$ or $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_{11}$. If $G \cong \mathbb{Z}_4 \times \mathbb{Z}_{11}$, then the subgraph of $\mathcal{D}(G)$ induced by the set $(P_1 \cup P_2) \setminus \{e\}$ contains a subgraph isomorphic to $K_{3,10}$. Consequently, $\gamma(\mathcal{D}(G)) \geq 2$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 4$. A genus 2 drawing of $\mathcal{D}(\mathbb{Z}_{44})$ is given in Figure 8.

If $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_{11}$, then $\mathcal{D}(G) \cong K_{3,10}$ (cf. Lemma 3.2) and so $\gamma(\mathcal{D}(G)) = 2$, $\bar{\gamma}(\mathcal{D}(G)) = 4$.

Subcase-4.2.6: $|P_1| = 4$ and $|P_2| = p_2$, where $p_2 \geq 13$. By Lemma 3.1, $\mathcal{D}(G)$ contains a subgraph isomorphic to $K_{3,12}$. Thus, $\gamma(\mathcal{D}(G)) \geq 3$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 5$.




FIGURE 7. Embedding of $\mathcal{D}(\mathbb{Z}_4 \times \mathbb{Z}_3 \times \mathbb{Z}_3)$ in \mathbb{S}_2 .




FIGURE 8. Embedding of $\mathcal{D}(\mathbb{Z}_{44})$ in \mathbb{S}_2 .

Subcase-4.2.7: $|P_1| = 4$ and $|P_2| = p_2^\alpha$, where $p_2 \geq 5$, $\alpha \geq 2$ and $\exp(P_2) = p_2$. By Lemma 3.1, $\mathcal{D}(G)$ contains a subgraph isomorphic to $K_{3,24}$. Thus, $\gamma(\mathcal{D}(G)) \geq 6$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 11$.

Subcase-4.2.8: $|P_1| = 4$ and $|P_2| = 3^\alpha$, where $\alpha \geq 3$. It implies that the minimum number of non-identity elements in P_1 and P_2 are 3 and 26, respectively. Consequently, by Lemma 3.1, $\mathcal{D}(G)$ contains a subgraph isomorphic to $K_{3,26}$. It follows that $\gamma(\mathcal{D}(G)) \geq 6$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 12$.

Subcase-4.3.1: $|P_1| = 2^\alpha$ and $|P_2| = 3^\beta$, where $\alpha \geq 3$, $\beta \geq 2$. By Lemma 3.1, the subgraph induced by the set $(P_1 \cup P_2) \setminus \{e\}$ contains a subgraph isomorphic to $K_{7,8}$. It follows that $\gamma(\mathcal{D}(G)) \geq 8$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 15$.

Subcase-4.3.2: $|P_1| = 2^\alpha$ ($\alpha \geq 3$) with $\exp(P_1) = 2$ and $|P_2| = 3$. By Theorem 2.14, the graph $\mathcal{D}(G)$ is planar and so $\gamma(\mathcal{D}(G)) = 0 = \bar{\gamma}(\mathcal{D}(G))$.

Subcase-4.3.3: $|P_1| = 2^\alpha$ ($\alpha \geq 3$) with $\exp(P_1) = 4$ and $|P_2| = 3$. Consider $P_2 = \langle x \rangle$. Further, suppose that P_1 has t (≥ 1) maximal cyclic subgroups of order 4 and s (≥ 0) maximal cyclic subgroups of order 2. Consider the maximal cyclic subgroups $M_i = \langle y_i \rangle$, where $1 \leq i \leq t$, of order 4. If $s \geq 1$, then consider $M'_j = \langle z_j \rangle$, where $1 \leq j \leq s$, as the maximal cyclic subgroup of order 2. Consequently, maximal cyclic subgroups of order 12 in G are of the form $M_i P_2 = \{e, x, x^2, y_i, y_i^2, y_i^3, y_i x, y_i^2 x, y_i^3 x, y_i x^2, y_i^2 x^2, y_i^3 x^2\}$. Also, maximal cyclic subgroups of order 6 are of the form $M'_j P_2 = \{e, x, x^2, z_j, z_j x, z_j x^2\}$ (see [10, Lemma 2.11]). Notice that P_2 contained in every maximal cyclic subgroup of G . Also, the identity element e and the generators of $M_i P_2$ and $M'_j P_2$ do not belong to the vertex set of $\mathcal{D}(G)$ (cf. Proposition 2.10).

If $t = 1$, then by Lemma 2.4, we have $P_1 = \mathbb{D}_8$. Thus, $G \cong \mathbb{D}_8 \times \mathbb{Z}_3$. By Theorem 2.14, $\mathcal{D}(G)$ becomes planar and so $\gamma(\mathcal{D}(G)) = 0 = \bar{\gamma}(\mathcal{D}(G))$. We may now suppose that $t > 1$. In view of Lemma 2.1, now we discuss this subcase into the following six further subcases.

Subcase-4.3.3(a): P_1 contains two maximal cyclic subgroups H and K of order 4 such that $|H \cap K| = 2$, and the intersection of any other pair of maximal cyclic subgroups of P_1 is trivial. Without loss of generality, assume that $H = M_1$ and $K = M_2$. Since $|M_1 \cap M_2| = 2$, we get $M_1 P_2 \cap M_2 P_2 = \{e, x, x^2, y_1^2, y_1^2 x, y_1^2 x^2\}$. Consider the set $S = \{y \in M_1 P_2 \cup M_2 P_2 : o(y) = 4\}$ and $T = \{z \in M_1 P_1 \cup M_2 P_2 : o(z) \in \{3, 6\}\}$. Let $x' \in S$ and $y' \in T$. Then $x', y' \in M_i P_2$ for some $i \in \{1, 2\}$. Thus, $x' \sim y'$ in $\mathcal{P}_E(G)$. Consequently, $x' \sim y'$ in $\mathcal{D}(G)$ (cf. Remark 2.9). Since $|S| = 4$ and $|T| = 4$, the subgraph induced by $S \cup T$ is isomorphic to $K_{4,4}$. Thus, $\gamma(\mathcal{D}(G)) \geq 1$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 2$. A genus 1 and cross-cap 2 drawings of $\mathcal{D}(G)$ are given in Figures 9 and 10, respectively. The graph G_3 , given in Figure 11, can be inserted in the face F of Figure 9 and 10, respectively, without edge crossings.




FIGURE 9. Embedding of $\mathcal{D}(P_1 \times P_2)$, where P_1 and P_2 are according to Subcase-4.3.3(a), in \mathbb{S}_1 .

Subcase-4.3.3(b): P_1 contains four maximal cyclic subgroups H_1, H_2, H_3 and H_4 of order 4 such that $|H_1 \cap H_2| = |H_3 \cap H_4| = 2$, and the intersection of any other pair of maximal cyclic subgroups of P_1 is trivial. Without loss of generality, assume that $H_i = M_i$ for $1 \leq i \leq 4$. Now similar to the Subcase-4.3.3(a), we get a subgraph Γ' of $\mathcal{D}(G)$, which is isomorphic to $K_{4,4}$. Moreover, $|M_3 \cap M_4| = 2$. It implies that $M_3 P_2 \cap M_4 P_2 = \{e, x, x^2, y_3^2, y_3^2 x, y_3^2 x^2\}$. Now to embed $\mathcal{D}(G)$ through Γ' in \mathbb{S}_1 , first we insert the vertices $y_3, y_3^3, y_4, y_4^3, y_3^2 x, y_3^2 x^2$ and their incident edges. Since $y_3 \sim y_3^2 x \sim y_3^3 \sim y_3^2 x^2 \sim y_4$ and $y_3^2 x \sim y_4^3 \sim y_3^2 x^2$; all these vertices must be inserted in the same face F' . Note that the vertices y_3, y_3^3, y_4 and y_4^3 are adjacent to both the vertices x and x^2 (see Proposition 2.10). Consequently, the face F' must contain the vertices x and x^2 . After inserting the vertices y_3, y_3^3, y_4, y_4^3 and their incident edges, it is impossible to insert the vertices $y_3^2 x, y_3^2 x^2$ without edge crossing (see Figure 13). Thus, $\gamma(\mathcal{D}(G)) \geq 2$. A genus 2 drawing of $\mathcal{D}(G)$ is given in Figure 12, and the subgraph G_5 , (given in Figure 11) can be inserted in the face F . By the similar argument used earlier in this subcase, any embedding of $\mathcal{D}(G)$ in \mathbb{N}_2 is also not possible without edge crossings. Hence, $\bar{\gamma}(\mathcal{D}(G)) \geq 3$.




FIGURE 10. Embedding of $\mathcal{D}(P_1 \times P_2)$, where P_1 and P_2 are according to Subcase-4.3.3(a), in \mathbb{N}_2 .






FIGURE 11. The subgraph G_i of $\mathcal{D}(G)$ induced by the set $(V(\mathcal{D}(G)) \setminus (\bigcup_{j=1}^{i-1} M_j P_2)) \cup \{x, x^2\}$.

Subcase-4.3.3(c): P_1 contains six maximal cyclic subgroups H_1, H_2, H_3, H_4, H_5 and H_6 of order 4 such that $|H_1 \cap H_2| = |H_3 \cap H_4| = |H_5 \cap H_6| = 2$, and the intersection of any three maximal cyclic subgroups of P_1 is trivial. Without loss of generality, assume that $H_i = M_i$ for each $i \in [6]$. Similar to the Subcase-4.3.3(b), we obtain a subgraph Γ'' of $\mathcal{D}(G)$, induced by the set $V(\mathcal{D}(G)) \cap (M_1 P_2 \cup M_2 P_2 \cup M_3 P_2 \cup M_4 P_2)$, which can not be embedded in \mathbb{S}_1 and \mathbb{N}_2 . It follows that $\gamma(\mathcal{D}(G)) \geq 2$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 3$. Also, we have $M_5 P_2 \cap M_6 P_2 = \{e, x, x^2, y_5^2, y_5^2 x, y_5^2 x^2\}$. Suppose $\gamma(\mathcal{D}(G)) = 2$. To embed $\mathcal{D}(G)$ in \mathbb{S}_2 , first we insert the vertices $y_5, y_5^3, y_6, y_6^3, y_5^2 x, y_5^2 x^2$ and their incident edges in genus 2 drawing of Γ' . By the similar argument used in Subcase-4.3.3(b) (by taking y_5 in place of y_3 , and y_4 in place of y_6) it is impossible to insert these vertices in \mathbb{S}_2 without edge crossings. Thus, $\gamma(\mathcal{D}(G)) \geq 3$.

FIGURE 12. Embedding of $\mathcal{D}(P_1 \times P_2)$, where P_1 and P_2 are according to Subcase-4.3.3(b), in \mathbb{S}_2 .FIGURE 13. The face F'

Subcase-4.3.3(d): P_1 contains three maximal cyclic subgroups H_1, H_2 and H_3 of order 4 such that $|H_1 \cap H_2 \cap H_3| = 2$, and the intersection of any other pair of maximal cyclic subgroups of P_1 is trivial. Without loss of generality, assume that $H_i = M_i$ for $1 \leq i \leq 3$. Since $|M_1 \cap M_2 \cap M_3| = 2$, we obtain $M_1P_2 \cap M_2P_2 \cap M_3P_2 = \{e, x, x^2, y_1^2, y_1^2x, y_1^2x^2\}$. Consider the set $S = \{y \in M_1P_2 \cup M_2P_2 \cup M_3P_2 : o(y) = 4\}$ and $T = \{z \in M_1P_2 \cup M_2P_2 \cup M_3P_2 : o(z) \in \{3, 6\}\}$. Note that the subgraph Γ' , induced by the set $S \cup T$ is isomorphic to $K_{6,4}$. Thus, $\bar{\gamma}(\mathcal{D}(G)) \geq 4$ and $\gamma(\mathcal{D}(G)) \geq 2$. Now we show that $\mathcal{D}(G)$ can be embedded in \mathbb{S}_2 without edge crossing. Since Γ' is a bipartite graph, it implies that each face of Γ' is of even length at least 4 in \mathbb{S}_2 . Consequently, each face must contain at least two vertices of each partite set of Γ' . It follows that there exists an embedding of Γ' in \mathbb{S}_2 such that the face F_2 contains the vertices x and x^2 . Now one can embed $\mathcal{D}(G)$ in \mathbb{S}_2 through Γ' by inserting the subgraph G_4 into F_2 . Therefore, $\gamma(\mathcal{D}(G)) = 2$.

Subcase-4.3.3(e): P_1 contains five maximal cyclic subgroups H_1, H_2, H_3, H_4 and H_5 of order 4 such that $|H_1 \cap H_2 \cap H_3| = 2 = |H_4 \cap H_5|$, and the intersection of any four maximal cyclic subgroups of P_1 is trivial. Without loss of generality, assume that $H_i = M_i$ for each $i \in [5]$. Similar to the Subcase-4.3.3(d), we obtain a subgraph Γ' of $\mathcal{D}(G)$, induced by the set $V(\mathcal{D}(G)) \cap (M_1P_2 \cup M_2P_2 \cup M_3P_2)$, which can not be embedded in \mathbb{S}_1 and \mathbb{N}_3 without edge crossing. It follows that $\gamma(\mathcal{D}(G)) \geq 2$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 3$. Moreover, $|M_4 \cap M_5| = 2$ implies that $M_4P_2 \cap M_5P_2 = \{e, x, x^2, y_4^2, y_4^2x, y_4^2x^2\}$. Suppose $\gamma(\mathcal{D}(G)) = 2$. Now to embed $\mathcal{D}(G)$ in \mathbb{S}_2 , first we insert the

vertices $y_4, y_4^3, y_5, y_5^3, y_4^2x, y_4^2x^2$ and their incident edges in genus 2 drawing of Γ' . By the similar argument used in the Subcase-4.3.3(b) (by taking y_4 in place of y_3 , and y_5 in place of y_6) it is impossible to insert these vertices in \mathbb{S}_2 without edge crossings. It follows that $\gamma(\mathcal{D}(G)) \geq 3$.

Subcase-4.3.3(f): P_1 contains four maximal cyclic subgroups H_1, H_2, H_3 and H_4 of order 4 such that $|H_1 \cap H_2 \cap H_3 \cap H_4| = 2$. Without loss of generality, assume that $H_i = M_i$ for each $i \in [4]$. Since $|M_1 \cap M_2 \cap M_3 \cap M_4| = 2$, we obtain $M_1P_2 \cap M_2P_2 \cap M_3P_2 \cap M_4P_2 = \{e, x, x^2, y_1^2, y_1^2x, y_1^2x^2\}$. Consider the set $S = \{y \in M_1P_2 \cup M_2P_2 \cup M_3P_2 \cup M_4P_2 : o(y) = 4\}$ and $T = \{z \in M_1P_2 \cup M_2P_2 \cup M_3P_2 \cup M_4P_2 : o(z) \in \{3, 6\}\}$. Note that the subgraph of $\mathcal{D}(G)$ induced by $S \cup T$ has a subgraph isomorphic to $K_{8,4}$. Thus, $\bar{\gamma}(\mathcal{D}(G)) \geq 3$ and $\gamma(\mathcal{D}(G)) \geq 6$.

Subcase-4.3.4: $|P_1| = 2^\alpha$ and $|P_2| = 3$, where $\alpha \geq 3$ and $\exp(P_1) \geq 8$. Then there exists an element $y \in P_2$ such that $o(y) = 8$. Suppose $x \in P_1$ such that $o(x) = 3$. Notice that $\langle xy \rangle$ is a cyclic subgroup of order 24 in G . Consider the sets $S = \{s \in \langle xy \rangle : o(s) = 8\}$ and $T = \{t \in \langle xy \rangle : o(t) \in \{3, 6, 12\}\}$. Let $x' \in S$ and $y' \in T$. Observe that, $x' \sim y'$ in $\mathcal{P}_E(G)$. Consequently, by Remark 2.9, $x' \sim y'$ in $\mathcal{D}(G)$. Thus, $\mathcal{D}(G)$ contains a subgraph isomorphic to $K_{|S|,|T|}$. Since $|S| = 4$ and $|T| = 8$, we obtain $\gamma(\mathcal{D}(G)) \geq 3$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 6$.

Subcase-4.3.5: $|P_1| = 2^\alpha$ and $|P_2| = p_2$ such that $\alpha \geq 3$, $p_2 \geq 5$. Then P_1 has at least 7 non-identity elements and P_2 has at least 4 non-identity elements. Consequently, by Lemma 3.1, $\mathcal{D}(G)$ contains a subgraph which is isomorphic to $K_{4,7}$. It follows that $\gamma(\mathcal{D}(G)) \geq 3$ and $\bar{\gamma}(\mathcal{D}(G)) \geq 5$.

This completes our proof.

DECLARATIONS

Funding: The first author gratefully acknowledge for providing financial support to CSIR (09/719(0110)/2019-EMR-I) government of India. The second author wishes to acknowledge the support of Core Research Grant (CRG/2022/001142) funded by SERB.

Conflicts of interest/Competing interests: There is no conflict of interest regarding the publishing of this paper.

Availability of data and material (data transparency): Not applicable.

Code availability (software application or custom code): Not applicable.

REFERENCES

- [1] G. Aalipour, S. Akbari, P. J. Cameron, R. Nikandish, and F. Shaveisi. On the structure of the power graph and the enhanced power graph of a group. *Electron. J. Combin.*, 24(3):3.16, 18, 2017.
- [2] M. Afkhami, M. Farrokhi D. G., and K. Khashyarmansh. Planar, toroidal, and projective commuting and noncommuting graphs. *Comm. Algebra*, 43(7):2964–2970, 2015.
- [3] S. Bera and A. K. Bhuniya. On enhanced power graphs of finite groups. *J. Algebra Appl.*, 17(8):1850146, 2018.
- [4] J. G. Berkovič. p -groups of finite order. *Sibirsk. Mat. Ž.*, 9:1284–1306, 1968.
- [5] S. Biswas, P. J. Cameron, A. Das, and H. K. Dey. On difference of enhanced power graph and power graph of a finite group. arXiv:2206.12422, 2022.
- [6] J. R. Britnell and N. Gill. Perfect commuting graphs. *J. Group Theory*, 20(1):71–102, 2017.
- [7] P. J. Cameron. The power graph of a finite group, II. *J. Group Theory*, 13(6):779–783, 2010.
- [8] P. J. Cameron. Graphs defined on groups. *Int. J. Group Theory*, 11(2):53–107, 2022.
- [9] P. J. Cameron and S. Ghosh. The power graph of a finite group. *Discrete Math.*, 311(13):1220–1222, 2011.
- [10] S. Chattopadhyay, K. L. Patra, and B. K. Sahoo. Minimal cut-sets in the power graphs of certain finite non-cyclic groups. *Comm. Algebra*, 49(3):1195–1211, 2021.
- [11] A. Doostabadi and M. Farrokhi D. G. Embeddings of (proper) power graphs of finite groups. *Algebra Discrete Math.*, 24(2):221–234, 2017.
- [12] D. S. Dummit and R. M. Foote. *Abstract algebra*. Prentice Hall, Inc., Englewood Cliffs, NJ, 1991.
- [13] M. Feng, X. Ma, and K. Wang. The structure and metric dimension of the power graph of a finite group. *European Journal of Combinatorics*, 43:82–97, 2015.
- [14] C. Godsil and G. Royle. *Algebraic graph theory*, volume 207 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2001.
- [15] S. Haji and S. M. Jafarian Amiri. On groups covered by finitely many centralizers and domination number of the commuting graphs. *Comm. Algebra*, 47(11):4641–4653, 2019.
- [16] U. Hayat, M. Umer, I. Gutman, B. Davvaz, and A. Nolla de Celis. A novel method to construct NSSD molecular graphs. *Open Math.*, 17(1):1526–1537, 2019.
- [17] A. Iranmanesh and A. Jafarzadeh. On the commuting graph associated with the symmetric and alternating groups. *J. Algebra Appl.*, 7(1):129–146, 2008.
- [18] I. M. Isaacs. *Character theory of finite groups*. AMS Chelsea Publishing, Providence, RI, 2006. Corrected reprint of the 1976 original.
- [19] A. V. Kelarev. *Graph algebras and automata*, volume 257. Marcel Dekker, Inc., New York, 2003.

- [20] A. V. Kelarev. Labelled Cayley graphs and minimal automata. *Australas. J. Combin.*, 30:95–101, 2004.
- [21] A. V. Kelarev and S. Quinn. A combinatorial property and power graphs of semigroups. *Comment. Math. Univ. Carolin.*, 45(1):1–7, 2004.
- [22] A. V. Kelarev, J. Ryan, and J. Yearwood. Cayley graphs as classifiers for data mining: the influence of asymmetries. *Discrete Math.*, 309(17):5360–5369, 2009.
- [23] A. Kulakoff. Über die Anzahl der eigentlichen Untergruppen und der Elemente von gegebener Ordnung in p -Gruppen. *Math. Ann.*, 104(1):778–793, 1931.
- [24] A. Kumar, L. Selvaganesh, P. J. Cameron, and T. Tamizh Chelvam. Recent developments on the power graph of finite groups—a survey. *AKCE Int. J. Graphs Comb.*, 18(2):65–94, 2021.
- [25] J. Kumar, S. Dalal, and V. Baghel. On the commuting graph of semidihedral group. *Bull. Malays. Math. Sci. Soc.*, 44(5):3319–3344, 2021.
- [26] X. Ma, M. Feng, and K. Wang. The rainbow connection number of the power graph of a finite group. *Graphs and Combinatorics*, 32(4):1495–1504, 2016.
- [27] X. Ma, A. Kelarev, Y. Lin, and K. Wang. A survey on enhanced power graphs of finite groups. *Electron. J. Graph Theory Appl. (EJGTA)*, 10(1):89–111, 2022.
- [28] X. Ma, G. L. Walls, and K. Wang. Power graphs of (non)orientable genus two. *Comm. Algebra*, 47(1):276–288, 2019.
- [29] A. Mahmoudifar and A. R. Moghaddamfar. Commuting graphs of groups and related numerical parameters. *Comm. Algebra*, 45(7):3159–3165, 2017.
- [30] G. A. Miller. An Extension of Sylow’s Theorem. *Proc. London Math. Soc. (2)*, 2:142–143, 1905.
- [31] M. Mirzargar, A. R. Ashrafi, and M. J. Nadjafi-Arani. On the power graph of a finite group. *Filomat*, 26(6):1201–1208, 2012.
- [32] Parveen, S. Dalal, and J. Kumar. Lambda number of the enhanced power graph of a finite group. arXiv:2208.00611, 2022.
- [33] Parveen, J. Kumar, and R. P. Panda. On the difference graph of power graphs of finite groups. arXiv:2212.07705, 2022.
- [34] S. Sarkar and M. Mishra. The lambda number of the power graph of a finite p -group. *J. Algebraic Combin.*, pages 1–10, 2022.
- [35] C. Thomassen. The graph genus problem is NP-complete. *J. Algorithms*, 10(4):568–576, 1989.
- [36] D. B. West. *Introduction to graph theory*. Prentice Hall, Inc., Upper Saddle River, NJ, 1996.
- [37] A. T. White. *Graphs, groups and surfaces*, volume 8 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, second edition, 1984.
- [38] L. Zhai, X. Ma, Y. Shao, and G. Zhong. Metric and strong metric dimension in commuting graphs of finite groups. *Comm. Algebra*, pages 1–11, 2022.

Parveen¹, Jitender Kumar¹

Addresses:

¹DEPARTMENT OF MATHEMATICS, BIRLA INSTITUTE OF TECHNOLOGY AND SCIENCE PILANI, PILANI 333031, INDIA
 Email address: p.parveenkumar144@gmail.com, jitenderarora09@gmail.com