

# STRONG EDGE-COLORING OF 2-DEGENERATE GRAPHS

GEXIN YU<sup>1</sup> AND RACHEL YU<sup>2</sup>

<sup>1</sup>*Department of Mathematics, William & Mary, Williamsburg, VA 23185, USA.*

<sup>2</sup>*Jamestown High School, Williamsburg, VA 23185, USA.*

ABSTRACT. A *strong edge-coloring* of a graph  $G$  is an edge-coloring in which every color class is an induced matching, and the strong chromatic index  $\chi'_s(G)$  is the minimum number of colors needed in strong edge-colorings of  $G$ . A graph is *2-degenerate* if every subgraph has minimum degree at most 2. Choi, Kim, Kostochka, and Raspaud (2016) showed  $\chi'_s(G) \leq 5\Delta + 1$  if  $G$  is a 2-degenerate graph with maximum degree  $\Delta$ . In this article, we improve it to  $\chi'_s(G) \leq 5\Delta - \Delta^{1/2-\epsilon} + 2$  when  $\Delta \geq 4^{1/(2\epsilon)}$  for any  $0 < \epsilon \leq 1/2$ .

## 1. INTRODUCTION

A *strong edge-coloring* of a graph  $G$  is an edge-coloring in which every color class is an induced matching; that is, there are no 2-edge-colored triangles or paths of three edges. The *strong chromatic index*  $\chi'_s(G)$  is the minimum number of colors in a strong edge-coloring of  $G$ . This notion was introduced by Fouquet and Jolivet [9] and one of the main open problems was proposed by Erdős and Nešetřil [8] during a seminar in Prague:

**Conjecture 1** (Erdős and Nešetřil, 1985). *If  $G$  is a simple graph with maximum degree  $\Delta$ , then  $\chi'_s(G) \leq 5\Delta^2/4$  if  $\Delta$  is even, and  $\chi'_s(G) \leq (5\Delta^2 - 2\Delta + 1)/4$  if  $\Delta$  is odd.*

This conjecture is true for  $\Delta \leq 3$ , see [1, 10]. For  $\Delta = 4$ , Huang, Santana and Yu [11] showed that  $\chi'_s(G) \leq 21$ , one more than the conjectured upper bound 20. Chung, Gyárfás, Trotter, and Tuza (1990, [6]) confirmed the conjecture for  $2K_2$ -free graphs. Using probabilistic methods, Molloy and Reed [14], Bruhn and Joos [3], Bonamy, Perrett, and Postle [2], and recently Hurley, Verclos, and Kang [12] showed that  $\chi'_s(G) \leq 1.772\Delta^2$  for sufficiently large  $\Delta$ .

Sparse graphs have also attracted a lot of attention. Interested readers may see the survey paper [7] for more details. In this article, we are interested in  $k$ -degenerate graphs when  $k = 2$ . A graph is  *$k$ -degenerate* if every subgraph has minimum degree at most  $k$ .

Let  $G$  be a 2-degenerate graph with maximum degree  $\Delta \geq 2$ . Chang and Narayanan [4] proved that  $\chi'_s(G) \leq 10\Delta - 10$ . Luo and Yu [13] improved it to  $\chi'_s(G) \leq 8\Delta - 4$ . For arbitrary values of  $k$ , Wang [15] improved the result of Yu [16] and showed the following result.

**Theorem 2.** *If  $G$  is a  $k$ -degenerate graph with maximum degree  $\Delta \geq k$ , then  $\chi'_s(G) \leq (4k - 2)\Delta - 2k^2 + 1$ .*

This implies that  $\chi'_s(G) \leq 6\Delta - 7$  for 2-degenerate graph  $G$ . Choi, Kim, Kostochka, and Raspaud [5] further improved it to  $\chi'_s(G) \leq 5\Delta + 1$  in 2016. Many believe that the optimal bound should be  $4\Delta + C$  for some constant  $C$ , but no progress has yet been made.

---

*E-mail address:* gyu@wm.edu.

In this article, we show that for any  $0 < \epsilon \leq 1/2$ , and  $\Delta \geq 4^{1/(2\epsilon)}$ ,  $\chi'_s(G) \leq 5\Delta - \Delta^{1/2-\epsilon} + 2$  for 2-degenerate graph  $G$  with maximum degree  $\Delta$ .

## 2. MAIN RESULT AND ITS PROOF

A *special vertex* is a vertex with at most two neighbors of degree more than two. Every 2-degenerate graph contains special vertices, which are the new  $2^-$ -vertices after we remove all vertices of degree at most two. Let  $G$  be a 2-degenerate graph and  $S$  be the set of special vertices of  $G$ . For each  $u \in S$ , there exists a set  $W_u$  of vertices such that each  $w \in W_u$  shares some 2-neighbors with  $u$ . The *capacity of special vertices* of  $G$  is the maximum number of common 2-neighbors that are shared by a vertex in  $S$  and other vertices in  $G$ . A *pendant edge* is an edge incident with a leaf (a vertex of degree one). Below is the main result of this article.

**Theorem 3.** *For any  $0 < \epsilon \leq 1/2$ , let  $D$  be a positive integer when  $D \geq 4^{1/(2\epsilon)}$ , for any 2-degenerate graph  $G$  with maximum degree  $\Delta$ , if a vertex  $u$  is adjacent to at least  $d(u) - D$  leaves when it has  $d(u) > D$ , and*

- $\Delta \leq D + 2$  when the capacity of special vertices is at least  $D^{1/2-\epsilon}$ , and
- $\Delta \leq D + D^{1/2-\epsilon}$  when the capacity of special vertices is less than  $D^{1/2-\epsilon}$ ,

then  $\chi'_s(G) \leq 5D - D^{1/2-\epsilon} + 2$ .

*Proof.* Let  $G$  be a counterexample with the fewest number of vertices of degree at least two.

Since  $G$  is 2-degenerate,  $G$  must have a set  $S$  of special vertices. Let  $u \in S$  and  $W_u$  be the set whose vertices share 2-neighbors with  $u$  such that the maximum number of common 2-neighbors of  $u$  and vertices of  $W_u$  is the capacity of special vertices of  $G$ . Let  $W_u = \{w_1, \dots, w_s\}$  and  $u_1, u_2$  be the two neighbors of  $u$  with degree more than 2. For each  $w_i \in W_u$ , let  $W_i = \{v_{i,1}, \dots, v_{i,t_i}\}$  be the common 2-neighbors of  $w_i$  and  $u$ . Then  $N(u) = \{u_1, u_2\} \cup \bigcup_{i=1}^s \{v_{i,1}, \dots, v_{i,t_i}\}$ . We assume that  $t_1 \geq t_2 \geq \dots \geq t_s \geq 1$ . Then  $t_1$  is the capacity of special vertices of  $G$ . It is not hard to see that  $u$  has no neighbors of degree one and  $w_i$  has degree at least two for each  $i$ .

For edge  $uv$ , let  $N_2(uv)$  be the set of edges  $xy$  such that  $x$  or  $y$  is adjacent to  $u$  or  $v$ . By definition,  $uv$  should have a color different from the colors on edges in  $N_2(uv)$  in a valid strong edge-coloring.

**Case 1.**  $t_1 < D^{1/2-\epsilon}$ . In this case, we have  $\Delta \leq D + D^{1/2-\epsilon}$ .

Let  $G'$  be the graph obtained from  $G - \{uv_{1,1}, \dots, uv_{s,t_s}\}$  by adding up to  $D^{1/2-\epsilon}$  pendant neighbors to each of  $\{w_1, \dots, w_s\}$  so that  $w_i$  has degree at most  $D + D^{1/2-\epsilon}$  and  $w_i$  has at least  $D^{1/2-\epsilon}$  pendant neighbors. Then the graph  $G'$  has fewer vertices of degree at least 2 and can be colored with  $5D - D^{1/2-\epsilon} + 2$  colors. We modify the coloring of  $G'$  to obtain a coloring of  $G$  according to the following algorithm.

- (1) Keep the colors of edges that appear in both  $G$  and  $G'$ , but if  $w_i v_{i,j}$  for some  $i, j$  in  $G$  has the same color as  $uu_1$  or  $uu_2$ , then we switch color of  $w_i v_{i,j}$  with a color on other pendant edges incident to  $w_i$  in  $G'$ .
- (2) For each  $i$ , if a color  $c$  appears on both a pendant edge incident to  $w_i$  in  $G'$  and an edge incident to  $u_1$  or  $u_2$  (not including  $uu_1$  and  $uu_2$ ), then we switch the color of  $w_i v_{i,j}$  for some  $j$  with the color  $c$ .
- (3) After (2), if a color  $c$  appears on pendant edges of two or more vertices in  $W_u$  in  $G'$ , then we switch the color of  $w_i v_{i,j}$  for some  $j$  with  $c$  for each such vertex  $w_i \in W_u$ .
- (4) After (2) and (3), we color the edges  $uv_{1,1}, \dots, uv_{s,t_s}$  in reverse order with colors available to them.

Now we show that the above algorithm gives a valid strong edge-coloring of  $G$ . To do that, we only need to show that each of the edges in  $\{uv_{1,1}, \dots, uv_{s,t_s}\}$  can be colored. Consider  $uv_{i,j}$  for  $1 \leq i \leq s$  and  $1 \leq j \leq t_i$ . It needs to get a color not on edges in  $N_2(uv_{i,j})$ . Note that  $N_2(uv_{i,j})$  contains the edges incident to  $u_1, u_2, w_i$  and the edges incident to the 2-neighbors of  $u$ ; So the number of colored edges in  $N_2(uv_{i,j})$ , denoted as  $n_2(uv_{i,j})$ , is at most

$$d(u_1) + d(u_2) + d(w_i) + d(u) - 3 + d(u) - 2 - t_i - \sum_{p=1}^{i-1} t_p - (j-1).$$

We assume that  $n_2(uv_{i,j}) \geq 5D - D^{1/2-\epsilon} + 2$ , for otherwise,  $uv_{i,j}$  can be colored. Because of the way the edges being colored, we have some repeated colors on edges incident to the 2-neighbors of  $u$ , namely  $v_{1,1}w_1, v_{1,2}w_1, \dots, v_{s,t_s}w_s$ . The number of colors on  $N_2(uv_{i,j})$  and edges incident to  $w_i$  for  $i \in [s]$  (with repetition) is  $n_2(uv_{i,j}) + D^{1/2-\epsilon} \cdot (s-1)$ . Thus  $n_2(uv_{i,j}) + D^{1/2-\epsilon}(s-1) - (5D - D^{1/2-\epsilon} + 2)$  colors are repeated. As each  $w_i$  may allow only one edge (for example,  $t_i = 1$ ) whose color is the same as other edges in  $N_2(uv_{i,j})$ , at least  $\frac{n_2(uv_{i,j}) + D^{1/2-\epsilon}(s-1) - (5D - D^{1/2-\epsilon} + 2)}{D^{1/2-\epsilon}}$  edges have the same colors as others. Since  $t_1 < D^{1/2-\epsilon}$  and  $s \geq \frac{d(u)-2}{t_1} \geq \frac{d(u)-2}{D^{1/2-\epsilon}}$ , the number of different colors in  $N_2(uv_{i,j})$  is at most

$$\begin{aligned} & n_2(uv_{i,j}) - \frac{n_2(uv_{i,j}) + D^{1/2-\epsilon}(s-1) - (5D - D^{1/2-\epsilon} + 2)}{D^{1/2-\epsilon}} \\ & \leq n_2(uv_{i,j}) \left(1 - \frac{1}{D^{1/2-\epsilon}}\right) - (s-1) + 5D^{1/2+\epsilon} - 1 + \frac{2}{D^{1/2-\epsilon}} \\ & \leq (d(u_1) + d(u_2) + 2d(u) - 5 + d(w_i) - t_1) \left(1 - \frac{1}{D^{1/2-\epsilon}}\right) - \frac{d(u)-2}{D^{1/2-\epsilon}} + 5D^{1/2+\epsilon} + \frac{2}{D^{1/2-\epsilon}} \\ & \leq 3\Delta \left(1 - \frac{1}{D^{1/2-\epsilon}}\right) + d(u) \left(2 - \frac{3}{D^{1/2-\epsilon}}\right) + 5D^{1/2+\epsilon} - 5 + \frac{9}{D^{1/2-\epsilon}} \\ & \leq 3(D + D^{1/2-\epsilon}) \left(1 - \frac{1}{D^{1/2-\epsilon}}\right) + D \left(2 - \frac{3}{D^{1/2-\epsilon}}\right) + 5D^{1/2+\epsilon} - 5 + \frac{9}{D^{1/2-\epsilon}} \\ & \leq 5D + 3D^{1/2-\epsilon} - D^{1/2+\epsilon} - 8 + \frac{9}{D^{1/2-\epsilon}} \leq 5D - D^{1/2-\epsilon} + 1. \end{aligned}$$

The last inequality holds because  $D^{1/2+\epsilon} \geq 4D^{1/2-\epsilon}$  and  $\frac{9}{D^{1/2-\epsilon}} \leq 9$  when  $D \geq 4^{\frac{1}{2\epsilon}}$ . Therefore, the edge  $uv_{i,j}$  can be colored, which implies that we can color all the uncolored edges.

**Case 2.**  $t_1 \geq D^{1/2-\epsilon}$ . In this case,  $\Delta \leq D + 2$ .

Let  $G'$  be the graph obtained from  $G$  by deleting the edge  $uv_{1,1}$  and adding up to two pendant edges notated  $\{w_1v', w_1v''\}$  so that  $w_1$  has at least three pendant edges and continues to have maximum degree at most  $D + 2$ .

Note that for  $1 \leq i \leq s$  and  $1 \leq j \leq t_s$ ,

$$(1) |N_2(uv_{i,j})| \leq d(u_1) + d(u_2) + d(w_i) + d(u) - 3 + d(u) - 2 - t_i \leq 3\Delta + 2D - 5 - t_i \leq 5D + 1 - t_i.$$

We observe that  $G'$  is 2-degenerate, has fewer vertices of degree at least 2, and contains at least  $d(v) - D$  vertices of degree one if  $d(v) > D$ . So,  $G'$  can be colored with  $5D - D^{1/2-\epsilon} + 2$  colors. Keep the colors of the edges in both  $G$  and  $G'$ , we try to get a valid coloring of  $G$ .

**Case 2.1.** the color of edge  $v_{1,1}w_1$  is different from ones on  $\{uu_1, uu_2, uv_{1,2}, \dots, uv_{s,t_s}\}$ . By (1), we have at most  $5D + 1 - D^{1/2-\epsilon}$  edges in  $N_2(uv_{1,1})$ . Since there are  $5D - D^{1/2-\epsilon} + 2$  different colors available, we can color  $uv_{1,1}$  with a color not on the edges in  $N_2(uv_{1,1})$ .

**Case 2.2.** the color of edge  $v_{1,1}w_1$  is the same as the color of edge  $uu_1$  (or edge  $uu_2$ ). In this case, we swap the color of  $v_{1,1}w_1$  with the color of  $w_1v'$  or  $w_1v''$ . By (1) again,  $N_2(uv_{1,1})$  has at most  $5D + 1 - D^{1/2-\epsilon}$  edges in  $N_2(uv_{1,1})$ . Since there are  $5D - D^{1/2-\epsilon} + 2$  different colors available, we can color  $uv_{1,1}$  with a color not on the edges in  $N_2(uv_{1,1})$ .

**Case 2.3.** the color of edge  $v_{1,1}w_1$  is the same as the color of edge  $uv_{i,j}$  for some  $2 \leq i \leq s$ . In this case, we uncolor the edges  $uv_{i,j}, uv_{1,1}, \dots, uv_{1,t_1}$ , and recolor them in the order. By (1), there are at most  $5D + 1 - t_i - (t_1 - 1) \leq 5D - D^{1/2-\epsilon} + 1$  colors on edges in  $N_2(uv_{i,j})$ , so  $uv_{i,j}$  can be colored. Similarly, for each  $j$ , there are at most  $5D - D^{1/2-\epsilon} + 1$  colors on edges in  $N_2(uv_{1,j})$ , so  $uv_{1,j}$  can be colored as well.

In any case,  $G$  can be colored with  $5D - D^{1/2-\epsilon} + 2$  colors. So  $\chi'_s(G) \leq 5D - D^{1/2-\epsilon} + 2$ .  $\square$

Observe that our main result follows from Theorem 3 with  $\Delta = D$ . Our proof does not work for corresponding result of list version.

#### REFERENCES

- [1] L. D. Andersen, The strong chromatic index of a cubic graph is at most 10, *Discrete Math.* 108 (1-3) (1992) 231–252.
- [2] M. Bonamy, T. Perrett and L. Postle, Colouring graphs with sparse neighbourhoods: Bounds and applications, *J. Combin. Theory Ser. B* 155 (2022) 278–317.
- [3] H. Bruhn and F. Joos, A stronger bound for the strong chromatic index, *Combin. Probab. Comput.* 27 (1) (2018) 21–43.
- [4] G. J. Chang and N. Narayanan, Strong chromatic index of 2-degenerate graphs, *J. Graph Theory* 73 (2) (2013) 119–126.
- [5] I. Choi, J. Kim, A. V. Kostochka and A. Raspaud, Strong edge-colorings of sparse graphs with large maximum degree, *European J. Combin.* 67 (2018) 21–39.
- [6] F. R. K. Chung, A. Gyarfas, Z. Tuza and W. T. Trotter, The maximum number of edges in  $2K_2$ -free graphs of bounded degree, *Discrete Math.* 81 (2) (1990) 129–135.
- [7] K. Deng, G. Yu and X. Zhou, Recent progress on strong edge-coloring of graphs, *Discrete Math. Algorithms Appl.* 11 (5) (2019) 1950062.
- [8] P. Erdős and J. Nešetřil, Irregularities of partitions, (G. Halasz, V.T. Sós (Eds.)), [Problem] (1989) 162–163.
- [9] J.-L. Fouquet and J.-L. Jolivet, Strong edge-colorings of graphs and applications to multi- $k$ -gons, *Ars Combin.* 16 (A) (1983) 141–150.
- [10] P. Horak, Q. He and W. T. Trotter, Induced matchings in cubic graphs, *J. Graph Theory* 17 (2) (1993) 151–160.
- [11] M. Huang, M. Santana and G. Yu, Strong chromatic index of graphs with maximum degree four, *Electron. J. Combin.* 25 (3) (2018) Paper #3.31.
- [12] E. Hurley, R. de Joannis de Verclos and R. J. Kang, An improved procedure for colouring graphs of bounded local density, *Adv. in Combinatorics*, (2022), 7, 33pp.
- [13] R. Luo and G. Yu, A note on strong edge-colorings of 2-degenerate graphs, arXiv:1212.6092, December 25, 2012, <https://arxiv.org/abs/1212.6092>.
- [14] M. Molloy and B. Reed, A bound on the strong chromatic index of a graph, *J. Combin. Theory Ser. B* 69 (2) (1997) 103–109.
- [15] T. Wang, Strong chromatic index of  $k$ -degenerate graphs, *Discrete Math.* 330 (2014) 17–19.
- [16] G. Yu, Strong edge-colorings for  $k$ -degenerate graphs, *Graphs Combin.* 31 (5) (2015) 1815–1818.