

# SOME DOUBLE SERIES FOR $\pi$ AND THEIR $q$ -ANALOGUES

GUOPING GU AND XIAOXIA WANG\*

ABSTRACT. In this paper, by applying the partial derivative operator on summation formulas of hypergeometric series and basic hypergeometric series, we establish several double series for  $\pi$  and their  $q$ -analogues.

## 1. INTRODUCTION

In 1914, Ramanujan [11] laid out 17 series for  $1/\pi$  without proof, and then all of them were proved by Borweins [2]. One of Ramanujan's formulas is stated as

$$\sum_{k=0}^{\infty} (6k+1) \frac{\left(\frac{1}{2}\right)_k^3}{k!^3 4^k} = \frac{4}{\pi},$$

where the shifted factorial  $(x)_n$  is defined as follows: for a complex number  $x$ ,

$$(x)_n = \begin{cases} x(x+1)(x+2)\cdots(x+n-1) & \text{if } n \in \mathbb{Z}^+; \\ 1 & \text{if } n = 0. \end{cases}$$

In addition,  $(x)_n$  can also be expressed by

$$(x)_n = \frac{\Gamma(x+n)}{\Gamma(x)}.$$

Here  $\Gamma(x)$  stands for the well-known Gamma function which is defined as

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt \quad \text{with } \operatorname{Re}(x) > 0.$$

The Gamma function has many important properties, for example:

$$\Gamma(x+1) = x\Gamma(x), \quad \Gamma(x)\Gamma(1-x) = \frac{\pi}{\sin(\pi x)}, \quad \lim_{n \rightarrow \infty} \frac{\Gamma(x+n)}{\Gamma(y+n)} n^{y-x} = 1,$$

which will be used without explanation in the following of this paper.

---

2010 *Mathematics Subject Classification.* Primary 33D15; Secondary 11A07, 11B65.

*Key words and phrases.* basic hypergeometric series;  $q$ -analogues; double series for  $\pi$ ; partial derivative operator.

Recently, Wei [14] certified the following two double series for  $\pi$  which were conjectured by Guo and Lian [7]:

$$\begin{aligned} \sum_{k=1}^{\infty} (6k+1) \frac{(\frac{1}{2})_k^3}{k!^3 4^k} \sum_{j=1}^k \left( \frac{1}{(2j-1)^2} - \frac{1}{16j^2} \right) &= \frac{\pi}{12}, \\ \sum_{k=1}^{\infty} (-1)^k (6k+1) \frac{(\frac{1}{2})_k^3}{k!^3 8^k} \sum_{j=1}^k \left( \frac{1}{(2j-1)^2} - \frac{1}{16j^2} \right) &= -\frac{\sqrt{2}\pi}{48}. \end{aligned} \quad (1.1)$$

Besides, Swisher [13] gave a congruence on truncated form of (1.1), which was conjectured by Long [10] in 2011. Furthermore, Wei also similarly gives another three double series for  $\pi$  in [15] and one of them is

$$\sum_{k=1}^{\infty} (-1)^k (4k+1) \frac{(\frac{1}{2})_k^3}{k!^3} \sum_{i=1}^{2k} \frac{(-1)^i}{i^2} = \frac{\pi}{12}, \quad (1.2)$$

with the  $q$ -analogue as

$$\sum_{k=1}^{\infty} (-1)^k q^{k^2} [4k+1] \frac{(q; q^2)_k^3}{(q^2; q^2)_k^3} \sum_{i=1}^{2k} (-1)^i \frac{q^i}{[i]^2} = \frac{(q, q^3; q^2)_{\infty}}{(q^2; q^2)_{\infty}^2} \sum_{j=1}^{\infty} \frac{q^{2j}}{[2j]^2}. \quad (1.3)$$

Here the  $q$ -integer  $[n]$  is defined as  $[n] = (1 - q^n)/(1 - q) = 1 + q + \cdots + q^{n-1}$  and the  $q$ -shifted factorial is stated as

$$(x; q)_{\infty} = \prod_{i=0}^{\infty} (1 - xq^i), \quad (x; q)_n = \frac{(x; q)_{\infty}}{(xq^n; q)_{\infty}}.$$

For convenience, we shall also adopt the following notation

$$(x_1, x_2, \dots, x_r; q)_m = (x_1; q)_m (x_2; q)_m \cdots (x_r; q)_m, \quad m \in \mathbb{Z}^+ \cup \{0, \infty\}.$$

Inspired by the work just mentioned, we deduce some double series for  $\pi$  and their  $q$ -analogues by applying the partial derivative operator. For more known series on  $\pi$ , we refer the reader to the papers [1, 3, 9, 12, 16].

For the completeness of this paper, we will introduce the following concept. For a multivariable function  $f(x_1, x_2, \dots, x_m)$ , define the partial derivative operator  $\mathcal{D}_{x_i}$  by

$$\mathcal{D}_{x_i} f(x_1, x_2, \dots, x_m) = \frac{d}{dx_i} f(x_1, x_2, \dots, x_m) \quad \text{with } 1 \leq i \leq m.$$

## 2. MAIN RESULTS

Applying the partial derivative operator on the summation formula of hypergeometric series, we get the following double series for  $\pi$ .

**Theorem 1.** *The following result is true.*

$$\sum_{k=1}^{\infty} (6k-1) \frac{\left(-\frac{1}{2}\right)_k^2}{4^k k! \left(\frac{3}{2}\right)_k} \sum_{i=1}^k \frac{1}{(2i-1)^2} = \frac{\pi^3}{144}. \quad (2.1)$$

With the help of Gasper and Rahman's quadratic summation formula [5, Equation (3.8.12)], we get the  $q$ -analogue of (2.1).

**Theorem 2.** *Let the complex number  $q$  with  $0 < |q| < 1$ . Then, we have*

$$\begin{aligned} & \sum_{k=1}^{\infty} [6k-1] \frac{(q^{-1}, q, q; q^2)_k (q^{-2}; q^4)_k}{(q^4, q^2, q^2; q^4)_k (q^3; q^2)_k} q^{(k+1)^2} \sum_{i=1}^k \left( \frac{q^{2i-1}}{[2i-1]^2} - \frac{q^{4i-2}}{[4i-2]^2} \right) \\ &= \frac{(q, q^4, q^4; q^4)_{\infty}}{(q^5, q^2, q^2; q^4)_{\infty}} \sum_{i=1}^{\infty} \left( \frac{q^{4i-2}}{[4i-2]^2} - \frac{q^{4i}}{[4i]^2} \right). \end{aligned} \quad (2.2)$$

In addition, we establish many other double series for  $\pi$  also from the summation formula of hypergeometric series.

**Theorem 3.** *The following results are true.*

$$\sum_{k=1}^{\infty} (4k+1) \frac{\left(-\frac{1}{2}\right)_k \left(\frac{1}{2}\right)_k^3}{(k+1)! k!^3} \sum_{i=1}^{2k} (-1)^{i-1} \frac{1}{i^2} = \frac{2}{3} - \frac{8}{\pi^2}, \quad (2.3)$$

$$\sum_{k=1}^{\infty} (4k+3) \frac{\left(-\frac{1}{2}\right)_k \left(\frac{1}{2}\right)_k^2 \left(\frac{3}{2}\right)_k}{k! (k+1)!^2 (k+2)!} \sum_{i=1}^k \left( \frac{1}{(2i-1)^2} - \frac{1}{4(i+1)^2} \right) = \frac{32}{27} - \frac{992}{81\pi^2}, \quad (2.4)$$

$$\sum_{k=1}^{\infty} (4k+3) \frac{\left(\frac{3}{2}\right)_k \left(\frac{1}{2}\right)_k^3}{k! (k+1)!^3} \sum_{i=1}^k \left( \frac{1}{(2i-1)^2} - \frac{1}{4(i+1)^2} \right) = \frac{8}{3} - \frac{24}{\pi^2}, \quad (2.5)$$

$$\sum_{k=1}^{\infty} (-1)^k (4k+3) \frac{\left(\frac{3}{2}\right)_k \left(\frac{1}{2}\right)_k^2}{k! (k+1)!^2} \sum_{i=1}^k \left( \frac{1}{(1+i)^2} - \frac{4}{(2i-1)^2} \right) = \frac{4\pi}{3} - \frac{8}{\pi}. \quad (2.6)$$

The  $q$ -analogues of the double series for  $\pi$  in Theorem 3 are expressed as follows.

**Theorem 4.** *Let the complex number  $q$  with  $0 < |q| < 1$ . Then, we have*

$$\begin{aligned} & \sum_{k=1}^{\infty} [4k+1] \frac{(q; q^2)_k^3 (q^{-1}; q^2)_k}{(q^2; q^2)_k^3 (q^4; q^2)_k} q^{2k} \sum_{i=1}^k \left( \frac{q^{2i}}{[2i]^2} - \frac{q^{2i-1}}{[2i-1]^2} \right) \\ &= \frac{(q^3; q^2)_{\infty}^3 (q; q^2)_{\infty}}{(q^2; q^2)_{\infty}^3 (q^4; q^2)_{\infty}} \sum_{i=1}^{\infty} \left( \frac{(-1)^{i+1}}{[i+1]^2} - \frac{q^{3i}(1-q)}{2[2i+1]^2} \right) q^{i+1}, \end{aligned} \quad (2.7)$$

$$\sum_{k=1}^{\infty} [4k+3] \frac{(q; q^2)_k^2 (q^3, q^{-1}; q^2)_k}{(q^4; q^2)_k^2 (q^2, q^6; q^2)_k} q^{4k} \sum_{i=1}^k \left( \frac{q^{2i+2}}{[2i+2]^2} - \frac{q^{2i-1}}{[2i-1]^2} \right)$$

$$= \frac{(q^3; q^2)_\infty (q^5; q^2)_\infty^2 (q^3; q^2)_\infty}{(1-q)(q^4; q^2)_\infty^3 (q^6; q^2)_\infty} \sum_{i=1}^{\infty} \left( \frac{(-1)^{i+1}}{[i+3]^2} - \frac{q^{3i+2}(1-q)}{2[2i+3]^2} \right) q^{i+3}, \quad (2.8)$$

$$\begin{aligned} & \sum_{k=1}^{\infty} [4k+3] \frac{(q; q^2)_k^3 (q^3; q^2)_k}{(q^4; q^2)_k^3 (q^2; q^2)_k} q^{2k} \sum_{i=1}^k \left( \frac{q^{2i+2}}{[2i+2]^2} - \frac{q^{2i-1}}{[2i-1]^2} \right) \\ &= \frac{(q^3; q^2)_\infty^3 (q^3; q^2)_\infty}{(1-q)(q^4; q^2)_\infty^3 (q^2; q^2)_\infty} \sum_{i=1}^{\infty} \left( \frac{(-1)^i}{[i+2]^2} + \frac{q^{3i}(1-q)}{2[2i+1]^2} \right) q^{i+2}, \end{aligned} \quad (2.9)$$

$$\begin{aligned} & \sum_{k=1}^{\infty} (-1)^k [4k+3] \frac{(q; q^2)_{k+1} (q; q^2)_k^2}{(q^2, q^4, q^4; q^2)_k} q^{-k(k+4)} \sum_{i=1}^k \left( \frac{q^{2i+2}}{[2i+2]^2} - \frac{q^{2i-1}}{[2i-1]^2} \right) \\ &= \frac{(q^3, q^3; q^2)_\infty}{(q^4, q^4; q^2)_\infty} \sum_{i=1}^{\infty} \frac{q^{2i+2}}{[2i+2]^2}. \end{aligned} \quad (2.10)$$

The structure of this paper is arranged as follows. We shall prove Theorem 1 in Section 3 and Theorem 2 in Section 4 by the partial derivative operator respectively. Finally, we will certify Theorems 3 and 4 in Section 5.

### 3. PROOF OF THEOREM 1

Following Gasper and Rahman [5], the hypergeometric series is defined as

$${}_{r+1}F_r \left[ \begin{matrix} a_1, & a_2, & \dots, & a_{r+1} \\ b_1, & b_2, & \dots, & b_r \end{matrix} ; z \right] = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k \dots (a_{r+1})_k}{(b_1)_k (b_2)_k \dots (b_r)_k} \frac{z^k}{k!}.$$

In order to prove Theorem 1, we recall the following  ${}_7F_6$ -summation formula [6, Equation (1.7)]:

$$\begin{aligned} & {}_7F_6 \left[ \begin{matrix} a, & 1 + \frac{a}{3}, & b, & 1 - b, & c, & \frac{1}{2} + a - c + n, & -n \\ & \frac{a}{3}, & \frac{2+a-b}{2}, & \frac{1+a+b}{2}, & 1 + a + 2n, & 1 + a - 2c, & 2c - a - 2n \end{matrix} \right] \\ &= \left[ \begin{matrix} \frac{1+a}{2}, & 1 + \frac{a}{2}, & \frac{1+a+b}{2} - c, & 1 + \frac{a-b}{2} - c \\ \frac{1+a+b}{2}, & 1 + \frac{a-b}{2}, & \frac{1+a}{2} - c, & 1 + \frac{a}{2} - c \end{matrix} \right]_n. \end{aligned} \quad (3.1)$$

*Proof of Theorem 1.* Applying the partial derivative  $\mathcal{D}_b$  on both sides of the  ${}_7F_6$ -summation formula (3.1), we subsequently obtain the following result

$$\begin{aligned} & \sum_{k=1}^n \frac{(a)_k (1 + \frac{a}{3})_k (b)_k (1 - b)_k (c)_k (\frac{1}{2} + a - c + n)_k (-n)_k}{(1)_k (\frac{a}{3})_k (\frac{2+a-b}{2})_k (\frac{1+a+b}{2})_k (1 + a + 2n)_k (1 + a - 2c)_k (2c - a - 2n)_k} A_k(a, b) \\ &= \frac{(\frac{1+a}{2})_n (1 + \frac{a}{2})_n (\frac{1+a+b}{2} - c)_n (1 + \frac{a-b}{2} - c)_n}{(\frac{1+a+b}{2})_n (1 + \frac{a-b}{2})_n (\frac{1+a}{2} - c)_n (1 + \frac{a}{2} - c)_n} B_n(a, b, c). \end{aligned} \quad (3.2)$$

where

$$A_k(a, b) = H_k(b-1) - H_k(-b) + \frac{1}{2}H_k\left(\frac{a-b}{2}\right) - \frac{1}{2}H_k\left(\frac{a+b-1}{2}\right),$$

$$B_n(a, b, c) = \frac{1}{2}H_n\left(\frac{a+b-1}{2} - c\right) - \frac{1}{2}H_n\left(\frac{a-b}{2} - c\right) - \frac{1}{2}H_n\left(\frac{a+b-1}{2}\right) + \frac{1}{2}H_n\left(\frac{a-b}{2}\right).$$

Here  $H_k(x)$  represents the generalized harmonic number, which is defined by

$$H_k(x) = \sum_{i=1}^k \frac{1}{x+i}.$$

More generally,  $H_k^{(m)}(x)$  represents the generalized harmonic number of order  $m$  as

$$H_k^{(m)}(x) = \sum_{i=1}^k \frac{1}{(x+i)^m}.$$

Taking  $m = 1$  and  $x = 0$  in  $H_k^{(m)}(x)$ , we get the famous classical harmonic number.

Now employing the partial derivative operator  $\mathcal{D}_b$  again on both sides of (3.2), we get

$$\begin{aligned} & \sum_{k=1}^n \frac{(a)_k \left(1 + \frac{a}{3}\right)_k (b)_k (1-b)_k (c)_k \left(\frac{1}{2} + a - c + n\right)_k (-n)_k}{(1)_k \left(\frac{a}{3}\right)_k \left(\frac{2+a-b}{2}\right)_k \left(\frac{1+a+b}{2}\right)_k (1+a+2n)_k (1+a-2c)_k (2c-a-2n)_k} \\ & \times \{A_k(a, b)^2 + C_k(a, b)\} \\ & = \frac{\left(\frac{1+a}{2}\right)_n \left(1 + \frac{a}{2}\right)_n \left(\frac{1+a+b}{2} - c\right)_n \left(1 + \frac{a-b}{2} - c\right)_n}{\left(\frac{1+a+b}{2}\right)_n \left(1 + \frac{a-b}{2}\right)_n \left(\frac{1+a}{2} - c\right)_n \left(1 + \frac{a}{2} - c\right)_n} \{B_n(a, b, c)^2 + D_n(a, b, c)\}. \end{aligned} \quad (3.3)$$

where

$$C_k(a, b) = -H_k^{(2)}(b-1) - H_k^{(2)}(-b) + \frac{1}{4}H_k^{(2)}\left(\frac{a-b}{2}\right) + \frac{1}{4}H_k^{(2)}\left(\frac{a+b-1}{2}\right),$$

$$D_n(a, b, c) = -\frac{1}{4}H_n^{(2)}\left(\frac{a+b-1}{2} - c\right) - \frac{1}{4}H_n^{(2)}\left(\frac{a-b}{2} - c\right) + \frac{1}{4}H_n^{(2)}\left(\frac{a+b-1}{2}\right) + \frac{1}{4}H_n^{(2)}\left(\frac{a-b}{2}\right).$$

Note that  $A_k(a, b) = B_n(a, b, c) = 0$  when  $a = -\frac{1}{2}, b = \frac{1}{2}, c \rightarrow -\frac{1}{2}$  in (3.3). Then, we subsequently arrive at Theorem 1 by performing the replacements  $a = -\frac{1}{2}, b = \frac{1}{2}, c \rightarrow -\frac{1}{2}$  and  $n \rightarrow \infty$  in (3.3). Here we have used the following property of Gamma function

$$\lim_{n \rightarrow \infty} \frac{\Gamma(x+n)}{\Gamma(y+n)} n^{y-x} = 1,$$

and Euler's formula

$$\sum_{i=1}^{\infty} \frac{1}{i^2} = \frac{\pi^2}{6}.$$

□

## 4. PROOF OF THEOREM 2

Following Gasper and Rahman [5], the basic hypergeometric series  ${}_{r+1}\phi_r$  is defined as

$${}_{r+1}\phi_r \left[ \begin{matrix} a_1, & a_2, & \dots, & a_{r+1} \\ b_1, & b_2, & \dots, & b_r \end{matrix} ; q, z \right] = \sum_{k=0}^{\infty} \frac{(a_1, a_2, \dots, a_{r+1}; q)_k}{(q, b_1, b_2, \dots, b_r; q)_k} z^k.$$

For the purpose of proving Theorem 2, we first present Gasper and Rahman's quadratic summation formula [5, Equation (3.8.12)] as follows:

$$\begin{aligned} & \sum_{k=0}^{\infty} \frac{1 - aq^{3k}}{1 - a} \frac{(a, b, q/b; q)_k (d, f, a^2q/df; q^2)_k}{(q^2, aq^2/b, abq; q^2)_k (aq/d, aq/f, df/a; q)_k} q^k \\ & \quad + \frac{(aq, f/a, b, q/b; q)_{\infty} (d, a^2q/df, fq^2/d, df^2q/a^2; q^2)_{\infty}}{(a/f, fq/a, aq/d, df/a; q)_{\infty} (aq^2/b, abq, fq/ab, bf/a; q^2)_{\infty}} \\ & \quad \times {}_3\phi_2 \left[ \begin{matrix} f, bf/a, fq/ab \\ fq^2/d, df^2q/a^2 \end{matrix} ; q^2, q^2 \right] \\ & = \frac{(aq, f/a; q)_{\infty} (aq^2/bd, abq/d, bdf/a, dfq/ab; q^2)_{\infty}}{(aq/d, df/a; q)_{\infty} (aq^2/b, abq, bf/a, fq/ab; q^2)_{\infty}}. \end{aligned} \quad (4.1)$$

*Proof of Theorem 2.* Performing the replacements  $d \rightarrow q^{-2n}$  and  $f \rightarrow c^2$  in (4.1), we get its truncated form as

$$\begin{aligned} & \sum_{k=0}^n \frac{1 - aq^{3k}}{1 - a} \frac{(a, b, q/b; q)_k (q^{-2n}, c^2, a^2q^{2n+1}/c^2; q^2)_k}{(q^2, aq^2/b, abq; q^2)_k (aq^{2n+1}, aq/c^2, c^2q^{-2n}/a; q)_k} q^k \\ & = \frac{(aq, aq^2, aq^2/bc^2, abq/c^2; q^2)_n}{(aq/c^2, aq^2/c^2, aq^2/b, abq; q^2)_n}. \end{aligned}$$

Applying the partial derivative  $\mathcal{D}_b$  on both sides of the above identity twice, then we subsequently obtain the following summation

$$\begin{aligned} & \sum_{k=1}^n \frac{1 - aq^{3k}}{1 - a} \frac{(a, b, q/b; q)_k (q^{-2n}, c^2, a^2q^{2n+1}/c^2; q^2)_k}{(q^2, aq^2/b, abq; q^2)_k (aq^{2n+1}, aq/c^2, c^2q^{-2n}/a; q)_k} q^k \{A_k^2(a, b) + C_k(a, b)\} \\ & = \frac{(aq, aq^2, aq^2/bc^2, abq/c^2; q^2)_n}{(aq/c^2, aq^2/c^2, aq^2/b, abq; q^2)_n} \{B_n^2(a, b, c) + D_n(a, b, c)\}. \end{aligned} \quad (4.2)$$

where

$$\begin{aligned} A_k(a, b) &= \sum_{i=1}^k \frac{-q^{i-1}}{1 - bq^{i-1}} + \sum_{i=1}^k \frac{q^i/b^2}{1 - q^i/b} - \sum_{i=1}^k \frac{aq^{2i}/b^2}{1 - aq^{2i}/b} - \sum_{i=1}^k \frac{-aq^{2i-1}}{1 - abq^{2i-1}}, \\ B_n(a, b, c) &= \sum_{i=1}^n \frac{aq^{2i}/b^2c^2}{1 - aq^{2i}/bc^2} + \sum_{i=1}^n \frac{-aq^{2i-1}/c^2}{1 - abq^{2i-1}/c^2} - \sum_{i=1}^n \frac{aq^{2i}/b^2}{1 - aq^{2i}/b} - \sum_{i=1}^n \frac{-aq^{2i-1}}{1 - abq^{2i-1}}, \end{aligned}$$

$$\begin{aligned}
C_k(a, b) &= \sum_{i=1}^k \frac{-q^{2i-2}}{(1-bq^{i-1})^2} + \sum_{i=1}^k \frac{q^i/b^3(q^i/b-2)}{(1-q^i/b)^2} - \sum_{i=1}^k \frac{aq^{2i}/b^3(aq^{2i}/b-2)}{(1-aq^{2i}/b)^2} \\
&\quad + \sum_{i=1}^k \frac{a^2q^{4i-2}}{(1-abq^{2i-1})^2}, \\
D_n(a, b, c) &= \sum_{i=1}^n \frac{aq^{2i}/b^3c^2(aq^{2i}/bc^2-2)}{(1-aq^{2i}/bc^2)^2} + \sum_{i=1}^n \frac{-a^2q^{4i-2}/c^4}{(1-abq^{2i-1}/c^2)^2} - \sum_{i=1}^n \frac{aq^{2i}/b^3(aq^{2i}/b-2)}{(1-aq^{2i}/b)^2} \\
&\quad + \sum_{i=1}^n \frac{a^2q^{4i-2}}{(1-abq^{2i-1})^2}.
\end{aligned}$$

Clearly,  $A_k(a, b) = B_n(a, b, c) = 0$  when  $a = q^{-\frac{1}{2}}, b = q^{\frac{1}{2}}, c = q^{-\frac{1}{2}}$ . We have the following identity by substituting  $a = q^{-\frac{1}{2}}, b = q^{\frac{1}{2}}$  and  $c = q^{-\frac{1}{2}}$  in (4.2)

$$\begin{aligned}
&\sum_{k=1}^n \frac{1-q^{2k+\frac{3}{2}}}{1-q^{\frac{3}{2}}} \frac{(q^{\frac{3}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{n+\frac{5}{2}}, q^{-n}; q)_k}{(q, q^2, q^2, q^2, q^{-n}, q^{n+\frac{5}{2}}; q)_k} q^k \sum_{i=1}^k \left( \frac{q^i}{(1-q^{i+1})^2} - \frac{q^{i-\frac{3}{2}}}{(1-q^{i-\frac{1}{2}})^2} \right) \\
&= \frac{(q^{\frac{5}{2}}, q^{\frac{3}{2}}, q^{\frac{3}{2}}, q^{\frac{3}{2}}; q)_n}{(q^2, q^2, q^2, q; q)_n} \left( \sum_{i=1}^{2n} (-1)^i \frac{q^{\frac{i}{2}}}{(1-q^{\frac{i}{2}+1})^2} + \frac{1}{2} \sum_{i=1}^n \frac{q^{2i} - q^{2i+\frac{1}{2}}}{(1-q^{i+\frac{1}{2}})^2} \right).
\end{aligned}$$

Finally, we get Theorem 2 by replacing  $q$  by  $q^2$  and letting  $n \rightarrow \infty$  in the above identity.  $\square$

## 5. PROOFS OF THEOREMS 3 AND 4

Since the double series for  $\pi$  in Theorem 3 are just the  $q \rightarrow 1$  cases of Theorem 4, the proof of Theorem 4 is enough. For proving Theorem 4, we first present the following Lemma which plays a key role in our proof.

**Lemma 1.** *Let the complex number  $q$  with  $0 < |q| < 1$ . Then, we have*

$$\begin{aligned}
&\sum_{k=1}^n \frac{1-aq^{2k}}{1-a} \frac{(a, b, c, q/b, a^2q^n/c, q^{-n}; q)_k}{(q, aq/b, aq/c, ab, cq^{1-n}/a, aq^{n+1}; q)_k} q^k \{A_k^2(a, b) + C_k(a, b)\} \\
&= \frac{(aq, aq/bc, a, ab/c; q)_n}{(aq/b, aq/c, ab, a/c; q)_n} \{B_n^2(a, b, c) + D_n(a, b, c)\}.
\end{aligned} \tag{5.1}$$

where

$$\begin{aligned}
A_k(a, b) &= \sum_{i=1}^k \frac{-q^{i-1}}{1-bq^{i-1}} + \sum_{i=1}^k \frac{q^i/b^2}{1-q^i/b} - \sum_{i=1}^k \frac{aq^i/b^2}{1-aq^i/b} + \sum_{i=1}^k \frac{aq^{i-1}}{1-abq^{i-1}}, \\
B_n(a, b, c) &= \sum_{i=1}^n \frac{aq^i/b^2c}{1-aq^i/bc} + \sum_{i=1}^n \frac{-aq^{i-1}/c}{1-abq^{i-1}/c} - \sum_{i=1}^n \frac{aq^i/b^2}{1-aq^i/b} + \sum_{i=1}^n \frac{aq^{i-1}}{1-abq^{i-1}},
\end{aligned}$$

$$\begin{aligned}
C_k(a, b) &= - \sum_{i=1}^k \frac{q^{2i-2}}{(1-bq^{i-1})^2} + \sum_{i=1}^k \frac{(q^i/b-2)q^i/b^3}{(1-q^i/b)^2} - \sum_{i=1}^k \frac{(aq^i/b-2)aq^i/b^3}{(1-aq^i/b)^2} \\
&\quad + \sum_{i=1}^k \frac{a^2q^{2i-2}}{(1-abq^{i-1})^2}, \\
D_n(a, b, c) &= \sum_{i=1}^n \frac{(aq^i/bc-2)aq^i/b^3c}{(1-aq^i/bc)^2} - \sum_{i=1}^n \frac{a^2q^{2i-2}/c}{(1-abq^{i-1}/c)^2} - \sum_{i=1}^n \frac{(aq^i/b-2)aq^i/b^3}{(1-aq^i/b)^2} \\
&\quad + \sum_{i=1}^n \frac{a^2q^{2i-2}}{(1-abq^{i-1})^2}.
\end{aligned}$$

Obviously, we can derive Wei's result (1.3) by replacing  $q \rightarrow q^2$  and letting  $n \rightarrow \infty$  after setting  $(a, b) = (q^{\frac{1}{2}}, q^{\frac{1}{2}})$  and  $c \rightarrow \infty$  in Lemma 1.

*Proof.* Recall Jackson's  $q$ -Dougall-Dixon formula [8] which can be stated as follows

$$\begin{aligned}
& {}_8\phi_7 \left[ \begin{matrix} a, qa^{\frac{1}{2}}, -qa^{\frac{1}{2}}, b, c, d, e, q^{-n} \\ a^{\frac{1}{2}}, -a^{\frac{1}{2}}, aq/b, aq/c, aq/d, aq/e, aq^{n+1} \end{matrix}; q, q \right] \\
&= \frac{(aq, aq/bc, aq/bd, aq/cd; q)_n}{(aq/b, aq/c, aq/d, aq/bcd; q)_n}, \quad \text{where } q^{n+1}a^2 = bcde.
\end{aligned} \tag{5.2}$$

Now, putting  $d \rightarrow q/b$  and  $e \rightarrow a^2q^n/c$  in (5.2), we arrive at

$$\sum_{k=0}^n \frac{1-aq^{2k}}{1-a} \frac{(a, b, c, q/b, a^2q^n/c, q^{-n}; q)_k}{(q, aq/b, aq/c, ab, cq^{1-n}/a, aq^{n+1}; q)_k} q^k = \frac{(aq, aq/bc, a, ab/c; q)_n}{(aq/b, aq/c, ab, a/c; q)_n}.$$

Then we subsequently conclude that the identity (1) holds by applying the partial derivative  $\mathcal{D}_b$  on both sides of the above identity twice.  $\square$

*Proof of Theorem 4.* We can subsequently get (2.7) by taking  $(a, b, c) = (q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{-\frac{1}{2}})$ , and then replacing  $q \rightarrow q^2$  and letting  $n \rightarrow \infty$  in Lemma 1. Similarly, letting  $(a, b, c) = (q^{\frac{3}{2}}, q^{\frac{1}{2}}, q^{-\frac{1}{2}})$ ,  $(a, b, c) = (q^{\frac{3}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{2}})$ ,  $(a, b) = (q^{\frac{3}{2}}, q^{\frac{1}{2}})$  and  $c \rightarrow \infty$ , we are led to the double summation identities (2.8), (2.9) and (2.10) respectively.  $\square$

In fact, we can also prove the truth of Theorem 3 by the following Dougall's summation formula [4, Corollary (3)] in the same way as we have done in the proof of Theorem 1.

$$\begin{aligned}
& {}_7F_6 \left[ \begin{matrix} a, 1 + \frac{a}{2}, & b, & c, & d, & e, & -n \\ & \frac{a}{2}, & 1+a-b, & 1+a-c, & 1+a-d, & 1+a-e, & 1+a+n \end{matrix} \right] \\
&= \left[ \begin{matrix} 1+a, & 1+a-b-c, & 1+a-b-d, & 1+a-c-d \\ 1+a-b, & 1+a-c, & 1+a-d, & 1+a-b-c-d \end{matrix} \right]_n.
\end{aligned}$$

## REFERENCES

- [1] A. Berkovich, H.H. Chan, M.J. Schlosser, Wronskians of theta functions and series for  $1/\pi$ , *Adv. Math.* 338 (2018), 266-304.
- [2] J.M. Borwein, P.B. Borwein,  $\pi$  and the AGM: A Study in Analytic Number Theory and Computational Complexity, Wiley, New York, 1987.
- [3] H.H. Chan, S.H. Chan, Z.-G. Liu, Domb's numbers and Ramanujan-Sato type series for  $1/\pi$ , *Adv. Math.* 186 (2004), 396-410.
- [4] W. Chu, X. Wang, The modified Abel lemma on summation by parts and terminating hypergeometric series identities, *Integral Transform. Spec. Funct.* 20 (2009), 93-118.
- [5] G. Gasper, M. Rahman, Basic Hypergeometric Series, Second Edition, Encyclopedia of Mathematics and Its Applications 96, Cambridge University Press, Cambridge, 2004.
- [6] I. Gessel, D. Stanton, Strange evaluations of hypergeometric series, *SIAM J. Math. Anal.* 13 (1982), 295-308.
- [7] V.J.W. Guo, X. Lian, Some  $q$ -congruences on double basic hypergeometric sums, *J. Difference Equ. Appl.* 27 (2021), 453-461.
- [8] F.H. Jackson, Summation of  $q$ -hypergeometric series, *Messenger Math.* 50 (1921), 101-112.
- [9] Z.-G. Liu, Gauss summation and Ramanujan type series for  $1/\pi$ , *Int. J. Number Theory* 8 (2012), 289-297.
- [10] L. Long, Hypergeometric evaluation identities and supercongruences, *Pacific J. Math.* 249 (2011), 405-418.
- [11] S. Ramanujan, Modular equations and approximations to  $\pi$ , *Quart. J. Math. (Oxford)* 45 (1914), 350-372.
- [12] Z.-W. Sun, A new series for  $\pi^3$  and related congruences, *Internat. J. Math.* 26 (2015), #1550055.
- [13] H. Swisher, On the supercongruence conjectures of van Hamme, *Res. Math. Sci.* 2 (2015), 1-21.
- [14] C. Wei, On two double series for  $\pi$  and their  $q$ -analogues, *Ramanujan J.* (2022). <https://doi.org/10.1007/s11139-022-00615-y>.
- [15] C. Wei, G. Ruan, Double series for  $\pi$  and their  $q$ -analogues, arXiv:2210.01331, 2022.
- [16] W. Zudilin, More Ramanujan-type formulae for  $1/\pi^2$ , *Russian Math. Surveys* 62 (2007), 634-636.

DEPARTMENT OF MATHEMATICS, SHANGHAI UNIVERSITY, SHANGHAI 200444, P. R. CHINA

*Email address:* \* Corresponding author. [guguoping0@163.com](mailto:guguoping0@163.com) (G. Gu), [xiaoxiawang@shu.edu.cn](mailto:xiaoxiawang@shu.edu.cn) (X. Wang).