

# CONTINUED FRACTIONS FOR PARTITION GENERATING FUNCTIONS

GEOFFREY B CAMPBELL

*Dedicated to Professor Rodney J Baxter on his 83rd birthday.*

ABSTRACT. We derive continued fractions for partition generating functions, utilizing both Euler's techniques and Ramanujan's techniques. Although our results are for integer partitions there is scope to extend this work to vector partitions, including for binary and n-ary partitions.

## 1. EULER'S CONTINUED FRACTION

Almost 290 years ago in 1737, Leonhard Euler wrote *De fractionibus continuis dissertatio*, which gave mathematics a first ever comprehensive account of the properties of continued fractions, and included the first proof that the number  $e$  is irrational. (See Sandifer [50]) Later, but still 275 years ago in 1748, Euler, in his *Introductio in analysin infinitorum Vol. I, Chapter 18* [28], proved:

- (a) the equivalence of his continued fraction to a generalized infinite series,
- (b) every rational number can be written as a finite continued fraction, and
- (c) the continued fraction of an irrational number is infinite.

---

2010 *Mathematics Subject Classification*. Primary: 11J70; Secondary: 05A15, 05E40, 11Y11, 11P21.

*Key words and phrases*. Continued fractions and generalizations. Exact enumeration problems, generating functions. Partitions of integers. Elementary theory of partitions. Combinatorial identities, bijective combinatorics. Lattice points in specified regions.

Thanks are due to Professor Dr Henk Koppelaar, whose discussions and suggestions have been very helpful for the book for which this paper is essentially a chapter.

Euler's continued fraction is the very nice identity, whose first few cases are:

$$\begin{aligned}
a_0 + a_0a_1 &= a_0/(1 - a_1/(1 + a_1)) \\
&= \frac{a_0}{1 - \frac{a_1}{1 + a_1}}; \\
a_0 + a_0a_1 + a_0a_1a_2 &= a_0/(1 - a_1/(1 + a_1 - a_2/(1 + a_2))) \\
&= \frac{a_0}{1 - \frac{a_1}{1 + a_1 - \frac{a_2}{1 + a_2}}}; \\
a_0 + a_0a_1 + a_0a_1a_2 + a_0a_1a_2a_3 &= a_0/(1 - a_1/(1 + a_1 - a_2/(1 + a_2 - a_3/(1 + a_3)))) \\
&= \frac{a_0}{1 - \frac{a_1}{1 + a_1 - \frac{a_2}{1 + a_2 - \frac{a_3}{1 + a_3}}}}.
\end{aligned}$$

Hence, we can state Euler's Continued Fraction in the following

**Theorem 1.1.** *If  $a_0, a_1, a_3, \dots, a_n$  are defined functions such that no denominator is zero in the following equations then*

$$\begin{aligned}
(1.1) \quad \sum_{k=0}^n \prod_{j=0}^k a_j &= a_0 + a_0a_1 + a_0a_1a_2 + \dots + a_0a_1\dots a_n \\
&= a_0/(1 - a_1/(1 + a_1 - a_2/(1 + a_2 - a_3/(1 + \dots a_{n-1}/(1 + a_{n-1} - a_n/(1 + a_n)))))). \\
&= \frac{a_0}{1 - \frac{a_1}{1 + a_1 - \frac{a_2}{1 + a_2 - \frac{a_3}{1 + a_3 - \dots \frac{a_{n-1}}{1 + a_{n-1} - \frac{a_n}{1 + a_n}}}}}}.
\end{aligned}$$

Obviously, this lends itself to many of the elementary series that arise in school and university analysis. However, we shall put this to good use in applying it to partition generating functions. The fact of this theorem involving a finite sum allows us to incrementally extend the number of terms until we can infer the infinite versions of the theorem.

Example 1: The exponential function is

$$\begin{aligned}
(1.2) \quad \exp(z) &= 1 + \frac{z}{1!} + \frac{z^2}{2!} + \frac{z^3}{3!} + \dots = 1 + \left(\frac{z}{1}\right) + \left(\frac{z}{1}\right)\left(\frac{z}{2}\right) + \left(\frac{z}{1}\right)\left(\frac{z}{2}\right)\left(\frac{z}{3}\right) + \dots \\
&= 1/\left(1 - z/\left(1 + z - \left(\frac{z}{2}\right)/\left(1 + \left(\frac{z}{2}\right) - \left(\frac{z}{3}\right)/\left(1 + \left(\frac{z}{3}\right) - \left(\frac{z}{4}\right)/\left(1 + \left(\frac{z}{4}\right) - \dots\right)\right)\right)\right)\right).
\end{aligned}$$

Applying an “equivalence transformation” that consists of clearing the fractions, this example is simplified to

$$\exp(z) = 1/(1 - z/(1 + z - z/(2 + z - 2z/(3 + z - 3z/(4 + z - \dots))))),$$

or the equivalent statement

$$\exp(z) = \frac{1}{1 - \frac{z}{1 + z - \frac{z}{2 + z - \frac{2z}{3 + z - \frac{3z}{4 + z - \dots}}}}}$$

and we know this continued fraction converges uniformly on every bounded domain in the complex plane because it is equivalent to the power series for  $\exp(z)$ .

Example 2: There is the well-known logarithmic function series

$$(1.3) \quad \log\left(\frac{1+z}{1-z}\right) = 2z\left(\frac{1}{1} + \frac{z^2}{3} + \frac{z^4}{5} + \dots\right) \\ = 2z\left(1 + \left(\frac{z^2}{3}\right) + \left(\frac{z^2}{3}\right)\left(\frac{3z^2}{5}\right) + \left(\frac{z^2}{3}\right)\left(\frac{3z^2}{5}\right)\left(\frac{5z^2}{7}\right) + \dots\right).$$

Applying Euler’s continued fraction formula to this expression shows that:

$$\log\left(\frac{1+z}{1-z}\right) \\ = 2z/(1 - (\frac{z^2}{3})/(1 + (\frac{z^2}{3}) - (\frac{3z^2}{5})/(1 + (\frac{3z^2}{5}) - (\frac{5z^2}{7})/(1 + (\frac{5z^2}{7}) - (\frac{7z^2}{9})/(1 + (\frac{7z^2}{9}) - \dots))))).$$

Applying the “equivalence transformation” this example is simplified to

$$\log\left(\frac{1+z}{1-z}\right) = 2z/(1 - z^2/(z^2 + 3 - (3z)^2/(3z^2 + 5 - (5z)^2/(5z^2 + 7 - (7z)^2/(7z^2 + 9 - \dots)))) \\ = \frac{2z}{1 - \frac{z^2}{z^2 + 3 - \frac{(3z)^2}{3z^2 + 5 - \frac{(5z)^2}{5z^2 + 7 - \frac{(7z)^2}{7z^2 + 9 - \dots}}}}}$$

Example 3: A continued fraction for  $\pi$ . We can use the previous example involving the principal branch of the natural logarithm function to construct a continued fraction representation of  $\pi$ . First we note that

$$(i+1)/(i-1) = i, \quad \text{so then} \quad \log((i+1)/(i-1)) = i\pi/2.$$

Setting  $z = i$  in the previous result, and remembering that  $i^2 = -1$ , we obtain immediately

$$\pi = \frac{4}{1 + \frac{1^2}{2 + \frac{3^2}{2 + \frac{5^2}{2 + \frac{7^2}{2 + \dots}}}}}$$

## 2. EULER'S CONTINUED FRACTION APPLIED TO PARTITIONS

In this section we will technically do no more than apply the previous section. However, the theory of partitions is full of generating functions that are amenable to the Euler continued fraction. In a subsequent section we will examine Ramanujan type continued fractions, but firstly we will gather some "low hanging fruit" from some elementary series-product identities.

We begin with the well-known telescoping identities:

If  $a_1, a_2, a_3, \dots, a_n$ , are functions chosen for nonzero denominators, then

$$(2.1) \quad 1 + \frac{a_1}{1 - a_1} + \frac{a_2}{(1 - a_1)(1 - a_2)} + \dots + \frac{a_n}{(1 - a_1)(1 - a_2)\dots(1 - a_n)}$$

$$= \frac{1}{(1 - a_1)(1 - a_2)(1 - a_3)\dots(1 - a_n)};$$

and

$$(2.2) \quad 1 + a_1 + a_2(1 + a_1) + a_3(1 + a_1)(1 + a_2) + \dots + a_n(1 + a_1)(1 + a_2)\dots(1 + a_{n-1})$$

$$= (1 + a_1)(1 + a_2)(1 + a_3)\dots(1 + a_n).$$

The series in (2.1) and (2.2) are already close to being in the required form to apply the Euler continued fraction since

$$(2.3) \quad 1 + \frac{a_1}{1 - a_1} + \frac{a_2}{(1 - a_1)(1 - a_2)} + \dots + \frac{a_n}{(1 - a_1)(1 - a_2)\dots(1 - a_n)}$$

$$= 1 + \frac{a_1}{1 - a_1} + \frac{a_1}{1 - a_1} \frac{a_2(1 - a_1)}{a_1(1 - a_2)} + \dots + \frac{a_1}{1 - a_1} \frac{a_2(1 - a_1)}{a_1(1 - a_2)} \dots \frac{a_n(1 - a_{n-1})}{a_{n-1}(1 - a_n)};$$

and

$$(2.4) \quad 1 + a_1 + a_2(1 + a_1) + a_3(1 + a_1)(1 + a_2) + \dots + a_n(1 + a_1)(1 + a_2)\dots(1 + a_{n-1})$$

$$= 1 + a_1 + a_1 \frac{a_2(1 + a_1)}{a_1} + a_1 \frac{a_2(1 + a_1)}{a_1} \frac{a_3(1 + a_2)}{a_2} + \dots + a_1 \frac{a_2(1 + a_1)}{a_1} \frac{a_3(1 + a_2)}{a_2} \dots \frac{a_n(1 + a_{n-1})}{a_{n-1}}.$$

Hence combining (2.1) with (2.3) and then (2.2) with (2.4) respectively, we obtain

$$(2.5) \quad \frac{1}{(1 - a_1)(1 - a_2)(1 - a_3)\dots(1 - a_n)}$$

$$= \frac{1}{1 - \frac{\frac{a_1}{1-a_1}}{1 + \frac{a_1}{1-a_1} - \frac{\frac{a_2(1-a_1)}{a_1(1-a_2)}}{1 + \frac{a_2(1-a_1)}{a_1(1-a_2)} - \frac{\frac{a_3(1-a_2)}{a_2(1-a_3)}}{1 + \frac{a_3(1-a_2)}{a_2(1-a_3)} - \dots - \frac{\frac{a_{n-1}(1-a_{n-2})}{a_{n-2}(1-a_{n-1})}}{1 + \frac{a_{n-1}(1-a_{n-2})}{a_{n-2}(1-a_{n-1})} - \frac{\frac{a_n(1-a_{n-1})}{a_{n-1}(1-a_n)}}{1 + \frac{a_n(1-a_{n-1})}{a_{n-1}(1-a_n)}}}}$$

and

$$(2.6) \quad (1 + a_1)(1 + a_2)(1 + a_3)\dots(1 + a_n)$$

$$= \frac{1}{1 - \frac{a_1}{1 + a_1 - \frac{\frac{a_2(1+a_1)}{a_1}}{1 + \frac{a_2(1+a_1)}{a_1} - \frac{\frac{a_3(1+a_2)}{a_2}}{1 + \frac{a_3(1+a_2)}{a_2} - \dots - \frac{\frac{a_{n-1}(1+a_{n-2})}{a_{n-2}}}{1 + \frac{a_{n-1}(1+a_{n-2})}{a_{n-2}} - \frac{\frac{a_n(1+a_{n-1})}{a_{n-1}}}{1 + \frac{a_n(1+a_{n-1})}{a_{n-1}}}}}}$$

After applying the “equivalence transformation” to both of (2.5) and then (2.6) to eliminate denominator terms, each continued fraction is simplified giving us the following two theorems.

**Theorem 2.1.** *If  $a_1, a_2, a_3, \dots, a_n$ , are functions chosen for nonzero denominators, then*

$$(2.7) \quad \frac{1}{(1 - a_1)(1 - a_2)(1 - a_3)\dots(1 - a_n)}$$

$$= \frac{1}{1 - \frac{a_1}{1 - \frac{a_2}{a_1 + a_2 - 2a_1a_2 - \frac{a_1a_3}{a_2 + a_3 - 2a_2a_3 - \dots - \frac{a_{n-2}a_n}{a_{n-1} + a_n - 2a_{n-1}a_n}}}}$$

At first glance we can see this theorem as being applicable to generating functions for unrestricted partitions of various kinds. Similarly the next theorem applies for partitions of various sorts into distinct parts.

**Theorem 2.2.** *If  $a_1, a_2, a_3, \dots, a_n$ , are functions chosen for nonzero denominators, then*

$$(2.8) \quad (1 + a_1)(1 + a_2)(1 + a_3)\dots(1 + a_n)$$

$$= \frac{1}{1 - \frac{a_1}{1 + a_1 - \frac{(1 + a_1)a_2}{a_1 + a_2 + a_1a_2 - \frac{(1 + a_2)a_3}{a_2 + a_3 + a_2a_3 - \dots \frac{(1 + a_{n-1})a_n}{a_{n-1} + a_n + a_{n-1}a_n}}}}}$$

There are many examples we could choose for substitution into theorems 2.2 and 2.2. So, let's start with the generating functions for unrestricted partitions, and for distinct partitions as follows.

**Corollary 2.1.** *If  $p_n(k)$ , is the number of unrestricted partitions of  $k$  into integers no greater than  $n$ , then*

$$(2.9) \quad \frac{1}{(1 - q^1)(1 - q^2)(1 - q^3)\dots(1 - q^n)} = \sum_{k=0}^{\infty} p_n(k)q^k$$

$$= \frac{1}{1 - \frac{q^1}{1 - \frac{q^2}{q^1 + q^2 - 2q^1q^2 - \frac{q^1q^3}{q^2 + q^3 - 2q^2q^3 - \dots \frac{q^{n-2}q^n}{q^{n-1} + q^n - 2q^{n-1}q^n}}}}}$$

**Corollary 2.2.** *If  $p_n(\mathfrak{D}, k)$ , is the number of distinct partitions of  $k$  into integers no greater than  $n$ , then*

$$(2.10) \quad (1 + q^1)(1 + q^2)(1 + q^3)\dots(1 + q^n) = \sum_{k=0}^{\infty} p_n(\mathfrak{D}, k)q^k$$

$$= \frac{1}{1 - \frac{q^1}{1 + q^1 - \frac{(1 + q^1)q^2}{q^1 + q^2 + q^1q^2 - \frac{(1 + q^2)q^3}{q^2 + q^3 + q^2q^3 - \dots \frac{(1 + q^{n-1})q^n}{q^{n-1} + q^n + q^{n-1}q^n}}}}}$$

Next we choose the odd integer powers substituted into the two theorems.

**Corollary 2.3.** *If  $p_n(\mathfrak{D}, k)$ , is the number of unrestricted partitions of  $k$  into odd integers no greater than  $2n - 1$ , then*

$$(2.11) \quad \frac{1}{(1 - q^1)(1 - q^3)(1 - q^5)\dots(1 - q^{2n-1})} = \sum_{k=0}^{\infty} p_n(\mathfrak{D}, k)q^k$$

$$= \frac{1}{1 - \frac{q^1}{1 - \frac{q^3}{1 - \frac{q^5}{q^1 + q^3 - 2q^1q^3 - \frac{q^1q^5}{q^3 + q^5 - 2q^3q^5 - \frac{\ddots}{q^{n-2}q^n}}}}}}}$$

**Corollary 2.4.** *If  $p_n(\mathfrak{D}\mathfrak{D}, k)$ , is the number of distinct partitions of  $k$  into odd integers no greater than  $2n - 1$ , then*

$$(2.12) \quad (1 + q^1)(1 + q^3)(1 + q^5)\dots(1 + q^{2n-1}) = \sum_{k=0}^{\infty} p_n(\mathfrak{D}\mathfrak{D}, k)q^k$$

$$= \frac{1}{1 - \frac{q^1}{1 + q^1 - \frac{(1 + q^1)q^3}{q^1 + q^3 + q^1q^3 - \frac{(1 + q^3)q^5}{q^3 + q^5 + q^3q^5 - \frac{\ddots}{(1 + q^{2n-3})q^{2n-1}}}}}}}$$

It is a well-known result due to Euler that  $p_{\infty}(\mathfrak{D}\mathfrak{D}, k) = p_{\infty}(\mathfrak{D}, k)$ . Explicitly, as  $n \rightarrow \infty$  equations (2.12) and (2.11) are equal to each other.

Next, let us give the cases covering binary partitions.

**Corollary 2.5.** *If  $b_n(2, k)$ , is the number of unrestricted binary partitions of  $k$  into non-negative powers of two no greater than  $2^n$ , then*

$$(2.13) \quad \frac{1}{(1 - q^1)(1 - q^2)(1 - q^4)\dots(1 - q^{2^n})} = \sum_{k=0}^{\infty} b_n(2, k)q^k$$

$$= \frac{1}{1 - \frac{q^1}{1 - \frac{q^2}{1 - \frac{q^4}{q^1 + q^2 - 2q^1q^2 - \frac{q^1q^4}{q^2 + q^4 - 2q^2q^4 - \frac{\ddots}{q^{2^{n-2}}q^{2^n}}}}}}}}$$

The following distinct binary partitions example is completely solvable.

**Corollary 2.6.** *If  $p_n(2\mathfrak{D}, k)$ , is the number of binary partitions of  $k$  into distinct non-negative powers of two no greater than  $2^n$ , then*

$$(2.14) \quad (1+q^1)(1+q^2)(1+q^4)\dots(1+q^{2^n}) = \frac{1-q^{2^{n+1}}}{1-q} = \sum_{k=0}^{2^{n+1}-1} p_n(2\mathfrak{D}, k)q^k$$

$$= \frac{1}{1 - \frac{q^1}{1 + q^1 - \frac{q^1 q^2}{q^1 + q^2 + q^1 q^2 - \frac{q^2 + q^4 + q^2 q^4 - \dots}{\dots \frac{(1 + q^{2^{n-1}})q^{2^n}}{q^{2^{n-1}} + q^{2^n} + q^{2^{n-1}} q^{2^n}}}}}}$$

Note that from (2.14) we have directly that

$$p_n(2\mathfrak{D}, k) = \begin{cases} 1, & \text{when } 0 \leq k < 2^{n+1}; \\ 0, & \text{when } k \geq 2^{n+1}. \end{cases}$$

The following distinct ternary partitions example is easily stated.

**Corollary 2.7.** *If  $p_n(3\mathfrak{D}, k)$ , is the number of ternary partitions of  $k$  into distinct non-negative powers of three no greater than  $3^n$ , then*

$$(2.15) \quad (1+q^1)(1+q^3)(1+q^9)\dots(1+q^{3^n}) = \sum_{k=0}^{3^n-1} p_n(3\mathfrak{D}, k)q^k$$

$$= \frac{1}{1 - \frac{q^1}{1 + q^1 - \frac{(1+q^1)q^3}{q^1 + q^3 + q^1 q^3 - \frac{q^3 + q^9 + q^3 q^9 - \dots}{\dots \frac{(1 + q^{3^{n-1}})q^{3^n}}{q^{3^{n-1}} + q^{3^n} + q^{3^{n-1}} q^{3^n}}}}}}$$

Note that from (2.15) we have directly that

$$p_n(3\mathfrak{D}, k) = \begin{cases} 1, & \text{for } 0 \leq k < 3^{n+1}; k \text{ is a sum of distinct powers of } 3. \\ 0, & \text{for } 0 \leq k < 3^{n+1}; k \text{ not a sum of distinct powers of } 3. \\ 0, & \text{for } k \geq 3^{n+1}. \end{cases}$$

Clearly this topic of Euler Continued Fractions applied to partition generating functions is an interesting elementary study for students, and a possible tool for researchers. The above results are old, and have probably been well-worked over time.



## 3. ROGERS-RAMANUJAN CONTINUED FRACTIONS FOR PARTITION FUNCTIONS

The fraction given here was mentioned by Ramanujan in his second letter to Hardy (see Adiga et al. [2, p. xxviii]); namely

$$(3.1) \quad R(a, b) = 1 + \frac{bq}{1 + aq + \frac{bq^2}{1 + aq^2 + \frac{bq^3}{1 + aq^3 + \frac{bq^4}{\ddots}}}}.$$

However, these now famous continued fractions, as with the Rogers-Ramanujan identities, were first discovered in 1894 by Rogers (see [49]). We define the functions  $G(q)$  and  $H(q)$  in the context of the Rogers-Ramanujan identities,

$$(3.2) \quad G(q) = \sum_{n=0}^{\infty} \frac{q^{n^2}}{(1-q)(1-q^2)\cdots(1-q^n)} = \sum_{n=0}^{\infty} \frac{q^{n^2}}{(q;q)_n} \\ = \frac{1}{(q;q^5)(q^4;q^5)} = \prod_{n=1}^{\infty} \frac{1}{(1-q^{5n-4})(1-q^{5n-1})},$$

and

$$(3.3) \quad H(q) = \sum_{n=0}^{\infty} \frac{q^{n^2+n}}{(1-q)(1-q^2)\cdots(1-q^n)} = \sum_{n=0}^{\infty} \frac{q^{n^2+n}}{(q;q)_n} \\ = \frac{1}{(q^2;q^5)(q^3;q^5)} = \prod_{n=1}^{\infty} \frac{1}{(1-q^{5n-3})(1-q^{5n-2})}.$$

The Rogers-Ramanujan continued fraction is then,

$$(3.4) \quad R(q) = \frac{q^{\frac{11}{60}H(q)}}{q^{\frac{-1}{60}G(q)}} = q^{\frac{1}{5}} \prod_{n=1}^{\infty} \frac{(1-q^{5n-4})(1-q^{5n-1})}{(1-q^{5n-3})(1-q^{5n-2})} \\ = 1 + \frac{q^{\frac{1}{5}}}{1 + \frac{q}{1 + \frac{q^2}{1 + \frac{q^3}{\ddots}}}}.$$

So, we note that  $R(0, 1)$  leads us to the celebrated Rogers-Ramanujan continued fraction, which has been researched by many (see Andrews [4, Chapter 7], for example). In the course of analyzing identities from Ramanujan's Lost Notebook [7], Andrews and Berndt have discussed the fraction  $R(a, b)$ , but mainly from the viewpoint of transformation formulas. Our emphasis here is on using (3.1) in a generalized approach to several partition identities, but there is a whole adjacent theory on particular values of these continued fractions determined from applying the theory of modular forms. Hence the examples, using  $\varphi$  as the golden ratio  $(\sqrt{5} + 1)/2$ ,

$$(3.5) \quad \frac{e^{-\frac{\pi}{5}}}{1 + \frac{e^{-\pi}}{1 + \frac{e^{-2\pi}}{1 + \frac{e^{-3\pi}}{\ddots}}}}} = \frac{1}{2}\varphi(\sqrt{5} - \varphi^{3/2})(\sqrt[4]{5} + \varphi^{3/2}),$$

$$(3.6) \quad \frac{e^{-\frac{2\pi}{5}}}{1 + \frac{e^{-2\pi}}{1 + \frac{e^{-4\pi}}{1 + \frac{e^{-6\pi}}{\ddots}}}}} = \sqrt[4]{5}\varphi^{1/2} - \varphi,$$

$$(3.7) \quad \frac{e^{-\frac{4\pi}{5}}}{1 + \frac{e^{-4\pi}}{1 + \frac{e^{-8\pi}}{1 + \frac{e^{-12\pi}}{\ddots}}}}} = \frac{1}{2}\varphi(\sqrt{5} - \varphi^{3/2})(-\sqrt[4]{5} + \varphi^{3/2}).$$

So next we examine the continued fraction  $R(a, b)$  of Ramanujan and consider various restricted partition functions. For further reading, a good reference is Alladi and Gordon [3]. We use the continued fraction to give results for several partition identities, some of which generalize results of Bressoud [12] and Göllnitz [34]. We also give a combinatorial interpretation for the coefficients in the power series expansion of the reciprocal  $\frac{1}{R(-a, -b)}$ , extending a result of Odlyzko and Wilf [42]. The full description of this approach would add several more pages to our work, but [3] covers all of this very nicely.

It turns out that **Lebesgue's identity** plays a major role in our analysis with respect to the numerators and denominators of the finite continued fractions we consider.

$$(3.8) \quad \sum_{k \geq 0} \frac{q^{k(k+1)/2} \prod_{j=1}^k (1 + bq^j)}{(1-q)(1-q^2)\dots(1-q^k)} = \prod_{m \geq 1} (1 + bq^{2m})(1 + q^m).$$

It is known that Lebesgue's identity implies Ramanujan's fraction  $R(a, b)$  has a product representation when  $a = 1$ . More precisely (3.14) and (3.15) (see below) yield

$$(3.9) \quad 1 + \frac{bq}{1 + q + \frac{bq^2}{1 + q^2 + \frac{bq^3}{1 + q^3 + \frac{bq^4}{\ddots}}}}} = \prod_{m=1}^{\infty} \frac{(1 + bq^{2m-1})}{(1 + bq^{2m})}.$$

A neat case of (3.9) is obtained from  $q \mapsto q^2$  and  $b \mapsto bq^{-1}$  so then

$$(3.10) \quad 1 + \frac{bq}{1 + q^2 + \frac{bq^3}{1 + q^4 + \frac{bq^5}{1 + q^6 + \frac{bq^7}{\ddots}}}} = \prod_{m=1}^{\infty} \frac{(1 + bq^{4m-3})}{(1 + bq^{4m-1})}.$$

For a continued fraction  $F$ , let  $P_n/Q_n$  denote its  $n$ th convergent, and suppose that  $\lim_{n \rightarrow \infty} P_n = P$ ,  $\lim_{n \rightarrow \infty} Q_n = Q$  in a suitable topology. We then say that  $F$  has numerator  $P$  and denominator  $Q$ , and write  $P = F^N$ ,  $Q = F^D$ . Consider the fraction

$$F(a, c) = 1 + a + \frac{acq}{1 + aq + \frac{acq^2}{1 + aq^2 + \frac{acq^3}{1 + aq^3 + \frac{acq^4}{\ddots}}}}.$$

This can be written in the form

$$F(a, c) = \frac{f(a, c)}{f(aq, c)}, \quad \text{where} \quad f(a, c) = \sum_{k \geq 0} A_k q^k.$$

We now compute the coefficients  $A_k = A_k(c, q)$ , observing that  $f(a, c)$  satisfies the recurrence

$$f(a, c) = (1 + a)f(aq, c) + acq f(aq^2, c).$$

Therefore the coefficients  $A_k$  satisfy

$$A_k = q^k A_k + q^{k-1} A_{k-1} q - cq^{2k-1} A_{k-1},$$

which is the same as

$$A_k = \frac{q^{k-1}(1 + cq^k)}{(1 - q^k)} A_{k-1}.$$

By iteration this yields

$$F(a, c) = \sum_{k \geq 0} \frac{a^k q^{\frac{k(k-1)}{2}} (-cq)_k}{(q)_k}.$$

Let  $c = a^{-1}b$ . Then

$$R(a, b) = \frac{f(a, a^{-1}b)}{f(aq, a^{-1}b)} - a$$

is Ramanujan's fraction (3.1).

**Lemma 3.1.** *For the fraction  $R(a, b)$ , the numerator is*

$$(3.11) \quad R^N(a, b) = \sum_{k \geq 0} \frac{a^k q^{k(k+1)/2} (-a^{-1}b)_k}{(q)_k},$$

and the denominator is

$$(3.12) \quad R^D(a, b) = \sum_{k \geq 0} \frac{a^k q^{k(k+1)/2} (-a^{-1}bq)_k}{(q)_k}.$$

*Proof:* The expansion (3.12) is an immediate consequence of

$$(3.13) \quad R^D(a, b) = f(aq, a^{-1}b).$$

The expansion (3.11) is more complicated. To obtain it, observe that

$$\begin{aligned} R^N(a, b) &= f(a, a^{-1}b) - a f(aq, a^{-1}b) \\ &= \sum_{k \geq 0} \frac{a^k q^{k(k-1)/2} (-a^{-1}bq)_k}{(q)_k} - \sum_{k \geq 0} \frac{a^{k+1} q^{k(k+1)/2} (-a^{-1}bq)_k}{(q)_k} \\ &= 1 + \sum_{k \geq 0} \frac{a^{k+1} q^{k(k+1)/2} (-a^{-1}bq)_k}{(q)_k} \left( \frac{1 + a^{-1}bq^{k+1}}{1 - q^{k+1}} - 1 \right) \\ &= 1 + \sum_{k \geq 0} \frac{a^{k+1} q^{(k+1)(k+2)/2} (-a^{-1}bq)_k (1 - a^{-1}b)}{(q)_{k+1}} \\ &= \sum_{k \geq 0} \frac{a^k q^{k(k+1)/2} (-a^{-1}b)_k}{(q)_k} \end{aligned}$$

as required. ■

Andrews (see [5] and [6]) considered the expansions in lemma 3.1 while discussing a transformation formula of Ramanujan [47] for  $R(a, b)$ . Our emphasis here is on the partition theorems that can be derived using  $R(a, b)$ , and for this the following lemma is crucial.

**Lemma 3.2.** *For the fraction  $R(a, b)$ , we also have the expansions*

$$(3.14) \quad R^N(a, b) = \sum_{i, j \geq 0} \frac{a^i b^j q^{(i^2+i)/2 + ij + j^2}}{(q)_i (q)_j},$$

and the denominator is

$$(3.15) \quad R^D(a, b) = \sum_{i, j \geq 0} \frac{a^i b^j q^{(i^2+i)/2 + ij + j^2 + j}}{(q)_i (q)_j}.$$

*Proof:* To obtain (3.14) and (3.15) from (3.12) and (3.13) we use the  $q$ -binomial theorem,

$$(-z)_k = \sum_{j=0}^k z^j q^{j(j-1)/2} \binom{k}{j}_q$$

with  $z = a^{-1}b$  and  $z = a^{-1}bq$ . (See Campbell [22] for the  $n$ -space  $q$ -binomial theorem.) Therefore

$$\begin{aligned} R^N(a, b) &= \sum_{k \geq 0} \frac{a^k q^{k(k+1)/2}}{(q)_k} \sum_{j=0}^k \frac{a^{-j} b^j q^{j(j-1)/2} (q)_k}{(q)_j (q)_{j-k}} \\ &= \sum_{i, j \geq 0} \frac{a^i b^j q^{(i+j)(i+j+1)/2}}{(q)_i (q)_j}, \end{aligned}$$

where  $i = k - j$ ; this is equivalent to (3.12). To obtain (3.13), observe that

$$(3.16) \quad R^D(a, b) = R^N(a, bq)$$

by comparing (3.14) and (3.15).

The following two theorems relate successively to the numerator and the denominator of the fraction (3.1), so then to (3.14) and (3.15). For a proof of these see Alladi and Gordon [3].

**Theorem 3.1. (Numerator)**

Let  $A^N(n; i, j)$  be the number of partitions of  $n$  into  $i + j$  distinct red parts and  $j$  distinct blue parts such that one of the blue parts may be zero and every blue part is  $\leq i + j - 1$ .

Let  $B^N(n; i, j)$  be the number of partitions of  $n$  into  $i$  distinct red parts and  $j$  distinct non-consecutive blue parts such that every red part is  $> j$ .

Let  $C^N(n; i, j)$  be the number of partitions of  $n$  into  $i$  red parts and  $j$  blue parts such that all parts are distinct and after each blue part there is a gap of at least 2. Then

$$A^N(n; i, j) = B^N(n; i, j) = C^N(n; i, j).$$

**Theorem 3.2. (Denominator)**

Let  $A^D(n; i, j)$  be as in  $A^N(n; i, j)$  except that every blue part is  $> 0$  and  $\leq i + j$ .

Let  $B^D(n; i, j)$  be as in  $B^N(n; i, j)$  except that part 1 cannot be blue.

Let  $C^D(n; i, j)$  be as in  $C^N(n; i, j)$  except that part 1 cannot be blue. Then

$$A^D(n; i, j) = B^D(n; i, j) = C^D(n; i, j).$$

So reprising (3.10) namely

$$1 + \frac{bq}{1 + q^2 + \frac{bq^3}{1 + q^4 + \frac{bq^5}{1 + q^6 + \frac{bq^7}{\ddots}}}}$$

we have interesting cancellations in numerator-denominator equations. That is, the numerator is given by

$$\begin{aligned} \sum_{k \geq 0} \frac{q^{k(k+1)}(-bq^{-1}; q^2)_k}{(q^2; q^2)_k} &= \prod_{m=1}^{\infty} (1 + bq^{4m-3})(1 + q^{2m}) \\ &= \prod_{m=1}^{\infty} (1 + bq^{4m-3})(1 + q^{4m-2})(1 + q^{4m}) \end{aligned}$$

and the denominator is given by

$$\sum_{k \geq 0} \frac{q^{k(k+1)}(-bq; q^2)_k}{(q^2; q^2)_k} = \prod_{m=1}^{\infty} (1 + bq^{4m-1})(1 + q^{4m-2})(1 + q^{4m})$$

with right sides having common factors that eliminate.

This leads in particular to the continued fraction identity

$$(3.17) \quad 1 + \frac{q}{1 + q^2 + \frac{q^3}{1 + q^4 + \frac{q^5}{1 + q^6 + \frac{q^7}{\ddots}}}} = \frac{\prod_{j \equiv 2, 3, 7 \pmod{8}} (1 - q^j)}{\prod_{j \equiv 1, 5, 6 \pmod{8}} (1 - q^j)}.$$

Göllnitz [34] states similar results, but (3.1) seems to have escaped attention. There is a continued fraction identity due to Gordon [33] and Göllnitz [34] which looks very similar to (3.17), namely

$$(3.18) \quad 1 + q + \frac{q^2}{1 + q^3 + \frac{q^4}{1 + q^5 + \frac{q^4}{1 + q^7 + \frac{bq^6}{\ddots}}}} = \frac{\prod_{j \equiv 3,4,5 \pmod{8}} (1 - q^j)}{\prod_{j \equiv 1,4,7 \pmod{8}} (1 - q^j)}.$$

However, this result first appears in Alladi and Gordon [3] almost 30 years after (3.1).

#### 4. RAMANUJAN'S THREE PARAMETER CONTINUED FRACTION

Ramanujan [45] obtained in addition to (3.1), the following continued fraction with three parameters  $a, b, q$  which has also a product representation

$$(4.1) \quad 1 - ab + \frac{(a - bq)(b - aq)}{(1 - ab)(1 + q^2) + \frac{(a - bq^3)(b - aq^3)}{(1 - ab)(1 + q^4) + \frac{(a - bq^5)(b - aq^5)}{(1 - ab)(1 + q^6) + \frac{(a - bq^7)(b - aq^7)}{\ddots}}}} = \prod_{m=1}^{\infty} \frac{(1 + a^2q^{4m-3})(1 + b^2q^{4m-3})}{(1 + a^2q^{4m-1})(1 + b^2q^{4m-1})}.$$

This was proved only in 1985 by the reviewers of Chapter 16 of Ramanujan's Second Notebook [2], 65 years after Ramanujan's death. If we put  $a = 0$  and replace  $b^2$  by  $-b$  in (4.1), we get (3.10). It seems there is still scope to study the combinatorial properties of the coefficients in the power series expansion of this fraction.

#### REFERENCES

- [1] ABRAMOWITZ, M., and STEGUN, I. Handbook of Mathematical Functions, Dover Publications Inc., New York, 1972.
- [2] ADIGA, C. BERNDT, B. C. BHARGAVA, S. AND WATSON, G. N. "Chapter 16 of Ramanujan's Second Notebook: Theta Functions and q-Series", Memoirs of the American Mathematical Society, Vol. 315, Amer. Math. Soc., Providence, RI, 1985.
- [3] ALLADI, K. and GORDON H., *Partition Identities and a Continued Fraction of Ramanujan*, Journal of Combinatorial Theory, Series A 63, 275-300 (1993)
- [4] ANDREWS, G.E. The Theory of Partitions, Addison-Wesley Publishing Company, Advanced Book Program, Reading, Massachusetts, 1976.
- [5] ANDREWS, G. E. *An introduction to Ramanujan's "lost" notebook*, Amer. Math. Monthly 86 (1979), 89-108.
- [6] ANDREWS, G. E. *Ramanujan's "Lost" Notebook. III. The Rogers-Ramanujan continued fraction*, Adv. Math. 41 (1981), 186-208.
- [7] ANDREWS, G. E., and BERNDT, B. C. *Ramanujan's Lost Notebook: Part V* Paperback (2018). Springer-Verlag, New York, ISBN-13: 978-3030085506.
- [8] ANDREWS, G.E. and ERIKSSON, K. *Integer Partitions*, Cambridge University Press, Cambridge, UK, New York, USA, Port Melbourne, Australia, Madrid, Spain, Cape Town, South Africa, 2004.

- [9] APOSTOL, T. *Introduction to Analytic Number Theory*, Springer-Verlag, New York, 1976.
- [10] BAXTER, R. J. *Exactly Solved Models in Statistical Mechanics*, Academic Press, New York, 1982.
- [11] BIRKHOFF, G. and MACLAINE, S. *A survey of modern algebra*, fourth ed., N.Y., Macmillan, 1977.
- [12] BRESSOUD, D.M. *On a partition theorem of Göllnitz*, *J. Reine Angew. Math.* **305** 215-217, (1979).
- [13] CAMPBELL, G. B. *Generalization of a Formula of Hardy*, Pure Math. Research Paper 79-5, La Trobe University, Melbourne, Australia, 1979.
- [14] CAMPBELL, G. B. *Multiplicative functions over Riemann zeta function products*, *J. Ramanujan Soc.* 7 No. 1, 1992, 52-63.
- [15] CAMPBELL, G. B. *Dirichlet summations and products over primes*, *Int. J. Math. Math. Sci.*, Vol 16, No 2, (1993) 359-372.
- [16] CAMPBELL, G. B. *A generalized formula of Hardy*, *Int. J. Math. Math. Sci.*, Vol 17, No 2, (1994) 369-378.
- [17] CAMPBELL, G. B. *A new class of infinite products, and Euler's totient*, *International Journal of Mathematics and Mathematical Sciences*, vol. 17, no. 3, pp. 417-422, 1994. <https://doi.org/10.1155/S0161171294000591>.
- [18] CAMPBELL, G. B. *Infinite products over visible lattice points*, *International Journal of Mathematics and Mathematical Sciences*, vol. 17, no. 4, pp. 637-654, 1994. <https://doi.org/10.1155/S0161171294000918>.
- [19] CAMPBELL, G. B. *Combinatorial identities in number theory related to  $q$ -series and arithmetical functions*, Doctor of Philosophy Thesis, School of Mathematical Sciences, The Australian National University, October 1997.
- [20] CAMPBELL, G. B. *A closer look at some new identities*, *International Journal of Mathematics and Mathematical Sciences*, vol. 21, no. 3, pp. 581-586, 1998. <https://doi.org/10.1155/S0161171298000805>.
- [21] CAMPBELL, G. B. *Infinite products over hyperpyramid lattices*, *International Journal of Mathematics and Mathematical Sciences*, vol. 23, no. 4, pp. 271-277, 2000. <https://doi.org/10.1155/S0161171200000764>.
- [22] CAMPBELL, G. B. *Some  $n$ -space  $q$ -binomial theorem extensions and similar identities*, arXiv:1906.07526v1 [math.NT], Jun 2019. (<https://arxiv.org/abs/1906.07526>)
- [23] CAMPBELL, G. B. *An interview with Rodney James Baxter*, *Aust. Math. Soc. Gazette*, Volume 47, No1, pp24-32, March 2020. (<https://austms.org.au/wp-content/uploads/2020/07/471Web.pdf>)
- [24] CAMPBELL, G. B. *Fun with numbers: Rational solutions to  $x^y y^x = v^w w^v$* , *Aust. Math. Soc. Gazette*, Volume 49, No5, pp210-211, November 2022. (<https://austms.org.au/publications/gazette/gazette495/>)
- [25] CAUCHY, A. *Mémoire sur les fonctions dont plusieurs . . .*, *C. R. Acad. Sci. Paris*, T. XVII, p. 523, *Oeuvres de Cauchy*, 1re série, T. VIII, Gauthier-Villars, Paris, 1893, 42- 50.
- [26] CHEEMA, M. S., *Vector partitions and combinatorial identities*, *Math. Comp.* 18, 1966 414-420.
- [27] CHEEMA, M. S. and MOTZKIN, T. S., *Multipartitions and multipermutations*, *Proc. Symp. Pure Math.* 19, 1971, 37-39.
- [28] EULER, L. *Introductio in analysin infinitorum*, Chapter 16. Marcum-Michaelum, Brousquet, Lausannae (1748).
- [29] GASPER, G. and RAHMAN, M. *Basic Hypergeometric Series*, *Encyclopedia of Mathematics and its Applications*, Vol 35, Cambridge University Press, (Cambridge - New York - Port Chester - Melbourne - Sydney), 1990.
- [30] GAUSS, C.F. *Disquisitiones generales circa seriem infinitam . . .*, *Comm. soc. reg. sci. Gött. rec.*, Vol II; reprinted in *Werke* 3 (1876), pp. 123-162.
- [31] GOLDFELD, D. *Beyond the last theorem*. *Math Horizons.* 4 (September): 26-34. (1996). doi:10.1080/10724117.1996.11974985. JSTOR 25678079.
- [32] GORDON, B. *Two theorems on multipartite partitions*, *J. London Math. Soc.* 38, 1963, 459-464.
- [33] GORDON, B. *Some continued fractions of the Rogers-Ramanujan type*, *Duke Math. J.* 32 (1965), 741-748.

- [34] GÖLLNITZ, H. *Partitionen mit Differenzenbedingungen*, J. Reine Angew. Math. 225 (1967), 154-190.
- [35] HARDY, G. H. *An extension of a theorem on oscillating series*, Collected Papers, Vol VI, Clarendon Press, Oxford, 1974, 500-506.
- [36] HARDY, G. H. *On certain oscillating series*, Collected Papers, Vol VI, Clarendon Press, Oxford, 1974, 146-167.
- [37] HARDY, G. H., and LITTLEWOOD, J. E. *A further note on the converse of Abel's theorem*. Collected Papers of Hardy, Vol VI, Clarendon Press, Oxford, 1974, 699-716.
- [38] HEINE, E. *Untersuchungen über die Reihe ...*, J. Reine angew. Math. 34, 1847, 285-328.
- [39] HEINE, E. *Handbuch der Kugelfunctionen, Theorie und Anwendungen*, Vol. 1, Reimer, Berlin, 1878.
- [40] MACDONALD, I. G. *Symmetric Functions And Hall Polynomials*, 2nd ed., Oxford : Clarendon Press ; New York : Oxford University Press, 1995.
- [41] MASSER, D. W. (1985). "Open problems". In Chen, W. W. L. (ed.). *Proceedings of the Symposium on Analytic Number Theory*. London: Imperial College.
- [42] ODLYZKO, A. M. and WILF, H. S. *n coins in a fountain*, Amer. Math. Monthly 95 (1988), 840-843.
- [43] OESTERLÉ, J. *Nouvelles approches du "théorème" de Fermat*, Astérisque, Séminaire Bourbaki exp 694 (161): 165–186, (1988), ISSN 0303-1179, MR 0992208.
- [44] RAMANUJAN, S. (1927) *Collected Papers of S. Ramanujan*, Cambridge University Press, Cambridge (1927); reprinted by Chelsea, New York, 1962.
- [45] RAMANUJAN, S. "Notebooks (Two Volumes)," Tata Institute, Bombay, 1957.
- [46] RAMANUJAN, S. *On certain trigonometrical sums and their application to the theory of numbers*, Collected Papers of S. Ramanujan, Cambridge University Press, Cambridge (1927), 179-199; reprinted by Chelsea, New York, 1962.
- [47] RAMANUJAN, S. "The Lost Notebook, and Other Unpublished Papers," Narosa, New Delhi, 1988.
- [48] RIEMANN, G. F. B. "Über die Anzahl der Primzahlen unter einer gegebenen Grösse." *Monatsber. Königl. Preuss. Akad. Wiss. Berlin*, 671-680, Nov. 1859.
- [49] ROGERS, L. J. (1894). "Second memoir on the expansion of certain infinite products". *Proc. London Math. Soc.* 25: 318-343.
- [50] SANDIFER, C. E. (2006). "Chapter 32: Who proved e is irrational?". *How Euler Did It* (PDF). Mathematical Association of America. pp. 185–190. ISBN 978-0-88385-563-8. LCCN 2007927658
- [51] SLOANE, N. J. A., *The On-Line Encyclopedia of Integer Sequences (OEIS) Euler transform*. [https://oeis.org/wiki/Euler\\_transform](https://oeis.org/wiki/Euler_transform).
- [52] SLOANE, N. J. A., *The On-Line Encyclopedia of Integer Sequences (OEIS) sequence A061159 Numerators in expansion of Euler transform of  $b(n)=1/2$*  <https://oeis.org/A061159>.
- [53] SLOANE, N. J. A., *The On-Line Encyclopedia of Integer Sequences (OEIS) sequence A061160 Numerators in expansion of Euler transform of  $b(n)=1/3$*  <https://oeis.org/A061160>.
- [54] SZPIRO, L. (1981). "Propriétés numériques du faisceau dualisant relatif". *Seminaire sur les pinceaux des courbes de genre au moins deux* (PDF). Astérisque. Vol. 86. pp. 44–78. Zbl 0517.14006.
- [55] SZPIRO, L. (1987), "Présentation de la théorie d'Arakelov", *Contemp. Math., Contemporary Mathematics*, 67: 279–293, doi:10.1090/conm/067/902599, ISBN 9780821850749, Zbl 0634.14012
- [56] WRIGHT, E. M. *Partitions of multipartite numbers*, *Proc. Amer. Math. Soc.* 28, 1956, 880-890.

MATHEMATICAL SCIENCES INSTITUTE, THE AUSTRALIAN NATIONAL UNIVERSITY, CANBERRA, ACT, 0200, AUSTRALIA

*Email address:* [Geoffrey.Campbell@anu.edu.au](mailto:Geoffrey.Campbell@anu.edu.au)