

ALGEBRAIC IDENTITIES ON Q-HARMONIC NUMBERS AND Q-BINOMIAL COEFFICIENTS

SAID ZRIAA AND MOHAMMED MOUÇOUF

ABSTRACT. The aim of this paper is to present a general algebraic identity. Applying this identity, we provide several formulas involving the q-binomial coefficients and the q-harmonic numbers. We also recover some known identities including an algebraic identity of D. Y. Zheng on q-Apéry numbers and we establish the q-analog of Euler's formula. The proposed results may have important applications in the theory of q-supercongruences.

1. INTRODUCTION

For an indeterminate x the q-shifted factorial is usually defined by

$$(x; q)_0 = 1 \quad \text{and} \quad (x; q)_n = (1-x)(1-qx)\cdots(1-q^{n-1}x) \quad \text{for } n = 1, 2, \dots$$

The Gaussian q-binomial coefficient is correspondingly given by

$$\begin{bmatrix} n \\ j \end{bmatrix} = \frac{(q; q)_n}{(q; q)_j (q; q)_{n-j}} \quad \text{for } j = 0, 1, 2, \dots, n,$$

The q-harmonic numbers are defined by

$$\mathcal{H}_n(q) = \sum_{k=1}^n \frac{1}{[k]_q} \quad \text{for } n = 1, 2, \dots,$$

where the q-numbers are given by

$$[k] \quad \text{or} \quad [k]_q := \frac{1-q^k}{1-q} = 1 + q + \cdots + q^{k-1}$$

Following Comtet [7], the complete Bell polynomials can be explicitly expressed as

$$\mathbf{B}_n(x_1, x_2, \dots, x_n) = \sum_{m_1+2m_2+\cdots+nm_n=n} \frac{n!}{m_1!m_2!\cdots m_n!} \left(\frac{x_1}{1!}\right)^{m_1} \left(\frac{x_2}{2!}\right)^{m_2} \cdots \left(\frac{x_n}{n!}\right)^{m_n}$$

During the last two decades, there has been an increasing interest in studying binomial sums and their q-analogues, one can consult recent papers, for example [6, 18, 19]. For a comprehensive account of the q-series and its applications to numbers theory, combinatorics and special functions, we refer the reader to the excellent monograph by G. Gasper and M. Rahman [8]. Recently, the study of the q-harmonic congruences turned into a very active research area (see e.g., [17] and the references therein).

There exist numerous combinatorial identities involving q-binomial coefficients in the mathematical literature (see e.g., [6, 12, 16, 18, 19]). Nowadays, there has been growing interest in deriving q-analogues of several combinatorial identities. This includes, for example, q-generalizations of some well known identities involving harmonic numbers.

The Apéry numbers are defined by the following binomial sum

$$A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$$

Key words and phrases. Algebraic identities, q-binomial coefficients, q-harmonic numbers, complete Bell polynomials.

These numbers have many interesting properties that make them extremely useful in the proof of the irrationality of $\zeta(3)$ (see [3] for further details). Also the Apéry numbers have remarkable arithmetic properties [9]. F. Beukers conjectured [4] that

$$A\left(\frac{p-1}{2}\right) \equiv a(p) \pmod{p^2}$$

where p is an odd prime and $a(n)$ is determined by

$$\sum_{k=1}^{\infty} a(k)q^k := q \prod_{k=1}^{\infty} (1 - q^{2k})^4 (1 - q^{4k})^4 = q - 4q^3 - 2q^5 + 24q^7 + \dots$$

Beukers' conjecture was later showed by S. Ahlgren, K. Ono [1], who reduce this statement in terms of the harmonic numbers H_n to the identity

$$\sum_{k=1}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left(1 + 2kH_{n+k} + 2kH_{n-k} - 4kH_k\right) = 0$$

In order to give a classical proof of the last identity, Chu [5] presented the following algebraic identity

$$\sum_{j=0}^n \binom{n}{j}^2 \binom{n+j}{j}^2 \left(\frac{-j}{(x+j)^2} + \frac{1 + 2j(H_{n+j} - H_j) + 2j(H_{n-j} - H_j)}{x+j} \right) = \frac{x(1-x)_n^2}{(x)_{n+1}^2}$$

which gives the desired formula in the limit.

In order to prove irrationality results on the q -analog of $\zeta(3)$:

$$\zeta_q(3) = \sum_{k=1}^{+\infty} \frac{q^k(1+q^k)}{(1-q^k)^3}$$

The authors of [13] introduced implicitly a q -analog of the Apéry numbers $A_q^{KRZ}(n)$ and they showed that

$$A_q^{KRZ}(n) = \sum_{k=0}^n \frac{a_q(n, k)}{q^k}$$

where $a_q(n, k)$ can be defined via the following q -partial fraction decomposition

$$\frac{(xq^{-n}; q)_n^2}{(x; q)_{n+1}^2} = \sum_{k=0}^n \left(\frac{a_q(n, k)}{(1-q^kx)^2} + \frac{b_q(n, k)}{1-q^kx} \right) \quad (1.1)$$

A. Straub [17] showed via the partial fraction decomposition technique that $A_q^{KRZ}(n)$ have the following explicit q -binomial representation

$$q^{n(2n+1)} A_q^{KRZ}(n) = \sum_{k=0}^n q^{(n-k)^2} \begin{bmatrix} n \\ k \end{bmatrix}^2 \begin{bmatrix} n+k \\ k \end{bmatrix}^2$$

which reduces to Apéry numbers when $q \rightarrow 1$.

D. Y. Zheng [20] has recently introduced the q -Apéry numbers

$$q^{n(n+1)} A_q^{KRZ}(n) = \sum_{k=0}^n q^{k(k-2n)} \begin{bmatrix} n \\ k \end{bmatrix}^2 \begin{bmatrix} n+k \\ k \end{bmatrix}^2$$

then he established an interesting algebraic identity

$$\frac{x^{2n}(q/x; q)_n^2}{(1-x)(xq; q)_n^2} = \frac{1}{1-x} + \sum_{j=1}^n \begin{bmatrix} n \\ j \end{bmatrix}^2 q^{j(j-2n)} \left(\frac{q^j - 1}{(1-xq^j)^2} + \frac{1 - 4[j]\mathcal{H}_j(q) + 2[j]\mathcal{H}_{n+j}(q) + 2q[j]\mathcal{H}_{n-j}(q^{-1})}{1-xq^j} \right) \quad (1.2)$$

which is a q -extension of the Chu's identity, and obtained the identity

$$\sum_{k=0}^n q^{k(k-2n)} \begin{bmatrix} n \\ k \end{bmatrix}^2 \begin{bmatrix} n+k \\ k \end{bmatrix}^2 \left(2\mathcal{H}_k(q) - \mathcal{H}_{n+k}(q) - q\mathcal{H}_{n-k}(q^{-1}) \right) = 0$$

The purpose of this paper is to establish and develop important algebraic identities involving q-harmonic numbers and q-binomial coefficients, which may have important applications in the theory of q-supercongruences.

2. SOME IDENTITIES OF Q-BINOMIAL COEFFICIENTS

We now state and prove one of the main results of this paper.

Theorem 2.1. *Let $\alpha_1, \alpha_2, \dots, \alpha_s$ be distinct elements of \mathbb{C} . For a positive integer m let $P(x) = (x - \alpha_1)^m (x - \alpha_2)^m \cdots (x - \alpha_s)^m$. For any polynomial $Q(x)$ such that $\deg(Q) < \deg(P)$, we have*

$$\frac{Q(x)}{P(x)} = \sum_{j=1}^s \sum_{i=0}^{m-1} \sum_{k=0}^{m-1-i} \frac{(-1)^k g_j(\alpha_j) \mathbf{B}_k(x_1, \dots, x_k) Q^{(i)}(\alpha_j)}{i! k! (x - \alpha_j)^{m-i-k}}. \quad (2.1)$$

where

$$x_l = m(l-1)! \sum_{i=1, i \neq j}^s \frac{1}{(\alpha_j - \alpha_i)^l} \quad \text{and} \quad g_j(x) = \prod_{i=1, i \neq j}^s (x - \alpha_i)^{-m}.$$

Proof. Following [14, Eq.4], we can write

$$Q(x) = \sum_{j=1}^s \sum_{i=0}^{m-1} \frac{1}{i!} Q^{(i)}(\alpha_j) L_{ji}(x)[P].$$

By virtue of Equation (2) of [14], it is clear that

$$L_{ji}(x)[P] = P(x) \sum_{k=0}^{m-1-i} \frac{g_j^{(k)}(\alpha_j)}{k! (x - \alpha_j)^{m-i-k}}$$

If we combine the two previous identities, we get

$$\frac{Q(x)}{P(x)} = \sum_{j=1}^s \sum_{i=0}^{m-1} \sum_{k=0}^{m-1-i} \frac{g_j^{(k)}(\alpha_j) Q^{(i)}(\alpha_j)}{i! k! (x - \alpha_j)^{m-i-k}}.$$

Since

$$g_j(x) = \phi(x) \circ f_j(x)$$

where $\phi(x) = \exp(mx)$ and $f_j(x) = \ln(\prod_{i=1, i \neq j}^s (x - \alpha_i)^{-1})$. Then $\phi^{(k)}(x) = m^k \exp(mx)$ and $f_j^{(k)}(x) = (-1)^k (k-1)! \mathcal{H}_{k, \alpha_s[j]}(x)$, where $\mathcal{H}_{l, \alpha_s[j]}(x) = \sum_{i=1, i \neq j}^s \frac{1}{(x - \alpha_i)^l}$. By using the Faà di Bruno formula, we can easily prove that

$$g_j^{(k)}(x) = (-1)^k g_j(x) \sum_{m_1+2m_2+\dots+km_k=k} \frac{k!}{m_1! m_2! \cdots m_k!} \prod_{l=1}^k \left(\frac{m(l-1)! \mathcal{H}_{l, \alpha_s[j]}(x)}{l!} \right)^{m_l}$$

In particular

$$g_j^{(k)}(\alpha_j) = (-1)^k g_j(\alpha_j) \mathbf{B}_k(x_1, \dots, x_k)$$

This gives the required result. \square

Taking $\alpha_i = q^{-i}$ in Theorem (2.1), we obtain after some minor manipulations the following theorem.

Theorem 2.2. *Let m and n positive integers and let $Q(x)$ be a polynomial such that $\deg(Q) < (n+1)m$, we have*

$$\frac{(q; q)_n^m Q(x)}{(x; q)_{n+1}^m} = \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^m q^{m \binom{j+1}{2}} \sum_{i=0}^{m-1} \sum_{k=0}^{m-1-i} \frac{(-1)^{mj+i} \mathbf{B}_k(x_1, x_2, \dots, x_k) Q^{(i)}(q^{-j})}{i! k! q^{j(i+k)} (1 - xq^j)^{m-i-k}}.$$

where

$$x_1 = \frac{q^j m}{1 - q} \left(\mathcal{H}_j(q) - q \mathcal{H}_{n-j}(q^{-1}) \right)$$

and

$$x_l = m(l-1)! \sum_{i=0, i \neq j}^n \frac{q^{jl}}{(1-q^{j-i})^l} \text{ for } l = 1, 2, \dots, m-1,$$

In view of Theorem (2.2), we establish interesting corollaries

Corollary 2.3. *Let n be a positive integer and let $Q(x)$ be a polynomial such that $\deg(Q) < 2(n+1)$. We have*

$$\frac{(q; q)_n^2 Q(x)}{(x; q)_{n+1}^2} = \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^2 q^{j(j+1)} \left(\frac{Q(q^{-j})}{(1-xq^j)^2} - \frac{q^{-j}Q'(q^{-j})}{(1-xq^j)} + \frac{2Q(q^{-j})}{(1-xq^j)} \left(\frac{\mathcal{H}_j(q) - q\mathcal{H}_{n-j}(q^{-1})}{1-q} \right) \right)$$

Identity 2.4. *Setting $Q(x) = 1$ in the last corollary, we obtain*

$$\frac{(q; q)_n^2}{(x; q)_{n+1}^2} = \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^2 q^{j(j+1)} \left(\frac{1}{(1-xq^j)^2} + \frac{2}{1-xq^j} \left(\frac{\mathcal{H}_j(q) - q\mathcal{H}_{n-j}(q^{-1})}{1-q} \right) \right)$$

In particular

$$(q; q)_n^2 = \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^2 q^{j(j+1)} \left(1 + 2 \left(\frac{\mathcal{H}_j(q) - q\mathcal{H}_{n-j}(q^{-1})}{1-q} \right) \right)$$

Identity 2.5.

$$\frac{x^{2n}(q/x; q)_n^2}{(1-x)(xq; q)_n^2} = \frac{1}{1-x} + \sum_{j=1}^n \begin{bmatrix} n \\ j \end{bmatrix}^2 q^{j(j-2n)} \left(\frac{q^j - 1}{(1-xq^j)^2} + \frac{1 - 4[j]\mathcal{H}_j(q) + 2[j]\mathcal{H}_{n+j}(q) + 2q[j]\mathcal{H}_{n-j}(q^{-1})}{1-xq^j} \right)$$

This identity recovers Zheng identity (1.2).

Proof. Let

$$Q(x) = (1-x)(x-q)^2 \cdots (x-q^n)^2 = (1-x)x^{2n}(q/x; q)_n^2$$

It is not difficult to verify that

$$\frac{Q(x)}{(x; q)_{n+1}^2} = \frac{x^{2n}(q/x; q)_n^2}{(1-x)(xq; q)_n^2}, \quad Q(q^{-j}) = q^{-j(2n+1)}(q^j - 1)(q; q)_n^2 \begin{bmatrix} n+j \\ j \end{bmatrix}^2$$

and

$$Q'(q^{-j}) = q^{-2nj}(q; q)_n^2 \begin{bmatrix} n+j \\ j \end{bmatrix}^2 \left(-1 - 2[j] \left(\mathcal{H}_{n+j}(q) - \mathcal{H}_j(q) \right) \right)$$

Applying Corollary (2.3), we get after some simplifications the desired identity. \square

Identity 2.6.

$$\frac{(xq^{-n}; q)_n^2}{(x; q)_{n+1}^2} = \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^2 \begin{bmatrix} n+j \\ j \end{bmatrix}^2 q^{j(j+1)-n(n+2j+1)} \left(\frac{1}{(1-xq^j)^2} + \frac{4\mathcal{H}_j(q) - 2q\mathcal{H}_{n-j}(q^{-1}) - 2\mathcal{H}_{n+j}(q)}{(1-q)(1-xq^j)} \right)$$

This identity gives the explicit representation of (1.1).

Proof. Let $Q(x) = (xq^{-n}; q)_n^2$. We have $Q(q^{-j}) = q^{-n(n+2j+1)} \begin{bmatrix} n+j \\ j \end{bmatrix}^2$ and

$$Q'(q^{-j}) = 2q^{-n(n+2j+1)+j} \begin{bmatrix} n+j \\ j \end{bmatrix}^2 \left(\frac{\mathcal{H}_{n+j}(q) - \mathcal{H}_j(q)}{1-q} \right)$$

Using Corollary (2.3), the identity follows. \square

Identity 2.7. *Letting $x = 0$ in the last identity, we obtain*

$$\sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^2 \begin{bmatrix} n+j \\ j \end{bmatrix}^2 q^{j(j+1)-2nj} \left(1 - q + 4\mathcal{H}_j(q) - 2q\mathcal{H}_{n-j}(q^{-1}) - 2\mathcal{H}_{n+j}(q) \right) = q^{n(n+1)}(1-q)$$

Corollary 2.8. Let m, n, l be tree positive integers such that $0 \leq l < (n+1)m$. Then we have

$$\frac{(q; q)_n^m x^l}{(x; q)_{n+1}^m} = \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^m q^{m \binom{j+1}{2}} \sum_{i=0}^{m-1} \sum_{k=0}^{m-1-i} \binom{l}{i} \frac{(-1)^{mj+i} \mathbf{B}_k(x_1, x_2, \dots, x_k)}{k! q^{j(k+l)} (1-xq^j)^{m-i-k}}.$$

where

$$x_l = m(l-1)! \sum_{i=0, i \neq j}^n \frac{q^{jl}}{(1-q^{j-i})^l}$$

Corollary 2.9. Let m and n be positive integers. We have

$$\frac{(q; q)_n^m}{(x; q)_{n+1}^m} = \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^m q^{m \binom{j+1}{2}} \sum_{k=0}^{m-1} \frac{(-1)^{mj} \mathbf{B}_k(x_1, x_2, \dots, x_k)}{k! q^{jk} (1-xq^j)^{m-k}}.$$

where

$$x_l = m(l-1)! \sum_{i=0, i \neq j}^n \frac{q^{jl}}{(1-q^{j-i})^l}$$

Corollary 2.10. Let $Q(x)$ be a polynomial such that $\deg(Q) < n+1$. Then we have

$$\frac{(q; q)_n Q(x)}{(x; q)_{n+1}} = \sum_{j=0}^n (-1)^j \begin{bmatrix} n \\ j \end{bmatrix} q^{\binom{j+1}{2}} \frac{Q(q^{-j})}{(1-xq^j)}.$$

The limiting case of Theorem (2.2), is the following

Theorem 2.11. Let m and n be two positive integers. Let $Q(x)$ be a polynomial of degree l with leading coefficient a_l . Then we have the following curious identity:

$$\sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^m q^{m \binom{j}{2}} \sum_{i=0}^{m-1} \frac{(-1)^{mj+i+1} \mathbf{B}_{m-1-i}(x_1, x_2, \dots, x_{m-1-i}) Q^{(i)}(q^{-j})}{i!(m-1-i)!} = \begin{cases} 0 & \text{if } 0 \leq l < m(n+1) - 1, \\ (-1)^{m(n+1)} (q; q)_n^m q^{-m \binom{n+1}{2}} a_l & \text{if } l = m(n+1) - 1. \end{cases}$$

where

$$x_l = m(l-1)! \sum_{i=0, i \neq j}^n \frac{q^{jl}}{(1-q^{j-i})^l}$$

When $m = 1$, the formula of Theorem (2.11) reads explicitly as

Corollary 2.12. Let n be a positive integer and $Q(x)$ be a polynomial of degree l with leading coefficient a_l . Then we have the following identity:

$$\sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix} q^{\binom{j}{2}} (-1)^j Q(q^{-j}) = \begin{cases} 0 & \text{if } 0 \leq l < n, \\ (-1)^n (q; q)_n q^{-\binom{n+1}{2}} a_l & \text{if } l = n. \end{cases}$$

Identity 2.13. Letting $Q(x) = (x-1)^l$ in the last corollary, we obtain, after some simple calculations, that

$$\sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix} q^{\binom{j}{2} - jl} (-1)^j [j]^l = \begin{cases} 0 & \text{if } 0 \leq l < n, \\ (-1)^n \frac{(q; q)_n}{(1-q)^n} q^{-\binom{n+1}{2}} & \text{if } l = n. \end{cases}$$

We remark that in the limiting case $q \rightarrow 1$ of this identity, we obtain the famous Euler's formula[2, 10, 11, 15]. Therefore, this identity is the q -analog of Euler's formula:

$$\sum_{j=0}^n \binom{n}{j} (-1)^j j^l = \begin{cases} 0 & \text{if } 0 \leq l < n, \\ (-1)^n n! & \text{if } l = n. \end{cases}$$

Opting $m = 2$ in Theorem (2.11), we gain the following important result

Corollary 2.14. *Let n be a positive integer and $Q(x)$ be a polynomial of degree l with leading coefficient a_l . Then the following identity holds:*

$$\sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^2 q^{j(j-1)} \left(Q'(q^{-j}) - \frac{2q^j Q(q^{-j})}{1-q} \left(\mathcal{H}_j(q) - q\mathcal{H}_{n-j}(q^{-1}) \right) \right) = \begin{cases} 0 & \text{if } 0 \leq l < 2n+1, \\ (q; q)_n^2 q^{-n(n+1)} a_l & \text{if } l = 2n+1. \end{cases}$$

Identity 2.15. *If $Q(x) = 1$, then the formula of Corollary (2.14) gives*

$$\sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^2 q^{j^2} \left(\mathcal{H}_j(q) - q\mathcal{H}_{n-j}(q^{-1}) \right) = 0$$

Identity 2.16. *Choosing $Q(x) = (1-x)(x-q)^2 \cdots (x-q^n)^2$ in Corollary (2.14), we obtain*

$$\sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}^2 \begin{bmatrix} n+j \\ j \end{bmatrix}^2 q^{j(j-1)-2nj} \left(1 + 2[j]\mathcal{H}_{n+j}(q) - 4[j]\mathcal{H}_j(q) + 2q\mathcal{H}_{n-j}(q^{-1}) \right) = q^{-n(n+1)}$$

It would be interesting to establish some identities by means of Corollary (2.10).

Theorem 2.17. *Let n be a positive integer and y any complex number. Then the following identity holds true:*

$$\sum_{j=0}^n (-1)^j \begin{bmatrix} n \\ j \end{bmatrix} q^{\binom{j+1}{2}-jn} \frac{(y; q)_{n+j}}{(y; q)_j (1-xq^j)} = \frac{(q; q)_n (y/x; q)_n x^n}{(x; q)_{n+1}}.$$

Proof. Let $Q(x)$ be the polynomial

$$Q(x) = (x-y)(x-xy) \cdots (x-yq^{n-1}) = (y/x; q)_n x^n$$

Since

$$Q(q^{-j}) = (yq^j; q)_n q^{-jn} \quad \text{and} \quad (yq^j; q)_n = \frac{(y; q)_{n+j}}{(y; q)_j}$$

we get from Corollary (2.10) the desired formula. \square

Letting $y = q$ in the last theorem, we obtain the following result

Theorem 2.18. *Let n be a positive integer, the following identities hold true*

(1)

$$\sum_{j=0}^n (-1)^j \begin{bmatrix} n \\ j \end{bmatrix} \begin{bmatrix} n+j \\ j \end{bmatrix} \frac{q^{\binom{j+1}{2}-jn}}{(1-xq^j)} = \frac{x^n (q/x; q)_n}{(x; q)_{n+1}}.$$

(2)

$$\sum_{j=0}^n (-1)^j \begin{bmatrix} n \\ j \end{bmatrix} \begin{bmatrix} n+j \\ j \end{bmatrix} \frac{q^{\binom{j+1}{2}-jn}}{(1-q^{j+l})} = 0.$$

for $l = 1, 2, \dots, n$.

(3)

$$\sum_{j=0}^{n-1} (-1)^{n+1+j} \begin{bmatrix} n \\ j \end{bmatrix} \begin{bmatrix} n+j \\ j \end{bmatrix} \frac{(1-q^{n+l})}{(1-q^{j+l})} q^{\binom{j+1}{2}+n^2-jn} = \begin{bmatrix} 2n \\ n \end{bmatrix} q^{\binom{n+1}{2}}.$$

for $l = 1, 2, \dots, n$.

(4)

$$q^{\binom{n+1}{2}} = \sum_{j=0}^n (-1)^{n-j} \begin{bmatrix} n \\ j \end{bmatrix} \begin{bmatrix} n+j \\ j \end{bmatrix} q^{\binom{j+1}{2}-jn}.$$

Setting $x = a$ in the first statement of Theorem (2.18), we recover Theorem 2 of [12]. The following result is an extension of Theorem 5 of [19].

Theorem 2.19. *Let n be a positive integer. For $m = 1, 2, \dots, n$, we have*

$$\sum_{j=0}^n (-1)^j \begin{bmatrix} n \\ j \end{bmatrix} \begin{bmatrix} n+j \\ j \end{bmatrix} q^{\binom{j}{2}-jn} (\mathcal{H}_{m+j,q}(x) - \mathcal{H}_{j,q}(x)) = (-1)^n q^{\binom{n+1}{2}} \sum_{i=1}^m q^i \frac{(xq^{i-n}; q)_n}{(xq^i; q)_{n+1}}.$$

where

$$\mathcal{H}_{n,q}(x) = \sum_{i=1}^n \frac{q^i}{1-xq^i}$$

Proof. By the first identity of Theorem (2.18), it is easy to check

$$\sum_{j=0}^n (-1)^j \begin{bmatrix} n \\ j \end{bmatrix} \begin{bmatrix} n+j \\ j \end{bmatrix} q^{\binom{j}{2}-jn} (\mathcal{H}_{m+j,q}(x) - \mathcal{H}_{j,q}(x)) = \sum_{i=1}^m q^i \frac{(xq^i)^n (q/xq^i; q)_n}{(xq^i; q)_{n+1}}.$$

By means of

$$(q/xq^i; q)_n = (-1)^n q^{n-in} x^{-n} q^{\binom{n}{2}} (xq^{i-n}; q)_n$$

we deduce the result. \square

DECLARATIONS

Ethical Approval. Not applicable.

Competing interests. No potential conflict of interest was reported by the authors.

Authors' contributions. The authors contributed equally.

Funding. This research received no funding.

Availability of data and materials. Not applicable.

REFERENCES

- [1] S. Ahlgren, K. Ono. A Gaussian hypergeometric series evaluation and Apéry number congruences, *J. Reine Angew. Math.* 518, 187-212 (2000).
- [2] H. Alzer and R. Chapman. On Boole's formula for factorials. *Australas. J. Combinatorics*, 99, 333-336, (2014).
- [3] R. Apéry. Irrationalité de $\zeta(2)$ et $\zeta(3)$ (French), *Astérisque* 61 (1979), 11–13. Luminy Conference on Arithmetic.
- [4] F. Beukers. Another congruence for Apéry numbers, *J. Number Theory* 25, 201-210 (1987).
- [5] W. Chu. A binomial coefficient identity associated with Beukers' conjecture on Apéry numbers, *Electron. J. Comb.* 11 (2004).
- [6] W. Chu and Y. You. Binomial symmetries inspired by Bruckman's problem. *Filomat* 24, 41–46 (2010).
- [7] L. Comtet, *Advanced combinatorics: the art of finite and infinite expansions*. Reidel, Dordrecht 1974.
- [8] G. Gasper, M. Rahman. *Basic hypergeometric series*, Cambridge University Press, Cambridge, 1990.
- [9] I. Gessel. Some congruences for Apéry numbers, *J. Number Theory* 14, 3, 362–368 (1982).
- [10] H. W. Gould. Euler's formula for the n th differences of powers, *Am. Math. Mon.* 85, 450-467 (1978).
- [11] E. A. Karatsuba. On an identity with binomial coefficients. *Mathematical Notes*, 105, 145-147, (2019).
- [12] E. Kiliç, H. Prodinger. Evaluation of sums involving products of Gaussian q -binomial coefficients with applications to Fibonomial sums. *Turj. J. Math.* 41, 707-716 (2017).
- [13] C. Krattenthaler, T. Rivoal, and W. Zudilin. Séries hypergéométriques basiques, q -analogues des valeurs de la fonction zêta et séries d'Eisenstein (French, with English and French summaries), *J. Inst. Math. Jussieu* 5 (2006), 1, 53–79.
- [14] M. Mouçouf, S. Zriaa, A new approach for computing the inverse of confluent Vandermonde matrices via Taylor's expansion, *Linear Multilinear Algebra*. (2021), DOI:10.1080/03081087.2021.1940807.
- [15] C. Phoata. Boole's formula as a consequence of Lagrange's interpolating polynomial theorem. *Integers*, 8, Article A23, (2008).
- [16] J. X. Sharon, J. Zeng. A q -analog of dual sequences with applications. *European Journal of Combinatorics*, 28, 214-227, (2007).
- [17] A. Straub. Supercongruences for polynomial analogs of the Apéry numbers. *Proc. Am. Math. Soc.* 147, 1023–1036 (2019).
- [18] X. Wang, W. Chu. Harmonic number sums and q -analogues. *International Journal of Computer Mathematics: Computer Systems Theory*, 4(1), 48-56 (2019).
- [19] Q. Yan, C. Wei, X. Fan. q -generalizations of Mortenson's identities and further identities. *Ramanujan J.* 35, 131-139, (2014).

[20] D. Y. Zheng. An algebraic identity on q -Apéry numbers. *Discrete. Math*, 311, 23-24 (2011).

SAID ZRIAA AND MOHAMMED MOUÇOUF, UNIVERSITY CHOUAIB DOUKKALI. DEPARTMENT OF MATHEMATICS,
FACULTY OF SCIENCE ELJADIDA, MOROCCO

Email address: `saidzriaa1992@gmail.com`

Email address: `moucou@hotmail.com`