

h^1 boundedness of Localized Operators and Commutators with bmo and lmo

Galia Dafni, Chun Ho Lau *

Abstract

We first consider two types of localizations of singular integral operators of convolution type, and show, under mild decay and smoothness conditions on the auxiliary functions, that their boundedness on the local Hardy space $h^1(\mathbb{R}^n)$ is equivalent. We then study the boundedness on $h^1(\mathbb{R}^n)$ of the commutator $[b, T]$ of an inhomogeneous singular integral operator with b in $\text{bmo}(\mathbb{R}^n)$, the nonhomogeneous space of functions of bounded mean oscillation. We define local analogues of the atomic space $H_b^1(\mathbb{R}^n)$ introduced by Pérez in the case of the homogeneous Hardy space and BMO, including a variation involving atoms with approximate cancellation conditions. For such an atom a , we prove integrability of the associated commutator maximal function and of $[b, T](a)$. For b in $\text{lmo}(\mathbb{R}^n)$, this gives h^1 to L^1 boundedness of $[b, T]$. Finally, under additional approximate cancellation conditions on T , we show boundedness to h^1 .

1 Introduction

The real Hardy spaces $H^p(\mathbb{R}^n)$ are a class of function spaces that has been extensively studied in harmonic analysis for over 50 years. In particular, while the boundedness of singular integral operators on $L^p(\mathbb{R}^n)$ for $1 < p < \infty$ does not extend to $0 < p \leq 1$, under sufficient smoothness and cancellation conditions such operators are bounded on $H^p(\mathbb{R}^n)$, $0 < p \leq 1$. At the other end of the scale, $p = \infty$, the space $\text{BMO}(\mathbb{R}^n)$ of functions of bounded mean oscillation introduced by John and Nirenberg [32] replaces L^∞ and is the dual of H^1 - see the fundamental paper of Fefferman and Stein [20] and more recent exposition in [46].

The focus of our paper is on a nonhomogeneous version of these spaces, introduced by Goldberg [23, 24] under the name “local Hardy spaces”, denoted $h^p(\mathbb{R}^n)$. Membership in $h^p(\mathbb{R}^n)$ does not require the global vanishing moment conditions necessary for $f \in H^p(\mathbb{R}^n)$ and so, unlike the case of $H^p(\mathbb{R}^n)$, important classes of smooth functions such as the Schwartz spaces $\mathcal{S}(\mathbb{R}^n)$ are contained (and are dense) in $h^p(\mathbb{R}^n)$. Moreover, $h^p(\mathbb{R}^n)$ is closed under multiplication by smooth cut-off functions, which is why these are sometimes called “localizable” Hardy spaces, and are more suitable to analysis on different settings such as

*The authors were partially supported by the Natural Sciences and Engineering Research Council (NSERC) of Canada, and the Centre de recherches mathématiques (CRM)

domains, manifolds and spaces of homogeneous type, as well as the study of partial differential equations and pseudo-differential operators.

We restrict ourselves to $h^1(\mathbb{R}^n)$, which satisfies $H^1(\mathbb{R}^n) \subsetneq h^1(\mathbb{R}^n) \subsetneq L^1(\mathbb{R}^n)$, and the nonhomogeneous version of $BMO(\mathbb{R}^n)$, denoted $bmo(\mathbb{R}^n)$ and identified by Goldberg as the dual of $h^1(\mathbb{R}^n)$. In addition to characterizations via maximal functions and atomic decomposition, Goldberg also identified $h^1(\mathbb{R}^n)$ as the space of functions in $L^1(\mathbb{R}^n)$ whose “local Riesz transforms” r_j , $j = 1, \dots, n$, are in $L^1(\mathbb{R}^n)$. The r_j are obtained by localizing the multipliers of the Riesz transforms R_j on the Fourier transform side via multiplication with a Schwartz function vanishing near the origin. We consider more general localizations of Calderón–Zygmund convolution-type singular integral operators, obtained either by multiplying the kernel by an appropriate function (as in the “truncated” Hilbert transform defined in [23]), or localizing on the Fourier transform side. We show, under mild decay and smoothness assumptions on the auxiliary functions, that the boundedness of different localizations of the same kernel on $h^1(\mathbb{R}^n)$ is equivalent (Theorem 3.2), allowing us to characterize h^1 by a broader class of localized Riesz transforms (Corollary 3.1).

We then turn to another important class of operators, the commutators of singular integral operators with multiplication operators: $[b, T]f := bT(f) - T(bf)$. Coifman, Rochberg and Weiss [10] proved that $[b, T]$, where $b \in BMO(\mathbb{R}^n)$ and T is a Calderón–Zygmund operator, is bounded on $L^p(\mathbb{R}^n)$ for all $1 < p < \infty$. They also proved a converse: for the Riesz transforms R_j , if $[b, R_j]$, $j = 1, \dots, n$, are all bounded on $L^p(\mathbb{R}^n)$ for some $1 < p < \infty$, then $b \in BMO(\mathbb{R}^n)$. Subsequently, Uchiyama [49] proved that $[b, T]$ is compact on $L^p(\mathbb{R}^n)$, $1 < p < \infty$, if and only if $b \in CMO(\mathbb{R}^n)$, where $CMO(\mathbb{R}^n)$ is the $BMO(\mathbb{R}^n)$ -closure of $C_c^\infty(\mathbb{R}^n)$. Janson [31] extended the boundedness of the commutator to Orlicz-type spaces. Recently, Hytönen [30] completely characterized $L^p \mapsto L^q$ boundedness.

The above results are all for L^p spaces with $1 < p < \infty$. We are interested in the endpoint case $p = 1$. For a Calderón–Zygmund operator T , we have weak-type $(1, 1)$ boundedness as well as from $H^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$. Does the commutator $[b, T]$ also satisfy this type of boundedness? The answer turns out to be no. Harboure, Segovia and Torrea [26] proved that there is no way to get boundedness of $[b, H]$ from $H^1(\mathbb{R})$ to $L^1(\mathbb{R})$, where H be the Hilbert transform, unless b is a constant function, and this also holds in the case of boundedness of $[b, H]$ from $L_c^\infty(\mathbb{R})$ to $BMO(\mathbb{R})$.

This means we need to seek a smaller subspace of H^1 to get the integrability of the commutator. In addition to proving $L \log L$ to weak- L^1 boundedness of the commutator $[b, T]$, Pérez [39] introduced a special type of H^1 atom that has extra cancellation against b , and showed that the commutator $[b, T](a) \in L^1(\mathbb{R}^n)$ for all such a . He then defined the space $H_b^1(\mathbb{R}^n)$ via atomic decompositions. The boundedness on infinite linear combinations of atoms, however, does not follow from the boundedness on one atom (see [7]), so another characterization of the space was needed.

Ky [34] introduced another space $H_b^1(\mathbb{R}^n)$ through a specially defined commutator maximal function, and proved that it is the largest subspace of $H^1(\mathbb{R}^n)$ such that the commutator $[b, T]$ is bounded from this space to $L^1(\mathbb{R}^n)$. As Pérez’s atomic space is continuously contained in Ky’s space, the boundedness on atoms could be extended to get boundedness on the whole atomic space. The reverse inclusion, namely whether every function in Ky’s maximal $H_b^1(\mathbb{R}^n)$

space has an atomic decomposition into Pérez atoms, was not shown. Ky also showed that under suitable assumptions, $[b, T]$ is bounded from H^1 to $h^1(\mathbb{R}^n)$ and also to $H^1(\mathbb{R}^n)$, and these results were extended to metric measure space of homogeneous type by Fu, D. Yang and S. Yang [21].

Unlike the case for $H^1(\mathbb{R}^n)$, commutators of the form $[b, T]$ with nonconstant b can be bounded on $h^1(\mathbb{R}^n)$. Hung and Ky [28] proved such boundedness for b with logarithmically vanishing mean oscillation and T in a class of pseudo-differential operators.

In this paper we introduce an $h^1(\mathbb{R}^n)$ version of the Pérez atomic space, as well as another atomic space which we call $h_{\text{atom},b}^1(\mathbb{R}^n)$, containing it, defined using new h_b^1 atoms which only satisfy approximate cancellation conditions. We then study the boundedness of $[b, T]$ for $b \in \text{bmo}(\mathbb{R}^n)$ and T an inhomogeneous singular integral operator, and show that such commutators take h_b^1 atoms to $L^1(\mathbb{R}^n)$ (Theorem 5.1). Moreover, imposing nonhomogeneous cancellation conditions on T by considering T^*1 and T^*b , we are able to show boundedness to $h^1(\mathbb{R}^n)$ (Theorem 5.3). However, as in the case of the Pérez atomic space H_b^1 , it is not possible to extend this boundedness to all of $h_{\text{atom},b}^1(\mathbb{R}^n)$ without a different characterization of the space. While we do not have a maximal characterization, we do show that a local analogue of Ky's commutator maximal function takes h_b^1 atoms to $L^1(\mathbb{R}^n)$.

Finally, restricting b to $\text{lmo}(\mathbb{R}^n)$, the functions of logarithmically vanishing mean oscillation in $\text{bmo}(\mathbb{R}^n)$, we are able to identify the space $h_{\text{atom},b}^1(\mathbb{R}^n)$ with $h^1(\mathbb{R}^n)$ (Theorem 4.6) and thus conclude the boundedness of the commutator on the whole space, generalizing the results of Hung and Ky [28].

Section 2 gives some background on h^1 , with particular emphasis on atomic decompositions and approximate cancellation conditions, as well as on bmo and lmo . Section 3 deals with the various localizations of Calderón–Zygmund operators of convolution type and their boundedness on h^1 , and ends with the definition of inhomogeneous singular integral operators. Section 4 introduces the commutator maximal functions and atomic spaces in the nonhomogeneous case, and proves relations between them. Finally, Section 5 contains the results on the boundedness of the commutators on the atoms.

2 Preliminaries

The Lebesgue measure of a set $A \subset \mathbb{R}^n$ is denoted by $|A|$, and the average of an integrable function f on a set E of positive finite measure by

$$f_E := \int_E f := \frac{1}{|E|} \int_E f(x) dx.$$

Balls will be denoted by $B(x_0, r)$ and the radius of a ball B by $r(B)$. The dilated ball $B(x_0, \delta r)$, $\delta > 0$, will be denoted by δB .

The constant C used in inequalities will may vary from line to line. We write $\alpha \lesssim \beta$ if there exists a constant C (which may depend on other factors but not on α and β) such that $\alpha \leq C\beta$, and $\alpha \approx \beta$ if $\alpha \lesssim \beta$ and $\beta \lesssim \alpha$.

We denote the Schwartz space by $\mathcal{S}(\mathbb{R}^n)$ and the seminorms defining its topology by

$$\|\varphi\|_{\alpha,\beta} := \sup_{x \in \mathbb{R}^n} |x^\alpha \partial_x^\beta \varphi(x)|, \quad \alpha, \beta \in \mathbb{Z}_+^n.$$

Its dual, $\mathcal{S}'(\mathbb{R}^n)$, is the space of tempered distributions.

2.1 Local Hardy spaces

Goldberg [23, 24] introduced nonhomogeneous versions of the real Hardy spaces $H^p(\mathbb{R}^n)$, called “local Hardy spaces” and denoted by $h^p(\mathbb{R}^n)$, and characterized them by means of maximal functions and atomic decomposition.

Definition 2.1 ([23]). For $0 < p \leq 1$, a (normalized) Goldberg h^p atom a is a function supported in a cube Q with

- (i) $\|a\|_{L^\infty} \leq |Q|^{-1}$;
- (ii) if $|Q| < 1$, $\int a(x)x^\alpha dx = 0$ for $|\alpha| \leq [n(1/p - 1)]$.

An atomic decomposition for h^p where the atoms need only have approximate moment conditions was introduced in [12] and later refined in [17, 18] for the case $p = 1$ in the more general setting of spaces of homogeneous type (see also [14, 15, 33] for $p < 1$).

Definition 2.2 ([18]). Fix $R > 0$, $1 < q \leq \infty$. We say a is an R -approximate $(1, q)$ atom if

- $\text{supp}(a) \subset B(x_0, r)$ for some $x_0 \in \mathbb{R}^n$ and $r > 0$;
- $\|a\|_{L^q} \leq |B(x_0, r)|^{\frac{1}{q}-1}$;
- $|\int a| \leq [\log(1 + \frac{R}{r})]^{-1}$.

Any of the following equivalent characterizations can be taken as the definition of $h^1(\mathbb{R}^n)$.

Proposition 2.1. *Let $f \in \mathcal{S}'(\mathbb{R}^n)$. Then the following are equivalent:*

1. For a fixed $\psi \in \mathcal{S}(\mathbb{R}^n)$ with $\int \psi \neq 0$,

$$\|f\|_{h^1, \psi} := \|M_\psi(f)\|_{L^1(\mathbb{R}^n)} < \infty,$$

where

$$M_\psi(f)(x) = \sup_{0 < t < T} |f * \psi_t(x)|, \quad \psi_t(y) = t^{-n} \psi(t^{-1}y).$$

2. For a fixed $T > 0$,

$$\|f\|_{h^1, T} := \|\mathcal{M}(f)\|_{L^1(\mathbb{R}^n)} < \infty,$$

where

$$\mathcal{M}(f)(x) := \sup_{\phi} |\langle f, \phi \rangle|,$$

and the supremum is taken over all $\phi \in C^1(\mathbb{R}^n)$ with $\text{supp}(\phi) \subset B(x, t)$, $0 < t < T$, $\|\phi\|_{L^\infty} \leq |B(x, t)|^{-1}$ and $\|\nabla \phi\|_{L^\infty} \leq [t|B(x, t)|]^{-1}$.

3. For N sufficient large

$$\|f\|_{h^1, \max} := \|\mathfrak{M}(f)\|_{L^1(\mathbb{R}^n)} < \infty,$$

where

$$\mathfrak{M}f(x) := \sup\{|f * \varphi_t(x)| : 0 < t < 1, \varphi \in \mathcal{F}_N\}$$

and

$$\mathcal{F}_N = \{\varphi \in \mathcal{S}(\mathbb{R}^n) : \|\varphi\|_{\alpha, \beta} \leq 1 \forall |\alpha|, |\beta| \leq N\}.$$

4.

$$\|f\|_{h^1, \text{atom}} := \inf \sum |\lambda_j| < \infty,$$

where the infimum is taken over all decompositions $f = \sum \lambda_j a_j$ in $\mathcal{S}'(\mathbb{R}^n)$ with $\{\lambda_j\} \in \ell^1$ and where the a_j are Goldberg h^1 atoms as in Definition 2.1.

5. For a fixed $R > 0$, $1 < q \leq \infty$,

$$\|f\|_{h^1, R} := \inf \sum |\lambda_j| < \infty,$$

where the infimum is taken over all decompositions $f = \sum \lambda_j a_j$ in $\mathcal{S}'(\mathbb{R}^n)$ with $\{\lambda_j\} \in \ell^1$ and where the a_j are R -approximate $(1, q)$ atoms as in Definition 2.2.

In case these conditions hold, we have

$$\|f\|_{h^1, \psi} \approx \|f\|_{h^1, T} \approx \|f\|_{h^1, \max} \approx \|f\|_{h^1, \text{atom}} \approx \|f\|_{h^1, R},$$

where the constants depend on the choices of T , R and N .

The equivalence of conditions (1), (3) and (4) of Proposition 2.1 is contained in [23, Theorem 1 and Lemma 5]. The equivalence of conditions (2), (4) and (5) is shown in [18, Section 7] on a doubling metric-measure space, with C^1 replaced by Lipschitz, for arbitrary T , R and q (see also [14, Lemma 1] for a proof that (4) implies (2) in the more general case of all $p \leq 1$). The maximal functions \mathcal{M} and \mathfrak{M} are often given the name *grand maximal function*. It is also possible to define nontangential versions of these (see Definition 4.1 below for the case of H^1) and get an equivalent characterization of h^1 (resp. H^1) - this is part of Goldberg's theorem (see [46, Section III.1] for the homogeneous Hardy spaces). Further characterizations of local Hardy spaces on spaces of homogeneous type were given in [27].

In what follows, we will use mostly condition (2) with $T = 1$, and abbreviate $\|f\|_{h^1, 1}$ to $\|f\|_{h^1}$, and condition (5) with $R = 1$ and $q = 2$, in which case we will call the 1 approximate $(1, 2)$ atoms simply h^1 atoms.

We will also need the notion of a molecule, which is a generalization of an atom not requiring compact support. Molecules are useful in showing boundedness of operators on h^p since, typically, singular integral operators take atoms to molecules. If we know that molecules are in h^p with bounded norm, then we have shown the boundedness of the operator on atoms.

Definition 2.3 ([14]). Let $1 < s < \infty$, $\lambda > n(s-1)$. A measurable function M on \mathbb{R}^n is called a (s, λ) molecule (for $h^1(\mathbb{R}^n)$) if there exists a ball $B = B(x_0, r)$ such that

$$\text{M1. } \|M\|_{L^s(B)} \leq r^{n(\frac{1}{s}-1)}$$

$$\text{M2. } \|M|\cdot -x_0|^{\frac{\Delta}{s}}\|_{L^s(B^c)} \leq r^{\frac{\Delta}{s}+n(\frac{1}{s}-1)}$$

$$\text{M3. } \left| \int M \right| \leq [\ln(1+r^{-1})]^{-1}.$$

Proposition 2.2 ([14]). *There exists a constant $C_{n,s,\lambda}$ such that $\|M\|_{h^1} \leq C_{n,s,\lambda}$ every (s, λ) molecule M .*

Conversely, for any $f \in h^1(\mathbb{R}^n)$, there exist a sequence $\{\lambda_j\} \in \ell^1$ and a sequence of h^1 molecules $\{M_j\}$ such that

$$f = \sum_{j=1}^{\infty} \lambda_j M_j \quad \text{in } h^1(\mathbb{R}^n).$$

Moreover,

$$\|f\|_{h^1} \approx \inf \sum |\lambda_j| < \infty,$$

where the infimum is taken over all such decompositions.

The following proposition (case $p = 1$ of [15, Proposition 1]) shows that the cancellation condition on atoms in Definition 2.2 is not only sufficient but also necessary.

Proposition 2.3 ([15]). *Suppose $g \in h^1(\mathbb{R}^n)$ with $\text{supp}(g) \subset B(x_0, r)$. Then*

$$\left| \int g \right| \lesssim \frac{\|g\|_{h^1}}{\log(1+r^{-1})}.$$

2.2 Nonhomogeneous BMO

Goldberg [24, Corollary1] identifies the dual of $h^1(\mathbb{R}^n)$ with the space $\text{bmo}(\mathbb{R}^n)$, a nonhomogeneous version of the John-Nirenberg space $\text{BMO}(\mathbb{R}^n)$ of functions of bounded mean oscillation. See Bourdaud [6] for an exposition of the properties of $\text{bmo}(\mathbb{R}^n)$. In the following definition, we use the notation introduced in [18], where it is shown that the definition is independent of the choice of scale used to distinguish large and small balls. Note that unlike $\text{BMO}(\mathbb{R}^n)$, we do not need to take the space modulo constants in order to get a Banach space.

Definition 2.4. Let $b \in L^1_{\text{loc}}(\mathbb{R}^n)$. We say $b \in \text{bmo}(\mathbb{R}^n)$ if

$$\|b\|_{\text{bmo}} := \sup_B \int_B |b(x) - c_B| dx < \infty,$$

where the supremum is taken over all balls, and for each ball B we set

$$c_B := \begin{cases} b_B := \int_B b, & \text{if } r(B) < 1, \\ 0, & \text{if } r(B) \geq 1. \end{cases} \quad (2.1)$$

Since there will only be one bmo function under discussion at any point in time, we will continue to use the notation c_B defined in (2.1) in the rest of the paper. We can then state a corresponding version of the John-Nirenberg

inequality for $b \in \text{bmo}(\mathbb{R}^n)$ (see [18, Theorem 3.1]): there exist $C, c > 0$ such that for any ball B ,

$$|\{x \in B : |b(x) - c_B| > \lambda\}| \leq C|B|e^{-\lambda c/\|b\|_{\text{bmo}}}$$

for all $\lambda > 0$. As consequence, one gets that

$$\|b\|_{\text{bmo}} \approx \|b\|_{\text{bmo},p} := \left(\sup_B \int_B |b(x) - c_B|^p dx \right)^{1/p} \quad (2.2)$$

for $1 \leq p < \infty$, with constants depending on p .

This gives us the following useful inequality (see the proof of [10, Lemma 2.4] for the case $\delta = 1$ and $b \in \text{BMO}(\mathbb{R}^n)$).

Lemma 2.1. *Let $b \in \text{bmo}(\mathbb{R}^n)$. Then for any $\delta > 0$, $p \geq 1$, there exists $C_{n,p,\delta} > 0$, independent of b , x_0 and r , such that*

$$r^\delta \int_{|x-x_0|>r} \frac{|b(x) - c_{B(x_0,r)}|^p}{|x-x_0|^{n+\delta}} \leq C_{n,p,\delta} \|b\|_{\text{bmo}}^p. \quad (2.3)$$

Proof. By translation invariance of $\text{bmo}(\mathbb{R}^n)$, we may assume $x_0 = 0$. Denote $B(0, 2^k r)$ by B_k , $k = 0, 1, 2, \dots$. Then using (2.2) and the fact that for $2^j r < 1$

$$|c_{B_j} - c_{B_{j-1}}| = |b_{B_j} - b_{B_{j-1}}| \leq 2^j \|b\|_{\text{bmo}} \quad (2.4)$$

and $c_B \lesssim \frac{1}{\log(1+r^{-1})}$ (see [18, Lemma 6.1]), we have

$$\begin{aligned} r^\delta \int_{\mathbb{R}^n \setminus B_0} \frac{|b(x) - c_{B_0}|^p}{|x|^{n+\delta}} &\lesssim \sum_{k=0}^{\infty} 2^{-k\delta} |B_k|^{-1} \int_{B_{k+1} \setminus B_k} |b(x) - c_{B_{k+1}}|^p + |c_{B_{k+1}} - c_{B_0}|^p dx \\ &\lesssim \sum_{k=0}^{\infty} 2^{-k\delta} \|b\|_{\text{bmo}}^p (1 + (\min\{k+1, \log_+ r^{-1}\})^p) dx \\ &\lesssim \|b\|_{\text{bmo}}^p. \end{aligned}$$

□

While bmo is the dual of the local Hardy space h^1 , it does not make sense to call it “local BMO” as it is actually strictly smaller. It is more appropriate to give that name to the space consisting of locally integrable functions b for which

$$\|b\|_{\text{BMO}_{\text{loc}}(\mathbb{R}^n)} := \sup_{r(B) < R} \int_B |b(x) - b_B| dx < \infty,$$

where R is some fixed constant (see [48]). This space is strictly larger than $\text{BMO}(\mathbb{R}^n)$ since it contains all Lipschitz functions, for example, and is independent of the choice of $R < \infty$. Taking those elements of BMO_{loc} for which $\sup_{r(B) \geq R} \int |b| < \infty$ gives $\text{bmo}(\mathbb{R}^n)$ (in [6] it is shown that it suffices to take $\sup_{r(B)=R} \int |b| < \infty$). In particular $\text{BMO}_{\text{loc}}(\mathbb{R}^n) \cap L^p(\mathbb{R}^n) \subset \text{bmo}(\mathbb{R}^n)$ for $1 \leq p < \infty$.

Vanishing mean oscillation refers to the property that

$$\lim_{R \rightarrow 0} \sup_{r(B) < R} \int_B |b(x) - b_B| dx = 0. \quad (2.5)$$

For $b \in \text{BMO}(\mathbb{R}^n)$, this holds if and only if $b \in \text{VMO}(\mathbb{R}^n)$, the closure of the uniformly continuous functions in $\text{BMO}(\mathbb{R}^n)$, as shown by Sarason [40] for $n = 1$. Analogously, we can denote by $\text{vmo}(\mathbb{R}^n)$ the space of functions in $\text{bmo}(\mathbb{R}^n)$ which have vanishing mean oscillation, or alternatively the closure of the bounded uniformly continuous functions in $\text{bmo}(\mathbb{R}^n)$. Finally, we use $\text{cmo}(\mathbb{R}^n)$ to denote the closure of the continuous (or smooth) functions with compact support in $\text{bmo}(\mathbb{R}^n)$. More about these spaces can be found in [6] and [13] (where the notation vmo is used for cmo). It was shown there that $h^1(\mathbb{R}^n)$ can be identified with the dual of $\text{cmo}(\mathbb{R}^n)$ and that $b \in \text{cmo}(\mathbb{R}^n)$ if and only if (2.5) holds and in addition

$$\lim_{R \rightarrow \infty} \sup_{\substack{|B| \geq 1 \\ B \subset (B(0,R))^c}} \int_B |b| = 0. \quad (2.6)$$

One can specify the rate at which the oscillation vanishes as the radius goes to zero in (2.5) (see [40, Chapter 5], [43]). A special class are those functions of *logarithmic mean oscillation*, usually denoted LMO (not to be confused with *bounded lower oscillation* denoted by BLO). Bounded functions in this class have been identified as the pointwise multipliers of H^1 and BMO on the circle [44], on the sphere [35] and on spaces of homogeneous type [9]. The analogous result in the case of bmo was shown in [5]. These spaces have also proved useful in the study of PDE. In [47] a logarithmic mean oscillation condition is imposed on the coefficients of a parabolic equation, while in [3, 4] a range of such conditions is imposed on the initial vorticity in the Euler and Navier-Stokes equations. Note that the latter articles use slightly different notation (for example bmo there refers to “local BMO” rather than Goldberg’s nonhomogeneous space) and an L^2 oscillation, which is equivalent to the L^1 one by the John-Nirenberg inequality for these spaces (see [43]). In [28], a space called LMO_∞ is introduced which consists of functions in BMO_{loc} which have logarithmic vanishing mean oscillation on small balls while the mean oscillation on large balls can grow as a power of the radius. This space is contained in $\text{LMO}_{\text{loc}}(\mathbb{R}^n)$, defined as follows.

Definition 2.5. Let $b \in L^1_{\text{loc}}(\mathbb{R}^n)$. We say $b \in \text{LMO}_{\text{loc}}(\mathbb{R}^n)$ if

$$\|b\|_{\text{LMO}_{\text{loc}}(\mathbb{R}^n)} := \sup_{r(B) < 1} \frac{[\log(1 + r(B)^{-1})]}{|B|} \int_B |b(x) - b_B| dx < \infty.$$

We also define $\text{lmo}(\mathbb{R}^n) := \text{LMO}_{\text{loc}}(\mathbb{R}^n) \cap \text{bmo}(\mathbb{R}^n)$ and

$$\|b\|_{\text{lmo}(\mathbb{R}^n)} := \|b\|_{\text{LMO}_{\text{loc}}(\mathbb{R}^n)} + \sup_{r(B) \geq 1} |b|_B < \infty.$$

Since $[\log(1 + r(B)^{-1})]^{-1} \rightarrow 0$ as $r \rightarrow 0$, we have that functions in $\text{LMO}_{\text{loc}}(\mathbb{R}^n)$ satisfy the vanishing mean oscillation condition (2.5) and therefore $\text{lmo}(\mathbb{R}^n) \subset \text{vmo}(\mathbb{R}^n)$. If we take $b \in \text{LMO}_{\text{loc}}(\mathbb{R}^n) \cap L^p(\mathbb{R}^n)$, $1 \leq p < \infty$, then b will also satisfy condition (2.6) since if $|B| \geq 1$ and $B \subset \mathbb{R}^n \setminus B(0, R) \subset \mathbb{R}^n$, then $\int_B |b| \leq \|b\|_{L^p(B(0,R)^c)} \rightarrow 0$ as $R \rightarrow \infty$. Thus $\text{LMO}_{\text{loc}}(\mathbb{R}^n) \cap L^p(\mathbb{R}^n) \subset \text{cmo}(\mathbb{R}^n)$.

3 Localized convolution kernels

In addition to the characterizations via maximal functions and atomic decomposition, as in Proposition 2.1, Goldberg also characterized $h^1(\mathbb{R}^n)$ by singular integral operators in an analogous way to the characterization of $H^1(\mathbb{R}^n)$ via the Riesz transforms R_j , $j = 1, \dots, n$, given by their multipliers as $\widehat{R_j f}(\xi) = i \frac{\xi_j}{|\xi|} \widehat{f}(\xi)$.

Theorem 3.1. [24, Theorem 2] *Fix a function $\varphi \in \mathcal{S}(\mathbb{R}^n)$ with $\varphi \equiv 1$ in a neighborhood of the origin. Then a distribution f belongs to $h^1(\mathbb{R}^n)$ if and only if $f \in L^1(\mathbb{R}^n)$ and $r_j f \in L^1(\mathbb{R}^n)$, $j = 1, \dots, n$, where r_j is defined by*

$$\widehat{r_j f}(\xi) = (1 - \varphi(\xi)) i \frac{\xi_j}{|\xi|} \widehat{f}(\xi).$$

See [38] for a generalization of this result to the case $0 < p \leq 1$, using φ of compact support.

For $n = 1$, in [23], Goldberg also claims a characterization of $h^1(\mathbb{R})$ via a localized version of the Hilbert transform. This can also be described as a “truncated” Hilbert transform (not to be confused with the usual truncation used to define the principal value integral). We set

$$\mathcal{H}_{\text{loc}}(f)(x) := p.v. \int_{\mathbb{R}} \frac{\eta(x-y)f(y)}{x-y} dy, \quad (3.1)$$

where $\eta \equiv 1$ on $(-1, 1)$ and is supported in $[-2, 2]$. In [16] it was shown that for smooth η , the local Hilbert transform \mathcal{H}_{loc} characterizes $h^1(\mathbb{R})$.

Note that the two localizations are different: \mathcal{H}_{loc} is obtained by localizing the kernel of the Hilbert transform near 0, while r_j is obtained by localizing the Fourier transform of R_j away from 0. This may cause some confusion. For example, in order to characterize weighted local Hardy spaces, higher-dimensional analogues of the local Hilbert transforms, namely “truncated” versions of the Riesz transforms where the kernel is multiplied by a smooth function of compact support, were defined in [48, Section 8] using the notation R_j . However, the result referred to as [48, Theorem A], which is [8, Corollary to Theorem 3.1], actually uses the same r_j as in Theorem 3.1, localized on the Fourier transform side.

In this section we consider these two types of localization for a more general convolution kernel, with minimal smoothness and decay conditions on the auxiliary functions used for the localization, and prove that the boundedness of the associated singular integral operators on $h^1(\mathbb{R}^n)$ does not depend on which type of localization we choose.

3.1 Calderón–Zygmund singular integral operators of convolution type

We assume that our kernel is δ -kernel, namely K is a measurable function on $\mathbb{R}^n \setminus \{0\}$ and there exist $\delta \in (0, 1]$ and $C > 0$ such that

$$|K(x)| \leq \frac{C}{|x|^\delta} \quad (3.2)$$

and

$$|K(x-y) - K(x)| \leq C \frac{|y|^\delta}{|x|^{n+\delta}} \quad \text{for } 2|y| \leq |x|. \quad (3.3)$$

We also assume that a cancellation condition holds in the following form:

$$\exists A < \infty \quad \sup_{0 < r < R} \left| \int_{r < |x| < R} K(x) dx \right| \leq A. \quad (3.4)$$

We then define the associated convolution operator $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ by the principal value integral

$$Tf(x) = \lim_{\varepsilon \rightarrow 0} \int_{|x-y| \geq \varepsilon} K(x-y) f(y) dy.$$

This extends to a bounded operator on $L^2(\mathbb{R}^n)$ (see [46, Section VII.3]). Furthermore, T is bounded on $L^p(\mathbb{R}^n)$, $1 < p < \infty$, and from $L^1(\mathbb{R}^n)$ to $L^{1,\infty}(\mathbb{R}^n)$. See [45, Chapter 2] for the classical theory of Calderón–Zygmund operators of convolution type and [25, Chapter 4] for more general kernels. A general Calderón–Zygmund operator can be written as a sum of an operator of this form plus an operator given by multiplication by a bounded function (see [46, Section I.7.4], [25, Proposition 4.1.11]), but since multiplication by a bounded function is typically not a bounded operator on h^1 , here we only consider the singular integral part.

We define two types of operators via localizations of the kernel K . On the one hand, we look at operators of the form

$$T^\psi(f) := T(f - \psi * f)$$

for suitable functions ψ . These are modelled on Goldberg’s localized Riesz transforms in Theorem 3.1, with $\psi = \check{\varphi}$.

On the other, based on the definition of the local Hilbert transform (3.1), we look at operators T_η , associated to the kernel $K\eta$, for a class of functions η .

3.2 The Operator T_η

We now show that under certain weak smoothness and decay conditions on η , the kernel $K\eta$ satisfies the same conditions as K and so we can associate to it an operator T_η in the same way that we associate T to K , and this operator enjoys the same boundedness properties as T .

Lemma 3.1. *Suppose that η is a bounded function and there exists a constant C for which*

$$|\eta(x-y) - \eta(x)| \leq C \frac{|y|^\delta}{|x|^\delta} \quad \text{for } 0 < 2|y| \leq |x|. \quad (3.5)$$

Then $K\eta$ satisfies conditions (3.2) and (3.3).

Proof. We have that $|K\eta| \leq \|\eta\|_{L^\infty} |x|^{-n}$, and for $0 < 2|y| \leq |x|$, since this

implies $|x - y| \geq |x|/2$, we get

$$\begin{aligned}
& |K(x - y)\eta(x - y) - K(x)\eta(x)| \\
& \leq |K(x - y)| |\eta(x - y) - \eta(x)| + |K(x - y) - K(x)| |\eta(x)| \\
& \lesssim \frac{1}{|x - y|^n} \frac{|y|^\delta}{|x|^\delta} + \frac{|y|^\delta}{|x|^{n+\delta}} \\
& \lesssim \frac{|y|^\delta}{|x|^{n+\delta}}.
\end{aligned}$$

□

The condition on η is a local δ -Lipschitz condition such that the Lipschitz constant in $B(x, |x|/2)$ decays like $|x|^{-\delta}$. In particular, if η has compact support, then we just require $\eta \in \text{Lip}_\delta$.

Next we show that $K\eta$ satisfies (3.4). For this we need further assumptions on η guaranteeing that η has a certain decay at infinity and $\eta - 1$ vanishes sufficiently fast at the origin. Note that we do not require that $\eta \equiv 1$ in a neighborhood of the origin. For example, if $\eta \in \text{Lip}_\delta$ then it suffices that $\eta(0) = 1$.

Lemma 3.2. *If η satisfies,*

$$\int_{0 < |x| < 1} \frac{|\eta(x) - 1|}{|x|^n} dx + \int_{|x| \geq 1} \frac{|\eta(x)|}{|x|^n} dx < \infty, \quad (3.6)$$

then (3.4) holds for $K\eta$.

Proof. For $0 < r < R$, by the assumptions on K and η ,

$$\begin{aligned}
& \left| \int_{r < |y| < R} K(y)\eta(y) dy \right| \\
& \leq \left| \int_{r < |y| < 1} K(y)[\eta(y) - 1] dy \right| + A + \left| \int_{1 \leq |y| < R} K(y)\eta(y) dy \right| \\
& \lesssim \int_{0 < |y| < 1} \frac{|\eta(y) - 1|}{|y|^n} dy + \int_{|y| \geq 1} \frac{|\eta(y)|}{|y|^n} dy + 1 \leq A',
\end{aligned}$$

where the constant A' is independent of r and R . □

If η satisfies the hypotheses of both lemmas, we get the boundedness of T_η on $L^2(\mathbb{R}^n)$. Once we have L^2 -boundedness, we can show that T_η is bounded on $L^p(\mathbb{R}^n)$ for $1 < p < \infty$ and from $L^1(\mathbb{R}^n)$ to weak- $L^1(\mathbb{R}^n)$.

3.3 Equivalent boundedness of T_η and T^ψ

For nice choices of ψ , one can study the boundedness of the localized operator T^ψ as a pseudo-differential operator. Alternatively, one can show that the kernel of T^ψ is sufficiently good. In particular, assuming that $\psi \in L^1$, T^ψ inherits from T the boundedness on L^p for $1 < p < \infty$ and the weak $(1, 1)$ boundedness. We are interested in the boundedness on h^1 .

Instead of working with T^ψ directly, we give sufficient conditions on ψ to be able to obtain the boundedness of T^ψ on h^1 from the boundedness of T_η . As

a consequence of the following equivalence, it suffices to prove the boundedness for one good η (e.g. a Lip_δ function of compact support) in order to obtain it for the whole class of η and ψ .

Theorem 3.2. *Suppose η satisfies the hypotheses of Lemmas 3.1 and 3.2, and ψ satisfies*

$$\psi \in L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n), \quad \int \psi = 1,$$

$$\int_{|x| \geq 1} \left[\frac{1}{|x|^n} \int_{|y| \geq |x|/2} |\psi(y)| dy + \frac{1}{|x|^{n+\delta}} \int_{|y| \leq |x|/2} |y|^\delta |\psi(y)| dy \right] dx < \infty \quad (3.7)$$

and

$$\int_{|x| \geq 1} \left[\int_{|y| \leq |x|/2} \frac{|\psi(x-y) - \psi(x)|}{|y|^n} dy \right] dx < \infty. \quad (3.8)$$

Then the operator T_η is bounded on $h^1(\mathbb{R}^n)$ if and only if T^ψ is bounded on $h^1(\mathbb{R}^n)$.

Note that condition (3.7) will hold, for example, if the function $|x|^\eta \psi(x)$ is integrable for some $\eta > 0$, while (3.8) will hold if $\psi \in \text{Lip}_\alpha$ for $0 < \alpha \leq 1$ and $|x|^\alpha \|\psi\|_{\text{Lip}_\alpha(B(x, |x|/2))}$ is integrable.

Proof. We will show that the operator $T_E := T_\eta - T^\psi$ is bounded on $L^1(\mathbb{R}^n)$, and from that deduce that T_E is bounded on $h^1(\mathbb{R}^n)$.

For $0 < \varepsilon < 1/2$, let $K_\varepsilon(x) = \chi_{|x| > \varepsilon} K(x)$ be the usual truncation of the kernel. Applying the same to $K\eta$, we consider $(K\eta)_\varepsilon - K_\varepsilon + K_\varepsilon * \psi$. Set

$$K_*(x) = \sup_{\varepsilon > 0} |(K\eta)_\varepsilon(x) - K_\varepsilon(x) + K_\varepsilon * \psi(x)|.$$

We claim that $K_* \in L^1(\mathbb{R}^n)$.

For the local estimate, we write

$$K_*(x) \leq |K(x)(\eta(x) - 1)| + T_*(\psi)(x),$$

where $T_*(\psi)(x) = \sup_{\varepsilon > 0} |K_\varepsilon * \psi(x)|$. Recall that the maximal operator T_* has the same boundedness as T (see [46, Section I.7], [25, Section 4.2.2]). Thus we have

$$\int_{|x| < 1} K_*(x) dx \lesssim \int_{|x| < 1} \frac{|\eta(x) - 1|}{|x|^n} dx + \|\psi\|_{L^2} < \infty$$

For $|x| \geq 1$, we can write

$$K_*(x) \leq |K\eta(x)| + \sup_{\varepsilon > 0} |K(x) - K_\varepsilon * \psi(x)|.$$

The first term is integrable for $|x| \geq 1$ by condition (3.6) on η , so it remains to bound the integral of the second term. We do this by fixing ε and obtaining a pointwise estimate on $|K(x) - K_\varepsilon * \psi(x)|$ in terms of ψ .

Using the hypotheses on ψ , we can write, for $|x| \geq 1$ and $\varepsilon < 1/2$,

$$\begin{aligned} \left| K(x) - \int_{|x-y|>\varepsilon} K(x)\psi(y)dy \right| &\leq |K(x)| \int_{|x-y|\leq\varepsilon} |\psi(y)|dy \\ &\lesssim |x|^{-n} \int_{|y|\geq|x|/2} |\psi(y)|dy. \end{aligned} \quad (3.9)$$

Thus it remains to consider

$$\begin{aligned} &\int_{|x-y|>\varepsilon} [K(x) - K(x-y)]\psi(y)dy \\ &= \int_{\left\{ \begin{array}{l} |y|\leq|x|/2 \\ |x-y|\geq|x|/2 \end{array} \right\}} + \int_{\left\{ \begin{array}{l} |y|\geq|x|/2 \\ |x-y|\geq|x|/2 \end{array} \right\}} + \int_{\varepsilon<|x-y|\leq|x|/2} [K(x) - K(x-y)]\psi(y)dy \\ &=: I_1 + I_2 + I_3. \end{aligned}$$

To estimate I_1 , note that $B(0, |x|/2) \subset B(x, |x|/2)^c$ so we can use the smoothness condition (3.3) on K to write

$$\begin{aligned} |I_1| &\leq \int_{|y|\leq|x|/2} |K(x) - K(x-y)| |\psi(y)| dy \\ &\lesssim \frac{1}{|x|^{n+\delta}} \int_{|y|\leq|x|/2} |y|^\delta |\psi(y)| dy. \end{aligned}$$

For I_2 , we use the decay bound (3.2) on K to get, as in (3.9),

$$\begin{aligned} |I_2| &\leq \int_{\left\{ \begin{array}{l} |y|\geq|x|/2 \\ |x-y|\geq|x|/2 \end{array} \right\}} (|K(x)| |\psi(y)| + |K(x-y)| |\psi(y)|) dy \\ &\lesssim \frac{1}{|x|^n} \int_{|y|\geq|x|/2} |\psi(y)| dy. \end{aligned}$$

Finally, for I_3 , we again use the decay bound (3.2) as well as the cancellation condition (3.4) on K to obtain

$$\begin{aligned} |I_3| &\leq \left| \int_{\varepsilon<|x-y|\leq|x|/2} [K(x)\psi(y) - K(x-y)(\psi(y) - \psi(x))] dy \right| \\ &\quad + \left| \int_{\varepsilon<|x-y|\leq|x|/2} K(x-y)\psi(x) dy \right| \\ &\lesssim \int_{|y|\geq|x|/2} \frac{|\psi(y)|}{|x|^n} dy + \int_{|x-y|\leq|x|/2} \frac{|\psi(y) - \psi(x)|}{|x-y|^n} dy + |\psi(x)| \\ &\lesssim \frac{1}{|x|^n} \int_{|y|\geq|x|/2} |\psi(y)| dy + \int_{|y'|\leq|x|/2} \frac{|\psi(x-y') - \psi(x)|}{|y'|^n} dy + |\psi(x)|. \end{aligned}$$

Since all these estimates are independent of ε , we can combine (3.9), the bounds on I_1 , I_2 and I_3 , and the integrability conditions (3.7) and (3.8) on ψ to get that

$$\int_{|x|\geq 1} \sup_{\varepsilon>0} |K(x) - K_\varepsilon * \psi(x)| < \infty$$

and as a result, $K_* \in L^1(\mathbb{R}^n)$. Let

$$K_E(x) = \lim_{\varepsilon \rightarrow 0} ((K\eta)_\varepsilon - K_\varepsilon + K_\varepsilon * \psi)(x) = K\eta(x) - K(x) + \lim_{\varepsilon \rightarrow 0} K_\varepsilon * \psi(x).$$

This limit exists for almost every $x \in \mathbb{R}^n$ by the properties of T_* (see [45, p. 45]). By the Dominated Convergence Theorem, $K_E \in L^1(\mathbb{R}^n)$ and

$$\lim_{\varepsilon \rightarrow 0} ((K\eta)_\varepsilon - K_\varepsilon + K_\varepsilon * \psi) * f = K_E * f \quad \text{for } f \in L^1(\mathbb{R}^n).$$

Let $f \in h^1(\mathbb{R}^n)$. Then $T_E f \in L^1(\mathbb{R}^n)$ with

$$\|T_E f\|_{L^1(\mathbb{R}^n)} \leq \|K_E\|_{L^1(\mathbb{R}^n)} \|f\|_{L^1(\mathbb{R}^n)}.$$

Moreover, the localized Riesz transforms r_j , $j = 1, \dots, n$ of Goldberg (see Theorem 3.1) commute with convolution with the L^1 function K_E , so

$$r_j(T_E(f)) = T_E(r_j(f)).$$

Since $r_j(f) \in L^1(\mathbb{R}^n)$, hence $T_E(r_j(f)) \in L^1(\mathbb{R}^n)$, we have

$$\begin{aligned} \|T_E f\|_{h^1(\mathbb{R}^n)} &\lesssim \|T_E f\|_{L^1(\mathbb{R}^n)} + \sum_{j=1}^n \|r_j(T_E(f))\|_{L^1(\mathbb{R}^n)} \\ &\leq \|K_E\|_{L^1(\mathbb{R}^n)} (\|f\|_{L^1(\mathbb{R}^n)} + \sum_{j=1}^n \|r_j(f)\|_{L^1(\mathbb{R}^n)}) \\ &\lesssim \|f\|_{h^1(\mathbb{R}^n)}. \end{aligned}$$

□

Starting with the kernels of the Riesz transforms R_j , $j = 1, \dots, n$, which satisfy conditions (3.2) and (3.3) with $\delta = 1$ and (3.4) with $A = 0$, and noting that the function $\psi = \tilde{\varphi} \in \mathcal{S}(\mathbb{R}^n)$ for φ as in Theorem 3.1 satisfies the hypotheses of Theorem 3.2, we can use both theorems to get the following.

Corollary 3.1. *Suppose that η and ψ satisfy assumptions in Theorem 3.2. For $j = 1, \dots, n$, define, for $f \in \mathcal{S}(\mathbb{R}^n)$,*

$$\mathcal{R}_{j,\eta} f(x) := c_n \lim_{\varepsilon \rightarrow 0} \int_{|x-y| \geq \varepsilon} \frac{(x_j - y_j) \eta(x-y)}{|x-y|^{n+1}} f(y) dy$$

and

$$\mathcal{R}_j^\psi f(x) := R_j(f - \psi * f).$$

Then

1. $\mathcal{R}_{j,\eta}$ and \mathcal{R}_j^ψ both map $h^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$;
2. for $f \in L^1(\mathbb{R}^n)$, $f \in h^1(\mathbb{R}^n) \iff \mathcal{R}_{j,\eta}(f) \in L^1(\mathbb{R}^n), j = 1, \dots, n \iff \mathcal{R}_j^\psi(f) \in L^1(\mathbb{R}^n), j = 1, \dots, n$.

As seen above, because convolution operators commute with the r_j , the boundedness from $h^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$ implies the boundedness on $h^1(\mathbb{R}^n)$, and extra cancellation conditions on the kernel are not needed (see also the remark following the proof of [24, Theorem 4]).

3.4 Inhomogeneous singular integral operators

We end this section by introducing a class of operators which are not necessarily of convolution type. As can be seen from the definition, however, for a convolution δ -kernel K and an η which decays sufficiently fast at infinity, the localized kernel $K\eta$ will be in this class.

Let $\Delta := \{(x, x) : x \in \mathbb{R}^n\}$ denote the diagonal in \mathbb{R}^{2n} .

Definition 3.3. Let $\delta \in (0, 1]$. We say T is an inhomogeneous singular integral operator associated with an inhomogeneous δ -kernel K if K is a locally integrable function on $\mathbb{R}^{2n} \setminus \Delta$ and there exist $\varepsilon > 0$ and C such that

1. $|K(x, y)| \leq C \min\{|x - y|^{-n}, |x - y|^{-n-\varepsilon}\}$ for all $(x, y) \in \mathbb{R}^{2n} \setminus \Delta$;
2. $|K(x, y) - K(x, z)| + |K(y, x) - K(z, x)| \leq C|y - z|^\delta |x - y|^{-n-\delta}$ for $2|y - z| \leq |x - y|$.

In addition, we assume that the operator T , defined by $Tf(x) := \int_{\mathbb{R}^n} K(x, y)f(y)dy$ for $f \in C_c^\infty(\mathbb{R}^n)$ and $x \notin \text{supp}(f)$, can be extended to a bounded operator on $L^2(\mathbb{R}^n)$.

This type of operator was introduced in [19] and a generalization was considered in [14], where it was shown that if certain cancellation conditions are imposed on T , then it is bounded from $h^p(\mathbb{R}^n)$ to $h^p(\mathbb{R}^n)$, $0 < p \leq 1$. If no cancellation conditions are imposed, we still have boundedness from $h^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$, as can be seen by noting that h^1 atoms are mapped to “pre-molecules” which are bounded in $L^1(\mathbb{R}^n)$ (see [15, Definition 3]). As T is a singular integral operator, T is also bounded from $L^p(\mathbb{R}^n)$ to itself for $1 < p < \infty$. We will consider the commutators of these operators with bmo functions in Section 5.

4 Local Hardy spaces related to commutators with bmo functions

In this section we introduce some “local” analogues of commutator maximal function defined in [34], and the atoms in [39], and consider the relationships between them.

4.1 Some Maximal Functions

Recall that the Hardy-Littlewood Maximal function defined for $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ by

$$Mf(x) := \sup_{B \ni x} |f|_B$$

is bounded on $L^p(\mathbb{R}^n)$ for $1 < p \leq \infty$ and also bounded from $L^1(\mathbb{R}^n)$ to $L^{1,\infty}(\mathbb{R}^n)$.

In connection with commutators, the following two maximal functions were introduced in [10, Section 6] in the one-dimensional case. Setting

$$M(b, f)(x) := \sup_{I \ni x} |(b(x) - b_I)f_I|,$$

it was shown in [10, Theorem IX] that $b \in \text{BMO}(\mathbb{R})$ if and only if $f \mapsto M(b, f)$ is bounded on $L^2(\mathbb{R})$. The proof uses [10, Theorem X] which states that for a different maximal function,

$$N(b, f)(x) := \sup_{I \ni x} |(bf)_I - b(x)f_I|,$$

$f \mapsto M(b, f)$ is bounded on $L^2(\mathbb{R})$ when $b \in \text{BMO}(\mathbb{R})$.

The commutator of M with multiplication by b , that is

$$[M, b](f)(x) := M(bf)(x) - b(x)Mf(x),$$

was studied in [1, 2, 37] and shown to be bounded on $L^p(\mathbb{R}^n)$ for $1 < p < \infty$ if and only if $b \in \text{BMO}(\mathbb{R}^n)$ and b is bounded below.

Finally, one has the following maximal function,

$$M_b f(x) = \sup_{B \ni x} \frac{1}{|B|} \int_B |b(x) - b(y)| |f(y)| dy,$$

which was studied in [1, 22, 42] and is sometimes denoted by C_b . In this case, as in [10], one has that M_b is bounded on $L^p(\mathbb{R}^n)$ for $1 < p < \infty$ if and only if $b \in \text{BMO}(\mathbb{R}^n)$. Moreover, for $b \in \text{BMO}(\mathbb{R}^n)$, [1, Corollary 1.11] gives the pointwise domination $M_b f(x) \lesssim \|b\|_{\text{BMO}} M^2 f(x)$ for $f \in L^1_{\text{loc}}(\mathbb{R}^n)$, and therefore

$$\|M_b\|_{L^p \rightarrow L^p} \lesssim \|b\|_{\text{BMO}}. \quad (4.1)$$

Ky [34] introduced the following commutator maximal function definition of the space H_b^1 .

Definition 4.1 ([34], Definition 2.2). Let $b \in \text{BMO}(\mathbb{R}^n)$ be nontrivial. The space $H_b^1(\mathbb{R}^n)$ consists of $f \in H^1(\mathbb{R}^n)$ such $[b, \mathfrak{M}_{\text{nt}}]f \in L^1(\mathbb{R}^n)$, where $[b, \mathfrak{M}_{\text{nt}}]f(x) := \mathfrak{M}_{\text{nt}}[b(x)f(\cdot) - b(\cdot)f(\cdot)](x)$ and \mathfrak{M}_{nt} is the non-tangential grand maximal functions defined by

$$\mathfrak{M}_{\text{nt}} f(x) := \sup\{|f * \phi_t(y)| : |y - x| < t, \varphi \in \mathcal{A}\}$$

with

$$\mathcal{A} = \{\phi \in \mathcal{S}(\mathbb{R}^n) : \|\phi\|_{\infty} + \|\nabla \phi\|_{\infty} \leq (1 + |x|^2)^{-n-1}\}.$$

The norm on H_b^1 is given by

$$\|f\|_{H_b^1} := \|f\|_{H^1} \|b\|_{\text{BMO}} + \|[b, \mathfrak{M}_{\text{nt}}]f\|_{L^1}.$$

Ky also defines a ‘‘local’’ version of the non-tangential grand maximal function \mathfrak{M}_{nt} by restricting t to $(0, 1)$. As pointed out following Proposition 2.1, this is equivalent to the maximal functions \mathcal{M} and \mathfrak{M} in terms of characterizing $h_b^1(\mathbb{R}^n)$. However, there is no definition of h_b^1 given in [34].

We choose the grand maximal function \mathcal{M} , using test functions of compact support, to give the analogue of Ky’s definition in the local case.

Definition 4.2. Given $b \in L^2_{\text{loc}}(\mathbb{R}^n)$, we define the maximal commutator function for $f \in L^2_{\text{loc}}(\mathbb{R}^n)$ to be

$$\mathcal{M}_b f(x) := [b, \mathcal{M}](x) := \mathcal{M}[b(x)f(\cdot) - b(\cdot)f(\cdot)](x).$$

Recall that in Proposition 2.1 (for $T = 1$) we defined, for $f \in \mathcal{S}'(\mathbb{R}^n)$,

$$\mathcal{M}(f)(x) := \sup_{\phi} |\langle f, \phi \rangle|,$$

where the supremum is taken over all $\phi \in C^1(\mathbb{R}^n)$ with $\text{supp}(\phi) \subset B(x, t)$ for some $0 < t < 1$, $\|\phi\|_{L^\infty} \leq |B(x, t)|^{-1}$ and $\|\nabla\phi\|_{L^\infty} \leq [t|B(x, t)|]^{-1}$. We therefore have that for $b, f \in L^2_{loc}(\mathbb{R}^n)$,

$$\mathcal{M}_b f(x) = \sup_{\phi} |\langle b(x)f(\cdot) - b(\cdot)f(\cdot), \phi \rangle| = \sup_{\phi} \left| \int_{\mathbb{R}^n} [b(x) - b(y)]f(y)\phi(y)dy \right|.$$

Since $(b(x) - b)f \in L^1_{loc}(\mathbb{R}^n)$ for almost every $x \in \mathbb{R}^n$, the integral is well-defined almost everywhere, and by the size and support conditions on ϕ ,

$$\mathcal{M}_b f(x) \lesssim M_b f(x). \quad (4.2)$$

The extension of this definition from $f \in L^2_{loc}(\mathbb{R}^n)$ to general $f \in h^1(\mathbb{R}^n)$, except in the special case where b is bounded, is not obvious. In fact, this is also the case for Ky's definition, Definition 4.1, and additional work, such as the sublinear decomposition [34, Theorem 3.1], is needed there.

4.2 Atomic commutator Hardy spaces

In [39], Pérez defined a restricted notion of atom with cancellation against a function $b \in \text{BMO}$.

Definition 4.3 ([39], Definition 1.3). For $b \in \text{BMO}(\mathbb{R}^n)$, a b -atom a such that for some cube Q

- (i) $\text{supp}(a) \subset Q$;
- (ii) $\|a\|_{L^\infty} \leq |Q|^{-1}$;
- (iii) $\int a = 0$ and $\int ab = 0$.

Pérez then proceeded to define an atomic space, which he called H_b^1 (Ky [34] uses \mathcal{H}_b^1 for this space) by taking all $f \in L^1$ which can be written as $f = \sum_{j=1}^{\infty} \lambda_j a_j$, where a_j are b -atoms and $\{\lambda_j\} \in \ell^1$. Note that since b -atoms are also H^1 atoms, such a decomposition will converge in the H^1 norm with $\|f\|_{H^1} \leq \sum |\lambda_j|$, and hence Pérez's space is contained in H^1 .

We give the following straightforward adaptation of Pérez's definition to the local case.

Definition 4.4. Fix $b \in \text{bmo}(\mathbb{R}^n)$. We say a is a Pérez h_b^1 atom if for some ball $B(x_0, r)$ we have

1. $\text{supp}(a) \subset B(x_0, r)$
2. $\|a\|_{L^2(\mathbb{R}^n)} \leq |B(x_0, r)|^{-\frac{1}{2}}$
3. $\int a = 0$ and $\int ab = 0$ whenever $r < 1$.

For $f \in L^1(\mathbb{R}^n)$, we say $f \in h_{\text{Pérez},b}^1(\mathbb{R}^n)$ if f can be written as $f = \sum_{j=1}^{\infty} \lambda_j a_j$, where a_j are Pérez h_b^1 atoms and $\{\lambda_j\} \in \ell^1$. For such functions, we define

$$\|f\|_{h_{\text{Pérez},b}^1} := \inf \sum_j |\lambda_j|,$$

where the infimum is taken over all such decompositions.

As in the homogeneous case, we note that a Pérez h_b^1 atom is a $(1, 2)$ atom in h^1 and therefore the decomposition converges in h^1 and we have $\|f\|_{h^1} \leq \sum |\lambda_j|$. Moreover, each Pérez h_b^1 atom supported in a ball $B = B(x_0, r)$ also satisfies that $a(b - c_B) \in h^1$, where c_B is as in (2.1). Indeed, we have that $a(b - c_B)$ is also supported in B , has integral zero whenever $r < 1$, and by (2.2)

$$\begin{aligned} \|a(b - c_B)\|_{L^s(\mathbb{R}^n)} &\leq \|a\|_{L^2(\mathbb{R}^n)} \|b - c_B\|_{L^p(\mathbb{R}^n)} \\ &\leq |B(x_0, r)|^{-\frac{1}{2}} \|b\|_{\text{bmo},p} |B(x_0, r)|^{\frac{1}{p}} = \|b\|_{\text{bmo},p} |B(x_0, r)|^{\frac{1}{s}-1} \end{aligned} \quad (4.3)$$

for $1 \leq s < 2$ and $\frac{1}{2} + \frac{1}{p} = \frac{1}{s}$; taking $s > 1$ makes $a(b - c_B)$ a multiple of a $(1, s)$ atom and we have

$$\|a(b - c_B)\|_{h^1} \lesssim \|b\|_{\text{bmo},p}.$$

We now give a variation on this definition which involves approximate cancellation conditions.

Definition 4.5. Fix $b \in \text{bmo}(\mathbb{R}^n)$ and let $C_b = \|b\|_{\text{bmo},2}$ as defined in (2.2). We say a is an approximate h_b^1 atom if for some ball $B(x_0, r)$ we have

1. $\text{supp}(a) \subset B(x_0, r)$;
2. $\|a\|_{L^2(\mathbb{R}^n)} \leq |B(x_0, r)|^{-\frac{1}{2}}$;
3. $\left| \int a \right| \leq \frac{1}{[\log(1 + r^{-1})]^2}$ and $\left| \int a(b - c_B) \right| \leq \frac{C_b}{\log(1 + r^{-1})}$.

For $f \in L^1(\mathbb{R}^n)$, we say $f \in h_{\text{atom},b}^1(\mathbb{R}^n)$ if f can be written as $f = \sum_{j=1}^{\infty} \lambda_j a_j$, where a_j are approximate h_b^1 atoms and $\{\lambda_j\} \in \ell^1$. For such functions, we define

$$\|f\|_{h_{\text{atom},b}^1} := \inf \sum_j |\lambda_j|,$$

where the infimum is taken over all such decompositions.

The choice of $C_b = \|b\|_{\text{bmo},2}$ guarantees that, by Cauchy-Schwarz, condition (2) implies condition (3) when $r \geq 1$. Thus every Pérez h_b^1 atom is an approximate h_b^1 atom, and

$$h_{\text{Pérez},b}^1(\mathbb{R}^n) \subset h_{\text{atom},b}^1(\mathbb{R}^n).$$

Moreover, comparing with Definition 2.2 for $R = 1$ and $q = 2$, we see, as in (4.3), that not only is every approximate h_b^1 atom a an approximate $(1, 2)$ atom (up to a factor of $\log 2$), but also $a(b - c_B)$ is a multiple of an approximate $(1, s)$ atom for $1 < s < 2$ and $\frac{1}{2} + \frac{1}{p} = \frac{1}{s}$, with

$$\|a(b - c_B)\|_{h^1} \lesssim \|b\|_{\text{bmo},p}. \quad (4.4)$$

Recall that in h^1 (or h^p), every atom (or molecule) with approximate cancellation conditions can be written as an infinite linear combination of atoms with exact cancellation - see [14, 18]. Is it possible to decompose an approximate h_b^1 atom into Pérez h_b^1 atoms with exact cancellation?

In the following proposition, we give a partial converse showing that every approximate h_b^1 atom can be written as a finite sum of atoms of integral zero, but we do not achieve the exact cancellation against b , and it is not clear if this can be done.

Proposition 4.1. *Suppose A is an approximate h_b^1 atom with respect to the ball $B(x_0, r)$, and $r < 1$. Then $A = \sum \lambda_j a_j$ where the sum is **finite**, a_j are approximate h_b^1 atoms supported in balls B_j , respectively, and in addition, when $r(B_j) < 1$,*

$$\int a_j = 0.$$

Moreover, for some constant depending only on n , $\sum |\lambda_j| \leq C_n$.

Proof. Set $\alpha = \int A \neq 0$ (otherwise we are done). Denote $B(0, 2^j r)$ by B_j , $j = 0, 1, 2, \dots, k$, where $2^{k-1}r < 1 \leq 2^k r$, and let $\eta_j = \frac{\chi_{B_j}}{|B_j|}$ be the normalized characteristic function of B_j . Then

$$A = A - \alpha \eta_0 + \sum_{j=1}^k \alpha (\eta_{j-1} - \eta_j) + \alpha \eta_k = \sum_{i=0}^{k+1} A_j$$

where

$$\begin{aligned} A_0 &:= A - \alpha \eta_0 \\ A_j &:= \alpha (\eta_{j-1} - \eta_j), \quad j = 1, \dots, k, \quad \text{and} \\ A_{k+1} &:= \alpha \eta_k. \end{aligned}$$

First observe that $\text{supp}(A_j) \subset B_j$ for $j = 1, \dots, k$, and $\text{supp}(A_{k+1}) \subset B_k$. Moreover, since $\int \eta_j = 1$ for all j , we have that $\int A_j = 0$ for $j = 0, \dots, k$.

Write $A_j = \lambda_j a_j$ where

$$a_0 := \frac{1}{2} A_0, \quad \lambda_0 := 2, \quad a_{k+1} := \alpha^{-1} A_{k+1}, \quad \lambda_{k+1} := \alpha,$$

and

$$a_j := \frac{A_j}{\alpha 2^n \log(1 + [2^j r]^{-1})}, \quad \lambda_j := \alpha 2^n \log(1 + [2^j r]^{-1}), \quad j = 1, \dots, k.$$

Since $|\alpha| \leq \|A\|_{L^1} \leq 1$, $\|\eta_j\|_{L^\infty} \leq |B_j|^{-1}$, and $2^n \log(1 + [2^j r]^{-1}) > 1$ for $j = 1, \dots, k$, we have that $\|a_0\|_{L^2} \leq |B_0|^{-\frac{1}{2}}$, $\|a_j\|_{L^\infty} \leq |B_j|^{-1}$ for $j = 1, \dots, k$, and $\|a_{k+1}\|_{L^\infty} \leq |B_k|^{-1}$. Thus, all the a_j satisfy Conditions (1) and (2) in Definition 4.5 and in addition $\int a_j = 0$ for $j = 0, \dots, k$. Since $r(B_k) = 2^k r \geq 1$, the latter condition is not required for a_{k+1} .

To verify Condition (3) in Definition 4.5, we need to check that the a_j satisfy the approximate cancellation condition against b . For $j = 0$, we get the approximate cancellation for a_0 from that of A :

$$\left| \int a_0 b \right| = \frac{1}{2} \left| \int A b - \alpha \int b \eta_0 \right| = \frac{1}{2} \left| \int A [b - b_B] \right| \leq \frac{C_b}{\log(1 + r^{-1})}.$$

For $j = 1, \dots, k$, we have, using (2.4),

$$\begin{aligned} \left| \int a_j b \right| &= \frac{1}{2^n \log(1 + [2^j r]^{-1})} \left| \int b(\eta_{j-1} - \eta_j) \right| \\ &= \frac{1}{2^n \log(1 + [2^j r]^{-1})} |b_{2^{j-1}B} - b_{2^j B}| \\ &\leq \frac{\|b\|_{\text{bmo}}}{\log(1 + [2^j r]^{-1})}, \end{aligned}$$

and for $j = k + 1$, since $r(B_k) = 2^k r \geq 1$,

$$\left| \int a_{k+1} b \right| = |b_{B_k}| \leq \frac{\|b\|_{\text{bmo}}}{\log(1 + (2^k r)^{-1})}.$$

If, as noted after Definition 4.5, we take $C_b = \|b\|_{\text{bmo},2} \geq \|b\|_{\text{bmo}}$, then we can conclude that a_j for $j = 0, \dots, k + 1$ are all approximate h_b^1 atoms.

Finally, we have

$$\begin{aligned} \sum |\lambda_j| &= 2 + \alpha 2^n \sum_{j=1}^k \log(1 + [2^j r]^{-1}) + \alpha \\ &\leq 3 + \frac{2^n}{[\log(1 + r^{-1})]^2} \sum_{j=1}^k \log(1 + [2^j r]^{-1}) \\ &\leq 3 + \frac{2^n k \log(1 + r^{-1})}{[\log(1 + r^{-1})]^2} \\ &\leq 3 + \frac{2^n (\log_2 r^{-1} + 1)}{[\log(1 + r^{-1})]} \\ &\leq C_n. \end{aligned}$$

□

4.3 Relations between the spaces

For the homogeneous case, Ky shows (see [34, Theorem 5.2]) that

$$\mathcal{H}_b^1(\mathbb{R}^n) \subset H_b^1(\mathbb{R}^n) \subset H^1(\mathbb{R}^n),$$

where \mathcal{H}_b^1 is Pérez atomic space and H_b^1 is the maximal commutator space, but does not discuss whether the inclusion is proper. The result of [26] shows that the last inclusion is proper, at least in dimension 1, unless b is a constant.

For the nonhomogeneous case, the discussion following the definitions of the atomic spaces in the previous section gave us the inclusions

$$h_{\text{Perez},b}^1(\mathbb{R}^n) \subset h_{\text{atom},b}^1(\mathbb{R}^n) \subset h^1(\mathbb{R}^n).$$

In the trivial case when b is a constant function, the cancellation conditions against b reduce to the conditions on the integral of the atoms. Thus every h^1 atom with exact cancellation is a Pérez h_b^1 atom, so

$$h_{\text{Perez},b}^1(\mathbb{R}^n) = h_{\text{atom},b}^1(\mathbb{R}^n) = h^1(\mathbb{R}^n). \quad (4.5)$$

In the next section will see later that the second equality can hold for b in a nontrivial subspace of bmo .

While we have not defined a maximal space, we can consider the action of the commutator maximal function \mathcal{M}_b on atoms.

Proposition 4.2. *Let $b \in \text{bmo}(\mathbb{R}^n)$. Then for every h_b^1 atom a ,*

$$\|\mathcal{M}_b a\|_{L^1} \lesssim \|b\|_{\text{bmo}}.$$

Proof. Let a be an approximate h_b^1 atom with support in $B = B(x_0, r)$. We want to bound $\mathcal{M}_b(a)$.

For the local estimate, using (4.2), we can apply the L^p boundedness of the maximal commutator operator $M_b f$ defined in Section 4.1, with the bound (4.1). Thus, using the L^2 size condition on a we have

$$\int_{2B} \mathcal{M}_b(a) \leq |2B|^{1/2} \|M_b a\|_{L^2(2B)} \lesssim |B|^{1/2} \|a\|_{L^2} \|M_b\|_{L^2 \rightarrow L^2} \lesssim \|b\|_{\text{bmo}}.$$

Next we handle the integral on $(2B)^c$. Note that

$$\left| \int_{B(x,t)} \phi(y) [b(x) - b(y)] a(y) dy \right| \neq 0$$

implies that there exists $y \in B(x, t) \cap B(x_0, r)$, which in turn implies

$$r \leq \frac{|x - x_0|}{2} \leq |x - x_0| - |x_0 - y| \leq |x - y| \leq t < 1. \quad (4.6)$$

This cannot happen when $r \geq 1$, so in that case the integral vanishes for all test functions ϕ and $\mathcal{M}_b(a) = 0$ on $(2B)^c$.

For $r < 1$, we may assume, by Proposition 4.1, that $\int a = 0$. Thus we have, for almost all $x \in (2B)^c$,

$$\begin{aligned} & \left| \int [b(x) - b(y)] a(y) \phi(y) dy \right| \\ & \leq \left| \int [b(x) - b(y)] a(y) [\phi(y) - \phi(x_0)] dy \right| + \left| \phi(x_0) \int b(y) a(y) dy \right| \\ & \leq |b(x) - b_B| \|\nabla \phi\|_{L^\infty} r \int |a(y)| dy + \|\nabla \phi\|_{L^\infty} r \int |b(y) - b_B| |a(y)| dy \\ & \quad + \|\phi\|_{L^\infty} \left| \int b(y) a(y) dy \right| \\ & \lesssim \frac{|b(x) - b_B| r}{|x - x_0|^{n+1}} + \frac{r \|b\|_{\text{bmo},2}}{|x - x_0|^{n+1}} + \frac{C_b}{|x - x_0|^n \log(1 + r^{-1})}, \end{aligned} \quad (4.7)$$

where in the last step we used the conditions on ϕ , (4.6), the L^1 estimate on $a(b - b_B)$ (see (4.3) with $s = 1, p = 2$), and condition (3) in Definition 4.5. Since the estimate above is independent of ϕ , it holds for $\mathcal{M}_b(a)(x)$ and therefore,

again using (4.6),

$$\begin{aligned}
\int_{(2B)^c} \mathcal{M}_b(a)(x)dx &= \int_{2r < |x-x_0| < 2} \mathcal{M}_b(a)(x)dx \\
&\lesssim \int_{(2B)^c} \frac{|b(x) - b_B|r}{|x-x_0|^{n+1}} + \int_{(2B)^c} \frac{r\|b\|_{\text{bmo},2}}{|x-x_0|^{n+1}} \\
&\quad + \int_{2r < |x-x_0| < 2} \frac{C_b}{|x-x_0|^n \log(1+r^{-1})} \\
&\lesssim \|b\|_{\text{bmo}}.
\end{aligned} \tag{4.8}$$

Here we have used (2.3), (2.2), and the fact that $C_b = \|b\|_{\text{bmo},2}$. \square

If we take an h^1 atom a without assuming any cancellation against b , the boundedness of the maximal function \mathcal{M}_b turns out to be equivalent to whether $a(b - c_B)$ belongs to h^1 , where B is the ball containing the support of a . Such boundedness gives us automatically the approximate cancellation against b .

Proposition 4.3. *Let $b \in \text{bmo}(\mathbb{R}^n)$, b nontrivial. Suppose a is an h^1 atom supported in a ball $B = B(x_0, r)$, and has vanishing integral when $r < 1$. Then*

$$\|\mathcal{M}_b a\|_{L^1} \lesssim \|a(b - c_B)\|_{h^1} + \|b\|_{\text{bmo}} \lesssim \|\mathcal{M}_b a\|_{L^1} + \|b\|_{\text{bmo}}.$$

Moreover,

$$\left| \int ab \right| \lesssim \min \left\{ \frac{\|\mathcal{M}_b a\|_{L^1} + \|b\|_{\text{bmo}}}{\log(1+r^{-1})}, \|b\|_{\text{bmo}} \right\}.$$

Proof. We assume that a satisfies an L^2 size condition. First note that by (4.3), $a(b - c_B) \in L^s(\mathbb{R}^n)$ for $1 < s < 2$, and it has compact support, $a(b - c_B)$ is in $h^1(\mathbb{R}^n)$. By Definition 4.2, we have, for almost every $x \in \mathbb{R}^n$,

$$\mathcal{M}_b(a)(x) = \mathcal{M}([b(x) - b]a)(x) \leq |b(x) - c_B| \mathcal{M}(a)(x) + \mathcal{M}([b - c_B]a)(x),$$

hence

$$\begin{aligned}
\|\mathcal{M}_b(a)\|_{L^1} &\leq \int |b(x) - c_B| \mathcal{M}(a)(x)dx + \|\mathcal{M}([b - c_B]a)\|_{L^1} \\
&\lesssim \int |b(x) - c_B| \mathcal{M}(a)(x)dx + \|a(b - c_B)\|_{h^1}.
\end{aligned}$$

Conversely,

$$\mathcal{M}([b - c_B]a)(x) \leq \mathcal{M}_b(a)(x) + |b(x) - c_B| \mathcal{M}(a)(x),$$

so

$$\|a(b - c_B)\|_{h^1} \leq \|\mathcal{M}_b(a)\|_{L^1} + \int |b(x) - c_B| \mathcal{M}(a)(x)dx.$$

It thus remains to show that the integral of $|b - c_B| \mathcal{M}(a)$ is controlled by $\|b\|_{\text{bmo}}$. The arguments are similar to those in the proof of Proposition 4.2. First, we have

$$\int_{2B} |b - c_B| \mathcal{M}(a) \leq \|b - c_B\|_{L^2(2B)} \|Ma\|_{L^2(2B)} \lesssim \|b\|_{\text{bmo}} |B|^{\frac{1}{2}} \|a\|_{L^2} \lesssim \|b\|_{\text{bmo}}.$$

Moreover, (4.6) implies that if $r \geq 1$, $\mathcal{M}(a)$ vanishes outside $2B$. For $r < 1$, we can use the cancellation condition on a to write, as in (4.7), for every test function ϕ , and for almost every $x \in (2B)^c$,

$$\begin{aligned} |b(x) - c_B| \left| \int a(y)\phi(y)dy \right| &\leq |b(x) - b_B| \|\nabla\phi\|_{L^\infty} r \int |a(y)|dy \\ &\lesssim \frac{|b(x) - b_B|r}{|x - x_0|^{n+1}}. \end{aligned}$$

The integral on $(2B)^c$ is then estimated as in the first term of (4.8).

By applying Proposition 2.3 to $a(b - c_B) \in h^1(\mathbb{R}^n)$, followed by the estimates above, we have

$$\left| \int ab \right| = \left| \int a(b - c_B) \right| \lesssim \frac{\|a(b - c_B)\|_{h^1}}{\log(1 + r^{-1})} \lesssim \frac{(\|\mathcal{M}_b a\|_{L^1} + \|b\|_{\text{bmo}})}{\log(1 + r^{-1})}.$$

The other upper bound follows from the duality of bmo and h^1 , or by Cauchy-Schwartz. \square

It is interesting to note that if we had not assumed exact cancellation on a in the hypotheses of Proposition 4.3 when $r < 1$, we would have ended up having to estimate an extra term of the form

$$\int_{2r < |x - x_0| < 2} \frac{|b(x) - b_B|}{|x - x_0|^n} dx \left| \int a \right|.$$

Looking at the proof of (2.3), if $\delta = 0$ and $p = 1$, the integral in x can be estimated by $\|b\|_{\text{bmo}}(\log \frac{1}{r})^2$. Therefore, we would need a cancellation condition of the form $|\int a| \leq [\log(1 + r^{-1})]^{-2}$, as in condition (3) of Definition 4.5.

Combining Propositions 4.1, 4.2 and 4.3, we get the following.

Corollary 4.1. *Let $f \in h^1(\mathbb{R}^n)$. Then the following are equivalent.*

1. *The function $f \in h_{\text{atom}, b}^1(\mathbb{R}^n)$.*
2. *There exists a sequence $\{\lambda_j\} \in \ell^1$ and a collection of h^1 atoms $\{a_j\}$, where B_j denotes the ball containing the support of a_j and a_j has vanishing integral when $r(B_j) < 1$, so that $f = \sum_j \lambda_j a_j$ and $\mathcal{M}_b a_j$ are uniformly bounded in $L^1(\mathbb{R}^n)$.*
3. *There exists a sequence $\{\lambda_j\} \in \ell^1$ and a collection of h^1 atoms $\{a_j\}$, where B_j denotes the ball containing the support of a_j and a_j has vanishing integral when $r(B_j) < 1$, so that $f = \sum_j \lambda_j a_j$ and*

$$\sum_j \lambda_j a_j (b - c_{B_j}) \text{ converges absolutely in } h^1(\mathbb{R}^n).$$

4.4 The case of b in lmo

We now come to the promised result which gives us the second equality in (4.5) even when b is not constant. Recall that the pointwise multipliers of $\text{bmo}(\mathbb{R}^n)$, and hence $h^1(\mathbb{R}^n)$, were identified as the elements of $L^\infty \cap \text{lmo}(\mathbb{R}^n)$ in [5]. It is therefore not surprising that we have the following results.

Theorem 4.6. *Let $b \in \text{lmo}(\mathbb{R}^n)$. Then $h_{\text{atom},b}^1(\mathbb{R}^n) = h^1(\mathbb{R}^n)$. Moreover,*

$$\|f\|_{h^1} \leq \|f\|_{h_{\text{atom},b}^1} \lesssim \frac{\|b\|_{\text{LMO}_{\text{loc}}}}{\|b\|_{\text{BMO}_{\text{loc}}}} \|f\|_{h^1}.$$

Proof. It was already observed following Definition 4.5 that for $b \in \text{bmo}$, h_b^1 atoms are approximate h^1 atoms (up to a factor of $\log 2$) and therefore $h_b^1(\mathbb{R}^n) \subset h^1(\mathbb{R}^n)$ with $\|f\|_{h^1} \lesssim \|f\|_{h_{\text{atom},b}^1}$. We only need the lmo condition for the reverse inclusion.

Let a be an h^1 atom with support in the ball $B = B(x_0, r)$, with L^2 size condition and such that $\int a = 0$. This means that conditions (1), (2), and the first part of (3) in Definition 4.5 are satisfied, and we only need to check the approximate cancellation against b . Since this holds automatically for $r \geq 1$, we assume $r < 1$. By the L^2 size condition on a ,

$$\begin{aligned} \left| \int_{\mathbb{R}^n} a(x)[b(x) - b_B] dx \right| &\leq \frac{1}{\log(1+r^{-1})} \frac{\log(1+r^{-1})}{|B|^{\frac{1}{2}}} \left(\int_B |b(x) - b_B|^2 dx \right)^{\frac{1}{2}} \\ &\leq \frac{\|b\|_{\text{LMO}_{\text{loc},2}}}{\log(1+r^{-1})}, \end{aligned}$$

where

$$\|b\|_{\text{LMO}_{\text{loc},2}} := \sup_{r(B) < 1} \log(1+r(B)^{-1}) \left(\int_B |b(x) - b_B|^2 dx \right)^{\frac{1}{2}}.$$

Since

$$\gamma := \frac{\log 2 \|b\|_{\text{BMO}_{\text{loc},2}}}{\|b\|_{\text{LMO}_{\text{loc},2}}} \leq 1$$

and $\log 2 \|b\|_{\text{BMO}_{\text{loc},2}} \leq \|b\|_{\text{bmo},2} = C_b$, we have that γa is an h_b^1 atom.

Finally, for $f \in h^1$ with an atomic decomposition $f = \sum_{j=1}^{\infty} \lambda_j a_j$, we can regard it as a decomposition using h_b^1 atoms multiplied by γ^{-1} , so $f \in h_{\text{atom},b}^1(\mathbb{R}^n)$ with

$$\|f\|_{h_{\text{atom},b}^1} \leq \gamma^{-1} \sum_{j=1}^{\infty} |\lambda_j|.$$

Taking the infimum over all such decompositions, we get $\|f\|_{h_{\text{atom},b}^1} \leq \gamma^{-1} \|f\|_{h^1}$. The inequality in the statement of the theorem is obtained by applying the John-Nirenberg inequality for LMO - see [43]. \square

We give a partial converse to the theorem.

Proposition 4.4. *Let $b \in L_{\text{loc}}^1(\mathbb{R}^n)$. Suppose there exists a constant C_b such that every h^1 atom a satisfies*

$$\left| \int ab \right| \leq \min \left\{ C_b, \frac{C_b \log(2)}{\log(1+r^{-1})} \right\},$$

where r is the radius of the ball B containing the support of a . Then $b \in \text{lmo}(\mathbb{R}^n)$.

Proof. We first handle a ball B with radius $r < 1$. We define $s(x) := \text{sgn}[b(x) - b_B]$ and $a(x) := \frac{s(x) - s_B}{|B|} \chi_B(x)$. Then a is an h^1 atom. Moreover, we can write

$$\begin{aligned} \log(1 + r^{-1}) \int_B |b(x) - b_B| dx &= \log(1 + r^{-1}) \int_B s(x)(b(x) - b_B) dx \\ &= \frac{\log(1 + r^{-1})}{|B|} \int_B (s(x) - s_B)(b(x) - b_B) dx \\ &= \log(1 + r^{-1}) \int_B a(x)b(x) dx \\ &= \log(1 + r^{-1}) \left| \int_B ab \right| \\ &\leq C_b. \end{aligned}$$

If $r \geq 1$, we define $a(x) := \frac{\text{sgn}[b(x)]}{|B|} \chi_B(x)$. A similar argument shows that

$$\frac{1}{|B|} \int_B |b(x)| dx = \left| \int_B a(x)b(x) dx \right| \leq C_b.$$

Therefore, we can conclude that $b \in \text{lmo}(\mathbb{R}^n)$ with $\|b\|_{\text{lmo}} \leq C_b$. \square

In a similar vein, we give a partial converse to Proposition 4.2

Proposition 4.5. *Let $b \in \text{bmo}(\mathbb{R}^n)$. Suppose there exists a constant κ_b such that every h^1 atom a satisfies*

$$\|\mathcal{M}_b a\|_{L^1} \leq \kappa_b.$$

Then b satisfies

$$\sup_{r(B) < 1} \frac{1}{|B|} \| |b - b_B| \chi_B \|_{h^1} + \sup_{r(B) \geq 1} \frac{1}{|B|} \| |b| \chi_B \|_{h^1} < \infty.$$

Proof. Fix a ball B with radius $r < 1$. As in the previous proof, set $s(x) := \text{sgn}[b(x) - b_B]$ and $a(x) := \frac{s(x) - s_B}{|B|} \chi_B(x)$. Then a is an h^1 atom, and from Proposition 4.3,

$$\begin{aligned} \| |b - b_B| \frac{\chi_B}{|B|} \|_{h^1} &= \| (b - b_B) s(x) \frac{\chi_B}{|B|} \|_{h^1} \\ &\leq \| a(b - b_B) \|_{h^1} + |s_B| \left\| \frac{(b - b_B) \chi_B}{|B|} \right\|_{h^1} \\ &\lesssim \| \mathcal{M}_b(a) \|_{L^1} + \| b \|_{\text{bmo}} + |s_B| \left\| \frac{(b - b_B) \chi_B}{|B|} \right\|_{h^1} \\ &\lesssim \| \mathcal{M}_b(a) \|_{L^1} + \| b \|_{\text{bmo}}. \end{aligned}$$

In the last step we used the fact that $|s_B| \leq 1$ and $\frac{(b - b_B) \chi_B}{|B| \|b\|_{\text{bmo}, 2}}$ is a $(1, 2)$ atom.

For B with radius $r \geq 1$, we again put $a(x) = \frac{\text{sgn}(b(x)) \chi_B(x)}{|B|}$ and use Proposition 4.3:

$$\| |b| \frac{\chi_B}{|B|} \|_{h^1} = \| a(b - c_B) \|_{h^1} \lesssim \| \mathcal{M}_b(a) \|_{L^1} + \| b \|_{\text{bmo}}.$$

\square

One interesting corollary from Proposition 4.2 and 4.5 is the following.

Corollary 4.2. *Let $b \in \text{bmo}(\mathbb{R}^n)$. Then the following are equivalent.*

1. *The function b is in $\text{lmo}(\mathbb{R}^n)$.*
2. *The function b satisfies*

$$A_b := \sup_{r(B) < 1} \frac{1}{|B|} \| |b - b_B| \chi_B \|_{h^1} + \sup_{r(B) \geq 1} \frac{1}{|B|} \| |b| \chi_B \|_{h^1} < \infty.$$

Proof. Suppose $b \in \text{lmo}(\mathbb{R}^n)$. Then from the proof of Theorem 4.6, for an h^1 atom a , γa is an h_b^1 atom, hence by Proposition 4.2

$$\| \mathcal{M}_b a \|_{L^1} \lesssim \gamma^{-1} \| b \|_{\text{bmo}} \approx \kappa_b := \frac{\| b \|_{\text{LMO}_{\text{loc}}}}{\| b \|_{\text{BMO}_{\text{loc}}}} \| b \|_{\text{bmo}},$$

which is exactly the hypothesis of Proposition 4.5.

Conversely, suppose $A_b < \infty$. Observe that from Proposition 2.3, we can estimate

$$\sup_{r(B) < 1} \frac{\log(1 + [r(B)]^{-1})}{|B|} \int_B |b(x) - b_B| dx \lesssim \sup_{r(B) < 1} \frac{1}{|B|} \| |b - b_B| \chi_B \|_{h^1}.$$

Meanwhile, using the fact that $\| f \|_{L^1(\mathbb{R}^n)} \leq \| f \|_{h^1(\mathbb{R}^n)}$, we have

$$\sup_{r(B) \geq 1} \frac{1}{|B|} \int_B |b(x)| dx \leq \sup_{r(B) \geq 1} \frac{1}{|B|} \| |b| \chi_B \|_{h^1}.$$

Thus $\| b \|_{\text{lmo}} \lesssim A_b < \infty$. □

5 Boundedness of the commutator of inhomogeneous singular integral operators with bmo functions

At the end of Section 3, we introduced a class of Calderón-Zygmund operators where the kernel has better decay away from the diagonal (see Definition 3.3). We will now show that the commutators of these operators with bmo functions take h_b^1 atoms to L^1 .

Theorem 5.1. *Suppose $b \in \text{bmo}(\mathbb{R}^n)$ and T is a inhomogeneous singular integral operator. Then $[b, T](a) \in L^1(\mathbb{R}^n)$ and*

$$\| [b, T] a \|_{L^1} \lesssim \| b \|_{\text{bmo}}$$

for all h_b^1 atoms a . Thus $[b, T] : h_{\text{finite}, b}^1(\mathbb{R}^n) \mapsto L^1(\mathbb{R}^n)$, where $h_{\text{finite}, b}^1(\mathbb{R}^n)$ denotes the space of finite linear combinations of h_b^1 atoms.

Proof. Let a be an h_b^1 atom with $\text{supp}(a) \subset B = B(x_0, r)$. Note that we can write the commutator acting on a as $[b, T](a) = (b - c_B)T(a) - T(a(b - c_B))$.

We first show that $T(a(b - c_B)) \in L^1(\mathbb{R}^n)$. By the L^p boundedness of T for $p = \frac{3}{2}$, we have

$$\begin{aligned} \|T(a(b - c_B))\|_{L^1(2B)} &\leq |2B|^{\frac{1}{3}} \|T(a(b - c_B))\|_{L^{\frac{3}{2}}(2B)} \\ &\lesssim |B|^{\frac{1}{3}} \|T\|_{L^{\frac{3}{2} \rightarrow L^{\frac{3}{2}}} \|a(b - c_B)\|_{L^{\frac{3}{2}}(\mathbb{R}^n)} \\ &\lesssim \|T\|_{L^{\frac{3}{2} \rightarrow L^{\frac{3}{2}}} |B|^{\frac{1}{3}} \|b - c_B\|_{L^6(B)} \|a\|_{L^2(B)} \\ &\lesssim \|T\|_{L^{\frac{3}{2} \rightarrow L^{\frac{3}{2}}} |B|^{-\frac{1}{6}} \|b - c_B\|_{L^6(B)} \lesssim \|T\|_{L^{\frac{3}{2} \rightarrow L^{\frac{3}{2}}} \|b\|_{\text{bmo},6}. \end{aligned}$$

Looking at $x \in (2B)^c$, first consider the case $r < 1$. From Proposition 4.1, we may assume $\int a = 0$. Since $x \notin \text{supp}(a(b - b_B))$, we can use the kernel representation to write

$$\begin{aligned} &\|T(a(b - b_B))\|_{L^1((2B)^c)} \\ &\leq \int_{(2B)^c} \left| \int_B K(x, y) a(y) (b(y) - b_B) dy \right| dx \\ &\leq \int_{(2B)^c} \left| \int_B [K(x, y) - K(x, x_0)] a(y) (b(y) - b_B) dy \right| dx + \int_{(2B)^c} |K(x, x_0)| \left| \int ab \right| dx \\ &=: I + II. \end{aligned}$$

As $2|y - x_0| < 2r \leq |x - x_0|$, we can use the smoothness of K (condition (2) in Definition 3.3) to estimate

$$\begin{aligned} I &\lesssim \int_{(2B)^c} \int_B \frac{|y - x_0|^\delta}{|x - x_0|^{n+\delta}} |a(y)| |b(y) - b_B| dy dx \\ &\lesssim \int_{(2B)^c} \frac{r^\delta}{|x - x_0|^{n+\delta}} \int_B |a(y)| |b(y) - b_B| dy dx \\ &\leq \int_{(2B)^c} \frac{Cr^\delta}{|x - x_0|^{n+\delta}} dx \|b\|_{\text{bmo},2} \\ &\lesssim \|b\|_{\text{bmo}}. \end{aligned}$$

Using the decay property of K (condition (2) in Definition 3.3) and the cancellation of a against b , recalling that we chose $C_b = \|b\|_{\text{bmo},2}$, we have

$$\begin{aligned} II &\leq \int_{(2B)^c} |K(x, x_0)| \left| \int ab \right| dx \\ &\lesssim \left(\int_{2r \leq |x - x_0| \leq 1} \frac{1}{|x - x_0|^n} dx + \int_{|x - x_0| \geq 1} \frac{1}{|x - x_0|^{n+\varepsilon}} dx \right) \frac{C_b}{\ln(1 + r^{-1})} \\ &\lesssim \|b\|_{\text{bmo},2} \frac{\ln(1 + r^{-1}) + 1}{\ln(1 + r^{-1})} \lesssim \|b\|_{\text{bmo}}. \end{aligned}$$

When $r \geq 1$, using the fact that $2|y - x_0| \leq |x - x_0|$ implies $|x - y| \geq \frac{1}{2}|x - x_0|$,

the decay of K again gives

$$\begin{aligned}
\|T(a(b - c_B))\|_{L^1((2B)^c)} &\leq \int_{(2B)^c} \left| \int_B K(x, y) a(y) b(y) dy \right| dx \\
&\lesssim \int_{|x-x_0| \geq 2r} \frac{1}{|x-x_0|^{n+\varepsilon}} \|b\|_{\text{bmo},2} \\
&\lesssim \int_{|x-x_0| \geq 2} \frac{1}{|x-x_0|^{n+\varepsilon}} \|b\|_{\text{bmo},2} \lesssim \|b\|_{\text{bmo}}.
\end{aligned}$$

Now we can handle the term $(b - c_B)T(a)$. Using the boundedness of T on L^2 , we have

$$\begin{aligned}
\|(b - c_B)T(a)\|_{L^1(2B)} &\leq \|b - c_B\|_{L^2(2B)} \|T(a)\|_{L^2(\mathbb{R}^n)} \\
&\lesssim \|b\|_{\text{bmo},2} |B|^{\frac{1}{2}} \|T\|_{L^2 \rightarrow L^2} \|a\|_{L^2(\mathbb{R}^n)} \\
&\lesssim \|b\|_{\text{bmo}} \|T\|_{L^2 \rightarrow L^2}.
\end{aligned}$$

When $r < 1$, we apply the cancellation of a , the smoothness of K and (2.3) to get

$$\begin{aligned}
\|(b - b_B)T(a)\|_{L^1((2B)^c)} &\leq \int_{(2B)^c} |b(x) - b_B| \left| \int_B [K(x, y) - K(x, x_0)] a(y) dy \right| dx \\
&\leq \int_{(2B)^c} |b(x) - b_B| \frac{r^\delta}{|x - x_0|^{n+\delta}} dx \lesssim \|b\|_{\text{bmo}}. \quad (5.1)
\end{aligned}$$

When $r \geq 1$, similar to the estimate of $\|T(a(b - c_B))\|_{L^1((2B)^c)}$, we have, again by (2.3),

$$\begin{aligned}
\|(b - c_B)T(a)\|_{L^1((2B)^c)} &\leq \int_{(2B)^c} |b(x)| \left| \int_B K(x, y) a(y) dy \right| dx \\
&\leq \int_{(2B)^c} \frac{|b(x)|}{|x - x_0|^{n+\varepsilon}} dx \\
&\leq \int_{(2B)^c} r^\varepsilon \frac{|b(x) - c_B|}{|x - x_0|^{n+\varepsilon}} dx \lesssim \|b\|_{\text{bmo}}. \quad (5.2)
\end{aligned}$$

We have thus shown $\|[b, T]a\|_{L^1} \lesssim \|b\|_{\text{bmo}}$ for all h_b^1 atoms a . Finally, $f \in h_{\text{finite},b}^1(\mathbb{R}^n)$ means it can be written as a finite linear combination $\sum_k \gamma_k a_k$ of h_b^1 atoms, so by linearity and taking the infimum over all such decompositions, we have

$$\|[b, T]f\| \lesssim \|b\|_{\text{bmo}} \inf \sum_k |\gamma_k|.$$

□

We cannot extend the conclusion to all of $h_{\text{atom},b}^1(\mathbb{R}^n)$ since we do not know whether for $f \in h_{\text{finite},b}^1(\mathbb{R}^n)$, the infimum $\inf \sum_k |\gamma_k|$ taken over all finite linear combinations of atoms is comparable to $\|f\|_{h_{\text{atom},b}^1}$, the infimum taken over all possible atomic decomposition.

When $h_{\text{atom},b}^1(\mathbb{R}^n) = h^1(\mathbb{R}^n)$, it is true that the infimum over the finite linear combinations of atoms is comparable to the h^1 norm, and it suffices to

verify the boundedness on one h^1 atom in order to get a bounded extension to the whole space (see [36, Theorem 3.1, Corollary 3.4] for the case of H^1 and [27, Proposition 7.1] for the case of h^1 on a space of homogeneous type). This gives us the following corollary.

Corollary 5.1. *Let $b \in \text{lmo}(\mathbb{R}^n)$, and T be a inhomogeneous singular integral operator. Then there exists a unique extension L_b of $[b, T]$ that is bounded from $h^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$.*

In particular, for $b \in \text{lmo}(\mathbb{R}^n)$, we can apply this corollary to show the boundedness of the commutators $[b, \mathcal{R}_{j,\eta}]$ and $[b, \mathcal{R}_j^\psi]$ defined in Corollary 3.1, provided the resulting localized convolution kernels have extra decay away from the origin. This requires some extra assumptions on η and ψ in addition to those in Theorem 3.2.

We can also compare this corollary with the results of [28]. While our assumption $b \in \text{lmo}$ is stronger than their assumption $b \in \text{LMO}_\infty$, which allows the oscillation to grow on large balls, and they obtain h^1 to h^1 boundedness, the operators they consider, which are pseudo-differential operators, have kernels which have much better decay and smoothness (see [28, Proposition 2.1]).

In order to get h^1 to h^1 boundedness of $[b, T]$, we need to impose some approximate cancellation conditions. We follow the ideas in [14, Definition 5.1].

Definition 5.2. Suppose T is an inhomogeneous singular integral operator. Given a ball B , denote by $L_0^2(\mathbb{R}^n)$ those $g \in L^2(B)$ with $\int g = 0$. For $b \in \text{bmo}$, define $T_B^*(b)$, relative to this ball B , in the distributional sense, by

$$\langle T_B^*(b), g \rangle := \int_{\mathbb{R}^n} (b - b_B) Tg(x) dx, \quad \forall g \in L_0^2(B). \quad (5.3)$$

As in (5.1), the conditions on the kernel and the vanishing integral of g guarantee that the integral on the right-hand-side of (5.3) converges absolutely and $T_B^*(b)$ is a bounded linear functional on $L_0^2(B)$ with norm bounded by a constant multiple of $\|b\|_{\text{bmo}}|B|^{1/2}$. Thus we can identify $T_B^*(b)$ with an equivalence class of functions in $L^2(B)$ modulo constants (similarly to the case $p = 1$ in [14, Remark 5.4]). In particular, we can impose the following condition, denoting $f := T_B^*(b)$, without ambiguity:

$$\left(\int_B |f - f_B|^2 \right)^{1/2} \leq \log(1 + r(B)^{-1}). \quad (5.4)$$

Similarly to what was pointed out in [14, Remark 5.4] for the case of $T^*((\cdot - x_B)^\alpha)$, this condition is not the same as requiring $T^*(b) \in \text{LMO}_{\text{loc}}(\mathbb{R}^n)$ since $T_B^*(b)$ is defined relative to a specific ball B , using the L^2 pairing, not the pairing of bmo and h^1 , and we are imposing condition (5.4) only for that ball.

While the local Campanato-type conditions used in [14, Theorem 5.3] to guarantee that inhomogeneous singular integral operators are bounded on $h^p(\mathbb{R}^n)$ are also expressed in terms of individual balls, in the case $p = 1$ we do not need to subtract any constant as the power of the monomial is $\alpha = 0$. Thus the condition becomes $T^*1 \in \text{LMO}_{\text{loc}}(\mathbb{R}^n)$, where T^* can be taken in the sense of duality of bmo and h^1 . In general, for b bounded, the pairing between bmo and h^1 can be given by integration, but for arbitrary $b \in \text{bmo}$, we need to subtract b_B in the integral in (5.3), unless we have $\int Tg = 0$. This is the case when T

is a convolution operator, as Tg inherits the cancellation from g (see also the remarks following Corollary 3.1).

Theorem 5.3. *Let $b \in \text{bmo}(\mathbb{R}^n)$. Suppose T is an inhomogeneous singular integral operator. If $T^*(1) \in \text{LMO}_{\text{loc}}(\mathbb{R}^n)$, and, for any ball B with $r(B) < 1$, $f := T_B^*(b)$ satisfies (5.4), then*

$$\|[b, T](a)\|_{h^1} \leq C(1 + \|b\|_{\text{bmo}})$$

for all h_b^1 atoms a .

Proof. Let a be an h_b^1 atom supported in a ball $B = B(x_0, r)$; as above, we may assume that when $r < 1$, $\int a = 0$. Again write $[b, T](a) = (b - c_B)T(a) - T(a(b - c_B))$.

As explained before the statement of the theorem, the assumption $T^*(1) \in \text{LMO}_{\text{loc}}(\mathbb{R}^n)$ allows us to apply [14, Theorem 5.3] get the boundedness of T on h^1 ; thus from (4.4) we get

$$\|T(a(b - c_B))\|_{h^1} \lesssim \|a(b - c_B)\|_{h^1} \lesssim \|b\|_{\text{bmo}}.$$

It remains to estimate the term $\|(b - c_B)T(a)\|_{h^1}$. We will show that $M = (b - c_B)T(a)$ is a multiple of an (s, λ) molecule (see Definition 2.3) associated to the ball $2B$, where we choose

$$s = \frac{3}{2}, \quad \lambda = \frac{n}{2} + \mu, \quad 0 < \mu < \frac{3}{2} \min(\delta, \varepsilon).$$

We first verify condition M1. Proceeding as in (4.3) and using the boundedness of T on L^2 , we have

$$\begin{aligned} \|(b - b_B)T(a)\|_{L^{\frac{3}{2}}(2B)} &\lesssim \|(b - b_B)\|_{L^6(2B)} \|T(a)\|_{L^2} \\ &\lesssim \|b - b_B\|_{L^6(2B)} \|a\|_{L^2} \\ &\lesssim \|b\|_{\text{bmo}, 6} r^{-\frac{3}{2}}. \end{aligned}$$

Next, we need to show condition M2: $\int_{(2B)^c} |M(x)|^{\frac{3}{2}} |x - x_0|^\lambda dx \lesssim r^\mu$. Suppose that $r < 1$. As in (5.1), using the cancellation on a and the smoothness of the kernel to bound $T(a)$, and (2.3), we have

$$\begin{aligned} &\int_{(2B)^c} |(b(x) - b_B)T(a)(x)|^{\frac{3}{2}} |x - x_0|^\lambda dx \\ &\lesssim \int_{(2B)^c} |b(x) - b_B|^{\frac{3}{2}} \frac{r^{\frac{3}{2}\delta}}{|x - x_0|^{\frac{3n}{2} + \frac{3\delta}{2}}} |x - x_0|^\lambda dx = r^{\frac{3}{2}\delta} \int_{(2B)^c} \frac{|b(x) - b_B|^{\frac{3}{2}}}{|x - x_0|^{n + \frac{3}{2}\delta - \mu}} dx \\ &\lesssim \|b\|_{\text{bmo}}^{\frac{3}{2}} r^\mu. \end{aligned}$$

Finally, condition (5.4) on $f = T_B^*(b)$ gives us

$$\begin{aligned} \left| \int_{\mathbb{R}^n} [b(x) - b_B]T(a)(x) dx \right| &= \left| \langle T_B^*(b), a \rangle \right| \leq \left(\int_B |f - f_B|^2 \right)^{1/2} \|a\|_{L^2(B)} \\ &\leq \log(1 + r(B)^{-1}). \end{aligned}$$

Now looking at the case $r \geq 1$, we can proceed as in (5.2) to write

$$\begin{aligned}
& \int_{(2B)^c} |(b(x) - c_B)T(a)(x)|^{\frac{3}{2}} |x - x_0|^\lambda dx \\
& \lesssim \int_{(2B)^c} \frac{|b(x) - c_B|^{\frac{3}{2}}}{|x - x_0|^{\frac{3}{2}n + \frac{3}{2}\varepsilon}} |x - x_0|^\lambda dx \\
& \leq r^{\frac{3}{2}\varepsilon} \int_{(2B)^c} \frac{|b(x) - c_B|^{\frac{3}{2}}}{|x - x_0|^{n + \frac{3}{2}\varepsilon - \mu}} dx \\
& \lesssim \|b\|_{\text{bmo}}^{\frac{3}{2}} r^\mu.
\end{aligned}$$

Here we have used (2.3) as well as the fact that $r \geq 1$, $\mu < \frac{3}{2}\varepsilon$. \square

Similarly to Corollary 5.1, when $h_b^1 = h^1$, the boundedness extends from one atom to all functions in the space.

Corollary 5.2. *Assuming the hypotheses of Theorem 5.3, if in addition $b \in \text{lmo}(\mathbb{R}^n)$, then $[b, T]$ is bounded from $h^1(\mathbb{R}^n)$ to $h^1(\mathbb{R}^n)$.*

References

- [1] Mujdat Agcayazi, Amiran Gogatishvili, Kerim Koca, and Rza Mustafayev, *A note on maximal commutators and commutators of maximal functions*, J. Math. Soc. Japan **67** (2015), no. 2, 581–593, DOI 10.2969/jmsj/06720581.
- [2] Jesús Bastero, Mario Milman, and Francisco J. Ruiz, *Commutators for the maximal and sharp functions*, Proc. Amer. Math. Soc. **128** (2000), no. 11, 3329–3334, DOI 10.1090/S0002-9939-00-05763-4.
- [3] Frédéric Bernicot, Tarek Elgindi, and Sahbi Keraani, *On the inviscid limit of the 2D Navier-Stokes equations with vorticity belonging to BMO-type spaces*, Ann. Inst. H. Poincaré C Anal. Non Linéaire **33** (2016), no. 2, 597–619, DOI 10.1016/j.anihpc.2014.12.001.
- [4] Frédéric Bernicot and Taoufik Hmidi, *On the global well-posedness for Euler equations with unbounded vorticity*, Dyn. Partial Differ. Equ. **12** (2015), no. 2, 127–155, DOI 10.4310/DPDE.2015.v12.n2.a3.
- [5] Aline Bonami and Justin Feuto, *Products of functions in Hardy and Lipschitz or BMO spaces*, Recent developments in real and harmonic analysis, Appl. Numer. Harmon. Anal., Birkhäuser Boston, Boston, MA, 2010, pp. 57–71, DOI 10.1007/978-0-8176-4588-5_4. MR2603175
- [6] Gérard Bourdaud, *Remarques sur certains sous-espaces de $BMO(\mathbb{R}^n)$ et de $\text{bmo}(\mathbb{R}^n)$* , Ann. Inst. Fourier (Grenoble) **52** (2002), no. 4, 1187–1218.
- [7] Marcin Bownik, *Boundedness of operators on Hardy spaces via atomic decompositions*, Proc. Amer. Math. Soc. **133** (2005), no. 12, 3535–3542, DOI 10.1090/S0002-9939-05-07892-5.
- [8] Huy Qui Bui, *Weighted Hardy spaces*, Math. Nachr. **103** (1981), 45–62, DOI 10.1002/mana.19811030106.
- [9] Der-Chen Chang and Song-Ying Li, *On the boundedness of multipliers, commutators and the second derivatives of Green’s operators on H^1 and BMO*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **28** (1999), no. 2, 341–356.
- [10] Ronald R. Coifman, Richard Rochberg, and Guido Weiss, *Factorization theorems for Hardy spaces in several variables*, Ann. of Math. (2) **103** (1976), no. 3, 611–635, DOI 10.2307/1970954. MR412721
- [11] Ronald R. Coifman and Guido Weiss, *Extensions of Hardy spaces and their use in analysis*, Bull. Amer. Math. Soc. **83** (1977), no. 4, 569–645, DOI 10.1090/S0002-9904-1977-14325-5.

- [12] Galia Dafni, *Hardy spaces on strongly pseudoconvex domains in \mathbb{C}^n and domains of finite type in \mathbb{C}^2* , 1993. Thesis (Ph.D.)–Princeton University.
- [13] ———, *Local VMO and weak convergence in h^1* , *Canad. Math. Bull.* **45** (2002), no. 1, 46–59, DOI 10.4153/CMB-2002-005-2.
- [14] Galia Dafni, Chun Ho Lau, Tiago Picon, and Claudio Vasconcelos, *Inhomogeneous cancellation conditions and Calderón-Zygmund type operators on h^p* , *Nonlinear Anal.* **225** (2022), Paper No. 113110, 22, DOI 10.1016/j.na.2022.113110.
- [15] ———, *Necessary cancellation conditions for the boundedness of operators on local Hardy spaces*, posted on 2022, DOI 10.48550/ARXIV.2210.05786.
- [16] Galia Dafni and Elijah Lifyand, *A local Hilbert transform, Hardy’s inequality and molecular characterization of Goldberg’s local Hardy space*, *Complex Anal. Synerg.* **5** (2019), no. 1, Paper No. 10, 9, DOI 10.1007/s40627-019-0036-2.
- [17] Galia Dafni, Ethan Mombourquette, and Hong Yue, *Some characterizations of local bmo and h^1 on spaces of homogeneous type*, *Harmonic analysis and nonlinear partial differential equations*, 2016, pp. 187–215.
- [18] Galia Dafni and Hong Yue, *Some characterizations of local bmo and h^1 on metric measure spaces*, *Anal. Math. Phys.* **2** (2012), no. 3, 285–318, DOI 10.1007/s13324-012-0034-5.
- [19] Wei Ding, YongSheng Han, and YuePing Zhu, *Boundedness of singular integral operators on local Hardy spaces and dual spaces*, *Potential Anal.* **55** (2021), no. 3, 419–441, DOI 10.1007/s11118-020-09863-3.
- [20] Charles Fefferman and Elias M. Stein, *H^p spaces of several variables*, *Acta Math.* **129** (1972), no. 3-4, 137–193, DOI 10.1007/BF02392215.
- [21] Xing Fu, Dachun Yang, and Sibe Yang, *Endpoint boundedness of linear commutators on local Hardy spaces over metric measure spaces of homogeneous type*, *J. Geom. Anal.* **31** (2021), no. 4, 4092–4164, DOI 10.1007/s12220-020-00429-8. MR4236554
- [22] José García-Cuerva, Eleonor O. Harboure, Carlos Segovia, and José Luis Torrea, *Weighted norm inequalities for commutators of strongly singular integrals*, *Indiana Univ. Math. J.* **40** (1991), no. 4, 1397–1420, DOI 10.1512/iumj.1991.40.40063.
- [23] David Goldberg, *Local Hardy spaces*, *Harmonic analysis in Euclidean spaces* (Proc. Sympos. Pure Math., Williams Coll., Williamstown, Mass., 1978), Proc. Sympos. Pure Math., XXXV, Part, Amer. Math. Soc., Providence, R.I., 1979, pp. 245–248. MR545262
- [24] ———, *A local version of real Hardy spaces*, *Duke Math. J.* **46** (1979), no. 1, 27–42. MR523600
- [25] Loukas Grafakos, *Modern Fourier analysis*, 3rd ed., Graduate Texts in Mathematics, vol. 250, Springer, New York, 2014. MR3243741
- [26] Eleonor Harboure, Carlos Segovia, and José L. Torrea, *Boundedness of commutators of fractional and singular integrals for the extreme values of p* , *Illinois J. Math.* **41** (1997), no. 4, 676–700. MR1468874
- [27] Ziyi He, Dachun Yang, and Wen Yuan, *Real-variable characterizations of local Hardy spaces on spaces of homogeneous type*, *Math. Nachr.* **294** (2021), no. 5, 900–955, DOI 10.1002/mana.201900320.
- [28] Ha Duy Hung and Luong Dang Ky, *An Hardy estimate for commutators of pseudo-differential operators*, *Taiwanese J. Math.* **19** (2015), no. 4, 1097–1109, DOI 10.11650/tjm.19.2015.5003. MR3384681
- [29] Duong Quoc Huy and Luong Dang Ky, *Weighted Hardy space estimates for commutators of Calderón-Zygmund operators*, *Vietnam J. Math.* **49** (2021), no. 4, 1065–1077, DOI 10.1007/s10013-020-00406-2.
- [30] Tuomas P. Hytönen, *The L^p -to- L^q boundedness of commutators with applications to the Jacobian operator*, *J. Math. Pures Appl.* (9) **156** (2021), 351–391, DOI 10.1016/j.matpur.2021.10.007 (English, with English and French summaries). MR4338459
- [31] Svante Janson, *Mean oscillation and commutators of singular integral operators*, *Ark. Mat.* **16** (1978), no. 2, 263–270, DOI 10.1007/BF02386000. MR524754
- [32] F. John and L. Nirenberg, *On functions of bounded mean oscillation*, *Comm. Pure Appl. Math.* **14** (1961), 415–426, DOI 10.1002/cpa.3160140317.

- [33] Yasuo Komori, *Calderón-Zygmund operators on $H^p(\mathbf{R}^n)$* , Sci. Math. Jpn. **53** (2001), no. 1, 65–73.
- [34] Luong Dang Ky, *Bilinear decompositions and commutators of singular integral operators*, Trans. Amer. Math. Soc. **365** (2013), no. 6, 2931–2958, DOI 10.1090/S0002-9947-2012-05727-8. MR3034454
- [35] Song-Ying Li, *Toeplitz operators on Hardy space $H^p(S)$ with $0 < p \leq 1$* , Integral Equations Operator Theory **15** (1992), no. 5, 807–824, DOI 10.1007/BF01200702.
- [36] Stefano Meda, Peter Sjögren, and Maria Vallarino, *On the H^1-L^1 boundedness of operators*, Proc. Amer. Math. Soc. **136** (2008), no. 8, 2921–2931, DOI 10.1090/S0002-9939-08-09365-9.
- [37] Mario Milman and Tomas Schonbek, *Second order estimates in interpolation theory and applications*, Proc. Amer. Math. Soc. **110** (1990), no. 4, 961–969, DOI 10.2307/2047743.
- [38] Marco M. Peloso and Silvia Secco, *Local Riesz transforms characterization of local Hardy spaces*, Collect. Math. **59** (2008), no. 3, 299–320, DOI 10.1007/BF03191189.
- [39] Carlos Pérez, *Endpoint estimates for commutators of singular integral operators*, J. Funct. Anal. **128** (1995), no. 1, 163–185, DOI 10.1006/jfan.1995.1027. MR1317714
- [40] Donald Sarason, *Functions of vanishing mean oscillation*, Trans. Amer. Math. Soc. **207** (1975), 391–405, DOI 10.2307/1997184.
- [41] ———, *Function theory on the unit circle*, Virginia Polytechnic Institute and State University, Department of Mathematics, Blacksburg, Va., 1978. Notes for lectures given at a Conference at Virginia Polytechnic Institute and State University, Blacksburg, Va., June 19–23, 1978.
- [42] Carlos Segovia and José L. Torrea, *Weighted inequalities for commutators of fractional and singular integrals*, Publ. Mat. **35** (1991), no. 1, 209–235, DOI 10.5565/PUBLMAT.35191.09. Conference on Mathematical Analysis (El Escorial, 1989).
- [43] Xian Liang Shi and Alberto Torchinsky, *Functions of vanishing mean oscillation*, Math. Nachr. **133** (1987), 289–296, DOI 10.1002/mana.19871330119.
- [44] David A. Stegenga, *Bounded Toeplitz operators on H^1 and applications of the duality between H^1 and the functions of bounded mean oscillation*, Amer. J. Math. **98** (1976), no. 3, 573–589, DOI 10.2307/2373807.
- [45] Elias M. Stein, *Singular Integrals and Differentiability Properties of Functions (PMS-30)*, Princeton University Press, 1970.
- [46] ———, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series, vol. 43, Princeton University Press, Princeton, NJ, 1993. With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III.
- [47] Lin Tang, *Interior h^1 estimates for parabolic equations with LMO coefficients*, Canad. J. Math. **62** (2010), no. 1, 202–217, DOI 10.4153/CJM-2010-011-1.
- [48] ———, *Weighted local Hardy spaces and their applications*, Illinois J. Math. **56** (2012), no. 2, 453–495.
- [49] Akihito Uchiyama, *On the compactness of operators of Hankel type*, Tohoku Math. J. (2) **30** (1978), no. 1, 163–171, DOI 10.2748/tmj/1178230105. MR467384

G. Dafni, CONCORDIA UNIVERSITY, DEPARTMENT OF MATHEMATICS AND STATISTICS, MONTRÉAL, QUÉBEC, H3G-1M8, CANADA

C. H. Lau, CONCORDIA UNIVERSITY, DEPARTMENT OF MATHEMATICS AND STATISTICS, MONTRÉAL, QUÉBEC, H3G-1M8, CANADA