# ON THE REAL-ROOTEDNESS OF THE EULERIAN TRANSFORMATION

CHRISTOS A. ATHANASIADIS

ABSTRACT. The Eulerian transformation is the linear operator on polynomials in one variable with real coefficients which maps the powers of this variable to the corresponding Eulerian polynomials. The derangement transformation is defined similarly. Brändén and Jochemko have conjectured that the Eulerian transforms of a class of polynomials with nonnegative coefficients, which includes those having all their roots in the interval [-1,0], have only real zeros. This conjecture is proven in this paper. More general transformations are introduced in the combinatorial-geometric context of uniform triangulations of simplicial complexes, where Eulerian and derangement transformations arise in the special case of barycentric subdivision, and are shown to have strong unimodality and gamma-positivity properties. General real-rootedness conjectures for these transformations, which unify various results and conjectures in the literature, are also proposed.

## 1. INTRODUCTION

Eulerian polynomials form one of the most important and well studied families of polynomials in mathematics, playing a prominent role in combinatorics and elsewhere; see, for instance, [21, 31] [38, Section 1.4]. The *n*th Eulerian polynomial is defined as

$$A_n(x) = \sum_{w \in \mathfrak{S}_n} x^{\operatorname{des}(w)} = \sum_{w \in \mathfrak{S}_n} x^{\operatorname{exc}(w)},$$

where  $\mathfrak{S}_n$  is the symmetric group of permutations of the set  $[n] := \{1, 2, \dots, n\}$  and

$$des(w) = \#\{i \in [n-1] : w(i) > w(i+1)\}$$
$$exc(w) = \#\{i \in [n-1] : w(i) > i\}$$

is the descent and excedance number of  $w \in \mathfrak{S}_n$ , respectively, with the convention  $A_0(x) := 1$ . A well known result, often attributed to Frobenius [24], states that  $A_n(x)$  has only real roots. Far less is known about the real-rootedness of linear combinations of Eulerian polynomials. Following the historical approach to study linear transformations which preserve real-rootedness properties (see the discussions in [14, Section 7.7] [15, Section 1]), Brändén and Jochemko [15] considered the linear operator  $\mathcal{A}^\circ : \mathbb{R}[x] \to \mathbb{R}[x]$ , defined by setting

(1) 
$$\mathcal{A}^{\circ}(x^{n}) = \begin{cases} 1, & \text{if } n = 0\\ xA_{n}(x), & \text{if } n \ge 1 \end{cases}$$

Date: May 16, 2023.

Research supported by the Hellenic Foundation for Research and Innovation (H.F.R.I.) under the '2nd Call for H.F.R.I. Research Projects to support Faculty Members & Researchers' (Project Number: HFRI-FM20-04537).

Key words and phrases. Eulerian polynomial, real-rooted polynomial, barycentric subdivision, triangulation, *h*-polynomial, local *h*-polynomial, gamma-positivity.

for  $n \in \mathbb{N} := \{0, 1, 2, ...\}$ , and named it the Eulerian transformation (the slight difference from [15] in the notation adopted here will be explained by our discussion in the sequel). They disproved a conjecture of Brenti [17, p. 32], stating that  $\mathcal{A}^{\circ}$  preserves the class of polynomials having only real and nonpositive roots, and conjectured [15, Conjecture 1] that  $\mathcal{A}^{\circ}(p(x))$  has only real roots for every polynomial p(x) which can be written as a nonnegative linear combination of the polynomials  $x^{n-k}(1+x)^k$  for  $k \in \{0, 1, ..., n\}$  for some  $n \in \mathbb{N}$ . This class, which we will denote by  $\mathcal{P}_n[x]$  throughout this paper, contains all polynomials of degree n with nonnegative coefficients which have all their roots in the interval [-1, 0]. To support their conjecture, Brändén and Jochemko showed [15, Theorem 3.2] that  $\mathcal{A}^{\circ}(p(x))$  has a unimodal symmetric decomposition with respect to n for every  $p(x) \in \mathcal{P}_n[x]$ ; in particular,  $\mathcal{A}^{\circ}(p(x))$  is unimodal, with a peak at position  $\lceil n/2 \rceil$  (undefined terminology on polynomials is explained in Section 2). They also studied other subtle properties of the Eulerian transformation.

This paper aims to prove [15, Conjecture 1], provide new combinatorial and geometric interpretations of the Eulerian transformation and place the latter in a much broader and natural combinatorial-geometric context. Our first main result proves the aforementioned conjecture of [15] and strengthens [15, Theorem 3.2]. The polynomial  $\mathcal{A}^{\circ}((1+x)^n)$  which appears in the statement is the *n*th binomial Eulerian polynomial; see Section 3 for more information.

**Theorem 1.1.** For every  $p(x) \in \mathcal{P}_n[x]$ :

- (a) The polynomial  $\mathcal{A}^{\circ}(p(x))$  has only real roots. Moreover, it interlaces  $xA_n(x)$  and it is interlaced by  $\mathcal{A}^{\circ}((1+x)^n)$ .
- (b) The polynomial  $\mathcal{A}^{\circ}(p(x))$  has a real-rooted and interlacing (in particular, unimodal and  $\gamma$ -positive) symmetric decomposition with respect to n.

For a simple nontrivial application to linear combinations of  $A_n(x)$ ,  $A_{n-1}(x)$  and  $A_{n-2}(x)$ , see Example 3.3.

The unimodality,  $\gamma$ -positivity and real-rootedness properties of Theorem 1.1 do not apply exclusively to the Eulerian transformation. We propose one way to generalize these results as follows. Our motivation comes from the fact that the right-hand side of Equation (1) can be interpreted as the interior *h*-polynomial of the barycentric subdivision of the (n-1)-dimensional simplex, denoted here by  $\sigma_n$  (basic definitions and terminology on the face enumeration of simplicial complexes are explained in Section 5). It seems natural to inquire about the behavior of the transformation obtained when barycentric subdivision is replaced by other types of triangulations.

A suitable context is provided by the theory of uniform triangulations, developed in [7]. A triangulation  $\Gamma$  of the simplex  $\sigma_n$  is called *uniform* if the *f*-vector of the restriction of  $\Gamma$  to a face of  $\sigma_n$  depends only on the dimension of that face. One can then consider the *h*-polynomial  $h_{\mathcal{F}}(\sigma_m, x)$  and the interior *h*-polynomial  $h^{\circ}_{\mathcal{F}}(\sigma_m, x)$  of the restriction of  $\Gamma$  to any (m-1)-dimensional face of  $\sigma_n$  (where  $\mathcal{F}$  records some combinatorial data determined by  $\Gamma$ ). We define the linear operator  $\mathcal{H}^{\circ}_{\mathcal{F}} : \mathbb{R}_n[x] \to \mathbb{R}_n[x]$  by setting

$$\mathcal{H}^{\circ}_{\mathcal{F}}(x^m) = h^{\circ}_{\mathcal{F}}(\sigma_m, x)$$

for  $m \in \{0, 1, ..., n\}$ , where  $\mathbb{R}_n[x]$  stands for the space of polynomials of degree at most nwith real coefficients. This map reduces to the restriction of the Eulerian transformation  $\mathcal{A}^\circ$  to  $\mathbb{R}_n[x]$  in the special case that  $\Gamma$  is the barycentric subdivision of  $\sigma_n$ . Following [8], we say that a triangulation  $\Gamma$  of the simplex  $\sigma_n = 2^V$  is theta unimodal (respectively, theta  $\gamma$ -positive) if  $\theta(\Gamma_F, x) := h(\Gamma_F, x) - h(\partial \Gamma_F, x)$  is unimodal (respectively,  $\gamma$ -positive) for every  $F \subseteq V$ , where  $\Gamma_F$  is the restriction of  $\Gamma$  to the face F of  $\sigma_n$ . For uniform triangulations, this means that

$$\theta_{\mathcal{F}}(\sigma_m, x) := h_{\mathcal{F}}(\sigma_m, x) - h_{\mathcal{F}}(\partial \sigma_m, x)$$

is unimodal (respectively,  $\gamma$ -positive) for every  $m \in \{0, 1, \ldots, n\}$ , where  $h_{\mathcal{F}}(\partial \sigma_m, x)$  stands for the *h*-polynomial of the restriction of  $\Gamma$  to the boundary complex of any (m-1)-dimensional face of  $\sigma_n$ . As discussed in [8, Section 5] (see [8, Theorem 5.1]), it follows from [3, Theorem 50] and the *g*-theorem for triangulations of spheres [1, 2, 30] that  $\Gamma$  is theta unimodal whenever  $\partial \Gamma_F$  is a vertex-induced subcomplex of  $\Gamma_F$  for every  $F \subseteq V$ . Moreover, several classes of triangulations of the simplex with interesting enumerative combinatorics are known to be theta  $\gamma$ -positive (see Section 6). For the barycentric subdivision  $\theta_{\mathcal{F}}(\sigma_m, x)$  is identically zero, and hence trivially  $\gamma$ -positive, for  $m \geq 1$ .

Our second main result significantly generalizes [15, Theorem 3.2] and supports a more general conjecture than [15, Conjecture 1] which can be stated in this context (see Conjecture 7.1).

**Theorem 1.2.** Let  $\Gamma$  be a uniform triangulation of the (n-1)-dimensional simplex.

- (a) If  $\Gamma$  is theta unimodal, then  $\mathcal{H}^{\circ}_{\mathcal{F}}(p(x))$  has a unimodal symmetric decomposition with respect to n for every  $p(x) \in \mathcal{P}_n[x]$ .
- (b) If  $\Gamma$  is theta  $\gamma$ -positive, then  $\mathcal{H}^{\circ}_{\mathcal{F}}(p(x))$  has a  $\gamma$ -positive symmetric decomposition with respect to n for every  $p(x) \in \mathcal{P}_n[x]$ .

The structure and other results of this paper can be described as follows. Section 2 provides basic definitions and background on real polynomials and their roots. Theorem 1.1 is proven in Section 3, after the combinatorics of the polynomials  $\mathcal{A}^{\circ}(x^{n-k}(1+x)^k)$  is sufficiently developed. The method extends that followed by Brändén and Jochemko in [15, Section 4] to obtain various partial results (although this is only one out of several possible approaches one may follow to attack [15, Conjecture 1]). Section 4 is concerned with a close relative of the Eulerian transformation, introduced and studied by Brändén and Solus [16, Section 3.2], termed here as the derangement transformation. Proposition 4.1 provides a common generalization of the gammapositivity of Eulerian and derangement polynomials; a result which easily implies Theorem 1.2 in the crucial special case of the barycentric subdivision. An analogue of Theorem 1.1 (b) for the derangement transformation (Corollary 4.2) is also derived in Section 4 from Proposition 4.1 and results of [16, Section 3.2].

Section 5 reviews basic enumerative combinatorics of triangulations (and uniform triangulations, in particular) of simplicial complexes. Given a uniform triangulation of the simplex  $\sigma_n$ , it introduces linear transformations  $\mathcal{H}^{\circ}_{\mathcal{F}}$  and  $\mathcal{L}_{\mathcal{F}}$  which reduce to the Eulerian and derangement transformations, respectively, in the special case of barycentric subdivision and provides a geometric interpretation of the polynomials  $\mathcal{H}^{\circ}_{\mathcal{F}}(x^{n-k}(1+x)^k)$  (see Proposition 5.2). Theorem 1.2 and its analogue for  $\mathcal{L}_{\mathcal{F}}$  (Corollary 6.6) is proven in Section 6. Essential ingredients of the proof are a family of polynomials interpolating between the *h*-polynomial and the local *h*-polynomial of a triangulation of a simplex (see Definition 6.1), properties of theta polynomials established in [8] and Proposition 4.1. Section 7 proposes generalizations of [15, Conjecture 1] for the transformations  $\mathcal{H}^{\circ}_{\mathcal{F}}$  and  $\mathcal{L}_{\mathcal{F}}$  and discusses supporting evidence and some further directions.

# 2. Recollections of polynomials and their roots

This section explains basic background and terminology on real polynomials which will be useful in the sequel. We recall that  $\mathbb{R}_n[x]$  stands for the space of polynomials  $p(x) \in \mathbb{R}[x]$  of degree at most n. We will denote by  $\mathcal{I}_n(p(x))$  the reciprocal  $x^n p(1/x)$  with respect to n of a polynomial  $p(x) \in \mathbb{R}_n[x]$ . A polynomial  $p(x) = a_0 + a_1x + \cdots + a_nx^n \in \mathbb{R}_n[x]$  is called

- symmetric, with center of symmetry n/2, if  $a_i = a_{n-i}$  for all  $0 \le i \le n$ ,
- unimodal, with a peak at position k, if  $0 \le a_0 \le a_1 \le \cdots \le a_k \ge a_{k+1} \ge \cdots \ge a_n \ge 0$ ,  $\gamma$ -positive, with center of symmetry n/2, if  $p(x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \gamma_i x^i (1+x)^{n-2i}$  for some nonnegative real numebrs  $\gamma_0, \gamma_1, \ldots, \gamma_{\lfloor n/2 \rfloor}$ ,
- real-rooted, if every root of p(x) is real, or  $p(x) \equiv 0$ .

Every  $\gamma$ -positive polynomial is symmetric and unimodal and every real-rooted and symmetric polynomial with nonnegative coefficients is  $\gamma$ -positive; see [5, 14, 25, 35] [31, Chapter 4] for more information on the connections among these concepts. Note that unimodal polynomials are assumed to have nonnegative coefficients in this paper.

A real-rooted polynomial p(x), with roots  $\alpha_1 \geq \alpha_2 \geq \cdots$ , is said to *interlace* a real-rooted polynomial q(x), with roots  $\beta_1 \ge \beta_2 \ge \cdots$ , if

$$\cdots \leq \alpha_2 \leq \beta_2 \leq \alpha_1 \leq \beta_1.$$

We then write  $p(x) \leq q(x)$ . By convention, the zero polynomial interlaces and is interlaced by every real-rooted polynomial and nonzero constant polynomials strictly interlace all polynomials of degree at most one. For real-rooted polynomials  $p(x), q(x) \in \mathbb{R}_n[x]$  with nonnegative coefficients,  $p(x) \preceq q(x) \Rightarrow q(x) \preceq xp(x)$  and  $p(x) \preceq q(x) \Rightarrow \mathcal{I}_n(q(x)) \preceq \mathcal{I}_n(p(x))$ . If two or more real-rooted polynomials with positive leading coefficients interlace (respectively, are interlaced by) a real-rooted polynomial p(x), then so does their sum. We will use these properties of interlacing in Section 3; standard references are [14, Section 7.8] [20].

A sequence  $(p_0(x), p_1(x), \ldots, p_m(x))$  of real-rooted polynomials is called *interlacing* if  $p_i(x) \leq 1$  $p_i(x)$  for  $0 \le i < j \le m$ . The following lemmas will be applied in Section 3. Part (b) of Lemma 2.2 is a special case of [27, Theorem 2.4] (another version appears in the proof of [15,Theorem 4.4]).

**Lemma 2.1.** ([39, Proposition 3.3]) Let  $p_0(x), p_1(x), \ldots, p_m(x)$  be nonzero real-rooted polynomials. If  $p_{i-1}(x) \leq p_i(x)$  for every  $i \in [m]$  and  $p_0(x) \leq p_m(x)$ , then  $(p_0(x), p_1(x), \ldots, p_m(x))$  is an interlacing sequence.

**Lemma 2.2.** Let  $(p_0(x), p_1(x), \ldots, p_m(x))$  be an interlacing sequence of real-rooted polynomials with positive leading coefficients.

- (a) Every nonnegative linear combination p(x) of  $p_0(x), p_1(x), \ldots, p_m(x)$  is real-rooted. Moroever,  $p_0(x) \leq p(x) \leq p_m(x)$ .
- (b) (cf. [27, Theorem 2.4]) The sequence  $(q_0(x), q_1(x), \ldots, q_{m+1}(x))$  defined by

$$q_0(x) = \alpha(x)p_0(x) + \sum_{i=1}^m p_i(x),$$

where  $\alpha(x) = 1$  or  $\alpha(x) = 1 + x$ , and

$$q_k(x) = x \sum_{i=0}^{k-1} p_i(x) + \sum_{i=k}^m p_i(x)$$

for  $k \in \{1, 2, \dots, m+1\}$  is also interlacing.

#### Example 2.3. Let

(2) 
$$p_{n,k}(x) = \sum_{w \in \mathfrak{S}_{n+1}: w(1)=k+1} x^{\operatorname{des}(w)} = \sum_{w \in \mathfrak{S}_{n+1}: w(n+1)=k+1} x^{\operatorname{asc}(w)}$$

(3) 
$$= \sum_{w \in \mathfrak{S}_{n+1}: w^{-1}(1) = k+1} x^{\operatorname{exc}(w)}$$

for  $k \in \{0, 1, ..., n\}$ , so that  $p_{n,0}(x) = A_n(x)$  and  $p_{n,n}(x) = xA_n(x)$ . The polynomials  $p_{n,k}(x)$  appeared in the work of Brenti and Welker [18] on *f*-vectors of barycentric subdivisions and, independently, in [19, Section 2.2], and have been studied intensely since then; see [8, Section 2] [12, Section 3] [14, Example 7.8.8] and the references given there. They satisfy the recurrence

(4) 
$$p_{n,k}(x) = x \sum_{i=0}^{k-1} p_{n-1,i}(x) + \sum_{i=k}^{n-1} p_{n-1,i}(x)$$

for  $k \in \{0, 1, ..., n\}$ . Hence, by Lemma 2.2 (b),  $(p_{n,k}(x))_{0 \le k \le n}$  is an interlacing sequence of real-rooted polynomials for every  $n \in \mathbb{N}$ .

Given  $p(x) \in \mathbb{R}_n[x]$ , there exist unique symmetric polynomials  $a(x) \in \mathbb{R}_n[x]$  and  $b(x) \in \mathbb{R}_{n-1}[x]$  with centers of symmetry n/2 and (n-1)/2, respectively, such that p(x) = a(x) + xb(x). This expression is known as the symmetric decomposition (or Stapledon decomposition) of p(x) with respect to n. Then, p(x) is said to have a nonnegative (respectively, unimodal,  $\gamma$ -positive or real-rooted) symmetric decomposition with respect to n if a(x) and b(x) have nonnegative coefficients (respectively, are unimodal,  $\gamma$ -positive or real-rooted); see [5, Section 5] [11, 16] for more information about these concepts. A nonnegative, real-rooted symmetric decomposition p(x) = a(x) + xb(x) is said to be interlacing if  $b(x) \leq a(x)$ . Then [16, Theorem 2.7], p(x) is real-rooted and is interlaced by each one of a(x), b(x) and  $\mathcal{I}_n(p(x))$ . Moreover,  $b(x) \leq a(x) \leq a(x) \leq p(x) \Leftrightarrow b(x) \leq p(x) \Leftrightarrow \mathcal{I}_n(p(x)) \leq p(x)$ , under the assumption that p(x) has a nonnegative symmetric decomposition with respect to n.

## 3. Combinatorics of the Eulerian transformation

This section studies the combinatorics of the Eulerian transformation and, especially, the polynomials  $\mathcal{A}^{\circ}(x^{n-k}(1+x)^k)$  for  $k \in \{0, 1, ..., n\}$ , and proves Theorem 1.1. For convenience, we consider the polynomials  $q_{n,k}(x)$  defined by

(5) 
$$\mathcal{I}_n(q_{n,k}(x)) = \mathcal{A}^\circ \left( x^{n-k} (1+x)^k \right)$$

for  $k \in \{0, 1, ..., n\}$  instead. They are shown on Table 1 for  $n \leq 4$ . We have  $q_{n,0}(x) = A_n(x)$ and  $q_{n,n}(x) = \widetilde{A}_n(x)$  for every  $n \in \mathbb{N}$ , where

$$\widetilde{A}_{n}(x) := \mathcal{A}^{\circ}((1+x)^{n}) = 1 + x \sum_{i=1}^{n} \binom{n}{i} A_{i}(x) = \sum_{i=0}^{n} \binom{n}{i} x^{n-i} A_{i}(x)$$

is the *n*th binomial Eulerian polynomial. This polynomial was shown to be symmetric and  $\gamma$ -positive, with center of symmetry n/2, in [32, Section 11] (see also [5, Section 2.1] [15, Section 3] [33]) and real-rooted in [27, Section 3] (see also [15, Section 4]). Moreover, as a special case of [15, Theorem 4.5],  $A_n(x)$  interlaces  $\widetilde{A}_n(x)$ . We will show that each  $q_{n,k}(x)$  is real-rooted and that the sequence  $(q_{n,k}(x))_{0 \le k \le n}$  is interlacing for every  $n \in \mathbb{N}$ .

CHRISTOS A. ATHANASIADIS

	k = 0	k = 1	k = 2	k = 3
n = 0	1			
n = 1	1	1+x		
n=2	1+x	1 + 2x	$1 + 3x + x^2$	
n = 3	$1 + 4x + x^2$	$1 + 5x + 2x^2$	$1 + 6x + 4x^2$	$1 + 7x + 7x^2 + x^3$
n = 4	$1 + 11x + 11x^2 + x^3$	$1 + 12x + 15x^2 + 2x^3$	$1 + 13x + 20x^2 + 4x^3$	$1 + 14x + 26x^2 + 8x^3$




TABLE 1. The polynomials  $q_{n,k}(x)$  for  $n \leq 4$ .

The following statement lists some recurrences and combinatorial formulas for the polynomials  $q_{n,k}(x)$ ; a geometric interpretation will be given in Section 5.

**Proposition 3.1.** (a) The polynomials  $q_{n,k}(x)$  satisfy the recurrence

$$q_{n,k+1}(x) = q_{n,k}(x) + xq_{n-1,k}(x)$$

for  $0 \le k < n$ . (b) We have

(6) 
$$q_{n,k}(x) = \sum_{i=0}^{k} \binom{k}{i} x^i A_{n-i}(x)$$

(7) 
$$= \sum_{w \in \mathfrak{S}_n} (1+x)^{\mathrm{fix}_k(w)} x^{\mathrm{exc}(w)}$$

(8) 
$$=\sum_{i=0}^{k} \binom{k}{i} p_{n-i,k-i}(x)$$

for  $0 \le k \le n$ , where  $fix_k(w)$  is the number of fixed points of  $w \in \mathfrak{S}_n$  not exceeding k and  $p_{n,k}(x)$  are the polynomials of Example 2.3.

*Proof.* For part (a) we note that  $x^{n-k-1}(1+x)^{k+1} = x^{n-k}(1+x)^k + x^{n-k-1}(1+x)^k$ , which implies that

$$x^{n}q_{n,k+1}(1/x) = \mathcal{A}^{\circ}\left(x^{n-k-1}(1+x)^{k+1}\right) = \mathcal{A}^{\circ}\left(x^{n-k}(1+x)^{k}\right) + \mathcal{A}^{\circ}\left(x^{n-k-1}(1+x)^{k}\right)$$
$$= x^{n}q_{n,k}(1/x) + x^{n-1}q_{n-1,k}(1/x).$$

Equation (6) follows from the recurrence of part (a) by induction on k, or by expanding the binomial  $(1 + x)^k$  in the defining Equation (5) for  $q_{n,k}(x)$  and applying the linearity of  $\mathcal{A}^\circ$ .

To derive Equation (7) from (6) we consider (as in [15, Section 3]) decorated permutations of [n], meaning that some of their fixed points (possibly all or none) are colored black. We denote by  $\mathfrak{S}_n^*$  the set of decorated permutations of [n] and for  $w \in \mathfrak{S}_n^*$  we denote by  $\operatorname{fix}^*(w)$  and  $\operatorname{Fix}^*(w)$ 

the number and the set of fixed points of w which are colored black, respectively, and by exc(w) the excedance number of the (undecorated) permutation which corresponds to w. Then,

$$\sum_{w \in \mathfrak{S}_n} (1+x)^{\operatorname{fix}_k(w)} x^{\operatorname{exc}(w)} = \sum_{w \in \mathfrak{S}_n^*: \operatorname{Fix}^*(w) \subseteq [k]} x^{\operatorname{fix}^*(w) + \operatorname{exc}(w)}$$
$$= \sum_{i=0}^k \binom{k}{i} x^i \sum_{u \in \mathfrak{S}_{n-i}} x^{\operatorname{exc}(u)} = \sum_{i=0}^k \binom{k}{i} x^i A_{n-i}(x)$$

To derive Equation (8) from Equation (6), we expand  $x^i = \sum_{j=0}^{i} {i \choose j} (x-1)^{i-j}$  on the righthand side of (6), apply the identity  ${k \choose i} {i \choose j} = {k \choose j} {k-j \choose i-j}$ , change the order of summation and use the formula

$$p_{n,k}(x) = \sum_{i=0}^{k} \binom{k}{i} (x-1)^{i} A_{n-i}(x),$$

which is easily derived from the identity [18, Equation (4)]

$$\sum_{m \ge 0} m^k (1+m)^{n-k} x^m = \frac{p_{n,k}(x)}{(1-x)^{n+1}}.$$

We omit the details, which are straightforward.

To prove the real-rootedness of  $q_{n,k}(x)$ , we introduce the polynomials

(9) 
$$q_{n,k,j}(x) = \sum_{w \in \mathfrak{S}_{n+1}: w^{-1}(1) = j+1} (1+x)^{\text{fix}_k(w)} x^{\text{exc}(w)}$$

for  $k \in \{0, 1, ..., n + 1\}$  and  $j \in \{0, 1, ..., n\}$ , where fix<sub>k</sub>(w) is as in Proposition 3.1. An application of the fundamental transformation [21, Section I.3] [38, Section 1.3], obtained when each cycle of a permutation  $w \in \mathfrak{S}_{n+1}$  is written with its smallest element last and cylces are arranged in increasing order of their smallest element, yields that

(10) 
$$q_{n,k,j}(x) = \sum_{w \in \mathfrak{S}_{n+1}: w(1)=j+1} (1+x)^{\operatorname{bad}_k(w)} x^{\operatorname{des}(w)}$$

(11) 
$$q_{n,k}(x) = \sum_{w \in \mathfrak{S}_n} (1+x)^{\operatorname{bad}_k(w)} x^{\operatorname{des}(w)}$$

where, for  $w \in \mathfrak{S}_n$ ,  $\operatorname{bad}_k(w)$  is the number of indices  $i \in [n]$  for which  $w(i) \leq k$  is a right-to-left minimum of w (meaning that  $w(i) \leq w(j)$  for  $i \leq j \leq n$ ) and either i = 1 or w(i-1) < w(i). We also set

(12) 
$$q_{n,k,j}^*(x) = \begin{cases} \frac{q_{n,k,j}(x)}{1+x}, & \text{if } j = 0 \text{ and } k \ge 1\\ q_{n,k,j}(x), & \text{otherwise} \end{cases}$$

for  $k \in \{0, 1, \dots, n+1\}$  and  $j \in \{0, 1, \dots, n\}$ .

**Proposition 3.2.** The following formulas hold for the  $q_{n,k,j}(x)$  and  $q_{n,k,j}^*(x)$ :

(a)  $q_{n,k,0}^*(x) = q_{n,k-1}(x)$  for  $k \in \{1, 2, \dots, n+1\}$ .

n-1

(b)

$$q_{n,k}(x) = \sum_{j=0} q_{n-1,k,j}(x) = \sum_{j=0} q_{n-1,k+1,j}^*(x)$$
  
for  $k \in \{0, 1, \dots, n-1\}$ .  
(c)  $q_{n,0,j}(x) = q_{n,0,j}^*(x) = q_{n,1,j}^*(x) = p_{n,j}(x)$  for  $j \in \{0, 1, \dots, n\}$ .  
(d)  $q_{n,k,k}(x) = q_{n,k+1,k}(x)$  for  $k \in \{1, 2, \dots, n\}$ .  
(e)  $q_{n,k,j}(x) = q_{n,k-1,j}(x) + xq_{n-1,k-1,j-1}(x)$  for  $2 \le k \le j \le n$ .  
(f) For  $k \ge 1$ ,

n-1

(13) 
$$q_{n,k,j}^*(x) = x \sum_{i=0}^{j-1} q_{n-1,k-1,i}^*(x) + \sum_{i=j}^{n-1} q_{n-1,k-1,i}^*(x)$$

for 
$$j \in \{1, 2, ..., k - 1\}$$
 and

(14) 
$$q_{n,k,j}^*(x) = x \sum_{i=0}^{j-1} q_{n-1,k,i}^*(x) + \sum_{i=j}^{n-1} q_{n-1,k,i}^*(x)$$

for 
$$j \in \{k, k+1, \dots, n\}$$
.  
(g)  $q_{n,k,n}^*(x) = xq_{n,k-1}(x)$  for  $k \in \{1, 2, \dots, n\}$ 

*Proof.* Part (a) and the first equality of part (b) are immediate consequences of Equation (7) and the defining equations (9) and (12) of  $q_{n,k,j}(x)$  and  $q_{n,k,j}^*(x)$ . For the second equality of part (b) we apply Equation (12), the first equality of part (b), part (a) and the recurrence of Proposition 3.1 (a), respectively, to get

$$\sum_{j=0}^{n-1} q_{n-1,k+1,j}^*(x) = \sum_{j=0}^{n-1} q_{n-1,k+1,j}(x) - xq_{n-1,k+1,0}^* = q_{n,k+1}(x) - xq_{n-1,k}(x) = q_{n,k}(x)$$

Similarly, part (c) follows from Equation (3) and the definitions of  $q_{n,k,j}(x)$  and  $q_{n,k,j}^*(x)$  and part (d) follows from definition of  $q_{n,k,j}(x)$ , since k+1 cannot be a fixed point of a permutation  $w \in \mathfrak{S}_{n+1}$  with  $w^{-1}(1) = k+1$ . To verify part (e), we split the right-hand side of Equation (9) into two sums, running over permutations  $w \in \mathfrak{S}_{n+1}$  with  $w^{-1}(1) = j+1$  for which k is a fixed point, or k is not a fixed point, respectively. We then note that the first sum is equal to  $(1+x)q_{n-1,k-1,j-1}(x)$  and the second to  $q_{n,k-1,j}(x) - q_{n-1,k-1,j-1}(x)$ .

Part (f) reduces to the recurrence (4) for the polynomials  $p_{n,k}(x)$  for k = 1 and follows from Equation (10) by standard arguments for  $k \ge 2$ . For part (g) we denote by  $\text{fix}'_k(w)$  the number of fixed points of w in  $\{2, 3, \ldots, k\}$  and compute from Equation (9) that

$$q_{n,k,n}^*(x) = x \sum_{w \in \mathfrak{S}_n} (1+x)^{\operatorname{fix}_k'(w)} x^{\operatorname{exc}(w)} = x \left( q_{n,k}(x) - x q_{n-1,k-1}(x) \right) = x q_{n,k-1}(x)$$

for  $k \in \{1, 2, \dots, n\}$ .

Proof of Theorem 1.1. We claim that  $(q_{n,k,j}(x))_{0 \le j \le n}$  is an interlacing sequence of real-rooted polynomials for all  $n \in \mathbb{N}$  and  $k \in \{0, 1, \ldots, n+1\}$ .

Given the claim, Proposition 3.2 (b) implies that  $q_{n,k}(x)$  is a real-rooted polynomial which is interlaced by  $q_{n-1,k+1,0}^*(x) = q_{n-1,k}(x)$  for  $k \in \{0, 1, \ldots, n-1\}$ . As a result,  $q_{n,k}(x) \preceq xq_{n-1,k}(x)$ and hence  $q_{n,k+1}(x) = q_{n,k}(x) + xq_{n-1,k}(x)$  is real-rooted and it is interlaced by  $q_{n,k}(x)$  for every

 $k \in \{0, 1, \ldots, n-1\}$ . Since, as already mentioned,  $q_{n,0}(x) = A_n(x) \leq \widetilde{A}_n(x) = q_{n,n}(x)$ , it follows from Lemma 2.1 that  $(q_{n,k}(x))_{0 \leq k \leq n}$  is an interlacing sequence of real-rooted polynomials for every  $n \in \mathbb{N}$ . Therefore, by Lemma 2.2 (a), for every polynomial  $p(x) = \sum_{k=0}^{n} c_k x^{n-k} (1+x)^k$ with  $c_0, c_1, \ldots, c_n \geq 0$ ,

$$\mathcal{I}_n(\mathcal{A}^{\circ}(p(x))) = \sum_{k=0}^n c_k(q_{n,k}(x))$$

is a real-rooted polynomial such that

$$A_n(x) = q_{n,0}(x) \preceq \mathcal{I}_n(\mathcal{A}^{\circ}(p(x))) \preceq q_{n,n}(x) = \widehat{A}_n(x).$$

Since  $A_n(x)$  and  $\widetilde{A}_n(x)$  are symmetric, with centers of symmetry (n-1)/2 and n/2, respectively, this implies that  $\widetilde{A}_n(x) \preceq \mathcal{A}^{\circ}(p(x)) \preceq xA_n(x)$  and proves part (a). Given that  $A_n(x) \preceq xA_n(x)$ , the interlacing relations

$$A_n(x) \preceq \mathcal{I}_n(\mathcal{A}^{\circ}(p(x))) \preceq A_n(x) \preceq \mathcal{A}^{\circ}(p(x)) \preceq x A_n(x)$$

and Lemma 2.1 imply that  $\mathcal{I}_n(\mathcal{A}^{\circ}(p(x))) \preceq \mathcal{A}^{\circ}(p(x))$ . Since  $\mathcal{A}^{\circ}(p(x))$  is already known to have a nonnegative (even unimodal) symmetric decomposition with respect to n [15, Theorem 3.2], the latter interlacing relation and [16, Theorem 2.7] imply part (b).

We now prove the claim (the proof reduces to part of the proof of [15, Theorem 4.4] in the special case k = n + 1). Proceeding by induction on n and k, we assume that  $n \ge 1$  and  $k \in \{0, 1, \ldots, n+1\}$  and that the result holds for all pairs which are lexicographically smaller than (n, k). By Proposition 3.2 (c), for  $k \in \{0, 1\}$  the claim reduces to the fact that  $(p_{n,k}(x))_{0 \le k \le n}$  is an interlacing sequence of real-rooted polynomials for every  $n \in \mathbb{N}$  (see Example 2.3). Suppose that  $k \ge 2$ . By parts (a) and (b) of Proposition 3.2 we have

$$q_{n,k,0}^*(x) = q_{n,k-1}(x) = \sum_{j=0}^{n-1} q_{n-1,k-1,j}(x) = (1+x)q_{n-1,k-1,0}^*(x) + \sum_{j=1}^{n-1} q_{n-1,k-1,j}^*(x).$$

Applying Lemma 2.2 (b) and the induction hypothesis to this expression, combined with those of Proposition 3.2 (f), we conclude that  $(q_{n,k,j}^*(x))_{0 \leq j \leq k-1}$  and  $(q_{n,k,j}^*(x))_{k \leq j \leq n}$  are interlacing sequences of real-rooted polynomials. Since the second sequence is empty for k = n+1, the claim holds in this case. Thus, we may assume that  $2 \leq k \leq n$ . By Lemma 2.1, it suffices to prove that  $q_{n,k,k-1}^*(x) \leq q_{n,k,k}^*(x)$  and  $q_{n,k,0}^*(x) \leq q_{n,k,n}^*(x)$ . The latter holds because  $q_{n,k,n}^*(x) = xq_{n,k,0}^*(x)$ by parts (a) and (g) of Proposition 3.2. For the former, recall from Proposition 3.2 (e) that

$$q_{n,k,k}^*(x) = q_{n,k-1,k}^*(x) + xq_{n-1,k-1,k-1}^*(x).$$

By Proposition 3.2 (d) and our induction hypothesis,  $q_{n,k,k-1}^*(x) = q_{n,k-1,k-1}^*(x) \preceq q_{n,k-1,k}^*(x)$ . Moreover, since  $(q_{n-1,k-1,j}(x))_{0 \le j \le n}$  is interlacing by induction, each term on the right-hand side of Equation (13) for j = k - 1 interlaces  $xq_{n-1,k-1,k-1}^*(x)$  and hence so does their sum  $q_{n,k,k-1}^*(x)$ . As a result,

$$q_{n,k,k-1}^*(x) \preceq q_{n,k-1,k}^*(x) + xq_{n-1,k-1,k-1}^*(x) = q_{n,k,k}^*(x)$$

and the proof follows.

**Example 3.3.** Applying Theorem 1.1 to  $p(x) = c_0 x^n + c_1 x^{n-1} (1+x) + c_2 x^{n-2} (1+x)^2$  shows that  $aA_n(x) + bA_{n-1}(x) + cA_{n-2}(x)$  is real-rooted whenever  $a \ge b - c \ge c \ge 0$ . Moreover, it has a real-rooted and interlacing symmetric decomposition with respect to n.

#### CHRISTOS A. ATHANASIADIS

#### 4. Combinatorics of the derangement transformation

The derangement transformation was introduced and studied by Brändén and Solus [16, Section 3.2]; it is the linear map  $\mathcal{D} : \mathbb{R}[x] \to \mathbb{R}[x]$  defined by setting  $\mathcal{D}(x^n) = d_n(x)$  for  $n \in \mathbb{N}$ , where

$$d_n(x) = \sum_{i=0}^n (-1)^i \binom{n}{i} A_{n-i}(x) = \sum_{w \in \mathfrak{S}_n: \operatorname{Fix}(w) = \emptyset} x^{\operatorname{exc}(w)}$$

is the *nth derangement polynomial* and Fix(w) is the set of fixed points of a permutation  $w \in \mathfrak{S}_n$ . This polynomial is known to be  $\gamma$ -positive (in particular, symmetric), with center of symmetry n/2, and real-rooted; see [5, Section 2.1.4] [16, Section 3.2] [27, Section 3.1] and references therein.

As shown in [16, Corollary 3.7], the derangement transformation satisfies the analogue of part (a) of Theorem 1.1. This section shows that the reciprocals of the polynomials

(15) 
$$d_{n,k}(x) = \mathcal{D}\left(x^k(1+x)^{n-k}\right)$$

have  $\gamma$ -positive (in fact, real-rooted and interlacing) symmetric decompositions with respect to n, for  $k \in \{0, 1, \ldots, n\}$ , a property that will be one of the ingredients of the proof of Theorem 1.2, and confirms the analogue of part (b) of Theorem 1.1 for  $\mathcal{D}$  (see Corollary 4.2). The polynomials  $d_{n,k}(x)$  are shown on Table 2 for  $n \leq 4$ .

	k = 0	k = 1	k = 2	k = 3	k = 4
n = 0	1				
n = 1	1	0			
n=2	1+x	x	x		
n = 3	$1 + 4x + x^2$	$3x + x^2$	$2x + x^2$	$x + x^2$	
n = 4	$1 + 11x + 11x^2 + x^3$	$7x + 10x^2 + x^3$	$4x + 9x^2 + x^3$	$2x + 8x^2 + x^3$	$x + 7x^2 + x^3$

TABLE 2. The polynomials  $d_{n,k}(x)$  for  $n \leq 4$ .

Part (b) of the following statement, which may be of independent interest, shows that  $d_{n,k}(x)$ can be written as a sum of two  $\gamma$ -positive polynomials with centers of symmetry n/2 and (n-1)/2. For  $k \in \{0, n\}$ , it reduces to known expressions (see, for instance, [5, Theorems 2.1 and 2.13]) for  $A_n(x) = d_{n,0}(x)$  and  $d_n(x) = d_{n,n}(x)$  which demonstrate their  $\gamma$ -positivity. The proof extends the proof of  $\gamma$ -positivity of  $d_n(x)$ , given in [10, Section 4]. We recall that a *decreasing run* of a permutation  $w \in \mathfrak{S}_n$  is defined as a maximal set of integers of the form  $\{a, a + 1, \ldots, b\} \subseteq [n]$ such that  $w(a) > w(a + 1) > \cdots > w(b)$ .

**Proposition 4.1.** For every positive integer n and every  $k \in \{0, 1, ..., n\}$ :

(a)

$$d_{n,k}(x) = \sum_{i=0}^{k} (-1)^{i} \binom{k}{i} A_{n-i}(x) = \sum_{w \in \mathfrak{S}_{n}: \operatorname{Fix}(w) \subseteq [n-k]} x^{\operatorname{exc}(w)},$$

$$d_{n,k}(x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \xi_{n,k,i}^+ x^i (1+x)^{n-2i} + \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} \xi_{n,k,i}^- x^i (1+x)^{n-1-2i}$$

where

- $\xi_{n,k,i}^+$  is equal to the number of permutations  $w \in \mathfrak{S}_n$  with w(1) > n k which have *i* decreasing runs and none of size one,
- $\xi_{n,k,i}^-$  is equal to the number of permutations  $w \in \mathfrak{S}_n$  with  $w(1) \leq n-k$  which have *i* decreasing runs and none, except possibly the first, of size one.
- (c) The polynomial  $\mathcal{I}_n(d_{n,k}(x))$  has a  $\gamma$ -positive symmetric decomposition with respect to n.

*Proof.* The first formula of part (a) follows from Equation (15) by expanding  $x^k(1+x)^{n-k}$  as  $x^k(1+x)^{n-k} = (1+x-1)^k(1+x)^{n-k} = \sum_{i=0}^k (-1)^i {k \choose i} (1+x)^{n-i}$ , applying the linearity of  $\mathcal{D}$  and recalling that

$$\mathcal{D}\left((1+x)^n\right) = \sum_{i=0}^n \binom{n}{i} d_i(x) = A_n(x)$$

for  $n \in \mathbb{N}$ . The second formula follows from the first by a standard inclusion-exclusion argument.

For part (b) we extend the proof of the special case k = n, given in [10, Section 4]. To sketch this argument, which is based on the idea of valley hopping [22, 23] (see also [5, Section 4.1] [31, Section 4.2]), we recall some notation and terminology. Let  $w = (w_1, w_2, \ldots, w_n) \in \mathfrak{S}_n$  be a permutation, written in one-line notation, so that  $w_i = w(i)$  for  $i \in [n]$ . An ascent (respectively, descent) of w is any index  $i \in [n-1]$  such that  $w_i < w_{i+1}$  (respectively,  $w_i > w_{i+1}$ ). The number of ascents (respectively, descents) of w is denoted by  $\operatorname{asc}(w)$  (respectively,  $\operatorname{des}(w)$ ). An entry  $w_j$ is said to be a left-to-right maximum of w if  $w_i < w_j$  for all i < j. We will denote by  $\mathcal{E}_{n,k}$  the set of all permutations  $w \in \mathfrak{S}_n$  such that j is a descent of w for every left-to-right maximum  $w_j > k$  of w.

The standard representation (essentially, the fundamental transformation, mentioned in Section 3) of permutations, as described in [38, p. 23] and [10, Section 4], and the second expression of Proposition 4.1 (a) for  $d_{n,k}(x)$  show that

$$d_{n,k}(x) = \sum_{w \in \mathcal{E}_{n,n-k}} x^{\operatorname{asc}(w)}.$$

We claim that

(16) 
$$\sum_{w \in \mathcal{E}_{n,n-k}: w(1) > n-k} x^{\operatorname{asc}(w)} = \sum_{i=0}^{\lfloor n/2 \rfloor} \xi^+_{n,k,i} x^i (1+x)^{n-2i}$$

(17) 
$$\sum_{w \in \mathcal{E}_{n,n-k}: w(1) \le n-k} x^{\operatorname{asc}(w)} = \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} \xi_{n,k,i}^{-} x^{i} (1+x)^{n-1-2i},$$

where  $\xi_{n,k,i}^+$  and  $\xi_{n,k,i}^-$  are as in the statement of the proposition. This implies part (b).

Let  $w = (w_1, w_2, \ldots, w_n) \in \mathcal{E}_{n,n-k}$  and let us assume first that  $w_1 > n - k$  or, equivalently, that all left-to-right maxima of w are larger that n - k. As a result, all left-to-right maxima of ware located at descents of w and, in particular,  $w_1 > w_2$ . We set  $w_0 = 0$  and  $w_{n+1} = n + 1$  and call  $i \in [n]$  a *double ascent* (respectively, *double descent*) of w if  $w_{i-1} < w_i < w_{i+1}$  (respectively,  $w_{i-1} > w_i > w_{i+1}$ ). Given a double ascent or double descent i of w, and setting  $w_i = a$ , we define the permutation  $\varphi_a(w) \in \mathfrak{S}_n$  as follows: if i is a double ascent of w, then  $\varphi_a(w)$  is the permutation obtained from w by moving  $w_i = a$  between  $w_j$  and  $w_{j+1}$ , where j is the largest index satisfying  $1 \leq j < i$  and  $w_j > w_i > w_{j+1}$  (such an index exists because i is not a descent

#### CHRISTOS A. ATHANASIADIS

of w and hence a is not a left-to-right maximum). Similarly, if i is a double descent of w, then  $\varphi_a(w)$  is the permutation obtained from w by moving  $w_i = a$  between  $w_j$  and  $w_{j+1}$ , where j is the smallest index satisfying  $i < j \leq n$  and  $w_j < w_i < w_{j+1}$  (such an index exists because  $w_{n+1} = n + 1$ ). We set  $\varphi_a(w) = w$  for all other  $a \in [n]$ .

We observe that w and  $\varphi_a(w)$  have the same left-to-right maxima (all larger that n-k) for all  $w \in \mathcal{E}_{n,n-k}$  and  $a \in [n]$  and conclude that the maps  $\varphi_a$  are pairwise commuting involutions on the set of elements of  $\mathcal{E}_{n,n-k}$  with first entry larger than n-k. Thus, they define a  $\mathbb{Z}_2^n$ -action on this set. Moreover, each orbit of this action has a unique element u having no double ascent (equivalently, all decreasing runs of u have size at least two). As in the proof of [10, Theorem 1.4] we find that

$$\sum_{w \in \operatorname{Orb}(u)} x^{\operatorname{asc}(w)} = x^{\operatorname{asc}(u)} (1+x)^{n-2\operatorname{asc}(u)},$$

where Orb(u) stands for the orbit of u. Summing over all orbits yields Equation (16).

Finally, we consider permutations  $w = (w_1, w_2, \ldots, w_n) \in \mathcal{E}_{n,n-k}$  such that  $w_1 \leq n-k$ . We now set  $w_0 = w_{n+1} = n+1$  and define  $\varphi_a(w)$  for  $a \in [n]$  using the same rules as before. We leave to the reader to verify that  $\varphi_a(w)$  has the same left-to-right maxima larger that n-k as w, but those which are less than n-k may differ; in particular, w and  $\varphi_a(w)$  may have different first entries. Nevertheless, the first entry of  $\varphi_a(w)$  is still no larger than n-k and the maps  $\varphi_a$ define a  $\mathbb{Z}_2^n$ -action on the set of elements of  $\mathcal{E}_{n,n-k}$  with first entry no larger than n-k. Again, each orbit of this action has a unique element  $u = (u_1, u_2, \ldots, u_n)$  having no double ascent, although it is now possible that  $u_1 < u_2$  (equivalently, all decreasing runs of u other than the first have size at least two). As in the special case k = 0 (see, for instance, [5, Section 4.1] [31, Section 4.2]) we find that

$$\sum_{w \in Orb(u)} x^{\operatorname{asc}(w)} = x^{\operatorname{asc}(u)} (1+x)^{n-1-2\operatorname{asc}(u)}$$

Summing over all orbits yields Equation (17). Part (c) follows from part (b).

u

As a consequence of [16, Theorem 3.6], each polynomial  $d_{n,k}(x)$  is real-rooted and the sequence  $(d_{n,k}(x))_{0 \le k \le n}$  is interlacing for every  $n \in \mathbb{N}$ . The following statement implies that the symmetric decompositions of Proposition 4.1 (c) are, in fact, real-rooted and interlacing.

**Corollary 4.2.** The polynomial  $\mathcal{I}_n(\mathcal{D}(p(x)))$  has a real-rooted and interlacing symmetric decomposition with respect to n for every  $p(x) \in \mathcal{P}_n[x]$ .

*Proof.* As already mentioned, it has been shown in [16, Corollary 3.7] that  $\mathcal{D}(p(x))$  is realrooted and that it is interlaces  $\mathcal{I}_n(\mathcal{D}(p(x)))$ . Thus, by [16, Theorem 2.7], it suffices to confirm that  $\mathcal{I}_n(\mathcal{D}(p(x)))$  has a nonnegative symmetric decomposition with respect to n. Since,

$$\mathcal{I}_n(\mathcal{D}(p(x))) = \sum_{k=0}^n c_k \,\mathcal{I}_n(d_{n,k}(x))$$

for every  $p(x) = \sum_{k=0}^{n} c_k x^k (1+x)^{n-k} \in \mathcal{P}_n[x]$ , this is guaranteed by Proposition 4.1 (c).  $\Box$ 

# 5. SIMPLICIAL COMPLEXES AND THEIR TRIANGULATIONS

This section includes background on simplicial complexes, their triangulations and their face enumeration which is essential to understand and prove Theorem 1.2. Moreover, the polynomials  $q_{n,k}(x)$  and  $d_{n,k}(x)$  of Sections 3 and 4 are generalized in the setting of uniform triangulations and basic properties of these generalizations are discussed.

5.1. Simplicial complexes. We assume familiarity with basic notions, such as abstract and geometric simplicial complexes and the correspondence between them, as explained, for instance, in [13, 28, 37]. All simplicial complexes considered here will be abstract and finite. Following [7], we denote by  $\sigma_n$  the abstract simplex  $2^V$  on an *n*-element vertex set *V*.

A fundamental enumerative invariant of a simplicial complex  $\Delta$  is the *h*-polynomial, defined by the formula

(18) 
$$h(\Delta, x) = \sum_{i=0}^{n} f_{i-1}(\Delta) x^{i} (1-x)^{n-i},$$

where  $f_i(\Delta)$  is the number of *i*-dimensional faces of  $\Delta$  and n-1 is its dimension. The *h*-polynomial has nonnegative coefficients if  $\Delta$  triangulates a ball or a sphere [37, Chapter II]. Moreover, it is symmetric in the later case, with center of symmetry n/2.

Suppose that  $\Delta$  triangulates the (n-1)-dimensional ball. The *interior* h-polynomial  $h^{\circ}(\Delta, x)$ of  $\Delta$  is defined by the right-hand side of Equation (18) when  $f_{i-1}(\Delta)$  is replaced by the number of (i-1)-dimensional interior faces of  $\Delta$ . Then,  $h^{\circ}(\Delta, x) = x^n h(\Delta, 1/x)$  (see, for instance, [34, Lemma 6.2]), a formula which the reader may wish to consider as the definition of  $h^{\circ}(\Delta, x)$ . The *theta polynomial* of  $\Delta$  is defined as

(19) 
$$\theta(\Delta, x) = h(\Delta, x) - h(\partial \Delta, x),$$

where  $\partial \Delta$  is the boundary complex of  $\Delta$ . This polynomial is symmetric, with center of symmetry n/2, and under some mild hypotheses it has nonnegative coefficients. For other basic properties of theta polynomials we refer the reader to [8], where their role in the enumerative theory of triangulations of simplicial complexes is also explained.

5.2. **Triangulations.** By the term *triangulation* of a simplicial complex  $\Delta$  we will always mean a geometric triangulation. Thus, a simplicial complex  $\Delta'$  is a triangulation of  $\Delta$  if there exists a geometric realization K' of  $\Delta'$  which geometrically subdivides a geometric realization K of  $\Delta$ . The restriction of  $\Delta'$  to a face  $F \in \Delta$  is a triangulation of the simplex  $2^F$  denoted by  $\Delta'_F$ .

The local h-polynomial of a triangulation  $\Gamma$  of a simplex  $2^V$  is defined [36, Definition 2.1] by the formula

(20) 
$$\ell_V(\Gamma, x) = \sum_{F \subseteq V} (-1)^{|V \setminus F|} h(\Gamma_F, x).$$

Stanley [36] showed that  $\ell_V(\Gamma, x)$  has nonnegative coefficients and that it is symmetric, with center of symmetry |V|/2. The significance of local *h*-polynomials stems from Stanley's Locality Formula [36, Theorem 3.2], which expresses the *h*-polynomial of a triangulation  $\Delta'$  of a pure simplicial complex  $\Delta$  in terms of the local *h*-polynomials of the restrictions of  $\Delta'$  to the faces of  $\Delta$  and the *h*-polynomials of the links of these faces in  $\Delta$ .

**Barycentric subdivision.** Barycentric subdivision is a prototypical example of uniform triangulation of a simplicial complex. The *barycentric subdivision* of a simplicial complex  $\Delta$ , denoted by sd( $\Delta$ ), is defined as the simplicial complex which consists of all chains of nonempty faces of  $\Delta$ . As is well known, sd( $\Delta$ ) can be realized as a triangulation of  $\Delta$ .

The antiprism construction. Every triangulation  $\Gamma$  of a simplex  $2^V$  can be extended to a triangulation of a sphere of the same dimension as follows. Let  $V = \{v_1, v_2, \ldots, v_n\}$  and pick an *n*-element set  $U = \{u_1, u_2, \ldots, u_n\}$  which is disjoint from the vertex set of  $\Gamma$ . The antiprism sphere over  $\Gamma$  [4, Section 4], denoted by  $\Delta_{\mathcal{A}}(\Gamma)$ , is defined as the collection of sets of the form  $E \cup G$ , where  $E = \{u_i : i \in I\}$  is a face of the simplex  $2^U$  for some  $I \subseteq [n]$  and G is a face of the restriction  $\Gamma_F$  of  $\Gamma$  to the face  $F = \{v_j : j \in [n] \setminus I\}$  of the simplex  $2^V$  which is complementary to E. The complex  $\Delta_{\mathcal{A}}(\Gamma)$  is a triangulation of the (n-1)-dimensional sphere which contains  $2^U$  and  $\Gamma$  as subcomplexes and naturally triangulates  $\Delta_{\mathcal{A}}(2^V)$ ; for other basic properties, see [4, Proposition 4.6] [6, Proposition 4.1] and [9, Section 3].

5.3. Uniform triangulations. Let  $\mathcal{F} = (f_{\mathcal{F}}(i, j))_{0 \le i \le j \le n}$  be a triangular array of nonnegative integers. A triangulation  $\Gamma$  of the simplex  $\sigma_n = 2^V$  is called  $\mathcal{F}$ -uniform [7] if for all  $0 \le i \le j \le n$ , the restriction of  $\Gamma$  to any face of  $\sigma_n$  of dimension j-1 has exactly  $f_{\mathcal{F}}(i, j)$  faces of dimension i-1, and uniform if it is  $\mathcal{F}$ -uniform for some  $\mathcal{F}$  (these definitions extend naturally to triangulations of any simplicial complex of dimension less than n). The array  $\mathcal{F}$  is called the *f*-triangle associated to  $\Gamma$ .

We define the maps  $\mathcal{H}_{\mathcal{F}}^{\circ}, \mathcal{L}_{\mathcal{F}}: \mathbb{R}_n[x] \to \mathbb{R}_n[x]$  by setting

(21) 
$$\mathcal{H}^{\circ}_{\mathcal{F}}(x^m) = h^{\circ}_{\mathcal{F}}(\sigma_m, x) = x^m h_{\mathcal{F}}(\sigma_m, 1/x)$$

(22) 
$$\mathcal{L}_{\mathcal{F}}(x^m) = \ell_{\mathcal{F}}(\sigma_m, x)$$

for  $m \in \{0, 1, ..., n\}$  and extending by linearity, where  $h_{\mathcal{F}}(\sigma_m, x)$ ,  $h^{\circ}_{\mathcal{F}}(\sigma_m, x)$  and  $\ell_{\mathcal{F}}(\sigma_m, x)$ are the *h*-polynomial, the interior *h*-polynomial and the local *h*-polynomial, respectively, of the restriction of  $\Gamma$  to any (m-1)-dimensional face of  $\sigma_n$ . In analogy with the cases of Eulerian and derangement transformations, we consider the polynomials  $q_{\mathcal{F},m,k}(x)$  and  $\ell_{\mathcal{F},m,k}(x)$  defined by

(23) 
$$\mathcal{I}_m(q_{\mathcal{F},m,k}(x)) = \mathcal{H}^{\circ}_{\mathcal{F}}\left(x^{m-k}(1+x)^k\right)$$

(24) 
$$\ell_{\mathcal{F},m,k}(x)) = \mathcal{L}_{\mathcal{F}}\left(x^k(1+x)^{m-k}\right)$$

for all integers  $0 \le k \le m \le n$ . We have  $q_{\mathcal{F},m,0}(x) = \ell_{\mathcal{F},m,0}(x) = h_{\mathcal{F}}(\sigma_m, x)$  and  $\ell_{\mathcal{F},m,m}(x) = \ell_{\mathcal{F}}(\sigma_m, x)$  for every  $m \in \{0, 1, \ldots, n\}$ .

**Example 5.1.** Let  $\Gamma$  be the barycentric subdivision of  $2^V$ . Then,

$$\mathcal{H}^{\circ}_{\mathcal{F}}(x^m) = x^m h_{\mathcal{F}}(\sigma_m, 1/x) = x^m A_m(1/x) = \begin{cases} 1, & \text{if } m = 0\\ x A_m(x), & \text{if } m \ge 1 \end{cases}$$

and

$$\mathcal{L}_{\mathcal{F}}(x^m) = \ell_{\mathcal{F}}(\sigma_m, x) = d_m(x)$$

for  $m \in \{0, 1, \ldots, n\}$ ; see, for instance, [5, Section 3.3.1] [36, Proposition 2.4]. Hence,  $\mathcal{H}_{\mathcal{F}}^{\circ}$  and  $\mathcal{L}_{\mathcal{F}}$  are the restrictions of  $\mathcal{A}^{\circ}$  and  $\mathcal{D}$  to  $\mathbb{R}_{n}[x]$  and  $q_{\mathcal{F},m,k}(x) = q_{m,k}(x)$ ,  $\ell_{\mathcal{F},m,k}(x) = d_{m,k}(x)$  for all m, k.

**Proposition 5.2.** Let  $\Gamma$  be a uniform triangulation of an (n-1)-dimensional simplex  $2^V$  with associated *f*-triangle  $\mathcal{F}$ .

(a) The polynomials  $q_{\mathcal{F},m,k}(x)$  and  $\ell_{\mathcal{F},m,k}(x)$  satisfy the recurrence

$$q_{\mathcal{F},m,k+1}(x) = q_{\mathcal{F},m,k}(x) + xq_{\mathcal{F},m-1,k}(x)$$
$$\ell_{\mathcal{F},m,k+1}(x) = \ell_{\mathcal{F},m,k}(x) - \ell_{\mathcal{F},m-1,k}(x)$$

for  $0 \le k < m \le n$ .

(b) We have

$$q_{\mathcal{F},m,k}(x) = \sum_{i=0}^{k} \binom{k}{i} x^{i} h_{\mathcal{F}}(\sigma_{m-i}, x)$$
$$\ell_{\mathcal{F},m,k}(x) = \sum_{i=0}^{k} (-1)^{i} \binom{k}{i} h_{\mathcal{F}}(\sigma_{m-i}, x)$$

for  $0 \leq k \leq m \leq n$ .

- (c) Let  $\Delta_{\mathcal{A}}(\Gamma)$  be the antiprism sphere over  $\Gamma$ , with new vertices  $u_1, u_2, \ldots, u_n$ . Then, for every  $k \in \{0, 1, \ldots, n\}$ ,  $q_{\mathcal{F},n,k}(x)$  is equal to the h-polynomial of the induced subcomplex of  $\Delta_{\mathcal{A}}(\Gamma)$  obtained by removing vertices  $u_{k+1}, \ldots, u_n$ .
- (d) The polynomial  $q_{\mathcal{F},n,n}(x)$  is symmetric, with center of symmetry n/2.

Proof. The proofs of (a) and (b) are identical to those of the corresponding statements of Proposition 3.1 for the  $q_{\mathcal{F},m,k}(x)$  and follow by similar reasoning for the  $\ell_{\mathcal{F},m,k}(x)$ . Given part (b), one can extend the proof of part (d) of [6, Proposition 4.1] (which corresponds to the special case k = n) to prove part (c). To sketch a more direct argument, let  $\Delta$  be the induced subcomplex of  $\Delta_{\mathcal{A}}(\Gamma)$  obtained by removing vertices  $u_{k+1}, \ldots, u_n$ . By the defining equation (18) of the *h*-polynomial we have

$$h(\Delta, x) = \sum_{F \in \Delta} x^{|F|} (1-x)^{n-|F|} = \sum_{I \subseteq [k]} \sum_{F \in \Delta_I} x^{|F|} (1-x)^{n-|F|},$$

where  $\Delta_I$  consists of those faces  $F \in \Delta$  for which  $F \cap U = \{u_i : i \in I\}$ . One then recognizes the inner sum as  $x^{|I|}$  times the *h*-polynomial of the restriction of  $\Gamma$  to the face  $\{v_j : j \in [n] \setminus I\}$  of  $2^V$  which is complementary to  $\{u_i : i \in I\}$  and concludes that

$$h(\Delta, x) = \sum_{\{v_{k+1}, \dots, v_n\} \subseteq F \subseteq V} x^{n-|F|} h(\Gamma_F, x) = \sum_{i=0}^k \binom{k}{i} x^i h_{\mathcal{F}}(\sigma_{n-i}, x).$$

By part (b), the latter sum is equal to  $q_{\mathcal{F},n,k}(x)$ . Part (d) follows from part (c), which implies that  $q_{\mathcal{F},n,n}(x) = h(\Delta_{\mathcal{A}}(\Gamma), x)$  is the *h*-polynomial of a triangulation of the (n-1)-dimensional sphere.

# 6. Gamma-positivity of the $\mathcal{H}_{\mathcal{F}}^{\circ}$ and $\mathcal{L}_{\mathcal{F}}$ transformations

This section proves Theorem 1.2. We first introduce a family of enumerative invariants of a triangulation  $\Gamma$  of a simplex  $2^V$  which provides a common generalization of the *h*-polynomial and the local *h*-polynomial of  $\Gamma$ . This generalization is different from the relative local *h*-polynomial  $\ell_V(\Gamma, E, x)$ , introduced in [4, Section 3].

**Definition 6.1.** Given a triangulation  $\Gamma$  of the simplex  $2^V$  and a face  $E \subseteq V$  of  $2^V$ , the local *h*-polynomial of  $\Gamma$  with respect to V and E is defined by the formula

(25) 
$$\ell_{V,E}(\Gamma, x) = \sum_{E \subseteq F \subseteq V} (-1)^{|V \smallsetminus F|} h(\Gamma_F, x).$$

**Remark 6.2.** (a) By definition, we have  $\ell_{V,\emptyset}(\Gamma, x) = \ell_V(\Gamma, x)$  and  $\ell_{V,V}(\Gamma, x) = h(\Gamma, x)$ . We also have  $\ell_{V,E}(\Gamma, x) = h(\Gamma, x) - h(\Gamma_E, x)$  for every facet (maximal with respect to inclusion face) E of  $\partial(2^V)$ .

(b) By inclusion-exclusion [38, Section 2.1], for given  $E \subseteq G \subseteq V$ ,

$$h(\Gamma_G, x) = \sum_{E \subseteq F \subseteq G} \ell_{F, E}(\Gamma_F, x).$$

(c) Combining the defining equation (25) of  $\ell_{V,E}(\Gamma, x)$  with the identity of part (b), applied for  $E = \emptyset$ , we get

$$\ell_{V,E}(\Gamma, x) = \sum_{E \subseteq G \subseteq V} (-1)^{|V \smallsetminus G|} h(\Gamma_G, x) = \sum_{E \subseteq G \subseteq V} (-1)^{|V \smallsetminus G|} \left( \sum_{F \subseteq G} \ell_F(\Gamma_F, x) \right)$$
$$= \sum_{F \subseteq V} \ell_F(\Gamma_F, x) \left( \sum_{E \cup F \subseteq G \subseteq V} (-1)^{|V \smallsetminus G|} \right) = \sum_{V \smallsetminus E \subseteq F \subseteq V} \ell_F(\Gamma_F, x)$$

for every  $E \subseteq V$ . This formula shows that  $\ell_{V,E}(\Gamma, x)$  has nonnegative coefficients.

**Example 6.3.** (a) By part (a) of Proposition 4.1, in the special case of barycentric subdivision we have  $\ell_{V,E}(\Gamma, x) = d_{|V|,|V \setminus E|}(x)$  for every  $E \subseteq V$ .

(b) More generally, let  $\Gamma$  be the *r*-fold edgewise subdivision of the barycentric subdivision (termed in [7, Section 3] as the *r*-colored barycentric subdivision) of  $2^V$ . Let n = |V| and  $\mathbb{Z}_r \wr \mathfrak{S}_n$  be the group of *r*-colored permutations of the set [n] (see [6, Section 5] for any undefined terminology about *r*-colored permutations). By [6, Proposition 5.1] and an application of the inclusion-exclusion principle we have

(26) 
$$\ell_{V,E}(\Gamma, x) = \sum_{w \in (\mathbb{Z}_r \wr \mathfrak{S}_n)^b \colon \operatorname{Fix}(w) \subseteq [k]} x^{\operatorname{fexc}(w)/r},$$

where fexc(w) is the flag excedance number of  $w \in \mathbb{Z}_r \wr \mathfrak{S}_n$ , Fix(w) is the set of fixed points of  $w \in \mathbb{Z}_r \wr \mathfrak{S}_n$  of zero color,  $(\mathbb{Z}_r \wr \mathfrak{S}_n)^b$  is the set of  $w \in \mathbb{Z}_r \wr \mathfrak{S}_n$  such that the sum of the colors of the coordinates of w is divisible by r and k = |E|. This polynomial reduces to  $d_{n,n-k}(x)$  for r = 1.

The following two formulas were exploited in [8] in order to investigate unimodality and  $\gamma$ -positivity properties of *h*-polynomials of triangulations of balls and local *h*-polynomials of triangulations of simplices:

(27) 
$$h(\Gamma, x) = \sum_{F \subseteq V} \theta(\Gamma_F, x) A_{|V \setminus F|}(x)$$

(28) 
$$\ell_V(\Gamma, x) = \sum_{F \subseteq V} \theta(\Gamma_F, x) \, d_{|V \smallsetminus F|}(x)$$

The second formula is equivalent to [29, Theorem 4.7] and the first follows from that and the formula  $h(\Gamma, x) = \sum_{F \subseteq V} \ell_F(\Gamma_F, x)$ ; see [8, Theorem 3.4] for a generalization. The following statement provides a common generalization of the two formulas.

**Proposition 6.4.** For every triangulation  $\Gamma$  of the simplex  $2^V$  and every  $E \subseteq V$ ,

(29) 
$$\ell_{V,E}(\Gamma, x) = \sum_{F \subseteq V} \theta(\Gamma_F, x) \, d_{|V \smallsetminus F|, |V \smallsetminus (E \cup F)|}(x).$$

*Proof.* Using the defining equation (25) for  $\ell_{V,E}(\Gamma, x)$  and Equation (27), we get

$$\ell_{V,E}(\Gamma, x) = \sum_{E \subseteq G \subseteq V} (-1)^{|V \smallsetminus G|} h(\Gamma_G, x)$$
  
$$= \sum_{E \subseteq G \subseteq V} (-1)^{|V \smallsetminus G|} \left( \sum_{F \subseteq G} \theta(\Gamma_F, x) A_{|G \smallsetminus F|}(x) \right)$$
  
$$= \sum_{F \subseteq V} \theta(\Gamma_F, x) \left( \sum_{E \cup F \subseteq G \subseteq V} (-1)^{|V \smallsetminus G|} A_{|G \smallsetminus F|}(x) \right)$$
  
$$= \sum_{F \subseteq V} \theta(\Gamma_F, x) d_{|V \smallsetminus F|, |V \smallsetminus (E \cup F)|}(x)$$

and the proof follows.

We recall that a triangulation  $\Gamma$  of the simplex  $2^V$  is called *theta unimodal* (respectively, *theta*  $\gamma$ -positive) [8] if  $\theta(\Gamma_F, x)$  is unimodal (respectively,  $\gamma$ -positive) for every  $F \subseteq V$ . The barycentric subdivision of any regular cell decomposition of the simplex [29, Theorem 4.6], and the *r*-fold edgewise subdivision (for  $r \geq n$ ), antiprism triangulation, interval triangulation and *r*-colored barycentric subdivision of any triangulation of the simplex  $\sigma_n$  (by [8, Corollary 3.9], combined with results of [7, Section 7] [9, Section 5] [11, Section 5]) are among those triangulations of  $\sigma_n$  which are known to be theta  $\gamma$ -positive.

**Corollary 6.5.** Given any theta unimodal (respectively, theta  $\gamma$ -positive) triangulation  $\Gamma$  of the (n-1)-dimensional simplex  $2^V$ , the polynomials  $x^n \ell_{V,E}(\Gamma, 1/x)$  and  $\sum_{E \subseteq G \subseteq V} h^{\circ}(\Gamma_G, x)$  have a unimodal (respectively,  $\gamma$ -positive) symmetric decomposition with respect to n for every  $E \subseteq V$ .

Proof. Since  $\theta(\Gamma_F, x)$  is assumed to be unimodal (respectively,  $\gamma$ -positive), with center of symmetry |F|/2, for every  $F \subseteq V$ , the statement about  $x^n \ell_{V,E}(\Gamma, 1/x)$  follows directly from Propositions 4.1 (b) and 6.4. Applying the formula of Remark 6.2 (b) for  $h(\Gamma_G, x) = x^{|G|} h^{\circ}(\Gamma, 1/x)$  gives

$$\sum_{E \subseteq G \subseteq V} h^{\circ}(\Gamma_G, x) = \sum_{E \subseteq G \subseteq V} \left( \sum_{E \subseteq F \subseteq G} x^{|G|} \ell_{F,E}(\Gamma_F, 1/x) \right)$$
$$= \sum_{E \subseteq F \subseteq V} x^{|F|} \ell_{F,E}(\Gamma_F, 1/x) \left( \sum_{F \subseteq G \subseteq V} x^{|G \smallsetminus F|} \right)$$
$$= \sum_{E \subseteq F \subseteq V} x^{|F|} \ell_{F,E}(\Gamma_F, 1/x) (1+x)^{n-|F|}.$$

The last sum has a unimodal (respectively,  $\gamma$ -positive) symmetric decomposition with respect to n for every  $E \subseteq V$ , since  $x^{|F|} \ell_{F,E}(\Gamma_F, 1/x)$  has such a decomposition with respect to |F| for  $E \subseteq F \subseteq V$ .

The following statement is the analogue of Theorem 1.2 for the  $\mathcal{L}_{\mathcal{F}}$  transformation.

**Corollary 6.6.** Let  $\Gamma$  be a uniform triangulation of the (n-1)-dimensional simplex with associated *f*-triangle  $\mathcal{F}$ .

- (a) If  $\Gamma$  is theta unimodal, then  $\mathcal{I}_n(\mathcal{L}_{\mathcal{F}}(p(x)))$  has a unimodal symmetric decomposition with respect to n for every  $p(x) \in \mathcal{P}_n[x]$ .
- (b) If  $\Gamma$  is theta  $\gamma$ -positive, then  $\mathcal{I}_n(\mathcal{L}_{\mathcal{F}}(p(x)))$  has a  $\gamma$ -positive symmetric decomposition with respect to n for every  $p(x) \in \mathcal{P}_n[x]$ .

Proof. By Proposition 5.2, for a uniform triangulation  $\Gamma$  we have  $\ell_{\mathcal{F},n,k}(x) = \ell_{V,E}(\Gamma, x)$  for every (n-k)-element subset E of V. Thus,  $x^n \ell_{\mathcal{F},n,k}(1/x)$  is unimodal (respectively,  $\gamma$ -positive) for every  $k \in \{0, 1, \ldots, n\}$  by Corollary 6.5 and therefore so is

$$\mathcal{I}_n(\mathcal{L}_{\mathcal{F}}(p(x))) = \sum_{k=0}^n c_k \, \mathcal{I}_n \, \mathcal{L}_{\mathcal{F}}\left(x^{n-k}(1+x)^k\right) = \sum_{k=0}^n c_k x^n \ell_{\mathcal{F},n,n-k}(1/x)$$

for every polynomial  $p(x) = \sum_{k=0}^{n} c_k x^{n-k} (1+x)^k$  with  $c_0, c_1, \dots, c_n \ge 0$ .

Proof of Theorem 1.2. Similarly, it suffices to show that  $\mathcal{H}^{\circ}_{\mathcal{F}}(x^{n-k}(1+x)^k)$  has a unimodal (respectively,  $\gamma$ -positive) symmetric decomposition with respect to n for every  $k \in \{0, 1, \ldots, n\}$ . Indeed,

$$\mathcal{H}_{\mathcal{F}}^{\circ}\left(x^{n-k}(1+x)^{k}\right) = \sum_{i=0}^{k} \binom{k}{i} \mathcal{H}_{\mathcal{F}}^{\circ}(x^{n-i}) = \sum_{i=0}^{k} \binom{k}{i} h_{\mathcal{F}}^{\circ}(\sigma_{n-i}, x)$$

and the result follows from Corollary 6.5.

# 7. On the real-rootedness of the $\mathcal{H}_{\mathcal{F}}^{\circ}$ and $\mathcal{L}_{\mathcal{F}}$ transformations

This section discusses possible generalizations and analogues of [15, Conjecture 1] in the framework of uniform triangulations. Let  $\Gamma$  be a uniform triangulation of an (n-1)-dimensional simplex  $\sigma_n$ , with associated *f*-triangle  $\mathcal{F}$ . Following [7, 11], we say that  $\Gamma$  has the *strong interlacing property* if for all  $2 \leq m < n$ ,

- (i)  $h_{\mathcal{F}}(\sigma_m, x)$  is real-rooted,
- (ii)  $\theta_{\mathcal{F}}(\sigma_m, x)$  is either identically zero, or a real-rooted polynomial of degree m-1 with nonnegative coefficients which is interlaced by  $h_{\mathcal{F}}(\sigma_{m-1}, x)$ .

These conditions imply strong real-rootedness properties for the *h*-polynomials of  $\mathcal{F}$ -uniform triangulations of simplicial complexes and their symmetric decompositions [7, Section 6] [11, Section 4].

We recall that the polynomials  $q_{\mathcal{F},n,k}(x)$  and  $\ell_{\mathcal{F},n,k}(x)$  have been defined in Section 5 for  $k \in \{0, 1, \ldots, n\}$ . By linearity of  $\mathcal{H}^{\circ}_{\mathcal{F}}$  and  $\mathcal{L}_{\mathcal{F}}$  we have

$$\mathcal{I}_n(\mathcal{H}^{\circ}_{\mathcal{F}}(p(x))) = \sum_{k=0}^n c_k q_{\mathcal{F},n,k}(x)$$
$$\mathcal{L}_{\mathcal{F}}(p(x))) = \sum_{k=0}^n c_k \ell_{\mathcal{F},n,n-k}(x)$$

for every polynomial  $p(x) = \sum_{k=0}^{n} c_k x^{n-k} (1+x)^k$ . Thus, the last claim in each part of the following conjecture is a consequence of the first.

**Conjecture 7.1.** Let  $\Gamma$  be a uniform triangulation of the (n-1)-dimensional simplex  $2^V$  with associated *f*-triangle  $\mathcal{F}$ .

- (a) If  $\Gamma$  has the strong interlacing property, then  $(q_{\mathcal{F},n,k}(x))_{0 \leq k \leq n}$  is an interlacing sequence of real-rooted polynomials. In particular,  $h(\Delta_{\mathcal{A}}(\Gamma), x)$  is real-rooted and so is  $\mathcal{H}^{\circ}_{\mathcal{F}}(p(x))$ for every  $p(x) \in \mathcal{P}_n[x]$ .
- (b) If  $\Gamma$  has the strong interlacing property, then  $(\ell_{\mathcal{F},n,k}(x))_{0 \leq k \leq n}$  is an interlacing sequence of real-rooted polynomials. In particular,  $\ell_V(\Gamma, x)$  is real-rooted and so is  $\mathcal{L}_{\mathcal{F}}(p(x))$  for every  $p(x) \in \mathcal{P}_n[x]$ .

Conjecture 7.1 unifies several results and conjectures in the literature. For instance, the important special case of barycentric subdivision of Conjecture 7.1 holds by Theorem 1.1 and [16, Theorem 3.6]. The real-rootedness of  $\ell_V(\Gamma, x)$  was proven for the *r*-colored barycentric subdivision in [16, Section 3.3] [26, Section 5] and was conjectured for the antiprism triangulation in [9, Conjecture 5.8]. Among other instances, part (a) has been proven by the author for edgewise subdivisions (the proof will appear elsewhere) and has been verified computationally for the interval triangulation for  $n \leq 10$  and for the antiprism triangulation for  $n \leq 15$ . Part (b) has been verified for the interval and antiprism triangulations for  $n \leq 30$ .

**Remark 7.2.** We conclude with some remarks on Conjecture 7.1.

(a) By replacing  $h_{\mathcal{F}}^{\circ}(\sigma_m, x)$  with  $h_{\mathcal{F}}(\sigma_m, x)$  in the definition of the  $\mathcal{H}_{\mathcal{F}}^{\circ}$  transformation one gets the linear operator  $\mathcal{H}_{\mathcal{F}}: \mathbb{R}_n[x] \to \mathbb{R}_n[x]$  defined by setting

(30) 
$$\mathcal{H}_{\mathcal{F}}(x^m) = h_{\mathcal{F}}(\sigma_m, x)$$

for  $m \in \{0, 1, \ldots, n\}$ . In the special case of barycentric subdivision one gets the linear transformation  $\mathcal{A} : \mathbb{R}[x] \to \mathbb{R}[x]$  for which  $\mathcal{A}(x^n) = A_n(x)$  for every  $n \in \mathbb{N}$ . Computations suggest that Conjecture 7.1 may still hold when the  $\mathcal{H}^{\circ}_{\mathcal{F}}$  transformation is replaced by  $\mathcal{H}_{\mathcal{F}}$ .

(b) One may define the type B analogues  $\mathcal{B}, \mathcal{D}^B : \mathbb{R}_n[x] \to \mathbb{R}_n[x]$  of the Eulerian and derangement transformations, respectively, by setting  $\mathcal{B}(x^n) = B_n(x)$  and

$$\mathcal{D}^B(x^n) = \sum_{i=0}^n (-1)^i \binom{n}{i} B_{n-i}(x)$$

for  $n \in \mathbb{N}$ , where  $B_n(x)$  is the standard *n*th Eulerian polynomial of type B [31, Section 11.4]. Then, computational evidence again suggests that the polynomials  $\mathcal{B}(p(x))$  and  $\mathcal{D}^B(p(x))$  have only real roots for every  $p(x) \in \mathcal{P}[x]$ .

(c) More generally, given any sequence  $(h_n(x))_{n \in \mathbb{N}}$  of polynomials with nonnegative coefficients, one may define polynomials  $h_{n,k}(x)$  and  $\ell_{n,k}(x)$  for  $k \in \{0, 1, \ldots, n\}$  recursively by

setting

$$h_{n,k+1}(x) = h_{n,k}(x) + xh_{n-1,k}(x)$$
$$\ell_{n,k+1}(x) = \ell_{n,k}(x) - \ell_{n-1,k}(x)$$

for  $0 \le k < n$ , or explicitly, by setting

$$h_{n,k}(x) = \sum_{i=0}^{k} \binom{k}{i} x^{i} h_{n-i}(x)$$
$$\ell_{n,k}(x) = \sum_{i=0}^{k} (-1)^{i} \binom{k}{i} h_{n-i}(x)$$

for  $k \in \{0, 1, ..., n\}$ . Under what conditions on the  $h_n(x)$  are  $(h_{n,k}(x))_{0 \le k \le n}$  and  $(\ell_{n,k}(x))_{0 \le k \le n}$  interlacing sequences of real-rooted polynomials for every  $n \in \mathbb{N}$ ?

Since  $h_{n,0}(x) = h_n(x)$  and  $h_{n,1}(x) = h_{n,0}(x) + xh_{n-1,0}(x)$ , a necessary condition is that each  $h_n(x)$  is real-rooted and it is reasonable to assume that  $h_{n-1}(x)$  interlaces  $h_n(x)$  for every  $n \ge 1$ . The example  $h_n(x) = (1+x)^n$  shows that this condition is not sufficient even if each  $h_n(x)$  is assumed to be symmetric of degree n and center of symmetry n/2. Indeed, if  $h_n(x) = (1+x)^n$  for every  $n \in \mathbb{N}$ , then  $h_{n,k}(x) = (1+x)^{n-k}(1+2x)^k$  and  $\ell_{n,k}(x) = x^k(1+x)^{n-k}$  for all n, k and the sequences  $(h_{2,0}(x), h_{2,1}(x), h_{2,2}(x))$  and  $(\ell_{2,0}(x), \ell_{2,1}(x), \ell_{2,2}(x))$  already fail to be interlacing. We note that  $h_n(x) = (1+x)^n$  can be expressed as the h-polynomial of a nonuniform triangulation of the n-dimensional simplex.

Acknowledgments. The author wishes to thank Katerina Kalampogia-Evangelinou for her help with all computations claimed in this article.

#### References

- [1] K. Adiprasito, Combinatorial Lefschetz theorems beyond positivity, arXiv:1812.10454.
- [2] K. Adiprasito, S.A. Papadakis and V. Petrotou, Anisotropy, biased pairings and the Lefschetz property for pseudomanifolds and cycles, arXiv:2101.07245.
- [3] K. Adiprasito and G. Yashfe, The Partition Complex: an invitation to combinatorial commutative algebra, in Surveys in Combinatorics 2021, London Math. Soc. Lecture Notes Ser. 470, Cambridge University Press, Cambridge, 2021, pp. 1–41.
- [4] C.A. Athanasiadis, Flag subdivisions and  $\gamma$ -vectors, Pacific J. Math. 259 (2012), 257–278.
- [5] C.A. Athanasiadis, Gamma-positivity in combinatorics and geometry, Sém. Lothar. Combin. 77 (2018), Article B77i, 64pp (electronic).
- [6] C.A. Athanasiadis, Binomial Eulerian polynomials for colored permutations, J. Combin. Theory Series A 173 (2020), Article 105214, 38pp.
- [7] C.A. Athanasiadis, Face numbers of uniform triangulations of simplicial complexes, Int. Math. Res. Notices 2022 (2022), 15756–15787.
- [8] C.A. Athanasiadis, Triangulations of simplicial complexes and theta polynomials, arXiv:2209.01674.
- [9] C.A. Athanasiadis, J-M. Brunink and M. Juhnke-Kubitzke, Combinatorics of antiprism triangulations, Discrete Comput. Geom. 68 (2022), 72–106.
- [10] C.A. Athanasiadis and C. Savvidou, The local h-vector of the cluster subdivision of a simplex, Sém. Lothar. Combin. 66 (2012), Article B66c, 21pp (electronic).
- [11] C.A. Athanasiadis and E. Tzanaki, Symmetric decompositions, triangulations and real-rootedness, Mathematika 67 (2021), 840–859.
- [12] M. Beck, K. Jochemko and E. McCullough, h<sup>\*</sup>-polynomials of zonotopes, Trans. Amer. Math. Soc. 371 (2019), 2021–2042.

- [13] A. Björner, Topological methods, in Handbook of combinatorics (R.L. Graham, M. Grötschel and L. Lovász, eds.), North Holland, Amsterdam, 1995, pp. 1819–1872.
- [14] P. Brändén, Unimodality, log-concavity, real-rootedness and beyond, in Handbook of Combinatorics (M. Bona, ed.), CRC Press, 2015, pp. 437–483.
- [15] P. Brändén and K. Jochemko, The Eulerian transformation, Trans. Amer. Math. Soc. 375 (2022), 1917–1931.
- [16] P. Brändén and L. Solus, Symmetric decompositions and real-rootedness, Int. Math. Res. Not. 2021 (2021), 7764–7798.
- [17] F. Brenti, Unimodal, log-concave and Pólya frequency sequences in combinatorics, Mem. Amer. Math. Soc. 81 (1989), no. 413, pp. viii+106.
- [18] F. Brenti and V. Welker, f-vectors of barycentric subdivisions, Math. Z. 259 (2008), 849–865.
- [19] S. Corteel, I.M. Gessel, C.D. Savage and H.S. Wilf, The joint distribution of descent and major index over restricted sets of permutations, Ann. Comb. 11 (2007), 375–386.
- [20] S. Fisk, Polynomials, roots, and interlacing, arXiv:0612833.
- [21] D. Foata and M.-P. Schützenberger, Théorie Géometrique des Polynômes Eulériens, Lecture Notes in Mathematics 138, Springer-Verlag, 1970.
- [22] D. Foata and V. Strehl, Rearrangements of the symmetric group and enumerative properties of the tangent and secant numbers, Math. Z. 137 (1974), 257–264.
- [23] D. Foata and V. Strehl, Euler numbers and variations of permutations, in Colloquio Internazionale sulle Teorie Combinatorie (Roma, 1973), Tomo I, Atti dei Convegni Lincei, No. 17, Accad. Naz. Lincei, Rome, 1976, pp. 119–131.
- [24] G. Frobenius, Uber die Bernoullischen Zahlen und die Eulerschen Polynome, Berl. Ber. 1910 (1910), 809–847.
- [25] S.R. Gal, Real root conjecture fails for five- and higher-dimensional spheres, Discrete Comput. Geom. 34 (2005), 269–284.
- [26] N. Gustafsson and L. Solus, Derangements, Ehrhart theory and local h-polynomials, Adv. Math 369 (2020), Article 107169.
- [27] J. Haglund and P.B. Zhang, Real-rootedness of variations of Eulerian polynomials, Adv. in Appl. Math. 109 (2019), 38–54.
- [28] T. Hibi, Algebraic Combinatorics on Convex Polytopes, Carslaw Publications, Australia, 1992.
- [29] M. Juhnke-Kubitzke, S. Murai and R. Sieg, Local h-vectors of quasi-geometric and barycentric subdivisions, Discrete Comput. Geom. 61 (2019), 364–379.
- [30] S.A. Papadakis and V. Petrotou, The characteristic 2 anisotropicity of simplicial spheres, arXiv:2012.09815.
- [31] T.K. Petersen, Eulerian Numbers, Birkhäuser Advanced Texts, Birkhäuser, 2015.
- [32] A. Postnikov, V. Reiner and L. Williams, Faces of generalized permutohedra, Doc. Math. 13 (2008), 207–273.
- [33] J. Shareshian and M.L. Wachs, Gamma-positivity of variations of Eulerian polynomials, J. Comb. 11 (2020), 1–33.
- [34] R.P. Stanley, Generalized h-vectors, intersection cohomology of toric varieties and related results, in Commutative Algebra and Combinatorics (N. Nagata and H. Matsumura, eds.), Adv. Stud. Pure Math. 11, Kinokuniya, Tokyo and North-Holland, Amsterdam-New York, 1987, pp. 187–213.
- [35] R.P. Stanley, Log-concave and unimodal sequences in algebra, combinatorics, and geometry, in Graph Theory and its Applications: East and West, Annals of the New York Academy of Sciences 576, New York Acad. Sci., New York, 1989, pp. 500–535.
- [36] R.P. Stanley, Subdivisions and local h-vectors, J. Amer. Math. Soc. 5 (1992), 805–851.
- [37] R.P. Stanley, Combinatorics and Commutative Algebra, second edition, Birkhäuser, Basel, 1996.
- [38] R.P. Stanley, Enumerative Combinatorics, vol. 1, Cambridge Studies in Advanced Mathematics 49, Cambridge University Press, second edition, Cambridge, 2011.
- [39] D.G. Wagner, Total positivity of Hadamard products, J. Math. Anal. Appl. 163 (1992), 459–483.

DEPARTMENT OF MATHEMATICS, NATIONAL AND KAPODISTRIAN UNIVERSITY OF ATHENS, PANEPISTIMIOUPO-LIS, 15784 ATHENS, GREECE

Email address: caath@math.uoa.gr