# On Nonlinear Black Hole Ringdowns from Gauge-Invariance and Measurements

A. Kehagias $^{1,2,*}$  and A. Riotto $^{3,4,\dagger}$ 

<sup>1</sup>Physics Division, National Technical University of Athens, Athens, 15780, Greece

 $^{2}$  CERN, Theoretical Physics Department, Geneva, Switzerland

<sup>3</sup>Département de Physique Théorique and Centre for Astroparticle Physics (CAP), Université de Genève, 24 quai E. Ansermet, CH-1211 Geneva, Switzerland

<sup>4</sup> Gravitational Wave Science Center (GWSC), Université de Genève, CH-1211 Geneva, Switzerland
(Dated: February 3, 2023)

### Abstract

It has been recently shown that nonlinear effects emerging at the time of the generation of the quasinormal modes are necessary to model ringdowns from black hole mergers. In this note we describe how nonlinearities also arise when defining gauge-invariant tensor modes and in the calculation of the observable measured in the interferometers beyond linear order.

<sup>\*</sup> kehagias@central.ntua.gr

 $<sup>^{\</sup>dagger}$ Antonio. Riotto<br/>@unige.ch

#### I. INTRODUCTION AND CONCLUSIONS

First-order BH perturbation theory is standardly adopted to study the Quasi Normal Modes (QNMs) generated by a perturbed Black Hole (BH) during the ringing down phase [1, 2]. QNMs are determined solely by the mass, the spin and the charge of the BH and, as such, they are fundamental in gravitational wave astronomy. It has been recently showed by BH-merger simulations [3, 4] (see also Refs. [5–7]), that not only first-order but also second-order effects are relevant to describe ringdowns. In particular, the nonlinear mode amplitude arising from the square of the fundamental  $(\ell, m) = (2, 2)$  mode is comparable to or it can be even larger than that of the linear mode (4, 4). Therefore, to correctly model the BH ringdown, the inclusion of nonlinear effects is unavoidable. It is worth noting that, in the case of Kerr BHs, such nonlinearities find their explanation in symmetry arguments [8].

In this simple note we would like to describe how second-order effects arise by two other sources when dealing with gravitational wave strains. The first source arises when working with gauge-invariant second-order tensor perturbations, see also Refs. [9–13]. Tensor modes are gauge-invariant at first-order in perturbation theory, but they are not at second-order. The construction of a gauge-invariant second-order tensor mode unavoidably introduces (first-order)<sup>2</sup> terms which are potentially relevant when dealing with comparison between observation and theoretical predictions.

Unfortunately, there exist infinite ways to render tensor modes gauge invariant at second-order, depending on the gauge one starts from. Which gauge one should adopt is in fact suggested by the measurement procedure and, in order to give a description of the response of the detector, the best choice seems to be the so-called TT frame [14]. As we will see, the expression of the time shifts measured in interferometers in the TT gauge contains as well second-order (first-order)<sup>2</sup> terms. Furthermore, analytical calculation of the QNMs are best performed in the so-called Regge-Wheeler (RW) gauge where Schwarzschild perturbations are solved through the RW and Zerilli equations. We will see that second-order effects arise necessarily when expressing gauge-invariant tensor modes constructed from the TT gauge through the first-order gravitational wave strains calculated in the RW gauge. Of course, a precise estimation of these effects goes beyond the scope of this note, but it is the natural step to take in the near future.

The note is organized as follows. In Section II we discuss how to construct gauge-invariant second-order tensor perturbations. In Section III we deal with the second-order effects introduced by the measurement operation, while in Section IV we devote our attention to the second-order effects from the gauge-invariance. Section V contains our conclusions. The paper is supplemented by an Appendix which offers a similarity with the Stückelberg mechanism.

## II. GAUGE-INVARIANT SECOND-ORDER TENSOR PERTURBATIONS

In this section we summarize how to construct second-order gauge-invariant tensor modes. The expert reader can skip this section.

Before launching ourselves in technicalities, let us start with some general remarks. Firstly, let us point out that it does not exist a unique way to construct gauge-invariant tensor modes. Gauge-invariant objects, not depending upon the coordinate definition in a given gauge, can be defined. For instance, the tensor modes at first-order are gauge independent, since they remain the same in all gauges. On the contrary, the gravitational potential is gauge-dependent since it changes in different time slicings. A gauge-invariant combination can be constructed from the gravitational potential, but it is not unique. There is an infinite number of ways of making a gravitational potential gauge-invariant and what is the best gauge one should start from to compute the actual observables depends on the measurement which is performed. Similarly, for the tensor modes at second-order there is not a unique way to render them gauge-invariant, the starting point being dependent upon the of observation one performs and, often, on the

comparison between the measured quantity and the theoretical prediction (analytical or numerical). As we argued in the introduction, the TT and the RW gauges play a special role, and in the following we will focus our attention on those gauges. Let us first though proceed in full generality.

## A. Metric transformation

Under a generic coordinate transformation of the form

$$x^{\mu} \to \widetilde{x}^{\mu} = x^{\mu} + \xi^{\mu} \quad \text{with} \quad \xi^{\mu} \equiv (\alpha, \xi^{i}),$$
 (II.1)

a generic metric transforms as

$$g_{\mu\nu} \to \widetilde{g}_{\mu\nu} = \bar{g}_{\mu\nu} + \delta g_{\mu\nu},$$
 (II.2)

where

$$\delta g_{\mu\nu} = \xi_{\mu;\nu} + \xi_{\nu;\mu},\tag{II.3}$$

and  $\bar{g}_{\mu\nu}$  is the background unperturbed metric. A semicolon denotes covariant differentiation. The coordinate transformations in Eq. (II.1) are not considered to be infinitesimal, but it can also be finite. In that case, they can be expanded in terms of a fictitious, bookkeeping parameter  $\epsilon$ , which we will suppress in the following. Therefore, Eq. (II.1) is written to second-order as [15]

$$x^{\mu} \to \widetilde{x}^{\mu} = x^{\mu} + \xi_1^{\mu} + \frac{1}{2} \left( \xi_{1,\nu}^{\mu} \xi_1^{\nu} + \xi_2^{\mu} \right),$$
 (II.4)

where we have expanded

$$\xi^{\mu} = \xi_1^{\mu} + \frac{1}{2}\xi_2^{\mu}. \tag{II.5}$$

Expanding the metric to second-order as

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + \delta g_{\mu\nu} \tag{II.6}$$

where

$$\delta g_{\mu\nu} = \delta_1 g_{\mu\nu} + \frac{1}{2} \delta_2 g_{\mu\nu},\tag{II.7}$$

we find that the first and second-order metric perturbations  $\delta_1 g_{\mu\nu}$  and  $\delta_2 g_{\mu\nu}$ , transform as

$$\widetilde{\delta}_1 g_{\mu\nu} = \delta_1 g_{\mu\nu} + \bar{g}_{\mu\nu,\lambda} \xi_1^{\lambda} + \bar{g}_{\kappa\nu} \xi_{1,\mu}^{\kappa} + \bar{g}_{\mu\lambda} \xi_{1,\nu}^{\lambda}, \tag{II.8}$$

$$\widetilde{\delta_2}g_{\mu\nu} = \delta_2 g_{\mu\nu} + \bar{g}_{\mu\nu,\lambda} \xi_2^{\lambda} + \bar{g}_{\mu\lambda} \xi_{2,\nu}^{\lambda} + \bar{g}_{\lambda\nu} \xi_{2,\mu}^{\lambda} + 2 \Big( \delta_1 g_{\mu\nu,\lambda} \xi_1^{\lambda} + \delta_1 g_{\mu\lambda} \xi_{1,\nu}^{\lambda} + \delta_1 g_{\lambda\nu} \xi_{1,\mu}^{\lambda} + \delta_1 g_{\lambda\nu} \xi_{1,\mu}^{\lambda} + \delta_1 g_{\lambda\nu} \xi_{2,\mu}^{\lambda} + \delta_1 g_{\mu\lambda} \xi_{2,\mu}$$

$$+ \bar{g}_{\mu\lambda,\alpha}\xi_{1}^{\alpha}\xi_{1}^{\lambda}{}_{,\nu} + \bar{g}_{\lambda\nu,\alpha}\xi_{1}^{\alpha}\xi_{1}^{\lambda}{}_{,\mu} + \bar{g}_{\lambda\alpha}\xi_{1}^{\lambda}{}_{,\mu}\xi_{1}^{\alpha}{}_{,\nu} + \bar{g}_{\mu\nu,\lambda\alpha}\xi_{1}^{\lambda}\xi_{1}^{\alpha} + \bar{g}_{\mu\nu,\lambda}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\alpha} + \bar{g}_{\mu\nu,\lambda}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1}^{\lambda}{}_{,\alpha}\xi_{1$$

Knowing the transformation properties of the metric perturbations allow us to construct gauge-invariant quantities. We will demonstrate this for the case of flat Minkowski spacetime with background metric

$$ds^2 = -dt^2 + \gamma_{ij} dx^i dx^j, \qquad (II.11)$$

where the  $x^i$  are generic curvilinear coordinates (which later on we will take to be polar coordinates). We can parametrise the perturbed metric  $\delta g_{\mu\nu}$  as

$$\delta g_{00} = -2\phi, \qquad \delta g_{0i} = B_i, \qquad \delta g_{ij} = 2C_{ij}. \tag{II.12}$$

In terms of the SO(3) subgroup of the full Poincaré isometry group of the Minkowski background, the perturbed metric is decomposed into scalar-vector-tensor (SVT) components by defining

$$B_i = B_{,i} - S_i, \qquad C_{ij} = -\psi \delta_{ij} + E_{,ij} + F_{(i,j)} + \frac{1}{2} h_{ij}.$$
 (II.13)

The tensor and vector degrees of freedom are defined to be divergence-free (transverse) and traceless and they satisfy the conditions

$$S_{i,i} = 0,$$
  $F_{i,i} = 0,$  and  $h_i^i = h_{ij,j} = 0.$  (II.14)

We now express all quantities at first and second-order perturbations around the background as

$$\phi = \phi_1 + \frac{1}{2}\phi_2 + \dots$$
 (II.15a)

$$\psi = \psi_1 + \frac{1}{2}\psi_2 + \dots \tag{II.15b}$$

$$B = B_1 + \frac{1}{2}B_2 + \dots {(II.15c)}$$

$$E = E_1 + \frac{1}{2}E_2 + \dots {(II.15d)}$$

$$S_i = S_{1i} + \frac{1}{2}S_{2i} + \dots {(II.15e)}$$

$$F_i = F_{1i} + \frac{1}{2}F_{2i} + \dots {(II.15f)}$$

$$h_{ij} = h_{1ij} + \frac{1}{2}h_{2ij} + \dots$$
 (II.15g)

We express the vector  $\xi^{\mu}$  as

$$\xi^{\mu} = \left(\alpha_1 + \frac{1}{2}\alpha_2, \xi_1^i + \frac{1}{2}\xi_2^i\right) \quad \text{with} \quad \xi_a^i = \beta_a,^i + \gamma_a^i,$$
 (II.16)

where  $a = \{1, 2\}$  and  $\gamma_a^i$  are divergence-free vectorial parameters  $(\gamma_{,i}^i = 0)$ . Using the transformation of the metric Eq. (II.9) for the first-order quantities, we find that the first-order gauge transformations are given by [16]

$$\widetilde{\phi_1} = \phi_1 + \dot{\alpha_1},\tag{II.17a}$$

$$\widetilde{\psi}_1 = \psi_1, \tag{II.17b}$$

$$\widetilde{B}_1 = B_1 - \alpha_1 + \dot{\beta}_1, \tag{II.17c}$$

$$\widetilde{E}_1 = E_1 + \beta_1, \tag{II.17d}$$

$$\widetilde{S_1}^i = S_1^i - \dot{\gamma_1}^i, \tag{II.17e}$$

$$\widetilde{F_1}^i = F_1^i + \gamma_1^i, \tag{II.17f}$$

$$\widetilde{h}_{1ij} = h_{1ij} \,. \tag{II.17g}$$

Similarly, using Eq. (II.10), we find that at second-order the gauge transformation of the components of the metric perturbation can be written as

$$\widetilde{\phi_2} = \phi_2 + \dot{\alpha}_2 + \alpha_1 \left( \ddot{\alpha}_1 + 2\dot{\phi}_1 \right) + 2\dot{\alpha}_1 \left( \dot{\alpha}_1 + 2\phi_1 \right) + \xi_{1k} \left( \dot{\alpha}_1 + 2\phi_1 \right)^k + \dot{\xi}_{1k} \left( \alpha_{1,}^k - 2B_{1k} - \dot{\xi}_1^k \right), \tag{II.18a}$$

$$\widetilde{\psi}_2 = \psi_2 - \frac{1}{4} \mathcal{X}_k^k + \frac{1}{4} \nabla^{-2} \mathcal{X}_{,ij}^{ij},$$
(II.18b)

$$\widetilde{B}_2 = B_2 - \alpha_2 + \dot{\beta}_2 + \nabla^{-2} \mathcal{X}_{\mathbf{B}_{-k}^k},\tag{II.18c}$$

$$\widetilde{E}_2 = E_2 + \beta_2 + \frac{3}{4} \nabla^{-2} \nabla^{-2} \mathcal{X}^{ij}_{,ij} - \frac{1}{4} \nabla^{-2} \mathcal{X}^k_{k}, \tag{II.18d}$$

$$\widetilde{S}_{2i} = S_{2i} - \dot{\gamma}_{2i} - \mathcal{X}_{B_i} + \nabla^{-2} \mathcal{X}_{B_i}^{k}, \tag{II.18e}$$

$$\widetilde{F}_{2i} = F_{2i} + \gamma_{2i} + \nabla^{-2} \mathcal{X}_{ik,}^{k} - \nabla^{-2} \nabla^{-2} \mathcal{X}^{kl}_{,kli}, \tag{II.18f}$$

$$\widetilde{h}_{2ij} = h_{2ij} + \mathcal{X}_{ij} + \frac{1}{2} \left( \nabla^{-2} \mathcal{X}^{kl}_{,kl} - \mathcal{X}^{k}_{k} \right) \delta_{ij} + \frac{1}{2} \nabla^{-2} \nabla^{-2} \mathcal{X}^{kl}_{,klij}$$

$$+\frac{1}{2}\nabla^{-2}\mathcal{X}_{k,ij}^{k} - \nabla^{-2}\left(\mathcal{X}_{ik, j}^{k} + \mathcal{X}_{jk, i}^{k}\right). \tag{II.18g}$$

We have defined the vector  $\mathcal{X}_{B_i}$  and the tensor  $\mathcal{X}_{ij}$ , which both depend only on the square of the first-order quantities

$$\mathcal{X}_{B_{i}} \equiv 2\left(\dot{B}_{1i}\alpha_{1} + B_{1i,k}\xi_{1}^{k} - 2\phi_{1}\alpha_{1,i} + B_{1k}\xi_{1,i}^{k} + B_{1i}\dot{\alpha}_{1} + 2C_{1ik}\dot{\xi}_{1}^{k}\right) - \alpha_{1,k}\xi_{1,i}^{k} 
+ \dot{\alpha}_{1}\left(\dot{\xi}_{1i} - 3\alpha_{1,i}\right) + \alpha_{1}\left(\ddot{\xi}_{1i} - \dot{\alpha}_{1,i}\right) + \dot{\xi}_{1}^{k}\left(\xi_{1i,k} + 2\xi_{1k,i}\right) + \xi_{1}^{k}\left(\dot{\xi}_{1i,k} - \alpha_{1,ik}\right), \tag{II.19}$$

and

$$\mathcal{X}_{ij} \equiv 4 \left( \alpha_1 \dot{C}_{1ij} + C_{1ij,k} \xi_1^{\ k} + C_{1ik} \xi_1^{\ k}_{\ ,j} + C_{1kj} \xi_1^{\ k}_{\ ,i} \right) + 2 \left( B_{1i} \alpha_{1,j} + B_{1j} \alpha_{1,i} \right) 
- 2 \alpha_{1,i} \alpha_{1,j} + 2 \xi_{1k,i} \xi_1^{\ k}_{\ ,j} + \alpha_1 \left( \dot{\xi}_{1i,j} + \dot{\xi}_{1j,i} \right) + \left( \xi_{1i,jk} + \xi_{1j,ik} \right) \xi_1^{\ k} + \xi_{1i,k} \xi_1^{\ k}_{\ ,j} 
+ \xi_{1j,k} \xi_1^{\ k}_{\ ,i} + \dot{\xi}_{1i} \alpha_{1,j} + \dot{\xi}_{1j} \alpha_{1,i}.$$
(II.20)

## B. Construction of second-order gauge-invariant tensor modes

Let us remind the reader how to construct gauge-invariant quantities from a particular gauge [16, 17]. Choosing a gauge is equivalent to pick up a vector  $\xi^{\mu}$  such that certain conditions are satisfied by the metric. This enforces the parameters  $\alpha$  and  $\xi^{i}$  to be expressed in terms of the perturbation fields  $(\delta g)$  or some combination thereof. Then, the particular form of  $\alpha(\delta g)$  and  $\xi^{i}(\delta g)$  used to fix the gauge can be employed to perform a general gauge transformation of the original fields so that the new transformed fields are now gauge-invariant quantities.

Let us illustrate how this procedure works for the first-order scalar potentials  $\phi_1$  (see also Appendix A). We can choose to set the parameters  $\alpha_1$  and  $\beta_1$  to go to a gauge where  $\widetilde{B}_1 = \widetilde{E}_1 = 0$ . From Eqs. (II.17c) and (II.17d), this will determine  $\beta_1 = -E_1$  and  $\alpha_1 = B_1 - \dot{E}_1$ . Inserting these choices into Eq. (II.17a) we find a gauge-invariant expression for the gravitational potential

$$\phi_1^{\text{GI}} \equiv \phi_1 + \dot{B}_1 - \ddot{E}_1. \tag{II.21}$$

A similar procedure can be used to define gauge-invariant second-order transverse-free and traceless perturbation which, as we will discuss later on, is to be identified with the tensor modes in the TT gauge. Using the gauge transformation properties of the tensor as in Eq. (II.18g) one finds [16, 18]<sup>1</sup>

$$h_{2ij}^{\text{GI}} \equiv h_{2ij} + \mathcal{X}_{ij}^{\text{GC}} + \frac{1}{2} \left( \nabla^{-2} \mathcal{X}_{,kl}^{\text{GC}kl} - \mathcal{X}_{k}^{\text{GC}k} \right) \delta_{ij} + \frac{1}{2} \nabla^{-2} \nabla^{-2} \mathcal{X}_{,klij}^{\text{GC}kl}$$

<sup>&</sup>lt;sup>1</sup> Non-local terms in the definition of the gauge-invariant second-order tensor modes are there to ensure that the modes are transverse and traceless. They disappear in the "projected" equation of motion.

$$+\frac{1}{2}\nabla^{-2}\mathcal{X}_{k,ij}^{GCk} - \nabla^{-2}\left(\mathcal{X}_{ik,\ j}^{GCk} + \mathcal{X}_{jk,\ i}^{GCk}\right),\tag{II.22}$$

where now

$$\mathcal{X}_{ij}^{\text{GC}} \equiv 4 \left( \alpha_{1}^{\text{GC}} \dot{C}_{1ij} + C_{1ij,k} \xi_{1}^{\text{GC}k} + C_{1ik} \xi_{1}^{\text{GC}k} + C_{1kj} \xi_{1}^{\text{GC}k} \right) + 2 \left( B_{1i} \alpha_{1,j}^{\text{GC}} + B_{1j} \alpha_{1,i}^{\text{GC}} \right) \\
- 2 \alpha_{1,i}^{\text{GC}} \alpha_{1,j}^{\text{GC}} + 2 \xi_{1k,i}^{\text{GC}} \xi_{1-j}^{\text{GC}k} + \alpha_{1}^{\text{GC}} \left( \dot{\xi}_{1i,j}^{\text{GC}} + \dot{\xi}_{1j,i}^{\text{GC}} \right) + \left( \xi_{1i,jk}^{\text{GC}} + \xi_{1j,ik}^{\text{GC}} \right) \xi_{1}^{\text{GC}k} \\
+ \xi_{1i,k}^{\text{GC}} \xi_{1-j}^{\text{GC}k} + \xi_{1j,k}^{\text{GC}k} \xi_{1-i}^{\text{GC}k} + \dot{\xi}_{1i}^{\text{GC}} \alpha_{1,j}^{\text{GC}} + \dot{\xi}_{1j}^{\text{GC}} \alpha_{1,i}^{\text{GC}} \right) \tag{II.23}$$

in terms of the fields  $\alpha_1^{\text{GC}}(\delta g)$  and  $\xi_{1i}^{\text{GC}}(\delta g)$ . The label (GC) reminds that the corresponding quantity is specified by solving a given (and arbitrary) gauge condition. Different gauge conditions give rise to different gauge-invariant quantities by using this procedure. However, there are also gauge-independent quantities, that is quantities that are independent of the gauge used. For example, the first-order transverse-traceless part  $h_{1ij}$  of the tensor perturbation  $C_{ij}$  (like  $\psi_1$ ) is gauge-independent: it is invariant in any gauge. On the other hand,  $h_{2ij}^{\text{GI}}$  is gauge-invariant, but it depends on the gauge used since it depends on the parameter in  $\xi^{\text{GC}\mu}(\delta g)$ .

What is more relevant for us is that the gauge-invariant second-order tensor mode automatically contains in its definition terms of the form (first-order)<sup>2</sup> which come from two sources: those contain in the intrinsically second-order quantity  $h_{2ij}$  which are determined by the dynamics (i.e. the merger of two BHs) and those which are explicitly present in the tensor  $\mathcal{X}_{ij}^{\text{GC}}$  whose introduction is necessary and unavoidable to define the gauge-invariant second-order tensor mode.

# III. THE NONLINEARITIES FROM THE MEASUREMENT OF GWS

In this section we discuss the presence of the nonlinearities coming from the measurement operation of the GWs. In interferometer experiments of arms of length L the measurement is done by sending photons to the mirrors located and observing the modulation in power recorded because of the different time shifts  $\Delta t_{A,B}$  acquired in the different travel paths along the arms A and B. Since for space-based observatories (like LISA) the frequency  $\omega$  of the GWs is such that  $\omega L = \mathcal{O}(1)$ , a single reference frame may not be adopted for which the whole apparatus is described by an (approximately) flat metric in the presence of the GW. On the contrary, a completely general relativistic framework has to be used. In this respect, the most suitable coordinate system turns out to be the TT frame where the coordinates in the positions of the mirrors (for a thorough discussion about the virtues of using the TT frame see Ref. [14]).

Here we briefly summarises how to define the TT gauge at first-order. The conditions to impose are  $\delta g_{00} = \delta g_{0i} = 0$ , which at first-order lead to

$$\alpha_1^{\text{TT}} = -\left[\int \phi_1 dt - \mathcal{C}_1(\boldsymbol{x})\right],\tag{III.1}$$

$$\beta_1^{\text{TT}} = \int \left(\alpha_1^{\text{TT}} - B_1\right) dt + \hat{\mathcal{C}}_1(\boldsymbol{x}), \tag{III.2}$$

$$\gamma_{1i}^{\text{TT}} = \int S_{1i} dt + \hat{\mathcal{C}}_{1i}(\boldsymbol{x}). \tag{III.3}$$

The time-slicing is fully determined once two arbitrary functions of the spatial coordinates  $C_1(x)$  and  $\hat{C}_1(x)$  are fixed. Furthermore, there is an arbitrary 3-vector  $\hat{C}_{1i}$  (with 2 independent components) depending upon the choice of spatial coordinates on an initial hypersurface. These extra four degrees of freedom can be further fixed by imposing the transverse-traceless (TT) condition

$$C_1{}^{ik}{}_{,k} = C_{1k}^k = 0.$$
 (III.4)

A similar procedure can be performed at second-order. In the TT gauge the second-order metric (by writing only the GW perturbation and using cartesian coordinates) reads

$$ds^{2} = -dt^{2} + \left(\delta_{ij} + h_{1ij}^{TT} + \frac{1}{2}h_{2ij}^{TT}\right)dx^{i}dx^{j},$$
(III.5)

where  $h_{ij}^{\text{\tiny TT}}$  is the transverse and traceless components of the  $C_{ij}$  in Eq. (II.12).

The effect of a GW passing through the interferometer is captured by measuring the proper times at the interferometers. Photons travel along the arms following the geodesic equation  $ds^2 = 0$  and we find up to second-order that the time shift reads (along two arms)

$$\Delta t_{A,B} = L_{A,B} - \int_{t_0}^{t_0 + 2L_{A,B}} dt \left( -\frac{1}{2} h_{1ij}^{TT} + \frac{3}{8} (h_{1ij}^{TT})^2 - \frac{1}{4} h_{2ij}^{TT} \right)_{i=j=1,2}.$$
 (III.6)

From these simple arguments we see that second-order effects of the QNMs enter not only through the intrinsically second-order quantity  $h_{2ij}^{\text{TT}}$ , but also through the explicit (first-order)<sup>2</sup> term  $3(h_{1ij}^{\text{TT}})^2/8$  in the time shifts. We expect that rendering the time shifts fully gauge independent will cause the appearance of other (first-order)<sup>2</sup> terms which must be taken into account in the gravitational waveform analysis.

#### IV. THE NONLINEARITIES FROM GAUGE INVARIANCE

In Section II we showed that at first-order the tensor part of the metric perturbation  $h_{1ij}$  is invariant under coordinate transformation and that its second-order counterpart is not. This means that at second-order any gauge-invariant tensor quantity will contain pieces which are (first-order)<sup>2</sup> which will eventually contribute through the QNMs to the time shifts measured in the interferometers.

We also learned in Section II that the TT gauge is preferable when dealing with observations. It is therefore natural to construct a gauge-invariant quantity starting from the TT gauge as we described in Section II. The procedure will be therefore the following. Starting from a fully generic gauge, one chooses the parameters of the coordinate transformations (II.16) by setting  $\delta g_{00} = \delta g_{0i} = 0$  as well as the transverse and traceless condition on the spatial part of the metric. With this procedure we arrive at the generic gauge-invariant expression for the tensor mode at second-order (II.22). The transverse and traceless condition, by construction, does not depend neither on  $h_{1ij}^{\rm TT}$  nor on  $h_{2ij}^{\rm TT}$ . Therefore, the quadratic pieces in the gauge-invariant second-order tensor mode does not contain explicit terms of the form  $\mathcal{O}[(h_{1ij}^{\rm TT})^2]$ . Terms quadratic in  $h_{1ij}^{\rm TT}$  will appear only in  $h_{2ij}^{\rm TT}$  once its dynamics is solved at second-order. In other words

$$h_{2ij}^{\text{GI}}\Big|_{\text{in the TT gauge}} = h_{2ij}^{\text{TT}}.$$
 (IV.1)

In order to compute  $h_{2ij}^{\rm GI}$  one has therefore various choices. Either one computes it in the TT gauge directly and writes an equation to calculate the evolution of  $h_{2ij}^{\rm TT}$ , or otherwise, one can calculate  $h_{2ij}^{\rm GI}$  in another gauge. Alternatively, one can adopt a sort of mixed procedure where one calculates the second-order evolution in a given gauge and then go to the TT gauge. We will pursue this third option here. The so-called RW gauge is convenient to solve for the Schwarzschild perturbations through the RW and Zerilli equations, which describe axial and polar perturbations, respectively. We first write the TT part of the perturbation in polar coordinates as

$$h_{1\mu\nu}^{\text{TT}}(t,r,\theta,\phi) = \sum_{\ell=2}^{\infty} \sum_{m=-\ell}^{\ell} h_{\ell m}^{\text{TT}}(t,r) \left(t_{\ell m}^{(E2)}\right)_{\mu\nu},\tag{IV.2}$$

where

 $Y_{\ell,m}$  are the spherical harmonics. Following Ref. [12], we can write the fundamental modes at infinity in the TT gauge at first- and second-order as (we set  $G_N = 1$ )

$$h_{1,(2,\pm 2)}^{\text{TT}} \simeq \frac{1}{r} \psi_{1,(2,\pm 2)}(t,r)$$

$$h_{2,(4,\pm 4)}^{\text{TT}} \simeq \frac{1}{r} \psi_{2,(4,\pm 4)}(t,r) - i \frac{\sqrt{70} \cdot 51}{1512 \cdot 2\sqrt{\pi}} \frac{(\omega_{2,\pm 2})}{r} \left(\psi_{1,(2,\pm 2)}(t,r)\right)^{2}, \tag{IV.4}$$

where the  $\psi_{1,2(\ell,m)}$  satisfy the corresponding first- and second-order Zerilli equation in the RW gauge and the second-order part has been properly regularized. At first-order, for instance, the equation reads

$$\left(-\frac{\partial^{2}}{\partial t^{2}} + \frac{\partial^{2}}{\partial r_{*}^{2}} - V_{Z}(r)\right) \psi_{1,(\ell,m)}(t,r) = 0,$$

$$V_{Z}(r) = \left(1 - \frac{2M}{r}\right) \frac{2\lambda^{2}(\lambda+1)r^{3} + 6\lambda^{2}Mr^{2} + 18\lambda M^{2}r + 18M^{3}}{r^{3}(\lambda r + 3M)^{2}}, \quad \lambda = \frac{1}{2}(\ell-1)(\ell+2),$$

$$r^{*} = r + 2M \ln\left(\frac{r}{2M} - 1\right),$$
(IV.5)

and the reader can find in Ref. [12] the equation for  $\psi_{2,(\ell,m)}$  with its corresponding (regularized) second-order source. Taking the approximation that the QNMs are produced around the peak of the (minus of the) Zerilli potential  $V_Z$ , at  $(r_{\rm pk}^{\ell=2}/M) \simeq 3.1$  and  $(r_{\rm pk}^{\ell=4}/M) \simeq 3.05$  and using the fact that for the n=0 mode the fundamental wavefunctions are

$$\begin{split} \psi_{1,(2,\pm 2)}(t,r) &\simeq A_{1,(2,\pm 2)} e^{-i\omega_{(2,\pm 2)}t} e^{-z^2/4}, \\ \psi_{1,(4,\pm 4)}(t,r) &\simeq A_{1,(4,\pm 4)} e^{-i\omega_{(4,\pm 4)}t} e^{-z^2/4}, \\ &z &\simeq (4k)^{1/4} e^{3\pi i/4} (r^* - r_{\rm pk}^*), \quad r_{\rm pk}^*(\ell=2) \simeq 1.9, \quad r_{\rm pk}^*(\ell=4) \simeq 1.76, \\ &k = -V_Z''(r_{\rm pk}^*)/2, \end{split} \tag{IV.6}$$

a saddle-point approximation specifies the corresponding amplitude to be

$$A_{2,(4,\pm 4)} = \frac{A_{1,(2,\pm 2)}^2}{M} \left[ 0.05 - 0.08(\omega_{2,\pm 2}M) + 0.24(\omega_{2,\pm 2}M)^2 - 2.9(\omega_{2,\pm 2}M)^4 \right].$$
 (IV.7)

Taking  $(\omega_{2,\pm 2}M) \simeq 0.37$ , one finally finds that  $A_{2,(4,\pm 4)} \simeq 0.06 A_{1,(2,\pm 2)}^2/M$ . Since one expects  $A_{1,(4,\pm 4)} = \mathcal{O}(0.1) A_{1,(2,\pm 2)}/M$  [3], we indeed see that the second-order mode for  $\ell = 4$  can indeed be of the same order of magnitude of the corresponding linear mode for sizeable amplitudes of the fundamental mode  $\ell = 2$ . It will be interesting to see if some hidden symmetries may explain the nonlinearties of the Schwarzschild BH along the lines of Ref. [8] for extremal Kerr BHs.

#### ACKNOWLEDGMENTS

We thank V. De Luca and G. Franciolini for very useful discussions. A.R. acknowledges financial support provided by the Boninchi Foundation.

## Appendix A: The Stückelberg procedure

The construction in Section II can more easily be illustrated for the case of Maxwell theory. Here the gauge transformations are the U(1) transformations of the vector potential  $A_{\mu}$ , which read

$$A_{\mu} \to \widetilde{A}_{\mu} = A_{\mu} + \phi_{,\mu}. \tag{A.1}$$

We may express  $A_{\mu}$  as

$$A_{\mu} = (A_0, C_i + F_{,i})$$
 with  $C_{,i}^i = 0,$  (A.2)

so that the gauge transformation is given by

$$\widetilde{A}_0 = A_0 + \dot{\phi},$$

$$\widetilde{F} = F + \phi,$$

$$\widetilde{C}_i = C_i.$$
(A.3)

Therefore, like the case of metric perturbations, the transverse part of the gauge potential does not change under U(1) gauge transformations. Similarly, the scalars  $A_0$  and F change according to (A.3). Let us now choose a particular gauge, for example the temporal gauge

$$\widetilde{A}_0 = 0. (A.4)$$

Under a gauge transformation we get that

$$\widetilde{A}_0 = A_0 + \dot{\phi} = 0, \tag{A.5}$$

from where we find that

$$\dot{\phi} = -A_0. \tag{A.6}$$

The solution of the above equation provides what we call  $\phi^{\text{GC}}$ , or in other words

$$\phi^{GC} = -\int^t A_0(\tau, \vec{x}) d\tau. \tag{A.7}$$

Having specified the gauge parameter  $\phi^{GC}$ , it is easy to construct gauge-invariant quantities. For example, following the discussion above, the gauge-invariant scalar is now

$$F^{\text{GI}} = F_{\text{tm}}^{\text{GI}} = F + \phi^{\text{GC}} = F - \int_{0}^{t} A_0(\tau, \vec{x}) d\tau.$$
 (A.8)

The scalar  $F^{\text{GI}}$  is indeed gauge-invariant since under a gauge transformation of the form (A.1) with a gauge parameter  $\chi$ , we get that

$$\widetilde{A}_0 = A_0 + \dot{\chi}, \qquad \widetilde{F} = F + \chi,$$
(A.9)

and therefore, from Eq. (A.8), the scalar  $F^{\text{GI}}$  is gauge-invariant. In addition, the transverse part  $C_i$  of  $A_i$  does not change at all. If we use another gauge, let say the axial  $A_3 = 0$ , we can repeat the above construction. In this case though, the gauge-invariant scalar will be

$$F_{\text{ax}}^{\text{GI}} = F - \int_{0}^{x_3} A_3(t, x_a, y) dy, \qquad a = (1, 2)$$
 (A.10)

but still  $\widetilde{C}_i = C_i$ . In other words, changing the gauge, the gauge-invariant quantities change accordingly, while gauge-independent quantities do not change from gauge to gauge. Of course, one recognises in the above construction the Stückelberg trick to build gauge-invariant quantities.

- [1] K.D. Kokkotas and B.G. Schmidt, Quasinormal modes of stars and black holes, Living Rev. Rel. 2 (1999) 2 [gr-qc/9909058].
- [2] E. Berti, V. Cardoso and A.O. Starinets, Quasinormal modes of black holes and black branes, Class. Quant. Grav. 26 (2009) 163001 [0905.2975].
- [3] K. Mitman et al., Nonlinearities in black hole ringdowns, 2208.07380.
- [4] M.H.-Y. Cheung et al., Nonlinear effects in black hole ringdown, 2208.07374.
- [5] L. London, D. Shoemaker and J. Healy, Modeling ringdown: Beyond the fundamental quasinormal modes, Phys. Rev. D 90 (2014) 124032 [1404.3197].
- [6] S. Ma, K. Mitman, L. Sun, N. Deppe, F. Hébert, L.E. Kidder et al., Quasinormal-mode filters: A new approach to analyze the gravitational-wave ringdown of binary black-hole mergers, Phys. Rev. D 106 (2022) 084036 [2207.10870].
- [7] M. Lagos and L. Hui, Generation and propagation of nonlinear quasi-normal modes of a Schwarzschild black hole, 2208.07379.
- [8] A. Kehagias, D. Perrone, A. Riotto and F. Riva, Explaining Nonlinearities in Black Hole Ringdowns from Symmetries, 2301.09345.
- [9] R.J. Gleiser, C.O. Nicasio, R.H. Price and J. Pullin, Gravitational radiation from Schwarzschild black holes: The Second order perturbation formalism, Phys. Rept. 325 (2000) 41 [gr-qc/9807077].
- [10] M. Campanelli and C.O. Lousto, Second order gauge invariant gravitational perturbations of a Kerr black hole, Phys. Rev. D 59 (1999) 124022 [gr-qc/9811019].
- [11] D. Brizuela, J.M. Martin-Garcia and G.A.M. Marugan, *High-order gauge-invariant perturbations of a spherical spacetime*, *Phys. Rev. D* **76** (2007) 024004 [gr-qc/0703069].
- [12] H. Nakano and K. Ioka, Second Order Quasi-Normal Mode of the Schwarzschild Black Hole, Phys. Rev. D 76 (2007) 084007 [0708.0450].
- [13] N. Loutrel, J.L. Ripley, E. Giorgi and F. Pretorius, Second Order Perturbations of Kerr Black Holes: Reconstruction of the Metric, Phys. Rev. D 103 (2021) 104017 [2008.11770].
- [14] M. Maggiore, *Gravitational Waves. Vol. 1: Theory and Experiments*, Oxford Master Series in Physics, Oxford University Press (2007).
- [15] M. Bruni, S. Matarrese, S. Mollerach and S. Sonego, Perturbations of space-time: Gauge transformations and gauge invariance at second order and beyond, Class. Quant. Grav. 14 (1997) 2585 [gr-qc/9609040].
- [16] K.A. Malik and D. Wands, Cosmological perturbations, Phys. Rept. 475 (2009) 1 [0809.4944].
- [17] N. Bartolo, E. Komatsu, S. Matarrese and A. Riotto, Non-Gaussianity from inflation: Theory and observations, Phys. Rept. 402 (2004) 103 [astro-ph/0406398].

[18] S. Matarrese, S. Mollerach and M. Bruni, Second order perturbations of the Einstein-de Sitter universe, Phys. Rev. D 58 (1998) 043504 [astro-ph/9707278].