

# ABELIANIZATION AND THE DUISTERMAAT–HECKMAN THEOREM

PETER CROOKS AND JONATHAN WEITSMAN

ABSTRACT. Fix a compact connected Lie group  $G$  with Lie algebra  $\mathfrak{g}$ , as well as a strong Gelfand–Cetlin datum on  $\mathfrak{g}^*$ . Let us also fix a connected symplectic manifold  $M$  endowed with an effective Hamiltonian  $G$ -action and proper moment map. We associate to such information a measure on  $\mathbb{R}^b$ , where  $b = \frac{1}{2}(\dim G + \text{rank } G)$ . We also express the Radon–Nikodym derivative of this measure in terms of the volumes of the symplectic quotients of  $M$  by  $G$ , and thereby prove a non-abelian version of the Duistermaat–Heckman theorem.

## CONTENTS

1. Introduction	1
1.1. The Duistermaat–Heckman theorem	1
1.2. Gelfand–Cetlin data and abelianization	2
1.3. A non-abelian Duistermaat–Heckman theorem	2
1.4. Recovering a description of a measure on $\mathfrak{t}_+^*$	3
1.5. Organization	3
Acknowledgements	3
2. Main results	3
2.1. Lie theory	3
2.2. Gelfand–Cetlin data and abelianization	4
2.3. Strong Gelfand–Cetlin data	5
2.4. Non-abelian Duistermaat–Heckman measures	5
2.5. Radon–Nikodym derivatives	6
References	8

## 1. INTRODUCTION

**1.1. The Duistermaat–Heckman theorem.** Let  $(M, \omega)$  be a  $2n$ -dimensional connected symplectic manifold. The volume form  $\frac{1}{n!}\omega^n$  then determines a measure  $\lambda_M$  on  $M$ . Let us also suppose that  $M$  comes equipped with an effective Hamiltonian action of a compact torus  $T$ , and that this action admits a proper moment map  $\mu : M \rightarrow \mathfrak{t}^*$ . One may endow  $T$  with the normalized Haar measure for which  $T$  has unit volume; this gives rise to a Lebesgue measure on  $\mathfrak{t}$ , which in turn induces a dual Lebesgue measure  $\lambda_{\mathfrak{t}^*}$  on  $\mathfrak{t}^*$ . A second measure on  $\mathfrak{t}^*$  may be obtained by pushing  $\lambda_M$  forward along  $\mu$ ; this new measure  $\text{DH}_{\mathfrak{t}^*} := \mu_*\lambda_M$  is defined by

$$\text{DH}_{\mathfrak{t}^*}(B) = \int_{\mu^{-1}(B)} \frac{1}{n!}\omega^n$$

for all Borel subsets  $B \subset \mathfrak{t}^*$ .

Duistermaat and Heckman [4] relate  $\text{DH}_{\mathfrak{t}^*}$  to  $\lambda_{\mathfrak{t}^*}$  by means of the symplectic quotients

$$M //_{\xi} T := \mu^{-1}(\xi)/T,$$

---

2020 *Mathematics Subject Classification.* 53D20 (primary); 17B80 (secondary).

*Key words and phrases.* Duistermaat–Heckman measure, Gelfand–Cetlin system, symplectic quotient.

where  $\xi \in \mathfrak{t}^*$  ranges over the regular values of  $\mu$ . Each such quotient is a compact symplectic orbifold, and as such has a well-defined symplectic volume  $\text{vol}(M//_{\xi} T)$ . One version of the Duistermaat–Heckman theorem is that

$$\text{DH}_{\mathfrak{t}^*} = \rho_{\mathfrak{t}^*} \lambda_{\mathfrak{t}^*},$$

where  $\rho_{\mathfrak{t}^*} : \mathfrak{t}^* \rightarrow [0, \infty)$  is a Radon–Nikodym derivative of  $\text{DH}_{\mathfrak{t}^*}$  with respect to  $\lambda_{\mathfrak{t}^*}$  satisfying

$$\rho_{\mathfrak{t}^*}(\xi) = \text{vol}(M//_{\xi} T)$$

for all regular values  $\xi \in \mathfrak{t}^*$  of  $\mu$ . The underlying argument is largely based on the existence of appropriate local normal forms for  $\mu$ , as presented in [4, Equations (1.7) and (2.1)].

**1.2. Gelfand–Cetlin data and abelianization.** Let  $G$  be a compact connected Lie group with Lie algebra  $\mathfrak{g}$ . Our paper [2] defines what it means for a continuous map  $\nu_{\text{big}} : \mathfrak{g}^* \rightarrow \mathbb{R}_{\text{big}}$  and open dense subset  $\mathfrak{g}_{\text{s-reg}}^* \subset \mathfrak{g}^*$  to form a *Gelfand–Cetlin datum* on  $\mathfrak{g}^*$ , where

$$\mathfrak{b} := \frac{1}{2}(\dim G + \text{rank } G) \quad \text{and} \quad \mathbb{R}_{\text{big}} := \mathbb{R}^{\mathfrak{b}}.$$

Among other things,  $\nu_{\text{big}}|_{\mathfrak{g}_{\text{s-reg}}^*}$  is a moment map for a Hamiltonian action of

$$\mathbb{T}_{\text{big}} := \text{U}(1)^{\mathfrak{b}}$$

on the open Poisson submanifold  $\mathfrak{g}_{\text{s-reg}}^* \subset \mathfrak{g}^*$ ; further details are provided in [2, Definition 1]. The works of Guillemin–Sternberg [7, 8] and Hoffman–Lane [9] imply that Gelfand–Cetlin on  $\mathfrak{g}^*$  exist.

Now suppose that  $M$  is a Hamiltonian  $G$ -space with moment map  $\mu : M \rightarrow \mathfrak{g}^*$ , and that  $(\nu_{\text{big}}, \mathfrak{g}_{\text{s-reg}}^*)$  is a Gelfand–Cetlin datum on  $\mathfrak{g}^*$ . The torus  $\mathbb{T}_{\text{big}}$  then acts in a Hamiltonian fashion on the open subset  $M_{\text{s-reg}} := \mu^{-1}(\mathfrak{g}_{\text{s-reg}}^*) \subset M$  with moment map  $(\nu_{\text{big}} \circ \mu)|_{M_{\text{s-reg}}}$ . The main result of [2] is the following *abelianization theorem* for generic symplectic quotients of  $M$  by  $G$ : there is a canonical isomorphism

$$M//_{\xi} G \cong M_{\text{s-reg}}//_{\nu_{\text{big}}(\xi)} \mathbb{T}_{\text{big}}$$

of stratified symplectic spaces [12] for each  $\xi \in \mathfrak{g}_{\text{s-reg}}^*$ .

**1.3. A non-abelian Duistermaat–Heckman theorem.** The purpose of this paper is to show that Gelfand–Cetlin data and the abelianization theorem from [2] give rise to a generalization of the Duistermaat–Heckman theorem. We take  $M$  to be a connected symplectic manifold equipped with an effective Hamiltonian  $G$ -action and proper moment map  $\mu : M \rightarrow \mathfrak{g}^*$ . We also fix a *strong* Gelfand–Cetlin datum (Definition 5) on  $\mathfrak{g}^*$ . The continuous map

$$\mu_{\text{big}} := \nu_{\text{big}} \circ \mu : M \rightarrow \mathbb{R}_{\text{big}}$$

is proper and gives rise to a measure

$$\text{DH}_{\text{big}} := (\mu_{\text{big}})_* \lambda_M$$

on  $\mathbb{R}_{\text{big}}$ . On the other hand, normalizing the Haar measure on  $\mathbb{T}_{\text{big}}$  to give a unit volume induces a Lebesgue measure  $\lambda_{\text{big}}$  on  $\mathbb{R}_{\text{big}} = \text{Lie}(\mathbb{T}_{\text{big}})^*$ .

To compare  $\text{DH}_{\text{big}}$  and  $\lambda_{\text{big}}$ , we let  $\mathfrak{g}_{\mu, \text{s-reg}}^*$  denote locus of the regular elements of  $\mu$  lying in  $\mathfrak{g}_{\text{s-reg}}^*$ . The subset

$$\text{U}_{\text{big}} := \nu_{\text{big}}(\mathfrak{g}_{\mu, \text{s-reg}}^*) \subset \mathbb{R}_{\text{big}}$$

is open, and allows us to state the following comparison between  $\text{DH}_{\text{big}}$  and  $\lambda_{\text{big}}$ ; this is the main result of our paper.

**Theorem.** *There is a Radon–Nikodym derivative  $\rho_{\text{big}} : U_{\text{big}} \rightarrow [0, \infty)$  of  $\text{DH}_{\text{big}}$  on  $U_{\text{big}}$  with respect to  $\lambda_{\text{big}}$ , and it satisfies*

$$\rho_{\text{big}}(\lambda_{\text{big}}(\xi)) = \text{vol}(M//_{\xi} G)$$

for all regular values  $\xi \in \mathfrak{g}_{\text{s-reg}}^*$  of  $\mu$ .

If  $G$  is a torus, then one can choose a strong Gelfand–Cetlin datum in such a way that  $\mathfrak{g}_{\text{s-reg}}^* = \mathfrak{g}^*$ ,  $\nu_{\text{big}} : \mathfrak{g}^* \rightarrow \mathbb{R}_{\text{big}}$  is a vector space isomorphism, and  $U_{\text{big}}$  is the locus of regular values of  $\mu$  in  $\mathbb{R}_{\text{big}} = \mathfrak{g}^*$ . This makes it clear that  $\rho_{\text{big}}$  coincides with Duistermaat and Heckman’s Radon–Nikodym derivative on regular values of  $\mu$ . In other words, our result may be interpreted as a generalization of the Duistermaat–Heckman theorem into the realm of non-abelian group actions.

**1.4. Recovering a description of a measure on  $\mathfrak{t}_+^*$ .** Let  $T \subset G$  be a maximal torus with Lie algebra  $\mathfrak{t} \subset \mathfrak{g}$ . By choosing positive roots and a  $G$ -invariant inner product on  $\mathfrak{g}$ , one obtains an inclusion  $\mathfrak{t}^* \subset \mathfrak{g}^*$  and fundamental Weyl chamber  $\mathfrak{t}_+^* \subset \mathfrak{t}^*$ . Restrict the measure  $\lambda_{\mathfrak{t}^*}$  from Section 1.1 to a measure  $\lambda_{\mathfrak{t}_+^*}$  on  $\mathfrak{t}_+^*$ . Let us also consider the adjoint quotient map  $\pi : \mathfrak{g}^* \rightarrow \mathfrak{t}_+^*$ .

Now let  $M$  be a connected symplectic manifold on which  $G$  acts effectively and in a Hamiltonian fashion with proper moment map  $\mu : M \rightarrow \mathfrak{g}^*$ . The measure

$$\text{DH}_{\mathfrak{t}_+^*} := (\pi \circ \mu)_* \lambda_M$$

on  $\mathfrak{t}_+^*$  is another non-abelian variant of the Duistermaat–Heckman measure, and has received some attention in the literature [1, 5, 6, 14].

We use the main result of this paper to prove that  $\text{DH}_{\mathfrak{t}_+^*}$  has a Radon–Nikodym derivative of

$$\rho_{\mathfrak{t}_+^*}(\xi) = \text{vol}(\mathcal{O}_{\xi}) \cdot \text{vol}(M//_{\xi} G)$$

with respect to  $\lambda_{\mathfrak{t}_+^*}$  on a suitable open subset of  $\mathfrak{t}_+^*$ , where  $\mathcal{O}_{\xi} \subset \mathfrak{g}^*$  denotes the coadjoint orbit of  $G$  through  $\xi \in \mathfrak{t}_+^*$  and  $\text{vol}(\mathcal{O}_{\xi})$  denotes its symplectic volume. This fact is presumably well-known and/or readily deducible from existing results.

**1.5. Organization.** The Lie-theoretic foundations of our paper are outlined in Section 2.1. This creates the appropriate context for Section 2.2, in which we recall the pertinent parts of [2]. Strong Gelfand–Cetlin data are then defined and briefly discussed in Section 2.3. In Section 2.4, we give some context for the non-abelian Duistermaat–Heckman measures  $\text{DH}_{\text{big}}$  and  $\text{DH}_{\mathfrak{t}_+^*}$  mentioned above. The Radon–Nikodym derivatives of  $\text{DH}_{\text{big}}$  and  $\text{DH}_{\mathfrak{t}_+^*}$  are then derived in Section 2.5.

**Acknowledgements.** The authors are grateful to Megumi Harada for encouraging us to pursue this work. P.C. is supported by a Utah State University startup grant, while J.W. is supported by Simons Collaboration Grant # 579801.

## 2. MAIN RESULTS

**2.1. Lie theory.** Let  $G$  be a compact connected Lie group of rank  $\ell$  with Lie algebra  $\mathfrak{g}$  and exponential map  $\exp : \mathfrak{g} \rightarrow G$ . Write  $G_{\xi} \subset G$  and  $\mathfrak{g}_{\xi} \subset \mathfrak{g}$  for the centralizers of  $\xi \in \mathfrak{g}^*$  with respect to the coadjoint representations of  $G$  and  $\mathfrak{g}$ , respectively, and  $\mathcal{O}_{\xi} \subset \mathfrak{g}^*$  for the coadjoint orbit of  $G$  through  $\xi$ . The latter centralizer is then the Lie algebra of the former. On the other hand, let us consider the locus of regular elements

$$\mathfrak{g}_{\text{reg}}^* := \{\xi \in \mathfrak{g}^* : \dim \mathfrak{g}_{\xi} = \ell\}.$$

One knows that  $\xi \in \mathfrak{g}^*$  is regular if and only if  $G_\xi$  is a maximal torus of  $G$ . In this case,

$$\Lambda_\xi := \frac{1}{2\pi} \ker \left( \exp|_{\mathfrak{g}_\xi} \right) \subset \mathfrak{g}_\xi$$

is a free  $\mathbb{Z}$ -submodule of rank  $\ell$ .

Fix a maximal torus  $T \subset G$  with Lie algebra  $\mathfrak{t} \subset \mathfrak{g}$ , and choose a set  $\Phi_+ \subset \mathfrak{t}^*$  of positive roots. One thereby obtains the fundamental Weyl chamber

$$\mathfrak{t}_+ := \{x \in \mathfrak{t} : \alpha(x) \geq 0 \text{ for all } \alpha \in \Phi_+\} \subset \mathfrak{t}.$$

Let us also choose a  $G$ -invariant inner product on  $\mathfrak{g}$ , and use it to identify the adjoint and coadjoint representations of  $G$ . This identification allows one to identify  $\mathfrak{t}_+ \subset \mathfrak{g}$  with a fundamental domain  $\mathfrak{t}_+^* \subset \mathfrak{g}^*$  for the coadjoint action. The *sweeping map*  $\pi : \mathfrak{g}^* \rightarrow \mathfrak{t}_+^*$  is then defined by the property that  $\{\pi(\xi)\} = \mathcal{O}_\xi \cap \mathfrak{t}_+^*$  for all  $\xi \in \mathfrak{g}^*$ ; its fibers are precisely the coadjoint orbits of  $G$ .

**2.2. Gelfand–Cetlin data and abelianization.** Let us set

$$u := \frac{1}{2}(\dim \mathfrak{g} - \ell) \quad \text{and} \quad b := \frac{1}{2}(\dim \mathfrak{g} + \ell),$$

and define the three tori

$$\mathbb{T}_{\text{small}} := \mathrm{U}(1)^\ell, \quad \mathbb{T}_{\text{int}} := \mathrm{U}(1)^u, \quad \text{and} \quad \mathbb{T}_{\text{big}} := \mathbb{T}_{\text{small}} \times \mathbb{T}_{\text{int}} \cong \mathrm{U}(1)^b.$$

By identifying  $\mathbb{R}$  with  $i\mathbb{R}$  in the usual way, we may regard

$$\mathbb{R}_{\text{small}} := \mathbb{R}^\ell, \quad \mathbb{R}_{\text{int}} := \mathbb{R}^u, \quad \text{and} \quad \mathbb{R}_{\text{big}} := \mathbb{R}_{\text{small}} \times \mathbb{R}_{\text{int}} \cong \mathbb{R}^b$$

as the duals of the Lie algebras of  $\mathbb{T}_{\text{small}}$ ,  $\mathbb{T}_{\text{int}}$ , and  $\mathbb{T}_{\text{big}}$ , respectively.

The following is a rephrased version of [2, Definition 1].

**Definition 1.** A *Gelfand–Cetlin datum*  $(\nu_{\text{big}}, \mathfrak{g}_{\text{s-reg}}^*)$  consists of a continuous map  $\nu_{\text{big}} = (\nu_1, \dots, \nu_b) : \mathfrak{g}^* \rightarrow \mathbb{R}_{\text{big}}$  and open dense subset  $\mathfrak{g}_{\text{s-reg}}^* \subset \mathfrak{g}_{\text{reg}}^*$  that satisfy the following conditions:

- (i)  $\nu_1, \dots, \nu_\ell$  are  $G$ -invariant on  $\mathfrak{g}^*$  and smooth on  $\mathfrak{g}_{\text{reg}}^*$ ;
- (ii) the set of differentials  $\{d_\xi \nu_1, \dots, d_\xi \nu_\ell\}$  is a  $\mathbb{Z}$ -basis of  $\Lambda_\xi \subset \mathfrak{g}_\xi$  for all  $\xi \in \mathfrak{g}_{\text{reg}}^*$ ;
- (iii)  $\nu_{\text{big}}|_{\mathfrak{g}_{\text{s-reg}}^*} : \mathfrak{g}_{\text{s-reg}}^* \rightarrow \mathbb{R}_{\text{big}}$  is a smooth submersion and moment map for a Poisson Hamiltonian  $\mathbb{T}_{\text{big}}$ -space structure on  $\mathfrak{g}_{\text{s-reg}}^*$ ;
- (iv)  $\nu_{\text{big}}|_{\mathfrak{g}_{\text{s-reg}}^*} : \mathfrak{g}_{\text{s-reg}}^* \rightarrow \nu_{\text{big}}(\mathfrak{g}_{\text{s-reg}}^*)$  is a principal  $\mathbb{T}_{\text{int}}$ -bundle;
- (v) if  $M$  is a Hamiltonian  $G$ -space with moment map  $\mu : M \rightarrow \mathfrak{g}^*$ , then

$$(\nu_{\text{big}} \circ \mu)|_{\mu^{-1}(\mathfrak{g}_{\text{s-reg}}^*)} : \mu^{-1}(\mathfrak{g}_{\text{s-reg}}^*) \rightarrow \mathbb{R}_{\text{big}}$$

is a moment map for a Hamiltonian  $\mathbb{T}_{\text{big}}$ -space structure on  $\mu^{-1}(\mathfrak{g}_{\text{s-reg}}^*)$ .

In this case, we will write

$$\begin{aligned} \nu_{\text{small}} &:= (\lambda_1, \dots, \lambda_\ell) : \mathfrak{g}^* \rightarrow \mathbb{R}_{\text{small}}, & \nu_{\text{int}} &:= (\lambda_{\ell+1}, \dots, \lambda_b) : \mathfrak{g}^* \rightarrow \mathbb{R}_{\text{int}}, \\ M_{\text{s-reg}} &:= \mu^{-1}(\mathfrak{g}_{\text{s-reg}}^*), & \mu_{\text{big}} &:= \nu_{\text{big}} \circ \mu : M \rightarrow \mathbb{R}_{\text{big}}. \end{aligned}$$

**Remark 2.** Suppose that  $G$  is a torus, i.e.  $G = T$  and  $\mathfrak{g} = \mathfrak{t}$ . Let  $\{\nu_1, \dots, \nu_\ell\}$  be a  $\mathbb{Z}$ -basis of  $\frac{1}{2\pi} \ker(\exp : \mathfrak{g} \rightarrow G) \subset \mathfrak{g}$ . It is straightforward to check that  $\nu_{\text{big}} := (\nu_1, \dots, \nu_\ell) : \mathfrak{g}^* \rightarrow \mathbb{R}^\ell$  and  $\mathfrak{g}_{\text{s-reg}}^* := \mathfrak{g}^*$  form a Gelfand–Cetlin datum.

**Remark 3.** The situation is considerably more subtle if one requires  $G$  to be simple. Examples of Gelfand–Cetlin data then include Guillemin and Sternberg’s Gelfand–Cetlin systems on the special unitary and special orthogonal Lie algebras [7, 8], as well as Hoffman and Lane’s recent generalizations of these systems to arbitrary Lie type [9].

The main result in [2] is the following comparison between the symplectic quotients of a Hamiltonian  $G$ -space  $M$  and those of the Hamiltonian  $\mathbb{T}_{\text{big}}$ -space  $M_{\text{s-reg}}$ .

**Theorem 4.** *Let  $(\nu_{\text{big}}, \mathfrak{g}_{\text{s-reg}}^*)$  be a Gelfand–Cetlin datum and  $M$  a Hamiltonian  $G$ -space. Given any  $\xi \in \mathfrak{g}_{\text{s-reg}}^*$ , there is a canonical isomorphism*

$$M //_{\xi} G \cong M_{\text{s-reg}} //_{\nu_{\text{big}}(\xi)} \mathbb{T}_{\text{big}}$$

of stratified symplectic spaces.

The definition of a *stratified symplectic space* is due to Sjamaar–Lerman [12]. This strict generalization of a symplectic manifold liberates us from making assumptions to ensure that the symplectic quotients  $M //_{\xi} G$  and  $M_{\text{s-reg}} //_{\nu_{\text{big}}(\xi)} \mathbb{T}_{\text{big}}$  are genuine symplectic manifolds.

**2.3. Strong Gelfand–Cetlin data.** In order to prove the main results of this manuscript, we will need the following strengthening of Definition 1.

**Definition 5.** A Gelfand–Cetlin datum  $(\nu_{\text{big}}, \mathfrak{g}_{\text{s-reg}}^*)$  will be called *strong* if it satisfies the following additional conditions:

- (vi) if  $\xi, \eta \in \mathfrak{g}^*$  and  $\nu_{\text{small}}(\xi) = \nu_{\text{small}}(\eta)$ , then  $\mathcal{O}_{\xi} = \mathcal{O}_{\eta}$ ;
- (vii)  $\nu_{\text{big}}$  is proper;
- (viii)  $\mathfrak{g}_{\text{s-reg}}^*$  is a union of fibers of  $\nu_{\text{big}}$ ;
- (ix) if  $\xi \in \mathfrak{g}_{\text{s-reg}}^*$ , then  $\mathcal{O}_{\xi} \cap \mathfrak{g}_{\text{s-reg}}^*$  is dense in  $\mathcal{O}_{\xi}$ .

**Remark 6.** Examples of strong Gelfand–Cetlin data include the data described in Remark 2, as well as Guillemin and Sternberg’s Gelfand–Cetlin systems on the special unitary and special orthogonal Lie algebras [7, 8].

**2.4. Non-abelian Duistermaat–Heckman measures.** Let  $M$  be a connected symplectic manifold carrying an effective Hamiltonian action of  $G$  and proper moment map  $\mu : M \rightarrow \mathfrak{g}^*$ . Define the  $G$ -invariant moment map to be the composition

$$\bar{\mu} := \pi \circ \mu : M \rightarrow \mathfrak{t}_+^*.$$

Let us also suppose that  $(\nu_{\text{big}}, \mathfrak{g}_{\text{s-reg}}^*)$  is a strong Gelfand–Cetlin datum. The following five measures naturally arise in this context:

- the measure  $\lambda_{\mathfrak{t}_+^*}$  on  $\mathfrak{t}_+^*$  induced by normalizing the Haar measure on  $T$  to give a unit volume, forming the resulting Lebesgue measure on  $\mathfrak{t}$ , taking the induced Lebesgue measure on  $\mathfrak{t}^*$ , and restricting to  $\mathfrak{t}_+^*$ ;
- the measure  $\lambda_{\mathbb{R}_{\text{big}}}$  on  $\mathbb{R}_{\text{big}}$  obtained by normalizing the Haar measure on  $U(1)^{\text{b}}$ , and proceeding analogously to the above to get a measure on  $\mathbb{R}_{\text{big}} = \text{Lie}(U(1)^{\text{b}})^*$ ;
- the Liouville measure  $\lambda_M$  on  $M$  associated to the volume form  $\frac{1}{n!}\omega^n$ , where  $\omega$  is the symplectic form on  $M$  and  $2n = \dim M$ ;
- the measure on  $\mathfrak{t}_+^*$  obtained by pushing  $\lambda_M$  forward along  $\bar{\mu} : M \rightarrow \mathfrak{t}_+^*$ , i.e.  $\text{DH}_{\mathfrak{t}_+^*} := \bar{\mu}_* \lambda_M$ ;
- the measure on  $\mathbb{R}_{\text{big}}$  obtained by pushing  $\lambda_M$  forward along  $\mu_{\text{big}} : M \rightarrow \mathbb{R}_{\text{big}}$ , i.e.  $\text{DH}_{\mathbb{R}_{\text{big}}} := (\mu_{\text{big}})_* \lambda_M$ ;

**Remark 7.** Suppose that  $G = T$  is a torus, and let all objects and notation be as discussed in Remark 2. Observe that  $\mathfrak{t}_+^* = \mathfrak{g}^*$ ,  $\mathbb{R}_{\text{big}} = \mathbb{R}^\ell$ , and that  $\nu_{\text{big}} : \mathfrak{g}^* \rightarrow \mathbb{R}_{\text{big}}$  is a vector space isomorphism. By means of the previous sentence, we may regard both  $\text{DH}_{\mathfrak{t}_+^*}$  and  $\text{DH}_{\text{big}}$  as measures on  $\mathbb{R}^\ell$ . These two measures are readily seen to coincide. This coincident measure is the subject of Duistermaat and Heckman’s work [4], and so often called the *Duistermaat–Heckman measure*.

The topological spaces  $\mathfrak{t}_+^*$  and  $\mathbb{R}_{\text{big}}$  are homeomorphic if and only if  $G$  is a torus. In particular, the measures discussed in Remark 7 are defined on different spaces in the case of our arbitrary compact connected Lie group  $G$ . One may therefore regard  $\text{DH}_{\mathfrak{t}_+^*}$  and  $\text{DH}_{\text{big}}$  as two different generalizations of the classical Duistermaat–Heckman measure, each into the realm of Hamiltonian actions by non-abelian groups. This motivates the following definition.

**Definition 8.** Let  $M$  be a connected Hamiltonian  $G$ -space with an effective  $G$ -action and proper moment map  $\mu : M \rightarrow \mathfrak{g}^*$ .

- (i) We call  $\text{DH}_{\mathfrak{t}_+^*} := \overline{\mu}_* \lambda_M$  the *non-abelian Duistermaat–Heckman measure* on  $\mathfrak{t}_+^*$  associated to  $M$ .
- (ii) Suppose that  $(\nu_{\text{big}}, \mathfrak{g}_{\text{s-reg}}^*)$  is a strong Gelfand–Cetlin datum. We call  $\text{DH}_{\text{big}} := (\mu_{\text{big}})_* \lambda_M$  the *non-abelian Duistermaat–Heckman measure* on  $\mathbb{R}_{\text{big}}$  associated to  $M$  and  $(\nu_{\text{big}}, \mathfrak{g}_{\text{s-reg}}^*)$ .

The measure  $\text{DH}_{\mathfrak{t}_+^*}$  and variants thereof have been studied quite extensively [1, 3, 5, 6, 10, 11, 13, 14].

**2.5. Radon–Nikodym derivatives.** Let  $(\nu_{\text{big}}, \mathfrak{g}_{\text{s-reg}}^*)$  be a strong Gelfand–Cetlin datum, and  $M$  a connected Hamiltonian  $G$ -space with proper moment map  $\mu : M \rightarrow \mathfrak{g}^*$  and effective  $G$ -action. Consider the open subset

$$\mathfrak{g}_{\mu, \text{s-reg}}^* := \{\xi \in \mathfrak{g}_{\text{s-reg}}^* : d_m \mu \text{ is surjective for all } m \in \mu^{-1}(\xi)\}$$

of regular values of  $\mu$  lying in  $\mathfrak{g}_{\text{s-reg}}^*$ .

**Lemma 9.** *The subset  $\mathfrak{g}_{\mu, \text{s-reg}}^*$  is a union of fibers of  $\nu_{\text{big}}$ . Each such fiber is contained in a regular coadjoint orbit of  $G$ .*

*Proof.* Let  $\xi \in \mathfrak{g}^*$  and  $\eta \in \mathfrak{g}_{\mu, \text{s-reg}}^*$  be such that  $\nu_{\text{big}}(\xi) = \nu_{\text{big}}(\eta)$ . Definition 5(viii) implies that  $\xi \in \mathfrak{g}_{\text{s-reg}}^*$ , while Definition 5(vi) tells us that  $\mathcal{O}_\xi = \mathcal{O}_\eta$ . Since  $\mu$  is  $G$ -equivariant and  $\eta$  is a regular value of  $\mu$ , the condition  $\mathcal{O}_\xi = \mathcal{O}_\eta$  forces  $\xi$  to be a regular value of  $\mu$ . We conclude that  $\xi \in \mathfrak{g}_{\mu, \text{s-reg}}^*$ . Our arguments also imply that  $\nu_{\text{big}}^{-1}(\nu_{\text{big}}(\eta)) \subset \mathcal{O}_\eta$ . These last two sentences suffice to complete the proof.  $\square$

Now consider the image

$$U_{\text{big}} := \nu_{\text{big}}(\mathfrak{g}_{\mu, \text{s-reg}}^*)$$

of the open subset  $\mathfrak{g}_{\mu, \text{s-reg}}^* \subset \mathfrak{g}_{\text{s-reg}}^*$  under  $\nu_{\text{big}}$ ; it is open in  $\mathbb{R}_{\text{big}}$  by Definition 1(iii). Given any  $x \in U_{\text{big}}$  and  $\xi, \eta \in \nu_{\text{big}}^{-1}(x)$ , Lemma 9 and the  $G$ -equivariance of  $\mu$  imply that  $M//_\xi G \cong M//_\eta G$  as symplectic orbifolds. It follows that the symplectic volume

$$\rho_{\text{big}}(x) := \text{vol}(M//_\xi G)$$

does not depend on the choice of  $\xi \in \nu_{\text{big}}^{-1}(x)$ . We thereby obtain a well-defined function  $\rho_{\text{big}} : U_{\text{big}} \rightarrow [0, \infty)$ ; it will turn out to be a Radon–Nikodym derivative of the non-abelian Duistermaat–Heckman measure  $\text{DH}_{\text{big}}$  with respect to  $\lambda_{\text{big}}$ , in the following sense.

**Theorem 10.** *The function  $\rho_{\text{big}}$  is a Radon–Nikodym derivative of  $\text{DH}_{\text{big}}$  on the open subset  $U_{\text{big}} \subset \mathbb{R}_{\text{big}}$  with respect to  $\lambda_{\text{big}}$ , i.e.  $\text{DH}_{\text{big}} = \rho_{\text{big}} \lambda_{\text{big}}$  on  $U_{\text{big}}$ .*

*Proof.* The properness of  $\mu : M \rightarrow \mathfrak{g}$  implies that of its pullback  $\mu^{-1}(\mathfrak{g}_{\mu, \text{s-reg}}^*) \rightarrow \mathfrak{g}_{\mu, \text{s-reg}}^*$  to  $\mathfrak{g}_{\mu, \text{s-reg}}^* \subset \mathfrak{g}^*$ . It is also clear that  $\mu^{-1}(\mathfrak{g}_{\mu, \text{s-reg}}^*) \rightarrow \mathfrak{g}_{\mu, \text{s-reg}}^*$  is a submersion. On the other hand, Definition 1(iv) and Lemma 9 tell us that  $\nu_{\text{big}}$  restricts to a proper submersion  $\mathfrak{g}_{\mu, \text{s-reg}}^* \rightarrow U_{\text{big}}$ . The last few sentences force the composite map

$$\mu^{-1}(\mathfrak{g}_{\mu, \text{s-reg}}^*) \rightarrow \mathfrak{g}_{\mu, \text{s-reg}}^* \rightarrow U_{\text{big}}$$

to be a proper submersion  $\mu^{-1}(\mathfrak{g}_{\mu, \text{s-reg}}^*) \rightarrow U_{\text{big}}$ . This proper submersion is precisely the pullback of  $\mu_{\text{big}} : M \rightarrow \mathbb{R}_{\text{big}}$  to  $U_{\text{big}} \subset \mathbb{R}_{\text{big}}$ . It is also the pullback of the  $\mathbb{T}_{\text{big}}$ -moment map  $\mu_{\text{big}}|_{M_{\text{s-reg}}} : M_{\text{s-reg}} \rightarrow \mathbb{R}_{\text{big}}$  to  $U_{\text{big}}$ . The local normal forms of moment maps used in the proof of the Duistermaat–Heckman theorem [4, Section 3] now yield

$$\text{DH}_{\text{big}} = (\mu_{\text{big}})_* \lambda_M = \phi \lambda_{\text{big}}$$

on  $U_{\text{big}}$ , where  $\phi : U_{\text{big}} \rightarrow [0, \infty)$  is defined by  $\phi(\eta) = \text{vol}(M_{\text{s-reg}} //_{\eta} \mathbb{T}_{\text{big}})$ . Theorem 4 and the definition of  $\rho_{\text{big}}$  immediately imply that  $\phi = \rho_{\text{big}}$ , completing the proof.  $\square$

Consider the pullback  $\pi_{\text{reg}} : \mathfrak{g}_{\text{reg}}^* \rightarrow (\mathfrak{t}_+^*)^\circ$  of  $\pi : \mathfrak{g}^* \rightarrow \mathfrak{t}_+^*$  to the interior  $(\mathfrak{t}_+^*)^\circ$  of  $\mathfrak{t}_+^*$ . Since  $\pi_{\text{reg}}$  is a submersion, the image

$$U_{\mathfrak{t}_+^*} := \pi(\mathfrak{g}_{\mu, \text{s-reg}}^*)$$

is an open subset of  $(\mathfrak{t}_+^*)^\circ$ . It is also clear that  $U_{\mathfrak{t}_+^*}$  consists of regular values of  $\mu$ . We may therefore define  $\rho_{\mathfrak{t}_+^*} : U_{\mathfrak{t}_+^*} \rightarrow [0, \infty)$  by

$$\rho_{\mathfrak{t}_+^*}(\xi) := \text{vol}(\mathcal{O}_\xi) \cdot \text{vol}(M //_\xi G),$$

where  $\text{vol}(\mathcal{O}_\xi)$  denotes the symplectic volume of  $\mathcal{O}_\xi$ .

**Corollary 11.** *The function  $\rho_{\mathfrak{t}_+^*}$  is a Radon–Nikodym derivative of  $\text{DH}_{\mathfrak{t}_+^*}$  on the open subset  $U_{\mathfrak{t}_+^*} \subset \mathfrak{t}_+^*$  with respect to  $\lambda_{\mathfrak{t}_+^*}$ , i.e.  $\text{DH}_{\mathfrak{t}_+^*} = \rho_{\mathfrak{t}_+^*} \lambda_{\mathfrak{t}_+^*}$  on  $U_{\mathfrak{t}_+^*}$ .*

*Proof.* Consider the pullbacks  $\bar{\mu}^{-1}(U_{\mathfrak{t}_+^*}) \rightarrow \pi^{-1}(U_{\mathfrak{t}_+^*})$  and  $\pi^{-1}(U_{\mathfrak{t}_+^*}) \rightarrow U_{\mathfrak{t}_+^*}$  of  $\mu : M \rightarrow \mathfrak{g}^*$  and  $\pi : \mathfrak{g}^* \rightarrow \mathfrak{t}_+^*$  to the open subsets  $\pi^{-1}(U_{\mathfrak{t}_+^*}) \subset \mathfrak{g}^*$  and  $U_{\mathfrak{t}_+^*} \subset \mathfrak{t}_+^*$ , respectively. The latter pullback map is evidently a proper submersion, while the former is clearly proper. To see that the former is also a submersion, recall that  $U_{\mathfrak{t}_+^*}$  consists of regular values of  $\mu$ . Let us also observe that  $\pi^{-1}(U_{\mathfrak{t}_+^*})$  is the union of the adjoint orbits through the elements of  $U_{\mathfrak{t}_+^*}$ . The  $G$ -equivariance of  $\mu$  then forces each element of  $\pi^{-1}(U_{\mathfrak{t}_+^*})$  to be a regular value of  $\mu$ . In particular,  $\bar{\mu}^{-1}(U_{\mathfrak{t}_+^*}) \rightarrow \pi^{-1}(U_{\mathfrak{t}_+^*})$  is also a proper submersion. The composite map

$$\bar{\mu}^{-1}(U_{\mathfrak{t}_+^*}) \rightarrow \pi^{-1}(U_{\mathfrak{t}_+^*}) \rightarrow U_{\mathfrak{t}_+^*}$$

is therefore a proper submersion. This amounts to stating that  $\bar{\mu}$  restricts to a proper submersion  $\bar{\mu}^{-1}(U_{\mathfrak{t}_+^*}) \rightarrow U_{\mathfrak{t}_+^*}$ .

Let us define a map  $\varphi : \nu_{\text{big}}(\pi^{-1}(U_{\mathfrak{t}_+^*})) \rightarrow U_{\mathfrak{t}_+^*}$  as follows:  $\varphi(x_1, \dots, x_b)$  is the unique element of  $U_{\mathfrak{t}_+^*}$  satisfying  $f_i(\varphi(x_1, \dots, x_b)) = x_i$  for all  $i \in \{1, \dots, \ell\}$ . This gives rise to a

commutative diagram

$$\begin{array}{ccc}
 & \nu_{\text{big}}(\pi^{-1}(U_{\text{t}_+^*})) & \\
 \nu_{\text{big}} \nearrow & & \searrow \varphi \\
 \pi^{-1}(U_{\text{t}_+^*}) & \xrightarrow{\pi} & U_{\text{t}_+^*} \\
 \mu \nwarrow & & \nearrow \bar{\mu} \\
 & \bar{\mu}^{-1}(U_{\text{t}_+^*}) &
 \end{array}$$

At the same time, Definition 5(ix) implies that the open subset  $\mathfrak{g}_{\mu, \text{s-reg}}^* \subset \pi^{-1}(U_{\text{t}_+^*})$  is dense. It follows that  $U_{\text{big}} = \nu_{\text{big}}(\mathfrak{g}_{\mu, \text{s-reg}}^*)$  is open and dense in  $\nu_{\text{big}}(\pi^{-1}(U_{\text{t}_+^*}))$ . We conclude that  $\text{DH}_{\text{t}_+^*}|_{U_{\text{t}_+^*}}$  is the push-forward of  $\text{DH}_{\text{big}}|_{U_{\text{big}}}$  along  $\varphi|_{U_{\text{big}}}$ . Theorem 10 now reduces us to proving that the fiber of  $\varphi|_{U_{\text{big}}}$  over each  $\xi \in U_{\text{t}_+^*}$  has Euclidean volume equal to the symplectic volume of  $\mathcal{O}_\xi$ . To this end, we have

$$\begin{aligned}
 \varphi|_{U_{\text{big}}}^{-1}(\xi) &= \varphi^{-1}(\xi) \cap U_{\text{big}} \\
 &= \{(f_1(\xi), \dots, f_\ell(\xi))\} \times \nu_{\text{int}}(\mathcal{O}_\xi \cap \mathfrak{g}_{\mu, \text{s-reg}}^*) \subset \mathbb{R}_{\text{small}} \times \mathbb{R}_{\text{int}} = \mathbb{R}_{\text{big}} \\
 &= \{(f_1(\xi), \dots, f_\ell(\xi))\} \times \nu_{\text{int}}(\mathcal{O}_\xi \cap \mathfrak{g}_{\text{s-reg}}^*) \subset \mathbb{R}_{\text{small}} \times \mathbb{R}_{\text{int}} = \mathbb{R}_{\text{big}}
 \end{aligned}$$

for all  $\xi \in U_{\text{t}_+^*}$ . It now suffices to observe that the Euclidean volume of  $\nu_{\text{int}}(\mathcal{O}_\xi \cap \mathfrak{g}_{\text{s-reg}}^*) \subset \mathbb{R}_{\text{int}}$  coincides with the symplectic volume of  $\mathcal{O}_\xi$ , as follows from Definition 5(ix) and [2, Proposition 3].  $\square$

## REFERENCES

- [1] CHRISTANDL, M., DORAN, B., KOUSIDIS, S., AND WALTER, M. Eigenvalue distributions of reduced density matrices. *Comm. Math. Phys.* 332, 1 (2014), 1–52.
- [2] CROOKS, P., AND WEITSMAN, J. Gelfand–Cetlin abelianizations of symplectic quotients. arXiv:2209.04978 (2022), 14pp.
- [3] DUISTERMAAT, J. J. On global action-angle coordinates. *Comm. Pure Appl. Math.* 33, 6 (1980), 687–706.
- [4] DUISTERMAAT, J. J., AND HECKMAN, G. J. On the variation in the cohomology of the symplectic form of the reduced phase space. *Invent. Math.* 69, 2 (1982), 259–268.
- [5] GUILLEMIN, V., LERMAN, E., AND STERNBERG, S. *Symplectic fibrations and multiplicity diagrams*. Cambridge University Press, Cambridge, 1996.
- [6] GUILLEMIN, V., AND PRATO, E. Heckman, Kostant, and Steinberg formulas for symplectic manifolds. *Adv. Math.* 82, 2 (1990), 160–179.
- [7] GUILLEMIN, V., AND STERNBERG, S. The Gel’fand–Cetlin system and quantization of the complex flag manifolds. *J. Funct. Anal.* 52, 1 (1983), 106–128.
- [8] GUILLEMIN, V., AND STERNBERG, S. On collective complete integrability according to the method of Thimm. *Ergodic Theory Dynam. Systems* 3, 2 (1983), 219–230.
- [9] HOFFMAN, B., AND LANE, J. Stratified gradient Hamiltonian vector fields and collective integrable systems. arXiv:2008.13656 (2022), 44pp.
- [10] JEFFREY, L. C., AND KIRWAN, F. C. Localization for nonabelian group actions. *Topology* 34, 2 (1995), 291–327.
- [11] PRATO, E., AND WU, S. Duistermaat–Heckman measures in a non-compact setting. *Compositio Math.* 94, 2 (1994), 113–128.
- [12] SJAMAAR, R., AND LERMAN, E. Stratified symplectic spaces and reduction. *Ann. of Math. (2)* 134, 2 (1991), 375–422.
- [13] WITTEN, E. Two-dimensional gauge theories revisited. *J. Geom. Phys.* 9, 4 (1992), 303–368.
- [14] ZHANG, L., JIANG, Y., AND WU, J. Duistermaat–Heckman measure and the mixture of quantum states. *J. Phys. A* 52, 49 (2019), 495203, 42.



(Peter Crooks) DEPARTMENT OF MATHEMATICS AND STATISTICS, UTAH STATE UNIVERSITY, 3900 OLD MAIN HILL, LOGAN, UT 84322, USA

*Email address:* `peter.crooks@usu.edu`

(Jonathan Weitsman) DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, 360 HUNTINGTON AVENUE, BOSTON, MA 02115, USA

*Email address:* `j.weitsman@northeastern.edu`