Extremal graphs for the odd prism^{*}

Xiaocong He, Yongtao Li^{*}, Lihua Feng

School of Mathematics and Statistics, HNP-LAMA, Central South University Changsha, Hunan, 410083, PR China

February 6, 2024

Abstract: The Turán number ex(n, H) of a graph H is the maximum number of edges in an n-vertex graph which does not contain H as a subgraph. The Turán number of regular polyhedrons was widely studied in a series of works due to Simonovits. In this paper, we shall present the exact Turán number of the prism C_{2k+1}^{\Box} , which is defined as the Cartesian product of an odd cycle C_{2k+1} and an edge K_2 . Applying a deep theorem of Simonovits and a stability result of Yuan [European J. Combin. 104 (2022)], we shall determine the exact value of $ex(n, C_{2k+1}^{\Box})$ for every $k \ge 1$ and sufficiently large n, and we also characterize the extremal graphs. Moreover, in the case of k = 1, motivated by a recent result of Xiao, Katona, Xiao and Zamora [Discrete Appl. Math. 307 (2022)], we will determine the exact value of $ex(n, C_3^{\Box})$ for every n instead of for sufficiently large n.

Keywords: Turán number, Extremal graphs, Cartesian product.

AMS classification: 05C35.

1 Introduction

In this paper, all graphs considered are undirected, finite and contain neither loops nor multiple edges. The vertex set of a graph G is denoted by V(G), the edge set of G by E(G), and the number of edges in G by e(G). Let K_n be the complete graph on n vertices, and $K_{s,t}$ be the complete bipartite graph with parts of sizes s and t. We write C_n and P_n for the cycle and path on n vertices, respectively. The Turán graph $T_r(n)$ is an n-vertex complete r-partite graph with each part of size n_i such that $|n_i - n_j| \leq 1$ for all $1 \leq i, j \leq r$, that is, n_i equals $\lfloor n/r \rfloor$ or $\lceil n/r \rceil$. Sometimes, we call $T_r(n)$ the balanced complete r-partite graph. Denote by $G_1 \cup G_2$ the vertex-disjoint union of two graphs G_1 and G_2 . For simplicity, we write kG for the vertex-disjoint union of k copies of G. Denote by $G_1 \otimes G_2$ the graph obtained from $G_1 \cup G_2$ by joining each vertex of G_1 to each vertex of G_2 . For a vertex $v \in V(G)$, and a subgraph $H \subseteq G$ (possibly $v \in V(H)$), we use $N_H(v)$ to denote the set of neighbors of v in V(H), and use $d_H(v)$ to denote $|N_H(v)|$. For two sets $A, B \subseteq V(G)$, we write G[A, B] for the induced bipartite graph of G with edges between A and B, and e(A, B)for the number of edges of G[A, B]. We write G[A] for the subgraph induced by A. We denote the minimum degree of G by $\delta(G)$, the complement of G by \overline{G} . In particular, \overline{K}_n is the empty graph, which is also called an independent set on n vertices. The chromatic number of G, denoted by $\chi(G)$, which is the minimum integer s such that there exists a coloring of the vertex set V(G)with s colors and the adjacent vertices have different colors.

^{*}E-mail addresses: hexc2018@qq.com (X. He), ytli0921@hnu.edu.cn (Y. Li, corresponding author), fenglh@163.com (L. Feng).

1.1 Turán number of regular polyhedrons

Let G and H be two simple graphs. We say that G is H-free if it does not contain a subgraph isomorphic to H. For example, every bipartite graph is triangle-free. The Turán number of H, denoted by ex(n, H), is the maximum number of edges in an n-vertex H-free graph. Moreover, an n-vertex H-free graph with maximum number of edges is called an *extremal graph* for H. In general, it is possible that there are many (not only one) extremal graphs for H. For instance, our result in this article shares this phenomenon. The problem of determining Turán numbers is one of the cornerstones of graph theory. Observe that the balanced complete r-partite graph $T_r(n)$ is K_{r+1} -free. In 1941, Turán [48] proved $ex(n, K_{r+1}) = e(T_r(n))$. Moreover, $T_r(n)$ is the unique K_{r+1} -free graph attaining the maximum number of edges. In particular, for r = 2, it gives

$$\operatorname{ex}(n, K_3) = \lfloor n^2/4 \rfloor,\tag{1}$$

which was early proved by Mantel [39]; see, e.g., the monograph [3, p. 294] for alternative proofs and extensions, and the surveys [27, 46] for more related results.

After Turán [48] determined the extremal graphs of the tetrahedron K_4 , he asked the following analogous problem: Determining the Turán number of a graph consisting of the vertices and edges of a regular polyhedron. For example, the cube Q_8 , the octahedron O_6 , the dodecahedron D_{20} and the icosahedron I_{12} ; see Figure 1.



Figure 1: The five regular polyhedrons

The Turán problem involving regular polyhedrons was widely studied in the literature; see, e.g., [19, 26, 32, 33, 41] for the cube Q_8 , [20] for the octahedron O_6 , [43] for the dodecahedron D_{20} , and [44] for the icosahedron I_{12} . In particular, although Erdős and Simonovits [19] showed that $ex(n, Q_8) \leq Cn^{8/5}$ for some constant C > 0, the lower bound on the Turán number of Q_8 remains open. Motivated by the above results, we shall consider the Turán problem of the prism, which can be viewed as a variant graph of the cube Q_8 . Recall that $\chi(H)$ denotes the chromatic number of H. The celebrated Erdős–Stone–Simonovits Theorem [17, 18] states that

$$ex(n,H) = \left(1 - \frac{1}{\chi(H) - 1}\right) {n \choose 2} + o(n^2).$$
 (2)

This result determines the asymptotic value of ex(n, H) for every non-bipartite graph H, and it reduces to $ex(n, H) = o(n^2)$ for every bipartite graph H. At the first glance, the formula (2) gives a satisfactory answer to the problem of estimating ex(n, H). Furthermore, it is an important problem in extremal combinatorics to obtain more accurate bound on the error term $o(n^2)$. In particular, one of the central problems is to determine the exact Turán number for a graph H.

There are many elegant results on finding the exact value of ex(n, H) for a specific graph H. For example, the earliest result due to Mantel [39] provided $ex(n, C_3) = \lfloor n^2/4 \rfloor$. Erdős, Füredi, Gould and Gunderson [15], Chen, Gould, Pfender and Wei [8] determined the exact Turán number for intersecting triangles and cliques, respectively. Liu [37] determined the Turán number for the edge-blow-ups of cycles and a large class of trees, and Ni, Kang, Shan and Zhu [40] investigated the edge-blow-ups of keyrings. In 2022, Yuan [54] studied the Turán number and extremal graphs for the edge-blow-ups of complete graphs and complete bipartite graphs. Applying the stability theorem of Erdős and Simonovits, Hou, Qiu and Liu [30, 31], and Yuan [52] independently determined the Turán number for flower graphs. Moreover, Dzido [12], Dzido and Jastrzebski [13], and Yuan [53] determined the Turán number for wheels. In 2022, Xiao, Katona, Xiao and Zamora [49], and Yuan [55] characterized the extremal graphs for the power of a path.

1.2 The prism and Cartesian product

Apart from the five regular polyhedrons in Figure 1, there are two well-known polyhedrons, called the pyramid and prism; see Figure 2. In fact, it is easy to observe that the pyramid is actually a wheel graph, which was studied by Dzido [12], Dzido and Jastrzebski [13] and Yuan [53]. Differing from the study of the pyramid (wheel), in this paper, we shall pay attention to the Turán number of the prism, and characterize the extremal graphs.



Figure 2: The quadrangular pyramid and triangular prism

One can consider the prism from the perspective of Cartesian product. The Cartesian product of graphs G and F, denoted by $G \Box F$, has vertex set $V(G) \times V(F)$, in which two distinct vertices (g_1, f_1) and (g_2, f_2) are adjacent in $G \Box F$ if either $g_1 = g_2$ and $f_1 f_2 \in E(F)$, or $f_1 = f_2$ and $g_1 g_2 \in E(G)$. In 2022, Bradač, Janzer, Sudakov and Tomon [4] studied the Turán number of the grid $P_t \Box P_t$, where P_t is the path on t vertices. More precisely, they proved that for $t \geq 2$, there exist positive numbers C_1 and C_2 depending only on t such that $C_1 n^{3/2} \leq \exp(n, P_t \Box P_t) \leq C_2 n^{3/2}$. In our paper, we shall consider the Turán number for the Cartesian product of an odd cycle and an edge. Recall that C_{2k+1} is the cycle on 2k + 1 vertices. Observe that the odd prism $C_{2k+1}^{\Box} := C_{2k+1} \Box K_2$ is the graph consisting of two vertex disjoint cycles of order 2k + 1 and a matching pairing the corresponding vertices of these two cycles. For example, the third and fourth graph in Figure 2 are the triangular prism. As promised, we shall study the Turán number of the prism C_{2k+1}^{\Box} . By Erdős–Stone–Simonovits' theorem in (2), we get an asymptotic value

$$ex(n, C_{2k+1}^{\Box}) = \frac{n^2}{4} + o(n^2)$$

Furthermore, one natural problem is to refine the error term and determine the exact Turán number of C_{2k+1}^{\Box} . Our result in this paper solves this problem and study the exact value of $ex(n, C_{2k+1}^{\Box})$. Moreover, we also characterize the extremal graphs for C_{2k+1}^{\Box} .

1.3 Main results

The first result in this paper can be stated as below.

Theorem 1.1. Let $k \ge 1$ be fixed and n be sufficiently large. Then

$$\exp(n, C_{2k+1}^{\square}) = \max_{n_a+n_b=n} \left\{ n_a(1+n_b) + \frac{1}{2}(j^2 - 3j) : j \in \{0, 1, 2\}, j \equiv n_a \pmod{3} \right\}.$$

Moreover, all extremal graphs for C_{2k+1}^{\square} are of the form of a complete bipartite graph K_{n_a,n_b} with an extremal graph for P_4 added to the part of size n_a .

There are a few kinds of graphs H for which ex(n, H) are known exactly for all integers n, including cliques, matchings, paths, odd cycles and some other special graphs. To be more specific, Turán [48] determined the value of $ex(n, K_{r+1})$ for all n, Erdős and Gallai [16] determined $ex(n, P_k)$ and $ex(n, M_{2k})$, where M_{2k} is the matching on 2k vertices. Füredi and Gunderson [23] determined $ex(n, C_{2k+1})$ for all n and $k \geq 2$. Bushaw and Kettle [6] determined $ex(n, kP_3)$ for all $n \geq 7k$, and later, Campos and Lopes [7], Yuan and Zhang [50], independently determined the value $ex(n, kP_3)$ for all integers n and k. Moreover, Bielak and Kieliszek [2] determined $ex(n, 2P_5)$ for all n, which was also proved by Yuan and Zhang [51] independently. In addition, Lan, Qin and Shi [36] determined $ex(n, 2P_7)$ for all n.

For the triangular prism in Figure 2, that is, the case k = 1 in Theorem 1.1, we will use a different method, and get rid of the condition that n is sufficiently large. More precisely, we will determine $ex(n, C_3^{\Box})$ for all n, and present all C_3^{\Box} -free graphs on n vertices with maximum number of edges. Our results will be presented in the following two theorems, where we provide the Turán number first, and give the corresponding extremal graphs later.

Theorem 1.2. The maximum number of edges in an n-vertex C_3^{\Box} -free graph $(n \neq 5)$ is

$$\operatorname{ex}(n, C_3^{\Box}) = \begin{cases} \lfloor \frac{n^2}{4} \rfloor + \lfloor \frac{n-1}{2} \rfloor, & \text{if } n \equiv 1, 2, 3 \pmod{6}; \\ \lfloor \frac{n^2}{4} \rfloor + \lceil \frac{n}{2} \rceil, & \text{otherwise.} \end{cases}$$

Before showing the extremal graphs in Theorem 1.2, we need to define some graphs, which originally appears in Xiao, Katona, Xiao and Zamora [49]. Let $K_{i,n-i}$ be the complete bipartite graph on parts X and Y with |X| = i and |Y| = n - i, respectively. On the one hand, if $3 \mid i$, then embedding $\frac{i}{3}$ vertex-disjoint triangles in the class X and keeping the class Y unchanged, we get a new graph on n vertices, and denoted by H_n^i . On the other hand, if $3 \nmid i$, then choosing an integer $j \in [1, i]$ such that $3 \mid (i - j)$, we define $F_n^{i,j}$ as the graph obtained from $K_{i,n-i}$ by embedding a star on j vertices, and $\frac{i-j}{3}$ vertex-disjoint triangles in the class X, where the star is disjoint from any triangle. Clearly, one can observe that both H_n^i and $F_n^{i,j}$ are C_{2k+1}^{\Box} -free for any i and j. In addition, for small n, we introduce three exceptional graphs G_1, G_2 and G_3 ; see Figure 3.



Figure 3: The graphs G_1, G_2 and G_3 .

The extremal graphs for C_3^{\Box} are presented in the following tables. **Theorem 1.3.** For $n \ge 6$, the extremal graphs for C_3^{\Box} are the following ones.

order n	6	7	8
extremal graphs	G_1, H_6^3	$G_2, F_7^{4,1}, F_7^{4,4}, H_7^3$	$G_3, F_8^{4,1}, F_8^{4,4}, F_8^{5,2}, F_8^{5,5}$

Moreover, for $n \geq 9$, we have the following extremal graphs.

order $n \pmod{6}$	0	1	2	3	4	5
extremal graphs	$H_n^{\frac{n}{2}}$	$F_n^{\left\lceil \frac{n}{2} \right\rceil,j}, H_n^{\left\lfloor \frac{n}{2} \right\rfloor}$	$F_n^{\frac{n}{2},j}, F_n^{\frac{n}{2}+1,j}$	$F_n^{\left\lceil \frac{n}{2} \right\rceil,j}, H_n^{\left\lceil \frac{n}{2} \right\rceil+1}$	$H_n^{\frac{n}{2}+1}$	$H_n^{\left\lceil \frac{n}{2} \right\rceil}$

where j can be chosen as any possible integer satisfying $1 \le j \le i$ and $3 \mid (i-j)$.

Outline of the paper. This paper is organized as follows. In Section 2, we shall present some preliminaries, including the Turán-type results on paths, and a deep structure theorem of Simonovits for graphs whose decomposition family contains a linear forest. In Section 3, we will give the proof of Theorem 1.1. In Section 4, in order to determine the *n*-vertex extremal graphs for C_3^{\Box} for every $n \ge 6$, we will provide some lemmas for our purpose. Later, the proofs of Theorems 1.2 and 1.3 are given in Section 5, respectively. Finally, some open problems are proposed in Section 6.

2 Preliminaries

2.1 The Turán-type results for paths

To begin with, we need to introduce a result of Erdős and Gallai [16].

Theorem 2.1 (Erdős and Gallai [16]). The maximum number of edges in an n-vertex P_k -free graph is $\frac{n(k-2)}{2}$, that is, $ex(n, P_k) \leq \frac{n(k-2)}{2}$, with equality if and only if (k-1)|n and the graph is a vertex disjoint union of $\frac{n}{k-1}$ copies of K_{k-1} .

Note that the bound in Theorem 2.1 is sharp only in the case (k-1)|n. Furthermore, for the general cases, say n = (k-1)t + r, this bound was refined by Faudree and Schelp [28], and independently by Kopylov [35] using a different method.

Theorem 2.2 (Faudree and Schelp [28], Kopylov [35]). Let n = (k-1)t + r with $t \ge 1$ and $0 \le r \le k-2$. Then

$$\operatorname{ex}(n, P_k) = t \binom{k-1}{2} + \binom{r}{2}.$$

Moreover, each n-vertex extremal graph for P_k is isomorphic to either

$$tK_{k-1} \cup K_r$$

or

$$(t-s-1)K_{k-1} \cup \left(K_{\frac{k-2}{2}} \otimes \overline{K}_{\frac{k}{2}+s(k-1)+r}\right)$$

for some $s \in [0, t-1]$ when k is even and $r \in \{\frac{k}{2}, \frac{k-2}{2}\}$.

In this paper, we shall use the case k = 4.

Corollary 2.3. Let G be a P_4 -free graph on n vertices. Then

$$e(G) \le n + \frac{1}{2}(j^2 - 3j).$$

where $j \in \{0, 1, 2\}$ and $j \equiv n \pmod{3}$. More precisely,

(a) if n = 3t, then $e(G) \leq 3t$, with equality if and only if $G = tC_3$;

(b) if n = 3t + 1, then $e(G) \leq 3t$, with equality if and only if

$$G \in \{tC_3 \cup K_1\} \cup \{(t-\ell-1)C_3 \cup K_{1,3\ell+3} : 0 \le \ell \le t-1\}$$

, (c) if n = 3t + 2, then $e(G) \le 3t + 1$, with equality if and only if

 $G \in \{tC_3 \cup K_2\} \cup \{(t-\ell-1)C_3 \cup K_{1,3\ell+4} : 0 \le \ell \le t-1\}.$

Recently, the stability result for (connected) P_k -free graphs was successively studied by Füredi, Kostochka, Luo and Verstraëte [24, 25], and Ma and Ning [38]. To prove Theorem 1.1, we need to apply the following variant due to Yuan [55]. In the original statement, it requires $t \ge 3$. We mention here that the result also holds for t = 2 by a similar argument.

Lemma 2.4 (Yuan [55]). Let c be a positive integer and n be sufficiently large. Given an n-vertex P_k -free graph G with $e(G) \ge ex(n, P_k) - c$. Let $\ell = \min\{n/(16c), n/(2k-2)\}$ and $t = \lfloor k/2 \rfloor \ge 2$. If k is odd, then G contains a copy of ℓK_{t-1} . If k is even, then G contains either a copy of ℓK_{k-1} or a copy of $K_{t-1,\ell}$.

2.2 A deep theorem of Simonovits

The decomposition family was firstly introduced by Simonovits [45], and it was used to study the Turán numbers of non-bipartite graphs. For a fixed graph H, the error term $o(n^2)$ in (2) can be refined by the decomposition family of H. Moreover, the decomposition family of a non-bipartite graph can help us to determine the finer structure of its extremal graphs. Now let us give the definition of the decomposition family of a graph.

Definition 2.5 (Simonovits). Given a graph L with $\chi(L) = p + 1 \ge 2$. Let $\mathcal{M}(L)$ be the family of minimal graphs M satisfying the following: there exists an integer $t \ge 1$ depending on L such that $L \subseteq (M \cup \overline{K}_t) \otimes T_{p-1}((p-1)t)$. We call $\mathcal{M}(L)$ the decomposition family of L.

In other words, a graph M belongs to $\mathcal{M}(L)$ if putting a copy of M (but not any of its proper subgraphs) into one vertex part of a large p-partite Turán graph $T_p(pt)$ will lead to a copy of L. For example, we have $\mathcal{M}(sK_{r+1}) = \{sK_2\}$ for each $s \geq 1$ and $r \geq 2$. Moreover, for the dodecahedron D_{20} and icosahedron I_{12} in Figure 1, it was proved in [43,44] that $\mathcal{M}(D_{20}) = \{6K_2\}$ and $\mathcal{M}(I_{12}) = \{P_6\}$, respectively. In addition, for the wheels (the pyramid) in Figure 2, it can be seen in [53] that $\mathcal{M}(W_{2k+1}) = \{K_{1,k}, C_{2k}\}$ and $\mathcal{M}(W_{2k}) = \{K_2\}$, since $\chi(W_{2k}) = 4$ and $\chi(W_{2k} - e) = 3$ for each edge e in the cycle.

The following lemma follows immediately from the definition.

Lemma 2.6. Given a graph L with $\chi(L) = p + 1 \ge 2$. If G is a graph obtained from the p-partite Turán graph $T_p(n)$ by embedding an extremal graph for $\mathcal{M}(L)$ into one part, then G is L-free. Consequently, we get $ex(n, L) \ge e(T_p(n)) + ex(\lceil n/p \rceil, \mathcal{M}(L))$.

In the proof of Theorem 1.1, we will use the following theorem, which attributes to a deep theorem of Simonovits [43, p. 371]. Roughly speaking, it states that if the decomposition family $\mathcal{M}(L)$ contains a linear forest (a graph in which each component is a path), then the extremal graphs for L have a very simple and symmetric structure.

Theorem 2.7 (Simonovits [43]). Let L be a given graph with $p = \chi(L) - 1$. If the decomposition family $\mathcal{M}(L)$ contains a linear forest, then there exist r = r(L) and $n_0 = n_0(r)$ such that if $n \ge n_0$, then each n-vertex extremal graph G for L belongs to the family of graphs satisfying the following. a at most n wanting of C the - amount C' entisfies (a) Afte . .

i) After deleting at most
$$r$$
 vertices of G , the remaining graph G' satisfies

$$G' = G^1 \otimes G^2 \otimes \cdots \otimes G^p,$$

where the graph G^i satisfies $||V(G^i)| - n/p| \leq r$ for each $1 \leq i \leq p$.

(b) For each $1 \le i \le p$, the graph G^i consists of vertex disjoint copies of non-isomorphic connected graphs $H_i^j s$ with $|V(H_i^j)| \leq r$ such that any two copies, say F_1 and F_2 , of $H_i^j s$ are symmetric subgraphs in G: there exists an isomorphism $\psi: F_1 \to F_2$ such that for every $x \in V(F_1)$ and $y \in G - F_1 - F_2$, xy is an edge if and only if $\psi(x)y$ is an edge.

Theorem 2.7 can benefit us to characterize the approximate structure of all extremal graphs for the prism C_{2k+1}^{\square} . We remark that this theorem was also used by Yuan [55] for treating the Turán problem of the power of a path. In particular, we refer the readers to [54] for graphs Lwhose decomposition family $\mathcal{M}(L)$ contains a matching.

3 Proof of Theorem 1.1

The following lemma is simple, but it plays an important role in our proof.

Lemma 3.1. The decomposition family $\mathcal{M}(C_{2k+1}^{\Box}) = \{P_4\}.$

Proof. The chromatic number $\chi(C_{2k+1}^{\square}) = 3$. For sufficiently large t, it is easy to see that embedding a P_4 into one partite of $K_{t,t}$ will lead to a copy of C_{2k+1}^{\square} , and embedding a proper subgraph of P_4 can not produce such a copy. Hence, $P_4 \in \mathcal{M}(C_{2k+1}^{\square})$. We claim that $\mathcal{M}(C_{2k+1}^{\square}) = \{P_4\}$. Suppose on the contrary that $\mathcal{M}(C_{2k+1}^{\square}) \neq \{P_4\}$. Let $H \in \mathcal{M}(C_{2k+1}^{\square})$ be a graph different from P_4 and Hdoes not contain a subgraph isomorphic to P_4 . Note that C_{2k+1}^{\square} is 3-regular. Then H is the disjoint union of some K_2 , $K_{1,2}$, $K_{1,3}$ and triangles. For sufficiently large t, let V_1 and V_2 be two partite sets of $K_{t,t}$. Embedding a copy of H into V_1 will lead to a C_{2k+1}^{\square} . We denote the new graph by $K_{t,t} + H$. Since V_2 is an independent set of $K_{t,t} + H$, we can see that H has no triangle components. Observe that there are exactly two copies of C_{2k+1} in C_{2k+1}^{\square} and V_2 is an independent set. Then H does not contain K_2 as a component. Since C_{2k+1} is an odd cycle, there must be a $K_{1,2}$ component in H, say uvw, such that uv is an edge of one C_{2k+1} and vw is an edge between two C_{2k+1} in C_{2k+1}^{\sqcup} . Let $x \notin \{u, w\}$ be a neighbor of v and $y, z \neq v$ be two neighbors of w. Then xy or xz is an edge of C_{2k+1}^{\square} . Observe that $\{x, y, z\} \subseteq V_2$ and V_2 is an independent set. Neither xy nor xz is an edge of $K_{t,t} + H$, a contradiction. Thus, such H does not exist and $\mathcal{M}(C_{2k+1}^{\square}) = \{P_4\}$.

Lemma 3.2. Let $G = G^1 \otimes G^2$ with $|V(G^i)| = n_i \ge 2k + 1$ for i = 1, 2.

- (a) Let G' be the graph obtained from G by adding a new vertex y with $|N_{G^i}(y)| \ge 3$ for i = 1, 2. If $G^1 \cup G^2$ has an edge incident with $N_G(y)$, then G' contains a copy of C_{2k+1}^{\square} .
- (b) If both G^1 and G^2 contain an edge, then G contains a copy of C_{2k+1}^{\square} .



Figure 4: The graph G'.

Proof. (a) Without loss of generality, we may assume that G^1 contains an edge e, which is incident with a vertex $x \in N_G(y)$; see Figure 4. Since $|N_{G^1}(y)| \ge 3$, let $v_1 \in V(G^1)$ be a neighbor of y such that v_1 is not an end vertex of e. Let $v_{2k} \in V(G^2)$ be a vertex adjacent to y. Then $yv_1v_2 \ldots v_{2k}$ and $xu_1u_2 \ldots u_{2k}$ are two copies of C_{2k+1} in G'; see the blue and red cycles in Figure 4, respectively. Observe that the above two cycles, together with the matching $\{xy, u_1v_1, u_2v_2, \ldots, u_{2k}v_{2k}\}$, form a copy of C_{2k+1}^{\square} . Thus, we get $C_{2k+1}^{\square} \subseteq G'$, as required.

(b) We may assume that $e_1 \in E(G^1)$ and $e_2 \in E(G^2)$. Similar with the case (a), one can find a red cycle C_{2k+1} containing the edge e_1 , and a blue cycle containing e_2 . Indeed, the odd prism C_{2k+1}^{\square} is contained in $K_{2k+1,2k+1} + \{e_1, e_2\}$. Combining with $(K_{2k+1,2k+1} + \{e_1, e_2\}) \subseteq G$, we then obtain a copy of C_{2k+1}^{\square} in G. The proof is completed. \square

The following well-known lemma proved by Erdős [14] and Simonovits [42] is a powerful tool in extremal graph problems; see, e.g., Corollary 4.3 in Chapter 6 of [3].

Lemma 3.3 (Erdős [14] and Simonovits [42]). Let H be a graph with $\chi(H) = p + 1 \ge 3$. If G is an extremal graph for H on n vertices, then $\delta(G) = (1 - 1/p)n + o(n)$.

Now, we are ready to give a proof of Theorem 1.1.

Proof of Theorem 1.1. Let G be a C_{2k+1}^{\Box} -free graph on n vertices with maximum number of edges. Our goal is to prove that G is obtained from a complete bipartite graph K_{n_a,n_b} (where $n_a + n_b = n$) by embedding an n_a -vertex extremal graph for P_4 into the part of size n_a . First of all, it follows from Lemmas 2.6 and 3.1 that

$$e(G) \ge \max\{n_a n_b + \exp(n_a, P_4) : n_a + n_b = n\}.$$
(3)

A simple calculation shows that

$$e(G) \ge e(T_2(n)) + \frac{n}{2} + O(1).$$

Now, we are going to apply Theorem 2.7. After deleting at most r vertices of G, the resulting graph G' can be written as the product $G' = G^1 \otimes G^2$ with $n/2 - r \leq |V(G^i)| \leq n/2 + r$ for each $i \in \{1, 2\}$. Moreover, each component of G^i is on at most r vertices. We denote by D the set of deleted vertices. Then $|D| \leq r$. By Lemma 3.2 (b), it is impossible that both G^1 and G^2 contain an edge. Then we may assume G^2 is an independent set of size n/2 + o(n). By Theorem 2.7, for each i = 1, 2, the isolated vertices of G^i are symmetric subgraphs of G. Thus, for any vertex $z \in D$, by the symmetry in Theorem 2.7, the isolated vertices of G^i either all are adjacent to z, or all are not adjacent to z.

We divide D into the following two sets: if $x \in D$ is not joint to vertices of G^2 , then let it belongs to D_2 , otherwise, let it belongs to D_1 . Let $A_i = V(G^i) \cup D_i$ and denote $|A_i| = n_i$ for i = 1, 2. Clearly, we have $V(G) = A_1 \cup A_2$. By the symmetry of Theorem 2.7, each vertex of A_1 is joint to all vertices of G^2 . By Lemma 3.3, we get $\delta(G) \ge n/2 + o(n)$. Note that G^2 is an empty graph and $|D_2| \le r$. Consequently, each vertex of G^2 is adjacent to n/2 + o(n) vertices of A_1 . Moreover, each vertex of D_2 is adjacent to n/2 - o(n) vertices of G^1 .

Claim 1. $G[A_i]$ does not contain a copy of P_4 for each $i \in \{1, 2\}$.

Proof of Claim 1. From the above discussion, we know that the edges between A_1 and $V(G^2)$ form a complete bipartite subgraph. By Definition 2.5 and Lemma 3.1, we have $G[A_1]$ is P_4 -free.

Assume on the contrary that $G[A_2]$ contains a copy of P_4 . Since $A_2 = V(G^2) \cup D_2$ and $E(G^2)$ is empty, the above copy of P_4 must be contained in $G[D_2]$. Choosing a set U consisting of 4kvertices of $A_2 \setminus V(P_4)$ arbitrarily, by Lemma 3.3, we know that these 4k vertices and the vertices of $V(P_4)$ have at least n/2 - o(n) common neighbors in G^1 . By Definition 2.5 and Lemma 3.1, there is a copy of C_{2k+1}^{\square} in the subgraph of G induced by the set $U \cup V(P_4)$ together with its common neighborhood, and so G contains a copy of C_{2k+1}^{\square} , a contradiction. The proof is completed. \square

Since $|D| \leq r$, by (3), we have $e(G[A_1]) \geq ex(|A_1|, P_4) - O(1)$. Thus, by Lemma 2.4, $G[A_1]$ contains a copy of either $K_{1,\ell}$ or ℓK_3 , where $\ell = \Theta(n)$. Next, we divide the proof into two cases. For convenience, we write $N(U) := \bigcap_{u \in U} N(u)$ for the common neighborhood of vertices of a set U and denote $N_A(U) := N(U) \cap A$.

Case 1. $G[A_1]$ contains a copy of ℓK_3 . We claim that A_2 is an independent set in G. Assume on the contrary that $G[A_2]$ contains an edge xy. Then xy is contained in $G[D_2]$ since $V(G^2)$ is an independent set and there are no edges between D_2 and $V(G^2)$. Let U_2 be a vertex set consisting of x, y and any 2k - 1 vertices of $A_2 \setminus \{x, y\}$. By Lemma 3.3, these 2k + 1 vertices of U_2 have at least $|A_1| - o(n)$ common neighbors in A_1 . In other words, there are o(n) vertices of A_1 which are outside of $N_{A_1}(U_2)$. Since $G[A_1]$ has ℓ disjoint copies of K_3 and $\ell = \Theta(n)$, we obtain that there exists a triangle located in $N_{A_1}(U_2)$. Consequently, we can choose a set U_1 consisting of 2k + 1vertices in $N_{A_1}(U_2)$ such that $G[U_1]$ contain a copy of K_3 . Note that $G[U_1]$ contains an edge and $G[U_2, U_1]$ forms a complete bipartite subgraph. By Lemma 3.2 (b), we obtain that G contains a copy of C_{2k+1}^{\Box} , a contradiction. Thus, there is no edge in $G[A_2]$. Denote $|A_1| = n_1$ and $|A_2| = n_2$. Therefore, it follows from Claim 1 that

$$e(G) = n_1 n_2 + e(A_1) \le \max \left\{ n_a n_b + \exp(n_a, P_4) : n_a + n_b = n \right\}.$$

Combining with (3), we obtain the desired bound on e(G). Moreover, the extremal graphs for the prism C_{2k+1}^{\square} are obtained from a complete bipartite graph K_{n_a,n_b} by embedding an n_a -vertex extremal graph for P_4 in the part of size n_a . Furthermore, the extremal graphs for P_4 are also characterized by applying Corollary 2.3.

Case 2. $G[A_1]$ contains a copy of $K_{1,\ell}$. Let u be the central vertex of $K_{1,\ell}$. Then u has at least $\ell = \Theta(n)$ neighbors in A_1 . Recall that each vertex of A_1 is joint to all vertices of G^2 . We denote $\ell_i = |N_{A_i}(u)|$ for each $i \in \{1, 2\}$. Then $\ell_i = \Theta(n)$.

Claim 2. There is no edge in $G[A_1 \setminus \{u\}]$ incident with a vertex of $N_{A_1}(u)$. Correspondingly, there is no edge in $G[A_2]$ incident with a vertex of $N_{A_2}(u)$.

Proof. If there is an edge in $G[A_1 \setminus \{u\}]$ incident with a vertex of $N_{A_1}(u)$, then $G[A_1]$ contains a copy of P_4 , which contradicts with Claim 1. Hence there is no edge in $G[A_1 \setminus \{u\}]$ incident with a vertex of $N_{A_1}(u)$. If there is an edge xy in $G[A_2]$ incident with a vertex of $N_{A_2}(u)$, then $x, y \in D_2$ since G^2 has no edge and there are no edges between D_2 and $V(G^2)$. Let E_2 be a vertex set consisting of x, y and any 2k vertices of $N_{G^2}(u)$. It follows that these 2k + 2 vertices of E_2 have $|A_1| - o(n)$ common neighbors in A_1 . Since u has $\ell_1 = \Theta(n)$ neighbors in A_1 , we know that

$$|N_{A_1}(E_2) \cap N_{A_1}(u)| \ge |N_{A_1}(E_2)| + |N_{A_1}(u)| - |A_1| \ge \ell_1 - o(n) \ge 2k + 1.$$

Thus, we can choose a set E_1 consisting of 2k + 1 vertices of $N_{A_1}(E_2) \cap N_{A_1}(u)$. Note that $|N_{E_1}(u)| = 2k + 1, |N_{E_2}(u)| \ge 2k + 1$ and $G[E_1, E_2]$ is a complete bipartite graph. By Lemma 3.2 (a), the sets E_1 , E_2 , and the vertex u, induce a graph containing a copy of C_{2k+1}^{\square} . This is a contradiction. Thus there is no edge in $G[A_2]$ incident with a vertex of $N_{A_2}(u)$.

From Claim 2, we obtain

$$e(A_1) = e(N_{A_1}(u) \cup \{u\}) + e(A_1 \setminus (N_{A_1}(u) \cup \{u\}))$$

$$\leq \ell_1 + \exp(n_1 - 1 - \ell_1, P_4) \leq \ell_1 + (n_1 - 1 - \ell_1) \leq \exp(n_1, P_4),$$
(4)

where the first inequality holds by Claim 1, and the others hold by Corollary 2.3. Applying Claim 2, we get

$$e(A_2) = e(A_2 \setminus N_{A_2}(u)) \le \exp(n_2 - \ell_2, P_4) \le n_2 - \ell_2.$$
(5)

Since u has $n_2 - \ell_2$ non-neighbors in A_2 , we bound the number of edges between A_1 and A_2 ,

$$e(A_1, A_2) \le n_1 n_2 - (n_2 - \ell_2). \tag{6}$$

Note that $e(G) = e(A_1, A_2) + e(A_1) + e(A_2)$. Combining (4), (5) and (6), we have

$$e(G) \le n_1 n_2 + \exp(n_1, P_4) \le \max\{n_a n_b + \exp(n_a, P_4) : n_a + n_b = n\}.$$
(7)

Hence, by (3), we have $e(G) = \max \{ \exp(n_a, P_4) + n_a n_b : n_a + n_b = n \}.$

Furthermore, using Corollary 2.3, we obtain

$$e(G) = \max_{n_a+n_b=n} \left\{ n_a(1+n_b) + \frac{1}{2}(j^2 - 3j) : j \in \{0, 1, 2\}, j \equiv n_a \pmod{3} \right\}.$$

Finally, we will characterize the extremal graphs. If the equality holds in (7), then from the above proof, we must have equalities in (4), (5) and (6). Denote $t_1 := n_1 - 1 - \ell_1$ and $t_2 := n_2 - \ell_2$. Then for i = 1, 2, we have

$$\operatorname{ex}(t_i, P_4) = t_i. \tag{8}$$

Moreover, attaining equality in (6) yields that each vertex of $A_1 \setminus \{u\}$ is adjacent to each vertex of A_2 . By Corollary 2.3, if $t_i > 0$, then $t_i \equiv 0 \pmod{3}$, and so $t_i \ge 3$. In addition, Corollary 2.3 also implies that an extremal graph for P_4 on at least 3 vertices contains a copy of P_3 . If both t_1 and t_2 are not zero, then by (8), both $G[A_1 \setminus \{u\}]$ and $G[A_2]$ contain a copy of P_2 . Hence by Lemma 3.2 (b), G contains a copy of C_{2k+1}^{\Box} , a contradiction. Thus, at most one of t_1 and t_2 is not zero. Firstly, if $t_1 \neq 0$ and $t_2 = 0$, then $A_2 = N_{A_2}(u)$ and A_2 is an independent set in G. Recall in Claim 1 that $G[A_1]$ is P_4 -free. Then G is obtained from K_{n_1,n_2} by embedding an extremal graph for P_4 into the

part A_1 . Secondly, if $t_1 = 0$ and $t_2 \neq 0$, then u is adjacent to all vertices of $A_1 \setminus \{u\}$. Moving the vertex u to the set A_2 , and we denote $B_1 := A_1 \setminus \{u\}$ and $B_2 := A_2 \cup \{u\}$. Recall in Claim 2 that there is no edge in $G[A_1 \setminus \{u\}]$ incident with a vertex of $N_{A_1}(u) = A_1 \setminus \{u\}$, so B_1 is an independent set. Moreover, $G[B_1, B_2]$ is a complete bipartite graph and $G[B_2]$ is P_4 -free, as desired. Finally, if $t_1 = t_2 = 0$, then $G[A_1, A_2]$ is a complete bipartite graph, $G[A_1]$ is a star K_{1,n_1-1} , and $G[A_2]$ is an empty graph, as required. To sum up, every extremal graph for C_{2k+1}^{\Box} is the join of an n_a -vertex extremal graph for P_4 and an n_b -vertex independent set.

4 Some Lemmas for the triangular prism

In this section, we shall consider the triangular prism. In the case k = 1 of Theorem 1.1, we will throw away the condition that requires the order n being sufficiently large. Using a quite different method, we will determine all n-vertex extremal graphs of the prism C_3^{\Box} for every integer $n \ge 6$.

4.1 A result for the power of a path

The *p*-th power of a path P_k on *k* vertices, denoted by P_k^p , is the graph obtained from P_k by joining each pair of vertices of P_k with distance at most *p*; see Figure 5.



Figure 5: The triangular prism C_3^{\Box} and the graph P_6^2 , respectively

In order to study $ex(n, C_3^{\Box})$, we need to use a result from Xiao, Katona, Xiao and Zamora [49] in which they determined the exact value of $ex(n, P_5^2)$ and $ex(n, P_6^2)$, respectively. For the power of a long path, see Yuan [55] for more results. In addition, the Turán number $ex(n, P_4^2)$ can be deduced from a general result of Dirac [11], which states that if $r \ge 3, n \ge r+1$ and G is an n-vertex graph with more than $e(T_{r-1}(n))$ edges, then G contains a copy of K_{r+1}^- , where K_{r+1}^- is the graph obtained from K_{r+1} by deleting any one edge. Setting r = 3 and observing that $P_4^2 = K_4^-$, we obtain $ex(n, P_4^2) = \lfloor n^2/4 \rfloor$ for each integer $n \ge 4$.

Before showing the proofs of Theorems 1.2 and 1.3, we need the following results.

Theorem 4.1 (Xiao et al. [49]). The maximum number of edges in an n-vertex P_6^2 -free graph $(n \neq 5)$ is

$$\operatorname{ex}(n, P_6^2) = \begin{cases} \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lfloor \frac{n-1}{2} \right\rfloor, & \text{if } n \equiv 1, 2, 3 \pmod{6}, \\ \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lceil \frac{n}{2} \right\rceil, & \text{otherwise.} \end{cases}$$

Theorem 4.2 (Xiao et al. [49]). Let $n \ge 6$. The extremal graphs for P_6^2 are the following ones. When $n \equiv 1 \pmod{6}$ then $F_n^{\lceil \frac{n}{2} \rceil, j}$ and $H_n^{\lfloor \frac{n}{2} \rfloor}$; when $n \equiv 2 \pmod{6}$ then $F_n^{\frac{n}{2}, j}$ and $F_n^{\frac{n}{2}+1, j}$; when $n \equiv 3 \pmod{6}$ then $F_n^{\lceil \frac{n}{2} \rceil, j}$ and $H_n^{\lceil \frac{n}{2} \rceil+1}$; when $n \equiv 0, 4, 5 \pmod{6}$ then $H_n^{\frac{n}{2}}$, $H_n^{\frac{n}{2}+1}$ and $H_n^{\lceil \frac{n}{2} \rceil}$, respectively, where j can have all the values satisfying the conditions $j \leq i$ and $3 \mid (i-j)$.

Comparing the above theorems with Theorems 1.2 and 1.3, it is surprising that the Turán numbers of graphs C_3^{\Box} and P_6^2 are completely the same, although their structures are different. In addition, it is noteworthy that the extremal graphs for C_3^{\Box} and P_6^2 are also the same except for three small graphs G_1, G_2 and G_3 , depicted in Figure 3.

In fact, it is a key relation between C_3^{\Box} and P_6^2 that $\mathcal{M}(C_3^{\Box}) = \mathcal{M}(P_6^2) = \{P_4\}$, where $\mathcal{M}(L)$ is the decomposition family of L stated in Definition 2.5. To a certain extent, the above phenomenon reveals that the decomposition family of a graph could determine the Turán number of a graph and also dominate the structures of the extremal graphs.

4.2 The C_3^{\Box} -free graphs containing P_6^2 as a subgraph

Before showing the proof of Theorems 1.2 and 1.3, we next present some useful lemmas for a special class of C_3^{\Box} -free graphs which contain the 2-power graph P_6^2 as a subgraph.

Lemma 4.3. Let G be a C_3^{\Box} -free graph on 6 vertices. If G contains a copy of P_6^2 , then $e(G) \leq 12$, with equality holding if and only if $G = G_1$.



Figure 6: Two drawings of the graph G_1

Proof. The proof is a standard argument on case-analysis. Since G has 6 vertices and P_6^2 is contained in G, we may assume that $V(G) = \{v_1, v_2, \ldots, v_6\}$, where v_1v_3, v_2v_4, v_3v_5 and v_4v_6 are edges of G. Clearly, we have $9 \le e(G) \le 15$. To prove $e(G) \le 12$, it is equivalent to show that there missed at least three edges in G whenever K_6 is C_3^{\Box} -free.

First of all, we claim that v_1v_6 can not be an edge of G. Otherwise, if $v_1v_6 \in E(G)$, then two triangles $v_1v_2v_3$ and $v_6v_4v_5$, together with three edges v_1v_6, v_2v_4, v_3v_5 form a copy of C_3^{\Box} , a contradiction. Thus, we get $2 \leq d_G(v_1) \leq 4$.

Case 1. $d_G(v_1) = 2$. More precisely, we have $v_1v_4, v_1v_5, v_1v_6 \notin E(G)$, and G misses at least three edges of K_6 . Clearly, we have $e(G) \leq 12$. If e(G) = 12, then v_2v_5, v_2v_6 and v_3v_6 are edges of G. In this case, we get $G = G_1$, as required.

Case 2. $d_G(v_1) = 3$. In other words, we have either $v_1v_4 \in E(G)$ or $v_1v_5 \in E(G)$. In the former case, if $v_1v_4 \in E(G)$, then $v_2v_6 \notin E(G)$. Otherwise, if $v_2v_6 \in E(G)$, then two triangles $v_1v_2v_3$ and $v_4v_6v_5$, together with edges v_1v_4, v_2v_6, v_3v_5 form a copy of C_3^{\Box} , a contradiction. Hence, we conclude that v_1v_6, v_1v_5 and v_2v_6 are not edges of G, which leads to $e(G) \leq 12$. If e(G) = 12, then v_1v_4, v_2v_5, v_3v_6 are edges of G, which together with two triangles $v_1v_2v_3$ and $v_4v_5v_6$ form a copy of C_3^{\Box} , a contradiction. Hence $e(G) \leq 11$.

In the later case, that is, $v_1v_5 \in E(G)$ and $v_1v_4, v_1v_6 \notin E(G)$, we can get $v_2v_6 \notin E(G)$ and $v_3v_6 \notin E(G)$. Otherwise, if $v_2v_6 \in E(G)$, then triangles $v_1v_2v_3$ and $v_5v_6v_4$, with edges v_1v_5, v_2v_6, v_3v_4 form a copy of C_3^{\Box} , a contradiction. If $v_3v_6 \in E(G)$, then triangles $v_1v_2v_3$ and $v_5v_4v_6$, with edges v_1v_5, v_2v_4, v_3v_6 form a copy of C_3^{\Box} , a contradiction. Therefore, we have $e(G) \leq 11$. **Case 3.** $d_G(v_1) = 4$. In this case, we have $v_1v_4, v_1v_5 \in E(G)$. Similar with the later case of Case 2, we get $v_2v_6 \notin E(G)$ and $v_3v_6 \notin E(G)$ as well. Recall that $v_1v_6 \notin E(G)$. Consequently, we obtain $e(G) \leq 12$. If e(G) = 12, then $v_2v_5 \in E(G)$. By reversing the order of $v_1v_2 \cdots v_6$, it is not hard to see that G is isomorphic to G_1 , as required.

In the following, we always assume that

$$P_6^2 \subseteq G. \tag{9}$$

Assume that the subgraph P_6^2 is obtained from $P_6 = v_1 v_2 \cdots v_6$ by joining each pair of vertices of $V(P_6)$ with distance at most 2. It can be checked that each vertex $v \in V(G) \setminus V(P_6^2)$ can be adjacent to at most 4 vertices of the copy of P_6^2 . Otherwise, if there is a vertex with 5 neighbors in P_6^2 , then we can find a copy of C_3^{\Box} in G, a contradiction.

The following claim holds immediately under the condition (9).

Claim 3. If $v \in V(G) \setminus V(P_6^2)$ is adjacent to exactly 4 vertices of P_6^2 , then there are exactly 4 adjacency relations (up to graph isomorphism) between v and P_6^2 ; see Figure 7.



Figure 7: Four types for the vertex v satisfying $|N(v) \cap V(P_6^2)| = 4$

Furthermore, the following lemma can be verified case by case.

Lemma 4.4. There are at most 2 vertices of $V(G) \setminus V(P_6^2)$ adjacent to exactly 4 vertices of $V(P_6^2)$. Moreover, if $x, y \in V(G) \setminus V(P_6^2)$ are such two vertices, then there are only 2 adjacency relations between $\{x, y\}$ and P_6^2 , and moreover $xy \notin E(G)$; see Figure 8.



Figure 8: The 8-vertex graphs H_1 and H_2 , respectively.

Proof. First of all, we show that if x, y are two vertices in the same type, then G contains a copy of C_3^{\Box} , except that they are of Type 1, drawing as H_2 in Figure 8. Indeed, if x, y are of Type 1, then either $N_{P_6^2}(x) = N_{P_6^2}(y) = \{v_1, v_2, v_3, v_4\}$, or $N_{P_6^2}(x) = \{v_1, v_2, v_3, v_4\}$ and $N_{P_6^2}(y) = \{v_6, v_5, v_4, v_3\}$. For the former case, the triangles v_1v_3x and v_2v_4y , together with three edges v_1y, v_3v_4, xv_2 , form a copy of C_3^{\Box} , a contradiction; For the latter case, it is possible, and stated as H_2 in Figure 8. If x, y are of Type 2, then the symmetry gives $N_{P_6^2}(x) = N_{P_6^2}(y) = \{v_2, v_3, v_4, v_5\}$. Then the triangles

 v_2v_3x and v_4v_5y , with the edges v_2y, v_3v_5, xv_4 form a copy of C_3^{\Box} . If x, y are of Type 3, then $N_{P_6^2}(x) = N_{P_6^2}(y) = \{v_1, v_3, v_4, v_6\}$. Thus, the triangles v_1v_3x and v_4v_6y form a copy of C_3^{\Box} since v_1y, v_3v_4, xv_6 are edges of G. If x, y are of Type 4, then either $N_{P_6^2}(x) = N_{P_6^2}(y) = \{v_1, v_3, v_5, v_6\}$, or $N_{P_6^2}(x) = \{v_1, v_3, v_5, v_6\}$ and $N_{P_6^2}(y) = \{v_6, v_4, v_2, v_1\}$. For the former case, we can see that triangles v_1v_3x and v_5v_6y form a copy of C_3^{\Box} since v_1y, xv_6, v_3v_5 are edges in G. For the latter case, it can be seen that G contains a copy of C_3^{\Box} as witnessed by triangles v_1v_3x and v_6v_4y with three edges v_1y, xv_6 and v_3v_4 .

Secondly, one consider the case that x, y are vertices with different types. More precisely, if x is of Type 1 and y is of Type 2, then G has a copy of C_3^{\Box} consisting of the triangles xv_1v_2, yv_3v_4 and the edges v_1v_3, v_2y, xv_4 . If x is of Type 1 and y is of Type 3, then the triangles xv_1v_2, yv_3v_4 and the edges v_1y, v_2v_3, xv_4 form a copy of C_3^{\Box} in G. If x is of Type 1 and y is of Type 4, then by symmetry, we get either $N_{P_6^2}(x) = \{v_1, v_2, v_3, v_4\}$ and $N_{P_6^2}(y) = \{v_1, v_3, v_5, v_6\}$, or $N_{P_6^2}(x) = \{v_6, v_5, v_4, v_3\}$ and $N_{P_6^2}(y) = \{v_1, v_3, v_5, v_6\}$. To avoid a copy of C_3^{\Box} in G, the former case is possible, which is stated as H_1 in Figure 8. For the latter case, one can find a copy of C_3^{\Box} on the triangles xv_4v_6, yv_3v_5 with the edges xv_3, v_4v_5, v_6y . If x is of Type 2 and y is of Type 3, then G contains a copy of C_3^{\Box} , which appears on the triangles xv_2v_4, yv_1v_3 and the edges xv_3, v_2v_1, v_4y . If x is of Type 2 and y is of Type 4, then G has a copy of C_3^{\Box} by combining the triangles xv_3v_4, yv_5v_6 with the edges v_3y, v_4v_6, xv_5 . If x is of Type 3 and y is of Type 4, then one can find a copy of C_3^{\Box} on the triangles xv_1v_3, yv_5v_6 with the edges xv_6, v_1y, v_3v_5 .

Moreover, it can be checked further that $V(G) \setminus V(P_6^2)$ does not contain three vertices which are adjacent to 4 vertices of P_6^2 . Otherwise, if such three vertices exist, i.e., we add one more vertex with degree 4 on P_6^2 to the graph H_1 or H_2 , then one can check that there is a copy of C_3^{\Box} in G. Finally, if xy is an edge of G, then either $H_1 + xy$ or $H_2 + xy$ can have a copy of C_3^{\Box} . Indeed, in both cases, the copy will appear on the triangles v_2v_4x and v_3v_5y , with the edges v_2v_3, v_4v_5, xy . Thus, we get $xy \notin E(G)$.

Lemma 4.5. If G contains H_1 as a subgraph, then $e(G[V(P_6^2)]) \leq 10$, with equality if and only if v_3v_6 is an edge in G.

Proof. Observe that $e(P_6^2) = 9$. In other words, we need to show that one can join at most one non-edge of P_6^2 to avoid the copy of C_3^{\Box} in G. Note that

$$E(K_6) \setminus E(P_6^2) = \{v_1v_4, v_1v_5, v_1v_6, v_2v_5, v_2v_6, v_3v_6\}.$$
(10)

If $v_1v_4 \in E(G[V(P_6^2)])$, then G contains two triangles v_1v_4x and v_3v_5y , which together with the matching $\{v_1y, v_4v_5, xv_3\}$, form a copy of C_3^{\Box} , a contradiction. If $v_1v_5 \in E(G[V(P_6^2)])$, then the triangles v_1v_2x and $v_3v_4v_5$, with the matching $\{v_1v_5, v_2v_3, xv_4\}$, form a copy of C_3^{\Box} , a contradiction. If $v_1v_6 \in E(G[V(P_6^2)])$, then there are two triangles v_1v_3x and $v_4v_5v_6$, combining with the matching $\{v_1v_6, v_3v_5, xv_4\}$, we find a copy of C_3^{\Box} , a contradiction. In fact, whenever $v_1v_6 \in E(G)$, one can find a copy of C_3^{\Box} without the use of the vertex x, which was showed in the second paragraph of the proof of Lemma 4.3. If $v_2v_5 \in E(G[V(P_6^2)])$, then triangles v_1v_2x and $v_3v_4v_5$, with the matching $\{v_1v_3, v_2v_5, xv_4\}$, form a copy of C_3^{\Box} , a contradiction. If $v_2v_6 \in E(G[V(P_6^2)])$, then G contains a copy of C_3^{\Box} , which consists of two triangles v_2v_3x and $v_4v_5v_6$, with the matching $\{v_2v_6, v_3v_5, xv_4\}$. Thus, the only possible case is $v_3v_6 \in E(G[V(P_6^2)])$, and then it follows that $e(G[V(P_6^2)]) \leq 10$. \Box

Similarly, we can prove the following lemma.

Lemma 4.6. If G contains H_2 as a subgraph, then $e(G[V(P_6^2)]) \leq 11$, and equality holds if and only if both v_1v_4 and v_3v_6 are edges in G.

Proof. We need to check each non-edge of $E(G[V(P_6^2)])$ stated in (10). If $v_1v_5 \in E(G[V(P_6^2)])$, then the triangles v_1v_2x and $v_3v_4v_5$, with the matching $\{v_1v_5, v_2v_4, xv_3\}$, form a copy of C_3^{\Box} , a contradiction. If $v_1v_6 \in E(G[V(P_6^2)])$, then the same argument in the proof of Lemma 4.5 can yield a copy of C_3^{\Box} , a contradiction. In addition, combining the triangles v_1v_3x, v_4v_6y with the matching $\{v_1v_6, v_3y, xv_4\}$, we find another copy of C_3^{\Box} . If $v_2v_5 \in E(G[V(P_6^2)])$, then the proof of Lemma 4.5 yields a copy of C_3^{\Box} , a contradiction. Additionally, there is another copy of C_3^{\Box} on two triangles v_2v_3x and v_4v_5y , joined by the matching $\{v_2v_5, v_3y, xv_4\}$. Finally, if $v_2v_6 \in$ $E(G[V(P_6^2)])$, then there is a copy of C_3^{\Box} in G, which consists of triangles v_2v_3x, v_4v_6y and the matching $\{v_2v_6, v_3y, xv_4\}$. Therefore, it is possible that $v_1v_4, v_3v_6 \in E(G[V(P_6^2)])$, and only in such a case, we have $e(G[V(P_6^2)]) = 11$.

5 Proofs of Theorems 1.2 and 1.3

In this section, we shall prove Theorems 1.2 and 1.3 as follows.

Proof of Theorem 1.2. For notational convenience, we denote

$$f(n) := \begin{cases} \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lfloor \frac{n-1}{2} \right\rfloor, & \text{if } n \equiv 1, 2, 3 \pmod{6}, \\ \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lceil \frac{n}{2} \right\rceil, & \text{otherwise.} \end{cases}$$

First of all, when $n \equiv 0, 4, 5 \pmod{6}$, we have $\exp(n, C_3^{\Box}) \ge \lfloor \frac{n^2}{4} \rfloor + \lceil \frac{n}{2} \rceil$, which follows from the constructions of $G_1, H_n^{\frac{n}{2}}, H_n^{\frac{n}{2}+1}$ and $H_n^{\lceil \frac{n}{2} \rceil}$, respectively. In addition, when $n \equiv 1, 2, 3 \pmod{6}$, the lower bound $\exp(n, C_3^{\Box}) \ge \lfloor \frac{n^2}{4} \rfloor + \lfloor \frac{n-1}{2} \rfloor$ follows from the definitions of G_2, G_3 and $F_n^{\lceil \frac{n}{2} \rceil, j}$. Thus, it remains to show the following inequality

$$\operatorname{ex}(n, C_3^{\square}) \le f(n). \tag{11}$$

Our proof of (11) is by induction on n. Let G be an n-vertex C_3^{\square} -free graph. We need to show $e(G) \leq f(n)$. Since our inductive step will go from n - 6 to n, we have to prove the base case in each residue class mod 6.

§ Base Case §

When $1 \le n \le 4$, it is clear that K_n is the C_3^{\square} -free graph with the maximum number of edges, and one can verify that $e(K_n) = f(n)$. Moreover, when n = 6, by Theorem 4.1, we have $e(G) \le 12$ whenever G is P_6^2 -free; by Lemma 4.3, we get $e(G) \le 12$ whenever G has a copy of P_6^2 . Then we always have $e(G) \le 12 = f(6)$, as needed. We remark here that when n = 5, we have

$$ex(5, C_3^{\perp}) = e(K_5) = 10 > 9 = f(5).$$
(12)

This is a counterexample. Thus we require $n \neq 5$.

Next, we are going to prove the statement for n = 11. First of all, if $P_6^2 \not\subseteq G$, then by Theorem 4.1, the results hold immediately. In what follows, we assume that $P_6^2 \subseteq G$. Under this constraint, our aim is to show that

$$e(G) < f(11) = 36. \tag{13}$$

Firstly, if $V(G) \setminus V(P_6^2)$ forms a copy of K_5 in G, then each $v_i \in V(P_6^2)$ is adjacent to at most 2 vertices of K_5 , otherwise, there is a copy of C_3^{\Box} in G. By Lemma 4.3, we get $e(G[V(P_6^2)]) \leq 12$. Therefore, we have $e(G) \leq 12 + 12 + 10 = 34 < 36 = f(11)$.

Secondly, if $V(G) \setminus V(P_6^2)$ forms a copy of K_5^- , then we assume that $V(K_5^-) = \{u_1, u_2, u_3, u_4, u_5\}$ and $u_4u_5 \notin E(K_5^-)$. The following two facts can be checked. The first fact asserts that for each $i \in \{1, \ldots, 6\}$, the vertex v_i is adjacent to at most 3 vertices of K_5^- , with equality holding if and only if $N_{K_5^-}(v_i) = \{u_1, u_2, u_3\}$, otherwise, it leads to $C_3^{\Box} \subseteq G$. The second fact states that there are at most 7 edges between $\{v_1, v_2, v_3\}$ and $V(K_5^-)$, otherwise, one can get $C_3^{\Box} \subseteq G$. The similar result holds between $\{v_4, v_5, v_6\}$ and K_5^- . Therefore, the number of edges between $V(P_6^2)$ and $V(K_5^-)$ is at most 14. Recall in Lemma 4.3 that $e(G[V(P_6^2)]) \leq 12$. Therefore, we get $e(G) \leq 12 + 14 + 9 = 35 < f(11)$.

Finally, we will consider the remaining case

$$e(V(G) \setminus V(P_6^2)) \le 8.$$

By Lemma 4.4, we proceed in the following three possible cases.

Case 1. There are exactly 2 vertices of $V(G) \setminus V(P_6^2)$, say x and y, each of them is adjacent to 4 vertices of $V(P_6^2)$. By Lemma 4.4, we obtain that either $H_1 \subseteq G$ or $H_2 \subseteq G$.

Subcase 1.1. If $H_1 \subseteq G$, then by Lemma 4.5, we know that $e(G[V(P_6^2)]) \leq 10$. Consequently, we have $e(G) \leq 10 + (4 + 4 + 3 + 3 + 3) + 8 = 35 < f(11)$, as desired.

Subcase 1.2. If $H_2 \subseteq G$, then by Lemma 4.6, we have $e(G[V(P_6^2)]) \leq 11$, equality holds if and only if both v_1v_4 and v_3v_6 are edges of G. If there is a vertex of $V(G) \setminus (V(P_6^2) \cup \{x, y\})$ adjacent to at most 2 vertices of $V(P_6^2)$, then $e(G) \leq 11 + (4 + 4 + 2 + 3 + 3) + 8 = 35 < f(11)$, as desired. If every vertex of $V(G) \setminus (V(P_6^2) \cup \{x, y\})$ is adjacent to 3 vertices of $V(P_6^2)$, then it can be seen further that $e(G[V(P_6^2)]) \leq 10$. Otherwise, if $e(G[V(P_6^2)]) = 11$, then Lemma 4.6 implies $v_1v_4, v_3v_6 \in E(G)$. Any vertex of $V(G) \setminus (V(P_6^2) \cup \{x, y\})$ with degree 3 on $V(P_6^2)$ can lead to a copy of C_3^{\Box} , a contradiction. Therefore, we have $e(G) \leq 10 + (4 + 4 + 3 \times 3) + 8 = 35 < f(11)$.

Case 2. There is exactly one vertex $v \in V(G) \setminus V(P_6^2)$, which is adjacent to 4 vertices of $V(P_6^2)$. By Claim 3, v is one of the four types in Figure 7. In either case, by Lemma 4.3, we can get $G[V(P_6^2)] \neq G_1$ and then $e(G[V(P_6^2)]) \leq 11$. Otherwise, if $G[V(P_6^2)] = G_1$, then v_2v_5, v_2v_6 and v_3v_6 are edges of G. Observe that the vertex set $\{v_2, v_3, \ldots, v_6\}$ forms a copy of the complete graph K_5 in G. Since v has 4 neighbors on P_6^2 , we know that there are at least 3 neighbors of v which are contained in the set $\{v_2, v_3, \ldots, v_6\}$. A trivial fact states that if v has at least 3 neighbors on K_5 , then $\{v\} \cup V(K_5)$ contains a copy of C_3^{\Box} . Thus G has a copy of C_3^{\Box} . Therefore, we have $e(G) \leq 11 + (4 + 3 \times 4) + 8 = 35 < f(11)$, as needed.

Case 3. All vertices of $V(G) \setminus V(P_6^2)$ are adjacent to at most 3 vertices of $V(P_6^2)$. By Lemma 4.3 again, it follows that $e(G) \le 12 + 15 + 8 = 35 < f(11)$. In conclusion, we complete the proof of (13) in the base case n = 11.

§ Inductive Step §

In what follows, we will show the inductive step. Suppose that (11) holds for all integers $\ell \leq n-1$ and $\ell \neq 5$. Now we consider the case $n \geq 7$. Assume that G is a C_3^{\square} -free graph on n vertices with maximum number of edges. We need to show that $e(G) \leq f(n)$. Our inductive step will start from n-6 to n. Upon on computations, we can obtain that

$$f(n) = f(n-6) + 3n - 6.$$
(14)

First of all, if $P_6^2 \not\subseteq G$, then by Theorem 4.1, we get $e(G) \leq f(n)$, as desired. In what follows, we assume that $P_6^2 \subseteq G$, and then we show a **stronger** result

$$e(G) \le f(n) - 1,\tag{15}$$

except for n = 7, 8. By Lemma 4.4, there are at most 2 vertices of $V(G) \setminus V(P_6^2)$ adjacent to exactly 4 vertices of $V(P_6^2)$. In the sequel, we shall present the proof in three cases.

Case A. If there are exactly 2 vertices of $V(G) \setminus V(P_6^2)$, say x and y, adjacent to 4 vertices of P_6^2 , then by Lemma 4.4, we get either $H_1 \subseteq G$ or $H_2 \subseteq G$.

Subcase A.1. Assume that $H_1 \subseteq G$. Then Lemma 4.5 gives $e(G[V(P_6^2)]) \leq 10$.

If $e(G[V(P_6^2)]) \leq 9$, then $e(G) \leq 9 + 8 + 3(n-8) + ex(n-6, C_3^{\Box}) \leq 3n-7 + f(n-6) < f(n)$, as needed. In what follows, we assume that $e(G[V(P_6^2)]) = 10$. Then Lemma 4.5 implies that $G[V(P_6^2)]$ consists of P_6^2 by adding the edge v_3v_6 .

If there exists a vertex of $V(G) \setminus (V(P_6^2) \cup \{x, y\})$ adjacent to at most 2 vertices of P_6^2 , then $e(G) \leq 10 + 8 + 2 + 3(n-9) + ex(n-6, C_3^{\Box}) \leq 3n-7 + f(n-6) < f(n)$, where the last inequality holds by (14).

If all vertices of $V(G) \setminus (V(P_6^2) \cup \{x, y\})$ are adjacent to 3 vertices of P_6^2 , then it is easy to check that they have the same neighbors on P_6^2 ; see H_3 in Figure 9. Next, we show that $V(G) \setminus V(P_6^2)$ is an independent set in G. Indeed, by Lemma 4.4, we know that $xy \notin E(G)$. In Figure 9, if xz is an edge of G, then combining with triangles $xv_3z, v_4v_5v_6$ with edges v_3v_6, xv_4, zv_5 , we see that Ghas a copy of C_3^{\Box} , a contradiction. If yz is an edge of G, then the triangles $v_3yz, v_4v_5v_6$ and edges v_3v_4, yv_5, zv_6 form a copy of C_3^{\Box} , a contradiction. If zw is an edge of G, then the copy of C_3^{\Box} will appear on the triangles $v_3zw, v_4v_5v_6$ and edges v_3v_4, zv_5, wv_6 , a contradiction. To sum up, we get that $G = H_3$; see Figure 9. Consequently, we have e(G) = 10 + 8 + 3(n - 8) < f(n), as desired.



Figure 9: The graphs H_3 and G_3 , respectively.

Subcase A.2. Assume that $H_2 \subseteq G$. By Lemma 4.6, we have $e(G[V(P_6^2)]) \leq 11$.

If $e(G[V(P_6^2)]) \le 9$, then $e(G) \le 9 + 8 + 3(n-8) + ex(n-6, C_3^{\Box}) \le 3n - 7 + f(n-6) < f(n)$, where the last inequality follows from (14).

If $10 \le e(G[V(P_6^2)]) \le 11$, then by the proof of Lemma 4.6, we know that $v_1v_4 \in E(G)$ or $v_3v_6 \in E(G)$. We need to prove the following claim.

Claim 4. If H_2 is a subgraph of G and $z \in V(G) \setminus (V(P_6^2) \cup \{x, y\})$ is a vertex which is adjacent to three vertices of P_6^2 , then $zx \notin E(G)$ and $zy \notin E(G)$.

Proof of Claim. The proof is straightforward.

If $V(G) \setminus (V(P_6^2) \cup \{x, y\})$ has a vertex, say z, which is adjacent to at most 2 vertices of P_6^2 , then $e(G) \leq 11+8+2+3(n-9)+ex(n-6, C_3^{\Box}) \leq 3n-6+f(n-6) = f(n)$. Next, we show further that e(G) < f(n). Otherwise, if the equality e(G) = f(n) holds, then the above inequalities must be equalities. More precisely, we get $e(G[V(P_6^2)]) = 11$, each vertex outside of $V(P_6^2) \cup \{x, y, z\}$ has exactly 3 neighbors on P_6^2 , and the induced subgraph $G[V(G) \setminus V(P_6^2)]$ is an (n-6)-vertex extremal graph for C_3^{\Box} . By Claim 4, we see that each vertex outside of $V(P_6^2) \cup \{x, y, z\}$ can not be adjacent to any vertex of $\{x, y\}$. Consequently, in the subgraph $G[V(G) \setminus V(P_6^2)]$, the vertex x(and y) has at most one neighbor, and the only possible neighbor is z. However, by induction, we can see that the extremal graphs of order at least 3 for C_3^{\square} do not contain any vertex with degree at most one. This is a contradiction. In other words, the subgraph $G[V(G) \setminus V(P_6^2)]$ can not be an extremal graph for C_3^{\square} . Then we get e(G) < f(n), as desired in (15).

If all vertices of $V(G) \setminus (V(P_6^2) \cup \{x, y\})$ are adjacent to 3 vertices of P_6^2 , then by Claim 4, there is no edge between $\{x, y\}$ and $V(G) \setminus (V(P_6^2) \cup \{x, y\})$. In other words, x, y are two isolated vertices in the induced subgraph $G[V(G) \setminus V(P_6^2)]$. Therefore, $e(G) \leq 11 + 8 + 3(n-8) + ex(n-8, C_3^{\Box}) \leq 3n - 5 + f(n-8) < f(n)$, where the second inequality follows by induction whenever $n \neq 13$, and the last inequality holds for $n \neq 8$ by a direct computation.

In particular, for n = 8, we get $V(G) = V(P_6^2) \cup \{x, y\}$ and

$$e(G) \le 11 + 8 = 19 = f(8), \tag{16}$$

where the equality holds if and only if G is obtained from H_2 by adding two edges v_1v_4 and v_3v_6 ; see G_3 in Figure 9. In the equality case, we get $G = G_3$ and $e(G_3) = 19 = f(8)$. In addition, for n = 13, recall in (12) that $ex(5, C_3^{\Box}) = e(K_5) = 10$, and then $e(G) \le 11 + 8 + 3 \times 5 + ex(5, C_3^{\Box}) = 44 < 48 = f(13)$, as needed.

Case B. If there is exactly one vertex $v \in V(G) \setminus V(P_6^2)$ adjacent to 4 vertices of P_6^2 , then by Claim 3, we know that v is one of the four types in Figure 7.

Firstly, we claim that $e(G[V(P_6^2)]) \leq 11$. Otherwise, if $e(G[V(P_6^2)]) = 12$, then Lemma 4.3 implies that v_2v_5, v_2v_6 and v_3v_6 are edges of G. Note that the vertex set $\{v_2, v_3, \ldots, v_6\}$ forms a copy of K_5 in G. Since v has 4 neighbors on P_6^2 , it follows that v has at least 3 neighbors on this copy of K_5 , which leads to a copy of C_3^{\Box} in G, a contradiction.

Subcase B.1. If v is of Type 2, Type 3 or Type 4 in Figure 7, then we can show further that $e(G[V(P_6^2)]) \leq 10$. More precisely, if v is of Type 2, then adding any non-edge of P_6^2 to G can make a copy of C_3^{\Box} , and so $e(G[V(P_6^2)]) = 9$; if v is of Type 3, then v_2v_5 is the only possible edge of G, and hence $e(G[V(P_6^2)]) \leq 10$; if v is of Type 4, then v_3v_6 is the only possible edge, and thus $e(G[V(P_6^2)]) \leq 10$. Therefore, we obtain $e(G) \leq 10 + 4 + 3(n-7) + ex(n-6, C_3^{\Box}) \leq 3n-7 + f(n-6) < f(n)$.

Subcase B.2. Assume that v is of Type 1 in Figure 7. In this type, we shall show that

$$v_1v_5, v_1v_6, v_2v_5, v_2v_6 \notin E(G).$$
 (17)

Indeed, if $v_1v_5 \in E(G)$, then G has a copy of C_3^{\Box} , which appears on the triangles $vv_1v_2, v_3v_4v_5$ with the edges v_1v_5, v_2v_3, vv_4 . If $v_1v_6 \in E(G)$, then the triangles $vv_1v_3, v_4v_5v_6$ and the edges v_1v_6, v_3v_5, vv_4 form a copy of C_3^{\Box} in G. If $v_2v_5 \in E(G)$, then G contains a copy of C_3^{\Box} consisting of the triangles $vv_1v_2, v_3v_4v_5$ and the edges v_1v_3, v_2v_5, vv_4 . If $v_2v_6 \in E(G)$, then $C_3^{\Box} \subseteq G$ by combining the triangles $vv_2v_3, v_4v_5v_6$ with the edges v_2v_6, v_3v_5, vv_4 . This completes the proof of (17). Therefore, to avoid a copy of C_3^{\Box} in G, the possible edges on $V(P_6^2)$ are v_1v_4 and v_3v_6 ; see Figure 10.



Figure 10: v is of Type 1 and $e(G[V(P_6^2)]) = 11$

First of all, if $e(G[V(P_6^2)]) \leq 10$, then $e(G) \leq 10 + 4 + 3(n-7) + ex(n-6, C_3^{\Box}) \leq 3n-7 + f(n-6) < f(n)$, as required. Next, we assume that $e(G[V(P_6^2)]) = 11$. By (17), we know that $v_1v_4, v_3v_6 \in E(G)$; see Figure 10.

If there exist a vertex of $V(G) \setminus (V(P_6^2) \cup \{v\})$ adjacent to at most 2 vertices of P_6^2 , then there are at most 4 + 2 + 3(n - 8) edges between $V(P_6^2)$ and $V(G) \setminus V(P_6^2)$. Thus, we get $e(G) \leq 11 + 4 + 2 + 3(n - 8) + ex(n - 6, C_3^{\Box}) \leq 3n - 7 + f(n - 6) < f(n)$.

If all vertices of $V(G) \setminus (V(P_6^2) \cup \{v\})$ are adjacent to 3 vertices of P_6^2 , then there are exactly 4 + 3(n-7) edges between $V(P_6^2)$ and $V(G) \setminus V(P_6^2)$. Since each vertex $x \in V(G) \setminus (V(P_6^2) \cup \{v\})$ has 3 neighbors on P_6^3 , we can check that $vx \notin E(G)$. Otherwise, one can find a copy of C_3^{\Box} in G. Therefore, v is an isolated vertex in the induced subgraph $G[V(G) \setminus V(P_6^2)]$. Then $e(G) \leq 11 + 4 + 3(n-7) + ex(n-7, C_3^{\Box}) \leq 3n-6 + f(n-7) < f(n)$, where the second inequality follows by induction whenever $n \neq 12$, and the last inequality holds for $n \neq 7$ by calculations.

In particular, for n = 7, we have $V(G) = V(P_6^2) \cup \{v\}$ and

$$e(G) \le 11 + 4 = 15 = f(7), \tag{18}$$

where equality holds if and only if G is a graph of Type 1 by adding edges v_1v_4 and v_3v_6 ; see Figure 10. In the equality case, we see that $G = G_2$ and $e(G_2) = 15 = f(7)$. Moreover, for n = 12, recall in (12) that $ex(5, C_3^{\Box}) = e(K_5) = 10$, then we have $e(G) \leq 11 + 4 + 3 \times 5 + ex(5, C_3^{\Box}) = 40 < 42 = f(12)$, as desired in (15).

Case C. Assume that $V(G) \setminus V(P_6^2)$ has no vertex which has 4 neighbors on P_6^2 . By Lemma 4.3, we have $e(G[V(P_6^2)]) \leq 12$. We show the proof in two subcases.

Subcase C.1. There is a vertex of $V(G) \setminus V(P_6^2)$ adjacent to at most 2 vertices of P_6^2 . Then the number of edges between $V(P_6^2)$ and $V(G) \setminus V(P_6^2)$ is at most 2 + 3(n-7) since the other vertices of $V(G) \setminus V(P_6^2)$ adjacent to at most 3 vertices of P_6^2 . Thus, we get $e(G) \leq 12 + 2 + 3(n-7) + ex(n-6, C_3^{\Box}) \leq 3n-7 + f(n-6) < f(n)$.

Subcase C.2. All vertices of $V(G) \setminus V(P_6^2)$ are adjacent to exactly 3 vertices of P_6^2 .

If $e(G[V(P_6^2)]) \le 11$, then $e(G) \le 11 + 3(n-6) + ex(n-6, C_3^{\Box}) \le 3n - 7 + f(n-6) < f(n)$, where the second inequality holds by induction.

If $e(G[V(P_6^2)]) = 12$, then by Lemma 4.3, we know that $G[V(P_6^2)] = G_1$, and so v_2v_5, v_2v_6 and v_3v_6 are edges of G. Note that $G[\{v_2, v_3, \ldots, v_6\}]$ forms a copy of K_5 ; see Figure 11. We next show that for every $y \in V(G) \setminus V(P_6^2)$,

$$N_{P_6^2}(y) = \{v_1, v_2, v_3\}.$$
(19)

Choosing any vertex $x \in V(G) \setminus V(P_6^2)$. If $N_{P_6^2}(x) \subseteq \{v_2, v_3, \ldots, v_6\}$, say $N_{P_6^2}(x) = \{v_3, v_4, v_5\}$ by symmetry, then G contains a copy of C_3^{\Box} , which consists of the triangles $xv_3v_4, v_2v_5v_6$ and the edges xv_5, v_3v_2, v_4v_6 , a contradiction. Hence, we get $N_{P_6^2}(x) \not\subseteq \{v_2, v_3, \ldots, v_6\}$. Without loss of generality, we may assume that $x \in V(G) \setminus V(P_6^2)$ is a vertex with $N_{P_6^2}(x) = \{v_1, v_2, v_3\}$. Moreover, if $V(G) \setminus V(P_6^2)$ contains a vertex y differing from x such that $N_{P_6^2}(y) \neq N_{P_6^2}(x)$, then by symmetry, we assume that either $N_{P_6^2}(y) = \{v_1, v_2, v_4\}$ or $N_{P_6^2}(y) = \{v_1, v_4, v_5\}$. In the former case, it is easy to see that the triangles xv_1v_3, yv_2v_4 and the edges xv_2, v_1y, v_3v_4 form a copy of C_3^{\Box} . In fact, one can find another copy without using vertex x by combining two triangles $yv_1v_2, v_3v_4v_5$ with edges yv_4, v_1v_3, v_2v_5 . In the latter case, we also see that G has a copy of C_3^{\Box} consisting of the triangles $v_1v_2v_3, yv_4v_5$ and the edges v_1y, v_2v_4, v_3v_5 . This is a contradiction. Consequently, we complete the proof of (19) and $H_4 \subseteq G$; see Figure 11.



Figure 11: The graph H_4 .

Since $v_1v_2v_3$ is a triangle, to avoid a copy of C_3^{\Box} in G, we obtain that $G[V(G) \setminus V(P_6^2)]$ is triangle-free. Hence, by Mantel's result (1), we have $e(G) \leq 12 + 3(n-6) + \lfloor \frac{(n-6)^2}{4} \rfloor = 3n-6 + \lfloor \frac{(n-6)^2}{4} \rfloor < f(n)$, as required. In a word, we complete the proof of (15).

In fact, our proof of Theorem 1.2 implies the following stronger result.

Theorem 5.1. Let $n \ge 9$ be an integer. If G is a graph with maximum number of edges over all C_3^{\Box} -free graphs on n vertices, then G is an extremal graph for P_6^2 .

Proof. Recalling the proof of (15), we are done.

Combining with Theorem 4.2, we give the extremal graphs for C_3^{\square} .

Proof of Theorem 1.3. Let G be an n-vertex C_3^{\Box} -free graph with maximum number of edges. In other words, $e(G) = ex(n, C_3^{\Box}) = f(n)$. For n = 6, if G contains P_6^2 as a subgraph, then by Lemma 4.3, we have e(G) = 12 and $G = G_1$; if G is P_6^2 -free, then by Theorem 4.2, we get $G = H_6^3$. For n = 7, if G contains P_6^2 as a subgraph, then the proof of (18) implies that $G = G_2$; if G is P_6^2 -free, then applying Theorem 4.2, we obtain $G \in \{F_7^{4,1}, F_7^{4,4}, H_7^3\}$. For n = 8, if G has a copy of P_6^2 , then the proof of (16) implies that $G = G_3$; if G is P_6^2 -free, then Theorem 4.2 gives that $G \in \{F_8^{4,1}, F_8^{4,4}, F_8^{5,2}, F_8^{5,5}\}$.

When $n \ge 9$, Theorem 5.1 asserts that G is an extremal graph for P_6^2 . Indeed, if $P_6^2 \subseteq G$, then by (15) we know that $e(G) \le f(n) - 1$, a contradiction. Therefore, using Theorem 4.2 again, we can obtain all extremal graph for C_3^{\Box} immediately.

6 Concluding remarks

In Theorem 1.1, we determined the Turán number $ex(n, C_{2k+1}^{\Box})$ and characterized the corresponding extremal graphs for fixed $k \geq 1$ and sufficiently large n. Note that C_{2k}^{\Box} is a bipartite graph, and it is a regular graph with degree 3. For such a spare bipartite graph, we can obtain that for every $k \geq 2$,

$$\exp(n, C_{2k}^{\Box}) = O(n^{5/3}).$$
(20)

Indeed, it can be derived from a result of Füredi [22] that if H is a bipartite graph with maximum degree at most d on one side of the bipartition, then $ex(n, H) = O(n^{2-1/d})$. In 2003, Alon, Krivelevich and Sudakov [1] provided an alternative proof of such a result as one of the applications of the celebrated dependent random choice method; see, e.g., [21] for a description of this powerful probabilistic technique. It seems extremely difficult to determine the magnitude of the Turán number of C_{2k}^{\Box} . For example, in the case k = 2, it reduces to the cube Q_8 , Simonovits [19] proved in 1970 that $ex(n, Q_8) = O(n^{8/5})$. Until now, it remains unknown whether $ex(n, Q_8) = \Theta(n^{8/5})$. Recently, Conlon and Lee [9] conjectured that if H is a bipartite graph with maximum degree at

most d on one side, and H is $K_{d,d}$ -free, then $ex(n, H) = O(n^{2-1/d-\varepsilon})$ holds for some $\varepsilon = \varepsilon(H) > 0$; see [9, 34] for proofs of the case d = 2, and [10, 47] for more related results. Observe that C_{2k}^{\Box} is $K_{3,3}$ -free. Motivated by the above works, we would like to propose the following conjecture, which generalized the bound in (20) exponentially.

Conjecture 6.1. For every $k \ge 2$, there exists $\varepsilon = \varepsilon(k) > 0$ such that

$$\exp(n, C_{2k}^{\square}) = O(n^{\frac{5}{3}-\varepsilon}).$$

After our paper announced on arXiv, Gao, Janzer, Liu and Xu [29] confirmed the conjecture in a stronger form $ex(n, C_{2k}^{\Box}) = \Theta(n^{3/2})$ for every $k \ge 4$. Extremal results for 'twisted' prisms can be found in Bradač, Methuku and Sudakov [5]. Finally, we conclude this paper with an interesting conjecture proposed by Bradač, Janzer, Sudakov and Tomon [4].

Conjecture 6.2 (See [4]). Let T, S be any two trees with at least one edge. There exist positive real numbers c and C such that $cn^{3/2} \leq ex(n, T \Box S) \leq Cn^{3/2}$.

Declaration of competing interest

The authors declare that they have no conflicts of interest to this work.

Data availability

No data was used for the research described in the article.

Acknowledgments

Thanks go to Dr. Zixiang Xu for pointing out the reference [33]. This work was supported by NSFC (Nos. 12271527, 12071484 and 11931002), Natural Science Foundation of Hunan Province (Nos. 2020JJ4675 and 2021JJ40707), the Fundamental Research Funds for the Central Universities of Central South University (Grant No. 2021zzts0034) and the Postdoctoral Fellowship Program of CPSF (No. GZC20233196).

References

- N. Alon, M. Krivelevich, B. Sudakov, Turán numbers of bipartite graphs and related Ramseytype questions, Combin. Probab. Comput. 12 (2003) 477–494.
- [2] H. Bielak, Kieliszek, The Turán number of the graph 2P₅, Discuss. Math. Graph Theory. 36 (3) (2016) 683–694.
- [3] B. Bollobás, Extremal Graph Theory, Academic Press, New York, 1978.
- [4] D. Bradač, O. Janzer, B. Sudakov, I. Tomon, The Turán number of the grid, Bull. London Math. Soc. 55 (2023) 194–204.
- [5] D. Bradač, A. Methuku, B. Sudakov, The extremal number of cycles with all diagonals, (2023), arXiv:2308.16163. See https://arxiv.org/abs/2308.16163.
- [6] N. Bushaw, N. Kettle, Turán numbers of multiple paths and equibipartite forests, Combin. Probab. Comput. 20 (2011) 837–853.

- [7] V. Campos, R. Lopes, A proof for a conjecture of Gorgol, Discrete Appl. Math. 245 (2018) 202-207.
- [8] G. Chen, R.J. Gould, F. Pfender, B. Wei, Extremal graphs for intersecting cliques, J. Combin. Theory. Ser. B 89 (2003) 159–171.
- [9] D. Conlon, J. Lee, On the extremal number of subdivisions, Int. Math. Res. Not. IMRN 12 (2021) 9122–9145.
- [10] D. Conlon, O. Janzer, J. Lee, More on the extremal number of subdivisions, Combinatorica 41 (4) (2021) 465–494.
- [11] G. Dirac, Extensions of Turán's theorem on graphs, Acta Math. Acad. Sci. Hungar. 14 (1963) 417–422.
- [12] T. Dzido, A note on Turán numbers for even wheels, Graphs and Combin. 29 (2013) 1305–1309.
- [13] T. Dzido, A. Jastrzebski, Turán Numbers for odd wheels, Discrete Math. 341 (4) (2018) 1150– 1154.
- [14] P. Erdős, On some new inequalities concerning extremal properties of graphs, stability problems, Theory Graphs (Proc. Colloq. Tihany, 1966) (1968) 77–81.
- [15] P. Erdős, Z. Füredi, R.J. Gould, D.S. Gunderson, Extremal graphs for intersecting triangles, J. Combin. Theory Ser. B 64 (1) (1995) 89–100.
- [16] P. Erdős, T. Gallai, On maximal paths, circuits of graphs, Acta Math. Acad. sci. Hungar. 10 (1959) 337–356.
- [17] P. Erdős, A.H. Stone, On the structure of linear graphs, Bull. Amer. Math. Soc. 52 (1946) 1087–1091.
- [18] P. Erdős, M. Simonovits, A limit theorem in graph theory, Studia Sci. Math. Hungar. 1 (1966) 51–57.
- [19] P. Erdős, M. Simonovits, Some extremal problems in graph theory. Combinatorial theory and its applications, I (Proc. Colloq., Balatonfüred, 1969), North-Holland, Amsterdam, 1970, pp. 377–390.
- [20] P. Erdős, M. Simonovits, An extremal graph problem, Acta Math. Acad. Sci. Hungar. 22 (1971/72) 275–282.
- [21] J. Fox, B. Sudakov, Dependent random choice, Random Structures Algorithms 38 (2011) 68– 99.
- [22] Z. Füredi, On a Turán type problem of Erdős, Combinatorica 11 (1991) 75–79.
- [23] Z. Füredi, D.S. Gunderson, Extremal numbers for odd cycles, Combin. Probab. Comput. 24 (4) (2015) 641–645.
- [24] Z. Füredi, A. Kostochka, J. Verstraëte, Stability in the Erdős–Gallai theorems on cycles and paths, J. Combin. Theory Ser. B 121 (2016) 197–228.
- [25] Z. Füredi, A. Kostochka, R. Luo, J. Verstraëte, Stability in the Erdős–Gallai theorem on cycles and paths II, Discrete Math. 341 (2018) 1253–1263.
- [26] Z. Füredi, On a theorem of Erdős and Simonovits on graphs not containing the cube, in Number Theory, Analysis, and Combinatorics, Proceedings of the Paul Turán Memorial Conference, Budapest, 2013, pp. 113–125. See https://arxiv.org/abs/1307.1062.
- [27] Z. Füredi, M. Simonovits, The history of degenerate (bipartite) extremal graph problems, in Erdős centennial, Bolyai Soc. Math. Stud., 25, János Bolyai Math. Soc., Budapest, 2013, pp. 169–264. See https://arxiv.org/abs/1306.5167v2.
- [28] R.J. Faudree, R.H. Schelp, Path Ramsey numbers in multicolorings, J. Combin. Theory. Ser.

B 19 (1975) 150–160.

- [29] J. Gao, O. Janzer, H. Liu, Z. Xu, Extremal number of graphs from geometric shapes, 2023, arXiv: 2303.13380. Israel Journal of Mathematics, to appear.
- [30] X. Hou, Y. Qiu, B. Liu, Extremal graph for intersecting odd cycles, Electron. J. Combin. 23 (2) (2016) P2.29.
- [31] X. Hou, Y. Qiu, B. Liu, Turán number and decomposition number of intersecting odd cycles, Discrete Math. 341 (2018) 126–137.
- [32] T. Jiang, A. Newman, Small dense subgraphs of a graph, SIAM J. Discrete Math. 31 (2017) 124–142.
- [33] T. Jiang, J. Ma, L. Yepremyan, On Turán exponents of bipartite graphs, Combin. Probab. Comput. 31 (2022) 333–344.
- [34] O. Janzer, Improved bounds for the extremal number of subdivisions, Electron. J. Combin. 26 (2019) Paper No. 3.3.
- [35] G.N. Kopylov, Maximal paths, cycles in a graph, Dokl. Akad. Nauk SSSR, 234 (1) (1977) 19–21. English translation: Soviet Mathh. Dokl. 18(3), (1977) 593–596.
- [36] Y. Lan, Z. Qin, Y. Shi, The Turán number for $2P_7$, Discuss. Math. Graph Theory. 39 (2019) 805–814.
- [37] H. Liu, Extremal graphs for blow-ups of cycles, trees, Electron. J. Combin. 20 (1) (2013), P65.
- [38] J. Ma, B. Ning, Stability results on the circumference of a graph, Combinatorica 40 (1) (2020) 105–147.
- [39] W. Mantel: Problem 28, soln. by H. Gouventak, W. Mantel, J. Teixeira de Mattes, F. Schuh, W.A. Wythoff. Wiskundige Opgaven 10 (1907) 60–61.
- [40] Z. Ni, L. Kang, E. Shan, H. Zhu, Extremal graphs for blow-ups of keyrings, Graphs and Combin. 36 (2020) 1827–1853.
- [41] R. Pinchasi, M. Sharir, On graphs that do not contain the cube and related problems, Combinatorica 25 (2005) 615–623.
- [42] M. Simonovits, A method for solving extremal problems in graph theory, stability problems, Theory of Graphs (Proc. Colloq., Tihany, 1966) (1968) 279–319.
- [43] M. Simonovits, Extremal graph problems with symmetrical extremal graphs, additionnal chromatic conditions, Discrete Math. 7 (1974) 349–376.
- [44] M. Simonovits, The extremal graph problem of the icosahedron, J. Combin. Theory Ser. B 17 (1974), 69–79.
- [45] M. Simonovits, Extremal problems, graph products, Studies in Pure Mathematics, pp. 669– 680, Akadémiai Kiadó, Birkhäuser Verlag 1982.
- [46] M. Simonovits, Paul Erdős' influence on extremal graph theory, in The Mathematics of Paul Erdős II, R.L. Graham, Springer, New York, 2013, pp. 245–311.
- [47] B. Sudakov, I. Tomon, Turán number of bipartite graphs with no $K_{t,t}$, Proc. Amer. Math. Soc. 148 (7) (2020) 2811–2818.
- [48] P. Turán, On an extremal problem in graph theory, Mat. Fiz. Lapok 48 (1941), pp. 436–452. (in Hungarian).
- [49] C. Xiao, G.O.H. Katona, J. Xiao, O. Zamora, The Turán number of the square of a path, Discrete Appl. Math. 307 (2022) 1–14.
- [50] L.-T. Yuan, X.-D. Zhang, The Turán number of disjoint copies of paths, Discrete Math. 340

(2) (2017) 132–139.

- [51] L.-T. Yuan, X.-D. Zhang, Turán numbers for disjoint paths, J. Graph Theory 98 (3) (2021) 499–524.
- [52] L.-T. Yuan, Extremal graphs for the k-flower, J. Graph Theory 89 (2018) 26–39.
- [53] L.-T. Yuan, Extremal graphs for odd wheels, J. Graph Theory 98 (2021) 691–707.
- [54] L.-T. Yuan, Extremal graphs for edge-blow up of graphs, J. Combin. Theory Ser. B 152 (2022) 379–398.
- [55] L.-T. Yuan, Extremal graphs of the *p*-th power of paths, European J. Combin. 104 (2022) 103548.