

CONJECTURES ABOUT VIRTUAL LEGENDRIAN KNOTS AND LINKS

VLADIMIR CHERNOV, RUSTAM SADYKOV

ABSTRACT. We formulate conjectures generalizing some known results to the category of virtual Legendrian knots. This includes statements relating virtual Legendrian knots to ordinary Legendrian knots, non-existence of positive virtual Legendrian self isotopies for the class of the fiber of ST^*M and the conjectural relation of virtual Legendrian isotopies to causality in generalized spacetimes. We prove the conjectures in the case of 2-dimensional M and $(2 + 1)$ -dimensional spacetimes. We also formulate and prove the version of the Arnold's 4 cusp conjecture for virtual isotopies.

1. MAIN RESULTS AND DEFINITIONS

We work in the smooth category, all maps and manifolds are assumed to be smooth and the word “smooth” means C^∞ . All manifolds, links and knots are assumed to be oriented and connected (unless we talk about links).

A *contact manifold* N^{2k+1} is an odd dimensional manifold equipped with a hyper plane distribution ξ that can locally be given as the kernel of a 1-form α with $\alpha \wedge d\alpha^k$ nowhere zero. A k -dimensional manifold L^k is *Legendrian* if it is everywhere tangent to the hyperplanes of the contact structure and it is *transverse* if it is nowhere tangent to the contact hyperplanes.

The standard examples of contact manifolds are the spherical cotangent bundle ST^*M and the first jet space J^1M which have natural contact structures. Note that a surgery on M resulting in a manifold M' induces a modification of contact manifolds $ST^*M \mapsto ST^*M'$.

A *virtual Legendrian isotopy* in ST^*M is a sequence of Legendrian isotopies, modifications of Legendrian submanifolds in ST^*M induced by surgery on M away from the front projection of the Legendrian submanifold, and contactomorphisms of ST^*M induced by diffeomorphisms of the manifold M . Virtual Legendrian knots were introduced in the work of Cahn and Levi [10].

The virtual Legendrian isotopy is *subcritical* if the index of the attached handles is less than half of the dimension of M and it is *allowable* if the index of the attached handles does not exceed half of the dimension of M . In particular, the virtual Legendrian isotopies of [10] that do not involve deleting handles are allowable. *All the conjectures we formulate are for the allowable virtual Legendrian isotopies.*

Note that a surgery on M induces a modification on J^1M and virtual Legendrian isotopies in J^1M are defined similarly to the virtual Legendrian isotopies on ST^*M . In a similar fashion one can define virtual transverse isotopies both for ST^*M and J^1M . We note that in contrast to virtual isotopies in ST^*M , in the case of virtual

2020 *Mathematics Subject Classification.* Primary 53C24; Secondary 53C50, 57D15, 83C75.

Key words and phrases. virtual knot, Legendrian knot, causality, spacetime, linking, Low conjecture, positive Legendrian isotopy, non-negative Legendrian isotopy.

Legendrian isotopies in J^1M the projection of a virtual Legendrian knot to M is of codimension 0. In both cases, the surgery of virtual Legendrian knots is performed on the complement in M to the projection of the virtual Legendrian knot.

Classical virtual knots were introduced by Kauffman [24] and they are ordinary knots in a thickened surface $F \times \mathbb{R}$ considered up to a sequence of isotopies and modifications of $F \times \mathbb{R}$ induced by surgery on F away from a projection of a knot (or a link). A Gauss diagram of a knot is an oriented chord diagram on a circle. The chords connect the preimages of double points of a diagram projection and are oriented from the preimage of an upper strand to the preimage of a lower strand. They are equipped with signs ± 1 which is the writhe of the corresponding diagram crossing. Polyak and Viro [32] noted that a Gauss diagram of a knot uniquely defines the knot on $S^2 \times \mathbb{R}$.

Gauss diagrams can be considered up to the formal diagrammatic version of the usual knot Reidemeister moves however using these moves it is easy to get a Gauss diagram which does not correspond to a planar knot diagram and hence is virtual. Carter, Kamada, Saito [11] proved that formal Gauss diagrams considered up to these formal Reidemeister moves give the same equivalence classes as the ones of virtual knots.

As it is well known the cooriented front projection of a Legendrian knot in ST^*M to M completely determines the Legendrian knot. Such front projections on a surface can be described by diagrams similar to Gauss diagrams with the symbols on chords denoting the crossing and cusp types. Cahn and Levi [10] proved that equivalence classes of such generalized ‘‘Gauss diagrams’’ for front projections to a surface coincide with the equivalence classes of virtual Legendrian knots in ST^*M .

A natural question is if a similar result holds for virtual Legendrian knots and links in J^1M or for ST^*M for higher dimensional manifold M closed or non-closed.

2. VIRTUAL POSITIVE LEGENDRIAN ISOTOPY

Recall that a Y_ℓ^x manifold is a Riemannian manifold (M, g) such that there is a point $x \in M$ and a positive number $\ell > 0$ such that all unit speed geodesics starting from x return back to x in time ℓ . For such manifolds the cogeodesic flow in ST^*M induces a positive Legendrian isotopy of the sphere fiber over x to itself. Recall that a Legendrian isotopy is *positive* if the evaluation of the contact form on all the velocity vectors of all the points and at all times of the deformed Legendrian submanifold is positive. *Non-negative* Legendrian isotopies are defined similarly.

The classical result of Berard-Bergery [3], [7] says that if M admits a Riemannian metric making it into a Y_ℓ^x manifold then the universal cover of M is compact (and hence M has finite $\pi_1(M)$) and the rational cohomology algebra of its universal cover is generated by one element. In the works of the first author and Nemirovski we proved [14, 15] that if M is such that there is a non-constant non-negative Legendrian isotopy of a fiber of ST^*M over a point of M back to itself then the universal cover of M is compact. So in this work we conjectured that if such a non-negative Legendrian isotopy exists, then the rational cohomology algebra of the universal cover is generated by one element.

A slightly stronger version of this conjecture for positive Legendrian isotopies was proved by Frauenfelder, Labrousse and Schlenk [20] who formally proved more than we conjectured and showed that the integer cohomology ring of the universal cover is the one of a CROSS (Compact Rank One Symmetric Space), i.e. a sphere or one

of the projective spaces over $\mathbb{C}, \mathbb{H}, \mathbb{C}a$. In particular this means that the conclusions of the Berard-Bergery theorem and of the Bott-Samelson theorem [9], [7] are the same. The result of [20] can be generalized to non-negative Legendrian isotopies by using the result of Nemirovski and the first author [16] saying that if there is a non-negative Legendrian isotopy then there is a positive Legendrian isotopy as well, even though this positive Legendrian isotopy is obtained not by deforming the non-negative Legendrian isotopy but rather by deforming a certain power of it. (See also Colin, Ferrand and Pushkar [19] and [14] for the case of positive isotopies of a fiber to itself in the case where M is coverable by an open domain in \mathbb{R}^m .)

Currently the only simply connected spaces known to us that satisfy these algebraic topology conditions but are not diffeomorphic to a CROSS are exotic spheres, see Besse [7]. So it could be that this algebraic topology condition means that the simply connected manifold is homeomorphic to a CROSS.

We define *strong virtual Legendrian isotopies* to be the virtual Legendrian isotopies during which the universal cover of the underlying manifold M is never homeomorphic to a CROSS.

Now we formulate our first conjecture

2.1. Conjecture. There exists no strong allowable positive (or non-constant non-negative) virtual Legendrian isotopy of the sphere fiber of ST^*M to itself.

2.2. Remark. A similar statement is false if we allow all positive virtual Legendrian isotopies rather than only the allowable ones. To see this consider a positive Legendrian isotopy of the fiber over a point of a surface M^2 to the knot whose front projection is a small outwards cooriented circle K . Let K_- and K_+ be circular fronts with the same center as K cooriented outwards with radii respectively a bit smaller and a bit larger than K . Now perform an index one surgery such that one of the feet of the glued handle is the disk bounded by K_- . Next perform the index two surgery along K_+ . As a result we get a small inwards cooriented front which clearly admits a positive Legendrian isotopy to the sphere fiber. Thus positive virtual Legendrian isotopies of a fiber of ST^*M to itself do exist if we do not require the virtual Legendrian isotopy to be allowable. It could however be that to avoid this phenomenon it is enough to prohibit only some of the surgeries involving handles of index more than half of the dimension of M^m (for example index m handles) rather than all of them.

2.3. Theorem. *The statement of Conjecture 2.1 is true in the case where M is a surface.*

Proof. Suppose that M is a closed surface. Assume to the contrary that a spherical fiber K_0 of ST^*M admits an allowable non-negative virtual Legendrian isotopy to another fiber. We may assume that the surface M is connected, and consider only surgeries of index 1 which we call stabilizations. Then an allowable non-negative virtual Legendrian isotopy is a composition of finitely many stabilizations of the surface and positive Legendrian isotopies.

Let M_k denote the surface obtained from M by k stabilizations. Suppose that the number of stabilizations is n , and n is minimal in the sense that there is no allowable non-negative virtual Legendrian isotopy with less than n stabilizations of K_0 to another fiber. Then a part of the allowable non-negative Legendrian isotopy takes K_0 to a Legendrian knot K_{n-1} in ST^*M_{n-1} that satisfies two properties:

- the Legendrian knot K_{n-1} does not admit a non-negative Legendrian isotopy to a fiber in ST^*M_{n-1} , and
- there is a non-negative Legendrian isotopy of the stabilization $K_n \subset ST^*M_n$ of the knot K_{n-1} to a fiber in ST^*M_n .

The surface M_n is obtained from the surface M_{n-1} by removing two discs $2D$ in the complement to $p(K_{n-1})$, where p is the projection of ST^*M_{n-1} to M_{n-1} , and attaching along the created boundary a handle $H = [-1, 1] \times S^1$. Let C denote the meridian $\{0\} \times S^1$ on the handle H . There is an infinite cover $\pi: M_n \setminus C \rightarrow M_n$ that restricts to the identity map on $M_n \setminus H$. Since the knot K_n is isotopic to a fiber in ST^*M_n , it admits a lift π^*K_n to a Legendrian knot in $\pi^*ST^*M_n$. Furthermore, the non-negative Legendrian isotopy of K_n to a fiber in ST^*M_n lifts to a non-negative Legendrian isotopy of π^*K_n to a fiber in $\pi^*ST^*M_n$.

On the other hand, the space $\pi^*(ST^*M_n)$ can be identified with the spherical cotangent bundle of $M_n \setminus C$, which in its turn is contactomorphic to the spherical cotangent bundle of $M_n \setminus H \cong M_{n-1} \setminus 2D$. Furthermore, we may choose a contactomorphism $\psi: \pi^*(ST^*M_n) \cong ST^*(M_{n-1} \setminus 2D)$ so that it covers a diffeomorphism $M_n \setminus C \rightarrow M_{n-1} \setminus 2D$ that restricts to the map that identifies the complement to a small neighborhood of H in M_n with the complement to a small neighborhood of $2D$ in M_{n-1} . In particular, the contactomorphism ψ takes the Legendrian knot K_n to the Legendrian knot K_{n-1} . Then the non-negative Legendrian isotopy of K_n to a fiber in ST^*M_n lifts to a non-negative Legendrian isotopy of K_{n-1} to a fiber in ST^*M_{n-1} .

This contradicts the minimality of n . Therefore the statement of Conjecture 2.1 is true for closed surfaces.

Suppose that the surface M is noncompact of finite genus. Then it is an open submanifold of a closed surface M' of finite genus. As above, we obtain surfaces M'_{n-1} and M'_n where M'_n is obtained from M'_{n-1} by a surgery of index 1. Again, let C be a meridian of the attached handle. Then there is a cover $\pi': M'_n \setminus C \rightarrow M'_n$. Similarly, we obtain open submanifolds $M_{n-1} \subset M'_{n-1}$ and $M_n \subset M'_n$. We may assume that $C \subset M_n$. Then there is a cover $\pi: M'' \rightarrow M_n$ obtained by restricting the map π' to $(\pi')^{-1}M_n$. On the other hand, the surface M'' can be identified with a subsurface of $M \setminus 2D$, where $2D$ is a disjoint union of two discs along which the 1-handle is attached. Therefore, the non-negative Legendrian isotopy of a knot in ST^*M_n lifts to a non-negative Legendrian isotopy of the corresponding knot in ST^*M_{n-1} .

Suppose now that M is of infinite genus. Then the trace of a non-negative virtual Legendrian isotopy is contained in a compact part of all the corresponding surfaces. Thus this case can be reduced to the case where M is of finite genus. \square

2.4. Corollary. *Let M be a surface and let L be a two component Legendrian link in ST^*M that admits a strong allowable virtual Legendrian isotopy to a pair of sphere fibers. Then there is **no** non-negative strong allowable virtual Legendrian isotopy of one component of L to the other.*

Proof. The proof is similar to the one of Theorem 2.3. Indeed, the argument in the proof of Theorem 2.3 shows that the existence of an allowable virtual Legendrian isotopy of L to a pair of sphere fibers implies the existence of a Legendrian isotopy of L to a pair of fibers in ST^*M . Consequently, there is a contactomorphism c_1 of ST^*M that takes L to a link L' that consists of two sphere fibers of ST^*M .

Similarly, the argument in the proof of Theorem 2.3 shows that the existence of a non-negative allowable virtual Legendrian isotopy from one component of L to the other implies the existence of a non-negative Legendrian isotopy φ_t from one component of L to the other in ST^*M . Then $c_1 \circ \varphi_t$ is a non-negative Legendrian isotopy from one sphere fiber in L' to the other, say a fiber F to the fiber F' .

There is a contactomorphism c_2 from ST^*M to itself that takes the fiber F to F' and takes F' to F . Then $c_2 \circ c_1 \circ \varphi_t$ is a non-negative Legendrian isotopy that takes the fiber F' to F .

Finally, the composition of non-negative Legendrian isotopies $c_1 \circ \varphi_t$ and $c_2 \circ c_1 \circ \varphi_t$ is a non-constant non-negative Legendrian isotopy from a fiber of ST^*M to itself. Consequently, M is a sphere, see [14]. However Corollary 2.4 is about **strong** allowable virtual Legendrian isotopy so M can not be a sphere. Thus, there is no non-negative strong allowable virtual Legendrian isotopy of one component to the other. \square

2.5. Conjecture. We conjecture that for M^m , $m \geq 3$, whose universal cover is not homeomorphic to CROSS and L that is a two component Legendrian link in ST^*M that admits an allowable virtual Legendrian isotopy to a pair of fibers, there is **no** non-negative **strong** allowable virtual Legendrian isotopy of one component of L to the other.

3. GENERALIZATION OF KUPERBERG RESULT TO VIRTUAL LEGENDRIAN KNOTS AND LINKS IN HIGHER DIMENSIONS

By Kuperberg theorem [25], if two ordinary virtual knots in $M^2 \times \mathbb{R}$ are virtually isotopic then they both can be simplified by a sequence of isotopies and modifications induced by surgeries on M^2 that decrease to the situation where two knots are related by an automorphism of $M^2 \times \mathbb{R}$ induced by an automorphism of M^2 . The similar result for virtual Legendrian knots **in ST^*M^2 , where M^2 is closed** was proved by us in [18].

In particular we got the following corollary: if two Legendrian knots in ST^*S^2 are virtual Legendrian isotopic then they are Legendrian isotopic. This follows immediately since the genus of S^2 can not be further decreased.

For higher dimensional virtual Legendrian knots it is not clear what surgeries decrease the complexity of the underlying manifold or even what this complexity is. However below we formulate the conjecture generalizing the above corollary in our work.

3.1. Conjecture. Two Legendrian links in ST^*S^m or ST^*CP^n are Legendrian isotopic if and only if they are virtually Legendrian isotopic. (This conjecture of the first author in the case of spheres was first formulated in [10].) The same conjecture is formulated for Legendrian links in J^1S^m and J^1CP^n . It is also interesting to know if a similar statement holds for the fibers of ST^*M where M is a CROSS.

4. VIRTUAL LEGENDRIAN ISOTOPIES AND CAUSALITY

Low [26, 27, 28, 29] formulated the conjecture relating topological linking of the skies to causality. Recall that a *sky* is the sphere of light rays through a point, and it is located in the contact manifold of all future directed unparameterized light rays (null geodesics). Recall that two points in a spacetime are causally related if

information traveling with the light or less can reach from one event point to the other, i.e. if there is a causal (non-spacelike) curve between these points.

Low's conjecture is for globally hyperbolic spacetimes which are defined as spacetimes without closed causal trajectories (time travel machines) and with the assumption that the intersection of causal future and causal past of any two points is compact (absence of naked singularities when a traveling particle disappears from the observable universe of the observer without crossing their line of sight), see [6], or [23] for a more traditional and equivalent definition. Globally hyperbolic spacetimes form the most important and studied class of spacetimes and one of the versions of Penrose's Strong Cosmic Censorship conjecture [31] says that all physically relevant spacetimes are globally hyperbolic. (The part of a spacetime located inside of black holes is not relevant to us since the particles can not escape the black hole horizons.)

Geroch [22] proved that globally hyperbolic spacetimes are homeomorphic to $\Sigma \times \mathbb{R}$ and Bernal-Sanchez [4, 5] proved that it is in fact diffeomorphic to $\Sigma \times \mathbb{R}$ with each $\Sigma \times t$ being a *spacelike Cauchy surface* i.e. a codimension one submanifold intersected at exactly one point by every unextendible causal curve and such that the restriction of the Lorentz metric to it is Riemannian.

As it was observed by Low for such spacetimes the contact manifold of all future directed unparameterized light rays (light geodesics) is contactomorphic to $ST^*\Sigma$ for every spacelike Cauchy surface Σ .

Low conjecture is for topological linking in $(2 + 1)$ -dimensional spacetimes and Σ being an open 2-dimensional surface. This conjecture was reformulated for Legendrian linking for $(3 + 1)$ -dimensional spacetimes with Σ homeomorphic to \mathbb{R}^3 by Nataro and Tod [30]. All these conjectures were proved in the works of first author and Nemirovski [14, 15, 13] and the current state of knowledge is that Legendrian linking is equivalent to causality for globally hyperbolic spacetimes of all dimensions provided that the universal cover of the Cauchy surface is not compact **or** that it is compact but the integer cohomology of the universal covering is not the one of a CROSS.

Antoniou, Kauffman and Lambropoulou [2] suggested that singular points of a Morse function on a spacetime can be considered as singularities corresponding to formation of black holes or something similar. The classical spacetimes are time oriented Lorentz manifolds. We generalize this notion as follows, a *generalized spacetime* is a manifold equipped with a bilinear symmetric form. Furthermore we assume that it can be equipped with a Morse function such that away from the critical points the form restricts to a Lorentz metric, we also assume that the Morse function is a timelike function away from the critical points, i.e. the velocity vector of the flow lines of the gradient of the Morse function dot product with itself is negative. We also assume that the non-critical level sets of the Morse function are spacelike Cauchy surfaces. This definition is in the spirit of Borde et al [8] and Sorkin [33] Morse spacetimes. (See also Garcia-Haveling [21] for some recent developments on the subject of Morse spacetimes.)

We do not quite agree with [2] on the interpretation of surgeries on a spacelike manifold as black hole formations, (because the topology of the level set of a Morse function does not have to change under passage through a black hole formation). However we observe that if the chunks of a generalized spacetime between critical points of a Morse function f are globally hyperbolic and the indexes of the critical

points of the Morse function $-f$ are such that they correspond to allowable surgeries on level sets; then a sphere of light rays through a point considered as a Legendrian knot in ST^*M (for M that is the spacelike level set of a noncritical level of the Morse timelike function) experiences an allowable virtual Legendrian isotopy when passing through the critical level in the direction from higher levels to lower levels. We call such generalized spacetimes *allowable*.

Clearly the definition of causality still makes sense for such generalized spacetimes and we formulate

4.1. Conjecture. Two points in a generalized allowable spacetime are causally related if and only if the Legendrian link of spheres of light rays through these two points is not strongly allowable virtually Legendrian isotopic to the link of two fibers of ST^*M_t of the level set manifold M_t . These Legendrian spheres can be considered in ST^*M_t for any M_t that is a non-critical level of the Morse function. We call such links non-trivial.

The last links correspond to causally unrelated events. (The reason why we have to consider strong allowable virtual Legendrian isotopy rather than usual virtual Legendrian isotopy is that Legendrian linking is not equivalent to causality when a Cauchy surface admits a Y_{ℓ}^x Riemann metric [14].)

Note also that if the two points are on the common light ray then the corresponding link of spheres of light rays has a double point and hence is singular. We call such singular links nontrivial as well the links described above.

4.2. Remark (How can one try to prove the above conjecture). This conjecture is closely related to the one about nonexistence of allowable strong non-negative non-constant virtual Legendrian isotopy of a fiber to itself. The relation is similar to the ones of the paper of Nemirovski and the first author [14]. Namely the transport of a point in a spacetime to the past with the speed less or equal than light (causal trajectory) induces a non-negative Legendrian isotopy of the sphere of all light rays through the point in the contact manifold of all light rays, see [14] for timelike trajectories and [17] for causal trajectories.

Given a Morse function f on a manifold M and any subset $I \subset \mathbb{R}$, let M_I denote the subset $f^{-1}(I)$ of M . To simplify notation, we will write M_t instead of $M_{\{t\}}$ in the case where $I = \{t\}$.

Let p and q be two events in a generalized spacetime M equipped with a Morse function f that is timelike in the complement to critical points. Suppose that $f(p) < f(q)$. We may slightly perturb the function f so that both events p and q belong to regular levels of the function f . Let $c_1 < \dots < c_n$ be all critical values in the interval $[f(p), f(q)]$. Choose regular values a_i so that

$$f(p) = a_0 < c_1 < a_1 < c_2 < \dots < a_{n-1} < c_n < a_n = f(q).$$

We observe that the manifold $M_{(c_i, c_{i+1})}$ is globally hyperbolic, while the manifold of all light rays \mathbf{L}_i on $M_{(c_i, c_{i+1})}$ can be canonically parametrized by the spherical cotangent bundle ST^*M_a of the level M_a for any $a \in (c_i, c_{i+1})$. These parametrizations are quite different. For example, a fiber of the cotangent bundle with respect to one canonical parametrization is not a fiber of the cotangent bundle with respect to another canonical parametrization. For a point x in $M_{[a_0, a_n]}$ we will denote the sphere of light rays in \mathbf{L}_i emanated from x by $S_i(x)$.

Suppose that the events p and q are causally unrelated. Choose a future directed timelike path q^t , where $t \in [a_0, a_n]$, such that $f(q^t) = t$, $q^{a_n} = q$, and the path q^t

does not pass through critical points of the function f . As t increases in the interval $[a_0, c_1)$, the sphere $S_0(q^t)$ is modified by a Legendrian isotopy in \mathbf{L}_0 . Similarly, as t increases in (c_i, c_{i+1}) for $i = 1, \dots, n-1$, the path q^t defines a Legendrian isotopy of $S_i(q^t)$ in \mathbf{L}_i , and as t increases in $(c_n, a_n]$ the path q^t defines a Legendrian isotopy of $S_n(q^t)$ in \mathbf{L}_n . Also as t varies in the considered intervals the sphere of light rays $S_i(p)$ in \mathbf{L}_i for $i = 0, \dots, n$ remains constant.

Recall that the manifold of light rays \mathbf{L}_i is canonically identified with the manifold $ST^*M_{a_i}$. In particular, every self-diffeomorphism of M_{a_i} induces a contactomorphism of $ST^*M_{a_i}$. Recall also that an elementary virtual Legendrian modification of a knot K in ST^*N consists of a modification of ST^*N induced by a surgery on N away from the front projection of K resulting in a manifold N' , together with a contactomorphism of ST^*N' induced by a self-diffeomorphism of N' .

It follows that up to a contactomorphism of \mathbf{L}_{i+1} induced by a diffeomorphism of the level $M_{a_{i+1}}$, the pair $(\mathbf{L}_{i+1}, S_{i+1}(q^{a_{i+1}}))$ is obtained from $(\mathbf{L}_i, S(q^{a_i}))$ by elementary virtual Legendrian modifications. We emphasize that at this step we do not identify light rays in $ST^*M_{a_{i+1}}$ with light rays in $ST^*M_{a_i}$. Instead, we make a discrete transformation using a handle decomposition of $M_{[a_i, a_{i+1}]}$ associated with the restricted Morse function $f|_{M_{[a_i, a_{i+1}]}}$. Each handle corresponds to one elementary virtual Legendrian modification. Under such a transformation, the complement to a neighborhood of the attaching sphere in M_{a_i} is identified with the complement to a neighborhood of the belt sphere in $M_{a_{i+1}}$. This identification extends to an identification of spherical cotangent bundles. Under this identification the fiber $S_{i+1}(q^{a_{i+1}})$ of the cotangent bundle $\mathbf{L}_{i+1} = ST^*M_{a_{i+1}}$ is identified with the fiber $S(q^{a_i})$ of the cotangent bundle $\mathbf{L}_i = ST^*M_{a_i}$.

Similarly, we claim that as t increases from $c_{i+1} - \varepsilon$ to $c_{i+1} + \varepsilon$ crossing a critical level c_i for $i = 1, \dots, n$, the sphere of light rays $S_i(p)$ changes to a sphere of light rays $S_{i+1}(p)$ by finitely many elementary virtual Legendrian modifications. Indeed, we may assume that the critical level c_i contains only one critical point. Then the complement to a neighborhood of the attaching sphere in $M_{c_{i+1}-\varepsilon}$ can be identified with the complement to a neighborhood of the belt sphere in $M_{c_{i+1}+\varepsilon}$ by means of the gradient flow of f . The above identification extends to an identification ψ of spherical cotangent bundles. Under this identification the sphere of light rays $S_{i+1}(p)$ in $\mathbf{L}_{i+1} = ST^*M_{c_{i+1}+\varepsilon}$ may not coincide with the image $\psi(S_i(p))$ of the sphere $S_i(p)$ in $\mathbf{L}_i = ST^*M_{c_{i+1}-\varepsilon}$ since the trajectories of the gradient flow of f in the definition of ψ are fairly arbitrary timelike trajectories that do not have much relation to the cones of light rays. However, this can be rectified by a contactomorphism of \mathbf{L}_{i+1} induced by a self-diffeomorphism of $M_{c_{i+1}+\varepsilon}$.

Thus, we have constructed a virtual Legendrian isotopy of the sphere $S_0(p)$ of light rays to the sphere $S_n(p)$ and a virtual Legendrian isotopy of the sphere $S_0(q^0)$ to $S_n(q)$. Furthermore, for each i , the sphere of light rays of p in \mathbf{L}_i is disjoint from the sphere of light rays of q in \mathbf{L}_i since otherwise there would exist a causal trajectory from p to q for causally unrelated events p and q . Thus, we have constructed a virtual Legendrian isotopy of the link of two fibers $(S_0(p), S_0(q^0))$ to the link $(S_n(p), S_n(q))$. The reverse isotopy is a strong allowable virtual Legendrian isotopy to a link of two fibers.

To prove the converse statement of the conjecture, suppose that p and q are causally related. Then there is a causal path q^t with $q^0 = p$ and $q^n = q$. The argument similar to the one above shows now that there is a non-trivial non-negative

Legendrian isotopy of the fiber $S_n(q^n)$ to the fiber $S_0(q^0)$. Indeed, as t varies in the past direction in one of the intervals $[a_0, c_1)$, (c_i, c_{i+1}) or $(c_n, a_n]$, the sphere $S_i(q^t)$ is modified by a non-trivial non-negative virtual Legendrian isotopy. As t passes a critical point c_{i+1} in the negative direction, the sphere $S_{i+1}(q^t)$ is modified by finitely many elementary virtual Legendrian modifications to a sphere $S_i(q^t)$. We emphasize that we never consider the sphere $S_n(q^t)$ for small values of t since the light rays of q^t may not reach the level a_n because of the critical points. As t becomes smaller than c_1 , the sphere $S_0(q^t)$ turns into the fiber $S_0(q^0)$ in $\mathbf{L}_0 = ST^*M_{a_0}$.

Next we postcompose the resulting non-trivial non-negative virtual Legendrian isotopy from $S_n(q^n)$ to $S_0(q^0)$ with the virtual Legendrian isotopy of $S_0(p) = S_0(q^0)$ to $S_n(p)$ that consists of elementary virtual Legendrian modifications and contactomorphisms of cotangent bundles induced by diffeomorphisms of underlying manifolds. The resulting isotopy φ is a non-trivial non-negative virtual Legendrian isotopy of one component of the link $(S_n(p), S_n(q))$ in \mathbf{L}_n to the other.

Assume now contrary to the statement of the conjecture that despite the existence of a causal path from p to q the link $(S_n(p), S_n(q))$ is an unlink, i.e., there is a contactomorphism ψ of \mathbf{L}_n that takes the link $(S_n(p), S_n(q))$ to the link that consists of two fibers. Then ψ takes φ to a non-negative non-trivial Legendrian isotopy that takes one fiber to another.

By Corollary 2.4, this implies the conjecture for $2 + 1$ -dimensional allowable generalized spacetimes.

For higher dimensional allowable generalized spacetimes the conjecture follows in a similar way provided that Conjecture 2.5 is true.

In particular we proved the following Theorem

4.3. Theorem. *The statement of conjecture 4.1 is true for $(2 + 1)$ -dimensional generalized allowable spacetimes.*

4.4. Remark. The proposed approach to the proof of Conjecture 4.1 is primarily based on the conjecture that there is no non-negative allowable virtual Legendrian isotopy of a sphere fiber of ST^*M to itself. Allowable virtual isotopies prohibit surgeries on M that involve gluing handles of index more than half of the dimension of M . However the only example we know where in the usual non-negative non-constant virtual Legendrian of a fiber to itself exists involves handles of index equal to the dimension of M . So it could be that these conjectures are true if one relaxes the definition of the allowable virtual Legendrian isotopy and prohibits only some of the surgeries involving handles of index more than half of the dimension of M^m (for example index m handles) rather than all of them.

5. A FEW WORDS ABOUT NON-EXISTENCE OF THE GENERALIZATION OF ARNOLD'S 4-CUSP CONJECTURE

Arnold [1] formulated the following conjecture. Consider two Legendrian knots in $ST^*\mathbb{R}^2$ that are the slightly perturbed circle-fibers that project to the inward and the outward cooriented oval fronts. Then every generic Legendrian isotopy between them involves a moment where the front projection has four cusps.

Chekanov and Pushkar [12] proved this conjecture.

Using standard covering techniques $\mathbb{R}^2 \rightarrow F^2$ it is clear that Arnold's conjecture is true for all surfaces other than a sphere. For the sphere the cogeodesic flow takes

the expanding circle around the North pole to the contracting circle around the South pole. This propagation does not involve any cusps and the statement of the conjecture would be false.

The conjecture also does not hold for virtual Legendrian knots.

Indeed, consider an outward growing small circular planar wave front and perform an index 1-surgery on the plane by gluing a handle one of whose feet is a disk inside of our front. Propagate the front a bit along the handle and do the reverse surgery via cutting along a circle located above the front. Now our front appears to be a contracting circle and no cusps appeared in this process at all.

However the following Theorem is true.

5.1. Theorem. *Let K_1 and K_2 be two Legendrian knots in $ST^*M_1^2$ and ST^*M_2 that correspond to the small circular inside and outside cooriented fronts respectively. Assume moreover that $M_i \neq S^2$. Then every generic strong allowable virtual Legendrian isotopy from K_1 to K_2 involves a moment when the front projection of the knot has at least four cusps.*

Proof. Take a strong allowable virtual Legendrian isotopy of K_1 to K_2 . It consists of

- elementary Legendrian modifications resulting in surfaces M_0, M_1, \dots, M_n ,
- contactomorphisms, which we omit to simplify notations, and
- isotopies $\varphi_t^{(i)}$, $t \in [0, 1]$ of Legendrian knots $K^{(i)} \subset ST^*M_i$ satisfying $K^{(0)} = K_1$, $\varphi_1^{(n)}(K^{(n)}) = K_2$, and the property that $K^{(i+1)}$ is the stabilization of $\varphi^{(i)}(K^{(i)})$.

For $i = 1, \dots, n$, the manifold M_i is obtained from M_{i-1} by attaching a handle. Let $S_i \subset M_i$ denote the meridian of the handle attached to M_{i-1} .

As in the proof of Theorem 2.3, we observe that there is a covering $M_i \setminus S_i \rightarrow M_i$, and that $M_i \setminus S_i$ can be identified with a subset of M_{i-1} . Therefore an isotopy in M_i of a knot in $M_i \setminus S_i$ can be lifted to an isotopy in M_{i-1} . Thus, starting with $i = n$, by induction with decreasing i , we may lift the family of isotopies φ_t^i to a Legendrian isotopy from K_1 to a knot with an inside cooriented circular front. By the Arnold's conjecture for surfaces, we deduce that for the resulting Legendrian isotopy there is a moment at which the front projection of the knot has at least four cusps. Therefore, the same is true for the virtual allowable Legendrian isotopy. \square

Acknowledgments. The authors are grateful to Roman Golovko for the many motivating questions and they are very thankful to the anonymous referee for many corrections and suggestions.

This work was partially supported by a grant from the Simons Foundation (#513272 to Vladimir Chernov).

REFERENCES

- [1] V. I. Arnold, *Topological invariants of plane curves and caustics*, (Univ. Lecture Ser., vol. 5) Amer. Math. Soc., Providence, RI (1994)
- [2] S. Antoniou, Stathis, L. Kauffman, S. Lambropoulou: *Black holes and topological surgery*. J. Knot Theory Ramifications **29** (2020), no. 10, 2042010, 6 pp.
- [3] L. Bérard-Bergery, *Quelques exemples de variétés riemanniennes où toutes les géodésiques issues d'un point sont fermées et de même longueur, suivis de quelques résultats sur leur topologie*, Ann. Inst. Fourier (Grenoble) **27** (1977), 231–249.
- [4] A. Bernal, M. Sánchez, *On smooth Cauchy hypersurfaces and Geroch's splitting theorem*, Comm. Math. Phys. **243** (2003), 461–470.

- [5] A. Bernal, M. Sánchez, *Smoothness of time functions and the metric splitting of globally hyperbolic space-times*, Comm. Math. Phys. **257** (2005), 43–50.
- [6] A. Bernal, M. Sánchez, *Globally hyperbolic spacetimes can be defined as “causal” instead of “strongly causal”*, Class. Quant. Grav. **24** (2007), 745–750.
- [7] A. L. Besse, *Manifolds all of whose geodesics are closed*, with appendices by D. B. A. Epstein, J.-P. Bourguignon, L. Bérard-Bergery, M. Berger and J. L. Kazdan. Ergebnisse der Mathematik und ihrer Grenzgebiete, 93. Springer-Verlag, Berlin-New York, 1978.
- [8] A. Borde, H. F. Dowker, R. S. Garcia, R. D. Sorkin, and S. Surya. *Causal continuity in degenerate spacetimes*. Classical Quantum Gravity, 16(11):3457–3481 (1999)
- [9] R. Bott, H. Samelson, *Applications of the theory of Morse to symmetric spaces*, American Journal of Mathematics, **80** (1958) pp. 964–1029
- [10] P. Cahn, A. Levi: *Vassiliev invariants of virtual Legendrian knots*. Pacific J. Math. **273** (2015), no. 1, 21–46
- [11] J. S. Carter, S. Kamada, M. Saito, *Stable Equivalence of Knots on Surfaces and Virtual Knot Cobordisms*, J. Knot Theory Ramifications 11 (2002), pp. 311–322.
- [12] Yu. V. Cherkanov and P. E. Pushkar: *Combinatorics of fronts of Legendrian links and the Arnold 4-conjectures*, Russian Math. Surveys 60:1 (2005) 95–149 and Uspekhi Mat. Nauk 60:1 (2005) 99–154
- [13] V. Chernov: *Causality and Legendrian linking for higher dimensional spacetimes*. J. Geom. Phys. **133** (2018), 26–29
- [14] V. Chernov, S. Nemirovski: *Legendrian links, causality, and the Low conjecture*. Geom. Funct. Anal. **19** (2010), no. 5, 1320–1333.
- [15] V. Chernov, S. Nemirovski: *Non-negative Legendrian isotopy in ST^*M* . Geom. Topol. **14** (2010), no. 1, 611–626
- [16] V. Chernov, S. Nemirovski: *Universal orderability of Legendrian isotopy classes*. J. Symplectic Geom. **14** (2016), no. 1, 149–170
- [17] V. Chernov, S. Nemirovski: *Interval Topology in Contact Geometry*, Commun. Contemp. Math. **22** (2020), 1950042
- [18] V. Chernov, R. Sadykov: *Virtual Legendrian isotopy*. Fund. Math. **234** (2016), no. 2, 127–137
- [19] V. Colin, Vincent, E. Ferrand, P. Pushkar, *Positive isotopies of Legendrian submanifolds and applications*. Int. Math. Res. Not. IMRN 2017, no. 20, 6231–6254.
- [20] U. Frauenfelder, C. Labrousse, F. Schlenk, *Slow volume growth for Reeb flows on spherizations and contact Bott–Samelson theorems*, J. Topol. Anal. **7** (2015), no. 3, 407–451.
- [21] L. García-Heveling, *Topology change with Morse functions: progress on the Borde-Sorkin conjecture*, arXiv:2202.09833 (2022)
- [22] R. P. Geroch, *Domain of dependence*, J. Math. Phys. **11** (1970), 437–449.
- [23] S. W. Hawking and G. F. R. Ellis, *The large scale structure of space-time* Cambridge Monographs on Mathematical Physics, Cambridge University Press (1973)
- [24] L. H. Kauffman, *Virtual knot theory*. European J. Combin. **20** (1999), no. 7, 663–690.
- [25] G. Kuperberg, *What is a virtual link?*. Algebraic and Geometric Topology. **3** (2003) 587–591
- [26] R. J. Low: *Causal relations and spaces of null geodesics*. PhD Thesis, Oxford University (1988)
- [27] R. J. Low: *Stable singularities of wave-fronts in general relativity*, J. Math. Phys. **39**(6), 3332 (1998)
- [28] R. J. Low: *The Space of Null Geodesics*. Proceedings of the Third World Congress of Nonlinear Analysts, Part 5 (Catania, 2000); Nonlinear Anal. **47**(5), 3005 (2001)
- [29] R. J. Low: *The Space of Null Geodesics (and a New Causal Boundary)*. Lecture Notes in Physics **692** (Springer, Berlin, Heidelberg, New York, 2006)
- [30] J. Nataro, P. Tod: *Linking, Legendrian linking and causality*. Proc. London Math. Soc. (**3**) **88** (2004), no. 1, 251–272
- [31] R. Penrose, *The question of cosmic censorship*, Black holes and relativistic stars (Chicago, IL, 1996), pp. 103–122, Univ. Chicago Press, Chicago, IL, 1998.
- [32] M. Polyak, M. O Viro, *Gauss diagram formulas for Vassiliev invariants*. Int. Math. Res. Notices 1994, no. 11, pp. 445–453
- [33] R. D. Sorkin, *Consequences of spacetime topology*. In Proceedings of the Third Canadian Conference on General Relativity and Relativistic Astrophysics, pages 137–163 (1989)

V. CHERNOV, MATHEMATICS DEPARTMENT, 6188 KEMENY, DARTMOUTH COLLEGE, HANOVER
NH 03755, USA

Email address: `Vladimir.Chernov@dartmouth.edu`

R. SADYKOV, MATHEMATICS DEPARTMENT 138 CARDWELL HALL 1228 N. MARTIN LUTHER
KING JR. DR. MANHATTAN, KS 66506-2602, USA

Email address: `Sadykov@math.ksu.edu`