

Prefixes of the Fibonacci word

Jeffrey Shallit

School of Computer Science

University of Waterloo

Waterloo, ON N2L 3G1

Canada

shallit@uwaterloo.ca

February 13, 2023

Abstract

Mignosi, Restivo, and Salemi (1998) proved that for all $\epsilon > 0$ there exists an integer N such that all prefixes of the Fibonacci word of length $\geq N$ contain a suffix of exponent $\alpha^2 - \epsilon$, where $\alpha = (1 + \sqrt{5})/2$ is the golden ratio. In this note we show how to prove an explicit version of this theorem with tools from automata theory and logic. Along the way we gain a better understanding of the repetitive structure of the Fibonacci word.

1 Introduction

The Fibonacci word $\mathbf{f}[0..\infty) = 01001010\dots$ is a well-studied infinite binary word with many interesting properties [2]. It can be defined in many ways; for example, as the limit of the sequence of finite words $(X_i)_{i \geq 1}$, with $X_1 = 1$, $X_2 = 0$, and $X_n = X_{n-1}X_{n-2}$ for $n \geq 3$.

Let $x = x[0..n-1]$ be a nonempty finite word. We say x has *period* p if $x[i] = x[i+p]$ for $0 \leq i < n-p$. The least positive period of a word x is called *the* period, and is denoted $\text{per}(x)$. The *exponent* of a word, denoted $\text{exp}(x)$, is defined to be $|x|/\text{per}(x)$. Thus **entente** has exponent $7/3$. We say a word x is an α -*power*, for $\alpha \geq 1$ a real number, if $|x| = \lceil \alpha \text{per}(x) \rceil$. Thus, for example, **underfund** is a $\sqrt{2}$ -power.

Let $\alpha = (1 + \sqrt{5})/2$ be the golden ratio. In 1997 this author conjectured, and Mignosi, Restivo, and Salemi later proved [5], that an infinite word \mathbf{x} has the property that every sufficiently long prefix has a suffix that is an α^2 -power if and only if \mathbf{x} is ultimately periodic. This is an example of the “local periodicity implies global periodicity” phenomenon.

Furthermore, they proved that the constant α^2 is best possible. Define $e(n)$ to be the largest exponent of a suffix of $\mathbf{f}[0..n-1]$ and for a real number γ define $M_\gamma = \{n : e(n) \geq \gamma\}$. Mignosi et al. proved that for all $\epsilon > 0$, the set $M_{\alpha^2 - \epsilon}$ contains all but finitely many natural numbers.

This latter result was proved as their Proposition 2, using a somewhat complicated induction. In this note we show how to prove an explicit version of the result using `Walnut`, a free theorem-prover for automatic sequences [6, 9]. We do not claim our proof to be simpler than the previous one, but it does give a more explicit version of the theorem, and is more straightforward in some sense. As a bonus, we gain a more detailed understanding of the repetitive properties of the prefixes of \mathbf{f} .

2 Prefixes

Define, as usual, the Fibonacci numbers by the recurrence $F_n = F_{n-1} + F_{n-2}$ with initial values $F_0 = 0$, $F_1 = 1$, and the Lucas numbers $L_n = L_{n-1} + L_{n-2}$ with initial values $L_0 = 2$, $L_1 = 1$. By the Binet formulas we have $F_n = (\alpha^n - \beta^n)/\sqrt{5}$, where $\beta = (1 - \sqrt{5})/2$, and $L_n = \alpha^n + \beta^n$, which also serve to define F_n and L_n for negative indices. Note that $F_{-n} = (-1)^{n+1}F_n$ and $L_{-n} = (-1)^{n+1}L_n$ for all integers n .

By a result of Currie and Saari [3, Cor. 4], we know that all (least) periods of the Fibonacci word are of length F_n for $n \geq 2$. (This can also be proved with `Walnut`; see [7, Thm. 3.15].) Thus, if we are to look for large repetitions, we can restrict our attention to periods of length a Fibonacci number.

We start by determining $G := M_{\alpha^2}$: that is, those “good” n for which $\mathbf{f}[0..n-1]$ has a suffix of exponent $> \alpha^2$. With `Walnut` we can characterize the set

$$G = \{13, 14, 22, 23, 24, 26, 27, 34, 35, 36, 37, 38, 39, 40, 43, \dots\}.$$

To do so, we need to phrase the assertions as a first-order logical formula in a language `Walnut` can understand. We represent numbers in the so-called Fibonacci (or Zeckendorf) numeration systems; see [4] or [10] for more details. Here is the `Walnut` code:

```
reg isfib msd_fib "0*10*":
reg evenfib msd_fib "0*1(00)*":
reg oddfib msd_fib "0*10(00)*":
reg adjfib msd_fib msd_fib "([0,0]*[1,1])|[0,0]*[1,0][0,1][0,0]*":
def ffactoreq "?msd_fib At (t<n) => F[i+t]=F[j+t]":
def suff "?msd_fib n>=x & x>=y & y>=1 & $ffactoreq(n-x,(n+y)-x,x-y)":
reg shift {0,1} {0,1} "([0,0]|[0,1][1,1]*[1,0])*":
def phi2n "?msd_fib (s=0 & n=0) | Ex,y $shift(n-1,x) &
  $shift(x,y) & s=y+2":
def good "?msd_fib Ex,y,z $suff(n,x,y) & $phi2n(y,z) & x>z":
```

The explanation is as follows:

- `isfib` asserts that its argument is a positive Fibonacci number.
- `evenfib` asserts that its argument is a positive even-indexed Fibonacci number.
- `oddfib` asserts that its argument is a positive odd-indexed Fibonacci number.

- `adjfib` asserts that its two arguments are F_{k+1} and F_k for some integer $k \geq 1$.
- `factoreq` asserts that $f[i..i+n-1] = f[j..j+n-1]$.
- `phi2n` is code from [9, p. 278] that asserts that $s = \lfloor \alpha^2 n \rfloor$.
- `suff` asserts that $f[0..n-1]$ has a suffix of length x with period $y \geq 1$.
- `good` asserts that $f[0..n-1]$ has a suffix with exponent $> \alpha^2$.

The last command produces a DFA (deterministic finite automaton) of 12 states accepting the Fibonacci representation of those n belonging to G , in Figure 1.




Figure 1: Fibonacci automaton recognizing the set of good n .

We claim that the only $n \geq 2$ that are not in G are either

- (1) of the form $n = F_i - F_j - 1$ for $i \geq 5$ and $3 \leq j \leq i - 2$; or
- (2) of the form $n = F_i - F_{2j+1}$ for $i \geq 5$ and $1 \leq j \leq (i - 3)/2$.

Let B_1 be the set of n satisfying (1) and B_2 be the set of n satisfying (2). Here are the first few terms of each set:

$$B_1 = \{2, 4, 5, 7, 9, 10, 12, 15, 17, 18, 20, 25, 28, 30, 31, 33, \dots\}$$

$$B_2 = \{3, 6, 8, 11, 16, 19, 21, 29, 32, 42, 50, 53, 55, 76, 84, 87, \dots\}.$$

We can verify the claim above as follows. For (1) we assert that $x = F_i$ and $y = F_j$ for some $i \geq j + 2$ and $j \geq 3$. This is the same as asserting that (x, y) are adjacent Fibonacci numbers with $x > 2y$ and $y \geq 2$. For (2) we let $x = F_i$ and $y = F_{2j+1}$ for some $i \geq 2j + 3$ and $j \geq 1$. This is the same as asserting that (x, y) are adjacent Fibonacci numbers, y is an odd-indexed Fibonacci number, $x > 2y$, and $y \geq 2$.

```
def b1 "?msd_fib $isfib(x) & $isfib(y) & x>2*y & y>=2 & n+1+y=x":
def b2 "?msd_fib $isfib(x) & $oddfib(y) & x>2*y & n>=y & y>=2 & n+y=x":
```

We now verify that for all $n \geq 2$ a number is in G if and only if it is in neither B_1 nor B_2 .

```
eval test "?msd_fib An (n>=2) => ( (~$good(n)) <=>
  (Ex,y $b1(n,x,y)|$b2(n,x,y)) )":
```

And Walnut returns TRUE.

For the n satisfying either condition (1) or (2), we need to locate a suffix of $\mathbf{f}[0..n-1]$ of large exponent. First we show

Lemma 1. *Suppose $n = F_i - F_j - 1$ for $i \geq 5$ and $3 \leq j \leq i - 2$. Then $\mathbf{f}[0..n-1]$*

(i) has period F_{i-2} , or

(ii) has a suffix of length $F_j - 1$ and period F_{j-2} .

Proof. The idea is to use x to represent F_{i-1} , y to represent F_{i-2} , w to represent F_{j-1} and z to represent F_{j-2} . We use the command

```
eval check1 "?msd_fib An,x,y,z,w (n>=2 & $b1(n,x+y,w+z) & $adjfib(x,y) &
  $adjfib(w,z) & z>=1 & x>w & n+w+z+1=x+y)
=> ($suff(n,w+z-1,z) & $suff(n,n,y))":
```

and Walnut returns TRUE. □

Lemma 2. *Suppose $n = F_i - F_{2j+1}$ for $i \geq 5$ and $1 \leq j \leq (i-3)/2$. Then $\mathbf{f}[0..n-1]$ has period F_{i-2} . If further $j \geq 2$ then $\mathbf{f}[0..n-1]$ has a suffix of length F_{2j+1} and period F_{2j-1} .*

Proof. The idea is to use x to represent F_{i-1} , y to represent F_{i-2} , w to represent F_{2j} and z to represent F_{2j-1} . We use the commands

```
eval check2a "?msd_fib An,x,y,z,w (n>=3 & $b2(n,x+y,w+z) & $adjfib(x,y)
  & $adjfib(w,z) & n+w+z=x+y) => $suff(n,n,y)":
eval check2b "?msd_fib An,x,y,z,w (n>=3 & $b2(n,x+y,w+z) & $adjfib(x,y)
  & $adjfib(w,z) & n+w+z=x+y & w>=2) => $suff(n,w+z,z)":
```

and Walnut returns TRUE for both. □

Our main result will now follow after some detailed estimates in the next section.

3 The main result

We will need a classic identity about Fibonacci numbers [1, p. 68], which can be easily proved with the Binet formula, namely,

$$F_a F_{b+2} - F_{a+2} F_b = (-1)^b F_{a-b} \quad (1)$$

for all integers a, b .

Next, we need some estimates.

Lemma 3. For $i \geq 1$ we have

$$\alpha^2 + \frac{(-1)^i}{F_{2i}} < \frac{F_{i+2}}{F_i} < \alpha^2 + \frac{(-1)^i}{F_{2i} - (-1)^i} \quad (2)$$

$$\alpha^2 + \frac{1}{F_{2i-1} + 2} < \frac{F_{2i+1} + 1}{F_{2i-1}} < \alpha^2 + \frac{1}{F_{2i-1}}. \quad (3)$$

Proof. Directly follows from the Binet formula. \square

Finally, we need some technical estimates.

Lemma 4.

(i) $(F_{k+1} + 1)^2 \leq 3F_{2k-1}$ for $k \geq 4$;

(ii) $F_{4k-2}^2 \geq 100F_{2k+1}$ for $k \geq 3$.

(iii) $F_{2k+2} \leq 6F_k^2$ for $k \geq 5$.

(iv) $F_{2k}^2 \geq 8F_{2k+2}$ for $k \geq 4$.

(v) $F_{12k-4}^2 \geq 10000F_{6k+3}$ for $k \geq 2$.

(vi) $F_{2k+1}^2 F_{6k+3} \leq 6F_{6k-2}^2$ for $k \geq 2$.

(vii) $F_{4k+2}^2 \geq 4F_{6k+5}$ for $k \geq 3$.

Proof. We prove only (i) and (iii); the others are simpler and are left to the reader. From the Binet form, we can verify that

$$3F_{2k-1} - (F_{k+1} + 1)^2 = \frac{1}{5}(8F_{2k-3} + 4F_{2k-2} - 10F_{k+1} - 5 - 2(-1)^k) \geq 0$$

for $k \geq 4$. This proves (i).

Similarly, we can verify that

$$6F_k^2 - F_{2k+2} = \frac{1}{5}(3F_{2k-1} - 4F_{2k-2} - 12(-1)^k) \geq 0$$

for $k \geq 5$. This proves (iii). \square

Theorem 5. For all $n \geq 1$ the prefix $\mathbf{f}[0..n-1]$ has a suffix of exponent $\geq \alpha^2 - 3n^{-1/2}$.

Proof. We can check the result for $1 \leq n \leq 21$. In what follows, then, we assume $n > 21$. There are three cases to consider:

Case 1: $n \in G$. In this case, we have already seen above that there is a suffix of $\mathbf{f}[0..n-1]$ of exponent $> \alpha^2$.

Case 2: $n \in B_1$, that is, $n = n_{i,j} = F_i - F_j - 1$ for $i \geq 5$ and $3 \leq j \leq i - 2$. Fix i ; using Lemma 1 we will find a lower bound for $e(n_{i,j})$ applicable to all j , $3 \leq j \leq i - 2$. Note that $n_{i,j} \leq F_i$. Define

$$\begin{aligned} f(i, j) &= (F_j - 1)/F_{j-2} \\ g(i, j) &= (F_i - F_j - 1)/F_{i-2} \\ \rho(i, j) &= (F_i - F_j - 1)F_{j-2} - (F_j - 1)F_{i-2}. \end{aligned}$$

Clearly $g(i, j) \geq f(i, j)$ iff $\rho(i, j) \geq 0$.

We now claim that f is increasing with increasing j . To see this, note that

$$\begin{aligned} f(i, j+1) - f(i, j) &= \frac{F_{j+1} - 1}{F_{j-1}} - \frac{F_j - 1}{F_{j-2}} \\ &= \frac{F_{j+1}F_{j-2} - F_{j-1}F_j + F_{j-1} - F_{j-2}}{F_{j-1}F_{j-2}} \\ &= \frac{(-1)^{j+1} + F_{j-3}}{F_{j-1}F_{j-2}} \geq 0 \end{aligned}$$

for $j \geq 3$. Here we have used Eq. (1) with $a = j - 2$ and $b = j - 1$.

Similarly, g is decreasing with increasing j . To see this, note that

$$g(i, j+1) - g(i, j) = \frac{F_j - F_{j+1}}{F_{i-2}} \leq 0$$

for $i \geq 3$. Hence, for each fixed $i \geq 3$, there is a ‘‘crossover point’’ where the graphs of $f(i, j)$ and $g(i, j)$ cross. This crossover point determines where $\min_j \max(f(i, j), g(i, j))$ occurs. See Figure 2 for an example for $i = 20$.




Figure 2: The crossover point for $f(20, j + 2)$ versus $g(20, j + 2)$.

Observe that $\rho(i, j)$ is increasing with increasing i :

$$\begin{aligned}\rho(i+1, j) - \rho(i, j) &= (F_{i+1} - F_j - 1)F_{j-2} - (F_j - 1)F_{i-1} - (F_i - F_j - 1)F_{j-2} + (F_j - 1)F_{i-2} \\ &= F_{i-1}F_{j-2} - (F_j - 1)F_{i-3} \\ &= F_{i-3} - (-1)^j F_{i-j-1} \geq 0.\end{aligned}$$

Here we used Eq. (1) with $a = i - 3$ and $b = j - 2$.

Using the Binet form one can verify that

$$\begin{aligned}\rho(2k+1, k+1) &= \frac{1}{5}(L_{2k-2} - 5F_{k-1} - 5F_{-k} - 3(-1)^k) \\ \rho(2k+2, k+2) &= \frac{1}{5}(-L_{2k-2} - 5F_k + 5F_{-k} + 3(-1)^k).\end{aligned}$$

It follows that $\rho(2k+2, k+1) \geq \rho(2k+1, k+1) > 0$ for $k \geq 4$ and $\rho(2k+1, k+2) \leq \rho(2k+2, k+2) < 0$ for $k \geq 1$. Set $j' = \lceil i/2 \rceil$. Hence $g(i, j') - f(i, j') \geq 0$ for $i \geq 6$ and $g(i, j'+1) - f(i, j'+1) < 0$ for $i \geq 0$. It follows that if $n \in B_1$, and $n = F_i - F_j - 1$ for some $i \geq 8$, then $e(n) \geq \min(g(i, j'), f(i, j'+1))$.

There are now two subcases to consider: i is odd and i is even.

Suppose i is odd. Then $i = 2k+1$ and $j' = k+1$ for some k . Then using the Binet form one can verify that

$$F_k F_{2k-1} (f(2k+1, k+2) - g(2k+1, k+1)) = \frac{1}{10}(-2(-1)^k + 2L_{2k-3} + 10F_k + 5F_{-k} + 5L_{-k}) > 0$$

for $k \geq 0$. Hence $e(n) \geq g(2k+1, k+1)$.

From Lemma 4 (i) we know that $(F_{k+1} + 1)^2 / F_{2k-1} \leq 3$ for $k \geq 4$. From Eq. (2) we know that $F_{2k+1} / F_{2k-1} \leq \alpha^2$. Multiplying these together and dividing by F_{2k+1} , we get

$$(F_{k+1} + 1)^2 / F_{2k-1}^2 \leq 3\alpha^2 F_{2k+1}^{-1} \leq 3\alpha^2 n^{-1}$$

for $k \geq 4$. Hence

$$(F_{k+1} + 1) / F_{2k-1} \leq \sqrt{3}\alpha n^{-1/2}$$

for $k \geq 3$. Then

$$\begin{aligned}e(n) \geq g(2k+1, k+1) &= \frac{F_{2k+1}}{F_{2k-1}} - \frac{F_{k+1} + 1}{F_{2k-1}} \\ &\geq \left(\alpha^2 - \frac{1}{F_{4k-2}}\right) - \sqrt{3}\alpha n^{-1/2} \\ &\geq \alpha^2 - \frac{1}{10}n^{-1/2} - \sqrt{3}\alpha n^{-1/2} \quad (\text{by Lemma 4 (ii)}) \\ &\geq \alpha^2 - 3n^{-1/2}\end{aligned}$$

for $k \geq 2$.

If i is even then $i = 2k + 2$ and $j' = k + 1$. Then using the Binet form one can verify that

$$F_k F_{2k} (g(2k + 2, k + 1) - f(2k + 2, k + 2)) = \frac{1}{10} (2(-1)^k + 2L_{2k-1} - 10F_k - 5F_{1-k} + 5L_{1-k}) \geq 0$$

for $k \geq 3$. Hence $e(n) \geq f(2k + 2, k + 2)$.

Now from Lemma 4 (iv) we get $-1/F_{2k} \geq -8^{-1/2} F_{2k+2}^{-1/2} \geq -8^{-1/2} n^{-1/2}$. Similarly from Lemma 4 (iii) we get $-1/F_k \geq -\sqrt{6} F_{2k+2}^{-1/2} \geq -\sqrt{6} n^{-1/2}$. Hence

$$\begin{aligned} e(n) \geq f(2k + 2, k + 2) &= \frac{F_{k+2} - 1}{F_k} \\ &\geq \left(\alpha^2 - \frac{1}{F_{2k}} \right) - \frac{1}{F_k} \\ &\geq \alpha^2 - 8^{-1/2} n^{-1/2} - \sqrt{6} n^{-1/2} \\ &\geq \alpha^2 - 3n^{-1/2}. \end{aligned}$$

Case 3: $n \in B_2$, that is, $n = n_{i,j} = F_i - F_{2j+1}$ for $i \geq 5$ and $1 \leq j \leq (i - 3)/2$. (For these n it turns out that $e(n) \geq \alpha^2 - O(n^{-2/3})$, but for our theorem it is not necessary to prove this stronger result.)

Using Lemma 2, we find lower bound for $e(n_{i,j})$ applicable to all j , $1 \leq j \leq (i - 3)/2$. Define

$$\begin{aligned} r(i, j) &= F_{2j+1}/F_{2j-1} \\ s(i, j) &= (F_i - F_{2j+1})/F_{i-2} \\ \psi(i, j) &= (F_i - F_{2j+1})F_{2j-1} - F_{i-2}F_{2j+1}. \end{aligned}$$

Again we need to determine the ‘‘crossover point’’, illustrated in Figure 3.

Clearly r is increasing with increasing j . To see this, note that

$$\begin{aligned} r(i, j + 1) - r(i, j) &= \frac{F_{2j+3}}{F_{2j+1}} - \frac{F_{2j+1}}{F_{2j-1}} \\ &= \frac{F_{2j+3}F_{2j-1} - F_{2j+1}^2}{F_{2j+1}F_{2j-1}} \\ &= \frac{1}{F_{2j+1}F_{2j-1}} > 0. \end{aligned}$$

Here we used Eq. (1) with $a = 2j - 1$ and $b = 2j + 1$.

Next, we show that s is decreasing with increasing j :

$$\begin{aligned} s(i, j + 1) - s(i, j) &= \frac{F_i - F_{2j+3}}{F_{i-2}} - \frac{F_i - F_{2j+1}}{F_{i-2}} \\ &= \frac{F_{2j+1} - F_{2j+3}}{F_{i-2}} < 0, \end{aligned}$$




Figure 3: The crossover point for $r(30, j)$ versus $s(30, j)$.

where we used Eq. (1) with $a = 2j - 1$ and $b = i - 3$.

Also $\psi(i, j)$ is increasing with increasing i , provided $i \geq 2j + 2$. We have

$$\begin{aligned}
 \psi(i + 1, j) - \psi(i, j) &= (F_{i+1} - F_{2j+1})F_{2j-1} - F_{i-1}F_{2j+1} - (F_i - F_{2j+1})F_{2j-1} - F_{i-2}F_{2j+1} \\
 &= F_{i-1}F_{2j-1} - F_{i-3}F_{2j+1} \\
 &= F_{i-2j-2} \geq 0.
 \end{aligned} \tag{4}$$

Using the Binet form we can easily check that

$$\begin{aligned}
 \psi(6k, k) &= (L_{4k-2} - 3)/5 \\
 \psi(6k + 5, k + 1) &= (-L_{4k} - 3)/5.
 \end{aligned}$$

This, combined with (4) above, gives

$$\psi(6k + 5, k) \geq \psi(6k + 4, k) \geq \psi(6k + 3, k) \geq \psi(6k + 2, k) \geq \psi(6k + 1, k) \geq \psi(6k, k) \geq 0$$

and

$$\begin{aligned}
 \psi(6k, k + 1) &\leq \psi(6k + 1, k + 1) \leq \psi(6k + 2, k + 1) \leq \psi(6k + 3, k + 1) \leq \psi(6k + 4, k + 1) \\
 &\leq \psi(6k + 5, k + 1) \leq 0.
 \end{aligned}$$

Now choose $j' = \lfloor i/6 \rfloor$. Then $r(i, j') \leq s(i, j')$ and $r(i, j' + 1) \geq s(i, j' + 1)$ for $i \geq 1$. So for $i \geq 1$ we have $e(n) \geq \min(s(i, j'), r(i, j' + 1))$.

There are now two cases to consider:

(a) $i = 6k + a$, $a \in \{0, 1, 2, 3\}$, and $j = k$, and

(b) $i = 6k + a$, $a \in \{4, 5\}$, and $j = k$.

In case (a) we have, for $i = 6k$, that

$$\begin{aligned} F_{2k+1}F_{6k-2}(r(6k, k+1) - s(6k, k)) &= F_{6k-2}F_{2k+3} - F_{2k+1}F_{6k} + F_{2k+1}^2 \\ &= \frac{1}{20}(7L_{4k-3} + 15F_{4k} + 8) \geq 0. \end{aligned}$$

For $i = 6k + 1$:

$$\begin{aligned} F_{2k+1}F_{6k-1}(r(6k+1, k+1) - s(6k+1, k)) &= F_{6k-1}F_{2k+3} - F_{2k+1}F_{6k+1} + F_{2k+1}^2 \\ &= \frac{1}{5}(-2L_{4k-3} + 5F_{4k} + 2) \geq 0. \end{aligned}$$

For $i = 6k + 2$:

$$\begin{aligned} F_{2k+1}F_{6k}(r(6k+2, k+1) - s(6k+2, k)) &= F_{6k}F_{2k+3} - F_{2k+1}F_{6k+2} + F_{2k+1}^2 \\ &= \frac{1}{10}(L_{4k-3} + 5F_{4k} + 4) \geq 0. \end{aligned}$$

For $i = 6k + 3$:

$$\begin{aligned} F_{2k+1}F_{6k+1}(r(6k+3, k+1) - s(6k+3, k)) &= F_{6k+1}F_{2k+3} - F_{2k+1}F_{6k+3} + F_{2k+1}^2 \\ &= \frac{1}{20}(-3L_{4k-3} + 5F_{4k} + 8) \geq 0. \end{aligned}$$

Hence

$$\begin{aligned} e(n) \geq s(6k+a, k) &= \frac{F_{6k+a} - F_{2k+1}}{F_{6k+a-2}} \\ &\geq \left(\alpha^2 - \frac{1}{F_{2k+2a-4}}\right) - \frac{F_{2k+1}}{F_{6k+a-2}} \\ &\geq \alpha^2 - (1/100)F_{6k+3}^{-1/2} - \sqrt{6}F_{6k+a}^{-1/2} \quad (\text{by Lemma 4 (vi)}) \\ &\geq \alpha^2 - (1/100 + \sqrt{6})n^{-1/2} \\ &\geq \alpha^2 - 3n^{-1/2}. \end{aligned}$$

In case (b) we have $a \in \{4, 5\}$. Then

$$\begin{aligned} e(n) \geq r(6k+a, k+1) &= \frac{F_{2k+3}}{F_{2k+1}} \\ &\geq \alpha^2 - \frac{1}{F_{4k+2}} \\ &\geq \alpha^2 - \frac{1}{2}F_{6k+5}^{-1/2} \\ &\geq \alpha^2 - \frac{1}{2}F_{6k+a}^{-1/2} \\ &\geq \alpha^2 - 2n^{-1/2} \quad (\text{by Lemma 4 (vii)}) \end{aligned}$$

for $k \geq 3$.

Thus in all cases $e(n) \geq \alpha^2 - 3n^{-1/2}$. This completes the proof. \square

4 Final remarks

For real numbers γ , the set M_γ can be quite complicated. For example, Rampersad [8] recently studied the set M_3 . In Walnut one can create a Fibonacci automaton accepting the set $M_{p/q}$, for natural numbers p, q , with a command of the form

```
def emmpq "?msd_fib Ex,y $suff(n,x,y) & p*y<=q*x":
```

where p is replaced by the specific number p and q is replaced by q .

Given a rational number $p/q < \alpha^2$, Walnut can also determine the largest n for which $e(n) < p/q$, as follows:

```
def has_suff "?msd_fib Ex,y $suff(n,x,y) & p*y<q*x":
def largest_index "?msd_fib (~$has_suff(n)) & Am (m>n) => $has_suff(m)":
test largest_index 1:
```

This code prints the Fibonacci representation of n . Again, one needs to replace p with the specific number p and likewise with q .

With more work, it should be possible to prove that, for $k \geq 6$, the largest n for which $e(n) < (F_{k+1} - 1)/F_{k-1}$ is $n = F_{2k-1} - F_k - 1$.

References

- [1] S. L. Basin and V. E. Hoggatt, Jr. A primer on the Fibonacci sequence—Part II. *Fib. Quart.* **1** (1963), 61–68.
- [2] J. Berstel. Fibonacci words—a survey. In G. Rozenberg and A. Salomaa, editors, *The Book of L*, pp. 13–27. Springer-Verlag, 1986.
- [3] J. D. Currie and K. Saari. Least periods of factors of infinite words. *RAIRO Info. Théor. Appl.* **43** (2008), 165–178.
- [4] C. G. Lekkerkerker. Voorstelling van natuurlijke getallen door een som van getallen van Fibonacci. *Simon Stevin* **29** (1952), 190–195.
- [5] F. Mignosi, A. Restivo, and S. Salemi. Periodicity and the golden ratio. *Theoret. Comput. Sci.* **204** (1998), 153–167.
- [6] H. Mousavi. Automatic theorem proving in Walnut. Arxiv preprint arXiv:1603.06017 [cs.FL], available at <http://arxiv.org/abs/1603.06017>, 2016.

- [7] H. Mousavi, L. Schaeffer, and J. Shallit. Decision algorithms for Fibonacci-automatic words, I: Basic results. *RAIRO Inform. Théor. App.* **50** (2016), 39–66.
- [8] N. Rampersad. Prefixes of the Fibonacci word that end with a cube. *C. R. Math. Acad. Sci. Paris* **361** (2023) 323–330.
- [9] J. Shallit. *The Logical Approach To Automatic Sequences: Exploring Combinatorics on Words with Walnut*, Vol. 482 of *London Math. Society Lecture Note Series*. Cambridge University Press, 2022.
- [10] E. Zeckendorf. Représentation des nombres naturels par une somme de nombres de Fibonacci ou de nombres de Lucas. *Bull. Soc. Roy. Liège* **41** (1972), 179–182.