

# Orthogonal cycle systems with cycle length less than 10

**Selda Küçükçifçi**

email: skucukcifci@ku.edu.tr

**Emine Şule Yazıcı**

email: eyazici@ku.edu.tr

Department of Mathematics, Koç University, Istanbul, Turkey

## Abstract

An  $H$ -decomposition of  $G$  is a partition of the edge-set of  $G$  into subsets, where each subset induces a copy of the graph  $H$ . A  $k$ -orthogonal  $H$ -decomposition of a graph  $G$  is a set of  $k$   $H$ -decompositions of  $G$ , such that any two copies of  $H$  in distinct  $H$ -decompositions intersect in at most one edge. When  $G = K_v$  we call the  $H$ -decomposition an  $H$ -system of order  $v$ . In this paper we consider the case  $H$  is an  $l$ -cycle and construct a pair of orthogonal  $l$ -cycle systems for all admissible orders when  $l = 5, 6, 7, 8$  or  $9$ , except  $(l, v) = (7, 7)$  and  $(l, v) = (9, 9)$ .

AMS classification: 05C38, 05B15, 05B30.

Keywords: Orthogonal cycle decompositions; Heffter arrays

## 1 Introduction

Orthogonal combinatorial structures especially; orthogonal Latin squares has a history dating back to the famous Euler's 36 officer problem. Please see for example the vast literature on orthogonal Latin squares in [13] and the survey of orthogonal factorizations in [2]. We will consider orthogonal cycle decompositions in this paper.

An  $H$ -decomposition of  $G$  is a partition of the edge-set of  $G$  into subsets, where each subset induces a copy of the graph  $H$ . A  $k$ -orthogonal  $H$ -decomposition of a graph  $G$  is a set of  $k$

$H$ -decompositions of  $G$ , such that any two copies of  $H$  in distinct  $H$ -decompositions intersect in at most one edge. When  $G = K_v$  we call the  $H$ -decomposition an  $H$ -system of order  $v$ . Copies of  $H$  are called blocks.

In [9], it was shown asymptotically that for sufficiently large  $v$ ,  $k$ -orthogonal  $K_r$ -decompositions of  $K_v$  exists whenever a  $K_r$ -decomposition of  $K_n$  exists. This result was generalized in [10] to the general graph  $H$  which we will refer as an  $H$ -system of order  $v$ . These structures are closely related to super-simple and completely-reducible super-simple designs. See [13] for definitions and related results.

It was established in [1, 6, 26] that when  $H$  is an  $l$ -cycle then a decomposition of order  $v$  exists if and only if  $v$  is odd,  $3 \leq l \leq v$  and  $v(v-1) \equiv 0 \pmod{2l}$ . Such a decomposition is called an  $l$ -cycle system of order  $v$ . When  $H$  is a 3-cycle, two 3-cycle systems are orthogonal if and only if the cycle systems are disjoint. Therefore, there are at most  $v-2$  mutually orthogonal 3-cycle systems of order  $v$ . A set of systems which meets this bound forms a large set of disjoint Steiner triple systems and they exist for all admissible  $v \neq 7$  [21, 22, 27].

**Example 1.1** *A pair of orthogonal Steiner triple systems of order 7 exists.*

Let  $V = \mathbb{Z}_7$ ,  $C_1$  and  $C_2$ , each, be a collection of triples obtained by developing the base triples  $(1, 2, 4)$  and  $(1, 3, 4)$ , respectively modulo 7. Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal Steiner triple systems of order 7.  $\square$

In [7], authors focus on the case  $v \equiv 1 \pmod{2l}$  when  $l$  is even and proved the following result.

**Theorem 1.2** [7] *Suppose  $\mu(v, l)$  represent the maximum possible number of mutually orthogonal  $l$  cycle systems of order  $v$ . Let  $l \geq 4$  is even,  $v \equiv 1 \pmod{2l}$  and  $N = \frac{v-1}{2l}$ . Then  $\mu(v, l) \geq \frac{N}{al+1} - 1$ , where  $(a, b) = (4, -2)$  if  $l \equiv 0 \pmod{4}$  and;  $(a, b) = (24, -18)$  if  $l \equiv 2 \pmod{4}$ .*

Also in the same paper it was noted that  $\mu(v, l) \leq v-2$  in general and  $\mu(v, l) \leq 1$  when  $2l^2 > v(v-1)$ .

On the other hand, orthogonal cycle systems naturally arise from face 2-colourable embeddings of graphs on surfaces, where each pair of faces share at most one edge and each edge

belongs to exactly two faces. So first results on determining the spectrum of 2-orthogonal cycle systems make use of Heffter arrays that are constructed for the purpose of obtaining these type of 2-colourable embeddings of the complete graph on high genus surfaces. A *Heffter array*  $H(m, n; s, l)$  is an  $m \times n$  array of integers, such that:

- each row contains  $s$  filled cells and each column contains  $l$  filled cells;
- the elements in every row and column sum to 0 (mod  $2ml + 1$ ); and
- for each integer  $1 \leq x \leq ms$ , either  $x$  or  $-x$  appears in the array.

A Heffter array is *square*, if  $m = n$  and necessarily  $s = l$ , and is denoted by  $H(m; l)$ . A square Heffter array  $H(m; l)$  is said to be a *simple array* if, for each row and column, the entries may be cyclically ordered so that all partial sums,

$$\alpha_i = \sum_{j=0}^i a_j \pmod{(2ml + 1)}$$

where  $a_0, \dots, a_{l-1}$  are the elements in that row or column, are distinct modulo  $2ml + 1$  for  $i = 0, \dots, l - 1$ . One may easily check that any Heffter array with  $l \leq 5$  is simple. The results in [3] implies the following theorem as part of a more general result; but it was stated explicitly in [16] including the case for the cocktail party graphs.

**Theorem 1.3** *If  $H(m; l)$  is a Heffter array with a simple ordering, then there exists a pair of orthogonal cyclic  $l$ -cycle systems of order  $2ml + 1$ .*

The  $H(m; 6)$ 's constructed in [11] are simple. Thus the following is implied by the existing literature on Heffter arrays.

**Theorem 1.4** [5, 8, 11, 16, 20] *Let  $m \geq l \geq 3$  be positive integers. Then there exists a pair of orthogonal  $l$ -cycle systems of order  $2ml + 1$  when*

- $3 \leq l \leq 6$ ;
- $l \in \{7, 9\}$  and  $ml \equiv 3 \pmod{4}$ ;
- $l \equiv 0 \pmod{4}$ ;

- $m \equiv 1 \pmod{4}$  and  $l \equiv 3 \pmod{4}$ ;
- $m \equiv 0 \pmod{4}$  and  $l \equiv 3 \pmod{4}$  (for large enough  $m$ ).

If they satisfy an extra condition on the orderings of the entries of the Heffter array, these orthogonal cycle systems in turn biembed to yield a face 2-colourable embedding on an orientable surface. (See for example [12, 19]). For other existence results and generalizations of Heffter arrays, see [4, 11, 14, 15, 17, 18, 23, 24, 25].

All the orthogonal  $l$ -cycle systems obtained using Heffter arrays given in Theorem 1.4 and the cyclic systems constructed in [7] are of order  $v = 2ml + 1$  for some integer  $m$ , hence only deals with the case  $v \equiv 1 \pmod{2l}$  for  $m \geq l$ . In this paper we will show that a pair of orthogonal  $l$ -cycle systems of order  $n$  exists whenever an  $l$ -cycle system of order  $n$  exists for the cycle lengths  $3 \leq l \leq 9$ .

When  $l = 3$ , as stated earlier a large set of Steiner triple systems exists for all  $v \neq 7$  and the unique  $STS(7)$  has an orthogonal pair hence a pair of orthogonal 3-cycle systems exist for all  $v \equiv 1, 3 \pmod{6}$ . When  $l = 4$ , a 4-cycle system exists if and only if  $v \equiv 1 \pmod{8}$ . A pair of orthogonal 4-cycle systems is obtained in [7] for each of these admissible orders. We will construct a pair of orthogonal  $l$ -cycle systems for all admissible orders when  $l = 5, 6, 7, 8$  or  $9$ , except  $(l, v) = (7, 7)$  and  $(l, v) = (9, 9)$ .

The followings are the definitions of the necessary structures for our constructions.

A  $k$ -GDD is a triple  $(V, \mathcal{G}, \mathcal{B})$ , where  $V$  is a finite set of vertices,  $\mathcal{G} = \{G_1, G_2, \dots, G_n\}$  is a partition of  $V$  into subsets, called *groups* and  $\mathcal{B}$  is a collection of isomorphic copies of  $K_k$ , which partitions the edges of  $K_{g_1, g_2, \dots, g_n}$  on  $V$ , where  $|G_i| = g_i$ . If for  $i = 1, 2, \dots, t$ , there are  $u_i$  groups of size  $g_i$ , we say that the  $k$ -GDD is of type  $g_1^{u_1} g_2^{u_2} \dots g_t^{u_t}$ .

We will denote all cycles with length  $l$  and edges  $\{a_1, a_2\}, \{a_2, a_3\}, \dots, \{a_{l-1}, a_l\}, \{a_l, a_1\}$  by  $(a_1, a_2, \dots, a_l)$ .

An automorphism of an  $STS(v)$   $(V, T)$  is a bijection  $\alpha : V \rightarrow V$  such that  $t = \{x, y, z\} \in T$  if and only if  $t\alpha = \{x\alpha, y\alpha, z\alpha\} \in T$ . An  $STS(v)$  is cyclic if it has an automorphism that is a permutation consisting of a single cycle of length  $v$ .

## 2 Orthogonal 5-cycle systems

The spectrum of 5-cycle systems; that is the set of all  $v$  such that a 5-cycle system of order  $v$  exists, is the set of all  $v \equiv 1$  or  $5 \pmod{10}$ . There exists a pair of orthogonal cyclic 5-cycle systems of order  $10k + 1$  for all  $k \geq 5$  [16]. But we will give a construction that works for both of the orders  $10k + 1$  and  $10k + 5$  for all  $k \geq 3$ .

Since a pair of orthogonal 5-cycle systems of order 11 is given in [7], we will start by giving the remaining pairs of orthogonal 5-cycle decompositions we will need in our constructions. All systems of small orders that are not used in our recursive constructions throughout the paper will be given in the Appendix.

**Example 2.1** *A pair of orthogonal 5-cycle systems of order 15 exists.*

Let  $V = \mathbb{Z}_{15}$  and

$$C_1 = \{(0, 1, 2, 3, 4), (0, 2, 12, 8, 5), (0, 3, 1, 4, 6), (0, 7, 1, 5, 9), (0, 8, 1, 6, 10), (0, 11, 1, 9, 12), (0, 13, 1, 10, 14), (1, 12, 3, 5, 14), (2, 4, 5, 6, 7), (2, 5, 7, 3, 6), (2, 8, 3, 9, 10), (2, 9, 4, 7, 11), (2, 13, 3, 11, 14), (3, 10, 4, 8, 14), (4, 11, 5, 10, 12), (4, 13, 5, 12, 14), (6, 8, 7, 9, 11), (6, 9, 8, 13, 14), (6, 12, 11, 10, 13), (7, 10, 8, 11, 13), (7, 12, 13, 9, 14)\},$$

$$C_2 = \{(0, 1, 3, 5, 2), (0, 3, 2, 4, 7), (0, 4, 1, 5, 8), (0, 5, 4, 3, 6), (0, 9, 1, 2, 10), (0, 11, 2, 6, 13), (0, 12, 1, 6, 14), (1, 7, 2, 8, 10), (1, 8, 3, 7, 11), (1, 13, 2, 9, 14), (2, 12, 3, 10, 14), (3, 9, 4, 6, 11), (3, 13, 5, 7, 14), (4, 8, 6, 5, 12), (4, 10, 5, 9, 13), (4, 11, 9, 8, 14), (5, 11, 8, 12, 14), (6, 7, 8, 13, 10), (6, 9, 7, 10, 12), (7, 12, 11, 14, 13), (9, 10, 11, 13, 12)\}.$$

Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 5-cycle systems of order 15.  $\square$

**Example 2.2** *A pair of orthogonal 5-cycle decompositions of  $K_{15} \setminus K_5$  exists.*

Let  $V = \mathbb{Z}_{15}$  with the hole of size 5 on  $\{0, 1, 2, 3, 4\}$  and

$$C_1 = \{(0, 5, 11, 6, 10), (0, 6, 5, 2, 14), (0, 7, 1, 5, 8), (0, 9, 1, 6, 12), (0, 11, 1, 8, 13), (1, 10, 2, 6, 13), (1, 12, 2, 7, 14), (2, 8, 3, 5, 9), (2, 11, 3, 7, 13), (3, 6, 4, 5, 10), (3, 9, 4, 7, 12), (3, 13, 4, 8, 14), (4, 10, 7, 5, 12), (4, 11, 7, 6, 14), (5, 13, 9, 12, 14), (6, 8, 10, 11, 9), (7, 8, 12, 10, 9), (8, 9, 14, 13, 11), (10, 13, 12, 11, 14)\},$$

$C_2 = \{(0, 5, 1, 6, 13), (0, 6, 2, 7, 8), (0, 7, 3, 5, 10), (0, 9, 2, 5, 11), (0, 12, 1, 8, 14), (1, 7, 4, 5, 9), (1, 10, 3, 8, 11), (1, 13, 3, 6, 14), (2, 8, 4, 6, 10), (2, 11, 4, 9, 12), (2, 13, 4, 10, 14), (3, 9, 6, 5, 14), (3, 11, 6, 8, 12), (4, 12, 7, 9, 14), (5, 7, 6, 12, 13), (5, 8, 9, 11, 12), (7, 10, 8, 13, 11), (7, 13, 10, 12, 14), (9, 10, 11, 14, 13)\}$ . Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 5-cycle decompositions of  $K_{15} \setminus K_5$ .  $\square$

**Lemma 2.3** *A pair of orthogonal 5-cycle systems of order  $v$  exists for all  $v \equiv 1$  or  $5 \pmod{10}$ .*

**Proof** The cases  $v = 11$  is given in [7],  $v = 15$  in Example 2.1 and  $v = 21, 25$  in the Appendix. The following construction works for all  $v = 10k + r$  and  $k \geq 3$ , when  $r = 1$  or  $5$ . Let  $(Q, \circ)$  be a commutative quasigroup of order  $2k$ , for  $k \geq 3$ , with holes  $H = \{h_1, h_2, \dots, h_k\}$ , where  $h_i = \{2i - 1, 2i\}$  for  $i = 1, 2, \dots, k$ . Let  $V = \bigcup_{j=1}^r \{\infty_j\} \cup (Q \times \{1, 2, 3, 4, 5\})$ , where  $r = 1$  or  $5$ . Let  $C_1$  and  $C_2$  be the following collections of 5-cycles on the set  $V$ :

(1) Consider a pair of orthogonal 5-cycle systems of order  $10 + r$ . On  $\bigcup_{j=1}^r \{\infty_j\} \cup (\{h_i\} \times \{1, 2, 3, 4, 5\})$ , when  $r = 1$  for  $i = 1, 2, \dots, k$  and when  $r = 5$  for  $i = 1$ , place a copy of the 5-cycles of the 5-cycle system of order  $10 + r$  in  $C_1$  and the 5-cycles of its orthogonal system in  $C_2$ .

(2) For the case  $r = 5$ , for each of the remaining holes  $h_i, 2 \leq i \leq k$ , consider a pair of orthogonal 5-cycle decompositions of  $K_{15} \setminus K_5$ . On  $\bigcup_{j=1}^5 \{\infty_j\} \cup (h_i \times \{1, 2, 3, 4, 5\})$ , place a copy of the 5-cycles in  $C_1$  and the 5-cycles of its orthogonal decomposition in  $C_2$ , being sure that the vertex set of  $K_5$  is  $\bigcup_{j=1}^5 \{\infty_j\}$ .

(3) If  $x$  and  $y$  belong to different holes of  $H$ , place the 5-cycles  $((x, i), (y, i), (x, i + 1), (x \circ y, i + 3), (y, i + 1)) \in C_1$  and  $((x, i), (y, i), (x, i + 2), (x \circ y, i + 3), (y, i + 2)) \in C_2$  for  $i = 1, 2, 3, 4, 5$  and addition is done in modulo 5.

Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 5-cycle systems of order  $v$ .  $\square$

### 3 Orthogonal 6-cycle systems

The spectrum of 6-cycle systems is the set of all  $v \equiv 1$  or  $9 \pmod{12}$ . For  $v \geq 73$  there exists a pair of orthogonal cyclic 6-cycle systems of order  $12k + 1$  [11, 16]. The constructions we will give, besides handling the case  $v \equiv 9 \pmod{12}$ , also take care of the missing small systems of the orders  $v \equiv 1 \pmod{12}$ .

**Example 3.1** *A pair of orthogonal 6-cycle systems of order 9 exists.*

Let  $V = \mathbb{Z}_9$  and

$$C_1 = \{(0, 1, 2, 3, 4, 5), (0, 2, 4, 1, 3, 6), (0, 3, 5, 1, 7, 8), (0, 4, 6, 8, 3, 7), (1, 6, 7, 2, 5, 8), (2, 6, 5, 7, 4, 8)\},$$

$$C_2 = \{(0, 1, 3, 5, 6, 7), (0, 2, 1, 5, 8, 4), (0, 3, 2, 7, 4, 6), (0, 5, 2, 6, 3, 8), (1, 4, 5, 7, 8, 6), (1, 7, 3, 4, 2, 8)\}.$$

Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 6-cycle systems of order 9. □

**Example 3.2** *A pair of orthogonal 6-cycle decompositions of  $K_{4,4,4}$  exists.*

Let  $V = \{0, 1, 2, \dots, 11\}$ , where  $\{0, 1, 2, 3\}$ ,  $\{4, 5, 6, 7\}$ ,  $\{8, 9, 10, 11\}$  be the parts of  $K_{4,4,4}$  and

$$C_1 = \{(0, 4, 1, 5, 2, 6), (0, 5, 3, 4, 2, 7), (0, 8, 1, 6, 3, 9), (0, 10, 1, 7, 3, 11), (1, 9, 5, 10, 4, 11), (2, 8, 7, 11, 6, 9), (2, 10, 6, 8, 5, 11), (3, 8, 4, 9, 7, 10)\},$$

$$C_2 = \{(0, 4, 2, 8, 1, 11), (0, 5, 1, 9, 2, 10), (0, 6, 8, 3, 5, 9), (0, 7, 3, 10, 5, 8), (1, 4, 3, 6, 9, 7), (1, 6, 2, 11, 7, 10), (2, 5, 11, 4, 8, 7), (3, 9, 4, 10, 6, 11)\}.$$

Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 6-cycle decompositions of  $K_{4,4,4}$ . □

**Example 3.3** *A pair of orthogonal 6-cycle systems of order 21 exists.*

Let  $V = \mathbb{Z}_7 \times \{1, 2, 3\}$ . Let  $B_1$  and  $B_2$  contain the 5 base 6-cycles that will be developed (modulo 7,  $-$ ) to produce the 35 6-cycles in  $C_1$  and  $C_2$ , respectively, such that  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 6-cycle systems of order 21.

$B_1 = \{((0, 2), (0, 1), (1, 1), (0, 3), (3, 1), (6, 1)), ((0, 3), (5, 2), (2, 2), (4, 1), (3, 2), (1, 2)), ((0, 1), (0, 3), (4, 3), (0, 2), (1, 3), (2, 3)), ((0, 1), (5, 1), (1, 2), (2, 2), (0, 3), (5, 3)), ((0, 1), (1, 3), (5, 2), (5, 3), (2, 1), (4, 2))\}.$

$B_2 = \{((0, 2), (1, 1), (3, 1), (0, 1), (6, 1), (0, 3)), ((0, 3), (2, 2), (3, 2), (5, 2), (1, 2), (4, 1)), ((0, 1), (4, 3), (1, 3), (0, 3), (2, 3), (0, 2)), ((0, 1), (1, 2), (0, 3), (4, 2), (2, 1), (2, 3)), ((1, 1), (4, 2), (1, 3), (0, 2), (2, 1), (0, 3))\}.$  □

**Lemma 3.4** *A pair of orthogonal 6-cycle systems of order  $v$  exists for all  $v \equiv 1, 9, 13$  or  $21$  (mod 24).*

**Proof** A pair of orthogonal 6-cycle systems of order 13 is given in [7]. The cases  $v = 9, 21$  are given in Examples 3.1, 3.3 and  $v = 45$  in the Appendix. The following construction works for all  $v = 24k + r$  when  $r = 1, 9$  or  $13$  for all  $k \geq 1$  and when  $r = 21$   $k \geq 2$ . Let  $V = \{\infty\} \cup (X \times \{1, 2, 3, 4\})$ , where  $|X| = \frac{v-1}{4}$ . Let  $(X, G, B)$  be a 3-GDD of type  $2^{(v-1)/8}$  for  $r = 1, 9$ , a 3-GDD of type  $3^{(v-1)/12}$  for  $r = 13$  and a 3-GDD of type  $5^1 3^{2k}$  for  $r = 21$  [13]. Let  $C_1$  and  $C_2$  be the following collection of 6-cycles on the set  $V$ :

(1) Consider a pair of orthogonal 6-cycle systems of order 9, 13 or 21 as needed. For each group  $g \in G$ , on  $V = \{\infty\} \cup (g \times \{1, 2, 3, 4\})$  place a copy of the 6-cycles of the needed system in  $C_1$  and the 6-cycles of its orthogonal system in  $C_2$ .

(2) For each triple  $\{x, y, z\} \in B$ , consider a pair of orthogonal 6-cycle decompositions of  $K_{4,4,4}$  (whose parts are  $x \times \{1, 2, 3, 4\}$ ,  $y \times \{1, 2, 3, 4\}$ ,  $z \times \{1, 2, 3, 4\}$ ) on the set  $\{x, y, z\} \times \{1, 2, 3, 4\}$ , and place a copy of the 6-cycles of the first system in  $C_1$  and the 6-cycles of its orthogonal system in  $C_2$ .

Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 6-cycle systems of order  $v$ . □

## 4 Orthogonal 7-cycle systems

The spectrum of 7-cycle systems is the set of all  $v \equiv 1$  or  $7 \pmod{14}$ . There exists a pair of orthogonal cyclic 7-cycle systems of order  $56k + 15$  for all  $k \geq 2$  [16]. Below we will give a construction that works for the orders  $14k + 1$  and  $14k + 7$  for all  $k \geq 3$ .



**Example 4.1** *A pair of orthogonal 7-cycle systems of order 21 exists.*

Let  $V = \mathbb{Z}_{20} \cup \{\infty\}$  and  $C_1$  contain the 7-cycles obtained by the two base blocks  $(0, 3, 19, 4, 16, 2, 13)$  and  $(\infty, 0, 1, 3, 13, 11, 10)$  modulo 20, where the second one is a short base block.

Let  $C_2 = \{(9, 14, 11, 13, 15, 18, 17), (10, 13, 14, 15, 12, 17, 19), (11, 15, 16, 19, 13, 18, \infty), (0, 8, 4, 11, 1, 10, 12), (0, 11, 2, 10, 3, 7, 15), (0, 14, 1, 9, 2, 12, 16), (0, 17, 1, 12, 3, 8, 18), (0, 19, 1, 13, 2, 14, \infty), (1, 15, 2, 16, 3, 9, 18), (1, 16, 4, 9, 5, 12, \infty), (2, 17, 3, 11, 5, 10, 18), (2, 19, 3, 13, 4, 15, \infty), (3, 14, 4, 12, 6, 7, 18), (3, 15, 5, 13, 6, 8, \infty), (4, 17, 5, 14, 6, 9, 19), (4, 18, 5, 16, 6, 10, \infty), (5, 19, 6, 11, 7, 9, \infty), (6, 15, 8, 7, 10, 11, 18), (6, 17, 7, 12, 8, 19, \infty), (7, 14, 8, 10, 9, 11, 19), (7, 16, 8, 11, 12, 13, \infty), (8, 13, 9, 15, 10, 16, 17), (9, 12, 14, 10, 17, 11, 16), (0, 1, 2, 3, 4, 5, 6), (0, 2, 4, 1, 3, 5, 7), (0, 3, 6, 1, 7, 4, 10), (0, 4, 6, 2, 5, 8, 9), (0, 5, 1, 8, 2, 7, 13), (12, 18, 16, \infty, 17, 14, 19), (13, 16, 14, 18, 19, 15, 17)\}$ . Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 7-cycle systems of order 21.  $\square$

**Example 4.2** *A pair of orthogonal 7-cycle decompositions of  $K_{21} \setminus K_7$  exists.*

Let  $V = \mathbb{Z}_{21}$ , where the vertex set of the hole  $K_7$  is  $V = \{0, 1, 2, 3, 4, 5, 6\}$  and

$C_1 = \{(0, 7, 1, 10, 6, 11, 8), (1, 8, 2, 11, 0, 12, 9), (2, 9, 3, 12, 1, 13, 10), (3, 10, 4, 13, 2, 7, 11), (4, 11, 5, 7, 3, 8, 12), (5, 12, 6, 8, 4, 9, 13), (6, 13, 0, 9, 5, 10, 7), (0, 14, 1, 17, 6, 18, 15), (1, 15, 2, 18, 0, 19, 16), (2, 16, 3, 19, 1, 20, 17), (3, 17, 4, 20, 2, 14, 18), (4, 18, 5, 14, 3, 15, 19), (5, 19, 6, 15, 4, 16, 20), (6, 20, 0, 16, 5, 17, 14), (0, 10, 8, 5, 15, 9, 17), (1, 11, 9, 6, 16, 7, 18), (2, 12, 7, 4, 14, 8, 19), (3, 13, 7, 8, 9, 10, 20), (7, 9, 14, 10, 11, 12, 15), (7, 14, 11, 13, 8, 15, 17), (7, 19, 9, 16, 8, 18, 20), (8, 17, 10, 12, 13, 14, 20), (9, 18, 10, 15, 11, 19, 20), (10, 16, 11, 17, 12, 14, 19), (11, 18, 12, 16, 17, 13, 20), (12, 19, 18, 13, 16, 15, 20), (13, 15, 14, 16, 18, 17, 19)\}$ ,

$C_2 = \{(0, 7, 2, 8, 3, 9, 10), (0, 8, 1, 11, 3, 7, 9), (0, 11, 4, 7, 1, 12, 13), (0, 12, 2, 9, 4, 10, 14), (0, 15, 1, 9, 5, 7, 16), (0, 17, 1, 10, 2, 11, 18), (0, 19, 1, 13, 2, 14, 20), (1, 14, 3, 10, 5, 8, 16), (1, 18, 2, 19, 4, 8, 20), (2, 15, 3, 12, 4, 13, 16), (2, 17, 5, 11, 6, 7, 20), (3, 13, 5, 14, 4, 15, 16), (3, 17, 6, 8, 7, 10, 19), (3, 18, 5, 12, 7, 11, 20), (4, 16, 6, 10, 8, 9, 20), (4, 17, 7, 13, 6, 9, 18), (5, 15, 7, 14, 6, 12, 19), (5, 16, 9, 11, 8, 12, 20), (6, 15, 8, 14, 9, 12, 18), (6, 19, 7, 18, 10, 13, 20), (8, 13, 9, 15, 10, 16, 18), (8, 17, 11, 13, 15, 18, 19), (9, 17, 10, 11, 14, 16, 19), (10, 12, 14, 18, 17, 16, 20),$

$(11, 12, 17, 14, 13, 19, 15), (11, 16, 12, 15, 20, 17, 19), (13, 17, 15, 14, 19, 20, 18)\}$ . Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 7-cycle decompositions of  $K_{21} \setminus K_7$ .  $\square$

**Lemma 4.3** *A pair of orthogonal 7-cycle systems of order  $v$  exists for all  $v \equiv 1$  or  $7 \pmod{14}$ , except  $v = 7$ .*

**Proof** A pair of orthogonal 7-cycle systems of order 7 does not exist since a 7-cycle system of order 7 has 3 blocks. A pair of orthogonal 7-cycle systems of order 15 is given in [7]. The case  $v = 21$  is given in Example 4.1 and the cases  $v = 29, 35$  are given in the Appendix. The following construction works for all  $v = 14k + r$  and all  $k \geq 3$ , where  $r = 1$  or  $7$ . Let  $(Q, \circ)$  be a commutative quasigroup of order  $2k$ , for  $k \geq 3$ , with holes  $H = \{h_1, h_2, \dots, h_k\}$ , where  $h_i = \{2i - 1, 2i\}$  for  $i = 1, 2, \dots, k$ . Let  $V = \bigcup_{j=1}^r \{\infty_j\} \cup (Q \times \{1, 2, 3, 4, 5, 6, 7\})$ , where  $r = 1$  or  $7$ . Let  $C_1$  and  $C_2$  be the following collections of 7-cycles on the set  $V$ :

(1) Consider a pair of orthogonal 7-cycle systems of order  $14 + r$ . On  $\bigcup_{j=1}^r \{\infty_j\} \cup (\{h_i\} \times \{1, 2, 3, 4, 5, 6, 7\})$  when  $r = 1$  for  $i = 1, 2, \dots, k$  and when  $r = 7$  for  $i = 1$ , place a copy of the 7-cycles of the system of order  $14 + r$  in  $C_1$  and the 7-cycles of its orthogonal system in  $C_2$ .

(2) When  $r = 7$ , for each of the remaining holes  $h_i$  with  $i = 2, 3, \dots, k$ , consider a pair of orthogonal 7-cycle decompositions of  $K_{21} \setminus K_7$ . On  $\bigcup_{j=1}^7 \{\infty_j\} \cup (\{h_i\} \times \{1, 2, 3, 4, 5, 6, 7\})$  for  $i = 2, 3, \dots, k$ , place a copy of the 7-cycles of the first system in  $C_1$  and the 7-cycles of its orthogonal system in  $C_2$ , being sure that the vertex set of  $K_7$  is  $\bigcup_{j=1}^7 \{\infty_j\}$ .

(3) If  $x$  and  $y$  belong to different holes of  $H$  with  $\{x, x'\} \in H$  and  $\{y, y'\} \in H$ . Then place the 7-cycles  $((x, i), (y, i), (x, i + 1), (y, i + 3), (x \circ y, i + 6), (x, i + 3), (y, i + 1)) \in C_1$  and  $((x, i), (y, i), (x', i + 3), (y, i + 4), (x \circ y, i + 6), (x, i + 4), (y', i + 3)) \in C_2$  for  $i = 1, 2, 3, 4, 5, 6, 7$  modulo 7.

Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 7-cycle systems of order  $v$ .  $\square$

## 5 Orthogonal 8-cycle systems

The spectrum of 8-cycle systems is the set of all  $v \equiv 1 \pmod{16}$ . There exists a pair of orthogonal 8-cycle systems of order  $16k + 1$  for all  $16k + 1 \geq 129$  [7].

The construction we will give to obtain the missing small orders work also for all  $v = 16k + 1 \geq 33$ .

**Example 5.1** *A pair of orthogonal 8-cycle decompositions of  $K_{16,16}$  exists.*

Let  $X = \{(i, 0) : i \in \mathbb{Z}_{16}\}$  and  $Y = \{(i, 1) : i \in \mathbb{Z}_{16}\}$  be the parts of  $K_{16,16}$ ,  $C_1$  and  $C_2$ , each, be a collection of 8-cycles obtained by developing the base 8-cycles  $((0, 0), (0, 1), (1, 0), (2, 1), (6, 0), (1, 1), (4, 0), (6, 1))$  and  $((0, 0), (0, 1), (6, 0), (7, 1), (4, 0), (3, 1), (7, 0), (5, 1))$ , respectively (mod 16,  $-$ ). Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 8-cycle decompositions of  $K_{16,16}$ .  $\square$

**Lemma 5.2** *A pair of orthogonal 8-cycle systems of order  $v$  exists for all  $v \equiv 1 \pmod{16}$ .*

**Proof** A pair of orthogonal 8-cycle systems of order 17 is given in [7]. For all  $v = 16k + 1 \geq 33$ , let  $X = \{\infty\} \cup \{(i, j) \mid 1 \leq i \leq k, 1 \leq j \leq 16\}$ .

(1) Consider a pair of orthogonal 8-cycle systems of order 17. For  $1 \leq i \leq k$ , on each set  $\{\infty\} \cup \{(i, j) \mid 1 \leq j \leq 16\}$ , place a copy of the 8-cycles of the first system in  $C_1$  and the 8-cycles of its orthogonal system in  $C_2$ .

(2) For each pair of  $x, y \in \{1, 2, \dots, k\}$ , consider a pair of orthogonal 8-cycle decompositions of  $K_{16,16}$  on  $\{(x, j) \mid 1 \leq j \leq 16\} \cup \{(y, j) \mid 1 \leq j \leq 16\}$  and place the 8-cycles of the first system in  $C_1$  and the 8-cycles of its orthogonal system in  $C_2$ .

Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 8-cycle systems of order  $v$ .  $\square$

## 6 Orthogonal 9-cycle systems

The spectrum of 9-cycle systems is the set of all  $v \equiv 1$  or  $9 \pmod{18}$ . There exists a pair of orthogonal cyclic 9-cycle systems of order  $72k + 55$  for all  $72k + 55 \geq 199$  [16]. The constructions we will give work for all  $v \equiv 1$  or  $9 \pmod{18}$  when  $v \geq 55$ .

**Example 6.1** *A pair of orthogonal 9-cycle systems of order 27 exists.*

Let  $V = \mathbb{Z}_{26} \cup \{\infty\}$  and

$C_1 = \{(i, 1 + i, 3 + i, 6 + i, 10 + i, 2 + i, 18 + i, 24 + i, 5 + i) : 0 \leq i \leq 26\} \cup \{(\infty, 2j, 9 + 2j, 20 + 2j, 6 + 2j, 19 + 2j, 2 + 2j, 13 + 2j, 25 + 2j) : 0 \leq j \leq 13\}$  modulo 26,

$C_2 = \{(i, 5+i, 3+i, 9+i, 25+i, 6+i, 14+i, 2+i, 11+i) : 0 \leq i \leq 26\} \cup \{(\infty, 2j, 1+2j, 5+2j, 8+2j, 21+2j, 18+2j, 14+2j, 13+2j) : 0 \leq j \leq 13\}$  modulo 26. Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 9-cycle systems of order 27.  $\square$

**Example 6.2** *A pair of orthogonal 9-cycle decompositions of  $K_{27} \setminus K_9$  exists.*

Let  $V = \mathbb{Z}_{27}$ , where the vertex set of the hole  $K_9$  is  $V = \{0, 1, 2, 3, 4, 5, 6, 7, 8\}$  and

$C_1 = \{(0, 9, 1, 12, 6, 13, 3, 15, 14), (1, 10, 2, 13, 7, 14, 4, 16, 15), (2, 11, 3, 14, 8, 15, 5, 17, 16), (3, 12, 4, 15, 0, 16, 6, 9, 17), (4, 13, 5, 16, 1, 17, 7, 10, 9), (5, 14, 6, 17, 2, 9, 8, 11, 10), (6, 15, 7, 9, 3, 10, 0, 12, 11), (7, 16, 8, 10, 4, 11, 1, 13, 12), (8, 17, 0, 11, 5, 12, 2, 14, 13), (0, 18, 1, 21, 6, 22, 3, 24, 23), (1, 19, 2, 22, 7, 23, 4, 25, 24), (2, 20, 3, 23, 8, 24, 5, 26, 25), (3, 21, 4, 24, 0, 25, 6, 18, 26), (4, 22, 5, 25, 1, 26, 7, 19, 18), (5, 23, 6, 26, 2, 18, 8, 20, 19), (6, 24, 7, 18, 3, 19, 0, 21, 20), (7, 25, 8, 19, 4, 20, 1, 22, 21), (8, 26, 0, 20, 5, 21, 2, 23, 22), (9, 19, 12, 23, 17, 21, 24, 20, 18), (10, 20, 13, 24, 9, 22, 25, 21, 19), (11, 21, 14, 25, 10, 23, 26, 22, 20), (12, 22, 15, 26, 11, 24, 18, 23, 21), (13, 23, 16, 18, 12, 25, 19, 24, 22), (14, 24, 17, 19, 13, 26, 20, 25, 23), (15, 25, 9, 20, 14, 18, 21, 26, 24), (16, 26, 10, 21, 15, 19, 22, 18, 25), (17, 18, 11, 22, 16, 20, 23, 19, 26), (0, 13, 9, 5, 18, 10, 12, 17, 22), (1, 14, 9, 11, 7, 20, 12, 15, 23), (2, 15, 9, 12, 8, 21, 13, 10, 24), (3, 16, 9, 23, 11, 13, 15, 17, 25), (4, 17, 10, 6, 19, 14, 16, 12, 26), (9, 21, 16, 11, 15, 10, 22, 14, 26), (10, 14, 11, 17, 20, 15, 18, 13, 16), (11, 19, 16, 24, 12, 14, 17, 13, 25)\},$

$C_2 = \{(0, 9, 19, 2, 13, 5, 21, 20, 23), (1, 10, 20, 3, 14, 6, 22, 21, 24), (2, 11, 21, 4, 15, 7, 23, 22, 25), (3, 12, 22, 5, 16, 8, 24, 23, 26), (4, 13, 23, 6, 17, 0, 25, 24, 18), (5, 14, 24, 7, 9, 1, 26, 25, 19), (6, 15, 25, 8, 10, 2, 18, 26, 20), (7, 16, 26, 0, 11, 3, 19, 18, 21), (8, 17, 18, 1, 12, 4, 20, 19, 22), (0, 14, 20, 25, 16, 9, 17, 1, 13), (1, 15, 21, 26, 17, 10, 9, 2, 14), (2, 16, 22, 18, 9, 11, 10, 3, 15), (3, 17, 23, 19, 10, 12, 11, 4, 16), (4, 9, 24, 20, 11, 13, 12, 5, 17), (5, 10, 25, 21, 12, 14, 13, 6, 9), (6, 11, 26, 22, 13, 15, 14, 7, 10), (0, 21, 2, 22, 11, 1, 25, 12, 15), (1, 22, 3, 23, 12, 2, 26, 13, 16), (2, 23, 4, 24, 13, 3, 18, 14, 17), (3, 24, 5, 25, 14, 4, 19, 15, 9), (4, 25, 6, 26, 15, 5, 20, 16, 10), (5, 26, 7, 18, 16, 6, 21, 17, 11), (6, 18, 8, 19, 17, 7, 22, 9, 12), (7, 19, 0, 20, 9, 8, 23, 10, 13), (8, 20, 1, 21, 10, 0, 24, 11, 14), (0, 12, 7, 11, 8, 13, 9, 23, 16), (0, 18, 11, 15, 16, 19, 12, 20, 22), (1, 19, 6, 24, 16, 21, 3, 25, 23), (2, 20, 7, 25, 9, 26, 4, 22, 24), (5, 18, 13, 17, 16, 14, 9, 21, 23), (8, 12, 18, 15, 23, 11, 19, 14, 26), (8, 15, 22, 10, 26, 12, 24, 19, 21), (10, 14, 22, 17, 25, 13, 19, 26, 24),$

$(10, 15, 17, 20, 13, 21, 14, 23, 18), (11, 16, 12, 17, 24, 15, 20, 18, 25)\}$ . Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 9-cycle decompositions of  $K_{27} \setminus K_9$ .  $\square$

**Example 6.3** *A pair of orthogonal 9-cycle systems of order 37 exists.*

Let  $V = \mathbb{Z}_{37}$ ,  $B_1$  and  $B_2$  contain the two 9-cycles that will be developed modulo 37 to produce the 74 9-cycles in  $C_1$  and  $C_2$ .

$$B_1 = \{(0, 2, 5, 9, 14, 20, 3, 10, 29), (0, 9, 19, 30, 5, 18, 32, 17, 1)\}.$$

$B_2 = \{(0, 2, 11, 14, 33, 6, 10, 24, 30), (0, 5, 16, 29, 14, 26, 10, 30, 1)\}$ . Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 9-cycle systems of order 37.  $\square$

**Example 6.4** *A pair of orthogonal 9-cycle systems of order 45 exists.*

Let  $V = \mathbb{Z}_{11} \times \{1, 2, 3, 4\} \cup \{\infty\}$ . Let  $B_1$  and  $B_2$  contain the 10 base 9-cycles that will be developed modulo  $(11, -)$  to produce the 110 9-cycles in  $C_1$  and  $C_2$ .

$$B_1 = \{(\infty, (1, 2), (8, 2), (0, 1), (7, 3), (10, 1), (0, 2), (5, 4), (8, 4)), (\infty, (8, 1), (3, 2), (0, 3), (10, 2), (6, 4), (0, 4), (4, 1), (5, 3)), ((0, 1), (1, 1), (1, 2), (2, 1), (2, 4), (0, 4), (5, 1), (8, 3), (5, 3)), ((0, 1), (4, 1), (2, 4), (1, 1), (3, 2), (5, 2), (5, 4), (1, 4), (10, 3)), ((0, 2), (7, 1), (10, 1), (6, 2), (1, 3), (8, 3), (7, 4), (10, 3), (10, 4)), ((0, 3), (10, 3), (10, 1), (1, 3), (1, 2), (6, 2), (4, 3), (9, 1), (1, 4)), ((0, 4), (4, 3), (0, 1), (2, 4), (10, 3), (8, 2), (6, 4), (2, 1), (10, 4)), ((0, 4), (7, 3), (4, 2), (1, 4), (8, 2), (10, 1), (5, 1), (7, 1), (5, 3)), ((0, 2), (6, 4), (8, 3), (1, 2), (3, 4), (9, 1), (3, 2), (7, 3), (1, 4)), ((0, 2), (8, 1), (7, 4), (4, 2), (1, 2), (2, 2), (7, 3), (1, 3), (10, 3))\}.$$

$B_2 = \{((1, 2), (8, 2), (5, 3), (8, 3), (4, 4), (5, 2), (10, 2), (8, 4), (4, 2)), ((3, 2), (0, 1), (8, 2), (2, 1), (4, 2), (10, 1), (5, 1), (3, 1), (5, 3)), ((0, 1), (7, 3), (6, 1), (9, 3), (9, 1), (2, 3), (7, 4), (8, 1), (6, 3)), (10, 1), (7, 3), (2, 1), (1, 3), (0, 2), (6, 3), (8, 2), (8, 3), (6, 4)), ((10, 1), (0, 2), (7, 4), (1, 1), (10, 4), (2, 3), (3, 3), (9, 4), (8, 2)), ((0, 2), (5, 4), (10, 4), (1, 4), (10, 3), (9, 4), (8, 3), (0, 4), (9, 2)), ((0, 1), (1, 1), (8, 2), (0, 3), (5, 3), (1, 2), (1, 4), (4, 4), (4, 1)), ( $\infty$ , (1, 1), (2, 4), (10, 1), (7, 4), (5, 1), (8, 1), (8, 2), (9, 2)), ((1, 4), (0, 4), (7, 1), (0, 2), (3, 4), (6, 2), (2, 3), (4, 3), (8, 3)), ((0, 1), (5, 4), (9, 4),  $\infty$ , (0, 3), (0, 4), (5, 2), (4, 3), (10, 2))\}$ . Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 9-cycle systems of order 45.  $\square$

**Example 6.5** *A pair of orthogonal 9-cycle decompositions of  $K_{9,9,9}$  exists.*

Let  $i, j$  be integers and  $\{(i, j) : 0 \leq i \leq 8\}$  be the groups of  $K_{9,9,9}$  for  $0 \leq j \leq 2$ . Let  $C_1 = \{((0 + i, 0 + j), (0 + i, 1 + j), (1 + i, 2 + j), (3 + i, 0 + j), (6 + i, 1 + j), (2 + i, 2 + j), (6 + i, 0 + j), (4 + i, 1 + j), (3 + i, 2 + j)) : 0 \leq i \leq 8 \text{ and } 0 \leq j \leq 2\}$  and  $C_2 = \{((0 + i, 0 + j), (0 + i, 1 + j), (2 + i, 2 + j), (7 + i, 0 + j), (5 + i, 1 + j), (6 + i, 2 + j), (3 + i, 0 + j), (7 + i, 1 + j), (1 + i, 2 + j)) : 0 \leq i \leq 8 \text{ and } 0 \leq j \leq 2\}$ , (mod 9, mod 3). Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 9-cycle decompositions of  $K_{9,9,9}$ .  $\square$

**Lemma 6.6** *A pair of orthogonal 9-cycle systems of order  $v$  exists  $v \equiv 1$  or  $9 \pmod{18}$ , except  $v = 9$ .*

**Proof** A pair of orthogonal 9-cycle systems of order 9 does not exist since a 9-cycle system of order 9 has 4 blocks. A pair of orthogonal 9-cycle systems of order 19 is given in [7]. The case  $v = 27, 37$  and  $45$  are given in Examples 6.1, 6.3 6.4. For all  $v = 18k + r$  and all  $k \geq 3$ , where  $r = 1$  or  $9$ , let  $V = \bigcup_{j=1}^r \{\infty_j\} \cup (X \times \{l : 1 \leq l \leq 9\})$ , where  $|X| = 2k$ . Let  $(X, G, B)$  be a  $3 - GDD$  of type  $2^k$  when  $2k \equiv 0$  or  $2 \pmod{6}$  and a  $3 - GDD$  of type  $4^1 2^{k-2}$  when  $2k \equiv 4 \pmod{6}$ . They exist for  $k \geq 3$  [13]. Let  $C_1$  and  $C_2$  be the following collection of 9-cycles on the set  $V$ :

(1) Consider a pair of orthogonal 9-cycle systems of order 19, 27, 37 and 45 as needed.

On  $\bigcup_{j=1}^r \{\infty_j\} \cup (g_1 \times \{l : 1 \leq l \leq 9\})$ , place a copy of the 9-cycles of the needed system in  $C_1$  and the 9-cycles of its orthogonal system in  $C_2$ .

(2) Consider a pair of orthogonal 9-cycle decompositions of  $K_{27} \setminus K_r$ . For each group  $g_i \in G$ , when  $i \geq 2$ , on  $\bigcup_{j=1}^r \{\infty_j\} \cup (\{g_i\} \times \{l : 1 \leq l \leq 9\})$ , place a copy of the 9-cycles of the needed decomposition in  $C_1$  and the 9-cycles of its orthogonal decomposition in  $C_2$ , being sure that the vertex set of  $K_9$  is  $\bigcup_{j=1}^9 \{\infty_j\}$ .

(3) For each triple  $\{x, y, z\} \in B$ , on  $\{x, y, z\} \times \{l : 1 \leq l \leq 9\}$ , consider a pair of orthogonal 9-cycle decompositions of  $K_{9,9,9}$  whose parts are  $x \times \{l : 1 \leq l \leq 9\}$ ,  $y \times \{l : 1 \leq l \leq 9\}$ ,  $z \times \{l : 1 \leq l \leq 9\}$ , and place a copy of the 9-cycles in  $C_1$  and the 9-cycles of its orthogonal system in  $C_2$ .

Then  $(V, C_1)$  and  $(V, C_2)$  form a pair of orthogonal 9-cycle systems of order  $v$ .  $\square$

Combining Lemmas 2.3, 3.4, 4.3, 5.2 and 6.6 gives the following result.

**Theorem 6.7** *A pair of orthogonal  $l$ -cycle systems of order  $v$  exists for all admissible  $v$  when  $l = 5, 6, 7, 8$  or  $9$ , except 7-cycle system of order 7 and 9-cycle system of order 9, except  $(l, v) = (7, 7)$  and  $(l, v) = (9, 9)$ .*

## References

- [1] B. Alspach and H. Gavlas, Cycle decompositions of  $K_n$  and  $K_n \setminus I$ , *J. Combin. Theory Ser. B*, **81** (2001), 77–99.
- [2] B. Alspach, K. Heinrich and G.Z. Liu, Orthogonal factorizations of graphs, In J.H. Dinitz and D.R. Stinson, (eds), *Contemporary Design Theory*, Wiley-Intersci. Ser. Discrete Math. Optim., Wiley-Intersci. Publ., Wiley, New York, (1992) 13-40.
- [3] D.S. Archdeacon, Heffter arrays and biembedding graphs on surfaces, *Electron. J. Combin.* **22** (2015), no 1, Paper 1.74, 14 pp.
- [4] D.S. Archdeacon, T. Boothby and J. Dinitz, Tight Heffter arrays exist for all possible values, *J. Combin. Des.* **25** (2017), 5–35.
- [5] D.S. Archdeacon, J.H. Dinitz, D.M. Donovan and E.S. Yazıcı, Square integer Heffter arrays with empty cells, *Des. Codes Cryptogr.* **77** (2015), 409–426.
- [6] M. Buratti, Rotational  $k$ -cycle systems of order  $v < 3k$ ; another proof of the existence of odd cycle systems, *J. Combin. Des.* **11** (2003), no. 6, 433–441.
- [7] A. Burgess, N. Cavenagh, D. Pike, Mutually orthogonal cycle systems, *Ars Math. Contemp.*, **23** (2023), no 2, Paper no 5, 20 pp.
- [8] K. Burrage, D. M. Donovan, N. J. Cavenagh, E. S. Yazıcı, Globally simple Heffter arrays  $H(n; k)$  when  $k \equiv 0, 3 \pmod{4}$ , *Discrete Math.*, **343** (2020), no 5, 111787, 17 pp.

- [9] Y. Caro and R. Yuster, Orthogonal decomposition and packing of complete graphs, *J. Combin. Theory Ser. A* **88** (1999), 93–111.
- [10] Y. Caro and R. Yuster, Orthogonal H-decompositions, *Bull. Inst. Combin. Appl.* **33** (2001), 42–48.
- [11] N. J. Cavenagh, J. H. Dinitz, D. M. Donovan, E. S. Yazıcı, The existence of square non-integer Heffter arrays, *Ars. Math. Contemp.*, **17** (2019), 369–395.
- [12] N. J. Cavenagh, D. Donovan, E. S. Yazıcı, Biembeddings of cycle systems using integer Heffter arrays. *J. Combin. Des.* **28** (2020), no. 12, 900–922.
- [13] C. J. Colbourn and J. H. Dinitz. The CRC Handbook of Combinatorial Designs (2nd edition), CRC Press, 2007.
- [14] S. Costa, S. Della Fiore, A. Pasotti, Non-zero sum Heffter arrays and their applications. *Discrete Math.* 345 (2022), no. 9, Paper no. 112952, 12 pp.
- [15] S. Costa, F. Morini, A. Pasotti, M.A. Pellegrini, A generalization of Heffter arrays, *J. Combin. Des.*, **28** (2020), no 3, 171–206.
- [16] S. Costa, F. Morini, A. Pasotti, M.A. Pellegrini, Globally simple Heffter arrays and orthogonally cyclic cycle decompositions, *Australas. J. Combin.* **72** (2018), 549–593.
- [17] S. Costa, A. Pasotti, On  $\lambda$ -fold relative Heffter arrays and biembedding multigraphs on surfaces. *European J. Combin.* 97 (2021), Paper No. 103370, 21 pp.
- [18] S. Costa, A. Pasotti, M.A. Pelegrini, Relative Heffter arrays and biembeddings, *Ars. Math. Contemp.* **18** (2020), no 2, 241–271.
- [19] J. Dinitz and A. Mattern, Biembedding steiner triple systems and  $n$ -cycle systems on orientable surfaces, *Australas. J. Comb.*, **67** (2017), 327–344.
- [20] J.H. Dinitz and I.M. Wanless, The existence of square integer Heffter arrays, *Ars. Math. Contemp.* **13** (2017), 81–93.



- [21] J. X. Lu, On large sets of disjoint Steiner triple systems. I, II, III, *J. Comb. Theory Ser. A*, **34** (1983), 140–182.
- [22] J. X. Lu, On large sets of disjoint Steiner triple systems. IV, V, VI, *J. Comb. Theory Ser. A*, **37** (1984), 136–192.
- [23] F. Morini, M. A. Pellegrini, On the existence of integer relative Heffter arrays, *Discrete Math.* 343 (2020), no. 11, 112088, 22 pp.
- [24] F. Morini, M. A. Pellegrini, Magic rectangles, signed magic arrays and integer  $\lambda$ -fold relative Heffter arrays, *Australas. J. Combin.* 80 (2021), 249–280.
- [25] F. Morini, M. A. Pellegrini, Rectangular Heffter arrays: a reduction theorem, *Discrete Math.* 345 (2022), no. 12, Paper No. 113073.
- [26] M. Sajna, Cycle decompositions. III. Complete graphs and fixed length cycles, *J. Comb. Des.* **10** (2002), 27–78.
- [27] L. Teirlinck, A completion of Lu’s determination of the spectrum for large sets of disjoint Steiner triple systems, *J. Comb. Theory Ser. A*, **57** (1991), 302–305.

## 7 Appendix

In the following cases  $(V, C_1)$  and  $(V, C_2)$  will form a pair of orthogonal  $l$ -cycle systems of order  $v$  for considered  $l$  and  $v$ .

**Example 7.1** *A pair of orthogonal 5-cycle systems of order 21 exists.*

Let  $V = \mathbb{Z}_{21}$ ,  $B_1$  and  $B_2$  contain the two 5-cycles that will be developed modulo 21 to produce the 5-cycles in  $C_1$  and  $C_2$ .

$$B_1 = \{(0, 1, 3, 6, 12), (0, 4, 9, 17, 10)\}.$$

$$B_2 = \{(0, 1, 4, 13, 15), (0, 4, 12, 2, 7)\}.$$

□

**Example 7.2** *A pair of orthogonal 5-cycle systems of order 25 exists.*

Let  $V = \mathbb{Z}_5 \times \mathbb{Z}_5$  and  $B_1$  and  $B_2$  each contains the 4 base 5-cycles that the first two base cycles will be developed modulo  $(5, 5)$  and the last two base cycles will be developed modulo  $(5, -)$  to produce the 60 5-cycles in each of  $C_1$  and  $C_2$ .

$$B_1 = \{((0, 0), (1, 0), (4, 1), (3, 3), (1, 1)), ((0, 0), (2, 0), (2, 1), (2, 3), (4, 1)), ((0, 0), (1, 2), (2, 4), (3, 1), (4, 3)), ((0, 0), (2, 1), (4, 2), (1, 3), (3, 4))\}.$$

$$B_2 = \{((0, 0), (1, 0), (4, 2), (3, 3), (1, 2)), ((0, 0), (2, 0), (2, 2), (2, 3), (4, 2)), ((0, 0), (1, 1), (2, 2), (3, 3), (4, 4)), ((0, 0), (2, 2), (4, 4), (1, 1), (3, 3))\}.$$

□

**Example 7.3** *A pair of orthogonal 6-cycle decompositions of  $K_{6,10}$  exists.*

Let  $X = \{0, 1, 2, 3, 4, 5\}$  and  $Y = \{6, 7, \dots, 15\}$  be the parts of  $K_{6,10}$  and

$$C_1 = \{(0, 6, 1, 7, 2, 8), (0, 7, 3, 6, 2, 9), (0, 10, 1, 8, 3, 11), (0, 12, 1, 9, 3, 13), (0, 14, 1, 11, 2, 15), (1, 13, 5, 7, 4, 15), (2, 10, 4, 9, 5, 12), (2, 13, 4, 6, 5, 14), (3, 10, 5, 11, 4, 12), (3, 14, 4, 8, 5, 15)\},$$

$$C_2 = \{(0, 6, 2, 10, 1, 13), (0, 7, 1, 8, 4, 12), (0, 8, 3, 6, 4, 14), (0, 9, 1, 6, 5, 10), (0, 11, 4, 7, 3, 15), (1, 11, 3, 9, 5, 15), (1, 12, 2, 7, 5, 14), (2, 8, 5, 12, 3, 13), (2, 9, 4, 13, 5, 11), (2, 14, 3, 10, 4, 15)\}.$$

□

**Example 7.4** *A pair of orthogonal 6-cycle systems of order 45 exists.*

Let  $V = \{\infty\} \cup \bigcup_{i=0}^3 X_i \cup \bigcup_{j=0}^1 Y_j$ , where  $X_i = \{x_{6i+1}, x_{6i+2}, \dots, x_{6i+6}\}$  and  $Y_j = \{y_{10j+1}, y_{10j+2}, \dots, y_{10j+10}\}$ . Place a pair of orthogonal 6-cycle systems of order 25 and 21 on  $\{\infty\} \cup \bigcup_{i=0}^3 X_i$  and  $\{\infty\} \cup \bigcup_{j=0}^1 Y_j$ , respectively.

Finally, for each  $i = 0, 1, 2, 3, j = 0, 1$ , place a pair of orthogonal 6-cycle decompositions of  $K_{6,10}$  on  $X_i \cup Y_j$ . □

**Example 7.5** *A pair of orthogonal 7-cycle systems of order 29 exists.*

Let  $V = \mathbb{Z}_{29}$ ,  $B_1$  and  $B_2$  contain the two 7-cycles that will be developed modulo 29 to produce the 58 7-cycles in  $C_1$  and  $C_2$ .

$$B_1 = \{(0, 1, 3, 6, 10, 15, 21), (0, 7, 16, 26, 15, 27, 14)\}.$$

$$B_2 = \{(0, 4, 12, 21, 7, 10, 17), (0, 1, 6, 16, 3, 5, 11)\}.$$
□

**Example 7.6** *A pair of orthogonal 7-cycle systems of order 35 exists.*

Let  $V = \mathbb{Z}_{34} \cup \{\infty\}$  and

$$C_1 = \{(i, 3+i, 10+i, 21+i, 19+i, 27+i, 14+i), (i, 1+i, 5+i, 10+i, 19+i, 31+i, 16+i) : 0 \leq i \leq 34\} \cup \{(\infty, 2j, 6+2j, 16+2j, 33+2j, 5+2j, 15+2j) : 0 \leq j \leq 17\} \text{ modulo } 34,$$

$$C_2 = \{(i, 7+i, 19+i, 30+i, 22+i, 28+i, 15+i), (i, 4+i, 6+i, 15+i, 20+i, 30+i, 16+i) : 0 \leq i \leq 34\} \cup \{(\infty, 2j, 1+2j, 4+2j, 21+2j, 18+2j, 17+2j) : 0 \leq j \leq 17\} \text{ modulo } 34. \quad \square$$